Supplementary Material: Learning-Aware Lyapunov Resource Control for Correlated Multi-Agent Markov Learning

Jialin Zhuang and Vincent K. N. Lau

I. Proof of the Same-Stream Actual-Risk Enclosure

Consider the stationary common factor in the main paper, with independent private observation noise and a deterministic probe $d = (q_d, b_d, n_d)$ ending at $T = n_d b_d$. The probe sigma-field $\mathcal{H}_d$ contains its observations; spatial contrasts contain no additional information about the common factor. Under true instance j, Gaussian filtering gives

$$C_T \mid \mathcal{H}_d \sim \mathcal{N}(m_j, P_j), \quad m_j \sim \mathcal{N}(0, s_j^2), \quad s_j^2 = \theta_j - P_j. \tag{S1}$$

The posterior variance P_j is deterministic because the probe schedule is fixed before data. Starting from variance θ_j, the observation update is $P \leftarrow P/(1+q_d P)$; between observations it is $P \leftarrow \lambda^{2b_d} P + (1-\lambda^{2b_d})\theta_j$. The last update has no subsequent prediction.

Fix either downstream program a, including its budget, initialization, sample spacing and deterministic weights. All learning observations occur strictly after T. With $r_a = \lambda^{b_a}$ and the notation of main Eq. (38), its unconditional stationary risk is

$$J_{j,a} = d_a^2 + \theta_j \sum_{t,u=1}^{N_a} \omega_{a,t}\omega_{a,u} r_a^{|t-u|} + q_a^{-1} \sum_{t=1}^{N_a} \omega_{a,t}^2. \tag{S2}$$

Conditional means of future common states are $\lambda^{b_a t} m_j$. Their conditional covariance is their stationary covariance minus $s_j^2 \lambda^{b_a(t+u)}$. Private learning noise is independent of the probe. Therefore

$$\mathbb{E}_j[Z_a^2 \mid \mathcal{H}_d] = J_{j,a} + 2 d_a k_a m_j + k_a^2 (m_j^2 - s_j^2). \tag{S3}$$

This identity includes the posterior mean, which is unbounded across probe histories; no uniform conditional Gaussian bound is invoked.

Let E_j be the event that the decision selects the wrong frozen program, and let $p_j = \Pr_j(E_j) \le \delta$. Subtracting Eq. (S3) for the correct and wrong programs gives

$$\mathbb{E}_j Z_{\widehat{a}}^2 = J_{j,a_c} + p_j \Delta J_j + \mathbb{E}_j[\mathbf{1}_{E_j} U_j], \tag{S4}$$

where

$$U_j = 2(d_{a_w} k_{a_w} - d_{a_c} k_{a_c}) m_j + (k_{a_w}^2 - k_{a_c}^2)(m_j^2 - s_j^2). \tag{S5}$$

The centered Gaussian first and second moments are orthogonal, with variances s_j^2 and $2s_j^4$. Consequently $\mathbb{E}_j U_j = 0$ and $\mathbb{E}_j U_j^2 = \Gamma_j^2$ as defined in the main paper. Cauchy–Schwarz gives $|\mathbb{E}_j[\mathbf{1}_{E_j} U_j]| \le \sqrt{p_j}\Gamma_j \le \sqrt{\delta}\Gamma_j$. Bounding the signed term $p_j \Delta J_j$ and using nonnegative squared risk proves main Theorem 5.

II. Exact Weights of the Implemented Learners

Let $M_a = \min\{\lfloor B'_m/(h+q_a)\rfloor, \lfloor B'_e/b_a\rfloor\}$ be the number of scheduled observations after subtracting probe costs. All M_a observations are paid. The FIFO withholding convention leaves $N_a = (M_a - D)_+$ usable observations, not a free replacement budget. For $N_a = 0$ the reported parameter is one, so $d_a = 1$ and the weight vector is empty.

A. Running mean

The Layer-A output is $(1+\sum_{t=1}^{N_a} \bar{X}_t)/(N_a+1)$. Thus $d_a = 1/(N_a+1)$ and every $\omega_{a,t} = 1/(N_a+1)$. Main Eq. (38) then gives

$$k_a = \frac{r_a(1 - r_a^{N_a})}{(N_a+1)(1-r_a)}. \tag{S6}$$

B. Averaged FIFO affine learner

In Layer B, an increment $\eta(\bar{X}_t - \mu w_{t-1})$ is computed before enqueueing and is applied after D opportunities. The original implementation fixes $\mu = 1/2$ and $\eta = 0.1/[\mu(D+1)]$, starts from parameter one with an empty queue, and averages the applied parameters. Indexing these applied parameters as z_n gives

$$z_n = z_{n-1} - \eta\mu z_{n-D-1} + \eta \bar{X}_n, \quad z_{-D} = \cdots = z_0 = 1. \tag{S7}$$

Define $h_0 = 1$, $h_t = 0$ for $t < 0$, and $h_t = h_{t-1} - \eta\mu h_{t-D-1}$ for $t \geq 1$. For $N = N_a > 0$, the output $N^{-1}\sum_{n=1}^{N} z_n$ has exact coefficients

$$\omega_{a,t} = \frac{\eta}{N}\sum_{v=0}^{N-t} h_v, \quad d_a = 1 - \mu\sum_{t=1}^{N}\omega_{a,t}. \tag{S8}$$

To verify the deterministic term, constant input $\bar{X}_n = \mu$ keeps the initial history at one; linearity then yields the identity for d_a. Eq. (S8) represents the average of applied iterates; all scheduled observations remain charged.

III. Shrinking-Risk and Budget-Shift Interface

Suppose the finite action catalogue has $|\lambda| \leq 1 - \gamma < 1$, bounded D, and strictly stable affine recursions. For the registered step rule this stability follows directly from the characteristic polynomial $z^{D+1} - z^D + t$, $t = \eta\mu$. At small $t > 0$, the root at one moves inside the unit disk and the other roots remain inside. Writing a first unit-circle crossing as $z = e^{i\omega}$ yields $t = 2\sin(\omega/2)$ and the smallest positive crossing $\omega = \pi/(2D+1)$. Thus $0 < t < 2\sin[\pi/(4D+2)]$ is stable; the registered $t = 0.1/(D+1)$ lies strictly inside this interval. The impulse response consequently decays geometrically, and $\sum_{v\geq 0} h_v = 1/(\eta\mu)$.

The interior weights in Eq. (S8) are $1/(\mu N)$ up to a summable end correction: their difference from these constant weights has ℓ_1 norm $O(N^{-1})$ and maximum absolute value $O(N^{-1})$. The homogeneous transient is summable, so its average d_a is $O(N^{-1})$. The AR(1) covariance kernel has uniformly bounded absolute row sums. Substitution into Eq. (S2) therefore gives

$$J_{j,a} = \frac{1}{\mu^2 N_a}\left[\frac{1}{q_a} + \theta_j\frac{1+r_a}{1-r_a}\right] + O(N_a^{-2}). \tag{S9}$$

The running mean has the same expansion with $\mu = 1$. These constants are uniform over the finite catalogue and bounded instance class. On a budget ray, $N_a = s/c_a + O(v+D+1)$ with $c_a = \max\{(h+q_a)/\beta_m, b_a/\beta_e\}$ and fixed probe charge v. Hence $J_{j,a} = K_{j,a}/s + O((v+D+1)/s^2)$, with

$$K_{j,a} = \frac{c_a}{\mu^2}\left[\frac{1}{q_a} + \theta_j\frac{1+r_a}{1-r_a}\right]. \tag{S10}$$

Subtracting a fixed probe charge changes N_a by at most $O(v+D+1)$, giving the claimed budget-shift remainder. Moreover $k_a = O(N_a^{-1})$ because the weights are uniformly $O(N_a^{-1})$ and $\sum_{t\geq 1}|r_a|^t$ is bounded. Since $s_j^2 \leq \theta_j$, main Eq. (39) yields

$$\sqrt{\delta}\Gamma_j = O(\sqrt{\delta}/s^2). \tag{S11}$$

For a pair of frozen actions satisfying the main phase theorem's leading-gap and wrong-decision assumptions, adding this remainder to admission changes only its second-order class constants. The same necessary-information and sufficient-scale argument then applies to the corrected actual-risk rule. This is an averaged, additive Gaussian affine result; it does not remove the nonzero noise floor of a fixed-step unaveraged learner or the conditional-history requirements of the general Markov theorem.

TABLE I
Population enclosures for mean information-only minus learning-aware risk, rounded outward. All means use the listed complete stratum.

Layer	Stratum	Cells	Lower	Upper
A	All	1184	.0506	.0706
A	Intermediate	576	.0770	.1180
A	Original practical	446	.1607	.1952
A	Search-censored	32	.4879	.4880
B	All	608	.0079	.0223
B	Intermediate	288	-.0049	.0252
B	Original practical	222	.0562	.0773
B	Search-censored	32	.1957	.1958

IV. Complete Existing-Evidence Reanalysis

The calculation uses all registered EXP-016B budget points, both regimes and all eight policies: 14,336 policy cells, each with its original 192-seed empirical mean. Layer A contains 104 scenario families and Layer B its original 56, including eight search-censored families in each layer. For fixed policies Eq. (S2) is exact. For probe policies main Eq. (40) supplies population-risk bounds. When information-only and learning-aware policies execute the same frozen program, their paired difference is identically zero, not the difference of two independent interval bounds. Otherwise subtract their one-sided bounds.

Below identification, all 208 Layer-A and 112 Layer-B comparisons are identical programs; above finite value thresholds the corresponding counts are 384 and 192. In the original practical subset, 323/446 Layer-A and 176/222 Layer-B cells have strictly positive improvement lower bounds, while 101 and 33 have strictly negative upper bounds; the remaining 22 and 13 are inconclusive. These include regime-specific losses even when the stratum-average improvement is positive. No new admission rule was executed and no old gate was replaced. The old probe-length routine used the sufficient exponent $\log(0.5/0.025)$, giving an equal-prior average-error target of 0.025 rather than a directional guarantee of 0.025. The reanalysis uses each probe's exact affinity $\exp(-I_B)$ as a valid bound on either directional error; over all 104 frozen probe designs these bounds range from 0.0352271 to 0.0497465. Failure of this sufficient directional certificate does not establish that the actual

directional errors exceed 0.025. No outcomes or probe lengths are changed to obtain a smaller error bound. This outcome-known reanalysis gives population enclosures, not new confidence intervals for empirical means.

V. Complete Threshold-Sandwich Scale Calculation

For the independently initialized Gaussian running mean, let $c_a = \max\{(h+q)/\beta_m, b/\beta_e\}$ and $r = \lambda^b$. The exact horizon is $N_a = s/c_a + O(1)$, with bounded terminal charges and rounding. Main Eq. (8) and Eq. (13) give $K_j(a) = c_a\{q^{-1}+\theta_j(1+r)/(1-r)\}$ and a uniform $O(s^{-2})$ remainder. Removing probe charge v changes N_a by $O(v+1)$ and hence risk by $O((v+1)/s^2)$ for $s \geq s_0(v+1)$; compact persistence and the finite catalogue make these constants uniform. Thus the phase-theorem risk conditions have a concrete Gaussian realization, separately from the constant-step affine certificate. Write $d_\delta = \mathrm{kl}(1-\delta,\delta)$, $r_0 = 1-\gamma < 1$, and $Q = \max \mathcal{Q}$. Under either true instance, predictable spacing gives $\|C_{S_k}\|_{L^2} \leq r_0\|C_{S_{k-1}}\|_{L^2} + \sqrt{\theta_+}$, hence a uniform bound $M_C = \sqrt{\theta_+}/(1-r_0)$. The normalized observation $Y_k/\sqrt{q_k}$ has L^2 norm at most $M_C + 1$. For either candidate filter, write its posterior mean as $(1-t_{j,k})m^-_{j,k} + t_{j,k}Y_k/\sqrt{q_k}$, with predictable $0 \leq t_{j,k} \leq 1$. Prediction multiplies this mean by a factor of magnitude at most r_0; the triangle inequality bounds both predictive means uniformly in L^2 by $M_F = r_0(M_C+1)/(1-r_0)$, starting from zero. Both predictive variances are at most θ_+, so $1 \leq V_{j,k} \leq V_+ = 1 + Q\theta_+$. Main Eq. (26) gives the expected, rather than pointwise, bound $\mathbb{E}_j\mathcal{I}_{j,1-j}(Z_k, A_k) \leq i_+$, where $i_+ = (\log V_+ + V_+ + 4QM_F^2)/2$ is finite. Extend a stopped rule predictably after stopping; nonnegativity and its deterministic budget bound $n_{\max}(s) \leq c_B s$ imply $\mathbb{E}_j \sum_{k\leq\tau} \mathcal{I}_{j,1-j} \leq i_+ n_{\max}(s)$. Together with main Eq. (30), this proves $B_N \geq c_N d_\delta$. The fixed probe has information at least $i_- > 0$, so $n_\delta = \lceil i_-^{-1}\log(1/\delta)\rceil$ observations suffice and cost at most $c_p n_\delta + D_{\max}$. Continuity of $\log(1/\delta)/d_\delta$ gives $n_\delta \leq c_0 + c_1 d_\delta \leq c_0 + c_2 B_N$. Write $u_\delta \leq c_p n_\delta + D_{\max}$ for the full probe charge. The assumed expansion and budget-shift inequality give a high-instance improvement over the full-budget fallback certificate of at least $g_{\min}/s - \{2M + L(u_\delta+1)\}/s^2 - \delta w_+/s$. It is strictly positive for $s > \{4M+2L(u_\delta+1)\}/g_{\min}$, also requiring $s \geq s_0(u_\delta+1)$. On instance zero, the assumed coefficient ordering gives $\epsilon_{\mathrm{safe}} \leq \{2M + L(u_\delta+1)\}/s^2 + \delta_0 w_+/s$. Thus $s \geq \sqrt{2\{2M + L(u_\delta+1)\}/\bar{\epsilon} + 2\delta_0 w_+/\bar{\epsilon}}$ suffices. Using $2\sqrt{xy} \leq x+y$, all three requirements are bounded by a class-constant multiple of $n_\delta + B_{\mathrm{oracle}} + B_\epsilon + 1$. Taking the maximum proves the upper bound in main Eq. (42); its lower bound follows because every admitted δ-correct probe obeys both information constraints.

VI. Robust-Certificate Sensitivity Details

This section supplies the proof of main Corollary 2. Write $\delta^+ = C_q(r_q^+)^b$. On the coverage event, $\delta^+ \geq \delta$ and monotonicity on $\mu - 2L\delta^+ > 0$ gives $a^+ \geq a$ and $\beta^+ \geq \beta$. Evaluating the delayed Young inequality at the shared $\lambda^+ = \sqrt{h/a^+}$ proves the upper recursion; separately optimized forcing terms need not be monotone. For $\Delta\delta = \delta^+ - \delta$, $\Delta K = K_q^+ - K_q$, and $\Delta\Omega = \Omega_q^+ - \Omega_q$,

$$\begin{aligned}
\Delta a &= 2\eta^2\Delta K + (2\eta L + 4\eta^2 L^2)\Delta\delta,\\
\Delta\beta &= 2\eta^2\Delta\Omega + 4\eta^2 G^2\Delta\delta\\
&\quad + 4\eta G^2\left[\frac{(\delta^+)^2}{\mu - 2L\delta^+} - \frac{\delta^2}{\mu - 2L\delta}\right],\\
\Delta c &= (\sqrt{a^+}+\sqrt{h})^2 - (\sqrt{a}+\sqrt{h})^2. \qquad \text{(S12)}
\end{aligned}$$

For $h > 0$,

$$\begin{aligned}
\Delta d &\leq (1+\sqrt{h/a^+})\Delta\beta + \sqrt{h}\left(\frac{1}{\sqrt{a}} - \frac{1}{\sqrt{a^+}}\right)\beta\\
&\quad + \frac{\sqrt{a^+}-\sqrt{a}}{\sqrt{h}}g, \qquad \text{(S13)}
\end{aligned}$$

and $\Delta d = \Delta\beta$ when $h = 0$. For $U(c,d) = d/(1-c) + c^n(R_0 - d/(1-c))_+$, $|\partial_d U| \leq (1-\bar{c})^{-1}$ and $|\partial_c U| \leq nR_0 + \bar{d}(1-\bar{c})^{-2}$. The mean-value bound proves main Eq. (44); applying it to the upper- and exact-score minimizers gives $U_{\hat{a}} \leq U_{a^\star} + 2\max_a \varepsilon_a$.

VII. Stochastic-Delay Interpretation

The controlled stochastic delay differential equation associated with one fixed resource action is

$$dE(t) = -A_q E(t-\tau(t))dt + \sqrt{\eta}\,\Sigma_q^{1/2}(E_t)dW(t). \qquad \text{(S14)}$$

Here $\Sigma_q(E_t)$ retains both additive innovation covariance and the state-dependent quadratic variation of random temporal-difference Jacobians. For $V(E) = \|E\|^2$, Itô's formula gives

$$\mathcal{L}_a V(E_t) = -2E(t)^\top A_q E(t-\tau(t)) + \eta\,\mathrm{tr}\,\Sigma_q(E_t). \qquad \text{(S15)}$$

Participation controls spatial covariance, spacing controls mixing, and η controls their scale; the discrete theorem retains finite steps, integer delays and rounding.

VIII. Constructing a Certificate from a Known Finite Model

Let Z be the finite augmented state containing all sources needed for the retained joint TD tuple. For each participation q, suppose its kernel M_q and stationary law π_q are known, and the marginal target policy and reward law are preserved. Tabulate $H_i(z)$ and $c_i(z)$, put $A = \mathbb{E}_{\pi_q}H_i$, $c = \mathbb{E}_{\pi_q}c_i$, and solve $Aw^\star = c$. Check that these means are common to all actors and

all q, and that $\mu = \lambda_{\min}((A + A^\top)/2) > 0$. With $\xi_i(z) = c_i(z) - H_i(z)w^\star$, compute

$$L = \max_{i,z} \|H_i(z)\|_2, \qquad G = \max_{i,z} \|\xi_i(z)\|,$$
$$K_q = \lambda_{\max}\left(\sum_z \pi_q(z)\bar{H}_q(z)^\top \bar{H}_q(z)\right),$$
$$\Omega_q = \sum_z \pi_q(z)\|\bar{\xi}_q(z)\|^2,$$
$$\beta_q(g) = \max_z \frac{1}{2}\sum_{z'} |M_q^g(z, z') - \pi_q(z')|. \tag{S16}$$

Here bars denote actor averages, not averages of squared norms. The integer g is the number of transitions of the augmented tuple chain between the conditioning state and the retained tuple; it must be derived from the declared physical schedule, not identified with a state lag or drain charge.

Proposition 1 (Constructive finite-model interface)**.** *If M_q is irreducible and aperiodic and the above common-mean and positivity checks pass, Assumption 1's boundedness, centering and conditional mixing inputs hold with Eq.* (S16) *and $\delta = \beta_q(g)$. For every action whose resulting μ_δ, a_δ and $c_{\mathrm{aff}} < 1$ pass the main paper's shield, the affine certificate is valid from any declared initial augmented state.*

Proof. The finite maxima bound every sample and stationarity gives $\mathbb{E}_{\pi_q}\xi_i = 0$. Conditional on any earlier history, the Markov property expresses the next tuple law as a mixture of rows of M_q^g. Convexity of total variation bounds its distance from π_q by $\beta_q(g)$, uniformly over that history. The drift proof uses only this total-variation distance, so the geometric envelope can be replaced by this exact finite-gap value. The stated positivity and contraction checks are precisely the remaining hypotheses of the affine bound. □

This is a construction, not a statistical estimator of an unknown kernel. A dense joint-state table may grow exponentially with participation; its offline construction cost is separate from the online scalar score. In the seven-state shared/private-source model, a product coupling gives the cheaper bound $\beta_q(g) \le \min\{1, (q+1)\beta_{\mathrm{pair}}(g)\}$, where $\beta_{\mathrm{pair}}(g) = \max_x \|M_{\mathrm{pair}}^g(x, \cdot) - \pi_{\mathrm{pair}}\|_{\mathrm{TV}}$. This uses the actual consecutive-pair chain and independent latent source chains; fresh selector coins are integrated by data processing. The existing implementation computes these pair kernels and the same-time aggregate moments directly. For a moving target policy, a frozen-kernel table does not certify mixed histories: either supply uniform bounds for the nonhomogeneous kernel products or reset the learner and exclude all pre-reset target updates.

IX. Executable Actual-Risk Gate and Saved-Result Reconstruction

For each fixed candidate probe, Gaussian filtering gives P_j, the formulas above give $(J_{j,a}, d_a, k_a)$, and the probe likelihood laws give a directional error bound δ_j under each hypothesis. The exact Gaussian affinity is one valid choice: a zero-threshold likelihood-ratio decision obeys $\Pr_j(\mathrm{error}) \le \int \sqrt{p_0 p_1}$ in each direction. An equal-prior average-error bound is not interchangeable with this directional bound. Compute both hypothetical-instance quantities before probing; the actual unknown instance is not a gate input.

Proposition 2 (Actual-risk admission)**.** *Fix the two hypotheses, noise law, learner initialization and deterministic linear program weights as in Theorem 5. Given a finite public probe catalogue, choose a feasible design satisfying Eq.* (41) *before any probe observations, with deterministic tie breaking. After its likelihood decision, execute its frozen selected program on the continuing stream. Then $\mathbb{E}_1 Z^2 < F_1$ and $\mathbb{E}_0 Z^2 \le F_0 + \bar{\epsilon}$. If none passes, no probe is executed and the actual risk is exactly F_j.*

Proof. For each fixed design Theorem 5 gives $\mathbb{E}_j Z^2 \le U_j(d)$. Pre-observation deterministic choice leaves its fixed-design hypothesis intact. The two gate inequalities imply the conclusions. In the rejection branch the stationary full-budget commitment defines F_j. □

Thus the procedure is: price the probe, compute the remaining horizons and exact weights for both branch programs, compute U_0, U_1, F_0, F_1, apply the two inequalities, and only then acquire data. An observation-adaptive choice of probe duration or of program weights is not covered by this fixed-design argument.

We reconstruct this gate on the existing EXP-016B outcomes, using its single registered information-only probe candidate and the original scenario-specific allowances 0.1 or 0.2. All 896 public layer/scenario/budget designs and both regimes are retained, including search-censored families. The corrected gate admits 480 designs, versus 505 without the carryover term; it removes 25 admissions. The outcome-known zero-allowance stress case admits none under either rule. It is retained separately: a paid probe can reduce the low-instance learning horizon even with perfect identification.

The gate is a deterministic function of public design quantities, so an admitted design takes the saved information-only outcome and a rejected design takes the saved no-probe fallback outcome. The simulator's potential observations are policy independent and shared across these branches; this reconstructs their empirical means without generating or resampling a trajectory. Table II is therefore a retrospective policy comparison,

TABLE II
SAVED-RESULT GATE RECONSTRUCTION: MEAN SQUARED RISK OVER ALL BUDGET/REGIME CELLS, WITHOUT NEW TRAJECTORIES. EACH ORIGINAL CELL MEAN CONTAINS ALL 192 SAVED SEEDS.

	A high	A low	B high	B low
Full-budget fallback	.63455	.09201	.82703	.54369
Information-only	.54814	.24062	.76422	.62515
Original learning-aware	.57637	.09304	.79121	.55825
No-carryover gate	.52548	.09488	.74204	.56252
Corrected gate	.52989	.09393	.74242	.56068

not fresh execution or independent confirmation. Against fallback, the corrected gate retains $13/592$ and $4/304$ empirically adverse high-instance cells in A and B; finite-sample cell means need not satisfy the population ordering. Its low-instance mean risks exceed fallback by $.00192$ and $.01699$, respectively, within the declared nonzero safety allowances, not a claim of zero-harm safety. The no-carryover ablation has slightly lower high-instance means but higher low-instance means; the correction trades admissions for a valid population-risk guarantee, not uniformly better observed risk. All eight original policies, per-cell costs, both replay variants and their complete cell tables are preserved. The new comparison was specified after earlier outcomes were known and supplies no new confidence interval.

X. NEAREST-WORK ASSUMPTIONS AND OBJECTIVES

The comparison uses the accessible full preprints, with theorem and proof locators recorded in the accompanying source-reading matrix; it is not an exhaustive priority claim. DASA [25], Assumption 3 and Theorem 1, combines independent agent chains with median-based update filtering to obtain average-delay dependence and parallel speedup. Its Lemma 2 uses cross-agent independence in the noise cross terms. Our bounded deterministic delay condition is narrower; our added object is covariance-dependent paid learning risk, not a stronger delay rate.

Perazzone et al. [20], Theorem 1 and Eqs. (12)–(23), link inverse participation probabilities to a nonconvex convergence bound, then optimize a time-average convergence/communication surrogate with power queues. Their solver and instantaneous-channel formulation address a different operational problem; our finite-budget probe debit and same-stream decision-event correction do not follow from that time-average objective. Yu et al. [26], Sections III–IV and Propositions 12–14, use an SDDE/arrival-process approximation to optimize worker activation, including a finite-service queue. That queue model and its delay law are not established by our fixed-delay experiment.

Saad et al. [32], Assumption 1, Protocol 1 and Theorems 4.1/5.1, adapt joint arm queries to unknown covariance with independent rounds and a fixed-confidence identification objective. Our dependence hypotheses are supplied, rather than learned nonparametrically; our comparison instead prices identification through the subsequent learner. Li et al. [35], Theorem 4.12, remove unknown mixing/eigenvalue inputs from a regularized TD step schedule. That reduces tuning requirements that our general confidence interface still has, but does not construct a calibrated multi-action terminal-risk shield. Li et al. [36], Assumptions 1–7 and Theorem 1, give shrinking-tube reliability for projected adaptive-kernel SA using uniform drift/minorization and kernel continuity. Their mixed-history result is broader than a frozen-kernel table; our Gaussian decision correction and paid finite horizon are different outputs, not a replacement for their regularity assumptions. Reference numbers in this supplement refer to the main paper's bibliography.

XI. COMPLETE TRACKING PROOF

Set $R_\star = d/(1-c)$ and $\Gamma_r = (Z_r - R_\star)_+$. The explicitly assumed uniform-history drift and the positive-part replacement argument give

$$\begin{aligned} \widetilde{Z}_{r+1} &\le R_\star + c\Gamma_r, \\ \Gamma_{r+1} &\le (1+\chi)c\Gamma_r + \chi R_\star + (1+\chi^{-1})v^2. \end{aligned} \tag{S17}$$

The second line follows by applying Young's inequality to the target shift and subtracting $R_\star$. Iteration and $\Gamma_0 \le Z_0$ prove Eq. (20) whenever $\bar{c} = (1+\chi)c < 1$. Finally, $\widehat{A}_w - A^\star = \{\gamma\phi(s') - \phi(s)\}^\top (w - w_n^\star)$ and $\|\gamma\phi(s') - \phi(s)\| \le (1+\gamma)\Phi$; Cauchy–Schwarz proves Eq. (21).

XII. COMPLETE OCCUPATION-MEASURE PROOF

Let $\Delta_j(z,a) \ge 0$ be the one-step opportunity loss relative to the instance-j certified oracle. For a stopped predictable policy, define

$$\nu_j(G,a) = \mathbb{E}_j^\pi \sum_{k \le \tau} \mathbf{1}\{Z_k \in G, A_k = a\}. \tag{S18}$$

Every δ-correct policy induces an occupation measure satisfying the controlled Kalman flow constraints,

$$\begin{aligned} \int \mathcal{I}_{j,1-j}(z,a)\nu_j(dz,da) &\ge \mathrm{kl}(1-\delta,\delta), \\ \int (h + q(a))\nu_j(dz,da) &\le B_{\mathrm{msg}}, \\ \int b(a)\nu_j(dz,da) + \ell\mathbb{P}_j(\tau > 0) &\le B_{\mathrm{env}}. \end{aligned} \tag{S19}$$

Let T_a be the belief transition and ζ_j the terminal belief law. For every bounded measurable f, predictability and stopped telescoping give

$$\int f(z)\zeta_j(dz) - f(z_1) = \int \Big[\int f(z')T_a(z, dz') - f(z) \Big] \nu_j(dz, da). \tag{S20}$$

Moreover,

$$\mathbb{E}_j^{\pi} \sum_{k \le \tau} \Delta_j(Z_k, A_k) = \int \Delta_j(z, a)\nu_j(dz, da). \tag{S21}$$

Eq. (29), Eq. (6), and Eq. (S20) place every admissible occupation measure in the controlled feasible set. Taking the infimum of its exact loss over that larger set proves Proposition 4.