

# Homework 3

MacMillan, Kyle

October 12, 2018

# Contents

Title

Table of Contents

1 Problem 6.5

2 Problem 6.8

i

1

5

## 1 Problem 6.5

Figure 1 shows the required plot.




Figure 1: Problem 6.5.a

Figure 2 shows the data points that were saved. The code is as follows:

```
def ddstep(self, xc, yc, qc, r, l, dt, w1, w2):
    xn = xc + (r * dt / 2.0) * (w1 + w2) * np.cos(qc)
    yn = yc + (r * dt / 2.0) * (w1 + w2) * np.sin(qc)
    qn = qc + (r * dt / (2.0 * l)) * (w1 - w2)
    return (xn, yn, qn)

def d():
    runs = 100
    iterations = int(5.0 / dt + 1)
    x = []
    y = []
    for i in range(runs):
        xerr = np.random.normal(mu, sigma, iterations)
        yerr = np.random.normal(mu, sigma, iterations)
        j = 0
        while (t < 5.0 - dt):
            xc, yc, qc = ddstep(xc + xerr[j],
                                yc + yerr[j],
                                qc,
                                r,
                                L,
                                dt,
                                w1 * t * t,
                                w2 * t)

            t = t + dt
            j += 1

        x.append(xc)
        y.append(yc)

    f = open("problem6-5-b.txt", "w")
    f.write('x, y\n')
    for i in range(len(x)):
        x = str(x[i])
        y = str(y[i])
        f.write(x + ' ' + y + '\n')
```




Figure 2: Problem 6.5.b

Figure 3 shows the data points that were saved.




Figure 3: Problem 6.5.c

Figure 4 shows the data points that were saved.




Figure 4: Problem 6.5.d

## 2 Problem 6.8

For this problem we have to take into account  $x$ ,  $y$ , and  $\theta$  derivatives with respect to time for the parametric form of an ellipse. We also make the assumption that velocity is constant. Parametric forms of  $x$ ,  $y$ , and  $\theta$  are as follows:

$$x(t) = 3 + 4\cos(t)$$

$$y(t) = 2 + 3\sin(t)$$

$$\theta(t) = \pi + \phi(t)$$

The derivatives are as follows:

$$\hat{x}(t) = -4\sin(t)$$

$$\hat{y}(t) = 3\cos(t)$$

$$\hat{\theta}(t) = \frac{12\sec^2(t)}{9\tan^2(t) + 16}$$

The only reason  $\hat{\theta}(t)$  is differentiable without chaos is because as the robot moves around, the angle is independent of the starting  $x$  and  $y$  position, so the equation used to calculate  $\hat{\theta}(t)$  are as follows:

$$\theta(t) = \pi + \arctan\left(\frac{3\sin(t)}{4\cos(t)}\right)$$

You then plug and chug those into the [inverse kinematic equations](#) to get wheel speed of each wheel at every time step.