


Measures of entanglement

Khoruzhii K.

05.07.2024



- Pure states $|\psi_{AB}\rangle$:
 - *Product state* if $|\psi_{AB}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$
 - A is entangled with B otherwise
- Mixed states $\hat{\rho}_{AB}$
 - *Separable state* if $\hat{\rho}_{AB} = \sum_k p_k \hat{\rho}_A^k \otimes \hat{\rho}_B^k$
 - A is entangled with B otherwise




Bipartite entanglement in pure states: SVD



Is this state entangled?

$$|\psi_{AB}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |01\rangle)$$

Bipartite entanglement in pure states: SVD




Is this state entangled?

$$|\psi_{AB}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |01\rangle)$$

Of course not:

$$|\psi_{AB}\rangle = |0\rangle \otimes \frac{1}{\sqrt{2}} (|0\rangle + |1\rangle)$$


Bipartite entanglement in pure states: SVD



What about this?

$$|\psi_{AB}\rangle = \frac{1}{2} (|00\rangle + |10\rangle - |01\rangle - |11\rangle)$$

Bipartite entanglement in pure states: SVD




What about this?

$$|\psi_{AB}\rangle = \frac{1}{2} (|00\rangle + |10\rangle - |01\rangle - |11\rangle)$$

After some SVD (or **Schmidt decomposition**):

$$|\psi_{AB}\rangle = \sum_{a \in A} \sum_{b \in B} \psi_{ab} |a\rangle \otimes |b\rangle = \sum_a \sum_b \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}_{ab} |ab\rangle$$

Bipartite entanglement in pure states: SVD



What about this?


$$|\psi_{AB}\rangle = \frac{1}{2} (|00\rangle + |10\rangle - |01\rangle - |11\rangle)$$

After some SVD (or **Schmidt decomposition**):

$$|\psi_{AB}\rangle = \sum_{a \in A} \sum_{b \in B} \psi_{ab} |a\rangle \otimes |b\rangle = \sum_a \sum_b \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}_{ab} |ab\rangle$$

$$U \begin{pmatrix} \sqrt{\lambda_1} & 0 \\ 0 & \sqrt{\lambda_2} \end{pmatrix} V = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & . \\ 1 & . \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & . \\ -1 & . \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}$$

Bipartite entanglement in pure states: SVD



What about this?

$$|\psi_{AB}\rangle = \frac{1}{2} (|00\rangle + |10\rangle - |01\rangle - |11\rangle)$$


After some SVD (or **Schmidt decomposition**):

$$|\psi_{AB}\rangle = \sum_{a \in A} \sum_{b \in B} \psi_{ab} |a\rangle \otimes |b\rangle = \sum_a \sum_b \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}_{ab} |ab\rangle$$

$$|\psi_{AB}\rangle = \frac{1}{\sqrt{2}} (|0\rangle + |1\rangle) \otimes \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle)$$

We can see, that it is separable.

Bipartite entanglement in pure states: SVD



Schmidt decomposition


$$|\psi_{AB}\rangle = \sum_{a \in A} \sum_{b \in B} \psi_{ab} |a\rangle \otimes |b\rangle = \sum_j \sqrt{\lambda_j} |\psi_{A,j}\rangle |\psi_{B,j}\rangle$$

deeply connected with the reduced density matrix $\rho_{A,B}$

$$\rho_{B,A} = \text{tr}_{A,B} (|\psi_{AB}\rangle\langle\psi_{AB}|) = \sum_i \lambda_i |\psi_{B,A}\rangle\langle\psi_{B,A}|.$$

If there is no entanglement, then

$$\rho_{A,B} = |\psi_{A,B}\rangle\langle\psi_{A,B}|$$




Which one is more entangled?

$$|\psi_{AB}^I\rangle = \frac{1}{\sqrt{4}} (|00\rangle + |10\rangle - |01\rangle - |11\rangle),$$

$$|\psi_{AB}^{II}\rangle = \frac{1}{\sqrt{3}} (|00\rangle + |01\rangle + |10\rangle),$$

$$|\psi_{AB}^{III}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle)$$

Von Neumann Entropy




Which one is more entangled? (singular values could help):

$$|\psi_{AB}^I\rangle = \frac{1}{\sqrt{4}} (|00\rangle + |10\rangle - |01\rangle - |11\rangle), \quad \lambda_{1,2}^I = \{1, 0\}$$

$$|\psi_{AB}^{II}\rangle = \frac{1}{\sqrt{3}} (|00\rangle + |01\rangle + |10\rangle), \quad \lambda_{1,2}^{II} = \{0.87, 0.13\}$$

$$|\psi_{AB}^{III}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle), \quad \lambda_{1,2}^{III} = \{0.5, 0.5\}$$

Von Neumann Entropy



Which one is more entangled? (singular values could help):

$$|\psi_{AB}^I\rangle = \frac{1}{\sqrt{4}} (|00\rangle + |10\rangle - |01\rangle - |11\rangle), \quad \lambda_{1,2}^I = \{1, 0\}$$


$$|\psi_{AB}^{II}\rangle = \frac{1}{\sqrt{3}} (|00\rangle + |01\rangle + |10\rangle), \quad \lambda_{1,2}^{II} = \{0.87, 0.13\}$$

$$|\psi_{AB}^{III}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle), \quad \lambda_{1,2}^{III} = \{0.5, 0.5\}$$

Measure of entanglement is fixed, if it has

- invariance under LUO $\Rightarrow E = E(\lambda)$
- continuity
- additivity $E(|\psi_{AB}\rangle \otimes |\varphi_{AB}\rangle) = E(|\psi_{AB}\rangle) + E(|\varphi_{AB}\rangle)$

Von Neumann Entropy



Which one is more entangled? (singular values could help):

$$|\psi_{AB}^I\rangle = \frac{1}{\sqrt{4}} (|00\rangle + |10\rangle - |01\rangle - |11\rangle), \quad \lambda_{1,2}^I = \{1, 0\}$$

$$|\psi_{AB}^{II}\rangle = \frac{1}{\sqrt{3}} (|00\rangle + |01\rangle + |10\rangle), \quad \lambda_{1,2}^{II} = \{0.87, 0.13\}$$

$$|\psi_{AB}^{III}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle), \quad \lambda_{1,2}^{III} = \{0.5, 0.5\}$$

Measure of entanglement is fixed, if it has

- invariance under LUO $\Rightarrow E = E(\lambda)$
- continuity
- additivity $E(|\psi_{AB}\rangle \otimes |\varphi_{AB}\rangle) = E(|\psi_{AB}\rangle) + E(|\varphi_{AB}\rangle)$

This is *the von Neumann entropy*

$$S(\rho_A) = S(\rho_B) = - \sum_j \lambda_j \ln \lambda_j, \quad S^I = 0, \quad S^{II} = 0.6, \quad S^{III} = 1.$$

Paired entanglement

The purity of the state represented by *one-tangle*

$$\tau_1[\rho_A] = 4 \det \rho_A = 1 - 4 (\langle S_x \rangle^2 + \langle S_y \rangle^2 + \langle S_z \rangle^2)$$

for $\rho_A = \text{tr}_B \rho_{AB}$.

Paired entanglement

The purity of the state represented by *one-tangle*

$$\tau_1[\rho_A] = 4 \det \rho_A = 1 - 4 (\langle S_x \rangle^2 + \langle S_y \rangle^2 + \langle S_z \rangle^2)$$

for $\rho_A = \text{tr}_B \rho_{AB}$.

For a pure state of two qubits we can define *concurrence*

$$C[|\psi_{AB}\rangle] = |\langle \psi^* | \sigma_A^y \otimes \sigma_B^y | \psi \rangle| = \sqrt{\tau_1}$$

Paired entanglement

The purity of the state represented by *one-tangle*

$$\tau_1[\rho_A] = 4 \det \rho_A = 1 - 4 (\langle S_x \rangle^2 + \langle S_y \rangle^2 + \langle S_z \rangle^2)$$

for $\rho_A = \text{tr}_B \rho_{AB}$.

For a pure state of two qubits we can define *concurrence*

$$C[|\psi_{AB}\rangle] = |\langle \psi^* | \sigma_A^y \otimes \sigma_B^y | \psi \rangle| = \sqrt{\tau_1}$$

Not all entanglement is stored in pairs:

$$\text{residual tangle} = \tau_{1,i} - \sum_{j \neq i} C_{ij}^2 \geq 0$$

Entanglement in mixed states: EoF



The Entanglement of Formation defined through *convex roof*:

$$E_F(\rho_{AB}) \stackrel{\text{def}}{=} \min_{\{p_j, \psi_j\}} \sum_j p_j S(\rho_{A,j}),$$

with $\rho_{AB} = \sum_j p_j |\psi_j\rangle\langle\psi_j|$ and $\rho_{A,j} = \text{tr}_B |\psi_j\rangle\langle\psi_j|$.

Entanglement in mixed states: EoF



The Entanglement of Formation defined through *convex roof*:

$$E_F(\rho_{AB}) \stackrel{\text{def}}{=} \min_{\{p_j, \psi_j\}} \sum_j p_j S(\rho_{A,j}),$$

with $\rho_{AB} = \sum_j p_j |\psi_j\rangle\langle\psi_j|$ and $\rho_{A,j} = \text{tr}_B |\psi_j\rangle\langle\psi_j|$.

For two qubits:

$$E_F(\rho) = - \sum_{\sigma=\pm} \frac{1}{2} \sqrt{1 + \sigma C^2(\rho)} \ln \frac{1}{2} \sqrt{1 + \sigma C^2(\rho)},$$

$C(\rho) \in [0, 1]$ – *concurrence*, that can be calculated from
 $R = \sqrt{\rho} \tilde{\rho} \sqrt{\rho} = \sqrt{\rho} (\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y)$ with $\lambda_1^2 \geq \dots \geq \lambda_4^2$

$$C = \max(\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0).$$

Concurrence through two-point spin correlation

In a spin-1/2 chain

$$C_{ij} = 2 \max\{0, C_{ij}^I, C_{ij}^{II}\},$$


where

$$C_{ij}^I = |g_{ij}^{xx} + g_{ij}^{yy}| - \sqrt{(1/4 + g_{ij}^{zz})^2 - M_z^2}$$
$$C_{ij}^{II} = |g_{ij}^{xx} - g_{ij}^{yy}| + g_{ij}^{zz} - 1/4,$$

with $g_{ij}^{\alpha\alpha} = \langle S_i^\alpha S_j^\alpha \rangle$ and $M_z = \langle S^z \rangle$.

Measurable example

Consider dimer system



With Hamiltonian

$$\hat{H} = -J \sum_{\sigma} (\hat{c}_{L\sigma} \hat{c}_{R\sigma} + \text{c.c.}) + U \sum_{j=L,R} \hat{n}_{j\downarrow} \hat{n}_{j\uparrow}$$

[2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, Nature Physics 15, no. 7 (2019)



Measurable example

$$\hat{H} = -J \sum_{\sigma} (\hat{c}_{L\sigma} \hat{c}_{R\sigma} + \text{c.c.}) + U \sum_{j=L,R} \hat{n}_{j\downarrow} \hat{n}_{j\uparrow}$$

What is available for us to measure?

$$\rho = \begin{pmatrix} \langle LL \rangle & \langle LR \rangle & \langle RL \rangle & \langle RR \rangle \\ \langle LL \rangle & \rho_{1,2} & \rho_{1,3} & \rho_{1,4} \\ \langle LR \rangle & \rho_{2,3} & \rho_{2,4} & \\ \langle RL \rangle & \rho_{3,4} & & \\ \text{h.c.} & \rho_{RL} & \rho_{3,4} & \rho_{RR} \end{pmatrix}$$

— Real-space populations
— Single-particle coherence
— Two-particle coherence



[2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, Nature Physics 15, no. 7 (2019)



Measurable example

$$\hat{H} = -J \sum_{\sigma} (\hat{c}_{L\sigma} \hat{c}_{R\sigma} + \text{c.c.}) + U \sum_{j=L,R} \hat{n}_{j\downarrow} \hat{n}_{j\uparrow}$$

What is available for us to measure?

$$\rho = \begin{pmatrix} \langle LL | & \langle LR | & \langle RL | & \langle RR | \\ \hline P_{LL} & \rho_{1,2} & \rho_{1,3} & \rho_{1,4} \\ P_{LR} & \rho_{2,3} & \rho_{2,4} & \rho_{3,4} \\ \hline P_{RL} & \rho_{3,4} & \rho_{2,3} & \rho_{1,2} \\ \hline \text{h.c.} & P_{RR} & P_{RL} & P_{LL} \end{pmatrix}$$

— Real-space populations
— Single-particle coherence
— Two-particle coherence



Lower bound for concurrence

$$C(\rho) \geq \max \left\{ 0, 2(|\rho_{1,4}| - \sqrt{P_{LR}P_{RL}}), 2(|\rho_{2,3}| - \sqrt{P_{LL}P_{RR}}) \right\}$$

[2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, Nature Physics 15, no. 7 (2019)

[3] M. Jafarpour et al., A Useful Strong Lower Bound on Two-Qubit Concurrence, Quantum Information Processing 11, no. 6 (2012)



Measurable example

$$\hat{H} = -J \sum_{\sigma} (\hat{c}_{L\sigma} \hat{c}_{R\sigma} + \text{c.c.}) + U \sum_{j=L,R} \hat{n}_{j\downarrow} \hat{n}_{j\uparrow}$$

What is available for us to measure?

$$\rho = \begin{pmatrix} \text{LL} & \text{LR} & \text{RL} & \text{RR} \\ \begin{pmatrix} P_{LL} & \rho_{1,2} & \rho_{1,3} & \rho_{1,4} \\ P_{LR} & \rho_{2,3} & \rho_{2,4} \\ P_{RL} & \rho_{3,4} \\ P_{RR} \end{pmatrix} \\ \text{h.c.} \end{pmatrix}$$

— Real-space populations
— Single-particle coherence
— Two-particle coherence




Lower bound for concurrence

$$C(\rho) \geq \max \left\{ 0, 2(|\rho_{1,4}| - \sqrt{P_{LR}P_{RL}}), 2(|\rho_{2,3}| - \sqrt{P_{LL}P_{RR}}) \right\}$$

[2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, *Nature Physics* 15, no. 7 (2019)


[3] M. Jafarpour et al., A Useful Strong Lower Bound on Two-Qubit Concurrence, *Quantum Information Processing* 11, no. 6 (2012)

Relative entropy of entanglement



[1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)

Relative entropy of entanglement




Measure of entanglement: «distance» from separable states

$$E(\rho) \stackrel{\text{def}}{=} \min_{\rho' \in \mathcal{D}} S(\rho || \rho'),$$

with $S(\rho || \rho') = \text{tr } \rho (\ln \rho - \ln \rho')$.

Relative entropy of entanglement




Measure of entanglement: «distance» from separable states

$$E(\rho) \stackrel{\text{def}}{=} \min_{\rho' \in \mathcal{D}} S(\rho || \rho'),$$

with $S(\rho || \rho') = \text{tr } \rho (\ln \rho - \ln \rho')$.

*reduces to $S(\rho)$ in the case of pure bi-partite states.


Multipartite entanglement measures



Multipartite entanglement: «distance» from product states

$$E_g(\Psi) \stackrel{\text{def}}{=} -\log_2 \max_{\Phi \in \mathcal{S}} |\langle \Psi | \Phi \rangle|^2,$$

Multipartite entanglement measures



Multipartite entanglement: «distance» from product states

$$E_g(\Psi) \stackrel{\text{def}}{=} -\log_2 \max_{\Phi \in \mathcal{S}} |\langle \Psi | \Phi \rangle|^2,$$

Or average purity

$$E_{\text{gl}} \stackrel{\text{def}}{=} 2 - \frac{2}{N} \sum_{j=1}^N \text{tr} \rho_j^2$$

Indistinguishable particles: example

Consider state represented by the Slater determinant

$$|\psi\rangle^{\text{SD}} = \frac{1}{\sqrt{2}} (|A, 0\rangle |B, 1\rangle - |B, 1\rangle |A, 0\rangle),$$

with A, B and $0, 1$ are the spatial and the internal degree of freedom.

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
- [4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)

Indistinguishable particles: example

Consider state represented by the Slater determinant

$$|\psi\rangle^{\text{SD}} = \frac{1}{\sqrt{2}} (|A, 0\rangle |B, 1\rangle - |B, 1\rangle |A, 0\rangle),$$

with A, B and $0, 1$ are the spatial and the internal degree of freedom.

This state results from antisymmetrizing the product state

$$|\psi\rangle^{\text{prod}} = |A, 0\rangle |B, 1\rangle,$$

so no entanglement actually.

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
[4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)

Indistinguishable particles: example

Consider state represented by the Slater determinant

$$|\psi\rangle^{\text{SD}} = \frac{1}{\sqrt{2}} (|A, 0\rangle |B, 1\rangle - |B, 1\rangle |A, 0\rangle),$$

with A, B and $0, 1$ are the spatial and the internal degree of freedom.

Meanwhile reduced density matrix is impure

$$\rho_{\text{f}}^{\text{SD}} = \frac{1}{2}|A, 0\rangle\langle A, 0| + \frac{1}{2}|B, 1\rangle\langle B, 1|.$$

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
[4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)

Indistinguishable particles: fermionic exchange correlations

In general for N identical fermions identified with a single Slater determinant reduced density matrix of M fermions

$$\mathrm{tr} (\rho_M^{\mathrm{SD}})^2 = \binom{N}{M}^{-1}.$$

Such correlations are compatible with the possibility of assigning a complete set of properties to the individual fermions.

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
- [4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)

Indistinguishable particles: fermionic exchange correlations

State that can not be represented by the Slater determinant

$$|\psi\rangle^{\text{SD}} = \frac{1}{2} (|A, 0\rangle |B, 1\rangle - |B, 1\rangle |A, 0\rangle + |A, 1\rangle |B, 0\rangle - |B, 0\rangle |A, 1\rangle),$$
$$|\psi\rangle^{\text{non-prod}} = \frac{1}{\sqrt{2}} (|A, 0\rangle |B, 1\rangle + |A, 1\rangle |B, 0\rangle)$$

And the purity of $\rho_f^{\text{non-SD}}$ is lower than the purity of ρ_f^{SD} .

This feature reveals the presence of correlations beyond exchange correlations.

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
[4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)

Indistinguishable particles: fermionic exchange correlations

State that can not be represented by the Slater determinant

$$|\psi\rangle^{\text{SD}} = \frac{1}{2} (|A, 0\rangle |B, 1\rangle - |B, 1\rangle |A, 0\rangle + |A, 1\rangle |B, 0\rangle - |B, 0\rangle |A, 1\rangle),$$
$$|\psi\rangle^{\text{non-prod}} = \frac{1}{\sqrt{2}} (|A, 0\rangle |B, 1\rangle + |A, 1\rangle |B, 0\rangle)$$

And the purity of $\rho_{\text{f}}^{\text{non-SD}}$ is lower than the purity of $\rho_{\text{f}}^{\text{SD}}$.

Entanglement criteria:


$$\text{tr } \rho_M^2 < \binom{N}{M}^{-1} \quad \Leftrightarrow \quad |\psi\rangle \text{ is entangled.}$$

In other words: how many Slater determinants you need to describe the state (*Slater rank*)?

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
- [4] A. P. Majtey et al., Indistinguishable Entangled Fermions: Basics and Future Challenges, Phil. Trans. R. Soc. A.381 (2023)


Entanglement witnesses

Some property, that differs for separable and entangled states.



Entanglement witnesses

Some property, that differs for separable and entangled states.




A state ρ_{AB} is entangled if and only if a positive map Λ exists:

$$(\mathbb{1}_A \otimes \Lambda_B) \rho_{AB} < 0.$$

Entanglement witnesses

Some property, that differs for separable and entangled states.



A state ρ_{AB} is entangled if and only if a positive map Λ exists:

$$(\mathbb{1}_A \otimes \Lambda_B) \rho_{AB} < 0.$$

Peres-Horodecki criterion of being separable:

$$\rho_{AB}^{T_B} \geq 0.$$

Entanglement witnesses: Peres-Horodecki criterion

Peres-Horodecki criterion of being separable (sufficient):

$$\rho_{AB}^{T_B} \geq 0.$$

Necessary for:

- two qubits
- two harmonic oscillator modes

$$\rho = \frac{1}{n} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & b & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & c & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{c} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{b} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{a} & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Entanglement witnesses: Peres-Horodecki criterion

Peres-Horodecki criterion of being separable (sufficient):

$$\rho_{AB}^{T_B} \geq 0.$$

Necessary for:

- two qubits
- two harmonic oscillator modes

$$\rho = \frac{1}{n} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & b & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & c & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{c} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{b} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{a} & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$


Measure of entanglement: *the logarithmic negativity*

$$E_N = \log_2(2N_{AB} + 1),$$

with N_{AB} is the absolute sum of the negative eigenvalues of $\rho_{AB}^{T_B}$.

Entanglement witnesses: example

We can measure correlations:




Construct the pair correlators

$$\xi(d = k_1 - k_2) = \frac{\int dk \langle n_\uparrow(k - d/2) n_\downarrow(k + d/2) \rangle}{\int dk \langle n_\uparrow(k - d/2) \rangle \langle n_\downarrow(k + d/2) \rangle}$$
$$\chi(s = k_1 + k_2) = \frac{\int dk \langle n_\uparrow(k + s/2) n_\downarrow(-k + s/2) \rangle}{\int dk \langle n_\uparrow(k + s/2) \rangle \langle n_\downarrow(-k + s/2) \rangle}$$

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
[2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, Nature Physics 15, no. 7 (2019)

Entanglement witnesses: example

The pair correlators as entanglement witnesses:



For separable states:

$$\xi_{\min} \leq \xi \leq \xi_{\max}, \quad \chi_{\min} \leq \chi \leq \chi_{\max}.$$

- [1] L. Amico et al., Entanglement in Many-Body Systems, Reviews of Modern Physics 80, no. 2 (2008)
- [2] A. Bergschneider et al., Experimental Characterization of Two-Particle Entanglement through Position and Momentum Correlations, Nature Physics 15, no. 7 (2019)



Random Unitaries

The second Rényi entropy

$$S_2(\rho_A) = -\log_2 \text{tr} \rho_A^2.$$

Bipartite entanglement exists between subsystems A and B of \mathcal{S} with reduced density matrices $\rho_A = \text{tr}_{\mathcal{S} \setminus A} \rho$ and $\rho_B = \text{tr}_{\mathcal{S} \setminus B} \rho$ **if**

$$S_2(\rho_A) > S_2(\rho_{AB}) \quad \text{or} \quad S_2(\rho_B) > S_2(\rho_{AB}).$$



[5] A. Elben et al., Statistical Correlations between Locally Randomized Measurements, Physical Review A 99, no. 5 (2019)

Random Unitaries


The second Rényi entropy

$$S_2(\rho_A) = -\log_2 \text{tr} \rho_A^2.$$

Bipartite entanglement exists between subsystems A and B of \mathcal{S} with reduced density matrices $\rho_A = \text{tr}_{\mathcal{S} \setminus A} \rho$ and $\rho_B = \text{tr}_{\mathcal{S} \setminus B} \rho$ if

$$S_2(\rho_A) > S_2(\rho_{AB}) \quad \text{or} \quad S_2(\rho_B) > S_2(\rho_{AB}).$$

Thank you for your attention!



[5] A. Elben et al., Statistical Correlations between Locally Randomized Measurements, Physical Review A 99, no. 5 (2019)