Coursera Page 1 of 6

Feedback — Quiz 2

Help Center

Thank you. Your submission for this quiz was received.

You submitted this quiz on Fri 13 Nov 2015 5:50 PM EST. You got a score of 8.00 out of 8.00.

Question 1

What is the variance of the distribution of the average an IID draw of n observations from a population with mean μ and variance σ^2 .

Your Answer		Score	Explanation
$\bigcirc \ 2\sigma/\sqrt{n}$			
$\bigcirc \sigma^2$			
$\odot \frac{\sigma^2}{n}$	~	1.00	
$\bigcirc \sigma/n$			
Total		1.00 / 1.00	

Question Explanation

$$\mathrm{Var}(ar{X}) = \sigma^2/n$$

Question 2

Suppose that diastolic blood pressures (DBPs) for men aged 35-44 are normally distributed with a mean of 80 (mm Hg) and a standard deviation of 10. About what is the probability that a random 35-44 year old has a DBP less than 70?

Your Answer	Score	Explanation	

Coursera Page 2 of 6



Question 3

Brain volume for adult women is normally distributed with a mean of about 1,100 cc for women with a standard deviation of 75 cc. What brain volume represents the 95th percentile?

Your Answer	Sco	ore	Explanation
O approximately 1247			
approximately 1223	✓ 1.00	0	
O approximately 1175			
○ approximately 977			
Total	1.00	0 / 1.00	

Question Explanation

qnorm(0.95, mean = 1100, sd = 75)

[1] 1223

Coursera Page 3 of 6

Question 4

Refer to the previous question. Brain volume for adult women is about 1,100 cc for women with a standard deviation of 75 cc. Consider the sample mean of 100 random adult women from this population. What is the 95th percentile of the distribution of that sample mean?

		Explanation
approximately 1112 cc	✓ 1.00	
approximately 1115 cc		
approximately 1088 cc		
) approximately 1110 cc		
otal	1.00 / 1.00	

Question 5

You flip a fair coin 5 times, about what's the probability of getting 4 or 5 heads?

Your Answer		Score	Explanation
○ 6%			
○ 12%			
○ 3%			
19%	~	1.00	
Total		1.00 / 1.00	

Coursera Page 4 of 6

Question Explanation

$${5 \choose 4} 2^{-5} + {5 \choose 5} 2^{-5} pprox 19\%$$

pbinom(3, size = 5, prob = 0.5, lower.tail = FALSE)

[1] 0.1875

Question 6

The respiratory disturbance index (RDI), a measure of sleep disturbance, for a specific population has a mean of 15 (sleep events per hour) and a standard deviation of 10. They are not normally distributed. Give your best estimate of the probability that a sample mean RDI of 100 people is between 14 and 16 events per hour?

Your Answer		Score	Explanation
O 95%			
68%	~	1.00	
○ 34%			
O 47.5%			
Total		1.00 / 1.00	

Question Explanation

The standard error of the mean is $10/\sqrt{100}=1$. Thus between 14 and 16 is with one standard deviation of the mean of the distribution of the sample mean. Thus it should be about 68%.

pnorm(16, mean = 15, sd = 1) - pnorm(14, mean = 15, sd = 1)

[1] 0.6827

Coursera Page 5 of 6

Question 7

Consider a standard uniform density. The mean for this density is .5 and the variance is 1 / 12. You sample 1,000 observations from this distribution and take the sample mean, what value would you expect it to be near?

Your Answer		Score	Explanation
⊚ 0.5	~	1.00	
○ 0.25			
○ 0.75			
○ 0.10			
Total		1.00 / 1.00	

Question Explanation

Via the LLN it should be near .5.

Question 8

The number of people showing up at a bus stop is assumed to be Poisson with a mean of 5 people per hour. You watch the bus stop for 3 hours. About what's the probability of viewing 10 or fewer people?

Your Answer		Score	Explanation
⊚ 0.12	~	1.00	
○ 0.06			
○ 0.08			
○ 0.03			
Total		1.00 / 1.00	

Question Explanation

Coursera Page 6 of 6

```
ppois(10, lambda = 15)

## [1] 0.1185
```