

## Useful formulas

$\binom{n}{k} = \frac{n!}{k!(n-k)!}$  — number of ways to choose  $k$  objects out of  $n$

$\binom{n+k-1}{k-1}$  — number of ways to choose  $k$  objects out of  $n$  with repetitions

$\left\lfloor \frac{n}{m} \right\rfloor$  — Stirling numbers of the first kind; number of permutations of  $n$  elements with  $k$  cycles

$$\left[ \begin{matrix} n+1 \\ m \end{matrix} \right] = n \left[ \begin{matrix} n \\ m \end{matrix} \right] + \left[ \begin{matrix} n \\ m-1 \end{matrix} \right]$$

$$(x)_n = x(x-1) \cdots x-n+1 = \sum_{k=0}^n (-1)^{n-k} \left[ \begin{matrix} n \\ k \end{matrix} \right] x^k$$

$\left\{ \begin{matrix} n \\ m \end{matrix} \right\}$  — Stirling numbers of the second kind; number of partitions of set  $1, \dots, n$  into  $k$  disjoint subsets.

$$\left\{ \begin{matrix} n+1 \\ m \end{matrix} \right\} = k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} + \left\{ \begin{matrix} n \\ k-1 \end{matrix} \right\}$$

$$\sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k = x^n$$

$$C_n = \frac{1}{n+1} \binom{2n}{n} - \text{Catalan numbers}$$

$$C(x) = \frac{1 - \sqrt{1-4x}}{2x}$$

## Binomial transform

If  $a_n = \sum_{k=0}^n \binom{n}{k} b_k$ , then  $b_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k$

- $a = (1, x, x^2, \dots), b = (1, (x+1), (x+1)^2, \dots)$
- $a_i = i^k, b_i = \left\{ \begin{matrix} n \\ i \end{matrix} \right\} i!$

## Burnside's lemma

Let  $G$  be a group of *action* on set  $X$  (Ex.: cyclic shifts of array, rotations and symmetries of  $n \times n$  matrix, ...)

Call two objects  $x$  and  $y$  *equivalent* if there is an action  $f$  that transforms  $x$  to  $y$ :  $f(x) = y$ .

The number of equivalence classes then can be calculated as follows:  $C = \frac{1}{|G|} \sum_{f \in G} |X^f|$ , where  $X^f$

is the set of *fixed points* of  $f$ :  $X^f = \{x | f(x) = x\}$

## Generating functions

Ordinary generating function (o.g.f.) for sequence

$$a_0, a_1, \dots, a_n, \dots \text{ is } A(x) = \sum_{n=0}^{\infty} a_n x^n$$

Exponential generating function (e.g.f.) for

$$\text{sequence } a_0, a_1, \dots, a_n, \dots \text{ is } A(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$B(x) = A'(x), b_{n-1} = n \cdot a_n$$

$$c_n = \sum_{k=0}^n a_k b_{n-k} \text{ (o.g.f. convolution)}$$

$c_n = \sum_{k=0}^n \binom{n}{k} a_k b_{n-k}$  (e.g.f. convolution, compute with FFT using  $\widetilde{a_n} = \frac{a_n}{n!}$ )

## General linear recurrences

If  $a_n = \sum_{k=1}^n b_k a_{n-k}$ , then  $A(x) = \frac{a_0}{1-B(x)}$ . We also can compute all  $a_n$  with Divide-and-Conquer algorithm in  $O(n \log^2 n)$ .

## Inverse polynomial modulo $x^l$

Given  $A(x)$ , find  $B(x)$  such that  $A(x)B(x) = 1 + x^l \cdot Q(x)$  for some  $Q(x)$

1. Start with  $B_0(x) = \frac{1}{a_0}$
2. Double the length of  $B_{k+1}(x) = (-B_k(x)^2 A(x) + 2B_k(x)) \bmod x^{2^{k+1}}$

## Fast subset convolution

Given array  $a_i$  of size  $2^k$ , calculate  $b_i = \sum_{j \& i = i} b_j$

```
for b = 0..k-1
  for i = 0..2^k-1
    if (i & (1 << b)) != 0:
      a[i + (1 << b)] += a[i]
```

## Hadamard transform

Treat array  $a$  of size  $2^k$  as  $k$ -dimensional array of size  $2 \times 2 \times \dots \times 2$ , calculate FFT of that array:

```
for b = 0..k-1
  for i = 0..2^k-1
    if (i & (1 << b)) != 0:
      u = a[i], v = a[i + (1 << b)]
      a[i] = u + v
      a[i + (1 << b)] = u - v
```