

Code No: 124DD**JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD****B.Tech II Year II Semester Examinations, May - 2017****MATHEMATICS – II****(Common to ME, MCT, MIE, MSNT)****Time: 3 Hours****Max. Marks: 75****Note:** This question paper contains two parts A and B.

Part A is compulsory which carries 25 marks. Answer all questions in Part A.

Part B consists of 5 Units. Answer any one full question from each unit. Each question carries 10 marks and may have a, b, c as sub questions.

PART- A**(25 Marks)**1.a) Show that $\nabla r^n = nr^{n-2}\vec{r}$, where $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and $r = |\vec{r}|$. [2]b) Find the values of a, b, c so that $\vec{F} = (x + 2y + az)\hat{i} + (bx - 3y - z)\hat{j} + (4x + cy + 2z)\hat{k}$ is irrotational. [3]

c) What are Dirichlet's conditions for the existence of Fourier series? [2]

d) Find the Fourier transform of $f(x) = e^{-|x|}$. [3]

e) Construct the forward difference table from the following data: [2]

x:	0	10	20	30
y:	0	0.174	0.347	0.518

f) Obtain the normal equations for fitting a straight line $y = ax + b$ to the data (x_i, y_i) , $i = 1, 2, \dots, n$. [3]g) If the first two approximations x_3 and x_4 for the root of $x^3 - 3x - 4 = 0$ are 2.125 and -3 respectively, find x_5 by the method of false position. [2]h) Find the LU decomposition for the matrix $A = \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}$. [3]i) Approximate $\int_0^{\pi} \sin x \, dx$ using the 2-point Gauss-Legendre formula. [2]j) Evaluate $\int_0^1 \frac{dx}{x}$ using Simpson's $\frac{1}{3}$ rule with $h = \frac{1}{4}$. [3]**PART-B****(50 Marks)**2.a) Find the values of a and b so that the surfaces $ax^2 - byz = (a+2)x$ and $4x^2y + z^3 = 4$ intersect orthogonally at the point $(1, -1, 2)$.b) Prove that $\nabla \times (\nabla \times \vec{F}) = \nabla(\nabla \cdot \vec{F}) - \nabla^2 \vec{F}$. [5+5]

- 3.a) Find the work done by the force $\vec{F} = (3x^2 - 6yz)\hat{i} + (2y + 3xz)\hat{j} + (1 - 4xyz^2)\hat{k}$ in moving a particle from (0, 0, 0) to (1, 1, 1) along the curve $C: x = t, y = t^2, z = t^3$.
- b) Use Green's theorem to evaluate $\oint_c (2xy - x^2)dx + (x^2 + y^2)dy$, where c is the boundary of the region enclosed by $y = x^2$ and $y^2 = x$. [5+5]

4. Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 2+x, & -2 \leq x \leq 0 \\ 2-x, & 0 < x \leq 2 \end{cases}, f(x+4) = f(x). \text{ Hence show that } \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}.$$

[10]

OR

- 5.a) Find the Fourier integral representation of $f(x) = \begin{cases} x, & |x| < 1 \\ 0, & |x| > 1 \end{cases}$.
- b) Find the inverse Fourier sine transform of $\frac{e^{-as}}{s}$. [5+5]
- 6.a) If $y_{20} = 24, y_{24} = 32, y_{28} = 35, y_{32} = 40$, find y_{25} using Gauss forward difference formula.
- b) Use Lagrange's interpolation formula to find a polynomial of least degree which suits the following data: [5+5]

x:	0	1	3	4
y:	5	6	50	105

OR

- 7.a) Fit a polynomial of second degree to the following data by the method of least squares:

x:	0	1	2
y:	1	6	17

- b) Fit a curve of the form $y = ae^{bx}$ for the following data: [5+5]

x:	1	2	3	4
y:	1.65	2.70	4.50	7.35

- 8.a) Find a root of the equation $e^x - x = 2$ using bisection method correct to 3 decimal Places.
- b) Compute $\sqrt{10}$ using Newton-Raphson method correct to 3 decimal places. [5+5]

OR

9. Solve the system of equations $10x + y + z = 12, 2x + 10y + z = 13, 2x + 2y + 10z = 14$ by Jacobi's iteration method and Gauss-Seidel iteration method. [10]

10.a) Evaluate $\int_0^1 \frac{dx}{1+x^2}$ using Trapezoidal rule with $h = \frac{1}{6}$.

b) Apply shooting method to solve the boundary value problem

$$y'' - 6y^2 = 0, y(0) = 1, y(0.5) = 0.44.$$

[5+5]

OR

11.a) Given that $\frac{dy}{dx} = 2 + \sqrt{xy}$, $y(1) = 1$. Find approximate value of y at $x = 2$ using Euler's modified method.

b) Find the largest eigen vector and the corresponding Eigen value of the matrix

$$A = \begin{pmatrix} 1 & 3 & -1 \\ 3 & 2 & 4 \\ -1 & 4 & 10 \end{pmatrix} \text{ by power method.}$$

[5+5]

---ooOoo---