

Zbrani zapiski za 1. letnik magistrskega študija

Patrik Žnidaršič

29. januar 2025

Opomba k notaciji: Včasih je produkt vektorja s skalarjem napisan kot $\lambda.x$, s piko. Tudi oprator $\vec{\nabla}$ včasih obravnavamo kot vektor, in pišemo gradient s piko spodaj, $\vec{\nabla}.f$. Divergenco označimo z $\vec{\nabla} \cdot f$.

Kazalo

1	Moderna fizika	5
1.1	Special relativity	6
1.1.1	The Michelson-Morley Experiment	6
1.1.2	Lorentz, the new Galileo	7
1.1.3	4-vector notation	9
2	Teorija grafov	11
2.1	Matchings	12
2.1.1	Tutte's theorem	15
2.1.2	Factors	16
2.2	Connectivity	17
2.2.1	Ear decomposition of a graph	22
2.2.2	x, y -cuts	23
2.3	Coloring	26
2.3.1	Mycielski's construction	27
2.3.2	Turán's theorem	28
2.3.3	Chordal graphs	29
2.3.4	Perfect graphs	30
2.3.5	Gallai-Roy-Vitaver theorem	32
2.4	Planar graphs	32
3	Teorija izračunljivosti	41
3.1	Introduction	42
3.1.1	Models of computation	46
3.2	Computability of natural numbers	46
3.3	Computable and computably enumerable sets	50
3.3.1	Varieties of non-computable sets	54
3.4	Computation with continuous data	56
3.4.1	Topological aspects of computing with ω -words	57
3.4.2	Computing with real numbers	59
3.5	Algorithmic information theory	60
3.6	Algorithmic randomness	62
4	Uvod v funkcionalno analizo	65
4.1	Normirani in Banachovi prostori	66
4.1.1	Napolnitve normiranih prostorov	67

4.1.2	Osnovne konstrukcije	68
4.2	Linearni funkcionali	70
4.2.1	Banachov izrek	72
4.2.2	Adjungirani operator in drugi dual	76
4.3	Temeljni izreki funkcionalne analize	77
4.4	Hilbertovi prostori	81
4.4.1	Ortonormirani sistemi	86
4.4.2	Stone-Weierstrassov izrek	91
4.5	Omejeni operatorji med Hilbertovimi prostori	93
4.5.1	Idempotenti in invariantni podprostori	99
4.5.2	Kompaktni operatorji	100
4.5.3	Izrek Arzela-Ascoli	102
4.5.4	Kompaktnost adjungiranega operatorja	104
4.6	Spektralna teorija	105
4.6.1	Spekter kompaktnega operatorja	110
5	Statistika 2	115
5.1	Ocenjevanje v linearnih modelih	116
5.1.1	Ocenjevanje v normalnem linearnem regresijskem modelu	117
5.2	Ocenjevanje za velike vzorce	118
5.2.1	Doslednost	121
5.2.2	Pristranske cenilke	122
5.2.3	Asimptotična normalnost	123
5.2.4	Konstrukcija cenilk	124
5.3	Preizkušanje domnev	125
5.3.1	Preizkušanje na podlagi razmerja verjetij	126

1 Moderna fizika

1.1 Special relativity


1.1.1 The Michelson-Morley Experiment

Suppose there is some fluid medium at rest in a frame S . Now, we create a plane wave in this medium, traveling in the \mathbf{e}_x direction, with a velocity of c_1 as measured in S . If we switch to a different frame S' with a Galilean transformation $\mathbf{x}' = \mathbf{x} + vt\mathbf{e}_x$, then we expect the plane wave to have a different velocity $c_1 - v$ in S' . This is true for waves in water, or sound waves in air, but, as famously measured by Michelson and Morley, it is not true for light. Their experiment, described below, showed that the speed of light c is the same in any inertial reference frame — a result that contradicts entirely with classical mechanics.

The Michelson-Morley experiment splits a beam of light into two beams traveling orthogonally to each other, which then bounce off a pair of mirrors, before being recombined and the interference pattern of this recombination being measured. Assume that S is the rest frame of the ether, in which light waves are traveling at c . Suppose that we now switch to a frame S' , with a relative velocity of v as measured in S . If one of the arms is parallel to the relative velocity vector, then the time of flight of a wave of light should be

$$t_1 = \frac{L}{c+v} + \frac{L}{c-v} = \frac{2cL}{c^2 - v^2},$$

where L is the distance between the beam splitter and the mirror. In S , the other beam of light travels a path that is not parallel to \mathbf{e}_x , as shown in figure 1.1.



Slika 1.1: Beam path in S

From Pythagoras' theorem, the length of the beam path is

$$ct_2 = 2\sqrt{L^2 + v^2 t_2^2 / 4}$$

so

$$t_2 = \frac{2L}{\sqrt{c^2 - v^2}}.$$

As mentioned, Michelson and Morley conducted this experiment, while using the Earth's rotation to perform many measurements in many directions. They found that their

experiment always gave the same result, no matter the relative velocity of the second frame and no matter the orientation of the testbed. This demonstrated beyond any doubt that Max Planck's advisor Philipp von Jolly was very wrong when he stated to Planck in 1874 that there was essentially nothing left to discover in theoretical physics,¹ since, as Feynman stated,

„It doesn't matter how beautiful your theory is, it doesn't matter how smart you are. If it doesn't agree with experiment, it's wrong.“

Classical mechanics and electromagnetism, as complete as they might seem, are wrong.

1.1.2 Lorentz, the new Galileo

Enter Einstein, who formulated special relativity with the following postulates:

1. The principle of relativity: The laws of physics are identical in all inertial frames. This is already a postulate of classical mechanics, but we now have an additional one.
2. The speed of light in vacuum is a law of physics, so it is also the same in all inertial reference frames.

Using these notions, we can find the expressions for the Lorentz transformations, the replacement of Galilean transformations in classical mechanics. Suppose we have two reference frames, one stationary and one moving with velocity $v\mathbf{e}_x$ relative to the first frame. Let $x' = f(x, t)$ be the position of an event in frame S' and $t' = g(x, t)$ be the time of that event, as measured by a clock in S' . Without loss of generality, we can limit our consideration to the case where $x = x' = 0$ at times $t = t' = 0$.

With some inspiration from dimensional analysis, we set $x' = \gamma(v)(x - vt)$ for some function γ . By the principle of relativity, rotating the system should have no effect on the value of γ , but if we rotate frame S' around, the same expression gives $x' = \gamma(-v)(x + vt)$. So γ must be an even function of v .

Now consider a light wave in S , moving with speed c towards the right. Then $x = ct$, and by Einstein's second principle, $x' = ct'$. The above transformation gives us

$$ct' = \gamma(v)(ct - vt).$$

The same transformation formula should also work in reverse, $ct = \gamma(v)(ct' + vt')$. Solving these two equations gives us

$$\gamma(v) = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

¹Source: <https://hsm.stackexchange.com/questions/2129/>

1 Moderna fizika

We still need to find $t'(x, t)$. For this, again solve the equations $x' = \gamma(x - vt)$ and $x = \gamma(x' + vt')$, but not for light. We get

$$t' = \gamma\left(t - \frac{v}{c^2}x\right),$$

which is the second part of a Lorentz transformation.

Remark. Note that if v is small compared to c , then $\gamma \approx 1$, so we recover Galilean transformations.

Remark. The transformation of time can be tricky to understand. Think about it this way: Time is what is measured by a clock. While in classical mechanics, every pair of clocks which ever agreed will always agree, in special relativity, this only holds for pairs of clocks which are stationary relative to each other (i.e. there exists an inertial reference frame S in which both clocks are stationary). Since the position of a clock doesn't matter, we can think of every reference frame as having a clock at its center, which measures the frame's time.

Definition 1.1.1. An EVENT is a point $(t, \mathbf{x}) \in \mathbb{R}^{1+3}$.

Definition 1.1.2. A WORLDLINE is a function $(t, \mathbf{x}) : \mathbb{R} \rightarrow \mathbb{R}^{1+3}$, for which $(t(\lambda), \mathbf{x}(\lambda))$ is an event for any $\lambda \in \mathbb{R}$. We call λ the AFFINE PARAMETER.

Example (time dilation). Let S be a rest frame and S' another frame moving with velocity v relative to S . Consider the time measured by a clock at $x' = 0$. Since $x = vt$, the Lorentz transformation of time gives us

$$t' = \gamma(v) \left(1 - \frac{v^2}{c^2}\right) t = \frac{1}{\gamma} t.$$

Note that $\gamma > 1$, so in the moving system, the clock shows a lower time than in the non-moving system.

Example (length contraction). Consider a stick of length l' in S' . We consider the ends of the stick to be two worldlines with $x'_1 = 0$ and $x'_2 = l'$. We wish to measure the length of the stick in the non-moving frame S . For this, we need to find two simultaneous events in S , one belonging to the first worldline and one belonging to the second.

We'll take the measurement at $t = 0$. From the Lorentz transformation of space, we then get

$$l' = \gamma l,$$


so the stick is shorter in S than in S' .

Note that the expression $s^2 = c^2 t^2 - \mathbf{x}^2$ is invariant under Lorentz transformations, as is the difference $\Delta s^2 = c^2 \Delta t^2 - \Delta \mathbf{x}^2$ between two events. This allows us to define Δs^2 as a sensible distance-like measure on \mathbb{R}^{1+3} , called the MINKOWSKI METRIC, though it is not a metric. Depending on the value of Δs^2 , we call a pair of events

- TIMELIKE if $\Delta s^2 > 0$,

- NULL or LIGHTLIKE if $\Delta s^2 = 0$, and
- SPACELIKE if $\Delta s^2 < 0$.

Lightlike events are those which can be connected with a light-speed signal. Two timelike events can then be causally connected, as there was enough time between them to transfer and process information. Spacelike events, on the other hand, cannot be causally connected, since no signal can travel faster than light, so there is no way for the information of the first event to reach the second. We can draw these events on a spacetime diagram, as shown in figure 1.2.



Slika 1.2: Spacetime diagram

1.1.3 4-vector notation

Further on, we will use 4-vectors to denote events and quantities derived from them. We will use the label X^μ to refer to either the μ -th component of the 4-vector X , or to the vector itself; it is generally defined as

$$X^\mu = \begin{bmatrix} ct \\ x \\ y \\ z \end{bmatrix}$$

for an event at time t and coordinate (x, y, z) . We also use the convention that indexing 4-vectors with Latin indices only refers to their space coordinates, ignoring the time coordinate. If we write X_μ with the index at the bottom, we mean

$$X_\mu = \eta_{\mu\nu} X^\nu$$

where the double index ν implies a summation, as per the Einstein convention. Above, $\eta_{\mu\nu}$ is the matrix

$$\eta_{\mu\nu} = \begin{bmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{bmatrix}.$$

This allows us to write the Minkowski dot product simply as $X^\mu X_\mu = X^\mu \eta_{\mu\nu} X^\nu$, so

$$ds^2 = \eta_{\mu\nu} dX^\mu dX^\nu.$$

We can now recognize Lorentz transformations as matrices $\Lambda^\mu{}_\nu$ which allow for transformations

$$X'^\mu = \Lambda^\mu{}_\nu X^\nu.$$

Not all matrices are correct here, as among other things, they need to satisfy

$$X^\rho X_\rho = X'^\mu X'_\mu$$

or

$$X^\rho \eta_{\rho\sigma} X^\sigma = X'^\mu \eta_{\mu\nu} X'^\nu = \eta_{\mu\nu} \Lambda^\mu{}_\rho X^\rho \Lambda^\nu{}_\sigma X^\sigma.$$

This gives us the constraint

$$\eta_{\rho\sigma} = \eta_{\mu\nu} \Lambda^\mu{}_\rho \Lambda^\nu{}_\sigma$$

which is what we will use for our definition of a Lorentz transformation, so a Lorentz transformation is any matrix $\Lambda^\mu{}_\nu$ which satisfies the property above.

2 Teorija grafov

2.1 Matchings

Definition 2.1.1. A vertex set $S \subseteq V$ is an **INDEPENDENT SET** of the graph $G = (V, E)$ if the induced subgraph $G[S]$ is empty. The maximum cardinality of an independent set is the **INDEPENDENCE NUMBER** $\alpha(G)$.

Definition 2.1.2. A vertex set $T \subseteq V$ is a **VERTEX COVER** if every edge has at least one of its endings in T . The maximum cardinality of a vertex cover is the **VERTEX COVER NUMBER** $\beta(G)$.

Definition 2.1.3. An edge set $M \subseteq E$ is a **MATCHING** if for every distinct $e_1, e_2 \in M$, edges e_1 and e_2 have no common ending. The maximum cardinality of a matching is the **MATCHING NUMBER** $\alpha'(G)$.

Definition 2.1.4. An edge set $C \subseteq E$ is an **EDGE COVER** if every vertex of G is covered by at least one edge from C . If the minimum degree $\delta(G)$ is at least 1, we can define the **EDGE COVER NUMBER** $\beta'(G)$ as the minimum cardinality of an edge cover.

Question 1. Define the independence number, the vertex and edge cover number, and the matching number.

Remark. The complement of an independent set is a vertex cover, so $\alpha(G) + \beta(G) = n(G)$ in every graph G . In a maximum matching, every edge must be covered by different vertices, so $\alpha'(G) \leq \beta(G)$. We can similarly argue that $\alpha'(G) \leq n(G)/2 \leq \beta'(G)$ and $\alpha(G) \leq \beta'(G)$.

Theorem 2.1.5 (Gallai). *If $\delta(G) \geq 1$, then $\alpha'(G) + \beta'(G) = n(G)$.*

Proof. Take a maximum matching M in G and let $V(M)$ be the vertices covered by M . For every vertex not covered by M , we can take an incident edge and add it to M . This gives an edge cover with

$$|M| + |\overline{V(M)}| = |M| + (n - 2|M|) = n - |M|$$

edges. Since $|M| = \alpha'(G)$, this implies $\beta'(G) + \alpha'(G) \leq n(G)$.

Now take a minimum edge cover C . We claim that for every edge in C , at least one end is covered only once by C . For suppose that $uv \in C$ is an edge and both u and v are covered by other edges in C . If we remove uv , then $C \setminus \{uv\}$ is a smaller cover, which is a contradiction.

The induced subgraph $G[C]$ is then a forest of stars. Suppose it consists of k components. We get $|C| = n - k$, since in a tree, the number of vertices is 1 more than the number of edges. A matching is obtained by choosing one edge from every star, which gives $\alpha'(G) + \beta'(G) \geq n(G)$, thus completing the proof. \square

Question 2. State and prove Gallai's theorem.

Definition 2.1.6. Let M be a matching. A path $v_1v_2 \dots v_k$ is an M -ALTERNATING PATH if the edges along the path alternate between M and \overline{M} . An M -alternating path is M -AUGMENTING if neither end of the path is covered by M .

Remark. Such a path cannot start or end with an edge from M , and the endpoints cannot be part of an edge in M .

Proposition 2.1.7. *If G is a graph, M is a matching and there exists an M -augmenting path P , then M is not a maximum matching.*

Proof. Suppose $P = v_1 \dots v_k$. We know the first and last edge are not in M , so $|E(P) \cap \overline{M}| = |E(P) \cap M| + 1$. Now let $M' = M \oplus E(P)$ be the symmetric difference of M and $E(P)$. This is clearly a matching. We know that $|M'| = |M| + 1$, so M cannot be maximum. \square

Question 3. How can you construct a larger matching from an augmenting path?

Definition 2.1.8. A KÖNIG-EGERVÁRY graph is a graph G with $\alpha'(G) = \beta(G)$.

Theorem 2.1.9 (König). *Let G be a bipartite graph. Then $\alpha'(G) = \beta(G)$. Additionally, if M is a matching in G and there is no M -augmenting path, M is a maximum matching.*

Proof. Let the partite classes of G be A and B . Suppose that M is a matching for which there is no M -augmenting path in G . Define X as the set of all vertices in A that are not covered by the matching, and Y the vertices in B not covered by M . Additionally, let B_1 be the set of vertices in B that can be reached by an M -alternating path from X , and similarly, let A_1 be the set of vertices of A which can be reached from X by an M -alternating path. Finally, define $B_2 = B \setminus (B_1 \cup Y)$ and $A_2 = A \setminus (A_1 \cup X)$.

Observe that on an M -alternating path from X , any edge from A to B is in \overline{M} and any edge from B to A is in M . Our matching provides a one-to-one mapping between A_1 and B_1 and between A_2 and B_2 , so $|A_1| = |B_1|$ and $|A_2| = |B_2|$. We also know that $|A_1| + |A_2| = |M|$. Now consider the possible edges between the defined vertex sets.

There is no edge between X and Y , since that would be a (trivial) M -augmenting path. There are also no edges between X and B_2 , since an edge xb is an M -alternating path, implying $b \in B_1$. So the only edges from X lead to B_1 .

There are also no edges between A_1 and Y , because we can construct an M -augmenting path with such an edge. If $a \in A_1$, then there is an alternating path from X to a , which we could extend with a a -to- Y edge to get an augmenting path. Finally, there are no edges between A_1 and B_2 , since that would give an alternating path from X to the B_2 -vertex as before.

Then $T = B_1 \cup A_2$ is a vertex cover with $|T| = |B_1| + |A_2| = |A_1| + |A_2| = |M|$. We have thus constructed a vertex cover with $|M|$ vertices, giving

$$\beta(G) \leq |T| = |M| \leq \alpha'(G).$$

The other inequality holds in the general case, so this completes the proof. \square

Question 4. State and prove König's theorem.

Corollary 2.1.10. *If G is a bipartite graph, then $\alpha(G) = \beta'(G)$.*

Definition 2.1.11. Let G be a bipartite graph with partite classes A and B . HALL'S CONDITION holds for the set A if for every $S \subseteq A$,

$$|S| \leq |N(S)| = \left| \bigcup_{u \in S} N(u) \right|.$$

Theorem 2.1.12 (Hall). *If G is a bipartite graph with partite classes A and B , then there exists a matching that covers A if and only if Hall's condition holds for A .*

Proof. Let M be a matching covering A and $S \subseteq A$. We can take the pairs matched by M

$$B_S = \{v \in B \mid v \text{ is covered by an edge in } M\}.$$

Clearly, $|S| = |B_S|$ and $B_S \subseteq N(S)$, so $|S| \leq |N(S)|$.

For the other implication, suppose there is no matching covering A . Divide the sets A and B as in the proof of König's theorem, using some matching M . Since the matching doesn't cover A , X is not empty. Now consider $S = A_1 \cup X$. All edges from S lead into B_1 , so $N(S) = B_1$, but $|S| = |A_1| + |X| > |B_1| = |N(S)|$. \square

Question 5. State and prove Hall's theorem.

Definition 2.1.13. A matching M is a PERFECT MATCHING if it covers all vertices.

Corollary 2.1.14. *In a bipartite graph G , there is a perfect matching if and only if $|A| = |B|$ and A satisfies Hall's condition.*

Definition 2.1.15. Let G be a bipartite graph with partite classes A, B , and $S \subseteq A$. The DEFICIENCY of S is $\text{def}(S) = |S| - |N(S)|$.

Theorem 2.1.16. *Let G be a bipartite graph with partite classes A and B . If M is a maximum matching in G , it covers*

$$\alpha'(G) = |A| - \max_{S \subseteq A} \text{def}(S)$$

vertices of A .

Theorem 2.1.17. *If G is a regular bipartite graph, then G has a perfect matching.*

Proof. The number of edges in the graph is $k \cdot |A| = k \cdot |B|$, so $|A| = |B|$. Let $S \subseteq A$. The number of edges between S and $N(S)$ is exactly $k \cdot |S|$. Every neighbour $u \in N(S)$ has exactly k neighbours, at most k are in S . So at most $k \cdot |N(S)|$ edges are between S and $N(S)$, implying $|S| \leq |N(S)|$. \square

Question 6. Show that a regular bipartite graph has a perfect matching.

Theorem 2.1.18. *Let M be a matching in G . Then there is an M -augmenting path in G if and only if M is not a maximum matching in G .*

Proof. We've already proved the right implication. Suppose there is a matching M' with $|M'| > |M|$. Consider the symmetric difference $M \Delta M'$ and denote $G' = G[M \Delta M']$. Clearly the maximum degree $\Delta(G') \leq 2$, from which we know that the components of G' are all paths or cycles.

Any cycle must alternate between edges from M and M' , so it is an even cycle, and contains the same number of edges from the two matchings. In any path of even length, there must be the same number of edges in M and in M' . And finally, in a path of odd length, one of the two sets has an extra edge compared to the other. Since $|M'| > |M|$, there must be a path with more edges in M' than in M . Label it G'_1 .

This component is an M -augmenting path in G , since if either of its endpoints are covered by M , they must also be covered by the same edge in M' , but then M' wouldn't be a matching. \square

Question 7. Show that any non-maximum matching in a graph admits an augmenting path.

We can find the maximum matching in polynomial time, with so-called “Blossom algorithms”, which find an augmenting path in $O(m\sqrt{n})$. This also means we can determine the edge-cover number $\beta'(G)$ in polynomial time.

2.1.1 Tutte's theorem

Definition 2.1.19. A component of a graph G is **ODD** if it has an odd number of vertices. We denote the number of odd components in G with $o(G)$.

Theorem 2.1.20 (Tutte). *A graph G has a perfect matching if and only if for any $S \subseteq V(G)$, $|S| \geq o(G - S)$ holds. This is called Tutte's condition.*

Proof. Left to right: Let $S \subseteq V(G)$ and M be a perfect matching in G . Let H_1, \dots, H_k be the components of $G - S$. If H_i is an odd component, then there exists at least one M -edge between $V(H_i)$ and S . Therefore $|S| \geq o(G - S)$.

Right to left: Suppose that Tutte's condition holds for a graph F but there is no perfect matching in F . Now add edges to F so that this still holds after the addition, and let G be a maximal such graph on $n(F)$ vertices. If we consider Tutte's condition on $S = \emptyset$, we see that there are no odd components in G , which means $n(G)$ is even.

Now take any edge in the complement of G . Since G is maximal, $G + e$ is not a counterexample, so it either has a perfect matching, or Tutte's condition does not hold for it. We know that for any $S \subseteq V(G)$, $|S| \geq o(G - S)$. After having added an edge, at

most one pair of components in $G - S$ is joined. If this was a pair of odd components, $o(G - S)$ has reduced, and in all other cases, it has remained the same. This means that $G + e$ must satisfy Tutte's condition, so it must have a perfect matching as it is not a counterexample.

We will consider two cases. Let U be the set of universal vertices in G , that is, the vertices adjacent to every other vertex, and let H_1, \dots, H_k be the components of $G - U$. In the first case, suppose every H_i induces a complete graph. We will construct a perfect matching. For the even components, we may create a matching within the component, and for the odd components, we may connect the one remaining vertex with U . We are left over with an even number of vertices in U (since n is even), which we may form a matching with, as $G[U]$ is a complete graph. So in this case, G is not a counterexample, which is a contradiction.

Now suppose there is a non-complete component H_i . Then there exist vertices x, y, z for which $xy \notin E(H_1)$ but both xz and yz are in $E(H_1)$. There is also another vertex $w \in V(G)$ for which $zw \notin E(G)$, since z is not a universal vertex. Let $G_1 = G + xy$ and $G_2 = G + zw$. We know both these graphs have perfect matchings, which must include xy and zw respectively. Denote the perfect matchings with M_1 and M_2 and consider $G[M_1 \triangle M_2]$. Note that every non-isolated vertex in this graph is of degree 2, since it is covered by both perfect matchings. These vertices must of course appear in even cycles.

If xy and zw belong to different components, then we may choose edges in each component, and the edges deleted by the symmetric difference, to form a perfect matching, and we can avoid taking xy or zw . This is a contradiction. Alternatively, if xy and zw belong to the same component, they appear in the same cycle. Without loss of generality they appear in the order $zwx y$ (but not necessarily adjacent). To form a new perfect matching, we will take the edge xz , and one edge set from each side of the cycle. This avoids both new edges xy and zw , so we have a contradiction. \square

Question 8. State and prove Tutte's theorem.

We also have the Berge-Tutte formula, which states that a maximum matching in G leaves uncovered exactly

$$\max_{S \subseteq V(G)} \{o(G - S) - |S|\}$$

vertices.

2.1.2 Factors

Definition 2.1.21. A **FACTOR** is a spanning subgraph (it contains all vertices). A k -**FACTOR** is a k -regular spanning subgraph.

Definition 2.1.22. A **CUBIC GRAPH** is a 3-regular graph.

Theorem 2.1.23 (Petersen). *Every bridgeless cubic graph has a 1-factor.*

Proof. We will prove that Tutte's condition holds. Take $S \subseteq V(G)$. Since the graph is 3-regular, the number of edges between S and \bar{S} is $|E(S, \bar{S})| \leq 3|S|$. If H_i is an odd component in $G - S$, then $E(V(H_i), S)$ contains at least one edge. Note that

$$2m(H_i) + |E(H_i, S)| = 3n(H_i) = \sum_{v \in H_i} \deg_G(v)$$

where $m(H_i)$ is the number of edges in H_i . Since the RHS is odd, $|E(H_i, S)|$ must be odd. Also, $|E(H_i, S)| \neq 1$, since that would be a cut edge (a bridge) otherwise. Therefore, $|E(H_i, S)| \geq 3$. Then $3|S| \geq |E(S, \bar{S})| \geq 3\sigma(G - S)$. \square

We can also improve this theorem, for example allowing for precisely one bridge in the graph, and the statement still holds. Or we could require that all cut edges lie on the same path, and it would still hold.

Question 9. State and prove Petersen's theorem for cubic graphs.

Theorem 2.1.24. *If G is a k -regular graph and k is even, then G has a 2-factor.*

Proof. We can limit ourselves to connected graphs. By Euler's theorem, there exists an Eulerian circuit C in G . We can fix a direction for this C and orient the edges according to it. From this, we obtain a directed graph \vec{G} . For every vertex v , C enters and exits v exactly $k/2$ times.

Define a bipartite graph F_G with partite classes $A = \{a_1, \dots, a_n\}$ and $B = \{b_1, \dots, b_n\}$ for $n = n(G)$ and the edge set

$$a_i b_j \in E(F_G) \Leftrightarrow v_i v_j \in E(\vec{G}).$$

Clearly, F_G is a $\frac{k}{2}$ -regular graph, so it has a perfect matching M . Define a spanning subgraph in G with the following edge set Q :

$$v_i v_j \in Q \Leftrightarrow a_i b_j \in M \vee a_j b_i \in M.$$

Every vertex v_i is covered exactly twice by Q . \square

Question 10. Show that every even-regular graph has a 2-factor.

2.2 Connectivity

Definition 2.2.1. The CONNECTIVITY NUMBER $\kappa(G)$ is the minimum number of vertices in $S \subseteq V(G)$ such that $G - S$ is either disconnected or contains only one vertex.

Definition 2.2.2. A graph G is k -CONNECTED if $\kappa(G) \geq k$ or if the removal of $k - 1$ vertices always results in a connected graph with at least two vertices.

Remark. In any graph, $\kappa(G) \leq \delta(G)$. If G is complete, then $\delta(G) = n - 1 = \kappa(G)$.

Remark. If A is an independent set, removing every other vertex gives a disconnected graph, so $\kappa(G) \leq n - \alpha(G) = \beta(G)$.

Question 11. Define k -connected graphs. Explain why $\kappa(G) \leq \beta(G)$.

Theorem 2.2.3. *The minimum number of edges in a k -connected graph of order n is $\lceil \frac{nk}{2} \rceil$, if $n > k \geq 2$.*

Proof. If the graph is k -connected, then $k \leq \kappa(G) \leq \delta(G)$, so

$$m(G) = \frac{1}{2} \sum_{v \in V} \deg_G(v) \geq \frac{nk}{2}.$$

We will show that there exists a k -connected graph of order n with the specified number of edges. For this, we define Harary graphs $H_{n,k}$ as follows.

- If k is even, $H_{n,k} = C_n^{k/2}$ (i.e. the graph we get by connecting all vertices from C_n which are at a distance of at most $k/2$).
- If k is odd and n is even, $H_{n,k}$ is $C_n^{(k-1)/2}$, along with all edges between two opposing vertices of the cycle.
- If both n and k are odd, then we again take $C_n^{(k-1)/2}$ and add the edges between i and $i + \frac{n-1}{2}$ (if we label the vertices $0, 1, \dots, n-1$).

All these graphs have $m(H_{n,k}) = \lceil \frac{nk}{2} \rceil$.

We will show that all these graphs are k -connected. For the first case, if k is even, let S be a vertex set with $|S| = k-1$. We define a big gap as $\frac{k}{2}$ consecutive vertices in S , and claim the following:

- If there is no big gap between u and v in a certain direction, then we may find a uv -path in that direction. This is clear from the construction.
- For any $u, v \in V(G) \setminus S$, there is a uv -path in $G - S$. Since there can be only one big gap between them, we can just avoid it by going in the other direction, so this is also clear.

If k is odd and n even, we define a big gap as $\frac{k-1}{2}$ consecutive missing vertices. Similarly as before, if there is no big gap on a path from u to v , we can find a path after removing S . But in this case, both paths may contain a big gap. Let P and Q be the two paths along the cycle. If there are big gaps along both, we know the length of both is at least $\frac{k-1}{2} + 1$. Let u' and v' be the opposite vertices of u and v . Suppose that Q is longer than P , and split it into paths Q_1 , Q_2 and Q_3 by v' and u' .

Note that by symmetry, the length of Q_2 (the center region) is also at least $\frac{k+1}{2}$, so the big gap in Q cannot cover both u' and v' . We can then find a u, v -path in $G - S$ using one of these vertices.

We can consider the case of odd k and odd n similarly, it's just more annoying to write down. In all cases, all graphs have precisely $\lceil \frac{kn}{2} \rceil$ edges. \square

Question 12. What is the minimum number of edges in a k -connected graph on n vertices? Prove the bound.

Definition 2.2.4. A set $F \subseteq E(G)$ is a DISCONNECTING SET if $G - F$ is disconnected.

Definition 2.2.5. An EDGE CUT of A is the set $E(A, \bar{A})$ of edges between A and \bar{A} .

Remark. An edge cut is a disconnecting set. A minimal disconnected set is an edge cut.

Definition 2.2.6. The EDGE-CONNECTIVITY number of G is the minimum number of edges in a disconnecting set. We denote it by $\kappa'(G)$. A graph is k -EDGE-CONNECTED if the removal of less than k edges always leaves a connected graph.

Theorem 2.2.7. Suppose G is a simple graph with $n(G) \geq 2$. Then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof. The second inequality is clear. Consider a minimum edge cut $E(A, \bar{A})$ in G . By definition $|E(A, \bar{A})| = \kappa'(G)$. We will show that there is a vertex cut with at most $\kappa'(G)$ vertices. For that, consider two cases. If $E(A, \bar{A})$ forms a complete bipartite graph, then

$$|E(A, \bar{A})| = |A| |\bar{A}| = |A| (n - |A|).$$

Since $0 < |A| < n$, we have $|E(A, \bar{A})| \geq n - 1$, so $\kappa'(G) \geq n - 1$, but $\kappa(G) \leq n - 1$ and $\kappa(G) \leq \kappa'(G)$.

In the second case, if there are vertices $x \in A$ and $y \in \bar{A}$ which are nonadjacent, then we may choose an endpoint different from x and y in each edge in the edge cut. This gives us a vertex cut with at most $|E(A, \bar{A})|$ vertices, in which x and y are not cut, and are disconnected. \square

Corollary 2.2.8. If G is k -connected, then G is k -edge-connected.

Question 13. Show that a k -connected graph is always k -edge-connected.

Corollary 2.2.9. The minimum number of edges in a k -edge-connected graph on n vertices is $\lceil \frac{kn}{2} \rceil$.

Proof. We know that $k \leq \kappa'(G) \leq \delta(G)$, so

$$m(G) = \frac{1}{2} \sum_{v \in V} \deg_G(v) \geq \frac{1}{2} n \delta(G)$$

which means $m(G) \geq \lceil \frac{nk}{2} \rceil$. For the other direction, note that the Harary graphs are k -edge-connected. \square

Theorem 2.2.10 (Whitney). *If G is a 2-connected graph, then for every $u, v \in V(G)$, there are two internally disjoint u, v -paths. The converse also holds.*

Proof. For the left implication, suppose $u, v \in V(G - x)$ for some vertex x . There are two disjoint paths from u to v in G , and at most one of them includes x , so there is a path from u to v in $G - x$. This means our graph has no cut vertex, so it is 2-connected.

Now consider the right implication. Let u, v be vertices in G . Induction on $d = d(u, v)$. For $d = 1$, we know that G is 2-edge-connected (since it is 2-connected), so there is no bridge in G . If we remove uv , then the graph must still be connected. Therefore, there is another u, v -path in G .

For a general d , let w be the neighbour of v on the shortest u, v -path. Then $d(u, w) = d - 1$. By the induction hypothesis, there are two internally disjoint paths P, Q from u to w . Consider two cases. If $v \in V(P)$ (or Q , symmetrically), then we have two internally disjoint u, v -paths in

$$u \xrightarrow{P} v, \quad u \xrightarrow{Q} w \rightarrow v.$$

Otherwise, if v is on neither path, then consider the graph $G - w$. It is still connected, so there is at least one u, v -path R in this graph. If R shares no internal vertex with P or Q , then we may choose $u \xrightarrow{P} w \rightarrow v$ and R as the two paths. Finally, if R intersects P and/or Q , identify the last intersection with either, and label it z . Without loss of generality, $z \in Q \cap R$. Then we have two paths

$$u \xrightarrow{P} w \rightarrow v, \quad u \xrightarrow{Q} z \xrightarrow{R} v.$$

□

Question 14. State and prove Whitney's theorem.

Theorem 2.2.11 (Expansion lemma). *If G is k -connected and we add a new vertex v and k incident edges to the graph, then we obtain a k -connected graph.*

Proof. Call the new vertex y , and let G' be the new graph. We will prove that every vertex cut in G' contains at least k vertices. For that, let S be a vertex cut in G' . Consider three cases.

- If $y \in S$, then let V_1 and V_2 be components of $G \setminus S$. Every path from V_1 to V_2 passes S , which is also true in G , since $y \in S$. In G , every vertex cut contains at least k vertices, so $S \setminus \{y\}$ contains at least k vertices, and S contains at least $k + 1$ vertices.
- If $y \notin S$ and $N(y) \subseteq S$, then y is its own component in $G' \setminus S$. This means $|S| \geq k$.

- Otherwise, there is a vertex y' which is a neighbour of y and belongs to the same component in $G' \setminus S$. If we remove y from G' , S remains a vertex cut in G , as no path exists between the components which avoids S . Note that we don't remove the component of y if we delete the vertex, as the component has at least one other vertex (y'). Since G is k -connected, every vertex cut contains at least k vertices, so $|S| \geq k$. \square

Question 15. State and prove the expansion lemma.

Theorem 2.2.12. *If G is a graph with $n(G) \geq 3$, then the following statements are equivalent:*

- G is 2-connected,
- G is connected and there is no cut vertex,
- for every $u, v \in V(G)$, there are at least two internally disjoint u, v -paths,
- for every $u, v \in V(G)$, there is a cycle through u and v ,
- $\delta(G) \geq 1$ and, for every two edges e_1, e_2 there is a cycle containing e_1 and e_2 .

Proof. We already know the first four statements are equivalent. Suppose G is 2-connected. Take edges $e, f \in E(G)$ and label $e = uv, f = u'v'$. Now add two vertices w, w' to G , w connected to u and v , and w' connected to u' and v' . Let G' be the resulting graph. By the expansion lemma, G' is 2-connected, and satisfies the fourth condition, so there is a cycle through w and w' . This cycle must contain all the new edges and the vertices $u, v, u'v'$. We can now replace the added edges with e and f .

Conversely, take vertices $u, v \in V(G)$. Now let $e = uu'$ and $f = vv'$ be two different edges. There is a cycle through e and f by assumption, so there is a cycle through u and v . \square

Question 16. Show that a graph is 2-connected if and only if it has no isolated vertices and there is a cycle through any pair of edges.

Proposition 2.2.13 (subdivision lemma). *Suppose G' is obtained from G by subdividing an edge $uv \in E(G)$ with a vertex w . Then G is 2-connected if and only if G' is 2-connected.*

Proof. Left to right: Consider two vertices $x, y \in V(G')$. If neither is equal to w , then we can take the cycle through xy in G . If it contains uv , we can replace it with the path uww . If one of the vertices x, y is w , then we find a cycle through u and v , and subdivide the edge.

Right to left: Any cycle in G' that contains w must also contain u and v . \square

Question 17. State the subdivision lemma.

2.2.1 Ear decomposition of a graph

Definition 2.2.14. In a graph G , a path P is an (OPEN) EAR if all internal vertices of P are of degree 2 in G , and for the end vertices of the path, the degree is at least 3.

Definition 2.2.15. An (OPEN) EAR DECOMPOSITION of G is a sequence P_0, P_1, \dots, P_k , where P_0 is a cycle in G , and every other P_i is an ear in the graph $G_i := P_0 \cup P_1 \cup \dots \cup P_i$. We also require $G_k = G$.

Theorem 2.2.16. A graph G is 2-connected if and only if it has an ear decomposition.

Proof. Right to left: We will prove that G_i is 2-connected for every i by induction. For $i = 0$, we know that a cycle is 2-connected. For the induction step, let u, v be the endpoints of P_{i+1} . Add an edge uv to G_i . It is still 2-connected, as adding an edge cannot decrease connectivity. Now we can repeatedly subdivide this edge, and the resulting graph is still 2-connected by the subdivision lemma.

Left to right: Since G is 2-connected, there exists a cycle C , which we can take as P_0 . We will construct the decomposition inductively. If G_i is obtained from P_0 by adding the ears P_1, \dots, P_i , then:

- If $G = G_i$, we're done.
- If G_i is not an induced subgraph of G , we may add an edge from $E(G) \setminus E(G_i)$ with both endpoints in $V(G_i)$. This is a valid ear (as since G_i is 2-connected, $\delta(G_i) \geq 2$), so we can continue the decomposition.
- If G_i is an induced subgraph of G , then there exists a vertex $v \in V(G) \setminus V(G_i)$ which is connected to a vertex $u \in V(G_i)$. Since G is 2-connected, there is a cycle C through the edge uv and some other edge $f \in E(G_i)$. Let u' be the first vertex on this cycle which is in G_i on a $u \rightarrow v \rightarrow \dots$ path. We can add this path between u and u' as P_{i+1} . □

Question 18. What is an open ear decomposition? Show that a graph is 2-connected if and only if it has one.

Proposition 2.2.17. A graph G is 2-edge-connected if and only if G is connected and there is no cut edge in G .

Definition 2.2.18. A CLOSED EAR in a graph G is a cycle for which all but one vertex has degree 2, and the exception has degree at least 4.

Definition 2.2.19. A CLOSED EAR DECOMPOSITION of G is a sequence P_0, P_1, \dots, P_k such that P_0 is a cycle and P_i is either an open or closed ear in $G_i = P_0 \cup P_1 \cup \dots \cup P_i$ for all $i \geq 1$.

Theorem 2.2.20. A graph G is 2-edge-connected if and only if it has a closed ear decomposition.

Proof. Right to left: Since P_0 is a cycle, it is 2-edge-connected. If G_i is 2-edge-connected, so is G_{i+1} : Take any edge $uv \in E(G_{i+1})$. If $uv \in E(G_i)$, then it is part of a cycle in G_i . If $uv \in E(P_{i+1})$, then either P_{i+1} is a closed ear, so a cycle containing uv , or it is an open ear between some vertices x, y , in which case there is an x, y -path in G_i , which forms a cycle with P_{i+1} .

Left to right: Start the closed ear decomposition with a cycle P_0 , and build inductively. If G_i is not an induced subgraph of G , then we can add edges with open ears. If it is, then let v be a vertex in $V(G) \setminus V(G_i)$ connected to a vertex $u \in V(G_i)$. We know uv is in a cycle in G . If u is the only vertex of the cycle in G_i , then we can add it as a closed ear. Otherwise, we can add it as an open ear as in the previous theorem. \square

Question 19. What is a closed ear decomposition? Show that a graph is 2-edge-connected if and only if it has one.

Definition 2.2.21. A STRONG ORIENTATION of an undirected graph G is a digraph \vec{G} which is strongly connected, and which you get from choosing an orientation for each edge in G .

Theorem 2.2.22 (Robbins). *An undirected graph G has a strong orientation if and only if it is 2-edge-connected.*

Proof. Left to right: Suppose that G is not 2-edge-connected, and consider two cases.

- If G is not connected, then there can be no strong orientation.
- If G has a cut edge $e = xy$, then for whatever orientation of e we choose, there is no path between those vertices in the opposite direction.

Right to left: If G is 2-edge-connected, then we have a closed ear decomposition P_0, \dots, P_k . We can orient the edges in P_0 consistently to get a strongly connected graph. Whenever you add an ear (open or closed) to \vec{G}_i , orient the new edges consistently. You can show via simple casework that the new digraph is still strongly connected. \square

Question 20. State and prove Robbins' theorem.

2.2.2 x, y -cuts

Definition 2.2.23. If x and y are nonadjacent vertices in the graph G , then $S \subseteq V(G)$ is an x, y -CUT if x and y belong to different components in $G - S$. We label the minimum size of an x, y -cut in G with $\kappa_G(x, y)$.

Definition 2.2.24. The maximum number of internally vertex-disjoint x, y -paths in a graph G is labeled with $\lambda_G(x, y)$.

Theorem 2.2.25 (Menger's theorem for vertex cuts). *If x and y are nonadjacent vertices in G , then $\kappa_G(x, y) = \lambda_G(x, y)$.*

Proof. Denote $\kappa = \kappa_G(x, y)$ and $\lambda = \lambda_G(x, y)$. We have $\lambda \leq \kappa$ since, if there are λ internally vertex-disjoint x, y -paths in G , we have to remove at least λ vertices to separate x and y , one on each path.

For the other direction, use induction on $n(G)$. If $n(G) = 2$, since x and y are not connected, they must be isolated, so $\kappa = \lambda$. For the induction step, consider two cases. In the first case, if there is a minimum x, y -cut in S such that $S \neq N(x)$ and $S \neq N(y)$, then at least one neighbour of x is not in S (as $N(x)$ is an x, y -cut and S is minimal). The same holds for y .

Consider x, S -paths (paths from x to a vertex in S which only hit S in one vertex). Let V_1 be the union of the vertex sets of all these paths, and let V_2 be the union of the vertex sets of all y, S -paths. We will prove that $V_1 \cap V_2 = S$. Let X_1 be the component of $G - S$ which includes x , and let X_2 be the component of $G - S$ which includes y . Note that every vertex in S must be adjacent to a vertex in X_1 and to a vertex in X_2 , as otherwise we could remove the offending vertex from S and get a smaller vertex cut.

Suppose that $w \in V_1 \cap V_2 \setminus S$. Then we have an x, w -path which does not intersect S , and an y, w -path which does not intersect S . This can't happen as S is a cut set, so $V_1 \cap V_2 \subseteq S$. The other inclusion clearly holds. As we've noted before, there is a neighbour of x not in S , so $|V_1 \setminus S| \geq 2$ and similarly $|V_2 \setminus S| \geq 2$.

Let

$$G_1 = G[V_1] \cup (\text{a vertex } y' \text{ adjacent to every vertex of } S).$$

We know $n(G_1) < n(G)$. It is easy to prove that S is a minimum x, y' -cut in G_1 , so $\kappa_{G_1}(x, y') = \kappa_G(x, y) = \kappa$, but by the induction hypothesis, $\lambda_{G_1}(x, y') = \kappa_{G_1}(x, y')$. So we have κ internally vertex-disjoint x, y' -paths in G_1 . We can similarly define G_2 as $G[V_2]$ with an added vertex x' , adjacent to every vertex of S , and find κ internally vertex-disjoint x', y -paths. This allows us to construct κ internally vertex-disjoint x, y -paths in G by just connecting the G_1 and G_2 paths which share a common vertex in S .

This concludes the first case, now consider the case where all minimum x, y -cuts are either $N(x)$ or $N(y)$. Consider three subcases.

- If there is a vertex $v \in N(x) \cap N(y)$, then $\kappa_{G-v}(x, y) = \kappa - 1$ and we can use the induction hypothesis for $G - v$.
- If there is a vertex $v \notin N[x] \cup N[y]$, then v does not belong to any minimum x, y -cuts, so $\kappa_{G-v}(x, y) = \kappa_G(x, y) = \kappa$. We can again use the induction hypothesis on $G - v$.
- If neither of the above hold, then $V(G) = N[x] \cup N[y]$ and $N[x] \cap N[y] = \emptyset$. Let F be the bipartite graph obtained by taking $V(F) = N(x) \cup N(y)$ and $E(F) = E(N(x), N(y))$. We may find an x, y -path by taking an edge in F and connecting x and y to the endpoints. If we have a matching in F , we can use it to construct that many internally vertex-disjoint paths, so $\lambda \geq \alpha'(F)$. To find an x, y -cut in

G , we must remove all edges in F , so we need a vertex cover T of F . By König's theorem, $\beta(F) = \alpha'(F)$, so $\kappa \leq \beta(F) = \alpha'(F) \leq \lambda$, which is what we were trying to prove. \square

Question 21. State and prove Menger's theorem for vertex cuts.

Definition 2.2.26. For two vertices x, y of G , a set $R \subseteq E(G)$ is an x, y -EDGE CUT if $G - R$ is disconnected and x, y belong to different components of $G - R$. We denote the minimum size of an x, y -edge cut in G by $\kappa'_G(x, y)$. We also define $\lambda'_G(x, y)$ to be the maximum number of edge-disjoint x, y -paths in G .

Theorem 2.2.27 (Menger's theorem for edge cuts). *Let $x, y \in V(G)$. Then $\kappa'_G(x, y) = \lambda'_G(x, y)$.*

Proof. Let G' be obtained by adding two vertices to G , u and v , and edges xu and yv . It is easy to see that a ux, yv -path in $L(G')$ corresponds to an x, y -path in G . By Menger's theorem for vertices,

$$\lambda'_G(x, y) = \lambda_{L(G')}(ux, yv) = \kappa_{L(G')}(ux, yv).$$

Clearly, a vertex cut in $L(G')$ that separates ux and yv corresponds to an edge cut in G that separates x and y , which finishes the proof. \square

Question 22. State and prove Menger's theorem for edge cuts.

Lemma 2.2.28. *Let $e \in E(G)$. Then $\kappa(G) - 1 \leq \kappa(G - e) \leq \kappa(G)$.*

Proof. Clearly $\kappa(G - e) \leq \kappa(G)$. Suppose that the strong inequality holds. Then let S be a minimum vertex cut in $G - e$ for $e = xy$, so $|S| = \kappa(G - e)$ and $G - e - S$ is disconnected. Since $|S| < \kappa(G)$, it is not a vertex cut in G , so x and y belong to different components of $G - e - S$. Let x be in the component X and y in the component Y . Consider three cases:

- if $|X| \geq 2$, then $S \cup \{x\}$ is a vertex cut in G ,
- if $|Y| \geq 2$, then $S \cup \{y\}$ is a vertex cut in G ,
- If $|X| = |Y| = 1$, then $n(G) - 2 = \kappa(G - e) = |S|$. We know $\kappa(G) \leq n(G) - 1$, so $\kappa(G - e) + 1 \geq \kappa(G)$. \square

Question 23. Show that for any $e \in E(G)$, we have $\kappa(G) - 1 \leq \kappa(G - e) \leq \kappa(G)$.

Theorem 2.2.29 (Menger). *In every graph G ,*

$$\kappa'(G) = \min_{x \neq y} \lambda'_G(x, y), \quad \kappa(G) = \min_{x \neq y} \lambda_G(x, y).$$

2 Teorija grafov

Proof. For any $x, y \in V(G)$, if S is an x, y -edge-cut, then it is an edge-cut in G , so

$$\min_{x \neq y} \kappa'_G(x, y) \geq \kappa'(G).$$

If S is an edge cut, then it separates two vertices, so

$$\kappa'(G) \geq \min_{x \neq y} \kappa'_G(x, y).$$

By Menger's theorem for edges, $\lambda'_G(x, y) = \kappa'_G(x, y)$.

For the second claim, we analogously show (if G is not complete)

$$\kappa(G) = \min_{x \neq y, xy \notin E(G)} \lambda_G(x, y).$$

It suffices to prove that for any two adjacent vertices x, y that $\lambda_G(x, y) \geq \kappa(G)$. For that, define $G' = G - xy$. Then $\lambda_{G'}(x, y) = \lambda_G(x, y) - 1$ since the single edge was a path. By Menger's theorem for G' , we have

$$\kappa_{G'}(x, y) = \lambda_{G'}(x, y) = \lambda_G(x, y) - 1,$$

and by the preceding lemma, $\kappa(G') \geq \kappa(G) - 1$, so

$$\lambda_G(x, y) = \kappa_{G'}(x, y) + 1 \geq \kappa(G).$$

□

Question 24. State Menger's theorem.

2.3 Coloring

Remark. In any graph, $\omega(G) \leq \chi(G) \leq \Delta(G) + 1$.

Remark. As the color classes are independent sets, the number of vertices in a color class is at most $\alpha(G)$, so

$$\chi(G) \geq \frac{n(G)}{\alpha(G)}.$$

Theorem 2.3.1 (Welsh-Powel). *If $d_1 \geq d_2 \geq \dots \geq d_n$ is the degree sequence of the vertices of G , then*

$$\chi(G) \leq 1 + \max_{i=1, \dots, n} \{\min\{d_i, i - 1\}\}.$$

Theorem 2.3.2 (Brooks). *If G is connected and not a complete graph or odd cycle, then $\chi(G) \leq \Delta(G)$.*

Proof. Let $k = \Delta(G)$. Consider two cases. First, if G is not k -regular, then there is a vertex v_n with degree at most $k - 1$. Define the following vertex order for a greedy coloring. Consider a breadth-first search tree, rooted in v_n . Order the vertices such that every child of the tree precedes its parent. This way, each vertex is preceded by at most $k - 1$ of its neighbours, so we can color the entire graph with at most k colors.

In the second case, if G is k -regular, we can quickly write off $k = 1$ as then $G = P_2$, which is a complete graph, and the case $k = 2$, where G must be a cycle. We have excluded odd cycles, and for even cycles, $\chi(G) = \Delta(G) = 2$. So let $k \geq 3$. We will consider three subcases.

If $\kappa(G) = 1$, then there exists a cut vertex $x \in V(G)$. Let V_1, V_2 be disconnected vertex sets in $G - x$ such that $V_1 \cup V_2 = V(G - x)$, and let $G_i = G[V_i \cup \{x\}]$. Since x has neighbours in both V_1 and V_2 , the degree of x in either G_i is at most $k - 1$ and G_1, G_2 are not regular. So we can color them with at most k colors. We can permute the colors in one of the colorings so that x has the same color in both.

In the second subcase, suppose $\kappa(G) = 2$. Then we have $x, y \in V(G)$ such that $G - \{x, y\}$ is disconnected. Define V_1 and V_2 as before and $G_i = G[V_i \cup \{x, y\}]$. We have $\deg_{G_i}(x) < \deg_G(x) = k$, so neither G_i is regular. Therefore, they are k -colorable, so we have colorings φ_1, φ_2 . If $\varphi_1(x) = \varphi_1(y)$ and $\varphi_2(x) = \varphi_2(y)$, or if both of these pairs are nonequal, then we can define a combined k -coloring.

If we cannot find such φ_1, φ_2 however, then without loss of generality every k -coloring of G_1 assigns the same color to x and y , and every k -coloring of G_2 assigns different colors to those vertices. Equivalently, $G_1 + xy$ is not k -colorable, and $G_2 + xy$ is. So $G_1 + xy$ must be k -regular, in which case, we will prove that G_2 can be k -colored with $\varphi_2(x) = \varphi_2(y)$. Let $G'_2 = G_2 - \{x, y\}$. It is k -colorable. Since $G_1 + xy$ is k -regular, x and y have precisely one neighbour each in G_2 , meaning we can choose a color different from the colors of those neighbours (as $k \geq 3$) and find a k -coloring for G_2 with $\varphi_2(x) = \varphi_2(y)$.

Finally, consider the case $\kappa(G) \geq 3$. We have vertices $u, v \in V(G)$ for which $d(u, v) = 2$, since G is not complete. Let z be a common neighbour for them. We want a vertex order such that in a greedy coloring, $\varphi(u) = \varphi(v)$ and the last vertex in the ordering is z . Let $G' = G - \{u, v\}$. Consider a breadth-first search in G' , rooted in z . Since $\kappa(G) \geq 3$, it will not stop immediately. Again, take the order in which every vertex is preceded by at most $k - 1$ children, and z is at the end. Now a greedy coloring $u, v, v_1, v_2, \dots, v_{n-2} = z$ of G will assign $\varphi(u) = \varphi(v) = 1$, and for v_1, \dots, v_{n-3} , at most $k - 1$ neighbours are colored before v_i , so it is assigned a color $\leq k$. We know that z has two neighbours with the same color, so we can find a k -coloring for G . \square

Question 25. State and prove Brooks' theorem.

2.3.1 Mycielski's construction

The Mycielskian $M(G)$ of a graph G is a graph defined as follows:

2 Teorija grafov

- label the vertices of G as v_1, \dots, v_n ,
- create $n + 1$ new vertices u_1, \dots, u_n, z ,
- add connections $u_i v_j$ for all pairs $v_i v_j \in E(G)$,
- add connections $u_i z$ for all i .

Label $V = \{v_1, \dots, v_n\}$ and $U = \{u_1, \dots, u_n\}$.

Theorem 2.3.3. *If G is a graph with at least one edge, then $\chi(M(G)) = \chi(G) + 1$ and $\omega(M(G)) = \omega(G)$.*

Proof. Since G is a subgraph of $M(G)$, we have $\omega(G) \leq \omega(M(G))$. If z is in a clique of $M(G)$, then this is a clique of order at most 2, which appears in G as well. If u_i is in a clique of $M(G)$, then v_i can't be in the same clique, so we can replace u_i with v_i , and find a clique in G of the same size. Therefore $\omega(M(G)) \leq \omega(G)$.

If we have a coloring of G , then we can paint u_i with the same color as v_i , and use a new color for z . So $\chi(M(G)) \leq \chi(G) + 1$. Now suppose that there exists a $\chi(G)$ -coloring of $M(G)$ and label it φ . Without loss of generality, $\varphi(z) = k := \chi(G)$. Then this color does not appear in U , so U is colored with $k - 1$ colors. But then since $\chi(G) = k$, the color k must appear in V . We can replace the colors for those v_i which have $\varphi(v_i) = k$ with the color of u_i , and get a proper $(k - 1)$ -coloring of G , which is impossible as $\chi(G) = k$.
—×— □

Question 26. Show that the difference between $\chi(G)$ and $\omega(G)$ can be arbitrarily large.

Theorem 2.3.4. *If G is a graph on n vertices and $\chi(G) = k$, then $m(G) \geq \binom{k}{2}$. This is sharp for any $n \geq k$.*

Proof. There is a partition of G into k color classes. Since $\chi(G) = k$, there must be at least one edge between any pair of color classes. This bound is sharp, as we can take K_k and add isolated vertices as required. □

Question 27. State and prove a lower bound for $m(G)$ in a graph with $\chi(G) = k$.

2.3.2 Turàn's theorem

Definition 2.3.5. A graph G is k -PARTITE if the vertex set can be partitioned into k classes V_1, V_2, \dots, V_k such that every edge is between different classes.

Remark. A graph G is k -partite if and only if G is k -colorable.

Definition 2.3.6. G is a COMPLETE k -PARTITE GRAPH if every edge between the partite classes is present in G .

Definition 2.3.7. THE TURÀN GRAPH $T_{n,k}$ is the complete k -partite graph on n vertices such that the partite classes are of nearly equal size, i.e. if $||V_i| - |V_j|| \leq 1$ for all i, j .

Theorem 2.3.8 (Turàn). *If G is a graph of order n and $\omega(G) \leq r$, then the number of edges in G is at most the number of edges in $T_{n,r}$.*

Proof. Induction on r . If $r = 1$, there are no edges, so $G = T_{n,1}$. For $r > 1$, let $k = \Delta(G)$ and let v be a vertex with degree k . Let $G' = G[N(v)]$. We know $\omega(G') \leq r - 1$, so by the induction hypothesis, $m(G') \leq m(T_{k,r-1})$.

Consider the following construction. Let H be the graph obtained by a complete join of a graph with $n - k$ independent vertices and the graph $T_{k,r-1}$. Clearly, $n(H) = n$ and $\omega(H) = r$, since $\omega(T_{k,r-1}) = r - 1$. Compute

$$m(H) = m(T_{k,r-1}) + (n - k)k \geq m(G') + (n - k)k$$

and $m(G) \leq m(G') + k(n - k)$, noting that in the remainder of the graph, every vertex has degree at most k , and there are $n - k$ vertices there.

We see that H is an r -partite graph. We claim that among the r -partite graphs on n vertices, $T_{n,r}$ has the most edges. Suppose that F is an r -partite graph with $n(F) = n$. If it is not a complete r -partite graph, then we may add edges until it is. If the sizes of the partite classes in F differ by at least 2, then take a vertex u from the larger class V_i and put it into the smaller class V_j . For this, we break $|V_j|$ edges and add $|V_i| - 1$ edges, so we have increased their total number. \square

Question 28. What is a Turàn graph? State and prove Turàn's theorem.

Remark. As $T_{n,r}$ satisfies the theorem's conditions, the bound is sharp. It is actually the unique graph with the maximum number of edges.

Remark. A similar result holds for graphs with $\chi(G) = r$.

2.3.3 Chordal graphs

Definition 2.3.9. A graph G is CHORDAL if there is no induced subgraph that is isomorphic to a cycle C_n for $n \geq 4$.

Definition 2.3.10. A vertex v is a SIMPLICIAL vertex in G if $N_G[v]$ induces a clique.

Lemma 2.3.11 (Voloshin). *If G is a chordal graph, then for every $x \in V(G)$ there exists a simplicial vertex among the vertices farthest from x .*

Proof. Induction on $n(G)$. If $n(G) = 1$, the statement is obvious. Let $n(G) > 1$. If x is a universal vertex in G , then remove x and apply the induction hypothesis to $G - x$. Otherwise, let T be the set of vertices farthest from x , and let H be a component in T . Take S to be the neighbours of vertices in H which are not themselves in H , so

$S = N(H) \setminus H$. Finally, let Q be the vertices in the component of $G - S$ which contains x .

We will prove that S induces a clique. Take $u, v \in S$. Then both vertices have neighbours in H and in Q , so there exist two u, v -paths through H and Q , respectively. Consider the shortest such paths P_H, P_Q . Combined, they form a cycle of order at least 4. There cannot be a chord in $\{u/v\} \cup H$ or $\{u/v\} \cup Q$, since that would give a shorter path, and there is also no edge between Q and H , so there must be a cord $uv \in E(G)$. So any two vertices in S are connected, meaning S is a clique. By the induction hypothesis on $G[S \cup H]$ with a vertex from S , there is a simplicial vertex in H . \square

Question 29. State and prove Voloshin's lemma.

Theorem 2.3.12. *A graph G is chordal if and only if there is a simplicial elimination ordering of the vertices of G .*

Proof. Right-to-left: If G is not chordal, then we will show there is no simplicial elimination ordering. Suppose there is one, v_1, v_2, \dots, v_n . Since G is not chordal, it has a cycle C_k with $k \geq 4$. Let v_i be the first vertex of this cycle in the ordering. Then v_i has two neighbours along the cycle with no edge between them, which contradicts the definition of the simplicial elimination ordering.

Left-to-right: By Voloshin's lemma, there is a simplicial vertex v_1 in G . Then $G - v_1$ is also a chordal graph, so it has a simplicial vertex v_2 . Then $G - v_1 - v_2$ is a chordal graph, and we keep going. We get a simplicial elimination ordering v_1, \dots, v_n . \square

Question 30. Characterize chordal graphs and prove the characterization.

Theorem 2.3.13. *If G is chordal, then $\chi(G) = \omega(G)$.*

Proof. In general $\omega(G) \leq \chi(G)$. Let v_1, \dots, v_n be a simplicial elimination ordering in G . Consider the greedy coloring in the reverse of that order. When we color a vertex v_i , then the neighbours that are already covered are $R_i = N(v_i) \cap \{v_{i+1}, \dots, v_n\}$. As it is a simplicial elimination ordering, R_i is a clique, and $R_i \cup \{v_i\}$ is also a clique. Since $|R_i \cup \{v_i\}| \leq \omega(G)$, at most $\omega(G) - 1$ neighbours of v_i are colored before v_i , so we can color v_i with one of $\{1, 2, \dots, \omega(G)\}$. Since this is true for any v_i , we have $\chi(G) \leq \omega(G)$. \square

2.3.4 Perfect graphs

Definition 2.3.14. A graph G is a PERFECT GRAPH if $\chi(H) = \omega(H)$ holds for every induced subgraph H of G .

Theorem 2.3.15. *Chordal graphs are perfect.*

Proof. An induced subgraph of a chordal graph is chordal. \square

Question 31. Show that chordal graphs are perfect.

Theorem 2.3.16. *Bipartite graphs are perfect.*

Proof. If G is bipartite, then either $E(G) = \emptyset$, in which case $\omega(G) = \chi(G) = 1$, or $E(G) \neq \emptyset$, in which case $\chi(G) = \omega(G) = 2$. Every subgraph of a bipartite graph is bipartite. \square

Theorem 2.3.17 (Vizing). *For every graph G , we have $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$.*

Theorem 2.3.18. *If G is bipartite, then $\chi'(G) = \Delta(G)$.*

Proof. Let $k = \Delta(G)$. Consider the following two cases. First, if G is k -regular, there exists a perfect matching M_1 in G . The graph $G - M_1$ is $(k - 1)$ -regular and bipartite, so there exists a perfect matching M_2 in $G - M_1$. We can continue this procedure to find matchings M_1, \dots, M_k . Now assign color i to M_i .

In the second case, if G is not k -regular, we will construct a k -regular bipartite graph which has G as a subgraph. We add edges to G such that the maximum degree remains k and the graph remains bipartite. If this is not possible but there are still vertices of degree $< k$, then we can follow the following construction.

- Create a copy G' of G , except it has partite classes switched.
- Create a new graph by combining G and G' , and adding connections $u \sim u'$ for $u \in G$ with $\deg(u) < k$.

This increases the minimum degree of G by at least one, and we still have a bipartite graph. If we continue this process, we will get a k -regular bipartite graph. \square

Question 32. Show that bipartite graphs are Vizing class one.

Theorem 2.3.19. *If G is bipartite, then its line graph $L(G)$ is perfect.*

Proof. For every graph, $\chi'(G) = \chi(L(G))$, as every edge coloring of G corresponds to a vertex coloring of $L(G)$. If G is bipartite, $\chi'(G) = \Delta(G)$. The maximal cliques in $L(G)$ corresponds to the vertices of G (since there are no triangles). If a vertex is of degree k , then the corresponding clique in $L(G)$ is also of size k , so $\Delta(G) = \omega(L(G))$. This means $\chi(L(G)) = \omega(L(G))$.

The induced subgraphs of $L(G)$ are still line graphs of bipartite graphs, as if $H = L(G)[S]$, then H is the line graph of the subgraph of G obtained by only keeping the edges corresponding to the vertices in S . \square

Question 33. Show that the line graph of a bipartite graph is perfect.

Theorem 2.3.20 (Perfect graph theorem). *A graph G is perfect if and only if its complement \overline{G} is perfect.*

Theorem 2.3.21 (Strong perfect graph theorem). *A graph G is perfect if and only if neither G nor \overline{G} have an induced odd cycle of size at least 5.*

Question 34. State the weak and strong perfect graph theorems.

Definition 2.3.22. A (β, α') -PERFECT GRAPH is a graph for which $\beta(H) = \alpha'(H)$ for every induced subgraph H .

Theorem 2.3.23. *A graph is (β, α') -perfect if and only if it is bipartite.*

2.3.5 Gallai-Roy-Vitaver theorem

Theorem 2.3.24. *If G is a simple graph and \vec{D} is an orientation of G , and $l(\vec{D})$ is the longest path in \vec{D} , then $\chi(G) \leq l(\vec{D}) + 1$. Further, there is an orientation for which the equality holds.*

Proof. Consider an arbitrary orientation \vec{D} of G . Let \vec{D}' be a maximal subdigraph of \vec{D} that does not contain any directed cycles. Define a vertex coloring c for G with $c(v)$ being one plus the length of the longest directed path in \vec{D}' that ends in v . This is a proper vertex coloring; consider an edge $\vec{uv} \in \vec{D}$. Consider two cases.

- If $\vec{uv} \in \vec{D}'$, then $l'(u) < l'(v)$, since v cannot be on the longest path ending at u (as there are no directed cycles).
- If $\vec{uv} \notin \vec{D}'$, then there is a directed path in \vec{D}' between v and u (which forms a cycle with \vec{uv}). Then the coloring c gives increasing colors to the vertices along this path, and $c(u) > c(v)$.

We skip the proof of the second statement. □

Question 35. State the Gallai-Roy-Vitaver theorem. Prove the $\chi(G) \leq$ inequality.

2.4 Planar graphs

Definition 2.4.1. If G is a graph, then the DRAWING of G into the plane is a function $h(x)$ defined on $V(G) \cup E(G)$ such that for every vertex v , $h(v)$ is a point and for every edge uv , $h(uv)$ is a $h(u), h(v)$ -curve.

Remark. We can assume without loss of generality that the image of an edge is a polygonal curve, so composed of finitely many line segments.

Definition 2.4.2. The PLANAR EMBEDDING of a graph G is a drawing where the curves corresponding to the edges of G do not intersect, except in common end vertices.

Definition 2.4.3. A PLANE GRAPH is a particular embedding of a planar graph.

Theorem 2.4.4 (Jordan). *Every closed simple curve in the plane divides it into exactly two regions, a bounded inside region and an unbounded outside region.*

Definition 2.4.5. If G is a plane graph, then a **FACE** of G is a maximal region that does not contain any points from the image of the embedding function f .

Definition 2.4.6. The **DUAL GRAPH** G^* of a plane graph G is obtained by switching the role of the vertices and faces of G , with two faces being connected if and only if they share a common border. If there are multiple common edges on the boundary between two faces, make that many edges. If there is an edge which borders twice on the same face, add a loop.

Remark. Two different embeddings of a planar graph may have non-isomorphic dual graphs.

Proposition 2.4.7. *Dual graphs are always connected.*

Proposition 2.4.8. *If G is a connected plane graph, then $(G^*)^* \cong G$.*

Remark. Note that dual graphs are always planar.

Definition 2.4.9. The **LENGTH** $l(F)$ of a face F in a plane graph G is the total length of the walk(s) along the boundary of F , in units of number of edges.

Remark. With this definition, cut edges are counted twice.

Remark. For any plane graph,

$$\sum_{F \text{ face}} l(F) = 2m(G).$$

Theorem 2.4.10. *Suppose G is a plane graph. The following statements are equivalent:*

- G is bipartite,
- every face of G has an even length,
- G^* is Eulerian (connected and all vertices are of even degree).

Proof. 1 to 2: The length of a face is the length of a cycle in G , and possibly some cut edges, which are counted twice.

2 to 1: Suppose G is not bipartite, so there is an odd cycle C . Consider the sum of the lengths of faces inside C . Every edge inside C (but not part of C) is counted twice, but every edge of C is counted only once. So the sum is odd. Then there must be a face F in the cycle with an odd length, which is a contradiction.

Equivalence between 2 and 3: Note that G^* is connected and $\deg_{G^*}(x) = l(x)$. □

Question 36. When is a plane graph bipartite? State two equivalences.

Theorem 2.4.11. *If G is a plane graph and $D \subseteq E(G)$, then D is the set of edges of a cycle if and only if the corresponding dual edge set D^* is a minimal edge cut in G^* .*

Proof. Consider three cases.

- If D is exactly the edge set of a cycle C , then D^* contains all edges between the inside and outside faces of C , so it is an edge cut in G^* and it is minimal, as the faces inside C are all connected in G^* .
- If D is a proper subset of the set of edges of a cycle, we can prove that D^* is an edge cut, but not minimal.
- If D does not contain the edge set of a cycle, we can show that D^* is not an edge cut. □

Definition 2.4.12. A planar graph G is OUTERPLANAR if there exists an embedding such that the boundary of the outer face contains all vertices.

Theorem 2.4.13. If G is a simple outerplanar graph, then $\delta(G) \leq 2$.

Proof. If $n \leq 4$, then this is simple casework. For $n \geq 4$, we will prove that additionally, there are at least two nonadjacent such vertices. Induction on $n = n(G)$. We covered the base case $n = 4$ before, just do the induction step, in which we consider two cases.

In the first case, if there is a cut vertex v , then $G - v$ splits into two graphs G_1, G_2 . Both $G_1 + v$ and $G_2 + v$, with the original edges from v , are outerplanar graphs. By the induction hypothesis, there are two nonadjacent vertices in $G_1 + v$ of degree at most 2, so there is a vertex $z_1 \neq v$ in $G_1 + v$ with degree at most 2 in $G_1 + v$. Similarly, there is a vertex z_2 in $G_2 + v$ for which the same holds. These vertices are of the same degree in G .

In the second case, if there is no cut vertex, then there is no cut edge, and the boundary of the outer face is a Hamiltonian cycle. If there is no chord in this cycle, the statement is proven. Otherwise, let xy be a chord. Consider the two cycles H_1, H_2 that the chord divides our Hamiltonian cycle into. Both are outerplanar, so using the induction hypothesis on each, we can find two nonadjacent vertices of degree at most 2. They can't be adjacent to a vertex in the other part of the Hamiltonian cycle, as that edge would cross xy . □

Question 37. Show that a simple outerplanar graph has $\delta(G) \leq 2$.

Theorem 2.4.14 (Euler). If G is a connected plane graph, then

$$n(G) + f(G) - m(G) = 2.$$

Definition 2.4.15. A SUBDIVISION of G is obtained by replacing some edges of G with internally vertex-disjoint paths.

Definition 2.4.16. A KURATOWSKI GRAPH is one which is a subdivision of K_5 or $K_{3,3}$.

Lemma 2.4.17. If G is a planar graph and $e \in E(G)$, then there is an embedding of G such that e is on the boundary of the outer face.

Proof. Consider an embedding of G and place a sphere on that plane, touching one of the faces adjacent to e . Project the entire drawing onto the sphere by taking the intersection of a line from a given point to the highest point on the sphere. Every face on the plane will correspond to a region of the sphere with the same set of boundary edges. Now project the surface of the sphere onto another plane above the sphere, by drawing a line through the bottom-most point of the sphere and another point of the sphere, and taking the intersection of that line with the new plane. \square

Question 38. Show that for any edge e of a planar graph, there is an embedding with e on the boundary of the outer face.

Lemma 2.4.18. *If G is a minimal nonplanar graph, then it is 2-connected.*

Proof. If G is disconnected, it cannot be minimal nonplanar, as components are proper subgraphs, which would all be planar if G was minimal. If G has a cut vertex v , then consider the v -lobes (the subgraphs of G induced by each of the components of $G - v$, combined with v). By the minimality of G , all v -lobes are planar subgraphs. Lemma 2.4.17 states that these graphs can be embedded in such a way that v is on the outer face. We can transform each picture such that they all fit in a portion of the plane, so we can combine the drawings to find G is planar. \square

Question 39. Show that a minimal nonplanar graph is 2-connected.

Lemma 2.4.19. *If $S = \{x, y\}$ is a minimum vertex cut in G and G is nonplanar, then $G - S$ contains a component G_i such that the S -lobe H_i , along with the edge xy , is nonplanar.*

Proof. Suppose to the contrary, that each $H_i + xy$ is planar. By the first lemma, we can embed all these graphs such that xy is on the outer face. We can construct an embedding for $G + xy$ in the following way:

- Start with the embedding of $H_1 + xy$
- Choose a face in $H_1 + xy$ that borders xy . Then we may attach the embedding of $H_2 + xy$ onto that face without intersecting $H_1 + xy$.
- Continue this process.

So $G + xy$ is planar, and G must be too. \square

Question 40. Show that in a nonplanar graph with a cut set $S = \{x, y\}$, there is a lobe H in $G - S$ such that $H + xy$ is nonplanar.

Lemma 2.4.20. *If G is a nonplanar graph without a Kuratowski subgraph and G has the minimum number of edges under this condition, then it is 3-connected.*

Proof. We know by lemma 2.4.18 that G is 2-connected. Suppose that $S = \{x, y\}$ is a vertex cut. Again as before consider the S -lobes H_1, H_2, \dots with an added edge xy . By lemma 2.4.19, we know there is a graph $H_i + xy$ which is not planar. By the minimality of $m(G)$, we know $H_i + xy$ has a Kuratowski subgraph, as it has fewer edges than G (note that S is a minimal vertex cut). Label that subgraph F . As it's not present in G by assumption, it must include the edge xy .

Consider another component G_j of $G - S$. Since S is a minimum vertex cut, both x and y have neighbours in G_j . Since G_j is connected, there is an x, y -path in H_j (not considering xy). We can replace xy in $H_i + xy$ with that path, and get a Kuratowski graph. But this is a subgraph of G . \square

Question 41. Suppose that G is the graph with the smallest number of edges among all nonplanar graphs without a Kuratowski subgraph. Show that G is 3-connected.

Theorem 2.4.21. *If G is a 3-connected graph with $n(G) \geq 5$, then there exists an edge $e \in E(G)$ such that $G \cdot e$ (G with e contracted) is 3-connected.*

Proof. Suppose there is no such edge. Let $e = xy \in E(G)$. Since $G \cdot e$ is not 3-connected, there is a vertex cut S with 2 vertices. Let w be the contracted vertex of $G \cdot e$. If $w \notin S$, then replacing w with the original edge still leaves $G - S$ disconnected, which can't happen. So $w \in S$. Let $S = \{w, z\}$. Then $S' = \{x, y, z\}$ is a vertex cut in G .

We have shown that every edge xy in G has a mate, so a vertex w such that $\{x, y, w\}$ is a disconnecting set. Let $f = uv$ be an edge with mate z such that $G - \{u, v, z\}$ has the largest component among all edge and mate vertex cuts. Let G_i be that largest component, and G_j be some other component in $G - \{u, v, z\}$. Since $\{u, v, z\}$ is minimal, there is a vertex z' of G_j which is adjacent to z . Let z^* be a mate for zz' and let H be the subgraph induced by $G_i \cup \{u, v\}$. Consider three cases.

- If $z^* \in V(H)$ and $H - z^*$ is disconnected, then $G - \{z, z^*\}$ is disconnected, as at least one component of $H - z^*$ has edges only to z , not u or v . But G is 3-connected, so this can't happen.
- If $z^* \in V(H)$, but $H - z^*$ is connected, then we can similarly show that $\{z, z'\}$ is a disconnecting set.
- If $z^* \notin V(H)$, then $G - \{z, z', z^*\}$ has a component which includes the entire H , but $n(H) > n(G_i)$, so our choice for $\{u, v, z\}$ wasn't optimal.

In each case, we got a contradiction. \square

Question 42. Show that in any 3-connected graph on at least 5 vertices, there is an edge which we can contract and still have a 3-connected graph.

Lemma 2.4.22. *If G contains no Kuratowski subgraphs and $e \in E(G)$, then $G \cdot e$ contains no Kuratowski subgraph.*

Proof. Let $e = xy$ and let w be the contracted vertex in $G \cdot e$. Suppose that $G \cdot e$ contains a Kuratowski subgraph F .

- If F doesn't contain w , then F is present in G .
- If $w \in V(F)$ and $\deg_F(w) = 2$, then we just uncontract the edge and find a Kuratowski subgraph in G .
- If $w \in V(F)$ and $\deg_F(w) \geq 3$, consider three subcases.
 - If $N_F(w) \subseteq N_G(x)$ or symmetrically for y , then we may replace w by x and find a Kuratowski subgraph in G .
 - If all but one neighbour of w is a neighbour of x in G , then we can subdivide the edge going into the neighbour of y and find a Kuratowski subgraph of G .
 - Otherwise, since F is Kuratowski, $\Delta(F) \leq 4$. In this case, we therefore have precisely two x -neighbours and two y -neighbours. Label them x_1, x_2, y_1, y_2 . In this case, F must be a subdivision of K_5 .

Consider the first branch vertices (those of degree at least 3) along the paths $wx_1 \dots x'_1, wx_2 \dots x'_2, wy_1 \dots y'_1, wy_2 \dots y'_2$. These paths are unique as possible intermediate vertices are of degree 2. Since they were part of a subdivision of K_5 in F , there are disjoint paths between any two of these vertices. We can use them to find a subdivision of $K_{3,3}$, with the partite classes being (x, y'_1, y'_2) and (y, x'_1, x'_2) . \square

Question 43. Show that edge contraction preserves the property that a graph has no Kuratowski subgraphs.

Theorem 2.4.23. *If G is a 3-connected graph without Kuratowski subgraphs, then there exists a convex embedding of G such that no 3 vertices are on a line.*

Proof. Use induction on $n(G)$. The base case is $n(G) = 4$, which is just K_4 . For the induction step, since G is 3-connected, there is an edge $e \in E(G)$ such that $G \cdot e$ is 3-connected. Let $e = xy$ and let z be the contracted vertex. Since G has no Kuratowski subgraph, $G \cdot e$ has no Kuratowski subgraph. By the induction hypothesis, there is a convex embedding for $G \cdot e$ with no 3 vertices on a line.

Since $G \cdot e$ is 3-connected, $G \cdot e - z$ is 2-connected, so if we remove the edges incident to z from $G \cdot e$, the region containing z must be a cycle. Let x_1, x_2, \dots, x_k be the neighbours of x on the cycle, labeled in cyclic order. Consider the following cases for the neighbours of y .

- If all neighbours of y are between x_i and x_{i+1} , then we can find a convex embedding for G by drawing y inside the $x_i x x_{i+1}$ angle, and very near x , while drawing x in z 's place. We can ensure that the resulting drawing has convex faces and no three points in a line.

- If x and y have at least 3 common neighbours, then they form a subdivision of K_5 along with x and y .
- If there are two neighbours x'_1, x'_2 of x and two neighbours of y , y_1, y_2 , which are four different vertices such that the cyclic order is $x'_1 y_1 x'_2 y_2$, this induces a subdivision of $K_{3,3}$, which is a Kuratowski subgraph. \square

Question 44. Show that a 3-connected graph without Kuratowski subgraphs can be embedded into the plane such that all interior faces are convex and no 3 vertices are on a line.

Theorem 2.4.24 (Kuratowski). *A graph is planar if and only if it contains no Kuratowski subgraph.*

Proof. Note that a subdivision of G is planar if and only if G is planar, which proves the left-to-right implication. For the other direction, suppose there exist nonplanar graphs without Kuratowski subgraphs. We've shown that the minimal such graph must be 3-connected, so by the previous theorem, it is planar. This is a contradiction. \square

Question 45. State and prove Kuratowski's theorem.

Definition 2.4.25. A graph H is a MINOR in G if we can obtain H after deleting and contracting edges in G .

Proposition 2.4.26. *If G contains a subdivision of a graph F as a subgraph, then F is a minor in G .*

Proof. We can contract the edges of between two subdivision vertices or between a subdivision vertex and a non-subdivision vertex. \square

Remark. The opposite implication is not true. The Petersen graph has K_5 as a minor, but no subgraph is a subdivision of K_5 .

Question 46. Show that if G contains a subdivision of F as a subgraph, then F is a minor in G , but that the reverse doesn't hold.

Theorem 2.4.27 (Wagner). *A graph G is planar if and only if neither K_5 nor $K_{3,3}$ are a minor of G .*

Question 47. State Wagner's theorem.

Theorem 2.4.28 (four-colour theorem). *If G is a planar graph, then $\chi(G) \leq 4$.*

Theorem 2.4.29. *If G is a planar graph, then $\chi(G) \leq 5$.*

Proof. If G is a planar graph, then by Euler's formula, we get $m(G) \leq 3n(G) - 6$, so $\delta(G) \leq 5$. Induction on $n(G)$. Consider a vertex $v \in V(G)$ of degree at most 5. By the induction hypothesis, there is a 5-colouring of $G - v$. If $N(v)$ uses fewer than 5 colours,

we can use a missing colour for v . Otherwise, $\deg_G(v) = 5$ and the neighbours v_1, \dots, v_5 , numbered cyclically, have colours $1, \dots, 5$.

Let $G_{1,3}$ be the subgraph of G induced by the vertices with colours 1 and 3. Consider two cases.

- If v_1 and v_3 are in different components of $G_{1,3}$, then we may switch the colours in one of the two components, and we get a 5-colouring of G where $N(v)$ is coloured with 4 colours.
- If v_1 and v_3 belong to the same component, there is a v_1, v_3 -path in $G_{1,3}$, where vertices alternate between colours 1 and 3. In the original graph, this path forms a cycle with v . One of v_2, v_4 must be inside this cycle, and the other outside. This means v_2, v_4 are in different components of $G_{2,4}$, since any v_2, v_4 -path would need to cross the cycle, so share a common vertex with it, which can't happen, as $G_{1,3}$ and $G_{2,4}$ have no common vertices. This means we can use the previous construction for $G_{2,4}$. \square

Question 48. Prove the 5-color theorem.

Definition 2.4.30. A graph G is a MAXIMAL PLANAR GRAPH if G is planar and for any $e \in E(\overline{G})$, $G + e$ is not a planar graph.

Definition 2.4.31. An embedding of G is a TRIANGULATION if every face has a boundary that is a 3-cycle.

Proposition 2.4.32. Let G be a planar graph with $n(G) \geq 3$. Then the following statements are equivalent.

- $m(G) = 3n(G) - 6$,
- every embedding of G is a triangulation,
- there is an embedding of G that is a triangulation,
- G is a maximal planar graph.

Proof. Note that

$$m = 3n - 6 \Leftrightarrow \sum_{F \text{ face}} l(F) = 3f \Leftrightarrow \text{every face has length 3.}$$

Since the embedding is arbitrary, this proves the equivalence of the first three statements.

1 to 4: If we add an edge, we will have more edges than we can in a planar graph.

4 to 2: Suppose there is an embedding that isn't a triangulation. Then we may add an edge to a face which is a cycle with at least 4 vertices (it may be a degenerate cycle). \square

Question 49. Characterize maximal planar graphs.

3 Teorija izračunljivosti

3.1 Introduction

A TURING MACHINE is defined to consist of the following components. There is an infinite tape divided into cells, each of which contains a symbol from the chosen alphabet Γ . This alphabet must include a **blank** symbol. At the start, only a finite number of cells in the tape have a character different than **blank**. The machine also possesses a read-write head positioned at some cell, and an internal control state, which determines the instruction to be followed.

Instructions are given as a TRANSITION (partial) function f , which maps

$$(\text{state}, \text{character}) \mapsto (\text{new state}, \text{new character}, \text{motion}).$$

To perform an action, a TM will look for rules matching its current control state and the character currently written at the position of the read-write head. When a matching rule is found, the machine switches to the defined new state, writes the specified character on the tape and moves according to the instruction. We limit its motion to three possibilities: one cell to the left or right, or no motion at all. More formally, we may restate the definition as follows:

Definition 3.1.1. A TURING MACHINE is specified by the following:

- a finite TAPE ALPHABET Γ with $\square \in \Gamma$,
- a finite set Q of STATES with **start** $\in Q$,
- a transition partial function

$$\delta : Q \times \Gamma \rightarrow Q \times \Gamma \times \{-1, 0, +1\}.$$

For a given input alphabet $\Sigma_1 \subseteq \Gamma$ and output alphabet $\Sigma_2 \subseteq \Gamma$, a TM specifies a partial function $f : \Sigma_1^* \rightarrow \Sigma_2^*$ if for any $w \in \Sigma_1^*$, running the TM on the input w results in the machine halting in the **halt** state and it has $f(w)$ as the word on the tape, with the head at the leftmost character. We also require $\square \notin \Sigma_1$ or Σ_2 . If the machine does not halt in the **halt** state, we say $f(w)$ is not defined.

Definition 3.1.2. A partial function $f : \Sigma_1^* \rightarrow \Sigma_2^*$ is COMPUTABLE if there exists a TM that computes it.

Question 1. Describe the basic working of a Turing machine. What does it mean for a function to be computable?

Definition 3.1.3. A LANGUAGE over an alphabet Σ is a subset $L \subseteq \Sigma^*$.

For language recognition we require a subset $\Sigma \subseteq \Gamma$ (the INPUT ALPHABET) and two distinguished states, **accept** and **reject**. A language-recognizing Turing machine accepts a word w if, when the machine is run on the input w , the computation halts in the **accept** state. Similarly, w is rejected if the machine halts in the **reject** state. We say

that a Turing machine M DECIDES or COMPUTES L if for any $w \in \Sigma^*$, $w \in L$ implies that M accepts w and $w \notin L$ implies that M rejects w .

If M correctly accepts words of the language, and does not accept words that are not part of the language, we say that M SEMIDECIDES, SEMICOMPUTES or RECOGNIZES L . A language is DECIDABLE or COMPUTABLE if there is a TM which decides it, and SEMIDECIDABLE, SEMICOMPUTABLE or COMPUTABLY INNUMERABLE if there is a TM which recognizes it. Clearly, every decidable language is also semidecidable.

Question 2. What is a decidable and what is a semidecidable language?

Given a k -tape machine (Γ, Q, δ) , we can simulate it by a single-tape machine. We will encode the different tapes by separating them with a special symbol $| \in \tilde{\Gamma}$ and encoding the positions of the read-write heads with another special symbol $\Delta \in \tilde{\Gamma}$. When simulating a computational step of the multi-tape machine, we scan for transitions and implement them manually.

Question 3. How can you simulate a multi-tape Turing machine with a single-tape machine?

Proposition 3.1.4. *There are languages that are not semidecidable.*

Proof. There are only countably many non-equivalent Turing machines, but an uncountable number of languages on any alphabet. \square

We can also give an example of a language that is semidecidable but not decidable. To construct it, we will encode a Turing machine $M = (\Gamma, Q, \delta)$ into a string. We encode it in $\langle M \rangle \in \Sigma_u^*$ for the alphabet

$$\Sigma_u = \{0, 1, -1, [,], \|, \cdot\}.$$

We encode every state $q \in Q$ as a word $\langle q \rangle \in \{0, 1, -1\}^l$, where $l \geq \log_3 |Q|$. We require that the encoding of the start state is 0^l , the encoding of the accepting state is 1^l , and the encoding of the rejecting state is $(-1)^l$. Every symbol $a \in \Gamma$ is encoded as $\langle a \rangle \in \{-1, 0, 1\}^m$ for $m \geq \log_3 |\Gamma|$. We require that the encoding of the blank symbol is 0^m .

Finally, to encode δ , consider an instruction $(q, a) \mapsto (q', b, d)$. This will be encoded as a word

$$[\langle q \rangle \cdot \langle a \rangle \| \langle q' \rangle \cdot \langle b \rangle \cdot d]$$

with $d \in \{0, 1, -1\}$. This encoding has length $2l + 2m + 7$, and allows us to encode the full Turing machine as the encoding of the start state, followed by a dot \cdot , followed by the encoding of the blank symbol, and then followed by the encodings of all transitions one after another. We can encode any word $w \in \Gamma^*$ as a sequence of characters, delimited by \cdot , so that we may define a language L_{accept} as follows: the language includes words of the form $\langle M \rangle \cdot \langle w \rangle$, where M is a single-tape Turing machine with tape alphabet $\Gamma \supseteq \Sigma_u$, and $w \in \Sigma_u^*$ is a word which M accepts.

Question 4. How is L_{accept} defined? Describe the universal encoding of a Turing machine.

Theorem 3.1.5. *The language L_{accept} is undecidable.*

Proof. Suppose that it is decidable, so there exists a Turing machine D which decides it. We define a new Turing machine N with input alphabet Σ_u . This machine reads its input string v and converts it to the string $v \cdot \langle v \rangle$. It then proceeds as D on this input, except it switches the accept and reject states.

Consider what N does when given an input of the form $v = \langle M \rangle$ for some Turing machine M . If D rejects $\langle M \rangle \cdot \langle \langle M \rangle \rangle$, then N terminates on the accepting state, and vice versa. Because D decides L_{accept} , we get from N :

- **accept** iff M does not accept $\langle M \rangle$, and
- **reject** iff M accepts $\langle M \rangle$.

Now run N on $\langle N \rangle$. We get **accept** iff N does not accept $\langle N \rangle$, and **reject** iff N accepts $\langle N \rangle$. This is a contradiction. \square

Question 5. Show that L_{accept} is undecidable.

Using Σ_u , we can also construct the UNIVERSAL TURING MACHINE.

Theorem 3.1.6. *There exists a Turing machine U over the tape alphabet $\Sigma_u \cup \{\square\}$ that exhibits the following behavior: If we run U on the string $\langle M \rangle \cdot \langle w \rangle$, then the resulting execution satisfies the following.*

- *It terminates if and only if M terminates on input w .*
- *It accepts if and only if M accepts w .*
- *It rejects if and only if M rejects w .*

The idea of the proof is to use a three-tape machine, putting the encoding of M on the first tape, the encoding of the state on the second, and the input on the third. Then simulate the execution of M .

Theorem 3.1.7. *The language L_{accept} is semidecidable.*

Proof. We construct a machine S that does the following. It first reads its input word $v \in \Sigma_u^*$ and checks whether v is of the form $\langle M \rangle \cdot \langle w \rangle$. If v is not of this form, then we reject immediately. Otherwise, we run the universal machine U on the input v and end in the end state of U . \square

Question 6. Show that L_{accept} is semidecidable.

Proposition 3.1.8. *If $f : \Sigma_1^* \rightarrow \Sigma_2^*$ and $g : \Sigma_2^* \rightarrow \Sigma_3^*$ are computable, then so is $g \circ f$.*

Note that the composite of two partial functions is defined as

$$(g \circ f)(w) \simeq \begin{cases} g(f(w)) & f(w) \downarrow \\ \uparrow & f(w) \uparrow \end{cases}$$

Definition 3.1.9. A k -tape Turing machine COMPUTES $f : \Sigma_1^* \times \dots \times \Sigma_k^* \rightarrow \Sigma^*$ if when we run the Turing machine on the configuration with a word on each tape, the machine terminates in the halt state if and only if for all (w_1, \dots, w_k) in the domain of f , it halts in the configuration with $f(w_1, \dots, w_k)$ on the first tape and all other tapes blank.

Example. To define the computability for partial functions $f : \mathbb{N} \rightarrow \mathbb{N}$, we can represent numbers using words over $\Sigma_b = \{0, 1\}$ and the representation

$$\gamma_{\mathbb{N}}(w) = \sum_{i=0}^{|w|-1} 2^{|w|-i-1} w_i.$$

If we allow for leading zeros, every number is represented by an infinite number of words. Additionally, zero is also represented by the empty word ε . A Turing machine M computes $f : \mathbb{N} \rightarrow \mathbb{N}$ if it computes $g : \Sigma_b^* \rightarrow \Sigma_b^*$ such that for all words w for which $\gamma_{\mathbb{N}}(g(w)) \simeq f(\gamma_{\mathbb{N}}(w))$. We say that f is COMPUTABLE if there is a Turing machine which computes it.

We could also restrict our representation, for example requiring words to begin with a 1 (we also allow the empty word). Alternatively, we could use e.g. a unary representation.

Definition 3.1.10. A REPRESENTATION of a set X by words over an alphabet Σ is a surjective partial function $\gamma : \Sigma^* \rightarrow X$.

Remark. Only countable sets can be represented.

Definition 3.1.11. Given representations $\gamma_1 : \Sigma_1^* \rightarrow X_1$ and $\gamma_2 : \Sigma_2^* \rightarrow X_2$, a partial function $f : X_1 \rightarrow X_2$ is $(\gamma_1 \rightarrow \gamma_2)$ -COMPUTABLE if there exists a computable partial function $g : \Sigma_1^* \rightarrow \Sigma_2^*$ such that for all words w in the domain of γ_1 , if $g(w)$ is defined, then $\gamma_2(g(w)) \simeq f(\gamma_1(w))$.

Definition 3.1.12. Two representations γ_1 and γ_2 of the same set X are EQUIVALENT if the identity function id_X is $(\gamma_1 \rightarrow \gamma_2)$ -computable and $(\gamma_2 \rightarrow \gamma_1)$ -computable.

Given representations $(\gamma_i : \Sigma_i^* \rightarrow X_i)_{i=1, \dots, k}$, we construct a product representation $\gamma : \Sigma^* \rightarrow X_1 \times \dots \times X_k$, where

$$\Sigma = (\Sigma_1 \cup \Sigma_2 \cup \dots \cup \Sigma_k) \amalg \{\cdot, \cdot\}$$

and

$$\gamma(w_1, \dots, w_k) = (\gamma_1(w_1), \dots, \gamma_k(w_k)).$$

Definition 3.1.13. A partial function $f : \mathbb{N} \rightarrow \mathbb{N}$ is COMPUTABLE if it is $(\gamma_{\mathbb{N}} \times \dots \times \gamma_{\mathbb{N}} \rightarrow \gamma_{\mathbb{N}})$ -computable.

Definition 3.1.14. Given a representation $\gamma : \Sigma^* \rightarrow X$, a subset $A \subseteq X$ is γ -DECIDABLE if there exists a Turing machine M such that for all w in the domain of γ , if $\gamma(w) \in A$, then M accepts w , and if $\gamma(w) \notin A$, then M rejects w . The same subset is γ -SEMIDECIDABLE if there exists a Turing machine M such that for all w in the domain of γ , M accepts w if and only if $\gamma(w) \in A$.

Proposition 3.1.15. *Given a representation γ of X and $A \subseteq X$, A is γ -decidable if and only if its characteristic function is $(\gamma \rightarrow \gamma_b)$ -computable, and A is γ -semidecidable if and only if its partial characteristic function is $(\gamma \rightarrow \gamma_b)$ -computable.*

Remark. Above, γ_b is the representation of $\{0, 1\}$ which maps $0 \mapsto 0$ and $1 \mapsto 1$.

3.1.1 Models of computation

We have many models of computation:

- partial recursive functions,
- λ -calculus,
- Turing machines,
- string rewriting systems,
- unrestricted grammars,
- cellular automata,
- nondeterministic Turing machines,
- quantum Turing machines,
- hypercomputation,
- finite/pushdown automata

Everything on this list, up until hypercomputation, is computationally equivalent. We say that a model is TURING COMPLETE if it can simulate any Turing machine.

There is also the Church-Turing thesis, which states that any Turing complete physically realizable computational model is equivalent to a Turing machine. Informally, the thesis states that the notion of what an algorithm is is equivalent to a Turing machine.

3.2 Computability of natural numbers

Definition 3.2.1. The COMPUTABLE PARTIAL FUNCTIONS is the set

$$\{f : \mathbb{N}^k \rightarrow \mathbb{N} \mid k \geq 0, f \text{ is computable}\}.$$

Definition 3.2.2. The PRIMITIVE RECURSIVE FUNCTIONS are the smallest collection $\mathcal{F} \subseteq \{\mathbb{N}^k \rightarrow \mathbb{N} \mid k \geq 0\}$ of partial functions that satisfies the following properties:

- \mathcal{F} contains the zero function $Z : \{\emptyset\} \rightarrow \mathbb{N}$, which maps $Z() = 0$, the successor function $S : \mathbb{N} \rightarrow \mathbb{N}$, which maps $x \mapsto x + 1$ and the projection functions: for any $k \geq 1$ and $1 \leq i \leq k$, $U_i^k : \mathbb{N}^k \rightarrow \mathbb{N}$ is defined as

$$U_i^k(x_1, \dots, x_k) = x_i.$$

- \mathcal{F} is closed under composition: if $f : \mathbb{N}^k \rightarrow \mathbb{N}$ and $g_1, \dots, g_k : \mathbb{N}^l \rightarrow \mathbb{N}$ are in \mathcal{F} , then $f \circ (g_1, \dots, g_k)$ is in \mathcal{F} .
- Primitive recursion: If $f : \mathbb{N}^k \rightarrow \mathbb{N}$ and $g : \mathbb{N}^{k+2} \rightarrow \mathbb{N}$ are both in \mathcal{F} , then so is $R_{fg} : \mathbb{N}^{k+1} \rightarrow \mathbb{N}$, defined with $R_{fg}(x_1, \dots, x_k, 0) \simeq f(x_1, \dots, x_k)$ and $R_{fg}(x_1, \dots, x_k, x + 1) \simeq g(x_1, \dots, x_k, x, R_{fg}(x_1, \dots, x_k, x))$.

Remark. Since all basic functions are total, every function in \mathcal{F} is total.

Remark. Not every total computable function is primitive recursive. We can show for example that the Ackermann function grows faster than any primitive recursive function.

Definition 3.2.3. The PARTIAL RECURSIVE FUNCTIONS are the smallest collection of partial functions $\mathcal{F} \subseteq \{\mathbb{N}^k \rightarrow \mathbb{N}\}_{k \geq 0}$, which satisfies the axioms of partial recursive functions, with an additional one:

- Minimization: If $f : \mathbb{N}^{k+1} \rightarrow \mathbb{N}$ is in \mathcal{F} , then so is $\mu f : \mathbb{N}^k \rightarrow \mathbb{N}$, with $\mu f(x_1, \dots, x_k)$ equal to the smallest number $n \in \mathbb{N}$ such that $f(x_1, \dots, x_k, n) = 0$ if it exists and $f(x_1, \dots, x_k, m)$ is defined for all $m < n$, and $\mu f(x_1, \dots, x_k)$ undefined otherwise.

Proposition 3.2.4. A partial function $f : \mathbb{N}^k \rightarrow \mathbb{N}$ is computable if and only if there exists a $(k + 1)$ -tape Turing machine such that for all $x_1, \dots, x_k \in \mathbb{N}$ and binary words w_1, \dots, w_k of x_1, \dots, x_k , if we run the Turing machine with w_1, \dots, w_k on the first k tapes, then it halts if and only if $f(x_1, \dots, x_k)$ is defined and it halts with the representation of $f(x_1, \dots, x_k)$ on the last tape, and w_1, \dots, w_k on the first k tapes.

Theorem 3.2.5. The partial recursive functions coincide with the computable partial functions.

Proof. A partial recursive function is computable: We will show that the family of computable functions satisfies the properties of partial computable functions. Since the partial recursive functions are the smallest such family, every partial recursive function is computable.

Clearly, computable partial functions satisfy composition and include the basic functions. Let's show they are closed under primitive recursion. Suppose we have a $(k + 1)$ -tape Turing machine M_f computing f and a $(k + 3)$ -tape Turing machine M_g computing g . We will find a $(k + 4)$ -tape Turing machine computing R_{fg} , which can then be compressed. On the first $k + 1$ tapes, put the arguments to R_{fg} . On the $(k + 2)$ -th tape, put a counter.

3 Teorija izračunljivosti

On the next tape, put the result of R_{fg} on the current iteration, and finally, on the last tape, put the result being computed. The Turing machine then proceeds as follows:

1. initialize tape $k + 2$ to 0
2. use M_f to compute $f(x_1, \dots, x_k)$ and write the result on tape $k + 4$
3. if the numbers on tapes $k + 1$ and $k + 2$ are equal, halt
4. copy tape $k + 4$ to tape $k + 3$
5. apply M_g on tapes $1, \dots, k, k + 2, k + 3$ and write the result on tape $k + 4$
6. increment tape $k + 2$
7. go to step 3

We may similarly construct a machine which performs minimization, which finishes this inclusion.

Before starting the proof of the other inclusion, consider the following. We can encode $\mathbb{N} \times \mathbb{N}$ using \mathbb{N} with the bijection $p(x, y) = \frac{1}{2}(x + y)(x + y + 1) + x$, which is also primitive recursive. The functions $q_1, q_2 : \mathbb{N} \rightarrow \mathbb{N}$ which reverse p (such that $q_1(p(x, y)) = x$ and $q_2(p(x, y)) = y$) are also primitive recursive.

We can also encode finite sequences with $\lceil \cdot \rceil : \mathbb{N}^* \rightarrow \mathbb{N}$, defined as $\lceil n_0, \dots, n_{k-1} \rceil = 2^{n_0} + 2^{n_0+n_1+1} + \dots + 2^{n_0+\dots+n_{k-1}+k-1}$, so that numbers are encoded in a binary sequence with the number of zeros between a pair of ones denoting the sequence. This is clearly a bijection. Additionally, the functions $\sigma : \mathbb{N}^2 \rightarrow \mathbb{N}$ mapping

$$\sigma(\lceil w \rceil, i) = \begin{cases} 0 & i \geq |w| \\ w_i + 1 & i < |w| \end{cases}$$

and $l : \mathbb{N} \rightarrow \mathbb{N}$ mapping

$$l(\lceil w \rceil) = |w|$$

are primitive recursive.

Now we can prove the other inclusion. Suppose $f : \mathbb{N}^k \rightarrow \mathbb{N}$ is computable and that it is computed by a Turing machine M . We assume M is a single-tape machine computing f via representations. Suppose M has a tape alphabet $\Gamma \supseteq \{\square, 0, 1, ,\}$ and state set $Q \supseteq \{\text{start}, \text{halt}\}$. Choose injective functions $r : \Gamma \rightarrow \{\text{odd numbers}\}$ and $s : Q \rightarrow \{\text{even numbers}\}$. Now suppose we are in a configuration C with a finitely many not necessarily blank symbols a_0, \dots, a_{k-1} , the tape head at a_i , and the current state equal to q . Define an encoding

$$\lceil C \rceil = \lceil r(a_0) \dots r(a_{i-1}) s(q) r(a_i) \dots r(a_{k-1}) \rceil.$$

The following functions are primitive recursive:

3.2 Computability of natural numbers

- $\text{step} : \mathbb{N} \rightarrow \mathbb{N}$, mapping $\lceil C \rceil$ to $\lceil C' \rceil$, which is the configuration we obtain by taking one step of M on configuration C . If the input of the function is not a valid configuration, it returns 0.
- $\text{run} : \mathbb{N}^2 \rightarrow \mathbb{N}$, mapping (n, x) to $\text{step}^n(x)$.
- $\text{extract} : \mathbb{N} \rightarrow \mathbb{N}$, mapping $\lceil s(\text{halt})r(w_0) \dots r(w_k) \rceil$ to n if w is a binary representation of n , and any other input to 0.
- $\text{halt?} : \mathbb{N} \rightarrow \mathbb{N}$ mapping $\lceil s(\text{halt})r(w_0) \dots r(w_{k+1}) \rceil$ to 0 and all other inputs to 1.
- $\text{init} : \mathbb{N}^k \rightarrow \mathbb{N}$, mapping (x_1, \dots, x_k) to $\lceil s(\text{start})y_1 \dots y_m \rceil$, where y_i is equal to the result of r on the i -th character of the sequence $(\text{bin}(x_1), \dots, \text{bin}(x_k))$.

Then, f is partial recursive because

$$f(x_1, \dots, x_n) \simeq \text{extract}(\text{run}(\mu(n \mapsto \text{halt?}(\text{run}(n, \text{init}(x_1, \dots, x_k))))), \text{init}(x_1, \dots, x_k)))$$

□

We will consider number representations $\gamma : \mathbb{N} \rightarrow X$. There is a canonical representation of \mathbb{N} , $n \mapsto n$. Given two representations γ_X of X and γ_Y of Y , we can define the product representation

$$\gamma_X \times \gamma_Y(n) = (\gamma_X(n_1), \gamma_Y(n_2)),$$

where $n = p(n_1, n_2)$ for the projection function p .

Given representations ρ_X of X and ρ_Y of Y , we say that a function $f : X \rightarrow Y$ is $(\rho_X \rightarrow \rho_Y)$ -computable if there is a computable partial function which computes between the representations.

As there are only a countable many partial recursive functions of any arity n , we can enumerate them. We will use the label ϕ_i^n to mean the i -th partial recursive function $\mathbb{N}^n \rightarrow \mathbb{N}$ in this enumeration. We require that all functions on the list are computable, and that every partial recursive function occurs on the list, but not necessarily with a unique index. The following definition satisfies these properties. We say that $\phi_e^n(x_1, \dots, x_n)$ is equal to y if there is a Turing machine M with $e = \lceil M \rceil$, which, when run on the input $\text{bin}(x_1), \dots, \text{bin}(x_n)$, halts with $\text{bin}(y)$ on the tape. Otherwise, we say that the expression is undefined.

Note that the function $\mathbb{N} \rightarrow \text{Comp}(\mathbb{N}^n \rightarrow \mathbb{N})$, defined with the expression

$$e \mapsto \phi_e^n$$

is a number representation of the set of all computable functions.

Proposition 3.2.6. *The function $h : \mathbb{N} \rightarrow \mathbb{N}$ defined below is not computable:*

$$h(e) = \begin{cases} \phi_e(e) + 1, & \phi_e(e) \downarrow, \\ 0, & \text{otherwise.} \end{cases}$$

3 Teorija izračunljivosti

Proof. Suppose that h is computable, so there is an index e such that $h = \phi_e$. Then $\phi_e(e) = h(e) = \phi_e(e) + 1$, which is a contradiction. \square

Theorem 3.2.7 (the universal function). *For any $n \geq 1$, the $(n+1)$ -arity function*

$$\phi_u^n(e, x_1, \dots, x_n) \simeq \phi_e^n(x_1, \dots, x_n)$$

exists and is computable.

Proposition 3.2.8. *The unary total function below is not computable.*

$$g(e) = \begin{cases} 1, & \text{if } \phi_e \text{ is a total function,} \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Suppose that g is computable and consider

$$h(x) = \begin{cases} \phi_x(x) + 1, & \text{if } \phi_x \text{ is total,} \\ 0, & \text{otherwise.} \end{cases}$$

Since g is computable, so is h (using the universal function). Then $h = \phi_e$ for some e , but $\phi_e(e) + 1 = h(e) = \phi_e(e)$, which is a contradiction. \square

Theorem 3.2.9 (S-M-N). *For every $n > m > 0$ there exists an $(m+1)$ -ary primitive recursive function $S_n^m : \mathbb{N}^{m+1} \rightarrow \mathbb{N}$ such that for all e, x_1, \dots, x_n :*

$$\phi_{S_n^m(e, x_1, \dots, x_n)}^{n-m}(x_{m+1}, \dots, x_n) \simeq \phi_e^n(x_1, \dots, x_n).$$

Theorem 3.2.10 (Kleene's normal form). *There exists a primitive recursive function $U : \mathbb{N} \rightarrow \mathbb{N}$ and for each $n \geq 1$ an $(n+2)$ -ary primitive recursive function $T^n : \mathbb{N}^{n+2} \rightarrow \mathbb{N}$ such that for all e, x_1, \dots, x_n :*

$$\phi_e^n(x_1, \dots, x_n) \simeq U((\mu T^n)(e, x_1, \dots, x_n)).$$

3.3 Computable and computably enumerable sets

Definition 3.3.1. A subset $A \subseteq \mathbb{N}$ is **COMPUTABLE** if the characteristic function $\chi_A : \mathbb{N} \rightarrow \mathbb{N}$ is computable. It is **COMPUTABLY ENUMERABLE** if the partial characteristic function $\chi_A^p : \mathbb{N} \rightarrow \mathbb{N}$ is computable as a partial function.

Definition 3.3.2. **KLEENE'S HALTING SET** is the set

$$K = \{e \in \mathbb{N} \mid \phi_e(e) \downarrow\}.$$

Proposition 3.3.3. *Kleene's halting set K is not computable.*

3.3 Computable and computably enumerable sets

Proof. Suppose it is. Then so is the partial function

$$h(e) \simeq \begin{cases} \uparrow, & \chi_K(e) = 1, \\ 0, & \chi_K(e) = 0. \end{cases}$$

Therefore there exists an index e such that $h = \phi_e$. Then $\phi_e(e) \downarrow \Leftrightarrow e \in K \Leftrightarrow h(e) \uparrow \Leftrightarrow \phi_e(e) \uparrow$. This is a contradiction. \square

Proposition 3.3.4. *A set $A \subseteq \mathbb{N}$ is computably enumerable if and only if it is the domain of some computable partial function.*

Proof. If a set A is computable enumerable, it is the domain of χ_A^p . Suppose that $A = \text{dom}(f)$ for some computable $f : \mathbb{N} \rightarrow \mathbb{N}$. We can compute χ_A^p by

$$\chi_A^p(n) \simeq \begin{cases} 1, & f(n) \downarrow, \\ \uparrow & \text{otherwise.} \end{cases} \quad \square$$

Proposition 3.3.5. *The halting set K is computably enumerable.*

Proof. It is the domain of the computable partial function $e \mapsto \phi_e(e) = \psi_u(e, e)$. \square

We can enumerate the computably enumerable sets by

$$W_e := \text{dom}(\phi_e).$$

Lemma 3.3.6. *Suppose that $A \subseteq \mathbb{N}$ is a set for which there exists a total computable function $t : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $x \in A$ if and only if there exists $z \in \mathbb{N}$ with $t(x, z) = 0$. Then A is computably enumerable. The reverse also holds.*

Proof. Suppose that A is computably enumerable, so by Kleene's normal form theorem, $x \in A$ if and only if $T(e, x, z) = 0$ for some z , and for $A = W_e$. We can take $t(x, z) = T(e, x, z)$.

As for the reverse, given a computable t , the partial function $\mu t : \mathbb{N} \rightarrow \mathbb{N}$ has A as its domain, so A is computably enumerable. \square

Theorem 3.3.7. *A set $A \subseteq \mathbb{N}$ is computable if and only if both A and \overline{A} are computably enumerable.*

Proof. The left-to-right implication is trivial. Suppose both A and \overline{A} are computably enumerable. By the lemma, there exist total computable $s, s' : \mathbb{N} \rightarrow \mathbb{N}$ such that $x \in A$ if and only if there exists $z \in \mathbb{N}$ for which $s(x, z) = 0$, and similarly $x \notin A$ if and only if there is a number z such that $s'(x, z) = 0$.

3 Teorija izračunljivosti

Then the following function is both computable and total:

$$g := \mu((x, z) \mapsto \min(s(x, z), s'(x, z))).$$

So

$$\chi_A(x) = \begin{cases} 1, & s(x, g(x)) = 0, \\ 0, & \text{otherwise,} \end{cases}$$

is computable. □

Corollary 3.3.8. *The set \overline{K} is not computably enumerable.*

Theorem 3.3.9. *The following are equivalent for a set $A \subseteq \mathbb{N}$:*

- *A is computably enumerable,*
- *$A = \emptyset$ or A is the range of a total computable function,*
- *A is the range of a computable partial function.*

Proof. One to two: Suppose $A \neq \emptyset$, and select any $a_0 \in A$. By the above lemma, there exists a computable $s : \mathbb{N}^2 \rightarrow \mathbb{N}$ satisfying

$$y \in A \Leftrightarrow \exists z. s(y, z) = 0.$$

Using the pairing function $p : \mathbb{N}^2 \rightarrow \mathbb{N}$, we have that A is the range of the total computable

$$x \mapsto \begin{cases} y, & s(y, z) = 0, \\ a_0 & \text{otherwise.} \end{cases}$$

We used $(y, z) = p^{-1}(x)$ above.

Two to three is trivial. Three to one: Suppose A is the range of ϕ_e . Let T and U be as in Kleene's normal form theorem. Then the total function $s : \mathbb{N}^2 \rightarrow \mathbb{N}$ is computable:

$$s(w, y) = \begin{cases} 0 & T(e, x, z) = 0 \wedge w = U(z), \\ 1 & \text{otherwise,} \end{cases}$$

where $y = p(x, z)$. Then $w \in A$ if and only if there exists a y such that $s(w, y) = 0$. □

Theorem 3.3.10. *The following are equivalent for a set $A \subseteq \mathbb{N}$:*

- *A is computable,*
- *$A = \emptyset$ or A is the range of an increasing total computable function,*
- *A is finite or A is the range of a strictly increasing computable total function.*

3.3 Computable and computably enumerable sets

Proof. We will only prove the equivalence of the first and third statement. One to three: Suppose A is computable and infinite. Then A is the image of the function which maps n to the n -th smallest element of A , by searching through $m = 0, 1, 2, \dots$ using χ_A .

Three to one: Suppose A is the range of $f : \mathbb{N} \rightarrow \mathbb{N}$, which is strictly increasing. Then we can compute $\chi_A(n)$ by checking whether n is in the image of $f(x)$ for $x = 0, 1, 2, \dots$, until we find either n or a number larger than it. \square

Corollary 3.3.11. *Every infinite computably enumerable set has an infinite computable subset.*

Proof. Let A be infinite and computably enumerable. Then A is the image of a computable total function f . Define $g(0) = f(0)$ and $g(n+1) = f(m)$ where m is the smallest number such that $f(m) > g(n)$. \square

We define E_e as the domain of ϕ_e , and

$$\mathcal{C} = \{f : \mathbb{N} \rightarrow \mathbb{N} \mid f \text{ is computable}\}.$$

Definition 3.3.12. A subset $B \subseteq \mathcal{C}$ is DECIDABLE if

$$I_B = \{e \in \mathbb{N} \mid \phi_e \in B\}$$

is a computable set. Also, B is SEMIDECIDABLE if I_B is computably enumerable.

Lemma 3.3.13 (Reduction lemma). *If A is reducible to B and B is computable, then so is A . If A is reducible to B and B is computably enumerable, then so is A .*

Theorem 3.3.14 (Rice). *A set $B \subseteq \mathcal{C}$ is decidable if and only if $B = \emptyset$ or $B = \mathcal{C}$.*

Proof. The right-to-left implication is trivial. Left-to-right: Assume B is neither \emptyset nor \mathcal{C} . We'll show that I_B is not computable. Without loss of generality, we assume the everywhere undefined partial function f_\emptyset is in $\mathcal{C} \setminus B$.

Choose some function $g \in B$. Define

$$f(x, y) \simeq \begin{cases} g(y), & x \in K \\ \uparrow, & \text{otherwise} \end{cases}$$

Clearly f is computable. By the s-m-n theorem, there is a total computable $S : \mathbb{N} \rightarrow \mathbb{N}$ such that $\phi_{s(x)}(y) \simeq f(x, y)$. Then for $x \in K$, $\phi_{s(x)}(y) \simeq g(y)$ and for $x \notin K$, $\phi_{s(x)}(y) \downarrow$. Since $g \in B$, we have $s(x) \in I_B$ for any $x \in K$. As f_\emptyset is in $\mathcal{C} \setminus B$, we have $\phi_{s(x)} \notin I_B$ for any $x \notin K$. So s is a reduction of K to I_B . But K is not computable, so I_B isn't either. \square

Definition 3.3.15. For partial functions $\mathbb{N} \rightarrow \mathbb{N}$, $f' \subseteq f$ if the graph of f' is a subset of the graph of f . We say that f is FINITE if its domain is finite.

Theorem 3.3.16 (Rice-Shapiro). *If $B \subseteq \mathcal{C}$ is semidecidable, then for all $f \in \mathcal{C}$, then $f \in B$ if and only if there exists $f' \in B$ such that f' is finite and $f' \subseteq f$.*

Proof. We prove that if either implication of the equivalence fails, then B is not semidecidable.

Suppose that the left-to-right implication fails, so there exists a function $f \in B$ such that all finite subfunctions of f aren't in B . Note that f is necessarily infinite. Because K is computably enumerable, there is some total computable $t : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $x \in K$ if and only if there exists a number z for which $t(x, z) = 0$. Define

$$g(x, z) \simeq \begin{cases} f(z), & \text{if } t(x, y) \neq 0 \text{ for all } 0 \leq y \leq z \\ \uparrow, & \text{otherwise} \end{cases}$$

By the s-m-n theorem, there is a total computable function $s : \mathbb{N} \rightarrow \mathbb{N}$ such that $\phi_{s(x)}(z) \simeq g(x, z)$. If $x \in K$, then $\phi_{s(x)}$ is finite and a subfunction of f , so by assumption $s(x) \notin I_B$ (since $\phi_{s(x)} \notin B$). If $x \notin K$, then $\phi_{s(x)} = f$, so $s(x) \in I_B$ because $f \in B$. Then s is a reduction of \overline{K} to I_B . By the reduction lemma, I_B is not computably enumerable (as \overline{K} isn't).

Now suppose that the right-to-left implication fails. Let f' be a finite subfunction of f such that $f' \in B$ and $f \notin B$. Define a computable partial function

$$g(x, z) \simeq \begin{cases} f(z), & \text{if } z \in \text{dom}(f') \text{ or } x \in K \\ \uparrow, & \text{otherwise} \end{cases}$$

By the s-m-n theorem, there is a total computable $s : \mathbb{N} \rightarrow \mathbb{N}$ such that $\phi_{s(x)}(z) = g(x, z)$. Since $f' \subseteq f$, we have:

- if $x \in K$, then $\phi_{s(x)} = f$, but $f \notin B$, so $s(x) \notin I_B$,
- if $x \notin K$, then $\phi_{s(x)} = f'$, but since $f' \in B$, so $s(x) \in I_B$.

So s is a reduction of \overline{K} to I_B , and I_B can't be computably enumerable. □

3.3.1 Varieties of non-computable sets

Observe that the following are equivalent for a set $A \subseteq \mathbb{N}$:

- A is not computably enumerable,
- for any $W_e \subseteq A$ there exists some $n \in A \setminus W_e$.

A set $A \subseteq \mathbb{N}$ is **PRODUCTIVE** if there exists a computable total function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $e \in \mathbb{N}$, if $W_e \subseteq A$, then $g(e) \in A \setminus W_e$. We call g an **OUTSIDER FINDER**.

Proposition 3.3.17. *Every productive set is not computably enumerable.*

Proposition 3.3.18. *The set \overline{K} is productive.*

3.3 Computable and computably enumerable sets

Proof. We show that $g = \text{id}_{\mathbb{N}}$ is an outsider finder for \overline{K} . Suppose that $W_e \subseteq \overline{K}$. Also suppose that $g(e) = e \notin \overline{K}$. Then $e \in K$, so $\phi_e(e) \downarrow$, which means $e \in W_e \subseteq \overline{K}$, which is a contradiction, meaning that $e \in \overline{K}$. Now if $e \in \overline{K}$, then $e \in \{f \mid f \notin W_f\}$, so $e \notin W_e$. All this implies $g(e) \in \overline{K} \setminus W_e$. \square

Lemma 3.3.19. *If A reduces to B and A is productive, then so is B .*

Proof. Let g be an outsider finder of A . We claim that $f \circ g \circ h$ is an outsider finder for B , where h is defined in the following way. Consider the map $(e, x) \mapsto \phi_e(f(x))$. By the s-m-n theorem, there is a computable $h : \mathbb{N} \rightarrow \mathbb{N}$ for which $\phi_{h(e)}(x) \simeq \phi_e(f(x))$. To prove $f \circ g \circ h$ is an outsider finder for B , suppose $W_e \subseteq B$. Then $f^{-1}(W_e) \subseteq A$, but

$$f^{-1}(W_e) = f^{-1}(\text{dom } \phi_e) = \text{dom } \phi_e \circ f = \text{dom } \phi_{h(e)} = W_{h(e)}.$$

So $f(g(h(e))) \subseteq B \setminus W_e$ because $g(h(e)) \in A \setminus W_{h(e)}$. \square

Definition 3.3.20. A set A is CREATIVE if it is computably enumerable and its complement is productive.

Theorem 3.3.21. *If $\emptyset \subsetneq B \subsetneq \mathcal{C}$ and the everywhere undefined function $f_{\emptyset} \in B$, then I_B is productive.*

Proof. In the proof of Rice's theorem, we reduced K to $\overline{I_B}$, so \overline{K} to I_B . Because \overline{K} is productive, so is I_B . \square

Theorem 3.3.22. *If $\emptyset \subsetneq B \subsetneq \mathcal{C}$ is such that I_B is computably enumerable, then is it creative.*

Theorem 3.3.23. *Every productive set has an infinite computably enumerable subset.*

Proof. Suppose that A is productive with outsider finder g . We will define a function $f : \mathbb{N} \rightarrow \mathbb{N}$ which satisfies $f(0) = e_0$ with $W_{e_0} = \emptyset$, and $f(n+1) = e_{n+1}$ with $W_{e_{n+1}} = W_{e_n} \cup \{g(e_n)\}$. By the s-m-n theorem, there exists a computable total function $h : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\phi_{h(x)}(y) \simeq \begin{cases} 0 & y \in W_x \vee y = g(x), \\ \uparrow & \text{otherwise.} \end{cases}$$

So $W_{h(e)} = W_e \cup \{g(e)\}$, and we can find an f such that $W_{f(0)} = \emptyset$ and $f(n+1) = h(f(n))$.

We then have $g \circ f$, a total computable function with an image that is an infinite subset of A . Then that image is a computably enumerable subset of A . \square

Definition 3.3.24. A set $A \subseteq \mathbb{N}$ is IMMUNE if it is infinite and it has no infinite computably enumerable subset.

Remark. If A is immune, it is not computably enumerable, and not productive.

Definition 3.3.25. A set $A \subseteq \mathbb{N}$ is SIMPLE if it is computably enumerable and its complement is immune.

Theorem 3.3.26 (Post). *There exists a simple set.*

Proof. Consider a partial function $f : \mathbb{N} \rightarrow \mathbb{N}$, defined as follows: $f(e) = \phi_e(z)$ if z is the smallest number such that $\phi_e(x) \downarrow$ for any $x \leq z$ and $\phi_e(z) \geq 2e$, and $f(e)$ is undefined if no such z exists. This is clearly computable, so its image is computably enumerable. Define $A = \text{im } f$. We will prove that A is simple.

When $f(e) = n$, we have $n \geq 2e$, so the numbers $\{0, 1, \dots, 2m-1\}$ can only appear as values $f(e)$ when $e < m$. So for every $m \geq 0$, at least m distinct numbers from that set belong to \overline{A} , meaning that \overline{A} is infinite.

Let B be an infinite computably enumerable set. We know that B must be the image of some total ϕ_e . Then $f(e) \downarrow$, since because B is infinite, it must contain some number larger than $2e$. So indeed $B \not\subseteq A$. \square

3.4 Computation with continuous data

A type 2 Turing machine (T2M for short) with k input tapes, n working tapes and one output tape is a Turing machine with $k + n + 1$ tapes where:

- Input tapes start with no blank symbols, are only infinite in one direction, and their heads are read-only and only move to the right.
- Working tapes start blank on all but a finite number of squares. They are infinite in both directions, and have regular heads.
- The output tape is initially blank and has a write-only head that can only move right.

Formally, a T2M is specified by

- the tape alphabet Γ with $\sqcup \in \Gamma$,
- an input/output alphabet $\Sigma \subseteq \Gamma \setminus \{\sqcup\}$,
- a finite set Q of control states,
- the transition function

$$\delta : Q \times \Sigma^k \times \Gamma^n \rightarrow Q \times \{0, 1\}^k \times \Gamma^n \times \{-1, 0, 1\}^n \times \{\Sigma \cup \sqcup\}.$$

We say that a T2M COMPUTES an infinite word $p \in \Sigma^\omega$ given input $(p^1, \dots, p^k) \in (\Sigma^\omega)^k$ if when we run the machine on input tapes containing p^1, \dots, p^k , it outputs p on the output tape. An ω -word is COMPUTABLE if it is computed by some T2M with not input tapes.

A T2M M COMPUTES a partial function $f : (\Sigma^\omega)^k \rightarrow \Sigma^\omega$ if:

- for any $(p^1, \dots, p^k) \in \text{dom } f$, M computes $f(p^1, \dots, p^k)$ given input (p^1, \dots, p^k) ,
- if M is given (p^1, \dots, p^k) as input, it will compute some $p \in \Sigma^\omega$ only if $(p^1, \dots, p^k) \in \text{dom } f$.

For any recognition machines for an ω -language, we assume distinguished halting states **accept** and **reject**. A single input tape T2M **ACCEPTS** $p \in \Sigma^\omega$ if, when run on input p , it halts in the accepting state. Similarly, it **rejects** p , if it halts in the rejecting state.

Given a word $p \in \Sigma^\omega$, we can define $\text{Prefix}(p) \subseteq \Sigma^*$ as the set of all prefixes of p .

Theorem 3.4.1. *The following are equivalent:*

- p is computable via a T2M,
- $\text{Prefix}(p)$ is decidable,
- $\text{Prefix}(p)$ is semidecidable.

Proof. 1 to 2: Suppose that there is a T2M M computing p . To decide whether w is in $\text{Prefix}(p)$ for some word w , we may run M until it produces $|w|$ characters, then compare that result to w .

2 to 3: Trivial.

3 to 1: Suppose S is an ordinary TM that semidecides $\text{Prefix}(p)$. We build a T2M that computes p as follows. Suppose we have already output n symbols of p . To find the next symbol, for each of the m symbols $b_1, \dots, b_m \in \Sigma$, we run S (in parallel) on the input $p_0 \dots p_{n-1} b_i$. Exactly one of these will halt in the accepting state, so when it does, output that symbol. \square

3.4.1 Topological aspects of computing with ω -words

Theorem 3.4.2. *If $L \subseteq \Sigma^\omega$ is semidecidable, then for any $p \in L$, there exists $n \geq 0$ such that for any infinite word $q \in \Sigma^\omega$, if $q|_n = p|_n$.*

Proof. Let M be a T2M that semidecides L . Consider any $p \in L$, and let n be 1 plus the position of the read head when M enters the accept state if run on p . Note that the read head can only move right, so M could only access the first n characters during its execution. If we give it another ω -word with the same n -prefix, it will take the same actions and accept. \square

Theorem 3.4.3. *If a partial function $f : \Sigma^\omega \rightarrow \Sigma^\omega$ is computable, then for every $p \in \text{dom } f$ and for every $n \geq 0$ there exists an $m \geq 0$ such that for all $q \in \text{dom } f$, if $q|_m = p|_m$, then $f(q)|_n = f(p)|_n$.*

3 Teorija izračunljivosti

Proof. Let M be a T2M that computes f . Consider any $p \in \text{dom } f$ and $n \geq 0$. Let m be 1 plus the position of the input head at the time M writes the n -th symbol to the output tape.

Now consider any $q \in \text{dom } f$ which agrees with p on the first m characters. Then the execution of M on q follows the same steps as on p , so it produces the same first n characters of output. \square

We introduce a topology on Σ^ω , which is just the infinite product topology of Σ (which is discrete). This topology is metrizable for the metric

$$d(p, q) = 2^{-i},$$

where i is the smallest number such that $p_i \neq q_i$. We of course take $d(p, p) = 0$. Also, for a word $p \in \Sigma^\omega$ and number $n \geq 0$, define the CYLINDER SET $\langle p|_n \rangle$, where for a finite word w ,

$$\langle w \rangle = \{q \in \Sigma^\omega \mid q|_{|w|} = w\}.$$

Note that the collection of cylinder sets is a countable basis for Σ^ω .

Remark. Theorem 3.4.2 states: A semidecidable language is an open set.

Remark. Theorem 3.4.3 states: Computable functions are continuous with respect to the subspace topology on $\text{dom } f$.

Proposition 3.4.4. *If $L \subseteq \Sigma^\omega$ is decidable, it is clopen.*

Proof. The complement is semidecidable. \square

Theorem 3.4.5. *An ω -language L is decidable if and only if it is clopen.*

Definition 3.4.6. A subset $Z \subseteq \Sigma^\omega$ is G_δ if it is a countable intersection of open sets.

Theorem 3.4.7. *If a partial function $f : \Sigma^\omega \rightarrow \Sigma^\omega$ is computable, then its domain of definition is a G_δ -subset of Σ^ω .*

Proof. Suppose that M is a T2M which computes f . For every $n \geq 0$, define

$$D_n = \{p \in \Sigma^\omega \mid M \text{ produces } \geq n \text{ output characters when run on } p\}.$$

Note that $\text{dom } f \subseteq D_n$ and that D_n is semidecidable (and hence open). Clearly $\text{dom } f$ is the intersection of all D_n . \square

Theorem 3.4.8. *The topological space Σ^ω is compact.*

Proof. Tychonoff. \square

Corollary 3.4.9. *The compact subsets of Σ^ω are exactly the closed sets.*

Lemma 3.4.10. *Every clopen set is decidable.*

Proof. Let L be a clopen set. Since the cylinder sets form a basis and L is open, we have

$$L = \bigcup \{ \langle w \rangle \mid w \in \Sigma^*, \langle w \rangle \subseteq L \}.$$

So this family of cylinders is an open cover of L . Because L is closed, it is compact, so there is a finite subcover

$$L = \langle w_1 \rangle \cup \dots \cup \langle w_k \rangle.$$

We can decide L by checking whether the prefix of a word is equal to any of w_1, \dots, w_k . \square

3.4.2 Computing with real numbers

We can represent real numbers as ω -words via infinite decimal expansions (or representations in other bases), so with the alphabet $\{0, 1, \dots, 9, -, .\}$, with at most one $-$ at the start, and exactly one decimal point. The problem is that this is a poor representation, as many useful algorithms cannot be written with it.

Definition 3.4.11. A TYPE 2 REPRESENTATION of a set X is a surjective partial function $\gamma : \Sigma^\omega \rightarrow X$. We say that p is a NAME for an element $x \in X$ if $\gamma(p) = x$, and that x is COMPUTABLE if it has a computable name.

We now introduce the Cauchy representation of \mathbb{R} . First, give a type 1 representation of the dyadic rationals \mathbb{Q}_d , which we represent by a finite word $\pm d_{m-1} \dots d_0 . d_{-1} \dots d_{-n}$ with every $d_i \in \{0, 1\}$ and $m, n \geq 0$. For a word u representing a dyadic rational, define $q_d(u)$ as its rational value, interpreted as we usually interpret binary numbers.

We can now define the Cauchy representation $\gamma_c : \Sigma_c^\omega \rightarrow \mathbb{R}$ for $\Sigma_c = \{-, ., 0, 1, ;\}$. The domain of γ_c consists of ω -words of the form

$$p = u_0; u_1; u_2; \dots,$$

where $u_i \in \text{dom } q_d$ and the sequence $(q_d(u_i))_i$ is a fast Cauchy sequence, i.e. it satisfies

$$|q_d(u_m) - q_d(u_n)| \leq \frac{1}{2^n}$$

for all $m \geq n \geq 0$. For such a name p , we define $\gamma_c(p) = \lim q_d(u_n)$.

Proposition 3.4.12. *The representation γ_c is surjective, and if $p \in \text{dom } \gamma_c$ is as above, then for any $n \geq 0$, $|q_d(u_n) - \gamma_c(p)| \leq 2^{-n}$.*

Proposition 3.4.13. *A real number is γ_c -computable if and only if it is computable with respect to the decimal or binary representation.*

Definition 3.4.14. A name p is CLOSE if for every $n \geq 0$, we have $|q_d(u_n) - \gamma_c(p)| \leq 2^{-(n+1)}$.

Lemma 3.4.15. *Every $x \in \mathbb{R}$ has a close name. If p is a close name for x , then for every $n \geq 0$, every x' with $|x' - x| < 2^{-(n+1)}$ has a name of the form*

$$u_0; u_1; u_2; \dots; u_n; u'_{n+1}; u'_{n+2}; \dots$$

Theorem 3.4.16 (continuity theorem). *If $f : \mathbb{R} \rightarrow \mathbb{R}$ is computable with respect to γ_c , then f is continuous on its domain.*

Proof. Suppose f is computable, so it has a realiser $g : \Sigma_c^\omega \rightarrow \Sigma_c^\omega$. Let $x \in \text{dom } f$ and $\varepsilon > 0$. We are searching for a suitable δ . Let $p = u_0; u_1; \dots$ be a close name for x , and let $r = g(p)$. Since g is a realiser for f , r is a name for $f(x)$. Therefore r has the form $r = v_0; v_1; \dots$

Let N be such that $2^{-N} < \varepsilon/2$. We have $|q_d(v_N) - f(x)| \leq 2^{-N} < \varepsilon/2$ by one of the preceding propositions. Let n be the length of the string $v_0; v_1; \dots; v_N$. By the topological continuity theorem, there exists an $m \geq 0$ such that for all $p' \in \text{dom } g$, if $p'|_m = p|_m$, then $g(p')|_n = v_0; v_1; \dots; v_N$. Now take $M \geq 0$ such that the prefix $u_0; u_1; \dots; u_M$ of p has length $m' \geq n$, and define $\delta = 2^{-(M+1)}$.

Then for $x' \in \text{dom } f$ with $|x' - x| < \delta$, we have a name for x' of the form $p' = u_0; \dots; u_M; u'_{M+1}; \dots$, since p is a close name for x . Since $x' \in \text{dom } f$, $g(p')$ is a name for $f(x')$, and $g(p')|_n = v_0; v_1; \dots; v_N$, so $|q_d(v_N) - f(x')| \leq 2^{-N} < \varepsilon/2$. Then $|f(x) - f(x')| < \varepsilon$. \square

3.5 Algorithmic information theory

A computable partial function $u : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is UNIVERSAL if for any computable partial function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ there exists a word $v_f \in \{0, 1\}^*$ such that for all $w \in \{0, 1\}^*$ we have $f(w) \simeq u(v_f w)$.

Proposition 3.5.1. *A universal computable partial function exists.*

Given a universal function u , the KOLMOGOROV COMPLEXITY $C_u(w)$ of a word w is

$$C_u(w) = \min\{|v| \mid u(v) = w\}.$$

Proposition 3.5.2. *Suppose u, u' are universal computable functions. Then for all words w , $C_{u'}(w) \leq C_u(w) + C$ for some constant C , equal for all words.*

Proposition 3.5.3. *For any $n \geq 0$, there exists a word w of length n such that $C_u(w) \geq n$.*

Proof. There are only at most 2^{n-1} words with complexity $< n$, but there are 2^n words of length n . \square

Definition 3.5.4. An infinite word $p \in \{0, 1\}^\omega$ is **K-INCOMPRESSIBLE** if for all $n \geq 0$, we have $C(p|_n) \geq n - O(1)$.

Theorem 3.5.5. No infinite sequence is *K-incompressible*.

Proof. Define a map

$$\lceil w \rceil = \sum_{i=0}^{|w|-1} w_i 2^i + 2^{|w|} - 1$$

which has an inverse word $\lfloor \cdot \rfloor : \mathbb{N} \rightarrow \{0, 1\}^*$. Let $p \in \{0, 1\}^\omega$. Consider any $d \geq 0$ and let $m = \lceil p|_d \rceil$. Define $h : \{0, 1\}^* \rightarrow \{0, 1\}^*$ by $h(w) = \text{word}(|w|)w$. Note that

$$p_0 p_1 \dots p_{d-1} p_d \dots p_{d+m-1} = p|_{d+m} = h(p_d \dots p_{d+m-1}) = u(v_h p_d \dots p_{d+m-1}),$$

so $C_u(p|_{d+m}) \leq |v_h| + \lceil p|_d \rceil$. \square

Definition 3.5.6. A partial function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is **PREFIX-FREE** if $\text{dom } f \subseteq \{0, 1\}^*$ satisfies the following property: for all $w, w' \in \text{dom } f$, w is not a proper prefix of w' .

Definition 3.5.7. A computable prefix-free function $u : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is **UNIVERSAL** if for any computable prefix-free function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ there exists $v_f \in \{0, 1\}^*$ such that for all w , $f(w) \simeq u(v_f w)$.

Definition 3.5.8. The **PREFIX-FREE COMPLEXITY** $K(w)$ is defined as

$$K(w) = \min\{|v| \mid u(v) = w\}$$

for a universal prefix-free function u .

Proposition 3.5.9. For any $d > 1$, there exists a constant C such that $K(w) \leq d|w| + C$ for any word w .

Proof. Outline. For any large N , we work with an encoding of $\{0, 1\}^*$ as words Σ^* where $|\Sigma| = 2^N - 1$. We map w to $\lceil w \rceil$, then represent Σ with binary words of length N . We are left with one non-represented word, which is 0^N without loss of generality. Then we assign to w the representation $\langle \lceil w \rceil \rangle$ followed by N zeros. \square

Definition 3.5.10. An infinite word p is **PREFIX-FREE INCOMPRESSIBLE** if there exists a constant c such that for all n , $K(p|_n) \geq n - c$.

Definition 3.5.11. Chaitin's halting probability Ω is the probability that u halts if given an input generated randomly by throwing coins, so

$$\Omega = \sum_{w \in \text{dom } u} 2^{-|w|}.$$

Then take p^Ω as the binary representation of Ω . If there are multiple options, choose the one with infinite ones.

Theorem 3.5.12. *Chaitin's halting probability p^Ω is prefix-free incompressible.*

Proof. Note that

$$\Omega = \sum_{i=0}^{\infty} p_i^\Omega 2^{-(i+1)}.$$

We will define a computable function $f : \{0,1\}^* \rightarrow \{0,1\}^*$ satisfying $f(p^\Omega|_n) \downarrow$ and $K(f(p^\Omega|_n)) > n$ for all n . If we do, then there is a number c such that for all $n \geq 0$,

$$K(f(p^\Omega|_n)) \leq K(p^\Omega|_n) + c,$$

so $K(p^\Omega|_n) > n - c$.

Let w_0, w_1, \dots be a computable enumeration of $\text{dom } u$ without repetitions. Given an input word $p_0 p_1 \dots p_{n-1}$, the algorithm of f finds the smallest $N \geq 0$ (if it exists) with the property

$$\sum_{i=0}^{N-1} 2^{-|w_i|} \geq \sum_{i=0}^{n-1} p_i 2^{-(i+1)}.$$

Then the algorithm returns the first $v \in \{0,1\}^*$ such that $v \notin \{u(w_0), \dots, u(w_{N-1})\}$. Define $f(p_0 \dots p_{n-1}) = v$.

Clearly $f(p^\Omega|_n)$ is defined for any $n \in \mathbb{N}$, as we have chosen a representation of p^Ω with infinite ones. From

$$\sum_{i=0}^{n-1} p_i^\Omega 2^{-(i+1)} \leq \sum_{i=0}^{N-1} 2^{-|w_i|} < \Omega < \sum_{i=0}^{n-1} p_i^\Omega 2^{-(i+1)} + 2^{-n}$$

we have

$$\Omega - \sum_{i=0}^{n-1} 2^{-|w_i|} < 2^{-n},$$

so every word w_i for $i \geq N$ is such that $|w_i| > n$. This means that for every finite word $v \notin \{u(w_0), \dots, u(w_{N-1})\}$, the smallest $w \in \text{dom } u$ with $u(w) = v$ has $|w| > n$ or $K(v) > n$. \square

3.6 Algorithmic randomness

The law of large numbers states that for a sequence, generated randomly by coin tosses,

$$P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} q_i = \frac{1}{2}\right) = 1.$$

We say that a sequence q satisfies the law of large numbers if

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} q_i = \frac{1}{2}.$$

If q does not satisfy the law, then there exists a $\delta > 0$ such that $|m_n(q) - \frac{1}{2}| < \delta$ for infinitely many n , where

$$m_n(q) = \frac{1}{n} \sum_{i=0}^{n-1} q_i.$$

Given δ , we then have the following non-randomness test. Define T_n as the set of all sequences $q \in \{0,1\}^\omega$ for which there are at least n different values of i where $|m_n(q) - \frac{1}{2}| > \delta$. Note that we can semidecide T_n .

Definition 3.6.1. A NAIVE NON-RANDOMNESS TEST is a sequence $(T_n)_n$ of subsets of $\{0,1\}^\omega$ such that

- every T_n is open,
- $\lim \lambda(T_n) = 0$ where $\lambda(T_n)$ is the probability that a fair randomly generated sequence lands in T_n .

Then $q \in 2^\omega$ SATISFIES $(T_n)_n$ if $q \in \bigcap T_n$.

Definition 3.6.2. A MARTIN-LÖF NON-RANDOMNESS TEST is a sequence $(T_n)_n$ of subsets of 2^ω such that

- $(T_n)_n$ is a computable sequence of computably open sets,
- $\lim \lambda(T_n) = 0$ with a computable rate of convergence.

A sequence q is MARTIN-LÖF RANDOM if it fails every ML test.

Proposition 3.6.3. *Computable sequences are not ML random.*

Proof. Let p be a computable sequence. Then $(\langle p|_n \rangle)_n$ is an ML test which p satisfies. \square

Definition 3.6.4. An ML test $(T_n^u)_n$ is UNIVERSAL if any non-ML random p satisfies $(T_n^u)_n$.

It holds that every open set $U \subseteq 2^\omega$ can be written as a disjoint union of cylinder sets $\langle w_i \rangle$ for some $(w_i)_{i \in I}$. Define $\lambda(\langle w \rangle) = 2^{-|w|}$. Then the probability of an open set is

$$\lambda(U) = \sum_{i \in I} \lambda(w_i)$$

for the disjoint union above. We can prove that $\lambda(U)$ is well-defined.

We represent open sets as infinite sequences w_0, w_1, \dots where every w_i is either a word in 2^* or the symbol \emptyset . We say that an open set is COMPUTABLE if it is represented by some computable infinite sequence w_0, w_1, \dots . Note that this definition coincides with the semidecidable sets.

Then a sequence of open sets $(T_n)_n$ is represented by a sequence of sequences, $T_i \sim (w_{ij})_j$. We say that $(T_n)_n$ is a COMPUTABLE SEQUENCE OF COMPUTABLE OPENS if the single sequence representing w_{ij} above, using the pairing function, is computable.

3 Teorija izračunljivosti

A computable total function $r : \mathbb{N} \rightarrow \mathbb{N}$ is a **COMPUTABLE RATE OF CONVERGENCE** for a Cauchy sequence $(x_n)_n \subseteq [0, 1]$ converging to 0 if for any n and $m \geq r(n)$, we have $x_m \leq 2^{-n}$.

Theorem 3.6.5 (Martin-Löf). *There exists a universal ML test.*

Proof. Outline. We can show the following:

- There is a computable function from \mathbb{N} to the representations of computable sequences of computably open sets, and it finds some representation of any sequence.
- There is a function from the computable sequences of computably open sets to ML tests, which preserves rapid ML tests in its domain (an ML test is rapid if $\lambda(T_n) \leq 2^{-n}$).
- There is a function from ML tests to rapid decreasing ML tests which preserves the set of sequences that satisfy the test (an ML test is decreasing if $T_{n+1} \subseteq T_n$ for all n).

So a sequence p is ML random if and only if it fails every rapid decreasing test. From all three points above, we have a computable enumeration of the rapid decreasing ML tests. It is surjective in the sense that for every rapid decreasing ML test, at least one representation is found. Let $(T_n^0)_n, (T_n^1)_n, \dots$ be this enumeration. Take

$$T_n^u = \bigcup_{i=0}^{\infty} T_{n+i+1}^i.$$

This is clearly a rapid decreasing sequence, and it is computable. □

Corollary 3.6.6. *The set R_{ML} of all ML random sequences is a countable union of closed sets (F_σ) . Additionally, $\lambda(R_{ML}) = 1$ and $|R_{ML}| \geq 2^{\aleph_0}$.*

Proof. The complement of R_{ML} is exactly the intersection of all T_n^u . □

Theorem 3.6.7. *The following are equivalent for any infinite sequence $q \in 2^\omega$.*

- q is ML random,
- q is prefix-free incompressible.

4 Uvod v funkcionalno analizo

4.1 Normirani in Banachovi prostori

Definicija 4.1.1. Naj bo X vektorski prostor nad poljem $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. Preslikava $\|\cdot\| : X \rightarrow \mathbb{R}$ je NORMA, če velja:

- $\|x\| \geq 0$,
- $\|x\| = 0 \Leftrightarrow x = 0$,
- $\|\lambda x\| = |\lambda| \|x\|$,
- $\|x + y\| \leq \|x\| + \|y\|$.

Opomba. Velja $|\|x\| - \|y\|| \leq \|x - y\|$, iz česar sledi, da je norma zvezna (celo Lipschitzova za $L = 1$).

Norma porodi metriko $d(x, y) = \|x - y\|$ na prostoru X , ki je invariantna na translacije, in za katero velja

$$d(\lambda x, \lambda y) = |\lambda| d(x, y).$$

Zaprto kroglo radija r s središčem v točki x označimo z $B(x, r)$, odprto kroglo pa z $\mathring{B}(x, r)$. Zaradi zveznosti norme je zaprtje odprte krogle natanko pripadajoča zaprta krogla.

Definicija 4.1.2. Normiran prostor je BANACHOV, če je poln za inducirano metriko.

Trditev 4.1.3. Seštevanje in množenje vektorjev s skalarjem sta zvezni operaciji.

Definicija 4.1.4. Algebra A je NORMIRANA ALGEBRA, če je normiran vektorski prostor in če velja $\|xy\| \leq \|x\| \|y\|$. Če ima normirana algebra enoto, zahtevamo še $\|e\| = 1$.

Primer. Naj bo X Hausdorffov topološki prostor in $\mathcal{C}_b(X)$ množica zveznih omejenih funkcij $X \rightarrow \mathbb{F}$. Če jo opremimo s supremum normo, postane normirana algebra za seštevanje in množenje po točkah. Preverimo lahko, da je celo Banachov prostor.

Posledica 4.1.5. Če je X kompakten Hausdorffov prostor, je $\mathcal{C}(X)$ Banachova algebra.

Trditev 4.1.6. Naj bo X normiran prostor in Y (vektorski) podprostor v X . Veljata naslednji točki.

- Če je Y poln, potem je Y zaprt v X .
- Če je X Banachov prostor, potem je Y Banachov natanko tedaj, ko je Y zaprt v X .

Dokaz. Prva točka: Naj bo $y \in \overline{Y}$. Obstaja zaporedje $(y_n)_n$ v Y , ki konvergira k y . To zaporedje je Cauchyjevo, torej ima limito, ki je enaka y . Sledi $y \in Y$, zato $Y = \overline{Y}$.

Druga točka: V desno smo ravno dokazali. V levo naj bo $(y_n)_n$ Cauchyjevo zaporedje v Y . Potem je Cauchyjevo tudi v X , kjer ima limito, saj je X poln. Ker je Y zaprt, je $y \in Y$, torej $y_n \rightarrow y$ v Y . \square

Vprašanje 1. Dokaži: če je X Banachov prostor in $Y \leq X$, je Y Banachov natanko tedaj, ko je zaprt v X .

Primer. Naj bo X lokalno kompakten Hausdorffov prostor ter $\mathcal{C}_0(X)$ množica vseh funkcij v $\mathcal{C}(X)$, za katere za vsak $\varepsilon > 0$ obstaja kompaktna $K_\varepsilon \subseteq X$, da je $|f|$ zunaj K_ε strogo manjša od ε .

Pri $X = \mathbb{R}$ so to natanko vse funkcije, katerih limita v obeh neskončnostih je enaka 0, pri $X = \mathbb{N}$ pa je to natanko prostor c_0 zaporedij, ki konvergirajo k 0.

Dokažemo lahko, da je $\mathcal{C}_0(X)$ zaprt dvostranski ideal v $\mathcal{C}_b(X)$ in zato Banachova algebra.

Primer. Množica c vseh konvergentnih zaporedij je Banachov prostor za supremum normo.

4.1.1 Napolnitve normiranih prostorov

Naj bo X normiran prostor, ki ni poln, in naj bo \tilde{X} množica vseh Cauchyjevih zaporedij v X . To je vektorski prostor za operacije po komponentah. Definiramo

$$\|(x_n)_n\| = \lim_{n \rightarrow \infty} \|x_n\|.$$

Ta izraz je dobro definiran, saj velja $\|x_n\| - \|y_n\| \leq \|x_n - y_n\|$, torej je zaporedje norm Cauchyjevo in konvergira v \mathbb{R} . To pa ni norma, ker obstaja veliko zaporedij z limito norm enako 0. Na \tilde{X} zato vpeljemo ekvivalenčno relacijo

$$(x_n)_n \sim (y_n)_n \Leftrightarrow x_n - y_n \xrightarrow{n \rightarrow \infty} 0.$$

Sedaj definiramo $\hat{X} = \tilde{X} / \sim$. V \hat{X} potem vpeljemo operaciji seštevanja in množenja s skalarjem, ki delujeta na predstavnikih, ter podobno vpeljemo normo.

Izrek 4.1.7. *Prostor $(\hat{X}, \|\cdot\|)$ je Banachov in vsebuje X kot gost podprostor.*

Dokaz. Dokaz o polnosti izpustimo. Definiramo vložitev $j : X \rightarrow \hat{X}$ s predpisom $x \mapsto [(x)_n]$. To je očitno linearna preslikava, za katero velja $\|j(x)\| = \|x\|$, torej je tudi izometrija. Pokazali bomo, da je $j(X)$ gost podprostor v \hat{X} .

Naj bo $[(x_n)_n] \in \hat{X}$. Za poljuben $\varepsilon > 0$ obstaja n_ε , da za vse $n, m \geq n_\varepsilon$ velja $\|x_n - x_m\| < \varepsilon$. Za $m = n_\varepsilon$ dobimo $j(x_{n_\varepsilon}) = [(x_{n_\varepsilon})_n]$, in velja

$$\|j(x_{n_\varepsilon}) - [(x_n)_n]\| = \lim_{n \rightarrow \infty} \|x_{n_\varepsilon} - x_n\| \leq \varepsilon. \quad \square$$

Vprašanje 2. Definiraj napolnitev normiranega prostora. Pokaži, da napolnitev vsebuje originalen prostor kot gost podprostor.

Posledica 4.1.8. *Prostor X je Banachov natanko tedaj, ko je $j(X) = \hat{X}$.*

Dokaz. Ideja. Izometrije ohranjajo polnost, polni podprostori so zaprti. Če so gosti, so enaki celoti. \square

4.1.2 Osnovne konstrukcije

Naj bo X vektorski prostor nad \mathbb{F} . Normi $\|\cdot\|_1$ in $\|\cdot\|_2$ sta EKVIVALENTNI, če obstajata $\alpha, \beta > 0$, da za vse $x \in X$ velja

$$\alpha \|x\|_1 \leq \|x\|_2 \leq \beta \|x\|_1.$$

Topologiji, ki jih normi porodita, sta enaki, zato je identiteta $(X, \|\cdot\|_1) \rightarrow (X, \|\cdot\|_2)$ linearni homeomorfizem.

Definicija 4.1.9. Normirana prostora X in Y sta IZOMORFNA, če obstaja linearni homeomorfizem med njima.

Vprašanje 3. Kdaj sta normi ekvivalentni? Kdaj sta normirana prostora izomorfna?

Če sta normi ekvivalentni, je $(X, \|\cdot\|_1)$ Banachov natanko tedaj, ko je $(X, \|\cdot\|_2)$ Banachov. Če je $Y \subseteq X$ podprostor in X normiran, je tudi Y normiran, če normo zožimo na Y . Vložitev Y v X je izometrija.

Lema 4.1.10. Če je $Y \subseteq X$ podprostor in X normiran, je \overline{Y} podprostor.

Dokaz. Naj bosta $x, y \in \overline{Y}$ in $\alpha, \beta \in \mathbb{F}$. Potem obstajata zaporedji $x_n \rightarrow x$ in $y_n \rightarrow y$. Velja $\alpha x_n + \beta y_n \rightarrow \alpha x + \beta y$. \square

Če je X normiran in $Y \leq X$, lahko na X vpeljemo relacijo $x_1 \sim x_2 \Leftrightarrow x_1 - x_2 \in Y$. V kvocientni prostor vpeljemo

$$\|x + Y\| = \inf\{\|x + y\| \mid y \in Y\}.$$

Trditev 4.1.11. Naj bo X normiran prostor in $Y \leq X$.

- $\|\cdot\|$ je polnorma na X/Y .
- $\|\cdot\|$ je norma na X/Y natanko tedaj, ko je Y zaprt v X .
- Če je X Banachov, je kvocient Banachov.

Dokaz.

- Točka je le vprašanje preverjanja; dokaz izpustimo.
- S sklicem na prvo točko moramo dokazati le, da je $\|x + Y\| = 0 \Leftrightarrow x \in Y$. Če je $\|x + Y\| = 0$, je $d(x, Y) = 0$, ker pa je Y zaprta, to pomeni $x \in Y$. Podobno v obratno smer.
- Naj bo $(x_n + Y)_n$ Cauchyjevo zaporedje v X/Y . Poiskali bomo podzaporedje $(x_{n_k} + Y)_k$, ki bo konvergiral. Ker je prvotno zaporedje Cauchyjevo, bo tudi konvergiral k isti limiti.

Vemo, da za poljuben $\varepsilon > 0$ obstaja $n_\varepsilon \in \mathbb{N}$, za katerega za vse $n, m \geq n_\varepsilon$ velja $\|x_n - x_m + Y\| < \varepsilon$. Sedaj induktivno konstruiramo podzaporedje $(x_{n_k} + Y)_k$, da

za vsak k velja $\|x_{n_{k+1}} - x_{n_k} + Y\| < 2^{-k}$. Ko to zaporedje imamo, po definiciji infimuma obstaja tak $y_k \in Y$, da je $\|x_{n_{k+1}} - x_{n_k} + y_k\| < 2^{-k}$. Definiramo $z_1 = 0$ in $z_{k+1} = z_k + y_k \in Y$. Tedaj

$$\|(x_{n_{k+1}} + z_{k+1}) - (x_{n_k} + z_k)\| = \|x_{n_{k+1}} - x_{n_k} + y_k\| < 2^{-k}.$$

Za $w_k = x_{n_k} + z_k$ velja $\|w_{k+1} - w_k\| < 2^{-k}$, hkrati pa je

$$\|w_{m+k} - w_m\| = \left\| \sum_{i=0}^{k-1} w_{m+i+1} - w_{m+i} \right\| \leq \sum_{i=0}^{k-1} \|w_{m+i+1} - w_{m+i}\| < \sum_{i=0}^{k-1} 2^{-m-i}$$

kar je manjše od 2^{1-m} . Sedaj je $(w_m)_m$ Cauchyjevo v X , torej obstaja limita $x \in X$. Zato je

$$\|(x_{n_i} + Y) - (x + Y)\| \leq \|x_{n_i} - x + z_i\| = \|w_i - x\| \rightarrow 0. \quad \square$$

Vprašanje 4. Definiraj normo na kvocientnem prostoru. Kaj mora veljati, da bo res norma? Kdaj je kvocient Banachov? Dokaži.

Trditev 4.1.12. *Naj bo Y zaprt podprostor normiranega prostora X . Tedaj je X/Y Banachov natanko tedaj, ko sta tako Y kot X/Y Banachova.*

Dokaz. V desno smo ravno dokazali. V levo: Naj bo $(x_n)_n$ Cauchyjevo zaporedje v X . Vemo, da je $\|x + Y\| \leq \|x\|$, torej je tudi $(x_n + Y)_n$ Cauchyjevo zaporedje in ima limito $x + Y$.

Za vsak $n \in \mathbb{N}$ obstaja tak y_n , da je

$$\|x_n - x + y_n\| < \|x_n - x + Y\| + \frac{1}{n}.$$

Ker velja

$$\|y_n - y_m\| \leq \|y_n + x_n - x\| + \|y_m + x_m - x\| + \|x_m - x_n\|,$$

je $(y_n)_n$ Cauchyjevo, torej ima limito $y \in Y$. Sledi $\lim x_n = x - y$. \square

Vprašanje 5. Pokaži: če sta Y in X/Y Banachova, je X Banachov.

Posledica 4.1.13. *Vsak končnorazsežen normiran prostor je Banachov.*

Dokaz. Indukcija na $d = \dim X$. Za $d = 1$ izberimo $x \in X$ z $\|x\| = 1$. Tedaj za vsak $y \in X$ velja $y = \lambda x$ in $\|y\| = |\lambda|$. Naj bo $(y_n)_n$ Cauchyjevo v X . Potem je $y_n = \lambda_n x$ in $\|y_n - y_m\| = |\lambda_n - \lambda_m|$, zato je $(\lambda_n)_n$ Cauchyjevo v \mathbb{F} in ima limito λ . Seveda $\lambda_n x \rightarrow \lambda x$.

Recimo, da so vsi končnorazsežni normirani prostori dimenzije $d - 1$ ali manj polni. Naj bo $Y \leq X$ poljuben enorazsežen prostor. Po predpostavki sta Y in X/Y Banachova. \square

Vprašanje 6. Pokaži: vsak končnorazsežen normiran prostor je Banachov.

Produkt $X \times Y$ lahko opremimo z eno od spodnjih norm:

- $\|(x, y)\|_\infty = \max\{\|x\|_X, \|y\|_Y\}$
- $\|(x, y)\|_p = (\|x\|_X^p + \|y\|_Y^p)^{1/p}$

Produkt je Banachov natanko tedaj, ko sta X in Y Banachova.

4.2 Linearni funkcionali

Izrek 4.2.1. Naj bo $T : X \rightarrow Y$ linearna preslikava med normiranimi prostoroma. Naslednje trditve so ekvivalentne:

- T je zvezna na X
- T je zvezna v $x_0 \in X$
- T je zvezna v 0
- obstaja $C > 0$, da za vse $x \in X$ velja $\|Tx\| \leq C \|x\|$
- T je Lipschitzova
- T je enakomerno zvezna

Vprašanje 7. Karakteriziraj zveznost linearnih preslikav med normiranimi prostori.

Za omejen operator $T : X \rightarrow Y$ definiramo

$$\|T\| = \inf\{C > 0 \mid \forall x. \|Tx\| \leq C \|x\|\}.$$

Ta infimum obstaja in je dejansko minimum. Potem velja

$$\|Tx\| \leq \|T\| \|x\|$$

za vsak $x \in X$. Velja

$$\|T\| = \sup_{\|x\|=1} \|Tx\| = \sup_{\|x\|\leq 1} \|Tx\| = \sup_{\|x\|<1} \|Tx\|.$$

Množico vseh omejenih linearnih operatorjev $X \rightarrow Y$ označimo z $B(X, Y)$, in jo opremo z zgornjo normo.

Vprašanje 8. Kako definiraš normo na (omejenem) linearnem operatorju? Povej še vsaj eno izražavo.

Trditev 4.2.2. Naj bodo X, Y, Z normirani prostori.

- $B(X, Y)$ je normiran prostor.

- Če $T \in B(X, Y)$ in $S \in B(Y, Z)$, potem je $ST \in B(X, Z)$ in $\|ST\| \leq \|S\| \|T\|$.

Dokaz. Samo druga točka. Ker je omejenost ekvivalentna zveznosti, je kompozitum v $B(X, Z)$. Za vsak $x \in X$ velja

$$\|STx\| \leq \|S\| \|Tx\| \leq \|S\| \|T\| \|x\|. \quad \square$$

Definicija 4.2.3. DUALNI PROSTOR prostora X je $X^* = B(X, \mathbb{F})$.

Izrek 4.2.4. Naj bo X normiran in Y Banachov prostor. Tedaj je $B(X, Y)$ Banachov prostor.

Dokaz. Naj bo $(T_n)_n$ Cauchyjevo v $B(X, Y)$. Izberimo $\varepsilon > 0$. Obstaja n_ε , da za $m, n \geq n_\varepsilon$ velja

$$\|(T_n - T_m)x\| \leq \|T_n - T_m\| \|x\| < \varepsilon \|x\|$$

za poljuben $x \in X$. Zaporedje $(T_n x)_n$ je Cauchyjevo, zato obstaja $Tx = \lim T_n x \in Y$. S tem dobimo po točkah definiran operator $T : X \rightarrow Y$.

Enostavno se prepričamo, da je T linearen. Ker je $|\|T_n\| - \|T_m\|| \leq \|T_n - T_m\|$, je tudi zaporedje norm Cauchyjevo, in zato omejeno. Torej je $\|T_n x\| \leq \|T_n\| \|x\| \leq M \|x\|$, in je T omejen.

Za konec za poljuben $x \in X$ in $n, m \geq n_\varepsilon$ dobimo $\|T_n x - T_m x\| < \varepsilon \|x\|$, in če vzamemo limito $n \rightarrow \infty$,

$$\|Tx - T_m x\| \leq \varepsilon \|x\|.$$

Torej je $\|T - T_m\| \leq \varepsilon$ za $m \geq n_\varepsilon$ in zato $T_m \rightarrow T$. \square

Vprašanje 9. Pokaži: če je Y Banachov, je $B(X, Y)$ Banachov.

Posledica 4.2.5. Dualni prostor je vedno Banachov.

Izrek 4.2.6. Naj bo X normiran prostor in $Y \leq X$. Naj bo $T : Y \rightarrow Z$ omejen operator in Z poln. Tedaj obstaja natanko en omejen linearen operator $S : \overline{Y} \rightarrow Z$, da je $S|_Y = T$. Velja še $\|S\| = \|T\|$.

Dokaz. Naj bo $x \in \overline{Y}$. Radi bi definirali Sx . Obstaja zaporedje $(x_n)_n$ v Y , da bo $x_n \rightarrow x$. Definiramo $Sx = \lim Tx_n$.

Če je tudi $(x'_n)_n$ zaporedje, ki konvergira k x , je

$$\lim(Tx_n - Tx'_n) = \lim T(x_n - x'_n) = 0,$$

ker je T zvezen. Torej je S dobro definiran. Očitno je tudi linearen, in velja $S|_Y = T$. Za enoličnost predpostavimo, da je tudi S' tak operator. Potem za $x \in \overline{Y}$ velja $Sx_n = S'x_n$, in sta limiti Sx in $S'x$ posledično tudi enaki.

Velja

$$\|Sx\| = \|\lim T x_n\| = \lim \|T x_n\| = \lim \|T\| \|x_n\| = \|T\| \|x\|,$$

torej je S omejen in $\|S\| \leq \|T\|$. Obrat je očitno. \square

Vprašanje 10. Dokaži: linearen operator $T : Y \rightarrow Z$, ki slika v poln prostor, lahko enolično razširimo do operatorja $\bar{Y} \rightarrow Z$.

Posledica 4.2.7. Naj bosta $S, T : X \rightarrow Y$ omejena operatorja, ki se ujemata na gostem podprostoru. Potem je $S = T$.

Posledica 4.2.8. Naj bo X normiran prostor, ki je gost podprostor v Banachovem prostoru Y . Tedaj sta \hat{X} in Y izometrično izomorfna.

Dokaz. Naj bosta ι_Y in $\iota_{\hat{X}}$ vložitvi.

$$\begin{array}{ccc} X & \xrightarrow{\iota_{\hat{X}}} & \hat{X} \\ & \nwarrow \iota_Y^{-1} & \uparrow \\ & & \iota_Y(X) \end{array}$$

Preslikavi ι_Y^{-1} in $\iota_{\hat{X}}$ sta izometriji, torej je tak tudi njun kompozitum. To preslikavo lahko enolično razširimo do izometrije prostorov Y in \hat{X} . \square

Definicija 4.2.9. Naj bo $T : X \rightarrow X$ omejen operator med normiranimi prostoroma. Tedaj je T OBRNLJIV, če je bijektiven in T^{-1} omejen.

Definicija 4.2.10. Operator $T : X \rightarrow Y$ je NAVZDOL OMEJEN, če obstaja $c > 0$, da je $\|Tx\| \geq c \|x\|$ za vsak $x \in X$.

Opomba. Vsak navzdol omejen operator je injektiven.

Trditev 4.2.11. Naj bo $T \in B(X, Y)$. Naslednji trditvi sta ekvivalentni.

- Obstaja operator $T^{-1} : TX \rightarrow X$.
- T je navzdol omejen.

4.2.1 Banachov izrek

Definicija 4.2.12. Naj bo X vektorski prostor. Preslikava $p : X \rightarrow \mathbb{R}$ je **SUBLINEARNI FUNKCIONAL**, če je $p(x + y) \leq p(x) + p(y)$ in $p(\lambda x) = \lambda p(x)$ za poljubne $x, y \in X$ ter $\lambda \geq 0$.

Vprašanje 11. Kaj je sublinearni funkcional?

Primer. Vsaka polnorma je sublinearni funkcional.

Izrek 4.2.13 (realni Hahn-Banachov izrek). *Naj bo $Y \leq X$ vektorski prostor in $p : X \rightarrow \mathbb{R}$ sublinearni funkcional. Naj bo $f : Y \rightarrow \mathbb{R}$ tak linearni funkcional, da za vsak $y \in Y$ velja $f(y) \leq p(y)$. Tedaj obstaja linearni funkcional $F : X \rightarrow \mathbb{R}$, da je $F|_Y = f$ in $F(x) \leq p(x)$ za vsak $x \in X$.*

Dokaz. Prvo obravnavajmo primer, kjer je $\dim X/Y = 1$. Tedaj je $X = Y \oplus \mathbb{R}x_0$ za neki $x_0 \in X \setminus Y$. Vse možne linearne razširitve f do F so oblike

$$F(x) = F(y + \lambda x_0) = F(y) + \lambda F(x_0),$$

torej je razširitev enolično določena z $F(x_0) =: \alpha$. Za $y_1, y_2 \in Y$ velja

$$f(y_1 + y_2) \leq p(y_1 + y_2) = p(y_1 + y_2 + x_0 - x_0) \leq p(y_1 + x_0) + p(y_2 - x_0)$$

oziroma

$$f(y_2) - p(y_2 - x_0) \leq p(y_1 + x_0) - f(y_1),$$

torej je

$$\sup_{y \in Y} (f(y) - p(y - x_0)) \leq \inf_{y \in Y} (p(y + x_0) - f(y)).$$

Za α lahko izberemo katerokoli vrednost med tema številoma. Potem definiramo $F(y + tx_0) = f(y) + t\alpha$ in ločimo primere.

- Če je $t = 0$, je $F(y) \leq p(y)$ po predpostavki, saj je $F(y) = f(y)$.
- Če je $t > 0$, je $\alpha \leq p(y/t + x_0) - f(y/t)$, kar množimo s t , in dobimo $f(y) + t\alpha \leq p(y + tx_0)$.
- Če je $t < 0$, je $\alpha \geq f(-y/t) - p(-y/t - x_0)$, kar množimo z $(-t)$ in zaključimo kot zgoraj.

Za splošen primer tvorimo množico

$$\mathcal{A} = \{(Y_1, f_1) \mid Y \leq Y_1 \leq X, f_1|_Y = f, \forall y_1 \in Y_1. f_1(y_1) \leq p(y_1)\}$$

in definiramo relacijo

$$(Y_1, f_1) \preceq (Y_2, f_2) \Leftrightarrow Y_1 \leq Y_2 \wedge f_2|_{Y_1} = f_1.$$

To je očitno delna urejenost. Naj bo $\{(Y_i, f_i)\}_{i \in I}$ neka veriga v \mathcal{A} . Vzemimo $Z = \bigcup_{i \in I} Y_i$. To je podprostor, ker je veriga urejena. Definiramo še preslikavo $g : Z \rightarrow \mathbb{R}$ z $g(z) = f_i(z)$ za nek $i \in I$, za katerega je $z \in Y_i$. To je dobro definiran funkcional, za katerega velja $g(z) \leq p(z)$ za vse $z \in Z$.

Očitno je (Z, g) zgornja meja za verigo. Po Zornovi lemi ima \mathcal{A} maksimalen element (\tilde{Y}, \tilde{f}) . Če je $\tilde{Y} \neq X$, potem obstaja $x \in X \setminus \tilde{Y}$. Po prvem koraku dokaza lahko \tilde{f} razširimo na linearno ogrinjačo množico $\{\tilde{Y}, x\}$, kar je protislovje z maksimalnostjo. Torej $X = \tilde{Y}$. \square

Vprašanje 12. Povej in dokaži realni Hahn-Banachov izrek.

Lema 4.2.14. Naj bo X kompleksen vektorski prostor. Potem veljajo naslednje točke.

- Če je $f : X \rightarrow \mathbb{R}$ \mathbb{R} -linearen funkcional, je $\tilde{f} : X \rightarrow \mathbb{C}$, definiran z $\tilde{f}(x) = f(x) - if(ix)$ \mathbb{C} -linearen funkcional, in velja $\operatorname{Re} \tilde{f} = f$.
- Če je $g : X \rightarrow \mathbb{C}$ \mathbb{C} -linearen funkcional in $\operatorname{Re} g = f$, potem je $g = \tilde{f}$.
- Če je p polnorma, potem je $|f(x)| \leq p(x)$ za vse $x \in X$ natanko tedaj, ko za vse $x \in X$ velja $|\tilde{f}(x)| \leq p(x)$.
- Če je X normiran in $f : X \rightarrow \mathbb{R}$ omejen, je \tilde{f} omejen in $\|\tilde{f}\| = \|f\|$.

Dokaz. Prva točka je enostavna. Za drugo točko le pogledamo $g(x) = f(x) + if_1(x)$ ter $ig(x) = g(ix)$, s čimer pokažemo $f_1(x) = -f(ix)$.

Za tretjo točko opazimo

$$|f(x)| = |\operatorname{Re} \tilde{f}(x)| \leq |\tilde{f}(x)| \leq p(x),$$

v drugo smer pa

$$|\tilde{f}(x)| = e^{i\varphi} \tilde{f}(x) = \tilde{f}(e^{i\varphi}x) = \operatorname{Re} \tilde{f}(e^{i\varphi}x)$$

za nek kot φ . Potem

$$|\tilde{f}(x)| = f(e^{i\varphi}x) \leq p(e^{i\varphi}x) = |e^{i\varphi}| p(x) = p(x).$$

Za zadnjo točko pogledajmo $|f(x)| \leq \|f\| \|x\|$. Potem je $p(x) = \|f\| \|x\|$ polnorma na X , za katero velja $|f(x)| \leq p(x)$. Po prejšnji točki velja $|\tilde{f}(x)| \leq p(x)$, torej je \tilde{f} omejen z $\|f\|$. Po drugi strani je $\|\tilde{f}(x)\| \leq q(x)$ za $q(x) = \|\tilde{f}\| \|x\|$, in zato po prejšnji točki $|f(x)| \leq q(x)$ in $\|f\| \leq \|\tilde{f}\|$. \square

Vprašanje 13. Pokaži, da lahko \mathbb{R} -linearni funkcional iz kompleksnega vektorskega prostora X enolično razširimo do \mathbb{C} -linearnega funkcionala, in da razširitev ohranja normo.

Izrek 4.2.15 (kompleksni Hahn-Banach). Naj bo X vektorski prostor nad \mathbb{F} , $Y \leq X$ in p polnorma na X . Če je $f : Y \rightarrow \mathbb{F}$ linearni funkcional, da za vse $y \in Y$ velja $|f(y)| \leq p(y)$, potem obstaja linearni funkcional $F : X \rightarrow \mathbb{F}$, za katerega je $F|_Y = f$ in $|F(x)| \leq p(x)$ za vsak $x \in X$.

Dokaz. Za $\mathbb{F} = \mathbb{R}$ po realni verziji izreka obstaja funkcional $F : X \rightarrow \mathbb{R}$, ki razširja f in za katerega je $F(x) \leq p(x)$ za $x \in X$. Velja $-F(x) = F(-x) \leq p(-x) = p(x)$, torej $|F(x)| \leq p(x)$.

Če pa je $\mathbb{F} = \mathbb{C}$, vzemimo \mathbb{R} -linearen funkcional $f_1 = \operatorname{Re} f$, za katerega po lemi velja $|f_1(y)| \leq p(y)$ za $y \in Y$. Tega lahko razširimo do $F_1 : X \rightarrow \mathbb{R}$, nato pa vzamemo funkcional \tilde{F} , za katerega velja $\operatorname{Re} \tilde{F} = F_1$. S pomočjo leme hitro vidimo, da \tilde{F} razširja f in $|\tilde{F}(x)| \leq p(x)$. \square

Vprašanje 14. Povej in dokaži kompleksni Hahn-Banachov izrek.

Izrek 4.2.16 (Hahn-Banachov izrek za normirane prostore). *Naj bo $Y \leq X$ podprostor normiranega prostora X in $f : Y \rightarrow \mathbb{F}$ omejen. Tedaj obstaja $F : X \rightarrow \mathbb{F}$, da je $F|_Y = f$ ter $\|F\| = \|f\|$.*

Vprašanje 15. Povej Hahn-Banachov izrek za normirane prostore.

Posledica 4.2.17. *Naj bo X normiran in $x \in X$ neničeln vektor. Tedaj obstaja $F \in X^*$, da je $\|F\| = 1$ in $F(x) = \|x\|$.*

Dokaz. Naj bo $Y = \mathbb{F} \cdot x$. Funkcional $g(\lambda x) = \lambda \|x\|$ je zvezen, linearen in definiran na Y z $\|g\| = 1$. Po Hahn-Banachu ga lahko razširimo na funkcional $X \rightarrow \mathbb{F}$. \square

Posledica 4.2.18. *Naj bo X normiran in $x \in X$. Tedaj je*

$$\|x\| = \max\{|f(x)| \mid f \in X^*, \|f\| = 1\}.$$

Vprašanje 16. Pokaži, da je $\|x\| = \max\{|f(x)| \mid f \in X^*, \|f\| = 1\}$.

Posledica 4.2.19. *Naj bo $Y \leq X$ zaprt podprostor normiranega prostora X . Naj bo $x_0 \in X \setminus Y$ ter $d = d(x_0, Y)$. Tedaj obstaja $f \in X^*$, da je $f(x_0) = 1$, $f|_Y = 0$ in $\|f\| = 1/d$.*

Dokaz. Naj bo $g \in (X/Y)^*$ tak, da je $\|g\| = 1$ in $g(x_0 + Y) = \|x_0 + Y\| = d$. Definirajmo $f = g \circ \pi$, kjer je π kvocientna projekcija. Za $y \in Y$ velja $f(y) = 0$, ker je $y + Y = 0$ in g linearna. Funkcional f je omejen, ker je kompozitum omejenih funkcionalov, velja $f(x_0) = d$. Izračunamo

$$\|f\| = \sup_{\|x\| < 1} |f(x)| = \sup_{\|x\| < 1} |g(\pi(x))| = \sup_{\|x+Y\| < 1} g(x+Y) = \|g\| = 1,$$

torej je iskani funkcional $\frac{1}{d}f$. \square

Vprašanje 17. Pokaži, da lahko z linearnim funkcionalom ločiš točko od zaprtega podprostora.

Izrek 4.2.20. *Naj bo $Y \leq X$ podprostor normiranega prostora X . Tedaj je*

$$\bar{Y} = \bigcap \{\ker f \mid f \in X^*, Y \subseteq \ker f\}.$$

Dokaz. Naj bo Z presek iz izreka. Če je $Y \subseteq \ker f$, je $\overline{Y} \subseteq \ker f$, saj je jedro zaprto. Torej je $\overline{Y} \subseteq Z$. Recimo, da \overline{Y} ni enako Z . Tedaj obstaja $z \in Z \setminus \overline{Y}$, torej po prejšnji posledici obstaja $f \in X^*$, da je $f(z) = 1$ in $f|_{\overline{Y}} = 0$. Velja $Y \subseteq \ker f$, torej $z \in \ker f$, kar je protislovje z $f(z) = 1$. \square

Vprašanje 18. Kako lahko izraziš zaprtje podprostora z jedri linearnih funkcionalov? Dokaži enakost.

4.2.2 Adjungirani operator in drugi dual

Definicija 4.2.21. Za normiran prostor X in $f \in X^*$ definiramo $\hat{x}(f) = f(x)$.

Trditev 4.2.22. Velja $\hat{x} \in X^{**}$ in $\|\hat{x}\| = \|x\|$.

Dokaz. Velja $|\hat{x}(f)| = |f(x)| \leq \|f\| \|x\|$, zato $\|\hat{x}\| \leq \|x\|$. Po Hahn-Banachu obstaja f z $\|f\| = 1$ in $f(x) = \|x\|$. Tedaj $\hat{x}(f) = \|x\|$, torej $\|\hat{x}\| \geq \|x\|$. \square

Prostor X vložimo v X^{**} s preslikavo $i(x) = \hat{x}$.

Trditev 4.2.23. Preslikava i je linearna izometrična vložitev.

Naj bo $A : X \rightarrow Y$ omejen linearen operator. Za $f \in Y^*$ definiramo adjungirani operator operatorja A v smislu Banachovih prostorov, A^* , z

$$A^*f = f \circ A.$$

Trditev 4.2.24. Naj bo $A : X \rightarrow Y$ omejen linearen operator med normiranimi prostori. Tedaj je $A^* : Y^* \rightarrow X^*$ tudi omejen in $\|A^*\| = \|A\|$.

Dokaz. Izračunamo

$$|A^*f(x)| = |f(Ax)| \leq \|f\| \|Ax\| \leq \|f\| \|A\| \|x\|,$$

torej $\|A^*f\| \leq \|A\| \|f\|$. Iz tega sledi, da je tudi A^{**} omejen z $\|A^{**}\| \leq \|A^*\| \leq \|A\|$. Vemo pa

$$\|Ax\| = \left\| \widehat{Ax} \right\| = \|A^{**}\hat{x}\| \leq \|A^{**}\| \|\hat{x}\| = \|A^{**}\| \|x\|$$

torej je $\|A\| \leq \|A^{**}\|$ in so norme enake. \square

Vprašanje 19. Kako je definiran adjungiran operator v smislu Banachovih prostorov? Pokaži, da adjungiranje ohranja normo operatorja.

Definicija 4.2.25. Normiran prostor je REFLEKSIVEN, če je i_X surjekcija.

Opomba. Refleksivni prostori so Banachovi, saj je X^{**} Banachov.

4.3 Temeljni izreki funkcionalne analize

Izrek 4.3.1 (Baire). *Naj bo (X, d) poln metrični prostor in $(U_n)_n$ števna družina odprtih gostih množic v X . Tedaj je presek $\bigcap_n U_n$ gost v X .*

Dokaz. Dokazujemo, da za vsak $x \in X$ in $r > 0$ velja

$$\mathring{B}(x, r) \cap \bigcap_{n \in \mathbb{N}} U_n \neq \emptyset.$$

Naj bosta $x \in X$ ter $r > 0$ poljubna. Induktivno bomo konstruirali zaporedji $(x_n)_n$ in $(r_n)_n$ z naslednjimi lastnostmi:

- $B(x_{n+1}, r_{n+1}) \subseteq U_n \cap \mathring{B}(x_n, r_n)$,
- $r_n \leq 1/n$.

Postavimo $x_1 = x$ in $r_1 = \min\{1, r\}$. Recimo, da smo že konstruirali zaporedji do n -tega člena. Ker je U_n gosta odprta množica, je $U_n \cap \mathring{B}(x_n, r_n)$ neprazna odprta množica, zato obstajata $r_{n+1} \leq 1/(n+1)$ in x_{n+1} , da je $\mathring{B}(x_{n+1}, 2r_{n+1}) \subseteq U_n \cap \mathring{B}(x_n, r_n)$. Prva lastnost sedaj velja, ker je $B(x_{n+1}, r_{n+1}) \subseteq \mathring{B}(x_{n+1}, 2r_{n+1})$.

S tem smo konstruirali želeni zaporedji. Ker je

$$x_n \in B(x_n, r_n) \subseteq U_{n-1} \cap \mathring{B}(x_{n-1}, r_{n-1}) \subseteq \mathring{B}(x_{n-1}, r_{n-1}) \subseteq \mathring{B}(x_m, r_m)$$

za $m < n$, je $d(x_m, x_n) \leq r_m \leq 1/m$. Torej je zaporedje Cauchyjevo in obstaja limita x_0 v X . Velja

$$x_0 \in \bigcap_{n \in \mathbb{N}} B(x_{n+1}, r_{n+1}) \subseteq \bigcap_{n \in \mathbb{N}} U_n \cap \mathring{B}(x_n, r_n) \subseteq \mathring{B}(x_1, r_1) \cap \bigcap_{n \in \mathbb{N}} U_n = \mathring{B}(x, r) \cap \bigcap_{n \in \mathbb{N}} U_n,$$

s čimer je dokaz zaključen. \square

Vprašanje 20. Povej in dokaži Bairov izrek.

Posledica 4.3.2. *Naj bo X poln metrični prostor in $(A_n)_n$ zaporedje zaprtih množic, da je $X = \bigcup_n A_n$. Tedaj obstaja $m \in \mathbb{N}$, da je $\mathring{A}_m \neq \emptyset$.*

Dokaz. Če je $\mathring{A}_j = \emptyset$ za vse j , potem je A_j^c odprta in gosta. Potem je $\bigcap_j A_j^c$ gost, torej $\bigcup_j A_j = \left(\bigcap_j A_j^c\right)^c \neq X$. \square

Vprašanje 21. Povej in dokaži posledico Bairovega izreka z zaprtimi množicami.

Naj bosta X in Y normirana prostora ter $\mathcal{F} \subseteq B(X, Y)$. Recimo, da obstaja $M \geq 0$, za katerega je $\|T\| \leq M$ za vse $T \in \mathcal{F}$. Tedaj za vsak $x \in X$ velja

$$\|Tx\| \leq \|T\| \|x\| \leq M \|x\|,$$

oziroma $\mathcal{F}x \subseteq B(0, M \|x\|)$. Pravimo, da je \mathcal{F} OMEJENA PO TOČKAH.

Izrek 4.3.3 (princip enakomerne omejenosti). *Naj bo X Banachov in Y normiran prostor. Naj bo $\mathcal{F} \subseteq B(X, Y)$ po točkah omejena družina. Potem je kot množica operatorjev enakomerno omejena v $B(X, Y)$.*

Dokaz. Definiramo

$$A_n = \{x \in X \mid \forall T \in \mathcal{F}. \|Tx\| \leq n\} = \bigcap_{T \in \mathcal{F}} f_T^{-1}([0, n])$$

za $f_T(x) = \|Tx\|$. Ker je A_n presek zaprtih množic, je zaprt. Ker je \mathcal{F} omejena po točkah, za poljuben $x \in X$ obstaja $m \in \mathbb{N}$, da je $\|Tx\| \leq m$ za poljuben $T \in \mathcal{F}$, oziroma $x \in A_m$. Torej je X enak uniji množic A_n , in po posledici Bairovega izreka obstaja tak $n_0 \in \mathbb{N}$, da ima A_{n_0} notranjo točko, in posledično vsebuje odprto kroglo $\mathring{B}(x_0, r)$.

Izberimo poljuben $x \in B(0, 1)$ ter definirajmo $y = x_0 + \frac{r}{2}x$. Velja $y \in \mathring{B}(x_0, r)$, torej je $\|Ty\| \leq n_0$ za poljuben $T \in \mathcal{F}$. Sledi

$$\|Tx\| = \left\| T \frac{2y - x_0}{r} \right\| \leq \frac{2}{r} \|Ty\| + \frac{1}{r} \|Tx_0\| \leq \frac{2}{r} n_0 + \frac{1}{r} \|Tx_0\| =: M.$$

Torej je $\|T\| \leq M$ za vsak $T \in \mathcal{F}$. □

Vprašanje 22. Povej in dokaži princip enakomerne omejenosti.

Izrek 4.3.4 (o šibki omejenosti). *Naj bo $A \subseteq X$ podmnožica normiranega prostora X . Potem je A omejena v X natanko tedaj, ko je za vsak $f \in X^*$ množica $\{f(x) \mid x \in A\}$ omejena v \mathbb{F} .*

Dokaz. V desno: Ker je A omejena, obstaja M , da je $\|x\| \leq M$ za vsak $x \in A$. Potem je $|f(x)| \leq \|f\| \|x\| \leq M \|f\|$ za $f \in X^*$.

V levo: Oglejmo si vložitev v drugi dual, $i(x) = \hat{x}$. Množica $\{f(x) \mid x \in A\}$ je omejena natanko tedaj, ko je omejena množica $\{\hat{x}(f) \mid \hat{x} \in i(A)\}$, ker pa je $i(A)$ omejena po točkah, je po principu enakomerne omejenosti $i(A)$ tudi enakomerno omejena, torej obstaja $M \geq 0$, da je $\|x\| = \|\hat{x}\| \leq M$ za vse $x \in A$. □

Vprašanje 23. Povej in dokaži izrek o šibki omejenosti.

Posledica 4.3.5. *Naj bo X Banachov prostor in Y normiran prostor. Naj bo $A \subseteq B(X, Y)$ taka, da za vsak $f \in Y^*$ in vsak $x \in X$ obstaja $M_{f,x} \geq 0$, da je $|f(Tx)| \leq M_{f,x}$ za vsak $T \in A$. Tedaj je A omejena.*

Dokaz. Po izreku o šibki omejenosti je množica $\{Ty \mid T \in A\}$ omejena v Y . Torej je A omejena po točkah, ker pa je X Banachov, je A omejena po principu enakomerne omejenosti. □

Lema 4.3.6. *Naj bo X Banachov prostor in $(x_n)_n$ zaporedje vektorjev, za katere velja $\sum \|x_n\| < \infty$. Tedaj vrsta $\sum x_n$ konvergira v X .*

Dokaz. Označimo $s_n = \sum_{i=1}^n x_i$. Za $n > m$ velja

$$\|s_n - s_m\| = \|x_n + x_{n-1} + \cdots + x_{m+1}\| \leq \|x_{m+1}\| + \cdots + \|x_n\| \leq \sum_{k \geq m} \|x_k\| \xrightarrow{m \rightarrow \infty} 0.$$

Torej je zaporedje $(s_n)_n$ Cauchyjevo in zato konvergentno. \square

Izrek 4.3.7 (o odprti preslikavi). *Naj bo T omejen surjektiven linearen operator med Banachovima prostoroma X in Y . Tedaj je T odprta preslikava.*

Dokaz. Dokaz poteka v štirih korakih. V prvem koraku dokažimo, da če odprta podmnožica $U \subseteq X$ vsebuje 0, ima $\overline{T(U)}$ notranjo točko. Ker je U odprta, obstaja $\delta > 0$, da je

$$\delta \mathring{B}(0, 1) = \mathring{B}(0, \delta) \subseteq U.$$

Poljuben $x \in X$ je v

$$x \in 2\|x\| \mathring{B}(0, 1) = \frac{2\|x\|}{\delta} \delta \mathring{B}(0, 1) \subseteq mU$$

za $m \geq \frac{2\|x\|}{\delta}$, torej je

$$X \subseteq \bigcup_{n \in \mathbb{N}} nU.$$

Velja

$$Y = TX = \bigcup_{n \in \mathbb{N}} T(nU) = \bigcup_{n \in \mathbb{N}} \overline{T(nU)}$$

in po Bairovem izreku obstaja n_0 , da ima $\overline{T(n_0U)}$ notranjo točko. Množenje s skalarjem je homeomorfizem, zato velja $\overline{T(n_0U)} = n_0 \overline{TU}$ in ima tudi \overline{TU} notranjo točko.

V drugem koraku pokažimo, da je ob isti predpostavki točka 0 notranja za \overline{TU} . Ker je U odprta v X , obstaja $\mathring{B}(0, \varepsilon) \subseteq U$. Vzemimo $V = \mathring{B}(0, \varepsilon/2)$. Potem je množica $V - V \subseteq \mathring{B}(0, \varepsilon)$, kjer smo vzeli definicijo

$$A - B := \{a - b \mid a \in A, b \in B\}.$$

Po dokazanem v prvem koraku ima \overline{TV} notranjo točko, torej obstaja odprta množica $W \subseteq \overline{TV}$. Množica

$$\bigcup_{w \in W} (W - \{w\}) = W - W \subseteq \overline{TV} - \overline{TV} \subseteq \overline{TV - TV} = \overline{T(V - V)} \subseteq \overline{TU}$$

je unija odprtih množic in vsebuje 0. Druga vključitev zgoraj velja, ker je funkcija $m : Y \times Y \rightarrow Y$, $m(y_1, y_2) = y_1 - y_2$, zvezna.

V tretjem koraku spet ob isti predpostavki pokažimo, da TU vsebuje odprto okolico za 0. Najprej pokažimo poseben primer, če je $U = \mathring{B}(0, \varepsilon)$. Definiramo $\varepsilon_0 = \varepsilon/2$ in zapišemo

$$\varepsilon_0 = \sum_{i=1}^{\infty} \varepsilon_i$$

za neko zaporedje $(\varepsilon_i)_i$ pozitivnih števil. Po drugem koraku za vsak i obstaja $\eta_i > 0$, da je $\mathring{B}(0, \eta_i) \subseteq T(\mathring{B}(0, \varepsilon_i))$. Sedaj izberimo $y \in \mathring{B}(0, \eta_0)$ in pokažimo $y = Tx$ za neki $x \in \mathring{B}(0, \varepsilon)$.

Če je $\|x\| < \varepsilon_i$, velja $\|Tx\| \leq \|T\| \varepsilon_i$, torej je

$$\mathring{B}(0, \eta_i) \subseteq T(\mathring{B}(0, \varepsilon_i)) \subseteq \mathring{B}(0, \varepsilon_i \|T\|).$$

Sledi $\eta_i \leq \varepsilon_i \|T\|$ in zaporedje η_i konvergira k 0. Po predpostavki je $y \in \mathring{B}(0, \eta_0) \subseteq T(\mathring{B}(0, \varepsilon_0))$. Po definiciji zaprtja η_1 -okolica za y seka $T(\mathring{B}(0, \varepsilon_0))$, torej obstaja $x_0 \in X$, da je $\|x_0\| < \varepsilon_0$ in $\|y - Tx_0\| < \eta_1$. Sedaj velja $y - Tx_0 \in \mathring{B}(0, \eta_1) \subseteq T(\mathring{B}(0, \varepsilon_1))$, in spet po definiciji zaprtja η_2 -okolica za $y - Tx_0$ seka $T(\mathring{B}(0, \varepsilon_1))$, torej obstaja $x_1 \in X$ z $\|x_1\| < \varepsilon_1$ ter $\|y - Tx_0 - Tx_1\| < \eta_2$.

Postopek ponavljamo, s čimer dobimo zaporedje $(x_n)_n$, za katerega velja $\|x_n\| < \varepsilon_n$ ter $\|y - T(x_1 + \dots + x_n)\| < \eta_{n+1}$. Ker je $\sum \|x_n\| < \sum \varepsilon_n < \infty$, vrsta $\sum x_n$ konvergira absolutno in zato v Banachovem prostoru X konvergira proti nekemu vektorju x . Ker je T omejen operator, konvergira tudi vrsta $\sum Tx_n$ in velja $Tx = \sum Tx_n$. To mora biti enako y , saj $\eta_i \rightarrow 0$. Dodatno je $\|x\| \leq \sum \|x_n\| < 2\varepsilon_0 = \varepsilon$, torej je $T(\mathring{B}(0, \varepsilon))$ okolica za 0.

Če pa U ni take oblike, obstaja $\varepsilon > 0$, da je $\mathring{B}(0, \varepsilon) \subseteq U$, in po ravno dokazanem T -slika te množice vsebuje odprto kroglo okoli 0. Zato jo vsebuje tudi TU .

V zadnjem koraku dokažimo, da je slika odprte $U \subseteq X$ odprta v Y . Naj bo $y \in TU$. Obstaja $x \in U$, da je $y = Tx$, torej je $V = U - x$ odprta okolica za 0 in je po prejšnjem koraku TV okolica za 0 v Y . Potem je $TU = TV + Tx$ okolica za y . \square

Vprašanje 24. Povej in dokaži izrek o odprti preslikavi.

Posledica 4.3.8. *Naj bo T omejen linearen bijektiven operator med Banachovima prostoroma. Potem je njegov inverz tudi omejen.*

Vprašanje 25. Povej posledico izreka o odprti preslikavi.

Posledica 4.3.9. *Naj bo X vektorski prostor in $\|\cdot\|_1, \|\cdot\|_2$ dve normi na X , za kateri je X Banachov prostor. Potem sta normi bodisi ekvivalentni bodisi neprimerljivi.*

Dokaz. Če je $\|x\|_1 \leq c_1 \|x\|_2$, je identiteta $I : (X, \|\cdot\|_2) \rightarrow (X, \|\cdot\|_1)$ omejena, in je njen inverz tudi omejen. \square

Vprašanje 26. Dokaži: dve normi, za kateri je X Banachov prostor, sta bodisi ekvivalentni bodisi neprimerljivi.

Lema 4.3.10. *Naj bosta X, Y normirana prostora in $f : X \rightarrow Y$ zvezna preslikava. Tedaj je $\Gamma_f^{\text{zap}} \subseteq X \times Y$, če ta produkt opremimo z normo $\|(x, y)\|_\infty = \max\{\|x\|, \|y\|\}$.*

Dokaz. Naj gre $(x_n, f(x_n)) \rightarrow (x, y)$. Velja $x_n \rightarrow x$ in $f(x_n) \rightarrow y$, ker pa je f zvezna, tudi $f(x_n) \rightarrow f(x)$. \square

Izrek 4.3.11 (o zaprtem grafu). *Naj bo $T : X \rightarrow Y$ linearna preslikava, X in Y Banachova prostora ter Γ_T zaprt v $X \times Y$. Potem je T omejena.*

Dokaz. Graf je zaprt, torej je Banachov. Oglejmo si spodnji diagram.

$$\begin{array}{ccc} X & \xrightarrow{(x, Tx)} & \Gamma_T \\ & \searrow T & \downarrow \text{pr}_2 \\ & & Y \end{array}$$

Projekcija $\text{pr}_1 : \Gamma_T \rightarrow X$ je omejena in bijektivna, torej je tudi njen inverz $x \mapsto (x, Tx)$ omejen po posledici izreka o odprti preslikavi. Velja $T = \text{pr}_2 \circ \text{pr}_1^{-1}$. \square

Vprašanje 27. Povej in dokaži izrek o zaprtem grafu.

4.4 Hilbertovi prostori

Omejimo se na primer $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.

Definicija 4.4.1. Naj bo X vektorski prostor nad \mathbb{F} . Preslikava $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{F}$ je SKALARNI PRODUKT, če zadošča

- $\langle x, x \rangle \geq 0$ (realno in nenegativno),
- $\langle x, x \rangle = 0$ natanko tedaj, ko je $x = 0$,
- $\langle \alpha x + \beta y, z \rangle = \alpha \langle x, z \rangle + \beta \langle y, z \rangle$,
- $\langle x, y \rangle = \overline{\langle y, x \rangle}$.

Trditev 4.4.2 (Paralelogramska enakost). *Naj bo X prostor s polskalarним produktom. Za $x, y \in X$ velja*

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2).$$

Trditev 4.4.3. *Skalarni produkt je zvezna preslikava.*

Izrek 4.4.4 (Jordan, von Neumann). *Če v normiranem prostoru velja paralelogramska enakost, je norma porojena s skalarnim produktom.*

Definicija 4.4.5. Prostor X s skalarnim produktom je HILBERTOV PROSTOR, če je za porojeno normo Banachov prostor.

Naj bo X prostor s skalarnim produktom in \hat{X} napolnitev X kot normiran prostor. Ker norma na X ustreza paralelogramski enakosti, zaradi zveznosti norme to velja tudi na \hat{X} in je norma na \hat{X} porojena s skalarnim produktom. Torej je Hilbertov prostor. Če $x_n \rightarrow x$ in $y_n \rightarrow y$, velja

$$\langle x, y \rangle = \lim_{n \rightarrow \infty} \langle x_n, y_n \rangle.$$

Definicija 4.4.6. Vektorja x in y sta PRAVOKOTNA, če $\langle x, y \rangle = 0$. Označimo $x \perp y$.

Definicija 4.4.7. Množici A in B sta PRAVOKOTNI, če je $\langle a, b \rangle = 0$ za vsak $a \in A$ ter $b \in B$.

Izrek 4.4.8 (Pitagora). *Naj bo X vektorski prostor s skalarnim produktom. Če sta vektorja x in y pravokotna, je $\|x\|^2 + \|y\|^2 = \|x + y\|^2$.*

Izrek 4.4.9. *Naj bo H Hilbertov prostor in K neprazna zaprta konveksna množica v H . Tedaj za vsak $x \in H$ obstaja natanko en $k \in K$, da je $d(x, K) = \|x - k\|$.*

Dokaz. Brez škode za splošnost lahko privzamemo $x = 0$. Označimo $d = \inf\{\|y\| \mid y \in K\}$. Po definiciji infimuma obstaja zaporedje $(k_n)_n$, da $\|k_n\| \rightarrow d$. Izberimo $\varepsilon > 0$. Potem obstaja $N \in \mathbb{N}$, da za $n \geq N$ velja $\|k_n\|^2 \leq d^2 + \varepsilon^2/4$. Po paralelogramski enakosti velja

$$\|k_n - k_m\|^2 = 2\|k_n\|^2 + 2\|k_m\|^2 - 4\left\|\frac{k_n + k_m}{2}\right\|^2 \leq 4d^2 + \varepsilon^2 - 4d^2 = \varepsilon^2,$$

torej je $(k_n)_n$ Cauchyjevo in ima limito $k \in \overline{K} = K$. Velja $\|k\| = d$.

Za enoličnost še enkrat uporabimo paralelogramsko enakost. Če je $k' \in K$ še en vektor s $\|k'\| = d$, potem

$$\|k - k'\|^2 = 2\|k\|^2 + 2\|k'\|^2 - 4\left\|\frac{k + k'}{2}\right\|^2 \leq 0,$$

saj je K konveksna in $k, k' \in K$. □

Vprašanje 28. Pokaži: v Hilbertovem prostoru obstaja enolična projekcija točke na konveksno zaprto množico.

Izrek 4.4.10. *Naj bo M zaprt podprostor Hilbertovega prostora H , $x \in H$ ter $x_0 \in M$. Tedaj velja*

$$x - x_0 \perp M \Leftrightarrow d(x, M) = \|x - x_0\|.$$

Dokaz. Recimo, da za $x_0 \in M$ velja $d(x, M) = \|x - x_0\|$ in da $x - x_0$ ni pravokoten na M . Tedaj obstaja $y \in M$, da je $\langle x - x_0, y \rangle \neq 0$. Brez škode za splošnost lahko privzamemo, da je ta skalarni produkt pozitiven, sicer y zavrtimo za potreben kot. Potem je

$$\|x - (x_0 + \varepsilon y)\|^2 = \|x - x_0\|^2 - 2\operatorname{Re}(\varepsilon \langle x - x_0, y \rangle) + |\varepsilon|^2 \|y\|^2$$

za poljuben $\varepsilon \in \mathbb{C}$, saj je $(x_0 + \varepsilon y) \in M$. Če izberemo dovolj majhen $\varepsilon \in \mathbb{R}$, dobimo

$$\|x - (x_0 + \varepsilon y)\|^2 < \|x - x_0\|^2,$$

kar je protislovje, saj je x_0 najbližji vektor iz M . Torej je $x - x_0 \perp M$. Tedaj za vsak $y \in M$ velja

$$\|x - y\|^2 = \|x - x_0\|^2 + \|x_0 - y\|^2 \geq \|x - x_0\|^2. \quad \square$$

Vprašanje 29. Pokaži: Če je M zaprt podprostor v Hilbertovem prostoru in $x_0 \in M$, potem je $x - x_0$ pravokoten na M natanko tedaj, ko je $d(x, M) = \|x - x_0\|$.

Definicija 4.4.11. Naj bo X prostor s skalarnim produktom. Za $x \in X$ definiramo $\{x\}^\perp = \{y \in X \mid x \perp y\}$, za $A \subseteq X$ pa

$$A^\perp = \bigcap_{x \in A} \{x\}^\perp.$$

Lema 4.4.12. Za $A \subseteq X$ je A^\perp vedno zaprt podprostor v X .

Dokaz. Dovolj je pokazati, da je $\{x\}^\perp$ zaprt podprostor za katerikoli $x \in X$. Očitno je podprostor. Za zaprtost vzemimo zaporedje $(y_n)_n$ v $\{x\}^\perp$, ki konvergira k $y \in \overline{\{x\}^\perp}$. Potem je

$$\langle y, x \rangle = \langle y - y_n, x \rangle + \langle y_n, x \rangle = \langle y - y_n, x \rangle$$

in zato

$$|\langle y, x \rangle| \leq \|y - y_n\| \|x\| \xrightarrow{n \rightarrow \infty} 0$$

po Cauchy-Schwarzu. Torej je $y \in \{x\}^\perp$. \square

Vprašanje 30. Pokaži: A^\perp je vedno zaprt podprostor.

Izrek 4.4.13. Naj bo M zaprt podprostor v Hilbertovem prostoru H . Za $x \in H$ definiramo $Px \in M$ kot tisti vektor, ki je najbližji x med vektorji iz M . Potem velja:

- P je linearen operator $H \rightarrow M$,
- $\|Px\| \leq \|x\|$,
- $P^2 = P$,
- $\text{im } P = M$ in $\ker P = M^\perp$,
- $H = M \oplus M^\perp$ in $M^{\perp\perp} = M$.

Dokaz. Prva točka: Vzemimo $z \in M$. Potem je

$$\langle (\alpha x + \beta y) - (\alpha Px + \beta Py), z \rangle = \alpha \langle x - Px, z \rangle + \beta \langle y - Py, z \rangle = 0,$$

torej po prejšnjem izreku $P(\alpha x + \beta y) = \alpha Px + \beta Py$.

4 Uvod v funkcionalno analizo

Za drugo točko izračunamo

$$\|x\|^2 = \|x - Px + Px\|^2 = \|x - Px\|^2 + \|Px\|^2 \geq \|Px\|^2.$$

Tretja točka je očitna. Za četrto je seveda $\operatorname{im} P = M$, za $x \in \ker P$ velja $x - Px \in M^\perp$, torej ($\ker Px = 0$) tudi $x \in M^\perp$. Če pa je $x \in M^\perp$, je $x = x - 0 \in M^\perp$, torej $Px = 0$ po definiciji P .

Za zadnjo točko razcepimo $x = Px + (x - Px)$. Ker za $A \subseteq H$ vedno velja $A \cap A^\perp \subseteq \{0\}$ in ker je $0 \in M$, je $H = M \oplus M^\perp$. Preslikava $I - P$ je pravokoten projektor na M^\perp . Velja $M^{\perp\perp} = \ker(I - P) = \operatorname{im} P = M$. \square

Vprašanje 31. Definiraj ortogonalni projektor na zaprt podprostor M v Hilbertovem prostoru H . Dokaži, da je linearen, idempotent, da velja $\|Px\| \leq \|x\|$ in da velja $H = M \oplus M^\perp$.

Idempotent P iz izreka je pravokotni projektor na M vzdolž M^\perp . Množica M^\perp se imenuje ORTOGONALNI KOMPLEMENT M .

Posledica 4.4.14. Za $A \subseteq H$ je $A^{\perp\perp} = \overline{\operatorname{Lin} A} =: [A]$.

Dokaz. Seveda je $A \subseteq [A]$. Velja $[A]^\perp \subseteq A^\perp$ in $A^{\perp\perp} \subseteq [A]^{\perp\perp} = [A]$ ter celo $[A]^\perp = A^\perp$, saj za $x \in A^\perp$ velja $x \perp A$ in bo x pravokoten tudi na linearno ogrinjačo A in njeno zaprtje. Torej $A^{\perp\perp} = [A]^{\perp\perp}$. \square

Vprašanje 32. Kaj je $A^{\perp\perp}$ za $A \subseteq H$?

Naj bo X prostor s skalarnim produktom in $y \in X$. Definiramo $f_y : X \rightarrow \mathbb{F}$ z

$$f_y(x) = \langle x, y \rangle.$$

Lema 4.4.15. Preslikava f_y je omejen linearen funkcional z $\|f_y\| = \|y\|$.

Dokaz. Očitno je linearen. Velja

$$\|f_y(x)\| = |\langle x, y \rangle| \leq \|x\| \|y\|$$

po Cauchy-Schwarzu. Če je x linearno odvisen od y , velja enakost. \square

Izrek 4.4.16 (Riesz). Naj bo H Hilbertov prostor in $f \in H^*$. Tedaj obstaja natanko en $y \in H$, da je $f(x) = \langle x, y \rangle$ in $\|f\| = \|y\|$.

Dokaz. Za enoličnost: če je $f_y = f_z$, potem za vsak x

$$\langle x, y - z \rangle = 0,$$

torej $y = z$.

Če je $f = 0$, vzamemo $y = 0$. Sicer $f \neq 0$, in je $\ker f$ zaprt podprostor (prasluka zaprte množice), torej $H = \ker f \oplus (\ker f)^\perp$. Obstaja $z \in (\ker f)^\perp$, da je $f(z) = 1$. Za $x \in H$ potem velja

$$x = \underbrace{x - f(x)z}_{\in \ker f} + \underbrace{f(x)z}_{\in (\ker f)^\perp},$$

torej $\langle x, z \rangle = \langle f(x)z, z \rangle = f(x) \langle z, z \rangle$. Potem je $f(x) = \langle x, z / \|z\|^2 \rangle$. \square

Vprašanje 33. Povej in dokaži Rieszov izrek.

Posledica 4.4.17. Za vsak $f \in (l^2)^*$ obstaja natanko ena $y \in l^2$, da je

$$f(x) = \sum_{n=1}^{\infty} x_n \overline{y_n}$$

Trditev 4.4.18. Preslikava $J : H \rightarrow H^*$, podana s predpisom $Jy = f_y$ za $f_y(x) = \langle x, y \rangle$, je poševno linearen izometrični izomorfizem.

Dokaz. Vemo $\|f_y\| = \|y\|$, torej je J izometrija. Po Rieszovem izreku je surjektivna in injektivna, poševna linearnost pa je preprost račun. \square

Izrek 4.4.19. Naj bo H Hilbertov prostor. Potem je tudi H^* Hilbertov s skalarnim produktom $\langle f, g \rangle_{H^*} = \langle y_g, y_f \rangle_H$.

Dokaz. Preprosto preverjanje. \square

Izrek 4.4.20. Naj bo H Hilbertov prostor in $K \leq H$ podprostor. Tedaj ima vsak $f \in K^*$ natanko eno Hahn-Banachovo razširitev na H .

Dokaz. Omejen funkcional $f : K \rightarrow \mathbb{F}$ lahko razširimo do $g : \overline{K} \rightarrow \mathbb{F}$, pri čemer se ohrani norma. Po Rieszovem izreku obstaja natanko ena $y \in \overline{K}$, da je $g(x) = \langle x, y \rangle$ za $x \in \overline{K}$. Definiramo $F(x) = \langle x, y \rangle$ za $x \in H$. Seveda je $F|_K = f$ in $\|F\| = \|y\| = \|f\|$.

Recimo, da je F' še ena Hahn-Banachova razširitev f . Po Rieszu obstaja natanko ena $y' \in H$, da je $F'(x) = \langle x, y' \rangle$. To velja tudi za $x \in \overline{K}$, torej $F'(x) = g(x)$, oziroma $\langle x, y' \rangle = \langle x, y \rangle$ za vse $x \in \overline{K}$. Sledi $y - y' \perp \overline{K}$. Potem je

$$\|F'\|^2 = \|y'\|^2 = \|y' - y + y\|^2 = \|y' - y\|^2 + \|y\|^2 = \|g\|^2 + \|y - y'\|^2.$$

Torej $\|y' - y\| = 0$. \square

Vprašanje 34. Pokaži, da so Hahn-Banachove razširitve v Hilbertovem prostoru enolične.

Posledica 4.4.21. Vsak Hilbertov prostor je refleksiven (vložitev v H^{**} je surjektivna).

4.4.1 Ortonormirani sistemi

Definicija 4.4.22. Naj bo X prostor s skalarnim produktom. Množica $E \subseteq X$ je ORTONORMIRAN SISTEM, če je $\|e\| = 1$ za vsak $e \in E$ ter $e \perp f$ za vsaka $e, f \in E$.

Opomba. Če velja le druga zahteva, je E ORTOGONALNA MNOŽICA.

Lema 4.4.23. Vsaka ortogonalna množica je linearno neodvisna.

Definicija 4.4.24. Naj bo H Hilbertov prostor. Ortonormiran sistem $E \subseteq H$ je KOMPLETEN ali BAZA Hilbertovega prostora H , če je maksimalen v množici vseh ortonormiranih sistemov (glede na \subseteq).

Vprašanje 35. Definiraj kompleten ortonormiran sistem.

Trditev 4.4.25. Vsak ortonormiran sistem v Hilbertovem prostoru lahko dopolnimo do kompletnega ortonormiranega sistema.

Dokaz. Če je $(F_\alpha)_\alpha$ veriga v množici vseh ortonormiranih sistemov, ki vsebujejo E , je

$$\bigcup_{\alpha} F_{\alpha}$$

zgornja meja za to verigo, ki je očitno ortonormiran sistem. Po Zornovi lemi obstaja kompleten ortonormiran sistem, ki vsebuje E . \square

Opomba. Kaj je rumeno in ekvivalentno aksiomu izbire? Zornova limona!

Posledica 4.4.26. Vsak Hilbertov prostor ima bazo.

Trditev 4.4.27. Naj bo $\{e_1, \dots, e_n\}$ ortonormiran sistem v Hilbertovem prostoru H . Naj bo P_n ortogonalna projekcija na $M_n = \text{Lin}\{e_1, \dots, e_n\}$. Tedaj za $x \in H$ velja

$$P_n x = \sum_{k=1}^n \langle x, e_k \rangle e_k.$$

Dokaz. Naj bo x_0 ta vsota. Tedaj za $1 \leq j \leq n$ velja

$$\langle x_0, e_j \rangle = \sum_{k=1}^n \langle x, e_k \rangle \langle e_k, e_j \rangle = \langle x, e_j \rangle,$$

torej je $x - x_0 \perp M_n$. Po definiciji je $x_0 = P_n x$. \square

Trditev 4.4.28 (Besselova neenakost). Naj bo $(e_n)_n$ števni ortonormiran sistem v prostoru s skalarnim produktom X . Tedaj za vsak $x \in X$ velja

$$\|x\|^2 \geq \sum_{n=1}^{\infty} |\langle x, e_n \rangle|^2.$$

Dokaz. Definiramo

$$x' = \sum_{k=1}^n \langle x, e_k \rangle e_k.$$

Enostavno lahko preverimo $x - x' \perp x'$, torej po Pitagori

$$\|x\|^2 = \|x - x'\|^2 + \|x'\|^2 \geq \|x'\|^2 = \sum_{k=1}^n |\langle x, e_k \rangle|^2.$$

To velja za vsak n , torej tudi v limiti. \square

Vprašanje 36. Povej in dokaži Besselovo neenakost.

Posledica 4.4.29. Naj bo X prostor s skalarnim produktom in $E \subseteq X$ ortonormiran sistem. Naj bo $x \in X$. Tedaj je $\{e \in E \mid \langle x, e \rangle \neq 0\}$ kvečjemu števna.

Dokaz. Definiramo

$$E_n = \left\{ e \in E \mid |\langle x, e \rangle| \geq \frac{1}{n} \right\}.$$

Potem je $\langle x, e \rangle \neq 0$ natanko tedaj, ko je $e \in E_n$ za vsak $n \in \mathbb{N}$. Trdimo, da so vsi E_n končni. Sicer obstaja $m \in \mathbb{N}$, da je E_m neskončna, torej vsebuje števno neskončno podmnožico $(e_k)_k$. Potem je

$$\|x\|^2 \geq \sum_{k=1}^{\infty} |\langle x, e_k \rangle|^2 = \infty,$$

ker je $|\langle x, e_k \rangle| \geq 1/m$ za vse k . \square

Vprašanje 37. Naj bo $\{e_i\}_i$ ONS in $x \in X$. Pokaži, da je kvečjemu števno mnogo vektorjev e_i pravokotnih na x .

Posledica 4.4.30. Če je $E \subseteq X$ kompleten ortonormiran sistem, za vsak $x \in X$ velja

$$\|x\|^2 \geq \sum_{e \in E} |\langle x, e \rangle|^2.$$

Dokaz. Po prejšnji posledici je $\langle x, e \rangle \neq 0$ za kvečjemu števno mnogo $e \in E$. Na njih uporabimo Besselovo neenakost. \square

Izrek 4.4.31. Za ortonormiran sistem $E \subseteq H$ so naslednje trditve ekvivalentne.

- E je kompleten,
- $E^\perp = \{0\}$,
- $[E] = H$ (zaprta linearna ogrinjača),

4 Uvod v funkcionalno analizo

- za vsak $x \in H$ velja

$$x = \sum_{e \in E} \langle x, e \rangle e,$$

- za poljubna $x, y \in H$ velja

$$\langle x, y \rangle = \sum_{e \in E} \langle x, e \rangle \langle y, e \rangle,$$

- (Parsevalova enakost) za vsak $x \in H$ velja

$$\|x\|^2 = \sum_{e \in E} |\langle x, e \rangle|^2.$$

Dokaz. 1 v 2: Recimo, da $E^\perp \neq \{0\}$. Vzamemo $x \in E^\perp \setminus \{0\}$ in definiramo $E \cup \{x/\|x\|\}$, kar je ortonormiran sistem, ki vsebuje E . —✕—

2 v 1: Recimo, da E ni KONS. Potem obstaja KONS E' , ki vsebuje E , in obstaja $x \in E' \setminus E$. Ampak $x \in E^\perp = \{0\}$. —✕—

Ekvivalentnost 2 in 3: Velja $[E] = H \Leftrightarrow E^\perp = [E]^\perp = H^\perp = \{0\}$.

2 v 4: Vzemimo $x \in H$. Vemo, da obstaja kvečjemu števno mnogo $e \in E$, da je $\langle x, e \rangle \neq 0$. Te vektorje oštevilčimo v $(e_n)_n$. Dokažimo, da je

$$x = \sum_{n=1}^{\infty} \langle x, e_n \rangle e_n.$$

Vrsta res konvergira, saj za delne vsote s_n in $m > n$ velja

$$\|s_m - s_n\|^2 = \left\| \sum_{k=n+1}^m \langle x, e_k \rangle e_k \right\|^2 = \sum_{k=n+1}^m |\langle x, e_k \rangle|^2 \leq \sum_{k=n+1}^{\infty} |\langle x, e_k \rangle|^2.$$

Ker ta vrsta konvergira po Besselovi neenakosti, je zaporedje $(s_n)_n$ Cauchyjevo v H . Zato $s_n \rightarrow x_0 \in H$. Ker je

$$\langle x_0, e_j \rangle = \sum_{k=1}^{\infty} \langle x, e_k \rangle \langle e_k, e_j \rangle = \langle x, e_j \rangle,$$

velja $x - x_0 \perp e_j$ za vsak j , torej $x = x_0$.

4 v 5: Preprost račun.

5 v 6: Preprost račun.

6 v 2: Če je $E^\perp \neq \{0\}$, obstaja $x \in E^\perp$, različen od 0. Po Parsevalu za x in E velja

$$0 \neq \|x\|^2 = \sum_{e \in E} |\langle x, e \rangle|^2 = 0,$$

kar je protislovno. —✕—

□

Vprašanje 38. Povej 5 trditev, ekvivalentnih dejstvu, da je $E \subseteq H$ KONS. Dokaži ekvivalence. Kaj je Parsevalova enakost?

Trditev 4.4.32. Poljubni ortonormirani bazi Hilbertovega prostora imata isto kardinalnost.

Dokaz. Naj bosta $E, F \subseteq H$ bazi. Če je $|E| < \infty$, rezultat vemo iz linearne algebre. Sicer za $e \in E$ tvorimo $F_e = \{f \in F \mid \langle e, f \rangle \neq 0\}$. Ta množica je kvečjemu števno neskončna, po drugi strani pa velja

$$F = \bigcup_{e \in E} F_e,$$

saj je E baza. Torej $|F| \leq |E| |\mathbb{N}| = |E|$ in podobno v drugo smer. \square

Vprašanje 39. Pokaži: če sta E, F neskončna KONS-a, imata isto kardinalnost.

Definicija 4.4.33. DIMENZIJA Hilbertovega prostora je enaka kardinalnosti katerekoli njene baze.

Lema 4.4.34. V separabilnem metričnem prostoru je vsaka družina paroma disjunktih odprtih krogel kvečjemu števno neskončna.

Dokaz. Naj bo $\{\dot{B}(x_i, \varepsilon_i) \mid i \in I\}$ družina paroma disjunktih odprtih krogel. Naj bo S števna gosta množica v danem metričnem prostoru. Zaradi gostosti je $\dot{B}(x_i, \varepsilon_i) \cap S \neq \emptyset$ za vse i . Izberimo $y_i \in \dot{B}(x_i, \varepsilon_i) \cap S$ in definiramo $\varphi : I \rightarrow S$ z $\varphi(i) = y_i$. Ker so krogle med seboj disjunktne, je φ injekcija, torej $|I| \leq |\mathbb{N}|$. \square

Trditev 4.4.35. Neskončnorazsežen Hilbertov prostor je separabilen natanko tedaj, ko je $\dim H = \aleph_0$.

Dokaz. V desno: Naj bo E KONS v H . Za $e, e' \in E$ velja

$$\|e - e'\|^2 = \langle e - e', e - e' \rangle = \|e\|^2 - 2 \operatorname{Re} \langle e, e' \rangle + \|e'\|^2 = 2.$$

Tvorimo $S = \{\dot{B}(e, \frac{\sqrt{2}}{2}) \mid e \in E\}$. Te krogle so paroma disjunktne, torej imamo injekcijo $\varphi : E \rightarrow S$. Po prejšnji lemi je S največ števna, torej $|E| \leq \aleph_0$.

V levo: Ker je $\dim H = \aleph_0$, obstaja števno KONS $(e_n)_n$. Vsak $x \in H$ lahko razvijemo v Fourierovo vrsto

$$x = \sum_{n=1}^{\infty} \langle x, e_n \rangle e_n.$$

Za poljuben $\varepsilon > 0$ potem obstaja n_ε , da za vse $n \geq n_\varepsilon$ velja

$$\left\| x - \sum_{k=1}^n \langle x, e_k \rangle e_k \right\| < \varepsilon.$$

Skalarne produkte $\langle x, e_k \rangle$ lahko aproksimiramo z $\lambda_k \in \mathbb{Q}$, če je $\mathbb{F} = \mathbb{R}$, oziroma $\lambda_k \in \mathbb{Q} + i\mathbb{Q}$, če je $\mathbb{F} = \mathbb{C}$. V obeh primerih zahtevamo $|\langle x, e_k \rangle - \lambda_k| < \varepsilon/n$. Tako dobimo števno gosto množico

$$\left\{ \sum_{k=1}^n \lambda_k e_k \mid n \in \mathbb{N}, \lambda_k \in \mathbb{Q} + i\mathbb{Q} \right\}. \quad \square$$

Vprašanje 40. Karakteriziraj separabilnost neskončnorazsežnega Hilbertovega prostora in dokaži karakterizacijo.

Definicija 4.4.36. Linearna preslikava $U : H \rightarrow K$ je **IZOMORFIZEM** Hilbertovih prostorov (tudi **UNITARNI OPERATOR**), če je surjektivna in če velja

$$\langle Ux, Uy \rangle = \langle x, y \rangle.$$

Trditev 4.4.37. Naj bo $U : X \rightarrow Y$ linearna izometrija med prostoroma s skalarnim produktom. Tedaj U ohranja skalarni produkt.

Dokaz. Računamo

$$\begin{aligned} \|U(x+y)\|^2 &= \|Ux\|^2 + \|Uy\|^2 + 2 \operatorname{Re} \langle Ux, Uy \rangle \\ \|U(x+y)\|^2 &= \|x+y\|^2 = \|x\|^2 + \|y\|^2 + 2 \operatorname{Re} \langle x, y \rangle \end{aligned}$$

torej $\operatorname{Re} \langle x, y \rangle = \operatorname{Re} \langle Ux, Uy \rangle$. Če namesto x pišemo ix , dobimo še $-\operatorname{Im} \langle x, y \rangle = -\operatorname{Im} \langle Ux, Uy \rangle$. \square

Vprašanje 41. Pokaži: linearna izometrija ohranja skalarni produkt.

Lema 4.4.38. Naj bo $U : H \rightarrow K$ izomorfizem Hilbertovih prostorov. Potem U slika **KONS** v **KONS**.

Dokaz. Naj bo $\{e_i\}_{i \in I}$ **KONS** za H . Potem je $\langle Ue_i, Ue_j \rangle = \langle e_i, e_j \rangle = \delta_{ij}$. Če je $y \in K$ tak, da je $y \perp Ue_i$ za vse i , potem je $0 = \langle y, Ue_i \rangle = \langle U^{-1}y, e_i \rangle$, torej $y = 0$. \square

Izrek 4.4.39. Hilbertova prostora H in K sta izomorfna natanko tedaj, ko je $\dim H = \dim K$.

Dokaz. V desno sledi iz leme. V levo: Dokazali bomo, da je $H \cong l^2(I)$, kjer je $(e_i)_{i \in I}$ **KONS** za H . Definiramo $U : H \rightarrow l^2(I)$ z $Ux = \hat{x}$, kjer je

$$\hat{x} : i \mapsto \langle x, e_i \rangle.$$

Potem je U linearna izometrija, saj

$$\|Ux\|^2 = \sum_{i \in I} |\hat{x}(i)|^2 = \sum_{i \in I} |\langle x, e_i \rangle|^2 = \|x\|^2$$

po Parsevalovi enakosti, hkrati pa je U tudi surjektivna, saj za $f \in l^2(I)$ lahko definiramo

$$x = \sum_{i \in I} f(i)e_i,$$

kar konvergira, ker je $f \in l^2(I)$. Seveda $\hat{x} = f$. Bijekcijo med indeksnima množicama lahko razširimo do izomorfizma Hilbertovih prostorov. \square

Vprašanje 42. Pokaži: Hilbertova prostora sta izomorfna natanko tedaj, ko imata isto dimenzijo.

Posledica 4.4.40. *Neskončnorazsežen separabilen Hilbertov prostor je izomorfen l^2 .*

Naj bosta H in K Hilbertova prostora. Produkt $H \times K$ opremimo s skalarnim produktom

$$\langle (x_1, y_1), (x_2, y_2) \rangle = \langle x_1, x_2 \rangle + \langle y_1, y_2 \rangle,$$

s čimer je prostor $H \times K$ poln, torej Hilbertov. Označimo $H \times K = H \oplus K$, konstrukciji pravimo ORTOGONALNA DIREKTNA VSOTA.

Za neskončne direktne vsote množico

$$\left\{ (x_n)_n \mid x_n \in H_n, \sum_{n=1}^{\infty} \|x_n\|^2 < \infty \right\},$$

kjer so H_n Hilbertovi prostori, opremimo s skalarnim produktom

$$\langle (x_n)_n, (y_n)_n \rangle = \sum_{n=1}^{\infty} \langle x_n, y_n \rangle.$$

Dobljen prostor označimo z

$$\bigoplus_{n=1}^{\infty} H_n,$$

in ga imenujemo ORTOGONALNA DIREKTNA VSOTA prostorov $(H_n)_n$. Je tudi Hilbertov prostor.

Vprašanje 43. Definiraj neskončno direktno vsoto Hilbertovih prostorov.

4.4.2 Stone-Weierstrassov izrek

Lema 4.4.41. *Naj bo X neprazna množica in V vektorski prostor funkcij na X , ki loči točke in vsebuje konstante. Tedaj za vsaka različna $x, y \in X$ in $\alpha, \beta \in \mathbb{R}$ obstaja $f \in V$, da je $f(x) = \alpha$ ter $f(y) = \beta$.*

Lema 4.4.42. *Naj bo K kompakten Hausdorffov prostor in $V \subseteq \mathcal{C}(K)$ realen vektorski prostor, ki je podmreža v $\mathcal{C}(K)$. Naj V vsebuje konstante in loči točke na K . Tedaj za vsak $g \in \mathcal{C}(K)$, $a \in K$ ter $\varepsilon > 0$ obstaja $f \in V$, da je $f(a) = g(a)$ in $f(x) > g(x) - \varepsilon$ za vse $x \in K$.*

Dokaz. Po prejšnji lemi za vsak $x \in K$ obstaja $f_x \in V$, da je $f_x(a) = g(a)$ in $f_x(x) = g(x)$. Zaradi zveznosti zato obstaja odprta okolica $U_x \ni x$, da je $f_x(y) > g(y) - \varepsilon$ za vse $y \in U_x$. Dobimo odprto pokritje K , zaradi kompaktnosti obstaja končno podpokritje $U_{x_1} \cup \dots \cup U_{x_n} = K$. Definiramo $f = \max\{f_{x_1}, \dots, f_{x_n}\}$. \square

Izrek 4.4.43 (mrežni Stone-Weierstrass). *Naj bo K kompakten Hausdorffov prostor ter V realen vektorski podprostor $\mathcal{C}(K)$, ki je podmreža, loči točke K in vsebuje konstante. Tedaj je V gost v $\mathcal{C}(K)$ glede na supremum normo.*

Dokaz. Naj bo $g \in \mathcal{C}(K)$ in $\varepsilon > 0$. Iščemo $f \in V$, da je $\|f - g\|_\infty < \varepsilon$, oziroma $f - g < \varepsilon$ in $g - f < \varepsilon$. Po prejšnji lemi za vsak $x \in K$ obstaja $f_x \in V$, da je $f_x > g - \varepsilon$ in $f_x(x) = g(x)$. Zaradi zveznosti obstaja odprta okolica V_x za x , da je $|f_x(y) - g(y)| < \varepsilon$ za vse $y \in V_x$. Množice V_x tvorijo pokritje za K , torej obstaja končno podpokritje $V_{x_1} \cup \dots \cup V_{x_n}$. Potem definiramo $f = \min\{f_{x_1}, \dots, f_{x_n}\}$. Za vsak $m = 1, \dots, n$ velja $f_{x_m} > g - \varepsilon$, torej $f > g - \varepsilon$. Za poljuben $y \in K$ obstaja m , da je $y \in V_m$, torej $f(y) \leq f_{x_m}(y) < g(y) + \varepsilon$. \square

Vprašanje 44. Povej in dokaži mrežni Stone-Weierstrassov izrek.

Lema 4.4.44. *Obstaja zaporedje polinomov, ki na $[0, 1]$ konvergira enakomerno proti funkciji \sqrt{x} .*

Dokaz. Funkcijo $x \mapsto 1 - \sqrt{1 - x}$ razvijemo v Taylorjevo vrsto na $[0, 1]$. Velja

$$1 - (1 - x)^{1/2} = 1 - \sum_{k=0}^{\infty} \binom{1/2}{k} (-1)^k x^k.$$

S $S_n(x)$ označimo n -to delno vsoto zgornje vrste. Za $x \in (0, 1)$ je zaporedje $(S_n(x))_n$ strogo naraščajoče in velja $\lim S_n(x) \leq 1$, saj je $1 - (1 - x)^{1/2} \leq 1$ za te x . Ker so S_n zvezni, je

$$S_n(1) = \lim_{x \rightarrow 1} S_n(x) \leq 1,$$

torej je zaporedje $(S_n(1))_n$ naraščajoče in navzgor omejeno. Definirajmo $f(x) = \lim S_n(x)$.

Na $[0, 1)$ se f in $1 - \sqrt{1 - x}$ ujemata, hkrati pa je za $x \in [0, 1]$ tudi

$$0 \leq f(x) - S_n(x) = \sum_{k=n+1}^{\infty} \binom{1/2}{k} (-1)^k x^k \leq \sum_{k=n+1}^{\infty} \binom{1/2}{k} (-1)^k = f(1) - S_n(1) \xrightarrow{n \rightarrow \infty} 0.$$

Torej S_n konvergira k f enakomerno na $[0, 1]$, in je zato f zvezna in se na tem intervalu ujema z $1 - \sqrt{1 - x}$. Dokazali smo, da obstaja zaporedje polinomov, ki konvergirajo k f enakomerno. Enostavno ga lahko transformiramo v zaporedje, ki enakomerno konvergira k \sqrt{x} . \square

Izrek 4.4.45 (Stone-Weierstrass, realna verzija). *Naj bo K kompakten Hausdorffov prostor in $A \subseteq \mathcal{C}(K)$ podalgebra, ki loči točke in vsebuje konstante. Tedaj je A gosta v $\mathcal{C}(K)$.*

Dokaz. Pokazali bomo, da je \overline{A} podmreža v $\mathcal{C}(K)$. Potem bo po mrežni različici izreka gosta v $\mathcal{C}(K)$. Ker je zaprta, bo $\overline{A} = \mathcal{C}(K)$.

Očitno je \overline{A} podalgebra. Ker velja $\max\{f, g\} = \frac{1}{2}(f + g + |f - g|)$ in $\min\{f, g\} = \frac{1}{2}(f + g - |f - g|)$, je dovolj pokazati, da je \overline{A} zaprta za absolutne vrednosti. Naj bo $f \in \overline{A}$. Oglejmo si $g = f / \|f\|_\infty$. Ker je $g^2 \leq 1$, in ker po lemi obstaja zaporedje polinomov $(p_n)_n$, ki konvergirajo enakomerno na $[0, 1]$ proti \sqrt{x} , velja $(p_n \circ g^2)(x) \rightarrow |g(x)|$ enakomerno. Torej $|g| \in \overline{A}$, ampak $|g| = |f| / \|f\|_\infty$, torej tudi $|f| \in \overline{A}$. \square

Vprašanje 45. Povej in pokaži realni Stone-Weierstrassov izrek.

Izrek 4.4.46 (Weierstrass). *Naj bo $K \subseteq \mathbb{R}$ kompaktna in $f : K \rightarrow \mathbb{R}$ zvezna. Tedaj za vsak $\varepsilon > 0$ obstaja polinom p , da je $|f(x) - p(x)| < \varepsilon$ za vse $x \in K$.*

Vprašanje 46. Povej Weierstrassov izrek.

Izrek 4.4.47 (Stone-Weierstrass, kompleksna verzija). *Naj bo K kompakten Hausdorffov prostor in $A \subseteq \mathcal{C}(K)$ podalgebra v algebri kompleksnih funkcij, ki*

- *loči točke K ,*
- *vsebuje konstante,*
- *je sebi-adjungirana: $f \in A$ pomeni $\overline{f} \in A$.*

Tedaj je A gosta v $\mathcal{C}(K)$.

Dokaz. Naj bo A_0 algebra realnih funkcij, ki so vsebovane v A . Očitno A_0 vsebuje konstante. Za $x, y \in K$ obstaja $f \in A$, da je $f(x) \neq f(y)$, torej ena od realnih funkcij $\frac{f+\overline{f}}{2}, \frac{f-\overline{f}}{2i} \in A$ loči točki x, y . Torej A_0 loči točke. Po realni verziji izreka je A_0 gosta v $\mathcal{C}(K, \mathbb{R})$.

Naj bo $f \in \mathcal{C}(K, \mathbb{C})$ poljubna in $\varepsilon > 0$. Pišimo $f = g + ih$ za realni g, h . Po ravno dokazanem obstajata $g_1, h_1 \in A_0$, da je $\|g - g_1\| < \varepsilon/2$ in $\|h - h_1\| < \varepsilon/2$. Potem je $\|f - (g_1 + ih_1)\| < \varepsilon$. \square

Vprašanje 47. Povej in dokaži kompleksni Stone-Weierstrassov izrek.

4.5 Omejeni operatorji med Hilbertovimi prostori

Definicija 4.5.1. Naj bosta H, K Hilbertova prostora. Preslikava $u : H \times K \rightarrow \mathbb{F}$ se imenuje SESKVILINEARNA FORMA, če velja

- $u(\alpha x + \beta y, z) = \alpha u(x, z) + \beta u(y, z),$

- $u(x, \alpha y + \beta z) = \overline{\alpha}u(x, y) + \overline{\beta}u(x, z)$.

Definicija 4.5.2. Seskvilinearna forma je ZVEZNA (ali OMEJENA), če obstaja $M \geq 0$, da $|u(x, y)| \leq M \|x\| \|y\|$.

Vprašanje 48. Kaj je seskvilinearna forma? Kdaj je omejena?

Primer. Za $A \in B(H, K)$ definiramo $u(x, y) = \langle Ax, y \rangle$. To je seskvilinearna forma, velja $|u(x, y)| \leq \|A\| \|x\| \|y\|$.

Izrek 4.5.3. Naj bo $u : H \times K \rightarrow \mathbb{F}$ omejena seskvilinearna forma. Potem obstajata natanko določeni $A \in B(H, K)$ in $B \in B(K, H)$, da

$$u(x, y) = \langle Ax, y \rangle = \langle x, By \rangle.$$

Dokaz. Fiksiramo $x \in H$ in definiramo $f_x : K \rightarrow \mathbb{F}$ z $f_x(y) = \overline{u(x, y)}$. To je očitno omejen linearen funkcional na K . Po Rieszovem izreku obstaja natanko določen $z_x \in K$, da je

$$\overline{u(x, y)} = f_x(y) = \langle y, z_x \rangle$$

za vsak $y \in K$. Definiramo $z_x = Ax$. To nam da preslikavo $A : H \rightarrow K$, da je $\overline{u(x, y)} = \langle y, Ax \rangle = \overline{\langle Ax, y \rangle}$. Enostavno lahko preverimo, da je A linearna, ker velja

$$\|Ax\| = \|z_x\| = \|f_x\| \leq M \|x\|,$$

pa je tudi omejena (tu M pride iz definicije omejene seskvilinearne forme). Če je $\langle Ax, y \rangle = \langle A'x, y \rangle$, potem velja $\langle Ax - A'x, y \rangle = 0$ za vsak y , torej $Ax = A'x$ za vse x . Sledi, da je A enolično določena. Podobno za B . \square

Vprašanje 49. Pokaži, da lahko vsako omejeno seskvilinearno formo izrazimo s skalarnim produktom.

Definicija 4.5.4. Naj bo A omejen operator $H \rightarrow K$. Tedaj operatorju $B : H \rightarrow K$, za katerega velja $\langle Ax, y \rangle = \langle x, By \rangle$, pravimo ADJUNGIRANI OPERATOR operatorja A . Označimo ga z A^* .

Trditev 4.5.5. Preslikava $U \in B(H, K)$ je izomorfizem Hilbertovih prostorov natanko tedaj, ko je U obrnljiv in $U^* = U^{-1}$.

Dokaz. V levo: Če je U obrnljiv, je surjektiven. Potem je

$$\langle Ux, Uy \rangle = \langle x, U^*Uy \rangle = \langle x, U^{-1}Uy \rangle = \langle x, y \rangle,$$

torej U ohranja skalarni produkt in je zato izomorfizem.

V desno: Če je U izomorfizem, je obrnljiv. Velja

$$\langle x, y \rangle = \langle Ux, Uy \rangle = \langle x, U^*Uy \rangle$$

za poljubna x, y , torej je $U^* = U^{-1}$. \square

Vprašanje 50. Pokaži: U je izomorfizem Hilbertovih prostorov natanko tedaj, ko je obrnljiv in $U^* = U^{-1}$.

Opomba. Če sta $A, B \in B(H, K)$, potem velja

- $(A + B)^* = A^* + B^*$,
- $(\alpha A)^* = \bar{\alpha} A^*$,
- $A^{**} = A$.

Zadnje je res, ker je

$$\langle Ax, y \rangle = \langle x, A^*y \rangle = \overline{\langle A^*y, x \rangle} = \overline{\langle y, A^{**}x \rangle} = \langle A^{**}x, y \rangle.$$

Opomba. Preslikava $i : A \mapsto A^*$ je involucija.

Trditev 4.5.6. Naj bosta $A \in B(H, K)$ in $B \in B(K, L)$ omejena operatorja. Tedaj $(BA)^* = A^*B^*$.

Dokaz. Preprost račun. □

Posledica 4.5.7. Operator $A \in B(H, K)$ je obrnljiv natanko tedaj, ko je $A^* \in B(K, H)$ obrnljiv. Če je A obrnljiv, velja $(A^*)^{-1} = (A^{-1})^*$.

Dokaz. Operator A je obrnljiv natanko tedaj, ko obstaja $B \in B(K, H)$, da velja $AB = I_K$ in $BA = I_H$. Potem je

$$\begin{aligned} (AB)^* &= B^*A^* = I_K, \\ (BA)^* &= A^*B^* = I_H, \end{aligned}$$

torej je A^* obrnljiv in $(A^*)^{-1} = B^* = (A^{-1})^*$. Podobno v drugo smer, kjer upoštevamo še $A^{**} = A$. □

Trditev 4.5.8. Če je $A \in B(H, K)$, je $\ker A^* = (\operatorname{im} A)^\perp$.

Dokaz. Velja

$$x \in \ker A^* \Leftrightarrow A^*x = 0 \Leftrightarrow \forall y. \langle A^*x, y \rangle = 0 \Leftrightarrow \forall y. \langle x, Ay \rangle = 0 \Leftrightarrow x \in (\operatorname{im} A)^\perp. \quad \square$$

Vprašanje 51. Kaj je $\ker A^*$?

Posledica 4.5.9. Za $A \in B(H, K)$ velja

- $\ker A = (\operatorname{im} A^*)^\perp$,
- $(\ker A)^\perp = \overline{\operatorname{im} A^*}$,
- $(\ker A^*)^\perp = \overline{\operatorname{im} A}$.

Dokaz. Prva točka je očitna. Če na njej uporabimo komplement in upoštevamo $X^{\perp\perp} = \overline{\text{Lin } X}$, dobimo drugo točko. Podobno za tretje. \square

Definicija 4.5.10. Operator $A \in B(H)$ je SEBI ADJUNGIRAN, če je $A^* = A$. Je NORMALEN, če je $A^*A = AA^*$. Je UNITAREN, če je $A^*A = AA^* = I$.

Opomba. Operator A je unitaren natanko tedaj, ko je A izomorfizem prostora H .

Opomba. Vsak unitaren operator je normalen.

Opomba. Sebi adjungiran operator je vedno normalen.

Če je $A \in B(H)$, lahko definiramo

$$\begin{aligned}\text{Re } A &= \frac{A + A^*}{2}, \\ \text{Im } A &= \frac{A - A^*}{2i},\end{aligned}$$

tako dobimo $A = \text{Re } A + i \text{Im } A$. Oba ta operatorja sta sebi adjungirana.

Trditev 4.5.11. Naj bo H kompleksen Hilbertov prostor in $A \in B(H)$. Tedaj je $A = A^*$ natanko tedaj, ko je $\langle Ax, x \rangle \in \mathbb{R}$ za vsak $x \in H$.

Dokaz. V desno: $\langle Ax, x \rangle = \langle x, Ax \rangle = \overline{\langle Ax, x \rangle}$.

V levo: Po predpostavki je $\langle A(x + y), x + y \rangle \in \mathbb{R}$. Velja

$$\langle A(x + y), x + y \rangle = \langle Ax, x \rangle + \langle Ay, y \rangle + \langle Ax, y \rangle + \langle Ay, x \rangle,$$

torej $\langle Ax, y \rangle + \langle Ay, x \rangle \in \mathbb{R}$ za vsaka x, y . Pišimo $\langle Ax, y \rangle = \alpha + i\beta$ in $\langle Ay, x \rangle = \gamma - i\beta$.

Če menjamo $y \rightarrow iy$, dobimo $i \langle Ay, x \rangle - i \langle Ax, y \rangle \in \mathbb{R}$. To je enako

$$i \langle Ay, x \rangle - i \langle Ax, y \rangle = i(\gamma - i\beta - \alpha - i\beta) = i(\gamma - \alpha) + 2\beta,$$

torej $\gamma = \alpha$. \square

Vprašanje 52. Pokaži: če je $\langle Ax, x \rangle \in \mathbb{R}$ za vsak x , je A sebi adjungiran.

Izrek 4.5.12. Naj bo H Hilbertov prostor in $A \in B(H)$ sebi adjungiran operator. Tedaj velja

$$\|A\| = w(A) := \sup_{\|x\|=1} |\langle Ax, x \rangle|.$$

Dokaz. Velja $|\langle Ax, x \rangle| \leq \|Ax\| \|x\| \leq \|A\| \|x\|^2$, torej je $w(A) \leq \|A\|$. Računamo

$$\begin{aligned}\langle A(x + y), x + y \rangle - \langle A(x - y), x - y \rangle &= 2 \langle Ax, y \rangle + 2 \langle Ay, x \rangle \\ &= 2 \langle y, Ax \rangle + 2 \langle Ax, y \rangle \\ &= 4 \text{Re } \langle Ax, y \rangle,\end{aligned}$$

torej

$$\begin{aligned} 4|\operatorname{Re}\langle Ax, y \rangle| &\leq |\langle A(x+y), x+y \rangle| + |\langle A(x-y), x-y \rangle| \\ &\leq w(A) (\|x+y\|^2 + \|x-y\|^2) \\ &= 2w(A) (\|x\|^2 + \|y\|^2) \end{aligned}$$

oziroma

$$|\operatorname{Re}\langle Ax, y \rangle| \leq \frac{1}{2}w(A) (\|x\|^2 + \|y\|^2).$$

Izberemo x z $\|x\| = 1$. Če je $A = 0$, izrek velja, sicer pa lahko vzamemo tak x , da $Ax \neq 0$. Potem nastavimo $y = \frac{Ax}{\|Ax\|}$, in dobimo

$$\left| \operatorname{Re} \left\langle Ax, \frac{Ax}{\|Ax\|} \right\rangle \right| \leq w(A),$$

zato $\|Ax\| \leq w(A)$ za vsak x z $\|x\| = 1$ in $Ax \neq 0$. Sedaj lahko naredimo supremum po enotski sferi. \square

Vprašanje 53. Kako lahko izraziš normo sebi adjungiranega operatorja? Dokaži.

Trditev 4.5.13. Naj bo H kompleksen Hilbertov prostor in $A \in B(H)$ tak, da je $\langle Ax, x \rangle = 0$ za vsak $x \in H$. Tedaj je $A = 0$.

Dokaz. Ker je H kompleksen, je $A = A^*$. Po izreku je

$$\|A\| = \sup_{\|x\|=1} |\langle Ax, x \rangle| = 0. \quad \square$$

Izrek 4.5.14. Za $A \in B(H)$ velja $\|A^*A\| = \|A\|^2$.

Dokaz. Računamo

$$\|A^*A\| = \sup_{\|x\|=1} |\langle A^*Ax, x \rangle| = \sup_{\|x\|=1} |\langle Ax, Ax \rangle| = \sup_{\|x\|=1} \|Ax\|^2 = \|A\|^2. \quad \square$$

Opomba. Tej enakosti pravimo C^* -aksiom.

Vprašanje 54. Pokaži, da v $B(H)$ velja C^* -aksiom.

Trditev 4.5.15. Za $A \in B(H)$ so naslednje trditve ekvivalentne:

- A je normalen,
- $\|A^*x\| = \|Ax\|$ za vsak $x \in H$,
- če je $\mathbb{F} = \mathbb{C}$, potem je $\operatorname{Re} A \operatorname{Im} A = \operatorname{Im} A \operatorname{Re} A$.

Dokaz. Ekvivalenca 1 in 2: Velja

$$\|A^*x\| = \|Ax\| \Leftrightarrow \langle A^*x, A^*x \rangle = \langle Ax, Ax \rangle \Leftrightarrow \langle AA^*x, x \rangle = \langle A^*Ax, x \rangle \Leftrightarrow \langle Bx, x \rangle = 0$$

za $B = AA^* - A^*A$. Zgornje velja za vse $x \in H$ natanko tedaj, ko je $B = 0$.

Ekvivalenca 1 in 3: Pišimo $A = H + iK$ za $H = \operatorname{Re} A$ in $K = \operatorname{Im} A$. Potem je A normalen natanko tedaj, ko je $(H - iK)(H + iK) = (H + iK)(H - iK)$, oziroma $HK = KH$. \square

Vprašanje 55. Karakteriziraj normalnost operatorja in dokaži karakterizacijo.

Posledica 4.5.16. Naj bo $A \in B(H)$ normalen in $\lambda \in \mathbb{F}$. Tedaj je $\ker(A^* - \lambda I) = \ker(A - \bar{\lambda}I)$.

Definicija 4.5.17. Operator $A \in B(H)$ je POZITIVNO SEMIDEFINITEN, če je sebi adjungiran in $\langle Ax, x \rangle \geq 0$ za vsak $x \in H$.

Opomba. Če je A pozitivno definiten, je injektiven in zato

$$\ker A = \ker A^* = (\operatorname{im} A)^\perp,$$

torej $\overline{\operatorname{im} A} = (\operatorname{im} A)^{\perp\perp} = (\ker A)^\perp = H$, torej je $\operatorname{im} A$ gost v H .

Lema 4.5.18. Naj bosta $A, B \in B(H)$ pozitivno semidefinitna. Potem je $A + B \geq 0$. Če je $A > 0$, je $A + B > 0$.

Definiramo množici

- $B(H)_{sa} = \{A \in B(H) \mid A = A^*\},$
- $B(H)_{sa}^+ = \{A \in B(H)_{sa} \mid A \geq 0\}.$

Trditev 4.5.19. Relacija $A \geq B \Leftrightarrow A - B \geq 0$ je delna urejenost na $B(H)_{sa}$.

Definicija 4.5.20. Podmnožica $C \subseteq X$ realnega vektorskega prostora X je STOŽEC, če velja

- $C + C \subseteq C,$
- $\lambda C \subseteq C$ za vsak $\lambda \geq 0,$
- $C \cap (-C) = \{0\}.$

Vprašanje 56. Kaj je stožec v realnem vektorskem prostoru?

Trditev 4.5.21. Množica $B(H)_{sa}^+$ je stožec v $B(H)_{sa}$.

Trditev 4.5.22. Naj bo $A \in B(H)$ sebi adjungiran in $B \in B(K, H)$.

- če $A \geq 0$, potem $B^*AB \in B(K)_{sa}^+,$
- velja $-\|A\|I \leq A \leq \|A\|I,$
- $B^*B \in B(K)_{sa}^+$ in $BB^* \in B(H)_{sa}^+.$

Izrek 4.5.23. Naj bo $A \in B(H)$. Tedaj obstaja natanko en $B \in B(H)_{sa}^+$, da je $A = B^2$.

4.5.1 Idempotenti in invariantni podprostor

Če je P idempotent, velja

$$x = \underbrace{x - Px}_{\in \ker P} + \underbrace{Px}_{\in \operatorname{im} P}.$$

Ker sta $\ker P$ in $\operatorname{im} P$ zaprta v H , velja $H = \operatorname{im} P \oplus \ker P$.

Definicija 4.5.24. Idempotent $P \in B(H)$ je ORTOGONALNI PROJEKTOR, če je $\ker P = (\operatorname{im} P)^\perp$.

Izrek 4.5.25. Naj bo $P \in B(H)$ neničelni idempotent. Naslednje trditve so ekvivalentne:

- P je ortogonalni projektor,
- $\|P\| = 1$,
- $P^* = P$,
- P je normalen,
- $\langle Px, x \rangle \geq 0$ za vsak $x \in H$.

Dokaz. 1 v 2: Vemo že, da je $\|Px\| \leq \|x\|$ za vsak $x \in H$. Če izberemo $x \in \operatorname{im} P$, je $Px = x$, torej $\|P\| = 1$.

2 v 1: Vzemimo $x \in (\ker P)^\perp$, da je $x - Px \in \ker P$. Sledi $\langle x - Px, x \rangle = 0$, oziroma $\langle Px, x \rangle = \|x\|^2$. Torej

$$\|x\|^2 = \langle Px, x \rangle \leq \|Px\| \|x\| \leq \|P\| \|x\|^2 = \|x\|^2,$$

iz česar sledi $\|x\| = \|Px\| = \sqrt{\langle Px, x \rangle}$. Potem je

$$\|x - Px\|^2 = \|x\|^2 + \|Px\|^2 - 2 \operatorname{Re} \langle x, Px \rangle = \|x\|^2 + \|Px\|^2 - 2 \langle Px, x \rangle = 0.$$

Torej $x = Px \in \operatorname{im} P$, oziroma $(\ker P)^\perp \subseteq \operatorname{im} P$.

Za drugo inkluzijo vzemimo $y \in \operatorname{im} P$ in zapišimo $y = y_1 + y_2$ za $y_1 \in \ker P$, $y_2 \in (\ker P)^\perp$. Potem je

$$y = Py = P(y_1 + y_2) = Py_2 = y_2 \in (\ker P)^\perp,$$

torej $\operatorname{im} P \subseteq (\ker P)^\perp$.

1 v 3: Za $x, y \in H$ zapišimo $x = x_1 + x_2$, $y = y_1 + y_2$, kjer sta $x_1, y_1 \in \ker P$ ter $x_2, y_2 \in (\ker P)^\perp$. Računamo

$$\begin{aligned} \langle P^*x, y \rangle &= \langle x, Py \rangle = \langle x_1 + x_2, Py_2 \rangle = \langle x_1 + x_2, y_2 \rangle = \langle x_2, y_2 \rangle \\ \langle Px, y \rangle &= \langle Px_2, y_1 + y_2 \rangle = \langle x_2, y_1 + y_2 \rangle = \langle x_2, y_2 \rangle. \end{aligned}$$

3 v 4 je očitno. 4 v 1: Operator P je normalen, torej je $(\operatorname{im} P)^\perp = (\ker P)^* = \ker P$. Torej $\operatorname{im} P = P^\perp$.

3 v 5: Računamo

$$\langle Px, x \rangle = \langle P^2x, x \rangle = \langle Px, P^*x \rangle = \langle Px, Px \rangle \geq 0.$$

5 v 1 izpustimo. □

Vprašanje 57. Kdaj je idempotent $P \in B(H)$ ortogonalni projektor? Karakteriziraj in dokaži.

Definicija 4.5.26. Zaprt podprostor M normiranega prostora X je INVARIANTEN za $A \in B(X)$, če je $AM \subseteq M$.

Trditev 4.5.27. Zaprt podprostor M je invarianten za $A \in B(H)$ natanko tedaj, ko je M^\perp invarianten za A^* .

Dokaz. Naj bo M invarianten za A . Če je $y \in M^\perp$, potem je $\langle A^*y, x \rangle = \langle y, Ax \rangle = 0$, in zato $A^*y \in M^\perp$. Drugo implikacijo dobimo, ko zamenjamo vlogi M in M^\perp ter A in A^* . □

Posledica 4.5.28. Naj bo $A \in B(H)_{sa}$. Potem je zaprt podprostor M invarianten za A natanko tedaj, ko je M^\perp invarianten za A .

4.5.2 Kompaktni operatorji

Definicija 4.5.29. Naj bo $T : X \rightarrow Y$ linearna preslikava med normiranimi prostoroma. Operator T je KOMPAKTEN, če je $T(B_X)$ kompakt v Y .

Opomba. $B_X = \{x \in X \mid \|x\| = 1\}$

Opomba. Vsak kompakten operator je omejen.

Označimo $K(X, Y) = \{T : X \rightarrow Y \mid T \text{ kompakten}\}$.

Izrek 4.5.30. Za metrični prostor M so naslednje trditve ekvivalentne:

- M je kompakten,
- vsako neskončno zaporedje v M ima stekališče,
- M je poln in povsem omejen (za vsak $\varepsilon > 0$ lahko M pokrijemo s končno mnogo krogli polmera ε).

Izrek 4.5.31. Naj bo $T : X \rightarrow Y$ linearen operator. Naslednje trditve so ekvivalentne:

- T je kompakten,
- T preslika omejene množice v relativno kompaktne množice,
- če je $(x_m)_m$ omejeno zaporedje v X , potem ima $(Tx_m)_m$ stekališče v Y .

Vprašanje 58. Definiraj in karakteriziraj kompaktnost linearnega operatorja.

Trditev 4.5.32. Naj bosta X, Y Banachova prostora. Potem velja:

- $K(X, Y)$ je zaprt podprostor v $B(X, Y)$,
- za vsake $K \in K(X, Y)$, $A \in B(X)$ ter $B \in B(Y)$ sta $KA, BK \in K(X, Y)$.

Dokaz. Samo prva točka. Naj bosta $K_1, K_2 \in K(X, Y)$ ter $\lambda, \mu \in \mathbb{F}$. Naj bo $(x_m)_m$ omejeno zaporedje v X . Ker je K_1 kompakten, obstaja podzaporedje $(x_{n_k})_k$, da $(K_1 x_{n_k})_k$ konvergira. Podobno imamo podzaporedje $(x_{n_{k_j}})_j$, da je $(K_2 x_{n_{k_j}})_j$ konvergentno. Potem $(\mu K_1 x_{n_{k_j}} + \lambda K_2 x_{n_{k_j}})_j$ konvergira v Y .

Naj bo sedaj $(K_n)_n$ zaporedje kompaktnih operatorjev, ki konvergira k nekemu K . Pokažimo, da je K kompakten. Ker je Y Banachov, je $\overline{K(B_X)}$ kompaktno natanko tedaj, ko je $K(B_X)$ povsem omejena, oziroma ko je $K(B_X)$ povsem omejena. Naj bo $\varepsilon > 0$. Obstaja $n \in \mathbb{N}$, da je $\|K - K_n\| < \varepsilon/3$. Ker je K_n kompakten, obstajajo $x_1, \dots, x_m \in B_X$, da je

$$K_n(B_X) \subseteq \bigcup_{i=1}^m \mathring{B}(Kx_i, \varepsilon/3).$$

Pokažimo še, da je

$$K(B_X) \subseteq \bigcup_{i=1}^m \mathring{B}(x_i, \varepsilon).$$

Če je $x \in B_X$, potem obstaja j , da je $\|K_n x_j - K_n x\| < \varepsilon/3$. Velja

$$\begin{aligned} \|Kx_j - Kx\| &\leq \|Kx_j - K_n x_j\| + \|K_n x_j - K_n x\| + \|K_n x - Kx\| \\ &\leq \|K - K_n\| (\|x_j\| + \|x\|) + \|K_n(x_j - x)\| \\ &< \varepsilon. \end{aligned}$$

□

Vprašanje 59. Pokaži, da je $K(X, Y)$ zaprt podprostor $B(X, Y)$ za Banachova X, Y .

Trditev 4.5.33. Naj bo $A \in B(X)$ operator z $\dim \operatorname{im} A < \infty$. Potem je A kompakten.

Dokaz. Velja $A(B_X) \subseteq \operatorname{im} A$. Ker je množica omejena v končnorazsežnem prostoru, je relativno kompaktna. □

Definicija 4.5.34. Operator $A \in B(X, Y)$ je KONČNEGA RANGA, če je $\dim \operatorname{im} A < \infty$. V tem primeru je $\operatorname{rang} A = \dim \operatorname{im} A$.

Definicija 4.5.35. Množico vseh operatorjev končnega ranga $X \rightarrow Y$ označimo z $F(X, Y)$.

Vprašanje 60. Pokaži, da je $F(X, Y) \subseteq K(X, Y)$.

Izrek 4.5.36. Naj bo X normiran prostor. Tedaj je B_X kompaktno natanko tedaj, ko je $\dim X < \infty$.

Posledica 4.5.37. *Identiteta $\text{id} : X \rightarrow X$ je kompakten operator natanko tedaj, ko je $\dim X < \infty$.*

Naj bo X neskončnorazsežen Banachov prostor. Tedaj je $\text{id} \in B(X) \setminus K(X)$. Tvorimo prostor $B(X)/K(X)$. Ker je X Banachov, je $B(X)/K(X)$ Banachov, ker pa je $K(X)$ zaprti ideal v Banachovi algebri $B(X)$, je tudi $B(X)/K(X)$ Banachova algebra. Pravimo ji CALKINOVA ALGEBRA.

Trditev 4.5.38. *Naj bo X normiran in Y Banachov. Naj bo $A \in K(X, Y)$ ter \bar{A} enolična razširitev A na \bar{X} po zveznosti. Tedaj je $\bar{A} \in K(\bar{X}, \bar{Y})$.*

4.5.3 Izrek Arzela-Ascoli

Definicija 4.5.39. Naj bo K kompakten Hausdorffov prostor. Množica $H \subseteq \mathcal{C}(K)$ je ENAKOZVEZNA, če za vsaka $x \in K$ in $\varepsilon > 0$ obstaja odprta množica $U_x \ni x$, da je $|f(y) - f(x)| < \varepsilon$ za vse $y \in U_x$ in $f \in H$.

Izrek 4.5.40 (Arzela-Ascoli). *Naj bo K kompakten Hausdorffov prostor in $H \subseteq \mathcal{C}(K)$ družina funkcij. Tedaj je H relativno kompaktna natanko tedaj, ko je enakozvezna in po točkah omejena.*

Vprašanje 61. Povej izrek Arzela-Ascoli.

Izrek 4.5.41. *Naj bo $(f_n)_n \subseteq \mathcal{C}([a, b])$ tako zaporedje, da za vse $\varepsilon > 0$ obstaja $\delta > 0$, da za vsak n velja*

$$|y - x| < \delta \implies |f_n(x) - f_n(y)| < \varepsilon.$$

Če je $(f_n)_n$ omejeno zaporedje, je relativno kompaktno.

Dokaz. Naj bo $(f_n)_n$ kot v izreku. Naj bo $(x_n)_n$ množica racionalnih števil na $[a, b]$, postavljena v zaporedje. Zaporedje $(f_n(x_1))_n$ je omejeno v \mathbb{C} , torej ima konvergentno podzaporedje $(f_n^{(1)}(x_1))_n$. Zaporedje $(f_n^{(1)}(x_2))_n$ je omejeno, torej ima konvergentno podzaporedje $(f_n^{(2)}(x_2))_n$. Postopek ponavljamo, da dobimo neskončno zaporedij $(f_n^{(k)}(x_k))_n$. Tvorimo $\tilde{f}_n = f_n^{(n)}$. Po konstrukciji $(\tilde{f}_n(x_j))_n$ konvergira za poljuben $j \in \mathbb{N}$.

Naj bo $\varepsilon > 0$. Obstaja $\delta > 0$, da velja

$$|y - x| < \delta \implies |f_n(y) - f_n(x)| < \frac{\varepsilon}{3}.$$

Ker je $[a, b]$ kompaktno, obstaja $p \in \mathbb{N}$, da je

$$[a, b] \subseteq \bigcup_{j=1}^p (x_j - \delta, x_j + \delta).$$

Obstaja $n_\varepsilon \in \mathbb{N}$, da za vse $m, n \geq n_\varepsilon$ velja $|\tilde{f}_n(x_j) - \tilde{f}_m(x_j)| < \varepsilon/3$ za vse $j = 1, \dots, p$. Za $x \in [a, b]$ potem obstaja j , da $|x_j - x| < \delta$, torej

$$|\tilde{f}_n(x) - \tilde{f}_m(x)| \leq |\tilde{f}_n(x) - \tilde{f}_n(x_j)| + |\tilde{f}_n(x_j) - \tilde{f}_m(x_j)| + |\tilde{f}_m(x_j) - \tilde{f}_m(x)| < \varepsilon,$$

torej je $\|\tilde{f}_n - \tilde{f}_m\| \leq \varepsilon$ za $m, n \geq n_\varepsilon$. Torej je zaporedje $(\tilde{f}_n)_n$ Cauchyjevo in zato konvergira. Limita je stekališče $(f_n)_n$. \square

Vprašanje 62. Pokaži: če za omejeno zaporedje $(f_n)_n$ zveznih funkcij na kompaktnem intervalu za poljuben $\varepsilon > 0$ obstaja $\delta > 0$, da za vsak n iz $|x - y| < \delta$ sledi $|f_n(x) - f_n(y)| < \varepsilon$, potem ima zaporedje stekališče.

Trditev 4.5.42. Naj bo $H \subseteq \mathcal{C}([a, b])$ družina zvezno odvedljivih funkcij. Če je $\sup\{\|f\|_\infty \mid f \in H\} < \infty$, potem je H enakozvezna.

Dokaz. Za $f \in H$ je

$$|f(y) - f(x)| = |f'(\xi)(y - x)| \leq \sup_{g \in H} \|g\|_\infty |y - x|$$

po Lagrangeovem izreku, za neki ξ . \square

Izrek 4.5.43. Naj bo $k \in \mathcal{C}([a, b] \times [a, b])$ in K integralski operator, dan s predpisom

$$(Kf)(x) = \int_a^b k(x, y)f(y)dy.$$

Tedaj sta $K : (\mathcal{C}([a, b]), \|\cdot\|_\infty) \rightarrow (\mathcal{C}([a, b]), \|\cdot\|_\infty)$ in $K : (\mathcal{C}([a, b]), \|\cdot\|_2) \rightarrow (\mathcal{C}([a, b]), \|\cdot\|_2)$ kompaktna operatorja.

Dokaz. Ker je k enakomerno zvezna, za vsak $\varepsilon > 0$ obstaja $\delta > 0$, da iz $|(x, y) - (z, w)| < \delta$ sledi $|k(x, y) - k(z, w)| < \varepsilon$. Potem je

$$\begin{aligned} |Kf(x) - Kf(y)| &\leq \int_a^b |k(x, z) - k(y, z)| |f(z)| dz \\ &< \varepsilon \int_a^b |f(z)| dz \\ &= \varepsilon \langle 1, f \rangle_{L^2} \\ &\leq \varepsilon \sqrt{b-a} \|f\|_2, \end{aligned}$$

torej je $K(B_{\mathcal{C}([a, b])})$ je enakozvezna. Je tudi omejena:

$$|Kf(x)| \leq \int_a^b |k(x, y)f(y)| dy \leq \|k\|_\infty \|f\|_\infty (b-a).$$

Po izreku Arzela-Ascoli je slika krogle relativno kompaktna glede na $\|\cdot\|_\infty$ normo. Za drugo normo vzemimo omejeno zaporedje $(f_n)_n \subseteq \mathcal{C}([a, b])$. Od prej vemo, da je $(Kf_n)_n$ enakozvezna, podobno pokažemo tudi, da je omejena. \square

Vprašanje 63. Pokaži, da je integralski operator z zveznih jedrom kompakten na $\mathcal{C}([a, b])$ za drugo ali neskončno normo.

4.5.4 Kompaktnost adjungiranega operatorja

Izrek 4.5.44. Naj bo $T \in B(H, K)$. Naslednje trditve so ekvivalentne:

- T je kompakten,
- T^* je kompakten,
- obstaja zaporedje $(T_n)_n \subseteq F(H, K)$, da $T_n \rightarrow T$.

Dokaz. 3 v 1 je jasno. 1 v 3: Množica $\overline{T(B_H)}$ je kompaktna v K , torej je separabilna in velja

$$\overline{\operatorname{im} T} \subseteq \bigcup_{n \in \mathbb{N}} \overline{nT(B_H)}.$$

Torej je tudi $\overline{\operatorname{im} T}$ separabilen Hilbertov prostor, in ima števno KONS $(e_n)_n$. Naj bo P_n ortogonalni projektor na linearno ogrinjačo prvih n vektorjev e_i . Projektor je končnega ranga in zato kompakten. Definiramo $T_n = P_n T$.

Če je $y \in \overline{\operatorname{im} T}$, potem je $y = \sum_m \langle y, e_m \rangle e_m$. Velja $P_n y \xrightarrow{n \rightarrow \infty} y$, torej za $x \in H$ velja $Tx \in \overline{\operatorname{im} T}$, torej $T_n x \xrightarrow{n \rightarrow \infty} Tx$. Torej $T_n \rightarrow T$ po točkah.

Pokazati moramo še, da $T_n \rightarrow T$ v $B(H, K)$. Ker je T kompakten, za vsak $\varepsilon > 0$ obstajajo $x_1, \dots, x_m \in B_H$, da je

$$T(B_H) \subseteq \bigcup_{j=1}^m \mathring{B}(Tx_j, \frac{\varepsilon}{3}).$$

Za $x \in B_H$ obstaja x_j , da je $\|Tx - Tx_j\| < \varepsilon/3$. Potem je

$$\|Tx - T_n x\| \leq \|Tx - Tx_j\| + \|Tx_j - T_n x_j\| + \|T_n x_j - T_n x\| < \varepsilon,$$

saj

$$\|T_n x_j - T_n x\| \leq \|P_n\| \|Tx_j - Tx\| < \frac{\varepsilon}{3}.$$

1 v 2: Najprej izračunamo

$$\|T^* x\|^2 = \langle T^* x, T^* x \rangle = \langle TT^* x, x \rangle \leq \|TT^* x\| \|x\|.$$

Ker je T kompakten, je TT^* kompakten. Naj bo $(x_n)_n$ omejeno v K , torej obstaja konvergentno podzaporedje $(x_{n_k})_k$. Velja

$$\|T^* x_{n_k} - T^* x_{n_j}\|^2 \leq \|TT^*(x_{n_k} - x_{n_j})\| \|x_{n_k} - x_{n_j}\|.$$

Prvi člen konvergiira k 0 za $k, j \rightarrow \infty$, drugi člen pa je omejen. Torej je zaporedje $(T^* x_{n_k})_k$ Cauchyjevo v H , ki je poln, zato konvergiira, in je T^* kompakten.

2 v 1: Velja $T^{**} = T$. □

Vprašanje 64. Pokaži: omejen operator T je kompakten natanko tedaj, ko je T^* kompakten, kar je natanko tedaj, ko obstaja zaporedje $(T_n)_n$ operatorjev končnega ranga, ki konvergiira k T .

4.6 Spektralna teorija

Naj bo A kompleksna Banachova algebra z enoto. Naj bo $a \in A$. Definiramo

$$\rho(a) = \{\lambda \in \mathbb{C} \mid \lambda - a \text{ obrnljiv v } A\}.$$

Opomba. Identificiramo $\lambda - a = \lambda \cdot 1 - a$.

Pravimo, da je $\rho(a)$ RESOLVENTA elementa a . Označimo

$$R(\lambda, a) = (\lambda - a)^{-1}.$$

Preslikava $\lambda \mapsto R(\lambda, a)$ je RESOLVENTNA FUNKCIJA. Množici $\sigma(a) = \mathbb{C} \setminus \rho(a)$ pravimo SPEKTER elementa a .

Trditev 4.6.1. Naj bo $A \in B(H)$. Tedaj je $\sigma(A^*) = \{\bar{\lambda} \mid \lambda \in \sigma(A)\}$.

Dokaz. Naj bo B tak, da je $(\lambda I - A)B = B(\lambda I - A) = I$. Če obe strani adjungiramo, dobimo $B^*(\bar{\lambda}I - A^*) = (\bar{\lambda}I - A^*)B^* = I$. \square

Definicija 4.6.2. Naj bo X kompleksen Banachov prostor in A omejen operator na X . Potem je λ v ZVEZNEM delu spektra ($\lambda \in \sigma_c(A)$), če je $\lambda I - A$ injektiven in $\overline{\text{im}(\lambda I - A)} = X$ ter $\text{im}(\lambda I - A) \neq X$. Pravimo, da je λ v RESIDUALNEM delu spektra ($\lambda \in \sigma_r(A)$), če je $\lambda I - A$ injektiven in $\text{im}(\lambda I - A)$ ni gosta. Pravimo, da je λ v TOČKASTEM delu spektra ($\lambda \in \sigma_p(A)$), če je λ lastna vrednost za A .

Vprašanje 65. Kaj je resolventa in kaj je spekter? Na katere tri dele se razdeli spekter omejenega operatorja na kompleksnem Banachovem prostoru?

Trditev 4.6.3. Naj bo $A \in B(H)$. Če je $A = A^*$, je $\sigma_p(A) \subseteq \mathbb{R}$. Če je A normalen, so lastni vektorji medseboj pravokotni.

Trditev 4.6.4. Naj bo H Hilbertov prostor in $A \in B(H)$. Če je $\lambda \in \sigma_r(A)$, je $\bar{\lambda} \in \sigma_p(A^*)$. Če je $\lambda \in \sigma_p(A)$, je $\bar{\lambda} \in \sigma_p(A^*) \cup \sigma_r(A^*)$.

Dokaz. Če $\text{im}(A - \lambda I)$ ni gost v H , potem je

$$\ker(A^* - \bar{\lambda}I) = \ker(A - \lambda I)^* = \text{im}(A - \lambda I)^\perp \neq \{0\},$$

torej obstaja lastni vektor za $\bar{\lambda}$ za A^* .

Za drugo trditev velja $\ker(\lambda I - A) \neq \{0\}$, če vzamemo pravokotni komplement obeh strani, dobimo $\overline{\text{im}(\bar{\lambda}I - A^*)} \neq H$. \square

Vprašanje 66. Kako sta povezana točkast in residualni spekter operatorja s spektrom adjungiranega operatorja? Dokaži.

Posledica 4.6.5. Če je $A \in B(H)$ normalen, je $\sigma_r(A) = \emptyset$.

4 Uvod v funkcionalno analizo

Dokaz. Če je $\lambda \in \sigma_r(A)$, potem je $\bar{\lambda} \in \sigma_p(A^*)$ in zato

$$\ker(A - \lambda I) = \ker(A - \lambda I)^* = \ker(A^* - \bar{\lambda} I) \neq \{0\},$$

kjer smo v prvem koraku uporabili normalnost. Sledi $\lambda \in \sigma_p(A)$. \dashv □

Vprašanje 67. Pokaži: normalen operator ima prazen residualni spekter.

Lema 4.6.6. *Naj bo X kompleksen Banachov prostor in $A \in B(X)$. Naj bo $\lambda \in \sigma_c(A)$. Tedaj obstaja $(x_n)_n$ z $\|x_n\| = 1$, da $Ax_n - \lambda x_n \rightarrow 0$.*

Izrek 4.6.7. *Naj bo $A \in B(H)$ sebi adjungiran operator. Tedaj $\sigma(A) \subseteq \mathbb{R}$.*

Dokaz. Ker je A sebi adjungiran, je normalen, zato $\sigma_r(A) = \emptyset$. Dokazati moramo torej le še $\sigma_c(A) \subseteq \mathbb{R}$. Če je $\lambda \in \sigma_c(A)$, obstaja zaporedje $(x_n)_n$ z $\|x_n\| = 1$ in $Ax_n - \lambda x_n \rightarrow 0$. Pišimo $\lambda = \alpha + i\beta$. Potem velja

$$|\langle (\lambda I - A)x_n, x_n \rangle| \leq \|x_n\| \|(\lambda I - A)x_n\| \xrightarrow{n \rightarrow \infty} 0,$$

skalarni produkt pa je enak

$$\langle (\lambda I - A)x_n, x_n \rangle = \langle \alpha x_n - Ax_n, x_n \rangle + i\beta \langle x_n, x_n \rangle = \langle \alpha x_n - Ax_n, x_n \rangle + i\beta,$$

torej $\beta = 0$. □

Vprašanje 68. Pokaži: spekter sebi adjungiranega operatorja je realen.

Izrek 4.6.8. *Naj bo A Banachova algebra z enico in naj bo $\|a\| < 1$ za nek $a \in A$. Tedaj je $1 - a$ obrnljiv in*

$$(1 - a)^{-1} = \sum_{n=0}^{\infty} a^n.$$

Dokaz. Označimo delne vsote z s_n . Če je $n > m$, velja

$$\|s_n - s_m\| = \|a^{m+1} + \dots + a^n\| \leq \|a\|^{m+1} + \dots + \|a\|^n \leq \|a\|^{m+1} \sum_{k=0}^{\infty} \|a\|^k = \frac{\|a\|^{m+1}}{1 - \|a\|},$$

kar konvergira k 0 za $n, m \rightarrow \infty$. Torej je zaporedje Cauchyjevo, in obstaja limita s . Velja $(1 - a)s_n = 1 - a^{n+1}$. Leva stran konvergira k $(1 - a)s$, desna pa k 1. □

Vprašanje 69. Pokaži: če je a element Banachove algebre in $\|a\| < 1$, je $1 - a$ obrnljiv.

Lema 4.6.9. *Naj bosta $\lambda, \mu \in \rho(a)$. Velja $R(\mu, a) - R(\lambda, a) = (\lambda - \mu)R(\lambda, a)R(\mu, a) = (\lambda - \mu)R(\mu, a)R(\lambda, a)$.*

Dokaz. Ker $(\lambda - a)$ in $(\mu - a)$ komutirata, komutirata tudi njuna inverza. Preostalo je preprost račun. □

Trditev 4.6.10. Naj bo $a \in A$, kjer je A kompleksna Banachova algebra z enico. Potem je $\rho(a)$ odprta v \mathbb{C} , $\sigma(a)$ pa kompaktna in vsebovana v $B(0, \|a\|)$.

Dokaz. Naj bo $\lambda_0 \in \rho(a)$. Iščemo $\varepsilon > 0$, da iz $|\lambda - \lambda_0| < \varepsilon$ sledi $\lambda \in \rho(a)$. Izrazimo lahko

$$\lambda - a = (\lambda_0 - a)(1 + (\lambda - \lambda_0)(\lambda_0 - a)^{-1}).$$

Če je $\|(\lambda - \lambda_0)(\lambda_0 - a)^{-1}\| \leq \varepsilon \|(\lambda_0 - a)^{-1}\| < 1$, potem je

$$1 + (\lambda - \lambda_0)(\lambda_0 - a)^{-1}$$

obrnljiv, torej $\lambda - a$ produkt dveh obrnljivih elementov.

Za drugo trditev je $\sigma(a) = \mathbb{C} \setminus \rho(a)$ zaprta množica. Če je $|\lambda| > \|a\|$, je

$$\lambda - a = \lambda(1 - \frac{a}{\lambda}),$$

ampak potem $\|a/\lambda\| < 1$, in je $\lambda \in \rho(a)$. Torej je $\sigma(a) \subseteq B(0, \|a\|)$. \square

Vprašanje 70. Pokaži: če je a element Banachove algebre, je $\sigma(a)$ kompaktno, vsebovan v $B(0, \|a\|)$.

Opomba. Če je $|\lambda| > \|a\|$, je $\lambda \in \rho(a)$ in

$$(\lambda - a)^{-1} = \sum_{n=0}^{\infty} \frac{a^n}{\lambda^{n+1}}.$$

Opomba. Če je $\lambda_0 \in \rho(a)$ in $|\lambda - \lambda_0| < \|(\lambda_0 - a)^{-1}\|^{-1}$, potem je $\lambda \in \rho(a)$ in velja

$$(\lambda - a)^{-1} = (\lambda_0 - a)^{-1} \sum_{n=0}^{\infty} (\lambda_0 - \lambda)^n (\lambda_0 - a)^{-n}.$$

Trditev 4.6.11. Naj bo $\mu \in \rho(a)$. Tedaj velja

$$\lim_{\lambda \rightarrow \mu} \frac{(\lambda - a)^{-1} - (\mu - a)^{-1}}{\lambda - \mu} = -(\mu - a)^{-2}.$$

Trditev 4.6.12. Resolventna funkcija je zvezna.

Dokaz. Naj bo $\lambda_0 \in \rho(a)$ in $|\lambda - \lambda_0| < \|R(\lambda_0, a)\|^{-1}$. Potem velja

$$R(\lambda, a) = \sum_{n=0}^{\infty} (\lambda - \lambda_0)^n R(\lambda_0, a)^{n+1},$$

torej je

$$R(\lambda, a) - R(\lambda_0, a) = \sum_{n=1}^{\infty} (\lambda_0 - \lambda)^n R(\lambda_0, a)^{n+1}.$$

Iz tega sledi

$$\begin{aligned} \|R(\lambda, a) - R(\lambda_0, a)\| &\leq |\lambda - \lambda_0| \|R(\lambda_0, a)\|^2 \sum_{n=0}^{\infty} |\lambda - \lambda_0|^n \|R(\lambda_0, a)\|^n \\ &= \frac{|\lambda_0 - \lambda| \|R(\lambda_0, a)\|^2}{1 - |\lambda_0 - \lambda| \|R(\lambda_0, a)\|} \\ &\xrightarrow{\lambda \rightarrow \lambda_0} 0. \end{aligned}$$

□

Vprašanje 71. Pokaži: resolventna funkcija je zvezna.

Izrek 4.6.13. Naj bo $a \in A$ in A kompleksna Banachova algebra z enico. Tedaj je $\sigma(a) \neq \emptyset$.

Dokaz. Recimo $\sigma(a) = \emptyset$. Potem je $\rho(a) = \mathbb{C}$, ker pa velja

$$\lim_{\lambda \rightarrow \mu} \frac{R(\lambda, a) - R(\mu, a)}{\lambda - \mu} = -R(\mu, a)^2,$$

imamo za vsak $\varphi \in A^*$

$$\varphi \left(\lim_{\lambda \rightarrow \mu} \frac{R(\lambda, a) - R(\mu, a)}{\lambda - \mu} \right) = \lim_{\lambda \rightarrow \mu} \frac{\varphi(R(\lambda, a)) - \varphi(R(\mu, a))}{\lambda - \mu} = -\varphi(R(\mu, a)^2),$$

torej je $\varphi \circ R(\cdot, a)$ cela holomorfná funkcija.

Če je $|\lambda| > \|a\|$, je

$$\|R(\lambda, a)\| \leq \sum_{n=0}^{\infty} \frac{\|a\|^n}{|\lambda|^{n+1}} = \frac{1}{|\lambda|} \frac{1}{1 - \|a\|/|\lambda|},$$

to pa konvergira k 0 za $|\lambda| \rightarrow \infty$. Torej je $R(\cdot, a)$ omejena, zato je tudi $\varphi \circ R(\cdot, a)$ omejena. Ker je cela holomorfná, je konstantna in velja $\varphi(R(\lambda, a)) = 0$ za vsak $\lambda \in \mathbb{C}$. Za $\lambda \in \rho(a)$ velja $(\lambda - a)R(\lambda, a) = 1$, torej $R(\lambda, a) \neq 0$, in smo prišli v protislovje s Hahn-Banachom. □

Vprašanje 72. Dokaži: spekter elementa Banachove algebre z enico je vedno neprazen.

Definicija 4.6.14. Naj bo $a \in A$. Definirajmo

$$r(a) = \sup_{\lambda \in \sigma(a)} |\lambda| = \max_{\lambda \in \sigma(a)} |\lambda|.$$

To število imenujemo **SPEKTRALNI RADIJ** elementa a .

Lema 4.6.15. če je $\lambda \in \sigma(a)$, je $\lambda^n \in \sigma(a^n)$.

Dokaz. Dokažemo obratno; če je $\lambda^n \in \rho(a^n)$, je $\lambda \in \rho(a)$. Računamo

$$\lambda^n - a^n = (\lambda - a)(\lambda^{n-1} + \dots + a^{n-1})$$

in desni člen produkta označimo z b . Obstaja $c \in A$, da je $(\lambda^n - a^n)c = c(\lambda^n - a^n) = 1$, torej $(\lambda - a)bc = 1$ in ima $\lambda - a$ desni inverz. Podobno pokažemo, da ima levi inverz. \square

Trditev 4.6.16. Naj bo $a \in A$ in p polinom. Tedaj je $\sigma(p(a)) = p(\sigma(a))$.

Izrek 4.6.17 (Gelfandova formula). Naj bo $a \in A$. Potem je

$$r(a) = \lim_{n \rightarrow \infty} \|a^n\|^{1/n} = \liminf_{n \rightarrow \infty} \|a^n\|^{1/n} = \inf_{n \in \mathbb{N}} \|a^n\|^{1/n}.$$

Dokaz. Naj bo $\lambda \in \sigma(a)$, da je $|\lambda| = r(a)$. Potem je $\lambda^n \in \sigma(a^n)$, torej $|\lambda|^n \leq \|a^n\|$ in velja $r(a) = |\lambda| \leq \|a^n\|^{1/n}$ za vsak $n \in \mathbb{N}$. Zaporedje je navzdol omejeno, torej ima infimum, ki je

$$r(a) \leq \inf_{n \in \mathbb{N}} \|a^n\|^{1/n} \leq \liminf_{n \rightarrow \infty} \|a^n\|^{1/n}.$$

Če pokažemo, da je $\limsup \|a^n\|^{1/n} \leq r(a)$, potem bo dokaz končan.

Naj bo $f \in A^*$. Definiramo $\tilde{f}(\lambda) = f(R(\lambda, a))$ kot preslikavo $\rho(a) \rightarrow \mathbb{C}$. To je holomorfná funkcija. Če je $|\lambda| > \|a\|$, lahko $R(\lambda, a)$ eksplicitno izrazimo in zato velja

$$\tilde{f}(\lambda) = \sum_{n=0}^{\infty} \frac{f(a^n)}{\lambda^{n+1}}.$$

To je Laurantov razvoj \tilde{f} v okolici 0 na komplement krogle $B(0, \|a\|)$. Razvoj lahko naredimo tudi na $B(0, r(a))^c$. Ker je enoličen, zgornji predpis velja za vsak $\lambda > r(a)$. Za $\varepsilon > 0$ in $\lambda = r(a) + \varepsilon$ torej konvergira vrsta

$$\sum_{n=0}^{\infty} \frac{f(a^n)}{(r(a) + \varepsilon)^{n+1}},$$

torej obstaja $M > 0$, da je $|f(a^n)/(r(a) + \varepsilon)^{n+1}| \leq M$ za vse $n \in \mathbb{N}$. Po principu šibke omejenosti je potem množica

$$\left\{ \frac{a^n}{(r(a) + \varepsilon)^{n+1}} \mid n \in \mathbb{N} \right\}$$

omejena in obstaja $\tilde{M} > 0$, da

$$\left\| \frac{a^n}{(r(a) + \varepsilon)^{n+1}} \right\| \leq \tilde{M}$$

za vse $n \in \mathbb{N}$. Sledi

$$\limsup_{n \rightarrow \infty} \|a^n\|^{1/n} \leq \lim_{n \rightarrow \infty} (\tilde{M}(r(a) + \varepsilon))^{1/n} (r(a) + \varepsilon) = r(a) + \varepsilon,$$

saj prvi člen produkta konvergira k 1. Torej je $\limsup \|a^n\|^{1/n} \leq r(a)$. \square

Vprašanje 73. Povej in dokaži Gelfandovo formulo.

Posledica 4.6.18. Naj bo A sebi adjungiran operator na Hilbertovem prostoru. Tedaj velja $r(A) = \|A\|$.

Dokaz. Po izreku velja $r(A) = \lim \|A^n\|^{1/n}$. Ker je $\|A^2\| = \|A^*A\| = \|A\|^2$, lahko z indukcijo pokažemo, da velja $\|A^{2^n}\| = \|A\|^{2^n}$. S tem dobimo podzaporedje zaporedja v limiti, ki bo konvergiral k $\|A\|$. \square

Vprašanje 74. Pokaži, da za sebi adjungiran operator velja $\|A\| = r(A)$.

4.6.1 Spekter kompaktnega operatorja

Primer. Naj bo $H = l^2$ in $(e_n)_n$ neka ortonormirana baza. Naj bo $(d_n)_n$ zaporedje omejenih števil. Definiramo diagonalni operator $D : l^2 \rightarrow l^2$ z

$$Dx = \sum_{n=0}^{\infty} d_n \langle x, e_n \rangle e_n.$$

Preverimo lahko naslednja dejstva:

- $\|D\| = \sup_n |d_n|$,
- D je sebi adjungiran natanko tedaj, ko je $d_n \in \mathbb{R}$ za vsak n ,
- D je normalen,
- D je unitaren natanko tedaj, ko je $|d_n| = 1$ za vsak n ,
- D je kompakten natanko tedaj, ko je $d_n \rightarrow 0$.

Lema 4.6.19 (Rieszova lema o pravokotnici). Naj bo X normiran prostor in Y zaprt podprostor v X . Tedaj za vsak $\varepsilon \in (0, 1)$ obstaja $x \in X$, da velja $\|x\| = 1$ in $d(x, Y) \geq \varepsilon$.

Vprašanje 75. Povej Rieszovo lemo o pravokotnici.

Opomba. Če je X končnorazsežen, obstaja $x \in X$ z $\|x\| = 1$ in $d(x, Y) = 1$. Skličemo se lahko na kompaktnost.

Trditev 4.6.20. Naj bo Y končnorazsežen podprostor v normiranem prostoru X . Tedaj obstaja zaprt podprostor $Z \subseteq X$, da je $X = Y \oplus Z$.

Dokaz. Naj bo $\{e_1, \dots, e_n\}$ baza za Y in $\{f_1, \dots, f_n\}$ dualna baza. Ker je dimenzija Y končna, je f_i omejen za vse $i = 1, \dots, n$. Naj bo F_i katerakoli Hahn-Banachova razširitev funkcije f_i . Definiramo

$$Z = \bigcap_{n=1}^{\infty} \ker F_i.$$

Ker je F_i omejen, je $\ker F_i$ zaprta množica, torej je presek zaprt.

Če je $x \in X$, lahko izrazimo

$$x = \underbrace{F_1(x)e_1 + \dots + F_n(x)e_n}_{\in Y} + \underbrace{x - F_1(x)e_1 - \dots - F_n(x)e_n}_{\in Z}.$$

Če je $x \in Y \cap Z$, velja $x = \alpha_1 e_1 + \dots + \alpha_n e_n$, ampak tudi $F_i(x) = 0 = \alpha_i$ za vse i , torej $x = 0$. \square

Vprašanje 76. Dokaži: za končnorazsežen podprostor $Y \leq X$ obstaja zaprt podprostor $Z \leq X$, da je $X = Y \oplus Z$.

Trditev 4.6.21. Naj bo X Banachov prostor in $K \in K(X)$. Če je $\lambda \neq 0$, je $\dim \ker(K - \lambda I) < \infty$. Dodatno je $\text{im}(K - \lambda I)$ zaprta v X .

Dokaz. Izrazimo lahko $K - \lambda I = \lambda(\frac{1}{\lambda}K - I)$, torej je $\ker(K - \lambda I) = \ker(\frac{1}{\lambda}K - I)$. Označimo drugo preslikavo z S . Velja $x \in \ker(S - I) \Leftrightarrow Sx = x$, torej je $S|_{\ker(S-I)}$ identiteta, in hkrati kompakten operator. Torej je jedro končnorazsežno.

Za drugo trditev zapišemo $X = \ker(K - \lambda I) \oplus Y$, kjer je Y zaprt podprostor X . Velja

$$\text{im}(K - \lambda I) = (K - \lambda I)(\ker(K - \lambda I) \oplus Y) = (K - \lambda I)Y = \text{im}(K - \lambda I|_Y).$$

Označimo $T = K - \lambda I$.

Dovolj je dokazati, da je $T|_Y$ navzdol omejen, saj bo potem topološki izomorfizem $Y \rightarrow \text{im } T$. Ker je Y zaprt, je Banachov, zato bo $\text{im } T$ Banachov, polni prostori pa so vedno zaprti.

Recimo, da $T|_Y$ ni navzdol omejen. Tedaj za vsak $n \in \mathbb{N}$ obstaja $y_n \in Y$, da $\|y_n\| = 1$ in $\|Ty_n\| \leq 1/n$. Ker je K kompakten, obstaja podzaporedje $(y_{n_k})_k$, da $(Ky_{n_k})_k$ konvergira. Velja

$$y_{n_k} = \lambda \frac{1}{\lambda} y_{n_k} = \frac{1}{\lambda} (\lambda y_{n_k} - Ky_{n_k} + Ky_{n_k}) \rightarrow y$$

za nek $y \in Y$ z $\|y\| = 1$ saj sta prva dva člena vsote del zaporedja Ty_n , ki konvergira. Enostavno lahko pokažemo, da velja $Ky = \lambda y$. To pomeni, da je $y \in \ker(K - \lambda I) \cap Y = \{0\}$, ampak $\|y\| = 1$. \nrightarrow \square

Vprašanje 77. Pokaži: če je K kompakten operator in $\lambda \neq 0$, je jedro $K - \lambda I$ končnorazsežno, slika $K - \lambda I$ pa je zaprta.

Lema 4.6.22. *Naj bo K kompakten operator na kompleksnem Banachovem prostoru X . Tedaj ima K za vsak $\varepsilon > 0$ le končno mnogo linearno neodvisnih lastnih vektorjev za lastne vrednosti λ z $|\lambda| \geq \varepsilon$.*

Dokaz. Recimo, da imamo $\varepsilon > 0$ in zaporedje $(x_n)_n$ linearno neodvisnih lastnih vektorjev, ki zaporedoma pripadajo lastnim vrednostim λ_n z $|\lambda_n| \geq \varepsilon$. Definirajmo X_n kot linearno ogrinjačo prvih n lastnih vektorjev. Velja $\dim X_n < \infty$, torej so ti prostori zaprti, hkrati pa velja

$$\{0\} \subsetneq X_1 \subsetneq X_2 \subsetneq X_3 \subsetneq \dots$$

Po Rieszovi lemi o pravokotnici obstaja $y_n \in X_n \setminus X_{n-1}$ z $\|y_n\| = 1$, da velja $d(y_n, X_{n-1}) \geq \frac{1}{2}$ za vse $n \geq 2$. Pišimo $y_n = \alpha_1 x_1 + \dots + \alpha_n x_n$ in $z_n = y_n / \lambda_n$. Potem velja $\|z_n\| = |\lambda_n|^{-1} \leq \frac{1}{\varepsilon}$ ter

$$Kz_n - y_n = \alpha_1 \left(\frac{\lambda_1}{\lambda_n} - 1 \right) x_1 + \dots + \alpha_{n-1} \left(\frac{\lambda_{n-1}}{\lambda_n} - 1 \right) x_{n-1} \in X_{n-1}.$$

Za $m < n$ izračunamo

$$\|Kz_n - Kz_m\| = \left\| \underbrace{y_n}_{\in X_n} - \underbrace{(y_n - Kz_n + Kz_m)}_{\in X_{n-1}} \right\| \geq \frac{1}{2}$$

torej $(Kz_n)_n$ nima Cauchyjevega podzaporedja, zato nima stekališča. To je protislovje s kompaktnostjo K . \square

Vprašanje 78. Pokaži, da ima kompakten operator le končno mnogo linearno neodvisnih lastnih vektorjev, ki pripadajo lastnim vrednostim z $|\lambda| > \varepsilon$.

Izrek 4.6.23. *Za kompakten operator K na kompleksnem Banachovem prostoru X veljajo naslednje trditve:*

- če je $\dim X = \infty$, je $0 \in \sigma(K)$,
- če je $\lambda \in \sigma(K) \setminus \{0\}$, je λ lastna vrednost K ,
- $\sigma(K)$ je končen ali števno neskončen,
- če je $\sigma(K)$ neskončen in so $(\lambda_n)_n$ lastne vrednosti, štete z geometrijsko večkratnostjo, potem $\lambda_n \rightarrow 0$.

Dokaz. Prva točka: Če $0 \notin \sigma(K)$, je K obrnljiv in je zato $\text{id} = KK^{-1}$ kompaktna, torej je $\dim X < \infty$.

Tretja točka: Računamo

$$\sigma(K) \setminus \{0\} = \bigcup_{n \in \mathbb{N}} \sigma(K) \cap \left\{ \lambda \in \mathbb{C} \mid |\lambda| \geq \frac{1}{n} \right\}.$$

To je števna unija končnih množic.

Četrta točka: Iz leme dobimo, da za vsak $\varepsilon > 0$ obstaja n_ε , za katerega za poljuben $n \geq n_\varepsilon$ velja $|\lambda_n| < \varepsilon$. Torej $\lambda_n \rightarrow 0$.

Druga točka: Recimo, da je $0 \neq \lambda \in \sigma(K)$. Recimo, da je $K - \lambda I$ injektiven. Če dokažemo, da je $K - \lambda I$ surjektiven, bo sledilo, da je obrnljiv z omejenim inverzom, oziroma $\lambda \in \rho(K)$.

Pokažimo prvo, da je $\text{im}(K - \lambda I)^n$ zaprta za vsak n . Trditev pove, da ima $K - \lambda I$ zaprto sliko. Izrazimo lahko

$$(K - \lambda I)^n = \sum_{k=0}^{\infty} \binom{n}{k} (-\lambda)^k K^{n-k} =: \tilde{K} + (-\lambda)^n I.$$

Operator \tilde{K} je kompakten, torej ima $(K - \lambda I)^n$ zaprto sliko.

Definirajmo $X_n = \ker(K - \lambda I)^n$ za $n \in \mathbb{N}_0$. Velja $X_0 = \{0\} \subseteq X_1 \subseteq X_2 \subseteq \dots$. Recimo, da so vse inkluzije stroge. Po Rieszovi lemi o pravokotnici za vsak $n \in \mathbb{N}$ obstaja $x_n \in X_n \setminus X_{n-1}$ z $\|x_n\| = 1$ in $d(x_n, X_{n-1}) \geq \frac{1}{2}$. Za $m < n$ je

$$\|Kx_m - Kx_n\| = \|\lambda x_n + Kx_n - \lambda x_n - Kx_m\| = |\lambda| \left\| x_n + \frac{1}{\lambda}(K - \lambda I)x_n - \frac{1}{\lambda}Kx_m \right\|.$$

Velja $x_m \in X_m$, torej $Kx_m \in X_m$ (saj K in $K - \lambda I$ komutirata), iz česar sledi $Kx_m \in X_{n-1}$, saj je $X_m \subseteq X_{n-1}$. Hkrati je $(K - \lambda I)x_n \in X_{n-1}$, torej $\frac{1}{\lambda}((K - \lambda I)x_n - Kx_m) \in X_{n-1}$. Velja torej $\|Kx_n - Kx_m\| \geq \frac{1}{2}|\lambda|$. Kot prej pridemo v protislovje s kompaktnostjo operatorja K . Torej mora obstajati n , za katerega je $X_n = X_{n+1}$.

Za konec recimo, da $K - \lambda I$ ni surjektiven. Potem $X \neq (K - \lambda I)X$, velja pa $(K - \lambda I)^n X = (K - \lambda I)^{n+1} X$. To je protislovje z injektivnostjo $(K - \lambda I)^n$. \square

Vprašanje 79. Kaj lahko poveš o spektru kompaktnega operatorja na kompleksnem Banachovem prostoru? Dokaži.

Izrek 4.6.24. *Naj bo $K : H \rightarrow H$ kompakten sebi adjungiran operator na kompleksnem Hilbertovem prostoru H . Potem obstajata zaporedje $(\lambda_n)_n \subseteq \mathbb{R}$ in ortonormiran sistem $(e_n)_n$ (lahko sta oba končna) z*

- $|\lambda_1| \geq |\lambda_2| \geq \dots$, $\lambda_n \neq 0$, in če je zaporedje neskončno, velja $\lambda_n \rightarrow 0$,
- $Ke_n = \lambda_n e_n$,
- če je $\lambda \in \sigma_p(K) \setminus \{0\}$, se λ pojavi v $(\lambda_n)_n$ natanko tolikokrat, kot je $\dim \ker(K - \lambda I)$,
- $Kx = \sum \lambda_n \langle x, e_n \rangle e_n$.

Dokaz. Ker je K sebi adjungiran, velja $r(K) = \|K\|$ in $\sigma(K) \subseteq [-\|K\|, \|K\|]$. Če je $K = 0$, nimamo nič za dokazati. Sicer lahko razvrstimo lastne vrednosti K v padajoče zaporedje po absolutni vrednosti, pri čemer jih štejemo glede na geometrijske večkratnosti. Če je $\lambda \in \sigma_p(K) \setminus \{0\}$, lahko izberemo ortonormirano bazo za $\ker(K - \lambda I)$. Če je

$\mu \neq \lambda$ tudi neničelna lastna vrednost, bo njena ONB pravokotna na ONB za λ . Postopek ponovimo za vsako lastno vrednost v zaporedju. Dobljene vektorje zložimo v $(e_n)_n$, da je $Ke_n = \lambda_n e_n$. Definiramo Y kot zaprtje linearne ogrinjače teh vektorjev. Potem je $(e_n)_n$ ONB za Y in $H = Y \oplus Y^\perp$.

Ker je $Ke_n = \lambda_n e_n$, je linearna ogrinjača $(e_n)_n$ invariantna na K , ki je zvezen, torej enako velja za Y . Iz sebi adjungiranosti sledi $K(Y^\perp) \subseteq Y^\perp$. Če je $K|_{Y^\perp} \neq 0$, je njegova norma neničelna lastna vrednost za K , kar je nemogoče, ker smo že našli vse lastne vrednosti. Torej za $x \in H$ velja $Kx = \sum \lambda_n \langle x, e_n \rangle e_n$. \square

Vprašanje 80. Kaj lahko poveš o spektru kompaktnega sebi adjungiranega operatorja na kompleksnem Hilbertovem prostoru? Dokaži.

5 Statistika 2

5.1 Ocenjevanje v linearnih modelih

Splošni linearni model je oblike $X = Z\beta + \varepsilon$, kjer je $Z \in \mathbb{R}^{n \times d}$ znana konstantna matrika, $\beta \in \mathbb{R}^d$ neznani parameter, ε pa je neopazljiv slučajni šum. Privzamemo, da velja $E(\varepsilon) = 0$. V splošnem za varianco ε ne privzamemo ničesar, v standardnih linearnih regresijskih modelih pa privzamemo, da je diagonalna.

Privzemimo splošni linearni model in naj bo B vektorski podprostor v \mathbb{R}^d . Naj bo $x \in \mathbb{R}^n$ realizacija slučajnega vektorja X . RESTRINGIRANA OCENA za $\beta \in B$ na podlagi x po metodi najmanjših kvadratov je tak vektor $\hat{\beta}_B$, za katerega je

$$\|x - Z\hat{\beta}_B\|^2 = \min_{b \in B} \|x - Zb\|^2.$$

Pišimo $\hat{\beta} = \hat{\beta}_B$. Vemo, da je $Z\hat{\beta}$ ravno pravokotna projekcija vektorja x na podprostor $ZB \subseteq \mathbb{R}^n$. Določena je z zahtevo $x - Z\hat{\beta} \perp ZB$. Za $B = \mathbb{R}^d$ to velja natanko v primeru $Z^T(X - Z\hat{\beta}) = 0$, torej $Z^T Z\hat{\beta} = Z^T x$. V primeru, ko je Z polnega ranga, je $Z^T Z$ obrnljiva in $\hat{\beta} = (Z^T Z)^{-1} Z^T x$. Če pa je jedro Z netrivialno, imamo rešitev več.

Če na stolpcih Z izvedemo prirejeno Gram-Schmidtovo ortogonalizacijo, lahko kljub temu poiščemo rešitev. Označimo s S_i rezultat ortogonalizacije na i -tem stolpcu Z . V primeru, ko je ta stolpec v linearni ogrinjači prejšnjih, nastavimo $S_i = 0$. S tem dobimo ortogonalne vektorje S_1, \dots, S_d . Dobimo razcep $Z = SP$, kjer je S matrika iz zloženih stolpcev S_i , P pa zgornje trikotna matrika s pozitivnimi števili na diagonalni, in zato obrnljiva. Velja $Z^T Z = P^T J P$, kjer je J diagonalna matrika, na diagonalni katere so kvadrati norm stolpcev S_i . S tem lahko definiramo posplošen inverz $(Z^T Z)' = P^{-1} J P^{-T}$. Potem $\hat{\beta} = (Z^T Z)' Z^T x$ reši enačbo $Z^T Z\hat{\beta} = Z^T x$.

Opomba. Vsaki matriki M , ki ustreza $Z^T Z M Z^T = Z^T$ pravimo POSPLOŠEN INVERZ matrike $Z^T Z$.

Izračunajmo

$$E(\hat{\beta}(X)) = (Z^T Z)' Z^T E(X) = (Z^T Z)' Z^T Z\beta = P^T J P^{-T} \beta.$$

Če Z nima polnega ranga, potem $\hat{\beta}$ ni nepristranska cenilka za β . V tem primeru pravzaprav ne obstaja nepristranska linearna cenilka za β , namreč, če je $U : \mathbb{R}^n \rightarrow \mathbb{R}^d$ taka, mora veljati $E(UX) = \beta$, hkrati pa $E(UX) = UZ\beta$, iz česar sledi, da je Z injektivna, torej matrika polnega ranga.

Trditev 5.1.1. *Naj bo $e(\beta, \varepsilon) = L\beta$ ocenjevana funkcija, kjer je $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ linearna preslikava. Dalje naj bo $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$ nepristranska linearna cenilka za $L\beta$. Tedaj je $L = UZ$ in je $UZ\hat{\beta}$ nepristranska linearna cenilka za $L\beta$.*

Dokaz. Kot zgoraj je $L\beta = UZ\beta$ za vse β . Izračunajmo

$$Z\hat{\beta}(X) = Z(Z^T Z)' Z^T X = S J S^T X = S S^T X.$$

Sledi

$$E(UZ\hat{\beta}(X)) = U S S^T Z\beta = U S S^T S P\beta = U S P\beta = UZ\beta. \quad \square$$

Izrek 5.1.2 (Gauss-Markov). *Privzemimo linearni regresijski model $X = Z\beta + \varepsilon$, kjer je $\text{var}(\varepsilon) = \sigma^2 I$. Naj bo $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$ linearna preslikava. Tedaj ima $UZ\hat{\beta}$ med vsemi nepristranskimi linearnimi cenilkami za $UZ\beta$ enakomerno najmanjšo disperzijo.*

Dokaz. Naj bo $W : \mathbb{R}^n \rightarrow \mathbb{R}^m$ druga nepristranska linearna cenilka za $UZ\beta$. Po prejšnji trditvi je $WZ = UZ$. Primerjati želimo $\text{var}(WX)$ in $\text{var}(UZ\hat{\beta}) = \text{var}(WZ\hat{\beta})$ upoštevaje $UZ = WZ$.

Velja $\text{var}(WX) = W \text{var}(X) W^T = \sigma^2 W W^T$ in podobno kot v prejšnjem dokazu

$$\text{var}(WZ\hat{\beta}) = \text{var}(W S S^T X) = \sigma^2 W S S^T W^T.$$

Trdimo, da za poljuben $\xi \in \mathbb{R}^m$ velja

$$\langle W W^T \xi, \xi \rangle \geq \langle W S S^T W^T \xi, \xi \rangle.$$

Upoštevaje $\langle W W^T \xi, \xi \rangle = \langle W^T \xi, W^T \xi \rangle$ in $\langle W S S^T W^T \xi, \xi \rangle = \langle S^T W^T \xi, S^T W^T \xi \rangle$ je dovolj za poljuben w pokazati

$$\langle w, w \rangle \geq \langle S^T w, S^T w \rangle.$$

To velja, ker je $\|S_i\| \in \{0, 1\}$. □

5.1.1 Ocenjevanje v normalnem linearnem regresijskem modelu

Privzamemo $X = Z\beta + \varepsilon$ za $\varepsilon \sim N(0, \sigma^2 I)$. To je parametričen model, vendar β in σ^2 porazdelitve ne določata enolično, če Z nima polnega ranga.

Če dodatno zahtevamo $x - Z\hat{\beta}(x) \perp \text{im } Z$, pa je vsaj $Z\hat{\beta}(x)$ enolično določen. Izračunajmo

$$\|x - Z\beta\|^2 = \|x\|^2 - 2\langle x, Z\beta \rangle + \|Z\beta\|^2 = \|x - Z\hat{\beta} + Z\hat{\beta}\|^2 - 2\langle x, Z\beta \rangle + \|Z\beta\|^2.$$

Upoštevaje pravokotnost je to enako

$$\|x - Z\beta\|^2 = \|x - Z\hat{\beta}\|^2 + \|Z\hat{\beta}\|^2 - 2\langle x, Z\beta \rangle + \|Z\beta\|^2$$

oziroma

$$\|x - Z\beta\|^2 = \|x - Z\hat{\beta}\|^2 + \|Z(Z^T Z)^{-1} Z^T x\|^2 - 2\langle Z^T x, \beta \rangle + \|Z\beta\|^2.$$

Ker je gostota porazdelitve enaka

$$f_X(x) = (2\pi\sigma^2)^{-n/2} \exp\left(\frac{-1}{2\sigma^2} \|x - Z\beta\|^2\right),$$

torej $\|x - Z\hat{\beta}\|^2$ in $Z^T x$ tvorita zadostno statistiko za dano porazdelitev. Označimo $T(x) = \left(Z^T x, \|x - Z\hat{\beta}\|^2\right)$. Drugemu členu dvojice označimo z $\text{SSR}(x)$.

Posledica 5.1.3. Statistike, ki so od vzorca odvisne le preko T , so avtomatično nepristranske cenilke z enakomerno najmanjšo disperzijo za svojo pričakovano vrednost.

Izrek 5.1.4. Statistika $U(X) = (Z\hat{\beta}(X), \frac{1}{n-r} \text{SSR}(X))$, kjer je $r = \text{rang } Z$, je nepristranska cenilka za $(Z\beta, \sigma^2)$, ki ima med vsemi nepristranskimi cenilkami enakomerno najmanjšo disperzijo. Dalje sta $Z\hat{\beta}(X)$ in $\text{SSR}(X)$ neodvisni, ter $\text{SSR}(X)/\sigma^2 \sim \chi_{n-r}^2$.

Dokaz. Vemo že, da je $Z\hat{\beta}$ nepristranska cenilka z enakomerno najmanjšo disperzijo. Če je $Y = (Y_1, \dots, Y_n) \sim N(\nu, \sigma^2 I)$, potem je vsaka funkcija (Y_1, \dots, Y_m) neodvisna od vsake funkcije (Y_{m+1}, \dots, Y_n) . Če je še $\nu_1 = \dots = \nu_m = 0$, je

$$\frac{1}{\sigma^2} \sum_{i=1}^m Y_i^2 \sim \chi_m^2.$$

Konstruirajmo ortogonalno matriko $\tilde{S} \in O(n)$ na sledeči način. Postavimo $\tilde{S}_i = S_i$ za tiste i , za katere je $\|S_i\| = 1$, za ostale stolpce pa te vrednosti dopolnimo do ortonormirane baze. Velja

$$\text{SSR}(X) = \|X - Z\hat{\beta}(X)\|^2 = \|\tilde{S}^T X - \tilde{S}^T S S^T X\|^2 = \|\tilde{S}^T X - \tilde{S}^T S S^T \tilde{S} \tilde{S}^T X\|$$

ker je \tilde{S} ortogonalna. Izračunamo

$$\text{SSR}(X) = \left\| \left(I - \begin{bmatrix} J \\ 0 \end{bmatrix} \begin{bmatrix} J & 0 \end{bmatrix} \right) \tilde{S}^T X \right\|^2 = \left\| \begin{bmatrix} I - J & \\ & I \end{bmatrix} \tilde{S}^T X \right\|^2,$$

torej je $\text{SSR}(X)/\sigma^2 \sim \chi_{n-r}^2$. Takoj se prepričamo, da je $Z\hat{\beta}(X) = S S^T X$ odvisen od preostalih r komponent vektorja $\tilde{S}^T X$, torej sta $Z\hat{\beta}(X)$ in $\text{SSR}(X)$ neodvisna slučajna vektorja. \square

5.2 Ocenjevanje za velike vzorce

Naj bodo X, X_1, X_2, \dots slučajni vektorji, definirani na nekem skupnem verjetnostnem prostoru.

- $(X_n)_n$ konvergira k X SKORAJ GOTOVO, če je $P(\lim X_n = X) = 1$.
- $(X_n)_n$ konvergira k X v VERJETNOSTI, če za vsak $\varepsilon > 0$ velja $\lim P(\|X_n - X\| > \varepsilon) = 0$.
- $(X_n)_n$ konvergira k X v L^2 , če je $\lim E(\|X_n - X\|_2^2) = 0$ ($\|\cdot\|_2$ je funkcijska norma).
- $(X_n)_n$ konvergira k X v PORAZDELITVI, če velja $\lim F_{X_n}(x) = F_X(x)$ za vsako točko x , v kateri je F_X zvezna.

Lema 5.2.1. Naj bodo X, X_1, X_2, \dots slučajni r -vektorji. Potem velja naslednje.

- X_n konvergira k X v verjetnosti natanko tedaj, ko ima vsako podzaporedje zaporedja $(X_n)_n$ nadaljnje podzaporedje, ki konvergira skoraj gotovo.
- X_n konvergira k X v porazdelitvi natanko tedaj, ko za vsak $\xi \in \mathbb{R}^r$ velja $\langle X_n, \xi \rangle \xrightarrow{d} \langle X, \xi \rangle$.

Trditev 5.2.2. Z enakimi oznakami, če X_n konvergira k X skoraj gotovo ali v L^2 smislu, potem konvergira tudi v verjetnosti. Če konvergira v verjetnosti, potem konvergira tudi v porazdelitvi. Če je X konstanten vektor, potem sta konvergenca v porazdelitvi in v verjetnosti ekvivalentni.

Trditev 5.2.3. Naj bodo X, X_1, X_2, \dots slučajni r -vektorji in naj bo $g : \mathbb{R}^r \rightarrow \mathbb{R}^s$ Borelova funkcija, ki je zvezna skoraj povsod glede na P_X . Potem, če X_n konvergira k X skoraj gotovo, ali v verjetnosti, ali v porazdelitvi, potem tudi $g(X_n)$ konvergira k $g(X)$ v enakem smislu.

Trditev 5.2.4 (Slucki). Naj bodo X, X_1, X_2, \dots in Y_1, Y_2, \dots slučajne spremenljivke. Dalje naj bo $c \in \mathbb{R}$. Privzemimo $X_n \xrightarrow{d} c$ in $Y_n \xrightarrow{d} c$. Tedaj $X_n + Y_n \xrightarrow{d} X + c$, $X_n Y_n \xrightarrow{d} cX$ in, če $c \neq 0$, $X_n/Y_n \xrightarrow{d} X/c$.

Remark. Trditev lahko posplošimo na slučajne vektorje.

Primer. Naj bodo X_1, X_2, \dots neodvisne enako porazdeljene slučajne spremenljivke z disperzijo σ^2 in pričakovano vrednostjo μ . Po centralnem limitnem izreku velja

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow[n \rightarrow \infty]{d} N(0, 1).$$

Potem po krepkem zakonu velikih števil

$$\left(\frac{1}{n} \sum_i X_i, \frac{1}{n} \sum_i X_i^2 \right) \xrightarrow[n \rightarrow \infty]{s.g.} (\mu, \sigma^2 + \mu^2).$$

Upoštevanje zveznosti potem velja

$$S^2 = \frac{n}{n-1} \left(\frac{1}{n} \sum_i X_i^2 - \bar{X}^2 \right) \xrightarrow[n \rightarrow \infty]{s.g.} \sigma^2$$

za

$$S := \sqrt{\frac{1}{n-1} \sum_i (X_i - \bar{X})^2}.$$

Po izreku Sluckega je potem

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \xrightarrow[n \rightarrow \infty]{d} N(0, 1).$$

Trditev 5.2.5. Naj bodo Y, X_1, X_2, \dots slučajni r -vektori, $c \in \mathbb{R}^r$ ter $(a_n)_n$ zaporedje pozitivnih realnih števil z $a_n \rightarrow \infty$. Naj bo $g: \mathbb{R}^r \rightarrow \mathbb{R}$ Borelova funkcija, diferenciable pri c . Če velja

$$a_n(X_n - c) \xrightarrow[n \rightarrow \infty]{d} Y$$

in $dg(c) \neq 0$, potem velja tudi

$$a_n(g(X_n) - g(c)) \xrightarrow[n \rightarrow \infty]{d} \langle \vec{\nabla} \cdot g(c), Y \rangle Y.$$

Dokaz. Spomnimo se: Za vsak $\varepsilon > 0$ obstaja $\delta > 0$, da iz $\|x - c\| \leq \delta$ sledi

$$|g(x) - g(c) - dg(c)(x - c)| \leq \varepsilon \|x - c\|.$$

Izberimo neki $\varepsilon > 0$ in mu pridružimo δ . Pišimo $Z_n = a_n(g(X_n) - g(c)) - a_n dg(c)(X_n - c)$. Pokazali bomo $Z_n \xrightarrow{p} 0$. Ko to vemo, lahko Z_n prištejemo $a_n dg(c)(X_n - c)$ in bo rezultat sledil.

Pokažimo, da za poljuben $\eta > 0$ velja $\lim P(|Z_n| > \eta) = 0$. Velja

$$\begin{aligned} P(|Z_n| > \eta) &= P(|Z_n| > \eta \wedge \|X_n - c\| > \delta) + P(|Z_n| > \eta \wedge \|X_n - c\| \leq \delta) \\ &\leq P(\|X_n - c\| > \delta) + P(|Z_n| > \eta \wedge |Z_n| \leq a_n \varepsilon \|X_n - c\|) \\ &\leq P(\|X_n - c\| > \delta) + P(a_n \varepsilon \|X_n - c\| > \eta). \end{aligned}$$

Ker $a_n^{-1} \rightarrow 0$ in $a_n(X_n - c) \xrightarrow{d} Y$, po Slutskem velja $X_n - c \xrightarrow{d} 0$. Sledi $P(\|X_n - c\| > \delta) \xrightarrow{p} 0$. Ker je norma zvezna, velja tudi $\|a_n(X_n - c)\| \xrightarrow{d} \|Y\|$, zato

$$\limsup P(|Z_n| > \eta) \leq 0 + \limsup P(\|a_n(X_n - c)\| > \eta/\varepsilon).$$

To je enako $1 - F_{\|Y\|}(\eta/\varepsilon)$, če je $F_{\|Y\|}$ zvezna v točki η/ε .

Točk nezveznosti je lahko največ števno mnogo. Izberemo lahko torej tako zaporedje $(\varepsilon_i)_i$, ki pada proti 0, za katero je η/ε_i točka zveznosti $F_{\|Y\|}$. Za vsak ε_i potem velja

$$\limsup P(|Z_n| > \eta) \leq P(\|Y\| > \eta/\varepsilon_i) \xrightarrow{i \rightarrow \infty} 0.$$

Torej limita $\lim P(|Z_n| > \eta)$ obstaja in je enaka 0. □

Posledica 5.2.6 (metoda δ). Z enakimi oznakami, če $a_n(X_n - c) \xrightarrow[n \rightarrow \infty]{d} N(0, \Sigma)$, potem $a_n(g(X_n) - g(c)) \xrightarrow[n \rightarrow \infty]{d} N(0, dg(c)\Sigma dg(c)^T)$.

Izrek 5.2.7 (krepki zakon velikih števil). Naj bodo X_1, X_2, \dots neodvisni enako porazdeljeni slučajni vektorji s pričakovano vrednostjo $\mu \in \mathbb{R}^r$. Potem

$$\frac{1}{n} \sum_i X_i \xrightarrow[n \rightarrow \infty]{s.g.} \mu.$$

Izrek 5.2.8 (centralni limitni izrek). Naj bodo X_1, X_2, \dots neodvisni enako porazdeljeni slučajni vektorji z variančno matriko $\Sigma > 0$ ter pričakovano vrednostjo $\mu \in \mathbb{R}^r$. Tedaj

$$\sqrt{n} \left(\frac{1}{n} \sum_i X_i - \mu \right) \xrightarrow[n \rightarrow \infty]{d} N(0, \Sigma).$$

5.2.1 Doslednost

Naj bodo X_1, X_2, \dots neodvisne replikacije slučajne spremenljivke $\Omega \rightarrow \mathbb{R}$. Spomnimo se, da tako zaporedje X_i lahko modeliramo na $S = \Omega^{\mathbb{N}}$ s predpisom $X_i(s) = X(\omega_i)$ za $s = (\omega_n)_n$. Seveda je P na S definirana s predpisom

$$P(A_1 \times A_2 \times \dots \times A_k \times \Omega \times \Omega \times \dots) = P(A_1) \dots P(A_k).$$

Porazdelitev X_i pripada privzetemu modelu dopustnih (enorazsežnih) porazdelitev \mathcal{P} . Naj bo $e : \mathcal{P} \rightarrow \mathbb{R}^r$ ocenjevana funkcija. Zaporedje cenilk $T_n : \mathbb{R}^n \rightarrow \mathbb{R}^r$ je za e

- KREPKO DOSLEDNO, če $T_n(X_1, \dots, X_n) \xrightarrow{s.g.} e(P_X)$,
- ŠIBKO DOSLEDNO, če $T_n(X_1, \dots, X_n) \xrightarrow{p} e(P_X)$,
- L^2 -DOSLEDNO ali SKN-DOSLEDNO, če $E(\|T_n(X_1, \dots, X_n) - e(P_X)\|^2) \rightarrow 0$

za vsako dopustno porazdelitev $P_X \in \mathcal{P}$ in vsako zaporedje $X_i \sim P_X$ neodvisnih cenilk.

Izrek 5.2.9 (šibki zakon velikih števil Markova). Naj bo X_1, X_2, \dots zaporedje nekoreliranih slučajnih spremenljivk z enako pričakovano vrednostjo μ in disperzijami σ_i^2 , za katere je $\sup \sigma_i^2 < \infty$. Tedaj za vsak $\varepsilon > 0$ velja $\lim P(|\bar{X} - \mu| > \varepsilon) = 0$.

Dokaz. Velja

$$P(|\bar{X} - \mu|^2 > \varepsilon^2) \leq \frac{E(|\bar{X} - \mu|^2)}{\varepsilon^2} = \frac{D(\bar{X})}{\varepsilon^2} = \frac{\sigma_1^2 + \dots + \sigma_n^2}{n^2 \varepsilon^2} \leq \frac{n}{n^2 \varepsilon^2} \sup_i \sigma_i^2$$

za disperzijo D . To konvergira k 0. □

Naj bodo $\mathcal{P}_1, \mathcal{P}_2, \dots$ družine dopustnih porazdelitvenih zakonov na $B(\mathbb{R}), B(\mathbb{R}^2), \dots$ za vektorje $X_1, (X_1, X_2), \dots$, ki so med seboj usklajeni; če je $P_{(X_1, \dots, X_{n+1})} \in \mathcal{P}_{n+1}$, je tudi $P_{(X_1, \dots, X_n)} \in \mathcal{P}_n$. Model je parametričen, če obstajajo usklajene bijektivne korespondence $P_n \rightarrow \Theta$.

Definicija 5.2.10. Naj bodo $e_n : \mathcal{P}_n \rightarrow \mathbb{R}^r$ ocenjevane funkcije in $T_n : \mathbb{R}^n \rightarrow \mathbb{R}^r$ Borelove preslikave. Zaporedje $(T_n)_n$ je za $(e_n)_n$

- ŠIBKO DOSLEDNO, če za vsak η

$$\lim_{n \rightarrow \infty} P(\|T_n(X^{(n)}) - e_n(P_n)\| > \eta) = 0,$$

- L^2 DOSLEDNO ali SKN DOSLEDNO, če

$$\lim_{n \rightarrow \infty} E \left(\left\| T_n(X^{(n)}) - e_n(P_n) \right\|^2 \right) = 0$$

za vsako zaporedje $(P_n \in \mathcal{P}_n)_n$ in vsak vektor $X = (X_1, \dots)$ z lastnostjo $P_{X^{(n)}} = P_n$ za vse n .

Izrek 5.2.11. Naj λ_{\min} označuje najmanjšo lastno vrednost matrike. Če $\lambda_{\min}(Z^T Z) \xrightarrow[n \rightarrow \infty]{d} \infty$, je zaporedje cenilk za regresijski parameter β SKN dosledno.

Trditev 5.2.12. Naj bo $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ statistika in X slučajni vektor z vrednostmi v \mathbb{R}^n . Tedaj je $E(\|TX - E(TX)\|^2) = \text{sl}(\text{var } TX)$.

Dokaz. Račun. Uporabimo dejstvo $\|t\|^2 = \text{sl}(tt^T)$. □

Izrek 5.2.13. Privzemimo naslednji linearni model:

$$\mathcal{P}_n = \{P_{X^{(n)}} \mid X^{(n)} = Z^{(n)}\beta + \varepsilon^{(n)}, \text{var } \varepsilon^{(n)} < \infty\}.$$

Naj bodo $L_n \in \mathbb{R}^{m \times d}$ fiksne realne matrike. Če velja

- $\sup_n \lambda_{\max}(\text{var } \varepsilon^{(n)}) < \infty$,
- $\lim \lambda_{\max}((Z^T Z)') = 0$,
- $\sup_n \max_i \|(L_n)_i\| < \infty$,
- $L_n = U_n Z^{(n)}$

za vse n , je zaporedje cenilk $T_n X^{(n)} = L_n(Z^T Z)' Z^T X$ SKN-dosledno za $e_n(\beta, \varepsilon^{(n)}) = L_n(Z^T Z)' Z^T Z \beta$.

Dokaz. Račun. Nujno naredi. □

5.2.2 Pristranske cenilke

Privzemimo parametrični model s prostorom parametrov $\Theta \subseteq \mathbb{R}^d$ in naj bo $e : \Theta \rightarrow \mathbb{R}^r$ ocenjevana funkcija. PRISTRANSKOST cenilke $T : \mathbb{R}^n \rightarrow \mathbb{R}^r$ za e je $b_\theta(T) = E_\theta(T(X)) - e(\theta)$. Kvaliteto take cenilke merimo s srednjo kvadratno napako $\text{SKN}(\theta) = E_\theta(\|T(X) - e(\theta)\|^2)$. Velja

$$\begin{aligned} \text{SKN}(\theta) &= E_\theta(\|T(X) - E_\theta(T(X)) + E_\theta(T(X)) - e(\theta)\|^2) \\ &= E_\theta(\|T(X) - E_\theta(T(X))\|^2) \\ &\quad + E_\theta(\langle T(X) - E_\theta(T(X)), E_\theta(T(X)) - e(\theta) \rangle) \\ &\quad + E_\theta(\|E_\theta(T(X)) - e(\theta)\|^2). \end{aligned}$$

Ker je $E_\theta(T(X) - E_\theta(T(X))) = 0$, sledi

$$\begin{aligned} \text{SKN}(\theta) &= E_\theta(\|T(X) - E_\theta(T(X))\|^2) + E_\theta(\|E_\theta(T(X) - e(\theta))\|^2) \\ &= \text{sl var}_\theta(T(X)) + \|b_\theta(T)\|^2. \end{aligned}$$

Definicija 5.2.14. Privzemimo model $(\mathcal{P}_n)_n$ za x_1, x_2, \dots in naj bo $e_n : \mathcal{P}_n \rightarrow \mathbb{R}^r$ zaporedje ocenjevanih funkcij ter $T_n : \mathbb{R}^n \rightarrow \mathbb{R}^r$ zaporedje cenilk za e_n . To zaporedje je NEPRISTRANSKO ZA e_n V LIMITI, če velja

$$\lim_{n \rightarrow \infty} E_{P_n}(T_n - e_n(P_n)) = 0$$

za vsako dopustno porazdelitev $(P_n)_n \in (\mathcal{P}_n)_n$.

Definicija 5.2.15. Naj bo sedaj $Y_{(\mathcal{P}_n)_n}$ družina porazdelitev. Dalje naj bo $(a_n)_n$ zaporedje pozitivnih realnih števil in $T_n : \mathbb{R}^n \rightarrow \mathbb{R}^r$ zaporedje statistik. Naj bo $e_n : \mathcal{P}_n \rightarrow \mathbb{R}^r$ zaporedje ocenjevanih funkcij. Če velja $\lim a_n \in (0, \infty]$ in

$$a_n(T_n(X^{(n)}) - e_n(P_n)) \xrightarrow[n \rightarrow \infty]{d} Y_{(P_n)_n}$$

za vse $(P_n)_n \in (\mathcal{P}_n)_n$ in vsak X z $X^{(n)} \sim P_n$, potem pravimo, da je T_n ASIMPTOTIČNO NEPRISTRANSKO ZAPOREDJE CENILK za $(e_n)_n$.

Primer. Če so T_n dosledne, velja $T_n(X^{(n)}) - e_n(P_n) \rightarrow 0$.

Primer. Iz CLI dobimo $\sqrt{n}(\bar{X} - \mu) \rightarrow N(0, \Sigma)$. V tem primeru so porazdelitve res odvisne od $(P_n)_n$ (zaradi Σ).

5.2.3 Asimptotična normalnost

V zgornjem kontekstu (definicija 5.2.14) pravimo, da je zaporedje $(T_n)_n$ ASIMPTOTIČNO NORMALNO zaporedje cenilk za $(e_n)_n$, če obstaja zaporedje funkcij $V_n : \mathcal{P}_n \rightarrow \text{SPD}(r)$ (simetrične pozitivno definitne matrike $r \times r$), za katerega velja

$$V_n(P_n)^{-1/2}(T_n(X^{(n)}) - e_n(P_n)) \xrightarrow[n \rightarrow \infty]{d} N(0, I)$$

za vsak $(P_n)_n \in (\mathcal{P}_n)_n$ in X s to porazdelitvijo.

Rečemo, da je $V_n(P_n)$ ASIMPTOTIČNA VARIANCA za $T_n(X^{(n)})$. Če je $V_n(P_n) = \frac{1}{n}A((P_n)_n)$ za družino simetričnih pozitivno definitnih matrik $A : (\mathcal{P}_n)_n \rightarrow \mathbb{R}^{r \times r}$, je to ASIMPTOTIČNA VARIANCA V OŽJEM SMISLU.

Opomba. Asimptotična varianca ni enolična, lahko jo npr. pomnožimo s poljubnim zaporedjem, ki konvergira k 1.

Izrek 5.2.16. Privzamemo model linearne regresije \mathcal{P}'_n od prej, $X = Z\beta + \varepsilon$. Privzemimo, da $\frac{1}{v}Z^T Z$ konvergira k neki simetrični pozitivno definitni matriki razsežnosti $d \times d$. Teda j imajo $Z^{(n)}$ poln rang za dovolj velika števila n in za $\hat{\beta} = (Z^T Z)^{-1}Z^T X$ velja

$$\frac{1}{\sigma}(Z^T Z)(\hat{\beta} - \beta) \xrightarrow[n \rightarrow \infty]{d} N(0, I).$$

5.2.4 Konstrukcija cenilk

Obravnavamo neodvisne in enako porazdeljene slučajne spremenljivke X , s pripadajočim modelom, parametriziranim z $\Theta^{\text{odp}} \subseteq \mathbb{R}^d$. Privzemimo, da obstajajo momenti $\mu_j = \mu_j(\theta) = e_j(\theta) = E_\theta(X^j)$ za $1 \leq j \leq d$, in da je funkcija $e = (e_1, \dots, e_d) : \Theta \rightarrow \mathbb{R}^d$ obrnljiva z inverzom $g : \text{im } e \rightarrow \Theta$. Za momente imamo standardne cenilke

$$\hat{\mu}_j = \frac{1}{n} \sum_{i=1}^n X_k^j,$$

ki so krepko dosledne cenilke za momente po krepkem zakonu velikih števil. Če je g zvezna, je $g(\hat{\mu}_1, \dots, \hat{\mu}_d)$ tudi krepko dosledna cenilka za θ .

Dodatno privzemimo obstoj momentov $E_\theta(X^j)$ za $j \leq 2d$. Vektorji $(X_i, X_i^2, \dots, X_i^d)$ so neodvisni in enako porazdeljeni s pričakovano vrednostjo (μ_1, \dots, μ_d) in variančno matriko $\Sigma = [\mu_{k+l} - \mu_k \mu_l]_{k,l}$. Privzemimo, da je Σ neizrojena, da po CLI velja

$$\sqrt{n}((\hat{\mu}_1, \dots, \hat{\mu}_d) - (\mu_1, \dots, \mu_d)) \xrightarrow[n \rightarrow \infty]{d} N(0, \Sigma).$$

Če je g diferenciable, potem

$$\sqrt{n}(g(\hat{\mu}_1, \dots, \hat{\mu}_d) - g(\mu_1, \dots, \mu_d)) \xrightarrow[n \rightarrow \infty]{d} N(0, Dg(\mu)\Sigma Dg^T(\mu)).$$

To pomeni, da je $g(\hat{\mu}_1, \dots, \hat{\mu}_d)$ asimptotično normalno zaporedje cenilk.

Alternativno lahko poiščemo cenilko po metodi največjega verjetja. Naj bo $X : \Omega \rightarrow \mathbb{R}^m$ proučevani slučajni vektor z modelom, parametriziranim z odprto množico $\Theta \subseteq \mathbb{R}^d$. Privzemimo gostote $f(\cdot, \theta)$, da je

$$P_\theta(X \in B) = \int_B f(x, \theta) d\nu(x).$$

Tu je ν neka σ -končna mera, ki dominira model $\{P_\theta \mid \theta \in \Theta\} \ll \nu$. Funkciji $L : \mathbb{R}^n \times \Theta \rightarrow [0, \infty)$, definirani z $L(x, \theta) = f(x, \theta)$, pravimo VERJETJE. Če za dano realizacijo x vektorja X obstaja $\hat{\theta} \in \mathbb{R}^d$, za katerega je $L(x, \hat{\theta})$ maksimum vrednosti $\{L(x, \theta) \mid \theta \in \bar{\Theta}\}$, mu pravimo OCENA PO MNV za x . Če $\hat{\theta} = \hat{\theta}(x)$ obstaja za ν -skoraj vse x , funkciji $\hat{\theta} : \mathbb{R}^d \rightarrow \bar{\Theta}$ pravimo CENILKA NAJVEČJEGA VERJETJA ZA θ .

Trditev 5.2.17. Če obstaja enolična $\hat{\theta}$ in je $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ zadostna statistika, velja $\hat{\theta}(x) = \hat{\theta}(Tx)$.

Dokaz. Po Fisher-Neymannu je $f(x, \theta) = g(Tx, \theta)h(x)$. Brez škode za splošnost je $h(x) > 0$, sicer $\hat{\theta}$ ni enolična. Maksimizacija se reducira na maksimizacijo funkcije $\theta \mapsto g(Tx, \theta)$. \square

Naj bo \mathcal{P} parametrični model z gostotami (glede na neko σ -končno mero) $\{f(\cdot, \theta) \mid \theta \in \Theta\}$, kjer je $\Theta^{\text{odp}} \subseteq \mathbb{R}^d$. Privzemimo dodatne regularnostne privzetke, med drugim da je množica $S = \{x \mid f(x, \theta) > 0\}$ neodvisna od θ . Velja

$$0 = \partial_{\theta_i} 1 = \partial_{\theta_i} \int f(x, \theta) d\nu(x) = \int \frac{\partial(\log f)}{\partial \theta_i} f(x, \theta) d\nu(x) = E_{\theta}(\partial_{\theta_i}(\log f(X, \theta))).$$

Funkciji

$$V_{\theta}(x) = \text{grad}_{\theta}(\log L)(x, \theta)$$

pravimo FUNKCIJA ZBIRA. Ker je $E_{\theta}(V_{\theta}(X)) = 0$ za vse θ , velja

$$0 \leq \text{var}_{\theta}(V_{\theta}(X)) = E_{\theta}(V_{\theta}(X)V_{\theta}(X)^T),$$

to matriko imenujemo FISHERJEVA INFORMACIJA in označimo s $\text{FI}(\theta)$. Če zgornje še enkrat odvajamo po θ_j , dobimo

$$0 = \int \frac{\partial^2(\log f)}{\partial \theta_i \partial \theta_j} f(x, \theta) d\nu(x) + \int \frac{\partial(\log f)}{\partial \theta_i} \frac{\partial(\log f)}{\partial \theta_j} f(x, \theta) d\nu(x)$$

kar je natanko

$$\text{FI}(\theta) = -E_{\theta}(H(\log L)(X, \theta)).$$

Če nam gostote f dopuščajo faktorizacijo

$$f(x, \theta) = h(x) \exp(-\psi(\theta) + \langle Q(\theta), T(x) \rangle),$$

kjer so $\psi : \Theta \rightarrow \mathbb{R}$, $Q : \Theta \rightarrow \mathbb{R}^m$, $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ in $h : \mathbb{R}^n \rightarrow [0, \infty)$ primerne funkcije, potem pravimo, da je model EKSPONENTEN. Pravimo, da je model NARAVNO PARAMETRIZIRAN, če je $Q = \text{id}$.

Pod nekaterimi regularnostnimi privzetki lahko zagotovimo obstoj zaporedja slučajnih vektorjev $\hat{\theta}^{(n)}$ z naslednjimi lastnostmi:

- verjetnost, da $\hat{\theta}^{(n)}$ reši logaritemsko enačbo verjetja, konvergira k 1,
- zaporedje je dosledno za θ ,
- velja $\sqrt{n}(\hat{\theta}^{(n)} - \theta) \rightarrow N(0, \text{FI}(\theta)^{-1})$.

5.3 Preizkušanje domnev

Obravnavamo model \mathcal{P} za slučajni vektor X . Privzemimo, da obstaja dekompozicija $\mathcal{P} = \mathcal{H} \cup \mathcal{A}$ na neprazni disjunktni množici. NERANDOMIZIRAN POIZKUS DOMNEVE \mathcal{H} proti \mathcal{A} je odločitveno pravilo $\phi : \mathbb{R}^n \rightarrow \{0, 1\}$, ki ga izvedemo tako:

- če je $\phi(x) = 1$, domnevo \mathcal{H} zavrnemo,
- če je $\phi(x) = 0$, domneve \mathcal{H} ne zavrnemo.

Če je \mathcal{P} parametriziran, torej v bijektivni korespondenci z množico Θ , potem označimo slike \mathcal{H} in \mathcal{A} pod to bijekcijo s H in A . Pravimo, da preizkušamo H proti A .

Poznamo dve vrsti napake. Če je v resnici $P_X \in H$, mi pa domnevo vseeno zavržemo, temu pravimo NAPAKA PRVE VRSTE, če pa $P_X \notin H$, a mi domneve ne zavržemo, je to NAPAKA DRUGE VRSTE. Pri primernih zveznostnih predpostavkah tipično velja

$$\sup\{P_\theta(\text{napaka prve vrste}) \mid \theta \in H\} + \sup\{P_\theta(\text{napaka druge vrste}) \mid \theta \in A\} = 1,$$

torej popolnega preizkusa ni. V praksi odločitev o zavrnitvi napravimo na podlagi neke testne statistike T v smislu

$$\phi(x) = \begin{cases} 1, & T(x) \in B \\ 0, & T(x) \notin B \end{cases}$$

za neko zavrnitveno območje B .

Zaradi komplementarnosti maksimalnih vrednosti napak obeh vrst izberemo pomembnejšo in jo skušamo omejiti. Po potrebi zamenjamo H in A , da je to napaka prve vrste. VELIKOST PREIZKUSA je potem

$$\sup_{\theta \in H} P_\theta(\text{napaka prve vrste}).$$

Pravimo, da ima preizkus STOPNJO ZNAČILNOSTI ali STOPNJO TVEGANJA α , če je njegova velikost največ α . Standardne izbire so $\alpha \in \{0.05, 0.1, 0.01\}$.

5.3.1 Preizkušanje na podlagi razmerja verjetij

Privzamemo parametrični model s prostorom parametrov $\Theta \subseteq \mathbb{R}^d$ in gladka verjetja $L(x, \theta)$. Naj bo $H \subseteq \Theta$ preizkušana domneva. RAZMERJE VERJETIJ je H je funkcija $\lambda : \mathbb{R}^n \rightarrow [0, 1]$, definirana z

$$\lambda(x) = \frac{\sup\{L(x, \theta) \mid \theta \in H\}}{\sup\{L(x, \theta) \mid \theta \in \Theta\}}.$$

Vedno lahko potem konstruiramo preizkus oblike

$$\phi(x) = \begin{cases} 1, & \lambda(x) < D \\ 0, & \lambda(x) \geq D \end{cases}$$

za neko primerno konstanto D .

Izrek 5.3.1 (Wilksov izrek o asimptotični porazdelitvi razmerja verjetij). *Naj bodo X_1, X_2, \dots neodvisni enako porazdeljeni slučajni vektorji z gostoto $f(x, \theta)$, kjer je $\theta \in \Theta$ in je Θ gladka končnorazsežna mnogoterost. Dalje privzemimo, da je H zaprta gladka podmnogoterost brez roba, tako da je $H \subseteq \Theta$ prava vložitev. Naj veljajo primerni gladkostni privzetki na gostote, ki zagotavljajo asimptotično normalnost cenilk največjega verjetja. Če dejanski parameter θ pripada H , potem*

$$-2 \log \lambda(X_1, \dots, X_n) \xrightarrow[n \rightarrow \infty]{d} \chi^2(\dim \Theta - \dim H).$$