

---

# A Computation and Communication Efficient Method for Distributed Nonconvex Problems in the Partial Participation Setting

---

Anonymous Author(s)

Affiliation

Address

email

## Abstract

1 We present a new method that includes three key components of distributed opti-  
2 mization and federated learning: variance reduction of stochastic gradients, partial  
3 participation, and compressed communication. We prove that the new method has  
4 optimal oracle complexity and state-of-the-art communication complexity in the  
5 partial participation setting. Regardless of the communication compression feature,  
6 our method successfully combines variance reduction and partial participation: we  
7 get the optimal oracle complexity, never need the participation of all nodes, and do  
8 not require the bounded gradients (dissimilarity) assumption.

## 9 1 Introduction

10 Federated and distributed learning have become very popular in recent years (Konečný et al., 2016;  
11 McMahan et al., 2017). The current optimization tasks require much computational resources and  
12 machines. Such requirements emerge in machine learning, where massive datasets and computations  
13 are distributed between cluster nodes (Lin et al., 2017; Ramesh et al., 2021). In federated learning,  
14 nodes, represented by mobile phones, laptops, and desktops, do not send their data to a server due to  
15 privacy and their huge number (Ramaswamy et al., 2019), and the server remotely orchestrates the  
16 nodes and communicates with them to solve an optimization problem.

17 As in classical optimization tasks, one of the main current challenges is to find **computationally**  
18 **efficient** optimization algorithms. However, the nature of distributed problems induces many other  
19 (Kairouz et al., 2021), including i) **partial participation** of nodes in algorithm steps: due to stragglers  
20 (Li et al., 2020) or communication delays (Vogels et al., 2021), ii) **communication bottleneck**: even  
21 if a node participates, it can be costly to transmit information to a server or other nodes (Alistarh  
22 et al., 2017; Ramesh et al., 2021; Kairouz et al., 2021; Sapio et al., 2019; Narayanan et al., 2019). It  
23 is necessary to develop a method that considers these problems.

## 24 2 Optimization Problem

25 Let us consider the nonconvex distributed optimization problem

$$\min_{x \in \mathbb{R}^d} \left\{ f(x) := \frac{1}{n} \sum_{i=1}^n f_i(x) \right\}, \quad (1)$$

26 where  $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$  is a smooth nonconvex function for all  $i \in [n] := \{1, \dots, n\}$ . The full  
27 information about function  $f_i$  is stored on  $i^{\text{th}}$  node. The communication between nodes is maintained  
28 in the parameters server fashion (Kairouz et al., 2021): we have a server that receives compressed

information from nodes, updates a state, and broadcasts an updated model.<sup>1</sup> Since we work in the nonconvex world, our goal is to find an  $\varepsilon$ -solution ( $\varepsilon$ -stationary point) of (1): a (possibly random) point  $\hat{x} \in \mathbb{R}^d$ , such that  $\mathbb{E}[\|\nabla f(\hat{x})\|^2] \leq \varepsilon$ .

We consider three settings:

**1. Gradient Setting.** The  $i^{\text{th}}$  node has only access to the gradient  $\nabla f_i : \mathbb{R}^d \rightarrow \mathbb{R}^d$  of function  $f_i$ . Moreover, the following assumptions for the functions  $f_i$  hold.

**Assumption 1.** *There exists  $f^* \in \mathbb{R}$  such that  $f(x) \geq f^*$  for all  $x \in \mathbb{R}$ .*

**Assumption 2.** *The function  $f$  is  $L$ -smooth, i.e.,  $\|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|$  for all  $x, y \in \mathbb{R}^d$ .*

**Assumption 3.** *The functions  $f_i$  are  $L_i$ -smooth for all  $i \in [n]$ . Let us define  $\widehat{L}^2 := \frac{1}{n} \sum_{i=1}^n L_i^2$ .<sup>2</sup>*

**2. Finite-Sum Setting.** The functions  $\{f_i\}_{i=1}^n$  have the finite-sum form

$$f_i(x) = \frac{1}{m} \sum_{j=1}^m f_{ij}(x), \quad \forall i \in [n], \quad (2)$$

where  $f_{ij} : \mathbb{R}^d \rightarrow \mathbb{R}$  is a smooth nonconvex function for all  $j \in [m]$ .

We assume that Assumptions 1, 2 and 3 hold and the following assumption.

**Assumption 4.** *The function  $f_{ij}$  is  $L_{ij}$ -smooth for all  $i \in [n], j \in [m]$ . Let  $L_{\max} := \max_{i \in [n], j \in [m]} L_{ij}$ .*

**3. Stochastic Setting.** The function  $f_i$  is an expectation of a stochastic function,

$$f_i(x) = \mathbb{E}_{\xi} [f_i(x; \xi)], \quad \forall i \in [n], \quad (3)$$

where  $f_i : \mathbb{R}^d \times \Omega_{\xi} \rightarrow \mathbb{R}$ . For a fixed  $x \in \mathbb{R}$ ,  $f_i(x; \xi)$  is a random variable over some distribution  $\mathcal{D}_i$ , and, for a fixed  $\xi \in \Omega_{\xi}$ ,  $f_i(x; \xi)$  is a smooth nonconvex function. The  $i^{\text{th}}$  node has only access to a stochastic gradients  $\nabla f_i(\cdot; \xi_{ij})$  of the function  $f_i$  through the distribution  $\mathcal{D}_i$ , where  $\xi_{ij}$  is a sample from  $\mathcal{D}_i$ . We assume that Assumptions 1, 2 and 3 hold and the following assumptions.

**Assumption 5.** *For all  $i \in [n]$  and for all  $x \in \mathbb{R}^d$ , the stochastic gradient  $\nabla f_i(x; \xi)$  is unbiased and has bounded variance, i.e.,  $\mathbb{E}_{\xi} [\nabla f_i(x; \xi)] = \nabla f_i(x)$ , and  $\mathbb{E}_{\xi} [\|\nabla f_i(x; \xi) - \nabla f_i(x)\|^2] \leq \sigma^2$ , where  $\sigma^2 \geq 0$ .*

**Assumption 6.** *For all  $i \in [n]$  and for all  $x, y \in \mathbb{R}$ , the stochastic gradient  $\nabla f_i(x; \xi)$  satisfies the mean-squared smoothness property, i.e.,  $\mathbb{E}_{\xi} [\|\nabla f_i(x; \xi) - \nabla f_i(y; \xi)\|^2] \leq L_{\sigma}^2 \|x - y\|^2$ .*

We compare algorithms using the *oracle complexity*, i.e., the number of (stochastic) gradients that each node has to calculate to get  $\varepsilon$ -solution, and the *communication complexity*, i.e., the number of bits that each node has to send to the server to get  $\varepsilon$ -solution.

## 2.1 Unbiased Compressors

We use the concept of unbiased compressors to alleviate the communication bottleneck. The unbiased compressors quantize and/or sparsify vectors that the nodes send to the server.

**Definition 1.** A stochastic mapping  $\mathcal{C} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is an *unbiased compressor* if there exists  $\omega \in \mathbb{R}$  such that

$$\mathbb{E}[\mathcal{C}(x)] = x, \quad \mathbb{E}[\|\mathcal{C}(x) - x\|^2] \leq \omega \|x\|^2, \quad (4)$$

for all  $x \in \mathbb{R}^d$ .

We denote a set of stochastic mappings that satisfy Definition 1 as  $\mathbb{U}(\omega)$ . In our methods, the nodes make use of unbiased compressors  $\{\mathcal{C}_i\}_{i=1}^n$ . The community developed a large number of unbiased

<sup>1</sup>Note that this strategy can be used in peer-to-peer communication, assuming that the server is an abstraction and all its algorithmic steps are performed on each node.

<sup>2</sup>Note that  $L \leq \widehat{L}$ ,  $\widehat{L} \leq L_{\max}$ , and  $\widehat{L} \leq L_{\sigma}$ .

Table 1: Summary of methods that solve the problem (1) in the stochastic setting (3). Abbr.: *VR* (Variance Reduction) = Does a method have the optimal oracle complexity  $\mathcal{O}\left(\frac{\sigma^2}{\varepsilon} + \frac{\sigma}{\varepsilon^{3/2}}\right)$ ? *PP* (Partial Participation) = Does a method support partial participation from Section 2.2? *CC* = Does a method have the communication complexity equals to  $\mathcal{O}\left(\frac{\omega}{\sqrt{n\varepsilon}}\right)$ ?

Method	VR	PP	CC	Limitations
<b>SPIDER, SARAH, PAGE, STORM</b> (Fang et al., 2018; Nguyen et al., 2017) (Li et al., 2021a; Cutkosky and Orabona, 2019)	✓	✗	✗	—
<b>MARINA</b> (Gorbunov et al., 2021)	✓	✗ <sup>(a)</sup>	✓ <sup>(b)</sup>	Suboptimal convergence rate (see (Tyurin and Richtárik, 2023)).
<b>FedPAGE</b> (Zhao et al., 2021b)	✗	✗ <sup>(a)</sup>	✗	Suboptimal oracle complexity $\mathcal{O}\left(\frac{\sigma^2}{\varepsilon^2}\right)$ .
<b>FRECON</b> (Zhao et al., 2021a)	✗	✓	✓	—
<b>FedAvg</b> (McMahan et al., 2017; Karimireddy et al., 2020b)	✗	✓	✗	Bounded gradients (dissimilarity) assumption of $f_i$ .
<b>SCAFFOLD</b> (Karimireddy et al., 2020b)	✗	✓	✗	Suboptimal convergence rate <sup>(e)</sup> .
<b>MIME<sup>(c)</sup></b> (Karimireddy et al., 2020a)	✗ <sup>(d)</sup>	✓	✗	Calculates full gradient. Bounded gradients (dissimilarity) assumption of $f_i$ . Suboptimal oracle compl. $\mathcal{O}\left(1/\varepsilon^{3/2}\right)$ in the setting (2).
<b>CE-LSGD (for Partial Participation)<sup>(c)</sup></b> (Patel et al., 2022) (concurrent work)	✓	✓	✗	Bounded gradients (dissimilarity) assumption of $f_i$ . Suboptimal oracle compl. $\mathcal{O}\left(1/\varepsilon^{3/2}\right)$ in the setting (2).
<b>DASHA</b> (Tyurin and Richtárik, 2023)	✓ ✗	✗ ✓	✓ ✓	—
<b>DASHA-PP</b> (new)	✓	✓	✓	—

<sup>(a)</sup> **MARINA** and **FedPAGE**, with a small probability, require the participation of all nodes so that they can not support partial participation from Section 2.2. Moreover, these methods provide suboptimal oracle complexities.

<sup>(b)</sup> On average, **MARINA** provides the compressed communication mechanism with complexity  $\mathcal{O}\left(\frac{\omega}{\sqrt{n\varepsilon}}\right)$ . However, with a small probability, this method sends non-compressed vectors.

<sup>(c)</sup> Note that **MIME** and **CE-LSGD** can not be directly compared with **DASHA-PP** because **MIME** and **CE-LSGD** consider the online version of the problem (1), and require more strict assumptions.

<sup>(d)</sup> Although **MIME** obtains the convergence rate  $\mathcal{O}\left(\frac{1}{\varepsilon^{3/2}}\right)$  of a variance reduced method, it requires the calculation of the full (exact) gradients.

<sup>(e)</sup> It can be seen when  $\sigma^2 = 0$ . Let us consider the  $s$ -nice sampling of the nodes, then **SCAFFOLD** requires  $\mathcal{O}\left(\frac{n^{3/2}}{\varepsilon s^{3/2}}\right)$  communication rounds to get  $\varepsilon$ -solution, while **DASHA-PP** requires  $\mathcal{O}\left(\frac{\sqrt{n}}{\varepsilon s}\right)$  communication rounds (see Theorem 4 with  $\omega = 0$ ,  $b = \frac{p_a}{2-p_a}$ , and  $p_a = \frac{s}{n}$ ).

compressors, including *RandK* (see Definition 5) (Beznosikov et al., 2020; Stich et al., 2018), Adaptive sparsification (Wangni et al., 2018) and Natural compression and dithering (Horváth et al., 2019a). We are aware of correlated compressors by Szlendak et al. (2021) and quantizers by Suresh et al. (2022) that help in the homogeneous regimes, but in this work, we are mainly concentrated on generic heterogeneous regimes, though, for simplicity, assume the independence of the compressors.

**Assumption 7.**  $\mathcal{C}_i \in \mathbb{U}(\omega)$  for all  $i \in [n]$ , and the compressors are statistically independent.

## 2.2 Nodes Partial Participation Assumptions

We now try to formalize the notion of partial participation. Let us assume that we have  $n$  events  $\{i^{\text{th}}$  node is *participating* $\}$  with the following properties.

**Assumption 8.** *The partial participation of nodes has the following distribution: exists constants  $p_a \in (0, 1]$  and  $p_{aa} \in [0, 1]$ , such that*

$$1. \quad \text{Prob}(i^{\text{th}} \text{ node is participating}) = p_a \quad \forall i \in [n],$$

$$2. \quad \text{Prob}(i^{\text{th}} \text{ and } j^{\text{th}} \text{ nodes are participating}) = p_{aa} \quad \forall i \neq j \in [n].$$

Table 2: Summary of methods that solve the problem (1) in the finite-sum setting (2). Abbr.: VR (Variance Reduction) = Does a method have the optimal oracle complexity  $\mathcal{O}\left(m + \frac{\sqrt{m}}{\varepsilon}\right)$ ? PP and CC are defined in Table 1.

Method	VR	PP	CC	Limitations
SPIDER, PAGE (Fang et al., 2018; Li et al., 2021a)	✓	✗	✗	—
MARINA (Gorbunov et al., 2021)	✓	✗ <sup>(a)</sup>	✓ <sup>(b)</sup>	Suboptimal convergence rate (see (Tyurin and Richtárik, 2023)).
ZeroSARAH (Li et al., 2021b)	✓	✓	✗	Only homogeneous regime, i.e., the functions $f_i$ are equal.
FedPAGE (Zhao et al., 2021b)	✗	✗ <sup>(a)</sup>	✗	Suboptimal oracle complexity $\mathcal{O}\left(\frac{m}{\varepsilon}\right)$ .
DASHA (Tyurin and Richtárik, 2023)	✓	✗	✓	—
DASHA-PP (new)	✓	✓	✓	—

<sup>(a)</sup>, <sup>(b)</sup> : see Table 1.

$$p_{aa} \leq p_a^2, \quad (5)$$

and these events from different communication rounds are independent.

We are not fighting for the full generality and believe that more complex sampling strategies can be considered in the analysis. For simplicity, we settle upon Assumption 8. Standard partial participation strategies, including  $s$ -nice sampling, where the server chooses uniformly  $s$  nodes without replacement ( $p_a = s/n$  and  $p_{aa} = s(s-1)/n(n-1)$ ), and independent participation, where each node independently participates with probability  $p_a$  (due to independence, we have  $p_{aa} = p_a^2$ ), satisfy Assumption 8. In the literature,  $s$ -nice sampling is one of the most popular strategies (Zhao et al., 2021a; Richtárik et al., 2021; Reddi et al., 2020; Konečný et al., 2016).

### 3 Motivation and Related Work

The main goal of our paper is to develop a method for the nonconvex distributed optimization that will include three key features: variance reduction of stochastic gradients, compressed communication, and partial participation. We now provide an overview of the literature (see also Table 1 and Table 2).

#### 1. Variance reduction of stochastic gradients

It is important to consider finite-sum (2) and stochastic (3) settings because, in machine learning tasks, either the number of local functions  $m$  is huge or the functions  $f_i$  is an expectation of a stochastic function due to the batch normalization (Ioffe and Szegedy, 2015) or random augmentation (Goodfellow et al., 2016), and it is infeasible to calculate the full gradients analytically. Let us recall the results from the nondistributed optimization. In the gradient setting, the optimal oracle complexity is  $\mathcal{O}(1/\varepsilon)$ , achieved by the vanilla gradient descent (GD) (Carmon et al., 2020; Nesterov, 2018). In the finite-sum setting and stochastic settings, the optimal oracle complexities are  $\mathcal{O}\left(m + \frac{\sqrt{m}}{\varepsilon}\right)$  and  $\mathcal{O}\left(\frac{\sigma^2}{\varepsilon} + \frac{\sigma}{\varepsilon^{3/2}}\right)$  (Fang et al., 2018; Li et al., 2021a; Arjevani et al., 2019), accordingly, achieved by methods SPIDER, SARAH, PAGE, and STORM from (Fang et al., 2018; Nguyen et al., 2017; Li et al., 2021a; Cutkosky and Orabona, 2019).

#### 2. Compressed communication

In distributed optimization (Ramesh et al., 2021; Xu et al., 2021), lossy communication compression can be a powerful tool to increase the communication speed between the nodes and the server. Different types of compressors are considered in the literature, including unbiased compressors (Alistarh et al., 2017; Beznosikov et al., 2020; Szlendak et al., 2021), contractive (biased) compressors (Richtárik et al., 2021), 3PC compressors (Richtárik et al., 2022). We will focus on unbiased compressors because methods DASHA and MARINA (Tyurin and Richtárik, 2023; Szlendak et al.,

2021; Gorbunov et al., 2021) that employ unbiased compressors provide the current theoretical state-of-the-art (SOTA) communication complexities.

Many methods analyzed optimization methods with the unbiased compressors (Alistarh et al., 2017; Mishchenko et al., 2019; Horváth et al., 2019b; Gorbunov et al., 2021; Tyurin and Richtárik, 2023). In the gradient setting, the methods MARINA and DASHA by Gorbunov et al. (2021) and Tyurin and Richtárik (2023) establish the current SOTA communication complexity, each method needs  $\frac{1+\omega/\sqrt{n}}{\varepsilon}$  communication rounds to get an  $\varepsilon$ -solution. In the finite-sum and stochastic settings, the current SOTA communication complexity is attained by the DASHA method, while maintaining the optimal oracle complexities  $\mathcal{O}\left(m + \frac{\sqrt{m}}{\varepsilon\sqrt{n}}\right)$  and  $\mathcal{O}\left(\frac{\sigma^2}{\varepsilon n} + \frac{\sigma}{\varepsilon^{3/2}n}\right)$  per node.

### 3. Partial participation

From the beginning of federated learning era, the partial participation has been considered to be the essential feature of distributed optimization methods (McMahan et al., 2017; Konečný et al., 2016; Kairouz et al., 2021). However, previously proposed methods have limitations: i) methods MARINA and FedPAGE from (Gorbunov et al., 2021; Zhao et al., 2021b) still require synchronization of all nodes with a small probability. ii) in the stochastic settings, methods FedAvg, SCAFFOLD, and FRECON with the partial participation mechanism (McMahan et al., 2017; Karimireddy et al., 2020b; Zhao et al., 2021a) provide results without variance reduction techniques from (Fang et al., 2018; Li et al., 2021a; Cutkosky and Orabona, 2019) and, therefore, get suboptimal oracle complexities. Note that FRECON and DASHA reduce the variance *only from compressors* (in the partial participation and stochastic setting). iii) in the finite-sum setting, the ZeroSARAH method by Li et al. (2021b) focuses on the homogeneous regime only (the functions  $f_i$  are equal). iv) The MIME method by Karimireddy et al. (2020a) and the CE-LSGD method (for Partial Participation) by the concurrent paper (Patel et al., 2022) consider the online version of the problem (1). Therefore, MIME and CE-LSGD (for Partial Participation) require stricter assumptions, including the bounded inter-client gradient variance assumption. In the finite-sum setting (2), MIME and CE-LSGD obtain a suboptimal oracle complexity  $\mathcal{O}(1/\varepsilon^{3/2})$  while, in the full participation setting, it is possible to get the complexity  $\mathcal{O}(1/\varepsilon)$ .

## 4 Contributions

We propose a new method DASHA-PP for the nonconvex distributed optimization.

- As far as we know, this is the first method that includes three key ingredients of federated learning methods: *variance reduction of stochastic gradients, compressed communication, and partial participation*.
- Moreover, this is the first method that combines *variance reduction of stochastic gradients and partial participation* flawlessly: i) it gets the optimal oracle complexity ii) does not require the participation of all nodes iii) does not require the bounded gradients assumption of the functions  $f_i$ .
- We prove convergence rates and show that this method has *the optimal oracle complexity and the state-of-the-art communication complexity in the partial participation setting*. Moreover, in our work, we observe a nontrivial side-effect from mixing the variance reduction of stochastic gradients and partial participation. It is a general problem not related to our methods or analysis that we discuss in Section C.

## 5 Algorithm Description and Main Challenges Towards Partial Participation

We now present DASHA-PP (see Algorithm 1), a family of methods to solve the optimization problem (1). When we started investigating the problem, we took DASHA as a baseline method for two reasons: the family of algorithms DASHA provides the current state-of-the-art communication complexities in the *non-partial participation* setting, and, unlike MARINA, it does not send non-compressed gradients and does not synchronize all nodes. Let us briefly discuss the main idea of DASHA, its problem in the *partial participation* setting, and why the refinement of DASHA is not an exercise.

In fact, the original DASHA method supports the partial participation of nodes *in the gradient setting*. Since the nodes only do the following steps (see full algorithm in Algorithm 6):

$$g_i^{t+1} = g_i^t + \mathcal{C}_i(\nabla f_i(x^{t+1})) - (1-a)\nabla f_i(x^t) - ag_i^t.$$

---

**Algorithm 1 DASHA-PP**

---

- 1: **Input:** starting point  $x^0 \in \mathbb{R}^d$ , stepsize  $\gamma > 0$ , momentum  $a \in (0, 1]$ , momentum  $b \in (0, 1]$ , probability  $p_{\text{page}} \in (0, 1]$  (only in **DASHA-PP-PAGE**), batch size  $B$  (only in **DASHA-PP-PAGE**, **DASHA-PP-FINITE-MVR** and **DASHA-PP-MVR**), probability  $p_a \in (0, 1]$  that a node is *participating*<sup>(a)</sup>, number of iterations  $T \geq 1$
  - 2: Initialize  $g_i^0 \in \mathbb{R}^d$ ,  $h_i^0 \in \mathbb{R}^d$  on the nodes and  $g^0 = \frac{1}{n} \sum_{i=1}^n g_i^0$  on the server
  - 3: Initialize  $h_{ij}^0 \in \mathbb{R}^d$  on the nodes and take  $h_i^0 = \frac{1}{m} \sum_{j=1}^m h_{ij}^0$  (only in **DASHA-PP-FINITE-MVR**)
  - 4: **for**  $t = 0, 1, \dots, T - 1$  **do**
  - 5:    $x^{t+1} = x^t - \gamma g^t$
  - 6:   Broadcast  $x^{t+1}, x^t$  to all *participating*<sup>(a)</sup> nodes
  - 7:   **for**  $i = 1, \dots, n$  in parallel **do**
  - 8:     **if**  $i^{\text{th}}$  node is *participating*<sup>(a)</sup> **then**
  - 9:       Calculate  $k_i^{t+1}$  using Algorithm 2, 3, 4 or 5
  - 10:        $h_i^{t+1} = h_i^t + \frac{1}{p_a} k_i^{t+1}$
  - 11:        $m_i^{t+1} = C_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right)$
  - 12:        $g_i^{t+1} = g_i^t + m_i^{t+1}$
  - 13:       Send  $m_i^{t+1}$  to the server
  - 14:     **else**
  - 15:        $h_{ij}^{t+1} = h_{ij}^t$  (only in **DASHA-PP-FINITE-MVR**)
  - 16:        $h_i^{t+1} = h_i^t, \quad g_i^{t+1} = g_i^t, \quad m_i^{t+1} = 0$
  - 17:     **end if**
  - 18:   **end for**
  - 19:    $g^{t+1} = g^t + \frac{1}{n} \sum_{i=1}^n m_i^{t+1}$
  - 20: **end for**
  - 21: **Output:**  $\hat{x}^T$  chosen uniformly at random from  $\{x^t\}_{k=0}^{T-1}$
- (a): For the formal description see Section 2.2.
- 

---

**Algorithm 2** Calculate  $k_i^{t+1}$  for **DASHA-PP** in the gradient setting. See line 9 in Alg. 1

---

- 1:  $k_i^{t+1} = \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))$
- 

---

**Algorithm 3** Calculate  $k_i^{t+1}$  for **DASHA-PP-PAGE** in the finite-sum setting. See line 9 in Alg. 1

---

- 1: Generate a random set  $I_i^t$  of size  $B$  from  $[m]$  *with replacement*
  - 2:  $k_i^{t+1} = \begin{cases} \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)), \\ \text{with probability } p_{\text{page}} \text{ on all } \textit{participating} \text{ nodes,} \\ \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)), \\ \text{with probability } 1 - p_{\text{page}} \text{ on all } \textit{participating} \text{ nodes} \end{cases}$
- 

---

**Algorithm 4** Calc.  $k_i^{t+1}$  for **DASHA-PP-FINITE-MVR** in the finite-sum setting. See line 9 in Alg. 1

---

- 1: Generate a random set  $I_i^t$  of size  $B$  from  $[m]$  *without replacement*
  - 2:  $k_{ij}^{t+1} = \begin{cases} \frac{m}{B} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))), & j \in I_i^t, \\ 0, & j \notin I_i^t \end{cases}$
  - 3:  $h_{ij}^{t+1} = h_{ij}^t + \frac{1}{p_a} k_{ij}^{t+1}$
  - 4:  $k_i^{t+1} = \frac{1}{m} \sum_{j=1}^m k_{ij}^{t+1}$
- 

---

**Algorithm 5** Calculate  $k_i^{t+1}$  for **DASHA-PP-MVR** in the stochastic setting. See line 9 in Alg. 1

---

- 1: Generate i.i.d. samples  $\{\xi_{ij}^{t+1}\}_{j=1}^B$  of size  $B$  from  $\mathcal{D}_i$ .
  - 2:  $k_i^{t+1} = \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1}) - b \left( h_i^t - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1}) \right)$
-



156 The partial participation mechanism (independent participation from Section 2.2) can be easily  
 157 implemented here if we temporally redefine the compressor and use another one instead:

$$\mathcal{C}_i^p := \begin{cases} \frac{1}{p} \mathcal{C}_i, & \text{with pr. } p_a, \\ 0, & \text{with pr. } 1 - p_a. \end{cases} \Rightarrow g_i^{t+1} = \begin{cases} g_i^t + \frac{1}{p_a} \mathcal{C}_i (\nabla f_i(x^{t+1}) - (1-a)\nabla f_i(x^t) - a g_i^t), & p_a \\ g_i^t, & \text{with pr. } 1 - p_a. \end{cases}$$

158 With probability  $1 - p$ , a node does not update  $g_i$  and does not send anything to the server. The main  
 159 observation is that we can do this trick since  $g_i^{t+1}$  depends only on the vectors  $x^{t+1}$ ,  $x^t$ , and  $g_i^t$ .

160 However, we focus our attention on partial participation *in the finite-sum and stochastic settings*.  
 161 Consider the nodes' steps in **DASHA-MVR** (see Algorithm 7) that is designed for the stochastic setting:

$$h_i^{t+1} = \nabla f_i(x^{t+1}; \xi_i^{t+1}) + (1-b)(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})), \quad (6)$$

$$g_i^{t+1} = g_i^t + \mathcal{C}_i (h_i^{t+1} - h_i^t - a(g_i^t - h_i^t)). \quad (7)$$

162 Even if we use the same trick for (7), we still have to update (6) in every iteration of the algorithm  
 163 since  $g_i^{t+1}$  additionally depends on  $h_i^{t+1}$  and  $h_i^t$ . In other words, if a node does not update  $g_i$  and  
 164 does not send anything to the server, it still has to update  $h_i$ , what is impossible without the points  
 165  $x^{t+1}$  and  $x^t$ . One of the main challenges was to “guess” how to generalize (6) and (7) to the partial  
 166 participation setting. We now provide a solution (**DASHA-PP-MVR** with the batch size  $B = 1$ ):

$$\begin{aligned} h_i^{t+1} &= h_i^t + \frac{1}{p_a} k_i^{t+1}, \quad k_i^{t+1} = \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - b(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})), \\ g_i^{t+1} &= g_i^t + \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \text{ with } p_a, \text{ and } h_i^{t+1} = h_i^t, g_i^{t+1} = g_i^t \text{ with } 1 - p_a. \end{aligned} \quad (8)$$

167 Now both control variables  $g_i^t$  and  $h_i^t$  do not change with the probability  $1 - p_a$ . However, when the  
 168  $i^{\text{th}}$  node participates, the update rule of  $g_i^{t+1}$  and  $h_i^{t+1}$  was modified to make the proof work. When  
 169  $p_a = 1$  (no partial participation), the updates rule from (8) reduce to (6) and (7).

170 The theoretical analysis of the new algorithm became more complicated: unlike (6) and (7), the  
 171 control variables  $h_i^{t+1}$  and  $g_i^{t+1}$  in (8) (see also main Algorithm 1) are coupled by the randomness  
 172 from the partial participation. Going deeper into details, one can compare Lemma I.2 from (Tyurin  
 173 and Richtárik, 2023) and Lemma 5. The former lemma does not use the knowledge about the  
 174 update rule of  $h_i^{t+1}$ , works with one expectation  $\mathbb{E}_C[\cdot]$ , uses only (4), (12), and (13). The latter  
 175 lemma additionally explicitly uses the structure of the update rule of  $h_i^{t+1}$ , surgically copes with  
 176 the expectations  $\mathbb{E}_C[\cdot]$  and  $\mathbb{E}_{p_a}[\cdot]$  (for instance, it is not trivial in each order one should apply the  
 177 expectations), and uses the sampling lemma (Lemma 1). The same reasoning applies to other part of  
 178 the analysis.

179 At the first reading of the proofs, we suggest the reader follow the proof of Theorem 2 in the gradient  
 180 setting, which takes a small part of the paper. Although the appendix seems to be dense and large, the  
 181 size is justified by the fact that we consider four different settings and PL-condition ( $4 \times 2$  tracks  
 182 of proofs. The theory is designed so that the proofs do not repeat steps of each other and use one  
 183 framework).

## 184 6 Theorems

185 We now present the convergence rates theorems of **DASHA-PP** in different settings. We will compare  
 186 the theorems with the results of the current state-of-the-art methods, **MARINA** and **DASHA**, that work  
 187 in the full participation setting. Suppose that **MARINA** or **DASHA** converges to  $\varepsilon$ -solution after  $T$   
 188 communication rounds. Then, ideally, we would expect the convergence of the new algorithms to  
 189  $\varepsilon$ -solution after up to  $T/p_a$  communication rounds due to the partial participation constraints<sup>3</sup>. The  
 190 detailed analysis of the algorithms under Polyak-Łojasiewicz condition we provide in Section F. Let  
 191 us define  $\Delta_0 := f(x^0) - f^*$ .

### 192 6.1 Gradient Setting

<sup>3</sup>We check this numerically in Section A.

193 **Theorem 2.** Suppose that Assumptions 1, 2, 3, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_a}{2-p_a}$ ,

$$\gamma \leq \left( L + \left[ \frac{48\omega(2\omega+1)}{np_a^2} + \frac{16}{np_a^2} \left( 1 - \frac{p_{aa}}{p_a} \right) \right]^{1/2} \hat{L} \right)^{-1},$$

194 and  $g_i^0 = h_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP), then  $\mathbb{E} [\|\nabla f(\hat{x}^T)\|^2] \leq \frac{2\Delta_0}{\gamma T}$ .

195 Let us recall the convergence rate of MARINA or DASHA, the number of communication rounds to get  
 196  $\varepsilon$ -solution equals  $\mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{\sqrt{n}} \hat{L} \right] \right)$ , while the rate of DASHA-PP equals  $\mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega+1}{p_a \sqrt{n}} \hat{L} \right] \right)$ .  
 197 Up to Lipschitz constants factors, we get the degeneration up to  $1/p_a$  factor due to the partial  
 198 participation. This is the perfect result since each worker sends useful information only with  
 199 the probability  $p_a$ .

## 200 6.2 Finite-Sum Setting

201 **Theorem 3.** Suppose that Assumptions 1, 2, 3, 4, 7, and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_{page} p_a}{2-p_a}$ ,  
 202 probability  $p_{page} \in (0, 1]$ ,

$$\gamma \leq \left( L + \left[ \frac{48\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{(1-p_{page})L_{\max}^2}{B} \right) + \frac{16}{np_a^2 p_{page}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{page})L_{\max}^2}{B} \right) \right]^{1/2} \right)^{-1}$$

203 and  $g_i^0 = h_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-PAGE) then  $\mathbb{E} [\|\nabla f(\hat{x}^T)\|^2] \leq$   
 204  $\frac{2\Delta_0}{\gamma T}$ .

205 We now choose  $p_{page}$  to balance heavy full gradient and light mini-batch calculations. Let us define  
 206  $\mathbb{1}_{p_a} := \sqrt{1 - \frac{p_{aa}}{p_a}} \in [0, 1]$ . Note that if  $p_a = 1$  then  $p_{aa} = 1$  and  $\mathbb{1}_{p_a} = 0$ .

207 **Corollary 1.** Let the assumptions from Theorem 3 hold and  $p_{page} = B/(m+B)$ . Then DASHA-PP-PAGE  
 208 needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{1}{p_a} \sqrt{\frac{m}{n}} \left( \frac{\mathbb{1}_{p_a} \hat{L}}{\sqrt{B}} + \frac{L_{\max}}{B} \right) \right] \right) \quad (9)$$

209 communication rounds to get an  $\varepsilon$ -solution and the expected number of gradient calculations per  
 210 node equals  $\mathcal{O}(m + BT)$ .

211 The convergence rate the rate of the current state-of-the-art method DASHA-PAGE without partial  
 212 participation equals  $\mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{\sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \sqrt{\frac{m}{n}} \frac{L_{\max}}{B} \right] \right)$ . Let us closer compare it with (9).  
 213 As expected, we see that the second term w.r.t.  $\omega$  degenerates up to  $1/p_a$ . Surprisingly, the third term  
 214 w.r.t.  $\sqrt{m/n}$  can degenerate up to  $\sqrt{B}/p_a$  when  $\hat{L} \approx L_{\max}$ . Hence, in order to keep degeneration up to  
 215  $1/p_a$ , one should take the batch size  $B = \mathcal{O}(L_{\max}^2/\hat{L}^2)$ . This interesting effect we analyze separately  
 216 in Section C. The fact that the degeneration is up to  $1/p_a$  we check numerically in Section A.

217 In the following corollary, we consider RandK compressors<sup>4</sup> (see Definition 5) and show that with  
 218 the particular choice of parameters, up to the Lipschitz constants factors, DASHA-PP-PAGE gets the  
 219 optimal oracle complexity and SOTA communication complexity. Indeed, comparing the following  
 220 result with (Tyurin and Richtárik, 2023)[Corollary 6.6], one can see that we get the degeneration  
 221 up to  $1/p_a$  factor, which is expected in the partial participation setting. Note that the complexities  
 222 improve with the number of workers  $n$ .

223 **Corollary 2.** Suppose that assumptions of Corollary 1 hold,  $B \leq \min \left\{ \frac{1}{p_a} \sqrt{\frac{m}{n}}, \frac{L_{\max}^2}{\frac{1}{p_a} \hat{L}^2} \right\}$ <sup>5</sup>, and we  
 224 use the unbiased compressor RandK with  $K = \Theta(Bd/\sqrt{m})$ . Then the communication complexity of  
 225 Algorithm 1 is  $\mathcal{O} \left( d + \frac{L_{\max} \Delta_0 d}{p_a \varepsilon \sqrt{n}} \right)$ , and the expected number of gradient calculations per node equals  
 226  $\mathcal{O} \left( m + \frac{L_{\max} \Delta_0 \sqrt{m}}{p_a \varepsilon \sqrt{n}} \right)$ .

227 The convergence rate of DASHA-PP-FINITE-MVR is provided in Section E.5.

<sup>4</sup>The choice of the compressor is driven by simplicity, and the following analysis can be used for other unbiased compressors.

<sup>5</sup>If  $\mathbb{1}_{p_a} = 0$ , then  $\frac{L_{\max}^2}{\frac{1}{p_a} \hat{L}^2} = +\infty$



### 6.3 Stochastic Setting

We define  $h^t := \frac{1}{n} \sum_{i=1}^n h_i^t$ .

**Theorem 4.** Suppose that Assumptions 1, 2, 3, 5, 6, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b \in \left(0, \frac{p_a}{2-p_a}\right]$ ,  $\gamma \leq \left(L + \left[\frac{48\omega(2\omega+1)}{np_a^2} \left(\hat{L}^2 + \frac{(1-b)^2 L_\sigma^2}{B}\right) + \frac{12}{np_a b} \left(\left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2 + \frac{(1-b)^2 L_\sigma^2}{B}\right)\right]^{1/2}\right)^{-1}$ , and  $g_i^0 = h_i^0$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-MVR). Then

$$\mathbb{E} \left[ \left\| \nabla f(\hat{x}^T) \right\|^2 \right] \leq \frac{1}{T} \left[ \frac{2\Delta_0}{\gamma} + \frac{2}{b} \left\| h^0 - \nabla f(x^0) \right\|^2 + \left( \frac{32b\omega(2\omega+1)}{np_a^2} + \frac{4 \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a} \right) \left( \frac{1}{n} \sum_{i=1}^n \left\| h_i^0 - \nabla f_i(x^0) \right\|^2 \right) \right] + \left( \frac{48b^2\omega(2\omega+1)}{p_a^2} + \frac{12b}{p_a} \right) \frac{\sigma^2}{nB}.$$

In the next corollary, we choose momentum  $b$  and initialize vectors  $h_i^0$  to get  $\varepsilon$ -solution.

**Corollary 3.** Suppose that assumptions from Theorem 4 hold, momentum  $b = \Theta \left( \min \left\{ \frac{p_a}{\sigma} \sqrt{\frac{n\varepsilon B}{\sigma^2}}, \frac{p_a n \varepsilon B}{\sigma^2} \right\} \right)$ ,  $\frac{\sigma^2}{n\varepsilon B} \geq 1$ , and  $h_i^0 = \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0)$  for all  $i \in [n]$ , and batch size  $B_{\text{init}} = \Theta \left( \frac{\sqrt{p_a B}}{b} \right)$ , then Algorithm 1 (DASHA-PP-MVR) needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon n}} \left( \frac{1 p_a \hat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right] + \frac{\sigma^2}{\sqrt{p_a n \varepsilon B}} \right)$$

communication rounds to get an  $\varepsilon$ -solution and the number of stochastic gradient calculations per node equals  $\mathcal{O}(B_{\text{init}} + BT)$ .

The convergence rate of the DASHA-SYNC-MVR, the state-of-the-art method without partial participation, equals  $\mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{\sqrt{\varepsilon n}} \frac{L_\sigma}{B} \right] + \frac{\sigma^2}{n\varepsilon B} \right)$ . Similar to Section 6.2, we see that in the regimes when  $\hat{L} \approx L_\sigma$  the third term w.r.t.  $1/\varepsilon^{3/2}$  can degenerate up to  $\sqrt{B}/p_a$ . However, if we take  $B = \mathcal{O}(L_\sigma^2/\hat{L}^2)$ , then the degeneration of the third term will be up to  $1/p_a$ . This effect we analyze in Section C. The fact that the degeneration is up to  $1/p_a$  we check numerically in Section A.

In the following corollary, we consider RandK compressors (see Definition 5) and show that with the particular choice of parameters, up to the Lipschitz constants factors, DASHA-PP-MVR gets the optimal oracle complexity and SOTA communication complexity of DASHA-SYNC-MVR method. Indeed, comparing the following result with (Tyurin and Richtárik, 2023)[Corollary 6.9], one can see that we get the degeneration up to  $1/p_a$  factor, which is expected in the partial participation setting. Note that the complexities improve with the number of workers  $n$ .

**Corollary 4.** Suppose that assumptions of Corollary 3 hold, batch size  $B \leq \min \left\{ \frac{\sigma}{p_a \sqrt{\varepsilon n}}, \frac{L_\sigma^2}{1 p_a \hat{L}^2} \right\}$ , we take RandK compressors with  $K = \Theta \left( \frac{B d \sqrt{\varepsilon n}}{\sigma} \right)$ . Then the communication complexity equals  $\mathcal{O} \left( \frac{d\sigma}{\sqrt{p_a} \sqrt{n\varepsilon}} + \frac{L_\sigma \Delta_0 d}{p_a \sqrt{n\varepsilon}} \right)$ , and the expected number of stochastic gradient calculations per node equals  $\mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} + \frac{L_\sigma \Delta_0 \sigma}{p_a \varepsilon^{3/2} n} \right)$ .

We are aware that the initial batch size  $B_{\text{init}}$  can be suboptimal w.r.t.  $\omega$  in DASHA-PP-MVR in some regimes (see also (Tyurin and Richtárik, 2023)). This is a side effect of mixing the variance reduction of stochastic gradients and compression. However, Corollary 4 reveals that we can escape these regimes by choosing the parameter  $K$  of RandK compressors in a particular way. To get the complete picture, we analyze the same phenomenon under PL condition (see Section F) and provide a new method DASHA-PP-SYNC-MVR (see Section G).

## References

Alistarh, D., Grubic, D., Li, J., Tomioka, R., and Vojnovic, M. (2017). QSGD: Communication-efficient SGD via gradient quantization and encoding. In *Advances in Neural Information Processing Systems (NIPS)*, pages 1709–1720.

Arjevani, Y., Carmon, Y., Duchi, J. C., Foster, D. J., Srebro, N., and Woodworth, B. (2019). Lower bounds for non-convex stochastic optimization. *arXiv preprint arXiv:1912.02365*.

Beznosikov, A., Horváth, S., Richtárik, P., and Safaryan, M. (2020). On biased compression for distributed learning. *arXiv preprint arXiv:2002.12410*.

Carmon, Y., Duchi, J. C., Hinder, O., and Sidford, A. (2020). Lower bounds for finding stationary points i. *Mathematical Programming*, 184(1):71–120.

Chang, C.-C. and Lin, C.-J. (2011). LIBSVM: a library for support vector machines. *ACM Transactions on Intelligent Systems and Technology (TIST)*, 2(3):1–27.

Cutkosky, A. and Orabona, F. (2019). Momentum-based variance reduction in non-convex SGD. *arXiv preprint arXiv:1905.10018*.

Fang, C., Li, C. J., Lin, Z., and Zhang, T. (2018). SPIDER: Near-optimal non-convex optimization via stochastic path integrated differential estimator. In *NeurIPS Information Processing Systems*.

Goodfellow, I., Bengio, Y., Courville, A., and Bengio, Y. (2016). *Deep learning*, volume 1. MIT Press.

Gorbunov, E., Burlachenko, K., Li, Z., and Richtárik, P. (2021). MARINA: Faster non-convex distributed learning with compression. In *38th International Conference on Machine Learning*.

Horváth, S., Ho, C.-Y., Horvath, L., Sahu, A. N., Canini, M., and Richtárik, P. (2019a). Natural compression for distributed deep learning. *arXiv preprint arXiv:1905.10988*.

Horváth, S., Kovalev, D., Mishchenko, K., Stich, S., and Richtárik, P. (2019b). Stochastic distributed learning with gradient quantization and variance reduction. *arXiv preprint arXiv:1904.05115*.

Ioffe, S. and Szegedy, C. (2015). Batch normalization: Accelerating deep network training by reducing internal covariate shift. In *International Conference on Machine Learning*, pages 448–456. PMLR.

Kairouz, P., McMahan, H. B., Avent, B., Bellet, A., Bennis, M., Bhagoji, A. N., Bonawitz, K., Charles, Z., Cormode, G., Cummings, R., et al. (2021). Advances and open problems in federated learning. *Foundations and Trends® in Machine Learning*, 14(1–2):1–210.

Karimireddy, S. P., Jaggi, M., Kale, S., Mohri, M., Reddi, S. J., Stich, S. U., and Suresh, A. T. (2020a). Mime: Mimicking centralized stochastic algorithms in federated learning. *arXiv preprint arXiv:2008.03606*.

Karimireddy, S. P., Kale, S., Mohri, M., Reddi, S., Stich, S., and Suresh, A. T. (2020b). Scaffold: Stochastic controlled averaging for federated learning. In *International Conference on Machine Learning*, pages 5132–5143. PMLR.

Konečný, J., McMahan, H. B., Yu, F. X., Richtárik, P., Suresh, A. T., and Bacon, D. (2016). Federated learning: Strategies for improving communication efficiency. *arXiv preprint arXiv:1610.05492*.

Li, T., Sahu, A. K., Zaheer, M., Sanjabi, M., Talwalkar, A., and Smith, V. (2020). Federated optimization in heterogeneous networks. *Proceedings of Machine Learning and Systems*, 2:429–450.

Li, Z., Bao, H., Zhang, X., and Richtárik, P. (2021a). PAGE: A simple and optimal probabilistic gradient estimator for nonconvex optimization. In *International Conference on Machine Learning*, pages 6286–6295. PMLR.

Li, Z., Hanzely, S., and Richtárik, P. (2021b). ZeroSARAH: Efficient nonconvex finite-sum optimization with zero full gradient computation. *arXiv preprint arXiv:2103.01447*.

Lin, Y., Han, S., Mao, H., Wang, Y., and Dally, W. J. (2017). Deep gradient compression: Reducing the communication bandwidth for distributed training. *arXiv preprint arXiv:1712.01887*.

308 McMahan, B., Moore, E., Ramage, D., Hampson, S., and y Arcas, B. A. (2017). Communication-  
309 efficient learning of deep networks from decentralized data. In *Artificial intelligence and statistics*,  
310 pages 1273–1282. PMLR.

311 Mishchenko, K., Gorbunov, E., Takáč, M., and Richtárik, P. (2019). Distributed learning with  
312 compressed gradient differences. *arXiv preprint arXiv:1901.09269*.

313 Narayanan, D., Harlap, A., Phanishayee, A., Seshadri, V., Devanur, N. R., Ganger, G. R., Gibbons,  
314 P. B., and Zaharia, M. (2019). PipeDream: generalized pipeline parallelism for dnn training. In  
315 *Proceedings of the 27th ACM Symposium on Operating Systems Principles*, pages 1–15.

316 Nesterov, Y. (2018). *Lectures on convex optimization*, volume 137. Springer.

317 Nguyen, L., Liu, J., Scheinberg, K., and Takáč, M. (2017). SARAH: A novel method for machine  
318 learning problems using stochastic recursive gradient. In *The 34th International Conference on*  
319 *Machine Learning*.

320 Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein,  
321 N., Antiga, L., et al. (2019). Pytorch: An imperative style, high-performance deep learning library.  
322 In *Advances in Neural Information Processing Systems (NeurIPS)*.

323 Patel, K. K., Wang, L., Woodworth, B., Bullins, B., and Srebro, N. (2022). Towards optimal commu-  
324 nication complexity in distributed non-convex optimization. In *Advances in Neural Information*  
325 *Processing Systems*.

326 Ramaswamy, S., Mathews, R., Rao, K., and Beaufays, F. (2019). Federated learning for emoji  
327 prediction in a mobile keyboard. *arXiv preprint arXiv:1906.04329*.

328 Ramesh, A., Pavlov, M., Goh, G., Gray, S., Voss, C., Radford, A., Chen, M., and Sutskever, I. (2021).  
329 Zero-shot text-to-image generation. *arXiv preprint arXiv:2102.12092*.

330 Reddi, S., Charles, Z., Zaheer, M., Garrett, Z., Rush, K., Konečný, J., Kumar, S., and McMahan,  
331 H. B. (2020). Adaptive federated optimization. *arXiv preprint arXiv:2003.00295*.

332 Richtárik, P., Sokolov, I., and Fatkhullin, I. (2021). EF21: A new, simpler, theoretically better, and  
333 practically faster error feedback. In *Neural Information Processing Systems, 2021*.

334 Richtárik, P., Sokolov, I., Fatkhullin, I., Gasanov, E., Li, Z., and Gorbunov, E. (2022). 3PC: Three  
335 point compressors for communication-efficient distributed training and a better theory for lazy  
336 aggregation. *arXiv preprint arXiv:2202.00998*.

337 Sapio, A., Canini, M., Ho, C.-Y., Nelson, J., Kalnis, P., Kim, C., Krishnamurthy, A., Moshref, M.,  
338 Ports, D. R., and Richtárik, P. (2019). Scaling distributed machine learning with in-network  
339 aggregation. *arXiv preprint arXiv:1903.06701*.

340 Stich, S. U., Cordonnier, J.-B., and Jaggi, M. (2018). Sparsified SGD with memory. *Advances in*  
341 *Neural Information Processing Systems*, 31.

342 Suresh, A. T., Sun, Z., Ro, J. H., and Yu, F. (2022). Correlated quantization for distributed mean  
343 estimation and optimization. *arXiv preprint arXiv:2203.04925*.

344 Szlendak, R., Tyurin, A., and Richtárik, P. (2021). Permutation compressors for provably faster  
345 distributed nonconvex optimization. *arXiv preprint arXiv:2110.03300*.

346 Tyurin, A. and Richtárik, P. (2023). DASHA: Distributed nonconvex optimization with commu-  
347 nication compression and optimal oracle complexity. *International Conference on Learning*  
348 *Representations (ICLR)*.

349 Vogels, T., He, L., Koloskova, A., Karimireddy, S. P., Lin, T., Stich, S. U., and Jaggi, M. (2021).  
350 RelaySum for decentralized deep learning on heterogeneous data. *Advances in Neural Information*  
351 *Processing Systems*, 34.

352 Wangni, J., Wang, J., Liu, J., and Zhang, T. (2018). Gradient sparsification for communication-  
353 efficient distributed optimization. *Advances in Neural Information Processing Systems*, 31.

- 354 Xu, H., Ho, C.-Y., Abdelmoniem, A. M., Dutta, A., Bergou, E. H., Karatsenidis, K., Canini, M.,  
355 and Kalnis, P. (2021). Grace: A compressed communication framework for distributed machine  
356 learning. In *2021 IEEE 41st International Conference on Distributed Computing Systems (ICDCS)*,  
357 pages 561–572. IEEE.
- 358 Zhao, H., Burlachenko, K., Li, Z., and Richtárik, P. (2021a). Faster rates for compressed federated  
359 learning with client-variance reduction. *arXiv preprint arXiv:2112.13097*.
- 360 Zhao, H., Li, Z., and Richtárik, P. (2021b). FedPAGE: A fast local stochastic gradient method for  
361 communication-efficient federated learning. *arXiv preprint arXiv:2108.04755*.

## 362 Contents

363	<b>1 Introduction</b>	<b>1</b>
364	<b>2 Optimization Problem</b>	<b>1</b>
365	2.1 Unbiased Compressors . . . . .	2
366	2.2 Nodes Partial Participation Assumptions . . . . .	3
367	<b>3 Motivation and Related Work</b>	<b>4</b>
368	<b>4 Contributions</b>	<b>5</b>
369	<b>5 Algorithm Description and Main Challenges Towards Partial Participation</b>	<b>5</b>
370	<b>6 Theorems</b>	<b>7</b>
371	6.1 Gradient Setting . . . . .	7
372	6.2 Finite-Sum Setting . . . . .	8
373	6.3 Stochastic Setting . . . . .	9
374	<b>A Numerical Verification of Theoretical Dependencies</b>	<b>15</b>
375	<b>B Original DASHA and DASHA-MVR Methods</b>	<b>16</b>
376	<b>C Problem of Estimating the Mean in the Partial Participation Setting</b>	<b>17</b>
377	<b>D Auxiliary facts</b>	<b>18</b>
378	D.1 Sampling Lemma . . . . .	18
379	D.2 Compressors Facts . . . . .	20
380	<b>E Proofs of Theorems</b>	<b>20</b>
381	E.1 Standard Lemmas in the Nonconvex Setting . . . . .	21
382	E.2 Generic Lemmas . . . . .	22
383	E.3 Proof for DASHA-PP . . . . .	25
384	E.4 Proof for DASHA-PP-PAGE . . . . .	29
385	E.5 Proof for DASHA-PP-FINITE-MVR . . . . .	40
386	E.6 Proof for DASHA-PP-MVR . . . . .	51
387	<b>F Analysis of DASHA-PP under Polyak-Łojasiewicz Condition</b>	<b>64</b>
388	F.1 Gradient Setting . . . . .	64
389	F.2 Finite-Sum Setting . . . . .	64
390	F.3 Stochastic Setting . . . . .	65
391	F.4 Proofs of Theorems . . . . .	66
392	F.4.1 Standard Lemma under Polyak-Łojasiewicz Condition . . . . .	66
393	F.4.2 Generic Lemma . . . . .	66

394	F.4.3 Proof for DASHA-PP under PL-condition . . . . .	68
395	F.4.4 Proof for DASHA-PP-PAGE under PL-condition . . . . .	70
396	F.4.5 Proof for DASHA-PP-MVR under PL-condition . . . . .	75
397	<b>G Description of DASHA-PP-SYNC-MVR</b>	<b>82</b>
398	G.1 Proof for DASHA-PP-SYNC-MVR . . . . .	84



## 399 A Numerical Verification of Theoretical Dependencies



Figure 1: Classification task with the *real-sim* dataset.

400 Our main goal is to verify the dependeces from the theory. We compare **DASHA-PP** with **DASHA**.  
 401 Clearly, **DASHA-PP** can not generally perform better than **DASHA**. In different settings, we verify  
 402 that the bigger  $p_a$ , the closer **DASHA-PP** is to **DASHA**, i.e., **DASHA-PP** converges no slower than  $1/p_a$   
 403 times.

In all experiments, we take the *real-sim* dataset with dimension  $d = 20,958$  and the number of samples equals 72,309 from LIBSVM datasets (Chang and Lin, 2011) (under the 3-clause BSD license), and randomly split the dataset between  $n = 100$  nodes equally, ignoring residual samples. In the finite-sum setting, we solve a classification problem with functions

$$f_i(x) := \frac{1}{m} \sum_{j=1}^m \left( 1 - \frac{1}{1 + \exp(y_{ij} a_{ij}^\top x)} \right)^2,$$

404 where  $a_{ij} \in \mathbb{R}^d$  is the feature vector of a sample on the  $i^{\text{th}}$  node,  $y_{ij} \in \{-1, 1\}$  is the corresponding  
 405 label, and  $m$  is the number of samples on the  $i^{\text{th}}$  node for all  $i \in [n]$ . In the stochastic setting, we  
 406 consider functions

$$f_i(x_1, x_2) := \mathbb{E}_{j \sim [m]} \left[ -\log \left( \frac{\exp(a_{ij}^\top x_1 x_2)}{\sum_{y \in \{1, 2\}} \exp(a_{ij}^\top x_y)} \right) + \lambda \sum_{y \in \{1, 2\}} \sum_{k=1}^d \frac{\{x_y\}_k^2}{1 + \{x_y\}_k^2} \right],$$

407 where  $x_1, x_2 \in \mathbb{R}^d$ ,  $\{\cdot\}_k$  is an indexing operation,  $a_{ij} \in \mathbb{R}^d$  is a feature of a sample on the  $i^{\text{th}}$  node,  
 408  $y_{ij} \in \{1, 2\}$  is a corresponding label,  $m$  is the number of samples located on the  $i^{\text{th}}$  node, constant  
 409  $\lambda = 0.001$  for all  $i \in [n]$ .

410 The code was written in Python 3.6.8 using PyTorch 1.9 (Paszke et al., 2019). A distributed  
 411 environment was emulated on a machine with Intel(R) Xeon(R) Gold 6226R CPU @ 2.90GHz and  
 412 64 cores.

413 We use the standard setting in experiments<sup>6</sup> where all parameters except step sizes are taken as  
 414 suggested in theory. Step sizes are finetuned from a set  $\{2^i \mid i \in [-10, 10]\}$ . We emulate the partial  
 415 participation setting using  $s$ -nice sampling with the number of nodes  $n = 100$ . We consider the  
 416  $\text{Rand}K$  compressor and take the batch size  $B = 1$ . We plot the relation between communication  
 417 rounds and values of the norm of gradients at each communication round.

418 In the finite-sum (Figure 1a) and in the stochastic setting (Figure 1b), we see that the bigger probability  
 419  $p_a = s/n$  to 1, the closer **DASHA-PP** to **DASHA**. Moreover, **DASHA-PP** with  $s = 10$  and  $s = 1$   
 420 converges approximately  $\times 10$  ( $= 1/p_a$ ) and  $\times 100$  ( $= 1/p_a$ ) times slower, accordingly. Our theory  
 421 predicts such behavior.

<sup>6</sup>Code: <https://github.com/mysteryresearcher/dasha-partial-participation>

## 422 B Original DASHA and DASHA-MVR Methods

423 To simplify the discussion and explanation from the main part, we present the algorithms from (Tyurin  
424 and Richtárik, 2023)

---

### Algorithm 6 DASHA

---

```

1: Input: starting point  $x^0 \in \mathbb{R}^d$ , stepsize  $\gamma > 0$ , momentum  $a \in (0, 1]$ , number of iterations  $T \geq 1$ 
2: Initialize  $g_i^0 \in \mathbb{R}^d$  on the nodes and  $g^0 = \frac{1}{n} \sum_{i=1}^n g_i^0$  on the server
3: for  $t = 0, 1, \dots, T - 1$  do
4:    $x^{t+1} = x^t - \gamma g^t$ 
5:   Broadcast  $x^{t+1}, x^t$  to all participating(a) nodes
6:   for  $i = 1, \dots, n$  in parallel do
7:      $m_i^{t+1} = \mathcal{C}_i(\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - a(g_i^t - \nabla f_i(x^t)))$ 
8:      $g_i^{t+1} = g_i^t + m_i^{t+1}$ 
9:     Send  $m_i^{t+1}$  to the server
10:  end for
11:   $g^{t+1} = g^t + \frac{1}{n} \sum_{i=1}^n m_i^{t+1}$ 
12: end for
13: Output:  $\hat{x}^T$  chosen uniformly at random from  $\{x^t\}_{k=0}^{T-1}$ 

```

---



---

### Algorithm 7 DASHA-MVR (with batch size $B = 1$ )

---

```

1: Input: starting point  $x^0 \in \mathbb{R}^d$ , stepsize  $\gamma > 0$ , momentums  $a, b \in (0, 1]$ , number of iterations  $T \geq 1$ 
2: Initialize  $g_i^0 \in \mathbb{R}^d$  on the nodes and  $g^0 = \frac{1}{n} \sum_{i=1}^n g_i^0$  on the server
3: for  $t = 0, 1, \dots, T - 1$  do
4:    $x^{t+1} = x^t - \gamma g^t$ 
5:   Broadcast  $x^{t+1}, x^t$  to all participating(a) nodes
6:   for  $i = 1, \dots, n$  in parallel do
7:      $h_i^{t+1} = \nabla f_i(x^{t+1}; \xi_i^{t+1}) + (1 - b)(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})), \quad \xi_i^{t+1} \sim \mathcal{D}_i$ 
8:      $m_i^{t+1} = \mathcal{C}_i(h_i^{t+1} - h_i^t - a(g_i^t - h_i^t))$ 
9:      $g_i^{t+1} = g_i^t + m_i^{t+1}$ 
10:    Send  $m_i^{t+1}$  to the server
11:  end for
12:   $g^{t+1} = g^t + \frac{1}{n} \sum_{i=1}^n m_i^{t+1}$ 
13: end for
14: Output:  $\hat{x}^T$  chosen uniformly at random from  $\{x^t\}_{k=0}^{T-1}$ 

```

---

## 425 C Problem of Estimating the Mean in the Partial Participation Setting

426 We now provide the example to explain why the only choice of  $B = \mathcal{O}\left(\min\left\{\frac{1}{p_a}\sqrt{\frac{m}{n}}, \frac{L_{\max}^2}{p_a L^2}\right\}\right)$  and  
 427  $B = \mathcal{O}\left(\min\left\{\frac{\sigma}{p_a\sqrt{\varepsilon n}}, \frac{L_{\sigma}^2}{p_a^2 \hat{\mathcal{L}}^2}\right\}\right)$  in **DASHA-PP-PAGE** and **DASHA-PP-MVR**, accordingly, guarantees  
 428 the degeneration up to  $1/p_a$ . This is surprising, because in methods with the variance reduction of  
 429 stochastic gradients (Li et al., 2021a; Tyurin and Richtárik, 2023) we can take the size of batch size  
 430  $B = \mathcal{O}(\sqrt{\frac{m}{n}})$  and  $B = \mathcal{O}\left(\frac{\sigma}{\sqrt{\varepsilon n}}\right)$  and guarantee the optimality. Note that the smaller the batch size  
 431  $B$ , the more the server and the nodes have to communicate to get  $\varepsilon$ -solution.

432 Let us consider the task of estimating the mean of vectors in the distributed setting. Suppose that we  
 433 have  $n$  nodes, and each of them contains  $m$  vectors  $\{x_{ij}\}_{j=1}^m$ , where  $x_{ij} \in \mathbb{R}^d$  for all  $i \in [n], j \in [m]$ .  
 434 First, let us consider that each node samples a mini-batch  $I^i$  of size  $B$  with replacement and sends it  
 435 to the server. Then the server calculates the mean of the mini-batches from nodes. One can easily  
 436 show that the variance of the estimator is

$$\begin{aligned} & \mathbb{E} \left[ \left\| \frac{1}{nB} \sum_{i=1}^n \sum_{j \in I^i} x_{ij} - \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m x_{ij} \right\|^2 \right] \\ &= \frac{1}{nB} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \left\| x_{ij} - \frac{1}{m} \sum_{j=1}^m x_{ij} \right\|^2. \end{aligned} \quad (10)$$

437 Next, we consider the same task in the partial participation setting with  $s$ -nice sampling, i.e., we  
 438 sample a random set  $S \subset [n]$  of  $s \in [n]$  nodes without replacement and receive the mini-batches  
 439 only from the sampled nodes. Such sampling of nodes satisfy Assumption 8 with  $p_a = s/n$  and  
 440  $p_a = s(s-1)/n(n-1)$ . In this case, the variance of the estimator (See Lemma 1 with  $r_i = 0$  and  
 441  $s_i = \sum_{j \in I^i} x_{ij}$ ) is

$$\begin{aligned} & \mathbb{E} \left[ \left\| \frac{1}{sB} \sum_{i \in S} \sum_{j \in I^i} x_{ij} - \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m x_{ij} \right\|^2 \right] \\ &= \underbrace{\frac{1}{sB} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \left\| x_{ij} - \frac{1}{m} \sum_{j=1}^m x_{ij} \right\|^2}_{\mathcal{L}_{\max}^2} \\ &+ \underbrace{\frac{n-s}{s(n-1)} \frac{1}{n} \sum_{i=1}^n \left\| \frac{1}{m} \sum_{j=1}^m x_{ij} - \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m x_{ij} \right\|^2}_{\hat{\mathcal{L}}^2}. \end{aligned} \quad (11)$$

442 Let us assume that  $s \leq n/2$ . Note that (10) scales with any  $B \geq 1$ , while (11) only scales when  
 443  $B = \mathcal{O}(\mathcal{L}_{\max}^2/\hat{\mathcal{L}}^2)$ . In other words, for large enough  $B$ , the variance in (11) does not significantly  
 444 improves with the growth of  $B$  due to the term  $\hat{\mathcal{L}}^2$ . In our proof, due to partial participation, the  
 445 variance from (11) naturally appears, and we get the same effect. As was mentioned in Sections 6.2  
 446 and 6.3, it can be seen in our convergence rate bounds.

## 447 **D Auxiliary facts**

448 We list auxiliary facts that we use in our proofs:

449 1. For all  $x, y \in \mathbb{R}^d$ , we have

$$\|x + y\|^2 \leq 2\|x\|^2 + 2\|y\|^2 \quad (12)$$

450 2. Let us take a *random vector*  $\xi \in \mathbb{R}^d$ , then

$$\mathbb{E} \left[ \|\xi\|^2 \right] = \mathbb{E} \left[ \|\xi - \mathbb{E}[\xi]\|^2 \right] + \|\mathbb{E}[\xi]\|^2. \quad (13)$$

### 451 **D.1 Sampling Lemma**

452 This section provides a lemma that we regularly use in our proofs, and it is useful for samplings that  
453 satisfy Assumption 8.

454 **Lemma 1.** *Suppose that a set  $S$  is a random subset of a set  $[n]$  such that*

455 1.  $\text{Prob}(i \in S) = p_a, \quad \forall i \in [n],$

456 2.  $\text{Prob}(i \in S, j \in S) = p_{aa}, \quad \forall i \neq j \in [n],$

457 3.  $p_{aa} \leq p_a^2,$

458 where  $p_a \in (0, 1]$  and  $p_{aa} \in [0, 1]$ . Let us take random independent vectors  $s_i \in \mathbb{R}^d$  for all  $i \in [n]$ ,  
459 nonrandom vector  $r_i \in \mathbb{R}^d$  for all  $i \in [n]$ , and random vectors

$$v_i = \begin{cases} r_i + \frac{1}{p_a} s_i, & i \in S, \\ r_i, & i \notin S, \end{cases}$$

460 then

$$\begin{aligned} & \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n v_i - \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n v_i \right] \right\|^2 \right] \\ &= \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E} \left[ \|s_i - \mathbb{E}[s_i]\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\mathbb{E}[s_i]\|^2 + \frac{p_{aa} - p_a^2}{p_a^2} \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E}[s_i] \right\|^2 \\ &\leq \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E} \left[ \|s_i - \mathbb{E}[s_i]\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\mathbb{E}[s_i]\|^2. \end{aligned}$$

461 *Proof.* Let us define additional constants  $p_{an}$  and  $p_{nn}$ , such that

462 1.  $\text{Prob}(i \in S, j \notin S) = p_{an}, \quad \forall i \neq j \in [n],$

463 2.  $\text{Prob}(i \notin S, j \notin S) = p_{nn}, \quad \forall i \neq j \in [n].$

464 Note, that

$$p_{an} = p_{aa} - p_a \quad (14)$$

465 and

$$p_{nn} = 1 - p_{aa} - 2p_{an}. \quad (15)$$

466 Using the law of total expectation and

$$\mathbb{E}[v_i] = p_a \left( r_i + \mathbb{E} \left[ \frac{1}{p_a} s_i \right] \right) + (1 - p_a) r_i = r_i + \mathbb{E}[s_i],$$

467 we have

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n v_i - \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n v_i \right] \right\|^2 \right] \\
&= \frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \|v_i - (r_i + \mathbb{E}[s_i])\|^2 \right] \\
&\quad + \frac{1}{n^2} \sum_{i \neq j}^n \mathbb{E} [\langle v_i - (r_i + \mathbb{E}[s_i]), v_j - (r_j + \mathbb{E}[s_j]) \rangle] \\
&= \frac{p_a}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \left\| r_i + \frac{1}{p_a} s_i - (r_i + \mathbb{E}[s_i]) \right\|^2 \right] \\
&\quad + \frac{1-p_a}{n^2} \sum_{i=1}^n \|r_i - (r_i + \mathbb{E}[s_i])\|^2 \\
&\quad + \frac{p_{aa}}{n^2} \sum_{i \neq j}^n \mathbb{E} \left[ \left\langle r_i + \frac{1}{p_a} s_i - (r_i + \mathbb{E}[s_i]), r_j + \frac{1}{p_a} s_j - (r_j + \mathbb{E}[s_j]) \right\rangle \right] \\
&\quad + \frac{2p_{an}}{n^2} \sum_{i \neq j}^n \mathbb{E} \left[ \left\langle r_i + \frac{1}{p_a} s_i - (r_i + \mathbb{E}[s_i]), r_j - (r_j + \mathbb{E}[s_j]) \right\rangle \right] \\
&\quad + \frac{p_{nn}}{n^2} \sum_{i \neq j}^n \langle r_i - (r_i + \mathbb{E}[s_i]), r_j - (r_j + \mathbb{E}[s_j]) \rangle.
\end{aligned}$$

468 From the independence of random vectors  $s_i$ , we obtain

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n v_i - \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n v_i \right] \right\|^2 \right] \\
&= \frac{p_a}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \left\| \frac{1}{p_a} s_i - \mathbb{E}[s_i] \right\|^2 \right] \\
&\quad + \frac{1-p_a}{n^2} \sum_{i=1}^n \|\mathbb{E}[s_i]\|^2 \\
&\quad + \frac{p_{aa}(1-p_a)^2}{n^2 p_a^2} \sum_{i \neq j}^n \langle \mathbb{E}[s_i], \mathbb{E}[s_j] \rangle \\
&\quad + \frac{2p_{an}(p_a-1)}{n^2 p_a} \sum_{i \neq j}^n \langle \mathbb{E}[s_i], \mathbb{E}[s_j] \rangle \\
&\quad + \frac{p_{nn}}{n^2} \sum_{i \neq j}^n \langle \mathbb{E}[s_i], \mathbb{E}[s_j] \rangle.
\end{aligned}$$

469 Using (14) and (15), we have

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n v_i - \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n v_i \right] \right\|^2 \right] \\
&= \frac{p_a}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \left\| \frac{1}{p_a} s_i - \mathbb{E}[s_i] \right\|^2 \right] \\
&\quad + \frac{1-p_a}{n^2} \sum_{i=1}^n \|\mathbb{E}[s_i]\|^2
\end{aligned}$$

$$\begin{aligned}
& + \frac{p_{aa} - p_a^2}{n^2 p_a^2} \sum_{i \neq j}^n \langle E[s_i], E[s_j] \rangle \\
& \stackrel{(13)}{=} \frac{1}{n^2 p_a} \sum_{i=1}^n E \left[ \|s_i - E[s_i]\|^2 \right] \\
& + \frac{1 - p_a}{n^2 p_a} \sum_{i=1}^n \|E[s_i]\|^2 \\
& + \frac{p_{aa} - p_a^2}{n^2 p_a^2} \sum_{i \neq j}^n \langle E[s_i], E[s_j] \rangle \\
& = \frac{1}{n^2 p_a} \sum_{i=1}^n E \left[ \|s_i - E[s_i]\|^2 \right] \\
& + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|E[s_i]\|^2 \\
& + \frac{p_{aa} - p_a^2}{p_a^2} \left\| \frac{1}{n} \sum_{i=1}^n E[s_i] \right\|^2.
\end{aligned}$$

470 Finally, using that  $p_{aa} \leq p_a^2$ , we have

$$\begin{aligned}
& E \left[ \left\| \frac{1}{n} \sum_{i=1}^n v_i - E \left[ \frac{1}{n} \sum_{i=1}^n v_i \right] \right\|^2 \right] \\
& \leq \frac{1}{n^2 p_a} \sum_{i=1}^n E \left[ \|s_i - E[s_i]\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|E[s_i]\|^2.
\end{aligned}$$

471

□

## 472 D.2 Compressors Facts

473 We define the *RandK* compressor that chooses without replacement  $K$  coordinates, scales them by a  
474 constant factor to preserve unbiasedness and zero-out other coordinates.

**Definition 5.** Let us take a random subset  $S$  from  $[d]$ ,  $|S| = K$ ,  $K \in [d]$ . We say that a stochastic mapping  $\mathcal{C} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is *RandK* if

$$\mathcal{C}(x) = \frac{d}{K} \sum_{j \in S} x_j e_j,$$

475 where  $\{e_i\}_{i=1}^d$  is the standard unit basis.

476 **Theorem 6.** If  $\mathcal{C}$  is *RandK*, then  $\mathcal{C} \in \mathbb{U} \left( \frac{d}{K} - 1 \right)$ .

477 See the proof in (Beznosikov et al., 2020).

## 478 E Proofs of Theorems

479 There are three different sources of randomness in Algorithm 1: the first one from vectors  $\{k_i^{t+1}\}_{i=1}^n$ ,  
480 the second one from compressors  $\{\mathcal{C}_i\}_{i=1}^n$ , and the third one from availability of nodes. We define  
481  $E_k[\cdot]$ ,  $E_C[\cdot]$  and  $E_{p_a}[\cdot]$  to be conditional expectations w.r.t.  $\{k_i^{t+1}\}_{i=1}^n$ ,  $\{\mathcal{C}_i\}_{i=1}^n$ , and availability,  
482 accordingly, conditioned on all previous randomness. Moreover, we define  $E_{t+1}[\cdot]$  to be a conditional  
483 expectation w.r.t. all randomness in iteration  $t+1$  conditioned on all previous randomness. Note,  
484 that  $E_{t+1}[\cdot] = E_k[E_C[E_{p_a}[\cdot]]]$ .

485 In the case of **DASHA-PP-PAGE**, there are two different sources of randomness from  $\{k_i^{t+1}\}_{i=1}^n$ .  
486 We define  $E_{p_{\text{page}}}[\cdot]$  and  $E_B[\cdot]$  to be conditional expectations w.r.t. the probabilistic switching and  
487 mini-batch indices  $I_i^t$ , accordingly, conditioned on all previous randomness. Note, that  $E_{t+1}[\cdot] =$   
488  $E_B[E_C[E_{p_a}[E_{p_{\text{page}}}[\cdot]]]]$  and  $E_{t+1}[\cdot] = E_B[E_{p_{\text{page}}}[E_C[E_{p_a}[\cdot]]]]$ .



## 489 E.1 Standard Lemmas in the Nonconvex Setting

490 We start the proof of theorems by providing standard lemmas from the nonconvex optimization.

491 **Lemma 2.** Suppose that Assumption 2 holds and let  $x^{t+1} = x^t - \gamma g^t$ . Then for any  $g^t \in \mathbb{R}^d$  and  
 492  $\gamma > 0$ , we have

$$f(x^{t+1}) \leq f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - \nabla f(x^t)\|^2. \quad (16)$$

493 *Proof.* Using  $L$ -smoothness, we have

$$\begin{aligned} f(x^{t+1}) &\leq f(x^t) + \langle \nabla f(x^t), x^{t+1} - x^t \rangle + \frac{L}{2} \|x^{t+1} - x^t\|^2 \\ &= f(x^t) - \gamma \langle \nabla f(x^t), g^t \rangle + \frac{L}{2} \|x^{t+1} - x^t\|^2. \end{aligned}$$

494 Next, due to  $-\langle x, y \rangle = \frac{1}{2} \|x - y\|^2 - \frac{1}{2} \|x\|^2 - \frac{1}{2} \|y\|^2$ , we obtain

$$f(x^{t+1}) \leq f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - \nabla f(x^t)\|^2.$$

495 □

496 **Lemma 3.** Suppose that Assumption 1 holds and

$$\mathbb{E} [f(x^{t+1})] + \gamma \Psi^{t+1} \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] + \gamma \Psi^t + \gamma C, \quad (17)$$

497 where  $\Psi^t$  is a sequence of numbers,  $\Psi^t \geq 0$  for all  $t \in [T]$ , constant  $C \geq 0$ , and constant  $\gamma > 0$ .  
 498 Then

$$\mathbb{E} [\|\nabla f(\hat{x}^T)\|^2] \leq \frac{2\Delta_0}{\gamma T} + \frac{2\Psi^0}{T} + 2C, \quad (18)$$

499 where a point  $\hat{x}^T$  is chosen uniformly from a set of points  $\{x^t\}_{t=0}^{T-1}$ .

500 *Proof.* By unrolling (17) for  $t$  from 0 to  $T - 1$ , we obtain

$$\frac{\gamma}{2} \sum_{t=0}^{T-1} \mathbb{E} [\|\nabla f(x^t)\|^2] + \mathbb{E} [f(x^T)] + \gamma \Psi^T \leq f(x^0) + \gamma \Psi^0 + \gamma TC.$$

501 We subtract  $f^*$ , divide inequality by  $\frac{\gamma T}{2}$ , and take into account that  $f(x) \geq f^*$  for all  $x \in \mathbb{R}$ , and  
 502  $\Psi^t \geq 0$  for all  $t \in [T]$ , to get the following inequality:

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} [\|\nabla f(x^t)\|^2] \leq \frac{2\Delta_0}{\gamma T} + \frac{2\Psi^0}{T} + 2C.$$

503 It is left to consider the choice of a point  $\hat{x}^T$  to complete the proof of the lemma. □

**Lemma 4.** If  $0 < \gamma \leq (L + \sqrt{A})^{-1}$ ,  $L > 0$ , and  $A \geq 0$ , then

$$\frac{1}{2\gamma} - \frac{L}{2} - \frac{\gamma A}{2} \geq 0.$$

504 The lemma can be easily checked with the direct calculation.

## 505 E.2 Generic Lemmas

506 **Lemma 5.** Suppose that Assumptions 7 and 8 hold and let us consider sequences  $g_i^{t+1}$ ,  $h_i^{t+1}$ , and  
 507  $k_i^{t+1}$  from Algorithm 1, then

$$\begin{aligned} & \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \\ & \leq \frac{2\omega}{n^2 p_a} \sum_{i=1}^n \|k_i^{t+1}\|^2 + \frac{a^2((2\omega + 1)p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2, \end{aligned} \quad (19)$$

508 and

$$\begin{aligned} & \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \\ & \leq \frac{2\omega}{p_a} \|k_i^{t+1}\|^2 + \left( \frac{a^2(2\omega + 1 - p_a)}{p_a} + (1-a)^2 \right) \|g_i^t - h_i^t\|^2 \quad \forall i \in [n]. \end{aligned} \quad (20)$$

509 *Proof.* First, we estimate  $\mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right]$ :

$$\begin{aligned} & \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \\ & = \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1} - \mathbb{E}_C [\mathbb{E}_{p_a} [g^{t+1} - h^{t+1}]]\|^2 \right] \right] + \|\mathbb{E}_C [\mathbb{E}_{p_a} [g^{t+1} - h^{t+1}]]\|^2, \end{aligned}$$

510 where we used (13). Due to Assumption 8, we have

$$\begin{aligned} & \mathbb{E}_C [\mathbb{E}_{p_a} [g_i^{t+1}]] \\ & = p_a \mathbb{E}_C \left[ g_i^t + \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right] + (1-p_a) g_i^t \\ & = g_i^t + p_a \mathbb{E}_C \left[ \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right] \\ & = g_i^t + k_i^{t+1} - a (g_i^t - h_i^t), \end{aligned}$$

511 and

$$\mathbb{E}_C [\mathbb{E}_{p_a} [h_i^{t+1}]] = p_a \mathbb{E}_C \left[ h_i^t + \frac{1}{p_a} k_i^{t+1} \right] + (1-p_a) h_i^t = h_i^t + k_i^{t+1}.$$

512 Thus, we can get

$$\begin{aligned} & \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \\ & = \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1} - \mathbb{E}_C [\mathbb{E}_{p_a} [g^{t+1} - h^{t+1}]]\|^2 \right] \right] + (1-a)^2 \|g^t - h^t\|^2. \end{aligned}$$

513 Due to the independence of compressors, we can use Lemma 1 with  $r_i = g_i^t - h_i^t$  and  $s_i =$

514  $p_a \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - k_i^{t+1}$ , and obtain

$$\begin{aligned} & \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \\ & \leq \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_C \left[ \left\| p_a \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - k_i^{t+1} - \mathbb{E}_C \left[ p_a \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - k_i^{t+1} \right] \right\|^2 \right] \\ & \quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \mathbb{E}_C \left[ p_a \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - k_i^{t+1} \right] \right\|^2 \end{aligned}$$

$$\begin{aligned}
& + (1-a)^2 \|g^t - h^t\|^2 \\
& = \frac{p_a}{n^2} \sum_{i=1}^n \mathbb{E}_{\mathcal{C}} \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right\|^2 \right] \\
& \quad + \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2.
\end{aligned}$$

515 From Assumption 7, we have

$$\begin{aligned}
& \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \\
& \leq \frac{\omega p_a}{n^2} \sum_{i=1}^n \left\| \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right\|^2 + \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \\
& = \frac{\omega}{n^2 p_a} \sum_{i=1}^n \|k_i^{t+1} - a (g_i^t - h_i^t)\|^2 + \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \\
& \stackrel{(12)}{\leq} \frac{2\omega}{n^2 p_a} \sum_{i=1}^n \|k_i^{t+1}\|^2 + \frac{a^2 ((2\omega + 1)p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2.
\end{aligned}$$

516 The second inequality can be proved almost in the same way:

$$\begin{aligned}
& \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \\
& = \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g_i^{t+1} - h_i^{t+1} - \mathbb{E}_{\mathcal{C}} [\mathbb{E}_{p_a} [g_i^{t+1} - h_i^{t+1}]]\|^2 \right] \right] + \|\mathbb{E}_{\mathcal{C}} [\mathbb{E}_{p_a} [g_i^{t+1} - h_i^{t+1}]]\|^2 \\
& = \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g_i^{t+1} - h_i^{t+1} - g_i^t + a (g_i^t - h_i^t) + h_i^t\|^2 \right] \right] + (1-a)^2 \|g_i^t - h_i^t\|^2 \\
& = p_a \mathbb{E}_{\mathcal{C}} \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \frac{1}{p_a} k_i^{t+1} + a (g_i^t - h_i^t) \right\|^2 \right] \\
& \quad + a^2 (1-p_a) \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g_i^t - h_i^t\|^2 \\
& \stackrel{(13)}{=} p_a \mathbb{E}_{\mathcal{C}} \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right\|^2 \right] \\
& \quad + a^2 \frac{(1-p_a)^2}{p_a} \|g_i^t - h_i^t\|^2 \\
& \quad + a^2 (1-p_a) \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g_i^t - h_i^t\|^2 \\
& \leq \frac{\omega}{p_a} \|k_i^{t+1} - a (g_i^t - h_i^t)\|^2 \\
& \quad + \frac{a^2 (1-p_a)}{p_a} \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g_i^t - h_i^t\|^2 \\
& \stackrel{(12)}{\leq} \frac{2\omega}{p_a} \|k_i^{t+1}\|^2 + \frac{a^2 (2\omega + 1 - p_a)}{p_a} \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g_i^t - h_i^t\|^2.
\end{aligned}$$

517

□

518 **Lemma 6.** Suppose that Assumptions 2, 7, and 8 hold and let us take  $a = \frac{p_a}{2\omega+1}$ , then

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{n p_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& \quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{n p_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] + \frac{4\gamma\omega(2\omega+1)}{n p_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

519 *Proof.* Due to Lemma 2 and the update step from Line 4 in Algorithm 1, we have

$$\begin{aligned}
& \mathbb{E}_{t+1} [f(x^{t+1})] \\
& \leq \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - \nabla f(x^t)\|^2 \right] \\
& = \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - h^t + h^t - \nabla f(x^t)\|^2 \right] \\
& \stackrel{(13)}{\leq} \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \left( \|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2 \right) \right].
\end{aligned}$$

520 Let us fix some constants  $\kappa, \eta \in [0, \infty)$  that we will define later. Combining the last inequality,  
521 bounds (19), (20) and using the law of total expectation, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] \\
& + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& = \mathbb{E} [\mathbb{E}_{t+1} [f(x^{t+1})]] \\
& + \kappa \mathbb{E} [\mathbb{E}_C [\mathbb{E}_{p_a} [\|g^{t+1} - h^{t+1}\|^2]]] + \eta \mathbb{E} \left[ \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \left( \|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2 \right) \right] \\
& + \kappa \mathbb{E} \left[ \frac{2\omega}{n^2 p_a} \sum_{i=1}^n \|k_i^{t+1}\|^2 + \frac{a^2((2\omega+1)p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \right] \\
& + \eta \mathbb{E} \left[ \frac{2\omega}{n p_a} \sum_{i=1}^n \|k_i^{t+1}\|^2 + \left( \frac{a^2(2\omega+1-p_a)}{p_a} + (1-a)^2 \right) \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& = \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( \gamma + \kappa(1-a)^2 \right) \mathbb{E} [\|g^t - h^t\|^2] \\
& + \left( \frac{\kappa a^2((2\omega+1)p_a - p_{aa})}{n p_a^2} + \eta \left( \frac{a^2(2\omega+1-p_a)}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left( \frac{2\kappa\omega}{n p_a} + \frac{2\eta\omega}{p_a} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

522 Now, by taking  $\kappa = \frac{\gamma}{a}$ , we can see that  $\gamma + \kappa(1-a)^2 \leq \kappa$ , and thus

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] \\
& + \frac{\gamma}{a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma}{a} \mathbb{E} [\|g^t - h^t\|^2] \\
& + \left( \frac{\gamma a((2\omega+1)p_a - p_{aa})}{n p_a^2} + \eta \left( \frac{a^2(2\omega+1-p_a)}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left( \frac{2\gamma\omega}{a n p_a} + \frac{2\eta\omega}{p_a} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

523 Next, by taking  $\eta = \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2}$  and considering the choice of  $a$ , one can show that  
 524  $\left( \frac{\gamma a((2\omega+1)p_a - p_{aa})}{np_a^2} + \eta \left( \frac{a^2(2\omega+1-p_a)}{p_a} + (1-a)^2 \right) \right) \leq \eta$ . Thus

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & + \left( \frac{2\gamma(2\omega+1)\omega}{np_a^2} + \frac{2\gamma((2\omega+1)p_a - p_{aa})\omega}{np_a^3} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right]. \end{aligned}$$

525 Considering that  $p_{aa} \geq 0$ , we can simplify the last term and get

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & + \frac{4\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right]. \end{aligned}$$

526 □

### 527 **E.3 Proof for DASHA-PP**

528 **Lemma 7.** Suppose that Assumptions 3 and 8 hold. For  $h_i^{t+1}$  and  $k_i^{t+1}$  from Algorithm 1 (DASHA-PP)  
 529 we have

1.

$$\begin{aligned} & \mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] \\ & \leq \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \|x^{t+1} - x^t\|^2 + \frac{2b^2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

2.

$$\begin{aligned} & \mathbb{E}_{p_a} [\|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2] \\ & \leq \frac{2(1-p_a)}{p_a} L_i^2 \|x^{t+1} - x^t\|^2 + \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n]. \end{aligned}$$

3.

$$\|k_i^{t+1}\|^2 \leq 2L_i^2 \|x^{t+1} - x^t\|^2 + 2b^2 \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n].$$

530 *Proof.* First, let us proof the bound for  $\mathbb{E}_k [\mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2]]$ :

$$\mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2]$$

$$= \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \mathbb{E}_{p_a} [h^{t+1}] \right\|^2 \right] + \left\| \mathbb{E}_{p_a} [h^{t+1}] - \nabla f(x^{t+1}) \right\|^2.$$

531 Using

$$\mathbb{E}_{p_a} [h_i^{t+1}] = h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))$$

532 and (13), we have

$$\begin{aligned} & \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \\ &= \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \mathbb{E}_{p_a} [h^{t+1}] \right\|^2 \right] + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2. \end{aligned}$$

533 We can use Lemma 1 with  $r_i = h_i^t$  and  $s_i = k_i^{t+1}$  to obtain

$$\begin{aligned} & \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \\ & \leq \frac{1}{n^2 p_a} \sum_{i=1}^n \left\| k_i^{t+1} - k_i^{t+1} \right\|^2 + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| k_i^{t+1} \right\|^2 + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\ & = \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\ & \stackrel{(12)}{\leq} \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \frac{2b^2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\ & \leq \frac{2(p_a - p_{aa}) \hat{L}^2}{n p_a^2} \left\| x^{t+1} - x^t \right\|^2 + \frac{2b^2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2. \end{aligned}$$

534 In the last in inequality, we used Assumption 3. Now, we prove the second inequality:

$$\begin{aligned} & \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \\ &= \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \mathbb{E}_{p_a} [h_i^{t+1}] \right\|^2 \right] + \left\| \mathbb{E}_{p_a} [h_i^{t+1}] - \nabla f_i(x^{t+1}) \right\|^2 \\ &= \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\ &= \frac{(1-p_a)^2}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \quad + (1-p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\ &= \frac{(1-p_a)}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\ & \leq \frac{2(1-p_a)}{p_a} L_i^2 \left\| x^{t+1} - x^t \right\|^2 + \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \left\| h_i^t - \nabla f_i(x^t) \right\|^2. \end{aligned}$$

535 Finally, the third inequality of the theorem follows from (12) and Assumption 3.  $\square$

536 **Theorem 2.** Suppose that Assumptions 1, 2, 3, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_a}{2-p_a}$ ,

$$\gamma \leq \left( L + \left[ \frac{48\omega(2\omega+1)}{n p_a^2} + \frac{16}{n p_a^2} \left( 1 - \frac{p_{aa}}{p_a} \right) \right]^{1/2} \hat{L} \right)^{-1},$$

537 and  $g_i^0 = h_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP), then  $\mathbb{E} \left[ \left\| \nabla f(\hat{x}^T) \right\|^2 \right] \leq \frac{2\Delta_0}{\gamma T}$ .

538 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 6, Lemma 7,  
539 and the law of total expectation, we obtain

$$\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \left\| g^{t+1} - h^{t+1} \right\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{n p_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \left\| g_i^{t+1} - h_i^{t+1} \right\|^2 \right]$$



$$\begin{aligned}
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& = \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \quad + \nu \mathbb{E} \left[ \mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] \right] + \rho \mathbb{E} \left[ \mathbb{E}_{p_a} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& \quad + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& \quad + \frac{4\gamma\omega(2\omega + 1)}{np_a^2} \mathbb{E} \left[ 2\hat{L}^2 \|x^{t+1} - x^t\|^2 + 2b^2 \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& \quad + \nu \mathbb{E} \left[ \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \|x^{t+1} - x^t\|^2 + \frac{2b^2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \right] \\
& \quad + \rho \mathbb{E} \left[ \frac{2(1-p_a)\hat{L}^2}{p_a} \|x^{t+1} - x^t\|^2 + \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

540 After rearranging the terms, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& \quad + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& \quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma\omega(2\omega + 1)\hat{L}^2}{np_a^2} - \nu \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} - \rho \frac{2(1-p_a)\hat{L}^2}{p_a} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& \quad + (\gamma + \nu(1-b)^2) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& \quad + \left( \frac{8b^2\gamma\omega(2\omega + 1)}{np_a^2} + \nu \frac{2b^2(p_a - p_{aa})}{np_a^2} + \rho \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

541 By taking  $\nu = \frac{\gamma}{b}$ , one can show that  $(\gamma + \nu(1-b)^2) \leq \nu$ , and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \quad + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& \quad + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma\omega(2\omega+1)\widehat{L}^2}{np_a^2} - \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} - \rho \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \mathbb{E} \left[ \|x^{t+1} - x^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b(p_a - p_{aa})}{np_a^2} + \rho \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

542 Note that  $b = \frac{p_a}{2-p_a}$ , thus

$$\begin{aligned}
& \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b(p_a - p_{aa})}{np_a^2} + \rho \left( \frac{2b^2(1-p_a)}{p_a} + (1-b)^2 \right) \right) \\
& \leq \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b(p_a - p_{aa})}{np_a^2} + \rho(1-b) \right).
\end{aligned}$$

543 And if we take  $\rho = \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}$ , then

$$\left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b(p_a - p_{aa})}{np_a^2} + \rho(1-b) \right) \leq \rho,$$

544 and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} \left[ \|\nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma\omega(2\omega+1)\widehat{L}^2}{np_a^2} - \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} \right. \\
& \quad \left. - \frac{16b\gamma\omega(2\omega+1)(1-p_a)\widehat{L}^2}{np_a^3} - \frac{4\gamma(p_a - p_{aa})(1-p_a)\widehat{L}^2}{np_a^3} \right) \mathbb{E} \left[ \|x^{t+1} - x^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

545 Let us simplify the last inequality. First, note that

$$\frac{16b\gamma\omega(2\omega+1)(1-p_a)\widehat{L}^2}{np_a^3} \leq \frac{16\gamma\omega(2\omega+1)\widehat{L}^2}{np_a^2},$$

546 due to  $b \leq p_a$ . Second,

$$\frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} \leq \frac{4\gamma(p_a - p_{aa})\widehat{L}^2}{np_a^3},$$

547 due to  $b \geq \frac{p_a}{2}$ . All in all, we have

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad - \left(\frac{1}{2\gamma} - \frac{L}{2} - \frac{24\gamma\omega(2\omega+1)\hat{L}^2}{np_a^2} - \frac{8\gamma(p_a - p_{aa})\hat{L}^2}{np_a^3}\right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
&\quad + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(\frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
\end{aligned}$$

548 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
&\mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
&\quad + \frac{\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(\frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
\end{aligned}$$

549 It is left to apply Lemma 3 with

$$\begin{aligned}
\Psi^t &= \frac{(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad + \frac{1}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(\frac{8b\omega(2\omega+1)}{np_a^2} + \frac{2(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right]
\end{aligned}$$

550 to conclude the proof.  $\square$

#### 551 **E.4 Proof for DASHA-PP-PAGE**

552 Let us denote

$$\begin{aligned}
k_{i,1}^{t+1} &:= \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)), \\
k_{i,2}^{t+1} &:= \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)), \\
h_{i,1}^{t+1} &:= \begin{cases} h_i^t + \frac{1}{p_a} k_{i,1}^{t+1}, & i^{\text{th}} \text{ node is participating,} \\ h_i^t, & \text{otherwise,} \end{cases} \\
h_{i,2}^{t+1} &:= \begin{cases} h_i^t + \frac{1}{p_a} k_{i,2}^{t+1}, & i^{\text{th}} \text{ node is participating,} \\ h_i^t, & \text{otherwise,} \end{cases}
\end{aligned}$$

553  $h_1^{t+1} := \frac{1}{n} \sum_{i=1}^n h_{i,1}^{t+1}$ , and  $h_2^{t+1} := \frac{1}{n} \sum_{i=1}^n h_{i,2}^{t+1}$ . Note, that

$$h^{t+1} = \begin{cases} h_1^{t+1}, & \text{with probability } p_{\text{page}}, \\ h_2^{t+1}, & \text{with probability } 1 - p_{\text{page}}. \end{cases}$$

554 **Lemma 8.** Suppose that Assumptions 3, 4, and 8 hold. For  $h_i^{t+1}$  and  $k_i^{t+1}$  from Algorithm 1  
 555 (DASHA-PP-PAGE) we have

1.

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\ & \leq \left( \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} + \frac{(1 - p_{\text{page}})L_{\max}^2}{np_a B} \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{page}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

2.

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\ & \leq \left( \frac{2(1 - p_a)L_i^2}{p_a} + \frac{(1 - p_{\text{page}})L_{\max}^2}{p_a B} \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \left( \frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n]. \end{aligned}$$

3.

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|k_i^{t+1}\|^2 \right] \right] \\ & \leq \left( 2L_i^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n]. \end{aligned}$$

556 *Proof.* First, we prove the first inequality of the theorem:

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\ & = p_{\text{page}} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_1^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_2^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right]. \end{aligned}$$

557 Using

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] = \\ & = p_a h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) + (1 - p_a) h_i^t \\ & = h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)). \end{aligned}$$

558 and

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] = \\ & = p_a h_i^t + \mathbb{E}_B \left[ \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) \right] + (1 - p_a) h_i^t \\ & = h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t), \end{aligned}$$

559 we obtain

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\ & \stackrel{(13)}{=} p_{\text{page}} \mathbb{E}_{p_a} \left[ \|h_1^{t+1} - \mathbb{E}_{p_a} [h_1^{t+1}]\|^2 \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_2^{t+1} - \mathbb{E}_{p_a} [h_2^{t+1}]\|^2 \right] \right] \end{aligned}$$

$$\begin{aligned}
& + p_{\text{page}} \left\| \mathbb{E}_{p_a} [h_1^{t+1}] - \nabla f(x^{t+1}) \right\|^2 + (1 - p_{\text{page}}) \left\| \mathbb{E}_B [\mathbb{E}_{p_a} [h_2^{t+1}]] - \nabla f(x^{t+1}) \right\|^2 \\
& = p_{\text{page}} \mathbb{E}_{p_a} \left[ \left\| h_1^{t+1} - \mathbb{E}_{p_a} [h_1^{t+1}] \right\|^2 \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \left\| h_2^{t+1} - \mathbb{E}_{p_a} [h_2^{t+1}] \right\|^2 \right] \\
& + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \left\| h^t - \nabla f(x^t) \right\|^2. \tag{21}
\end{aligned}$$

560 Next, we consider  $\mathbb{E}_{p_a} \left[ \left\| h_1^{t+1} - \mathbb{E}_{p_a} [h_1^{t+1}] \right\|^2 \right]$ . We can use Lemma 1 with  $r_i = h_i^t$  and  $s_i = k_{i,1}^{t+1}$   
561 to obtain

$$\begin{aligned}
& \mathbb{E}_{p_a} \left[ \left\| h_1^{t+1} - \mathbb{E}_{p_a} [h_1^{t+1}] \right\|^2 \right] \\
& \leq \frac{1}{n^2 p_a} \sum_{i=1}^n \left\| k_{i,1}^{t+1} - k_{i,1}^{t+1} \right\|^2 + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| k_{i,1}^{t+1} \right\|^2 \\
& = \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\
& \stackrel{(12)}{\leq} \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{page}}^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

562 From Assumption 3, we have

$$\begin{aligned}
& \mathbb{E}_{p_a} \left[ \left\| h_1^{t+1} - \mathbb{E}_{p_a} [h_1^{t+1}] \right\|^2 \right] \\
& \leq \frac{2(p_a - p_{aa})\hat{L}^2}{n p_a^2} \left\| x^{t+1} - x^t \right\|^2 + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{page}}^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2. \tag{22}
\end{aligned}$$

563 Now, we prove the bound for  $\mathbb{E}_B \left[ \left\| h_2^{t+1} - \mathbb{E}_{p_a} [h_2^{t+1}] \right\|^2 \right]$ . Considering that mini-  
564 batches in the algorithm are independent, we can use Lemma 1 with  $r_i = h_i^t$  and  $s_i = k_{i,2}^{t+1}$   
565 to obtain

$$\begin{aligned}
& \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_2^{t+1} - \mathbb{E}_{p_a} [h_2^{t+1}] \right\|^2 \right] \right] \\
& \leq \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - \mathbb{E}_{p_a} [k_{i,2}^{t+1}] \right\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \mathbb{E}_B [k_{i,2}^{t+1}] \right\|^2 \\
& = \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
& + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\
& = \frac{1}{n^2 p_a B^2} \sum_{i=1}^n \mathbb{E}_B \left[ \sum_{j \in I_i^t} \left\| (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
& + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\
& = \frac{1}{n^2 p_a B m} \sum_{i=1}^n \sum_{j=1}^m \left\| (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \\
& + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\
& \leq \frac{1}{n^2 p_a B m} \sum_{i=1}^n \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) \right\|^2 + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

566 Next, we use Assumptions 3 and 4 to get

$$\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_2^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_2^{t+1} \right] \right] \right\|^2 \right] \right] \leq \left( \frac{L_{\max}^2}{np_a B} + \frac{(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2. \quad (23)$$

567 Applying (22) and (23) into (21), we get

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \right] \\ & \leq p_{\text{page}} \left( \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \|x^{t+1} - x^t\|^2 + \frac{2(p_a - p_{aa}) b^2}{n^2 p_a^2 p_{\text{page}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right) + \\ & \quad + (1 - p_{\text{page}}) \left( \frac{L_{\max}^2}{np_a B} + \frac{(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \|h^t - \nabla f(x^t)\|^2 \\ & \leq \left( \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} + \frac{(1 - p_{\text{page}}) L_{\max}^2}{np_a B} \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \frac{2(p_a - p_{aa}) b^2}{n^2 p_a^2 p_{\text{page}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

568 The proof of the second inequality almost repeats the previous one:

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \right] \\ & = p_{\text{page}} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,2}^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \\ & \stackrel{(13)}{=} p_{\text{page}} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,2}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] \right\|^2 \right] \right] \\ & \quad + p_{\text{page}} \left\| \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] - \nabla f_i(x^{t+1}) \right\|^2 + (1 - p_{\text{page}}) \left\| \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] - \nabla f_i(x^{t+1}) \right\|^2 \\ & = p_{\text{page}} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \right] + (1 - p_{\text{page}}) \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,2}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] \right\|^2 \right] \right] \\ & \quad + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2. \quad (24) \end{aligned}$$

569 Let us consider  $\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \right]$ :

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \right] \\ & = \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \\ & = p_a \left\| h_i^t + \frac{1}{p_a} k_{i,1}^{t+1} - \left( h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right) \right\|^2 \\ & \quad + (1 - p_a) \left\| h_i^t - \left( h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right) \right\|^2 \\ & = \frac{(1 - p_a)^2}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \quad + (1 - p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & = \frac{1 - p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2. \end{aligned}$$



570 Considering (12) and Assumption 3, we obtain

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,1}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right] \right\|^2 \right] \right] \\ & \leq \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{2(1-p_a)b^2}{p_a p_{\text{page}}^2} \|h_i^t - \nabla f_i(x^t)\|^2. \end{aligned} \quad (25)$$

571 Next, we obtain the bound for  $\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,2}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] \right\|^2 \right] \right]$ :

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{i,2}^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_{i,2}^{t+1} \right] \right] \right\|^2 \right] \right] \\ & = p_a \mathbb{E}_B \left[ \left\| h_i^t + \frac{1}{p_a} k_{i,2}^{t+1} - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\ & \quad + (1-p_a) \mathbb{E}_B \left[ \left\| h_i^t - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\ & = p_a \mathbb{E}_B \left[ \left\| \frac{1}{p_a} k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\ & \quad + (1-p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\ & \stackrel{(13)}{=} \frac{1}{p_a} \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] + \frac{(1-p_a)^2}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\ & \quad + (1-p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\ & = \frac{1}{p_a} \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\ & \leq \frac{1}{p_a} \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] + \frac{(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2, \end{aligned} \quad (26)$$

572 where we used Assumption 3. By plugging (25) and (26) into (24), we get

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \right] \\ & \leq p_{\text{page}} \left( \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{2(1-p_a)b^2}{p_a p_{\text{page}}^2} \|h_i^t - \nabla f_i(x^t)\|^2 \right) \\ & \quad + (1-p_{\text{page}}) \left( \frac{1}{p_a} \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] + \frac{(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 \right) \\ & \quad + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\ & \leq \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{1-p_{\text{page}}}{p_a} \mathbb{E}_B \left[ \left\| k_{i,2}^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\ & \quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2. \end{aligned}$$

573 From the independence of elements in the mini-batch, we obtain

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \right] \\ & \leq \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{1-p_{\text{page}}}{p_a} \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\ & \quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \end{aligned}$$

$$\begin{aligned}
&= \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{1-p_{\text{page}}}{p_a B^2} \mathbb{E}_B \left[ \sum_{j \in I_i^t} \|(\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))\|^2 \right] \\
&\quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\
&= \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{1-p_{\text{page}}}{m p_a B} \sum_{j=1}^m \|(\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))\|^2 \\
&\quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\leq \frac{2(1-p_a)L_i^2}{p_a} \|x^{t+1} - x^t\|^2 + \frac{1-p_{\text{page}}}{m p_a B} \sum_{j=1}^m \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)\|^2 \\
&\quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\leq \left( \frac{2(1-p_a)L_i^2}{p_a} + \frac{(1-p_{\text{page}})L_{\max}^2}{p_a B} \right) \|x^{t+1} - x^t\|^2 \\
&\quad + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2,
\end{aligned}$$

574 where we used Assumption 4. Finally, we prove the last inequality:

$$\begin{aligned}
&\mathbb{E}_B \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|k_i^{t+1}\|^2 \right] \right] \\
&= p_{\text{page}} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-p_{\text{page}}) \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) \right\|^2 \right] \\
&\stackrel{(13)}{=} p_{\text{page}} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{page}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-p_{\text{page}}) \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + (1-p_{\text{page}}) \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
&\stackrel{(12)}{\leq} 2p_{\text{page}} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\quad + (1-p_{\text{page}}) \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + (1-p_{\text{page}}) \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
&\leq 2 \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2
\end{aligned}$$

$$+ (1 - p_{\text{page}}) \mathbb{E}_B \left[ \left\| \frac{1}{B} \sum_{j \in I_i^t} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right].$$

575 Using the independence of elements in the mini-batch, we have

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|k_i^{t+1}\|^2 \right] \right] \\ & \leq 2 \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2 \\ & \quad + \frac{1 - p_{\text{page}}}{B^2} \mathbb{E}_B \left[ \sum_{j \in I_i^t} \|(\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))\|^2 \right] \\ & = 2 \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2 \\ & \quad + \frac{1 - p_{\text{page}}}{Bm} \sum_{j=1}^m \|(\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))\|^2 \\ & \leq 2 \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2 \\ & \quad + \frac{1 - p_{\text{page}}}{Bm} \sum_{j=1}^m \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)\|^2 \end{aligned}$$

576 It is left to consider Assumptions 3 and 4 to get

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|k_i^{t+1}\|^2 \right] \right] \\ & \leq \left( 2L_i^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{p_{\text{page}}} \|h_i^t - \nabla f_i(x^t)\|^2. \end{aligned}$$

577

□

578 **Theorem 3.** Suppose that Assumptions 1, 2, 3, 4, 7, and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_{\text{page}}p_a}{2-p_a}$ ,  
579 probability  $p_{\text{page}} \in (0, 1]$ ,

$$\gamma \leq \left( L + \left[ \frac{48\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) + \frac{16}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right]^{1/2} \right)^{-1}$$

580 and  $g_i^0 = h_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-PAGE) then  $\mathbb{E} [\|\nabla f(\hat{x}^T)\|^2] \leq$   
581  $\frac{2\Delta_0}{\gamma T}$ .

582 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 6, Lemma 8,  
583 and the law of total expectation, we obtain

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & \quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
= & \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_{\text{page}}} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \right] \right] \\
& + \nu \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \right] \\
& + \rho \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{page}}} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \right] \\
\leq & \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \left( 2\hat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{p_{\text{page}}} \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \nu \mathbb{E} \left( \left( \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} + \frac{(1-p_{\text{page}})L_{\max}^2}{np_a B} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{page}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \|h^t - \nabla f(x^t)\|^2 \right) \\
& + \rho \mathbb{E} \left( \left( \frac{2(1-p_a)\hat{L}^2}{p_a} + \frac{(1-p_{\text{page}})L_{\max}^2}{p_a B} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right)
\end{aligned}$$

584 After rearranging the terms, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
\leq & \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} \left[ \|\nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( 2\hat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right)
\end{aligned}$$

$$\begin{aligned}
& -\nu \left( \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} + \frac{(1 - p_{\text{page}})L_{\max}^2}{np_a B} \right) - \rho \left( \frac{2(1 - p_a)\widehat{L}^2}{p_a} + \frac{(1 - p_{\text{page}})L_{\max}^2}{p_a B} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( \gamma + \nu \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \right) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b^2\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{2\nu(p_a - p_{aa})b^2}{np_a^2 p_{\text{page}}} \right. \\
& \quad \left. + \rho \left( \frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Due to  $b = \frac{p_{\text{page}} p_a}{2 - p_a} \leq p_{\text{page}}$ , one can show that  $\left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \leq 1 - b$ . Thus, if we take  $\nu = \frac{\gamma}{b}$ , then

$$\left( \gamma + \nu \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \right) \leq \gamma + \nu(1 - b) = \nu,$$

585 therefore

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega + 1)}{np_a^2} \left( 2\widehat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \frac{\gamma}{b} \left( \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} + \frac{(1 - p_{\text{page}})L_{\max}^2}{np_a B} \right) - \rho \left( \frac{2(1 - p_a)\widehat{L}^2}{p_a} + \frac{(1 - p_{\text{page}})L_{\max}^2}{p_a B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b^2\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{page}}} \right. \\
& \quad \left. + \rho \left( \frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Next, with the choice of  $b = \frac{p_{\text{page}} p_a}{2 - p_a}$ , we ensure that

$$\left( \frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \leq 1 - b.$$

If we take  $\rho = \frac{8b^2\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}}$ , then

$$\left( \frac{8b^2\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{page}}} + \rho \left( \frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1 - p_{\text{page}}) \right) \right) \leq \rho,$$

586 therefore

$$\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right]$$

$$\begin{aligned}
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( 2\hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \right. \\
& \quad \left. - \frac{\gamma}{bnp_a} \left( 2 \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \right. \\
& \quad \left. - \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^3 p_{\text{page}}} + \frac{2\gamma \left( 1 - \frac{p_{\text{aa}}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2(1-p_a) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \right] \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Let us simplify the inequality. First, due to  $b \geq \frac{p_{\text{page}} p_a}{2}$ , we have

$$\frac{\gamma}{bnp_a} \left( 2 \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \leq \frac{4\gamma}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right).$$

587 Second, due to  $b \leq p_a p_{\text{page}}$  and  $p_{\text{aa}} \leq p_a^2$ , we get

$$\begin{aligned}
& \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^3 p_{\text{page}}} + \frac{2\gamma \left( 1 - \frac{p_{\text{aa}}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2(1-p_a) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \\
& \leq \left( \frac{8\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma \left( 1 - \frac{p_{\text{aa}}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2 \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \\
& \leq \frac{16\gamma\omega(2\omega+1)}{np_a^2} \left( \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \\
& \quad + \frac{4\gamma \left( 1 - \frac{p_{\text{aa}}}{p_a} \right)}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \\
& \leq \frac{16\gamma\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right) \\
& \quad + \frac{4\gamma}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{\text{aa}}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\text{max}}^2}{B} \right).
\end{aligned}$$

588 Combining all bounds together, we obtain the following simplified inequality:

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{24\gamma\omega(2\omega+1)}{np_a^2} \left( \widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \frac{8\gamma}{np_a^2 p_{\text{page}}} \left( \left(1 - \frac{p_{\text{aa}}}{p_a}\right) \widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

589 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

590 It is left to apply Lemma 3 with

$$\begin{aligned}
\Psi^t &= \frac{(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{1}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2(p_a - p_{\text{aa}})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]
\end{aligned}$$

591 to conclude the proof.  $\square$

592 **Corollary 1.** Let the assumptions from Theorem 3 hold and  $p_{\text{page}} = B/(m+B)$ . Then DASHA-PP-PAGE  
593 needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{p_a \sqrt{n}} \left( \widehat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{1}{p_a} \sqrt{\frac{m}{n}} \left( \frac{1}{\sqrt{B}} + \frac{L_{\max}}{B} \right) \right] \right) \quad (9)$$

594 communication rounds to get an  $\varepsilon$ -solution and the expected number of gradient calculations per  
595 node equals  $\mathcal{O}(m + BT)$ .

596 *Proof.* In the view of Theorem 3, it is enough to do

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \sqrt{\frac{\omega^2}{np_a^2} \left( \widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right)} + \frac{1}{np_a^2 p_{\text{page}}} \left( \left(1 - \frac{p_{\text{aa}}}{p_a}\right) \widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right] \right)$$

597 steps to get  $\varepsilon$ -solution. Using the choice of  $p_{\text{mega}}$  and the definition of  $\mathbb{I}_{p_a}$ , we can get (9).

598 Note that the expected number of gradients calculations at each communication round equals  $p_{\text{mega}}m +$   
599  $(1 - p_{\text{mega}})B = \frac{2mB}{m+B} \leq 2B$ .  $\square$

600 **Corollary 2.** Suppose that assumptions of Corollary 1 hold,  $B \leq \min \left\{ \frac{1}{p_a} \sqrt{\frac{m}{n}}, \frac{L_{\max}^2}{\frac{1}{p_a} L^2} \right\}$ <sup>7</sup>, and we  
601 use the unbiased compressor RandK with  $K = \Theta(Bd/\sqrt{m})$ . Then the communication complexity of  
602 Algorithm 1 is  $\mathcal{O} \left( d + \frac{L_{\max} \Delta_0 d}{p_a \varepsilon \sqrt{n}} \right)$ , and the expected number of gradient calculations per node equals  
603  $\mathcal{O} \left( m + \frac{L_{\max} \Delta_0 \sqrt{m}}{p_a \varepsilon \sqrt{n}} \right)$ .

<sup>7</sup>If  $\mathbb{I}_{p_a} = 0$ , then  $\frac{L_{\max}^2}{\frac{1}{p_a} L^2} = +\infty$

604 *Proof.* The communication complexity equals

$$\mathcal{O}(d + KT) = \mathcal{O}\left(d + \frac{\Delta_0}{\varepsilon} \left[ KL + K \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + K \frac{1}{p_a} \sqrt{\frac{m}{n}} \left( \frac{\mathbb{1}_{p_a} \hat{L}}{\sqrt{B}} + \frac{L_{\max}}{B} \right) \right]\right).$$

605 Since  $B \leq \frac{L_{\max}^2}{\mathbb{1}_{p_a}^2 \hat{L}^2}$ , we have  $\frac{\mathbb{1}_{p_a} \hat{L}}{\sqrt{B}} + \frac{L_{\max}}{B} \leq \frac{2L_{\max}}{B}$  and

$$\mathcal{O}(d + KT) = \mathcal{O}\left(d + \frac{\Delta_0}{\varepsilon} \left[ KL + K \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + K \frac{1}{p_a} \sqrt{\frac{m}{n}} \frac{L_{\max}}{B} \right]\right).$$

606 Note that  $K = \Theta\left(\frac{Bd}{\sqrt{m}}\right) = \mathcal{O}\left(\frac{d}{p_a \sqrt{n}}\right)$  and  $\omega + 1 = \frac{d}{K}$  due to Theorem 6, thus

$$\begin{aligned} \mathcal{O}(d + KT) &= \mathcal{O}\left(d + \frac{\Delta_0}{\varepsilon} \left[ \frac{d}{p_a \sqrt{n}} L + \frac{d}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{d}{p_a \sqrt{n}} L_{\max} \right]\right) \\ &= \mathcal{O}\left(d + \frac{L_{\max} \Delta_0 d}{p_a \varepsilon \sqrt{n}}\right). \end{aligned}$$

607 Using the same reasoning, the expected number of gradient calculations per node equals

$$\begin{aligned} \mathcal{O}(m + BT) &= \mathcal{O}\left(m + \frac{\Delta_0}{\varepsilon} \left[ BL + B \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + B \frac{1}{p_a} \sqrt{\frac{m}{n}} \left( \frac{\mathbb{1}_{p_a} \hat{L}}{\sqrt{B}} + \frac{L_{\max}}{B} \right) \right]\right) \\ &= \mathcal{O}\left(m + \frac{\Delta_0}{\varepsilon} \left[ BL + B \frac{d}{K p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + B \frac{1}{p_a} \sqrt{\frac{m}{n}} \frac{L_{\max}}{B} \right]\right) \\ &= \mathcal{O}\left(m + \frac{\Delta_0}{\varepsilon} \left[ \frac{1}{p_a} \sqrt{\frac{m}{n}} L + \frac{\sqrt{m}}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{1}{p_a} \sqrt{\frac{m}{n}} L_{\max} \right]\right) \\ &= \mathcal{O}\left(m + \frac{L_{\max} \Delta_0 \sqrt{m}}{p_a \varepsilon \sqrt{n}}\right). \end{aligned}$$

608 □

## 609 E.5 Proof for DASHA-PP-FINITE-MVR

610 **Lemma 9.** Suppose that Assumptions 3, 4, and 8 hold. For  $h_i^{t+1}$ ,  $h_{ij}^{t+1}$  and  $k_i^{t+1}$  from Algorithm 1  
611 (DASHA-PP-FINITE-MVR) we have

1.

$$\begin{aligned} &\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \\ &\leq \left( \frac{2L_{\max}^2}{np_a B} + \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \\ &\quad + \frac{2(p_a - p_{aa}) b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \frac{2b^2}{n^2 p_a B m} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\ &\quad + (1 - b)^2 \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

2.

$$\begin{aligned} &\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \\ &\leq \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1 - p_a) L_i^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \\ &\quad + \frac{2b^2}{p_a B m} \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \left( \frac{2(1 - p_a) b^2}{p_a} + (1 - b)^2 \right) \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n]. \end{aligned}$$



3.

$$\begin{aligned}
& \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1}) \right\|^2 \right] \right] \\
& \leq \frac{2 \left( 1 - \frac{p_a B}{m} \right) L_{\max}^2}{\frac{p_a B}{m}} \left\| x^{t+1} - x^t \right\|^2 \\
& \quad + \left( \frac{2 \left( 1 - \frac{p_a B}{m} \right) b^2}{\frac{p_a B}{m}} + (1-b)^2 \right) \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2, \quad \forall i \in [n], \forall j \in [m].
\end{aligned}$$

4.

$$\begin{aligned}
& \mathbb{E}_B \left[ \left\| k_i^{t+1} \right\|^2 \right] \\
& \leq \left( \frac{2L_{\max}^2}{B} + 2L_i^2 \right) \left\| x^{t+1} - x^t \right\|^2 \\
& \quad + \frac{2b^2}{Bm} \sum_{j=1}^m \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2 + 2b^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2, \quad \forall i \in [n].
\end{aligned}$$

612 *Proof.* We start by proving the first inequality. Note that

$$\begin{aligned}
& \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h_i^{t+1} \right] \right] \\
& = p_a \left( h_i^t + \frac{1}{p_a} \mathbb{E}_B \left[ k_i^{t+1} \right] \right) + (1-p_a) h_i^t \\
& = h_i^t + \frac{1}{m} \sum_{j=1}^m \frac{B}{m} \cdot \frac{m}{B} \left( \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b \left( h_{ij}^t - \nabla f_{ij}(x^t) \right) \right) + \left( 1 - \frac{B}{m} \right) \cdot 0 \\
& = \nabla f_i(x^{t+1}) + (1-b) \left( h_i^t - \nabla f_i(x^t) \right),
\end{aligned}$$

613 thus

$$\begin{aligned}
& \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \\
& \stackrel{(13)}{=} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ h^{t+1} \right] \right] \right\|^2 \right] \right] + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

614 We can use Lemma 1 with  $r_i = h_i^t$  and  $s_i = k_i^{t+1}$  to obtain

$$\begin{aligned}
& \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \\
& \leq \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_B \left[ \left\| k_i^{t+1} - \mathbb{E}_B \left[ k_i^{t+1} \right] \right\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \mathbb{E}_B \left[ k_i^{t+1} \right] \right\|^2 \\
& \quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\
& = \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_B \left[ \left\| \frac{1}{m} \sum_{j=1}^m k_{ij}^{t+1} - \left( \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b \left( h_i^t - \nabla f_i(x^t) \right) \right) \right\|^2 \right] \\
& \quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b \left( h_i^t - \nabla f_i(x^t) \right) \right\|^2 \\
& \quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

615 Next, we again use Lemma 1 with  $r_i = 0$ ,  $s_i = \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b \left( h_{ij}^t - \nabla f_{ij}(x^t) \right)$ ,

616  $p_a = \frac{B}{m}$ , and  $p_{aa} = \frac{B(B-1)}{m(m-1)}$ :

$$\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right]$$

$$\begin{aligned}
&\leq \frac{1}{n^2 p_a} \sum_{i=1}^n \left( \frac{m-B}{Bm(m-1)} \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t)) \right\|^2 \right) \\
&\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\
&\leq \frac{1}{n^2 p_a Bm} \sum_{i=1}^n \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t)) \right\|^2 \\
&\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\
&\stackrel{(12)}{\leq} \frac{2}{n^2 p_a Bm} \sum_{i=1}^n \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) \right\|^2 + \frac{2b^2}{n^2 p_a Bm} \sum_{i=1}^n \sum_{j=1}^m \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2 \\
&\quad + \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

617 Due to Assumptions 3 and 4, we have

$$\begin{aligned}
&\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \\
&\leq \left( \frac{2L_{\max}^2}{np_a B} + \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \right) \left\| x^{t+1} - x^t \right\|^2 \\
&\quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 + \frac{2b^2}{n^2 p_a Bm} \sum_{i=1}^n \sum_{j=1}^m \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

618 Let us get the bound for the second inequality:

$$\begin{aligned}
&\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \\
&\stackrel{(13)}{=} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - (\nabla f_i(x^{t+1}) + (1-b)(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \right] \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&= p_a \mathbb{E}_B \left[ \left\| h_i^t + \frac{1}{p_a} k_i^{t+1} - (\nabla f_i(x^{t+1}) + (1-b)(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + (1-p_a) \left\| h_i^t - (\nabla f_i(x^{t+1}) + (1-b)(h_i^t - \nabla f_i(x^t))) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\stackrel{(13)}{=} \frac{1}{p_a} \mathbb{E}_B \left[ \left\| k_i^{t+1} - \mathbb{E}_B[k_i^{t+1}] \right\|^2 \right] \\
&\quad + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

619 Let us use Lemma 1 with  $r_i = 0$ ,  $s_i = \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))$ ,  $p_a = \frac{B}{m}$ , and

620  $p_{aa} = \frac{B(B-1)}{m(m-1)}$ :

$$\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right]$$

$$\begin{aligned}
&\leq \frac{1}{p_a} \left( \frac{m-B}{Bm(m-1)} \sum_{j=1}^m \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))\|^2 \right) \\
&\quad + \frac{1-p_a}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))\|^2 \\
&\quad + (1-b)^2 \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\leq \frac{1}{p_a B m} \sum_{j=1}^m \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))\|^2 \\
&\quad + \frac{1-p_a}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))\|^2 \\
&\quad + (1-b)^2 \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\stackrel{(12)}{\leq} \frac{2}{p_a B m} \sum_{j=1}^m \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)\|^2 + \frac{2(1-p_a)}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
&\quad + \frac{2b^2}{p_a B m} \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\
&\leq \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)L_i^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \\
&\quad + \frac{2b^2}{p_a B m} \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \|h_i^t - \nabla f_i(x^t)\|^2,
\end{aligned}$$

621 where we used Assumptions 3 and 4. We continue the proof by considering

$$\begin{aligned}
622 \quad &\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \right] : \\
&\mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \right] \\
&\stackrel{(13)}{=} \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h_{ij}^{t+1} - (\nabla f_{ij}(x^{t+1}) + (1-b)(h_{ij}^t - \nabla f_{ij}(x^t)))\|^2 \right] \right] \\
&\quad + (1-b)^2 \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\
&= \frac{p_a B}{m} \mathbb{E}_B \left[ \left\| h_{ij}^t + \frac{m}{B p_a} (\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))) - (\nabla f_{ij}(x^{t+1}) + (1-b)(h_{ij}^t - \nabla f_{ij}(x^t))) \right\|^2 \right] \\
&\quad + \left( 1 - \frac{p_a B}{m} \right) \|h_{ij}^t - (\nabla f_{ij}(x^{t+1}) + (1-b)(h_{ij}^t - \nabla f_{ij}(x^t)))\|^2 \\
&\quad + (1-b)^2 \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\
&= \frac{\left( 1 - \frac{p_a B}{m} \right)^2}{\frac{p_a B}{m}} \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))\|^2 \\
&\quad + \left( 1 - \frac{p_a B}{m} \right) \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))\|^2 \\
&\quad + (1-b)^2 \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\
&= \frac{\left( 1 - \frac{p_a B}{m} \right)}{\frac{p_a B}{m}} \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t))\|^2 \\
&\quad + (1-b)^2 \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\
&\stackrel{(12)}{\leq} \frac{2 \left( 1 - \frac{p_a B}{m} \right)}{\frac{p_a B}{m}} \|\nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t)\|^2 + \left( \frac{2 \left( 1 - \frac{p_a B}{m} \right) b^2}{\frac{p_a B}{m}} + (1-b)^2 \right) \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2.
\end{aligned}$$

623 It is left to consider Assumption 4:

$$\begin{aligned} & \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \left\| h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1}) \right\|^2 \right] \right] \\ & \leq \frac{2 \left( 1 - \frac{p_a B}{m} \right) L_{\max}^2}{\frac{p_a B}{m}} \|x^{t+1} - x^t\|^2 + \left( \frac{2 \left( 1 - \frac{p_a B}{m} \right) b^2}{\frac{p_a B}{m}} + (1-b)^2 \right) \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2. \end{aligned}$$

624 Finally, we obtain the bound for the last inequality of the lemma:

$$\begin{aligned} & \mathbb{E}_B \left[ \left\| k_i^{t+1} \right\|^2 \right] \\ & \stackrel{(13)}{=} \mathbb{E}_B \left[ \left\| k_i^{t+1} - \mathbb{E}_B \left[ k_i^{t+1} \right] \right\|^2 \right] \\ & \quad + \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2. \end{aligned}$$

625 Using Lemma 1, we get

$$\begin{aligned} & \mathbb{E}_B \left[ \left\| k_i^{t+1} \right\|^2 \right] \\ & \leq \frac{m-B}{Bm(m-1)} \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t)) \right\|^2 \\ & \quad + \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \leq \frac{1}{Bm} \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) - b(h_{ij}^t - \nabla f_{ij}(x^t)) \right\|^2 \\ & \quad + \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \stackrel{(12)}{\leq} \frac{2}{Bm} \sum_{j=1}^m \left\| \nabla f_{ij}(x^{t+1}) - \nabla f_{ij}(x^t) \right\|^2 + 2 \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 \\ & \quad + \frac{2b^2}{Bm} \sum_{j=1}^m \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2 + 2b^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\ & \leq \left( \frac{2L_{\max}^2}{B} + 2L_i^2 \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \frac{2b^2}{Bm} \sum_{j=1}^m \left\| h_{ij}^t - \nabla f_{ij}(x^t) \right\|^2 + 2b^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2, \end{aligned}$$

626 where we used Assumptions 3 and 4. □

**Theorem 7.** Suppose that Assumptions 1, 2, 3, 4, 7, and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{\frac{p_a B}{m}}{2 - \frac{p_a B}{m}}$ ,

$$\gamma \leq \left( L + \sqrt{\frac{148\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{L_{\max}^2}{B} \right) + \frac{72m}{np_a^2 B} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{L_{\max}^2}{B} \right)} \right)^{-1},$$

627  $g_i^0 = h_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  and  $h_{ij}^0 = \nabla f_{ij}(x^0)$  for all  $i \in [n], j \in [m]$  in Algorithm 1

628 (DASHA-PP-FINITE-MVR) then  $\mathbb{E} \left[ \left\| \nabla f(\hat{x}^T) \right\|^2 \right] \leq \frac{2\Delta_0}{\gamma^T}$ .

629 *Proof.* Let us fix constants  $\nu, \rho, \delta \in [0, \infty)$  that we will define later. Considering Lemma 6, Lemma 9,  
630 and the law of total expectation, we obtain

$$\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \left\| g^{t+1} - h^{t+1} \right\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \left\| g_i^{t+1} - h_i^{t+1} \right\|^2 \right]$$

$$\begin{aligned}
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
\leq & \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega + 1)}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
= & \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega + 1)}{np_a^2} \mathbb{E} \left[ \mathbb{E}_B \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \right] \\
& + \nu \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\
& + \rho \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\
& + \delta \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \right] \right] \\
\leq & \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega + 1)}{np_a^2} \mathbb{E} \left[ \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{Bmn} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \frac{2b^2}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \nu \mathbb{E} \left( \left( \frac{2L_{\max}^2}{np_a B} + \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \frac{2b^2}{n^2 p_a B m} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \\
& \quad \left. + (1 - b)^2 \|h^t - \nabla f(x^t)\|^2 \right)
\end{aligned}$$

$$\begin{aligned}
& + \rho \mathbb{E} \left( \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \frac{2b^2}{p_a B n m} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right) \\
& + \delta \mathbb{E} \left( \frac{2 \left(1 - \frac{p_a B}{m}\right) L_{\max}^2}{\frac{p_a B}{m}} \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \left( \frac{2 \left(1 - \frac{p_a B}{m}\right) b^2}{\frac{p_a B}{m}} + (1-b)^2 \right) \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right).
\end{aligned}$$

Due to  $b = \frac{\frac{p_a B}{m}}{2 - \frac{p_a B}{m}} \leq \frac{p_a}{2 - p_a}$ , we have

$$\left( \frac{2 \left(1 - \frac{p_a B}{m}\right) b^2}{\frac{p_a B}{m}} + (1-b)^2 \right) \leq 1 - b$$

and

$$\left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \leq 1 - b.$$

631 Moreover, we consider that  $1 - \frac{p_a B}{m} \leq 1$ , therefore

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{Bmn} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + \frac{2b^2}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \nu \mathbb{E} \left( \left( \frac{2L_{\max}^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \frac{2b^2}{n^2 p_a B m} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right. \\
& \quad \left. + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \right) \\
& + \rho \mathbb{E} \left( \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{2b^2}{p_a B n m} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 + (1-b) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \Big) \\
& + \delta \mathbb{E} \left( \frac{2mL_{\max}^2}{p_a B} \|x^{t+1} - x^t\|^2 + (1-b) \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right).
\end{aligned}$$

632 After rearranging the terms, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \right. \\
& \quad \left. - \nu \left( \frac{2L_{\max}^2}{np_a B} + \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\hat{L}^2}{p_a} \right) - \delta \frac{2mL_{\max}^2}{p_a B} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + (\gamma + \nu(1-b)^2) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\nu(p_a - p_{aa})b^2}{np_a^2} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{2\nu b^2}{np_a B} + \frac{2\rho b^2}{p_a B} + \delta(1-b) \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right].
\end{aligned}$$

633 Thus, if we take  $\nu = \frac{\gamma}{b}$ , then  $\gamma + \nu(1-b)^2 \leq \nu$  and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \right.
\end{aligned}$$

$$\begin{aligned}
& - \left( \frac{2\gamma L_{\max}^2}{bn p_a B} + \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bn p_a^2} \right) - \rho \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) - \delta \frac{2mL_{\max}^2}{p_a B} \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{2\gamma b}{np_a B} + \frac{2\rho b^2}{p_a B} + \delta(1-b) \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right].
\end{aligned}$$

Next, if we take  $\rho = \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}$ , then

$$\left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho(1-b) \right) = \rho,$$

634 therefore

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \delta \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \right) \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) \\
& - \left( \frac{2\gamma L_{\max}^2}{bn p_a B} + \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bn p_a^2} \right) - \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \\
& - \delta \frac{2mL_{\max}^2}{p_a B} \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{2\gamma b}{np_a B} + \frac{16b^3\gamma\omega(2\omega+1)}{np_a^3 B} + \frac{4b^2\gamma(p_a - p_{aa})}{nBp_a^3} + \delta(1-b) \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right].
\end{aligned}$$

635 Due to  $b \leq p_a$  and  $\frac{p_a - p_{aa}}{p_a} \leq 1$ , we have

$$\begin{aligned}
& \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{2\gamma b}{np_a B} + \frac{16b^3\gamma\omega(2\omega+1)}{np_a^3 B} + \frac{4b^2\gamma(p_a - p_{aa})}{nBp_a^3} \\
& \leq \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{2\gamma b}{np_a B} + \frac{16b^2\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{4\gamma b}{np_a B}
\end{aligned}$$



$$= \frac{24b^2\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma b}{np_aB}.$$

636 Let us take  $\delta = \frac{24b\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma}{np_aB}$ . Thus

$$\left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2B} + \frac{2\gamma b}{np_aB} + \frac{16b^3\gamma\omega(2\omega+1)}{np_a^3B} + \frac{4b^2\gamma(p_a - p_{aa})}{nBp_a^3} + \delta(1-b) \right) \leq \delta$$

637 and

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \frac{\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma}{np_aB} \right) \mathbb{E}\left[\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\ & - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \right. \\ & \quad \left. - \left( \frac{2\gamma L_{\max}^2}{bnp_aB} + \frac{2\gamma(p_a - p_{aa})\hat{L}^2}{bnp_a^2} \right) - \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \left( \frac{2L_{\max}^2}{p_aB} + \frac{2(1-p_a)\hat{L}^2}{p_a} \right) \right. \\ & \quad \left. - \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma}{np_aB} \right) \frac{2mL_{\max}^2}{p_aB} \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\ & + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] \\ & + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\ & + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma}{np_aB} \right) \mathbb{E}\left[\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2\right]. \end{aligned}$$

638 Let us simplify the term near  $\mathbb{E}[\|x^{t+1} - x^t\|^2]$ . Due to  $b \leq p_a$ ,  $\frac{p_a - p_{aa}}{p_a} \leq 1$ , and  $1 - p_a \leq 1$ , we

639 have

$$\begin{aligned} & \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \\ & + \left( \frac{2\gamma L_{\max}^2}{bnp_aB} + \frac{2\gamma(p_a - p_{aa})\hat{L}^2}{bnp_a^2} \right) \\ & + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \left( \frac{2L_{\max}^2}{p_aB} + \frac{2(1-p_a)\hat{L}^2}{p_a} \right) \\ & + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2B} + \frac{6\gamma}{np_aB} \right) \frac{2mL_{\max}^2}{p_aB} \\ & \leq \frac{12\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\hat{L}^2 \right) \end{aligned}$$

$$\begin{aligned}
& + \left( \frac{6\gamma L_{\max}^2}{bnp_a B} + \frac{6\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} \right) \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \frac{2mL_{\max}^2}{p_a B}
\end{aligned}$$

640 Considering that  $b \leq \frac{p_a B}{m}$  and  $b \geq \frac{p_a B}{2m}$ , we obtain

$$\begin{aligned}
& \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) \\
& + \left( \frac{2\gamma L_{\max}^2}{bnp_a B} + \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} \right) \\
& + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \left( \frac{2L_{\max}^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \frac{2mL_{\max}^2}{p_a B} \\
& \leq \frac{36\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) + \left( \frac{18\gamma L_{\max}^2}{bnp_a B} + \frac{6\gamma(p_a - p_{aa})\widehat{L}^2}{bnp_a^2} \right) \\
& \leq \frac{36\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) + \left( \frac{36m\gamma L_{\max}^2}{np_a^2 B^2} + \frac{12m\gamma(p_a - p_{aa})\widehat{L}^2}{Bnp_a^3} \right).
\end{aligned}$$

641 All in all, we have

$$\begin{aligned}
& \mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \mathbb{E}\left[\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2\right] \\
& \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{36\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2L_{\max}^2}{B} + 2\widehat{L}^2 \right) - \left( \frac{36m\gamma L_{\max}^2}{np_a^2 B^2} + \frac{12m\gamma(p_a - p_{aa})\widehat{L}^2}{Bnp_a^3} \right) \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \mathbb{E}\left[\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2\right].
\end{aligned}$$

642 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right]$$

$$\begin{aligned}
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^{t+1} - \nabla f_{ij}(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{24b\gamma\omega(2\omega+1)}{np_a^2 B} + \frac{6\gamma}{np_a B} \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right].
\end{aligned}$$

643 It is left to apply Lemma 3 with

$$\begin{aligned}
\Psi^t &= \frac{(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{1}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b\omega(2\omega+1)}{np_a^2} + \frac{2(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{24b\omega(2\omega+1)}{np_a^2 B} + \frac{6}{np_a B} \right) \mathbb{E} \left[ \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \|h_{ij}^t - \nabla f_{ij}(x^t)\|^2 \right]
\end{aligned}$$

644 to conclude the proof.  $\square$

## 645 E.6 Proof for DASHA-PP-MVR

646 Let us denote  $\nabla f_i(x^{t+1}; \xi_i^{t+1}) := \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1})$ .

647 **Lemma 10.** Suppose that Assumptions 3, 5, 6 and 8 hold. For  $h_i^{t+1}$  and  $k_i^{t+1}$  from Algorithm 1  
648 (DASHA-PP-MVR) we have

1.

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] \right] \\
& \leq \frac{2b^2\sigma^2}{np_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa}) \widehat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \\
& + \frac{2(p_a - p_{aa}) b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2.
\end{aligned}$$

2.

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} [\|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2] \right] \\
& \leq \frac{2b^2\sigma^2}{p_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) L_i^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \\
& + \left( \frac{2(1-p_a) b^2}{p_a} + (1-b)^2 \right) \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n].
\end{aligned}$$

3.

$$\mathbb{E}_k \left[ \|k_i^{t+1}\|^2 \right] \leq \frac{2b^2\sigma^2}{B} + \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2L_i^2 \right) \|x^{t+1} - x^t\|^2 + 2b^2 \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n].$$

649 *Proof.* First, let us proof the bound for  $\mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right]$ :

$$\begin{aligned} & \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \\ &= \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \mathbb{E}_k [\mathbb{E}_{p_a} [h^{t+1}]]\|^2 \right] \right] + \|\mathbb{E}_k [\mathbb{E}_{p_a} [h^{t+1}]] - \nabla f(x^{t+1})\|^2. \end{aligned}$$

650 Using

$$\mathbb{E}_k [\mathbb{E}_{p_a} [h_i^{t+1}]] = h_i^t + \mathbb{E}_k [k_i^{t+1}] = h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))$$

651 and (13), we have

$$\begin{aligned} & \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \\ &= \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \mathbb{E}_k [\mathbb{E}_{p_a} [h^{t+1}]]\|^2 \right] \right] + (1-b)^2 \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

652 We can use Lemma 1 with  $r_i = h_i^t$  and  $s_i = k_i^{t+1}$  to obtain

$$\begin{aligned} & \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \\ &\leq \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|k_i^{t+1} - \mathbb{E}_k [k_i^{t+1}]\|^2 \right] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\mathbb{E}_k [k_i^{t+1}]\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \\ &= \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - b(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})) \right. \\ &\quad \left. - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)))\|^2 \right] \\ &\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))\|^2 \\ &\quad + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \\ &\stackrel{(12)}{\leq} \frac{2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|b(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1}))\|^2 \right] \\ &\quad + \frac{2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|(1-b)(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)))\|^2 \right] \\ &\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))\|^2 \\ &\quad + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \\ &= \frac{2b^2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1})\|^2 \right] \\ &\quad + \frac{2(1-b)^2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \|\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))\|^2 \right] \\ &\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))\|^2 \\ &\quad + (1-b)^2 \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

$$\begin{aligned}
&= \frac{2b^2}{n^2 p_a B^2} \sum_{i=1}^n \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
&\quad + \frac{2(1-b)^2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

653 In the last equality, we use the independence of elements in the mini-batches. Due to Assumption 5,  
654 we get

$$\begin{aligned}
&\mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \\
&\leq \frac{2b^2 \sigma^2}{n p_a B} \\
&\quad + \frac{2(1-b)^2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2 \\
&\stackrel{(12)}{\leq} \frac{2b^2 \sigma^2}{n p_a B} \\
&\quad + \frac{2(1-b)^2}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2. \\
&= \frac{2b^2 \sigma^2}{n p_a B} \\
&\quad + \frac{2(1-b)^2}{n^2 p_a B^2} \sum_{i=1}^n \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^t; \xi_{ij}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\quad + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2,
\end{aligned}$$

655 where we use the independence of elements in the mini-batches. Using Assumptions 3 and 6, we  
656 obtain

$$\begin{aligned}
&\mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h^{t+1} - \nabla f(x^{t+1}) \right\|^2 \right] \right] \\
&\leq \frac{2b^2 \sigma^2}{n p_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{n p_a B} + \frac{2(p_a - p_{aa}) \widehat{L}^2}{n p_a^2} \right) \left\| x^{t+1} - x^t \right\|^2 \\
&\quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \left\| h_i^t - \nabla f_i(x^t) \right\|^2 + (1-b)^2 \left\| h^t - \nabla f(x^t) \right\|^2.
\end{aligned}$$

657 Now, we prove the second inequality:

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \\
&= \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ h_i^{t+1} \right] \right] \right\|^2 \right] \right] \\
&\quad + \left\| \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ h_i^{t+1} \right] \right] - \nabla f_i(x^{t+1}) \right\|^2 \\
&= \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \right] \\
&\quad + \left\| h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) - \nabla f_i(x^{t+1}) \right\|^2 \\
&= \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \right] \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&= p_a \mathbb{E}_k \left[ \left\| h_i^t + \frac{1}{p_a} k_i^{t+1} - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + (1-p_a) \left\| h_i^t - (h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&= p_a \mathbb{E}_k \left[ \left\| \frac{1}{p_a} k_i^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + (1-p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\stackrel{(13)}{=} \frac{1}{p_a} \mathbb{E}_k \left[ \left\| k_i^{t+1} - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + \frac{(1-p_a)^2}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&= \frac{1}{p_a} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - b(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&= \frac{1}{p_a} \mathbb{E}_k \left[ \left\| b(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1})) + (1-b)(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2 \\
&\stackrel{(12)}{\leq} \frac{2b^2}{p_a} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
&\quad + \frac{2(1-b)^2}{p_a} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

658 Considering the independence of elements in the mini-batch, we obtain

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \\
&= \frac{2b^2}{p_a B^2} \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
&\quad + \frac{2(1-b)^2}{p_a B^2} \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^t; \xi_{ij}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{1-p_a}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\quad + (1-b)^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2. \\
&\stackrel{(12)}{\leq} \frac{2b^2}{p_a B^2} \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
&\quad + \frac{2(1-b)^2}{p_a B^2} \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^t; \xi_{ij}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + \frac{2(1-p_a)}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \left\| h_i^t - \nabla f_i(x^t) \right\|^2
\end{aligned}$$

659 Next, we use Assumptions 3, 6, 5, to get

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \left\| h_i^{t+1} - \nabla f_i(x^{t+1}) \right\|^2 \right] \right] \\
&\leq \frac{2b^2 \sigma^2}{p_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) L_i^2}{p_a} \right) \left\| x^{t+1} - x^t \right\|^2 \\
&\quad + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \left\| h_i^t - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

660 It is left to prove the bound for  $\mathbb{E}_k \left[ \left\| k_i^{t+1} \right\|^2 \right]$ :

$$\begin{aligned}
& \mathbb{E}_k \left[ \left\| k_i^{t+1} \right\|^2 \right] \\
&= \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - b(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})) \right\|^2 \right] \\
&\stackrel{(13)}{=} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - b(h_i^t - \nabla f_i(x^t; \xi_i^{t+1})) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&= \mathbb{E}_k \left[ \left\| b(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1})) + (1-b)(\nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))) \right\|^2 \right] \\
&\quad + \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - b(h_i^t - \nabla f_i(x^t)) \right\|^2 \\
&\stackrel{(12)}{\leq} 2b^2 \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
&\quad + 2(1-b)^2 \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_i^{t+1}) - \nabla f_i(x^t; \xi_i^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
&\quad + 2 \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) \right\|^2 + 2b^2 \left\| h_i^t - \nabla f_i(x^t) \right\|^2.
\end{aligned}$$

661 Using Assumptions 3, 6, 5 and the independence of elements in the mini-batch, we get

$$\mathbb{E}_k \left[ \left\| k_i^{t+1} \right\|^2 \right]$$

$$\leq \frac{2b^2\sigma^2}{B} + \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2L_i^2 \right) \|x^{t+1} - x^t\|^2 + 2b^2 \|h_i^t - \nabla f_i(x^t)\|^2.$$

662

□

663 **Theorem 4.** Suppose that Assumptions 1, 2, 3, 5, 6, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  
 664  $b \in \left(0, \frac{p_a}{2-p_a}\right]$ ,  $\gamma \leq \left( L + \left[ \frac{48\omega(2\omega+1)}{np_a^2} \left( \widehat{L}^2 + \frac{(1-b)^2L_\sigma^2}{B} \right) + \frac{12}{np_ab} \left( \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2 + \frac{(1-b)^2L_\sigma^2}{B} \right) \right]^{1/2} \right)^{-1}$ , and  
 665  $g_i^0 = h_i^0$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-MVR). Then

$$\begin{aligned} \mathbb{E} \left[ \|\nabla f(\widehat{x}^T)\|^2 \right] &\leq \frac{1}{T} \left[ \frac{2\Delta_0}{\gamma} + \frac{2}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{32b\omega(2\omega+1)}{np_a^2} + \frac{4 \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a} \right) \left( \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \right] \\ &+ \left( \frac{48b^2\omega(2\omega+1)}{p_a^2} + \frac{12b}{p_a} \right) \frac{\sigma^2}{nB}. \end{aligned}$$

666 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 6, Lemma 10,  
 667 and the law of total expectation, we obtain

$$\begin{aligned} &\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ &+ \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ &\leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ &+ \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ &+ \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \\ &+ \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ &= \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ &+ \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ &+ \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \mathbb{E}_k \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \right] \\ &+ \nu \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] \right] \right] \\ &+ \rho \mathbb{E} \left[ \mathbb{E}_B \left[ \mathbb{E}_{p_a} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\ &\leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ &+ \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ &+ \frac{4\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \frac{2b^2\sigma^2}{B} + \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 + 2b^2 \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \end{aligned}$$



$$\begin{aligned}
& + \nu \mathbb{E} \left( \frac{2b^2\sigma^2}{np_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \right) \\
& + \rho \mathbb{E} \left( \frac{2b^2\sigma^2}{p_a B} + \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) \hat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right).
\end{aligned}$$

668 After rearranging the terms, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \right. \\
& \quad \left. - \nu \left( \frac{2(1-b)^2 L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) - \rho \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) \hat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + (\gamma + \nu(1-b)^2) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\nu(p_a - p_{aa})b^2}{np_a^2} + \rho \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \nu \frac{2b^2}{np_a} + \rho \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

669 By taking  $\nu = \frac{\gamma}{b}$ , one can show that  $(\gamma + \nu(1-b)^2) \leq \nu$ , and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \right. \\
& \quad \left. - \frac{\gamma}{b} \left( \frac{2(1-b)^2 L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) - \rho \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) \hat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2]
\end{aligned}$$

$$\begin{aligned}
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b}{np_a} + \rho \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

670 Note that  $b \leq \frac{p_a}{2-p_a}$ , thus

$$\begin{aligned}
& \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \right) \\
& \leq \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho(1-b) \right).
\end{aligned}$$

671 And if we take  $\rho = \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2}$ , then

$$\left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})b}{np_a^2} + \rho(1-b) \right) \leq \rho,$$

672 and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \right. \\
& \quad \left. - \frac{\gamma}{np_a b} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^3} + \frac{2\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2} \right) \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2(1-p_a) \hat{L}^2 \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b}{np_a} + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

673 Let us simplify the inequality. First, due to  $b \leq p_a$  and  $(1-p_a) \leq \left( 1 - \frac{p_{aa}}{p_a} \right)$ , we have

$$\begin{aligned}
& \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^3} + \frac{2\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2} \right) \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2(1-p_a) \hat{L}^2 \right) \\
& = \frac{8b\gamma\omega(2\omega+1)}{np_a^3} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2(1-p_a) \hat{L}^2 \right) \\
& \quad + \frac{2\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2(1-p_a) \hat{L}^2 \right)
\end{aligned}$$

$$\leq \frac{8\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \\ + \frac{2\gamma}{np_a b} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 \right),$$

674 therefore

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{12\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \right. \\ & \quad \left. - \frac{3\gamma}{np_a b} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\ & + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\ & + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b}{np_a} + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B} \\ & = \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\ & + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{24\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \hat{L}^2 \right) \right. \\ & \quad \left. - \frac{6\gamma}{np_a b} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\ & + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\ & + \left( \frac{8b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma b}{np_a} + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}. \end{aligned}$$

675 Also, we can simplify the last term:

$$\begin{aligned} & \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \\ & = \frac{16b^3\gamma\omega(2\omega+1)}{np_a^3} + \frac{4b^2\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2} \\ & \leq \frac{16b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{4b\gamma}{np_a}, \end{aligned}$$

676 thus

$$\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right]$$

$$\begin{aligned}
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{24\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
& \quad \left. - \frac{6\gamma}{np_a b} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \left( 1 - \frac{p_{aa}}{p_a} \right) \widehat{L}^2 \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{24b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{6\gamma b}{np_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

677 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{24b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{6\gamma b}{np_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

678 It is left to apply Lemma 3 with

$$\begin{aligned}
\Psi^t &= \frac{(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&+ \frac{1}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{8b\omega(2\omega+1)}{np_a^2} + \frac{2(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]
\end{aligned}$$

679 and  $C = \left( \frac{24b^2\omega(2\omega+1)}{p_a^2} + \frac{6b}{p_a} \right) \frac{\sigma^2}{nB}$  to conclude the proof.  $\square$

680 **Corollary 3.** Suppose that assumptions from Theorem 4 hold, momentum  $b =$   
681  $\Theta \left( \min \left\{ \frac{p_a}{\omega} \sqrt{\frac{n\varepsilon B}{\sigma^2}}, \frac{p_a n \varepsilon B}{\sigma^2} \right\} \right)$ ,  $\frac{\sigma^2}{n\varepsilon B} \geq 1$ , and  $h_i^0 = \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0)$  for all  $i \in [n]$ ,  
682 and batch size  $B_{\text{init}} = \Theta \left( \frac{\sqrt{p_a B}}{b} \right)$ , then Algorithm 1 (DASHA-PP-MVR) needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{p_a \sqrt{n}} \left( \widehat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon n}} \left( \frac{1_{p_a} \widehat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right] + \frac{\sigma^2}{\sqrt{p_a n \varepsilon B}} \right)$$

683 communication rounds to get an  $\varepsilon$ -solution and the number of stochastic gradient calculations per  
684 node equals  $\mathcal{O}(B_{\text{init}} + BT)$ .

685 *Proof.* Using the result from Theorem 4, we have

$$\begin{aligned} & \mathbb{E} \left[ \left\| \nabla f(\hat{x}^T) \right\|^2 \right] \\ & \leq \frac{1}{T} \left[ 2\Delta_0 \left( L + \sqrt{\frac{48\omega(2\omega+1)}{np_a^2}} \left( \hat{L}^2 + \frac{(1-b)^2 L_\sigma^2}{B} \right) + \frac{12}{np_a b} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1-b)^2 L_\sigma^2}{B} \right) \right) \right. \\ & \quad \left. + \frac{2}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{32b\omega(2\omega+1)}{np_a^2} + \frac{4 \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a} \right) \left( \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \right] \\ & \quad + \left( \frac{48b^2\omega(2\omega+1)}{p_a^2} + \frac{12b}{p_a} \right) \frac{\sigma^2}{nB} \end{aligned}$$

686 We choose  $b$  to ensure  $\left( \frac{48b^2\omega(2\omega+1)}{p_a^2} + \frac{12b}{p_a} \right) \frac{\sigma^2}{nB} = \Theta(\varepsilon)$ . Note that  $\frac{1}{b} =$

687  $\Theta \left( \max \left\{ \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{n\varepsilon B}}, \frac{\sigma^2}{p_a n \varepsilon B} \right\} \right) \leq \Theta \left( \max \left\{ \frac{\omega^2}{p_a}, \frac{\sigma^2}{p_a n \varepsilon B} \right\} \right)$ , thus

$$\begin{aligned} & \mathbb{E} \left[ \left\| \nabla f(\hat{x}^T) \right\|^2 \right] \\ & = \mathcal{O} \left( \frac{1}{T} \left[ \Delta_0 \left( L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \right. \right. \\ & \quad \left. \left. + \frac{1}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{b\omega^2}{np_a^2} + \frac{1}{np_a} \right) \left( \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \right] + \varepsilon \right), \end{aligned}$$

688 where  $\mathbb{1}_{p_a} = \sqrt{1 - \frac{p_{aa}}{p_a}}$ . It enough to take the following  $T$  to get  $\varepsilon$ -solution.

$$\begin{aligned} T & = \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \right. \right. \\ & \quad \left. \left. + \frac{1}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{b\omega^2}{np_a^2} + \frac{1}{np_a} \right) \left( \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \right] \right). \end{aligned}$$

689 Let us bound the norms:

$$\begin{aligned} \mathbb{E} \left[ \|h^0 - \nabla f(x^0)\|^2 \right] & = \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0) - \nabla f(x^0) \right\|^2 \right] \\ & = \frac{1}{n^2 B_{\text{init}}^2} \sum_{i=1}^n \sum_{k=1}^{B_{\text{init}}} \mathbb{E} \left[ \left\| \nabla f_i(x^0; \xi_{ik}^0) - \nabla f_i(x^0) \right\|^2 \right] \\ & \leq \frac{\sigma^2}{n B_{\text{init}}}. \end{aligned}$$

690 Using the same reasoning, one can get  $\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[ \|h_i^0 - \nabla f_i(x^0)\|^2 \right] \leq \frac{\sigma^2}{B_{\text{init}}}$ . Combining all inequalities, we have

$$T = \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \right. \right. \\ \left. \left. + \frac{\sigma^2}{bnB_{\text{init}}} + \frac{b\omega^2 \sigma^2}{np_a^2 B_{\text{init}}} + \frac{\sigma^2}{np_a B_{\text{init}}} \right] \right).$$

692 Using the choice of  $B_{\text{init}}$  and  $b$ , we obtain

$$T = \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \right. \right. \\ \left. \left. + \frac{\sigma^2}{\sqrt{p_a} n B} + \frac{b^2 \omega^2 \sigma^2}{np_a^{5/2} B} + \frac{b\sigma^2}{p_a^{3/2} n B} \right] \right) \\ = \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \right. \right. \\ \left. \left. + \frac{\sigma^2}{\sqrt{p_a} n B} + \frac{\varepsilon}{\sqrt{p_a}} \right] \right) \\ = \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right] + \frac{\sigma^2}{\sqrt{p_a} n \varepsilon B} + \frac{1}{\sqrt{p_a}} \right).$$

693 Using  $\frac{\sigma^2}{n\varepsilon B} \geq 1$ , we can conclude the proof of the inequality. The number of stochastic gradients that  
694 each node calculates equals  $B_{\text{init}} + 2BT = \mathcal{O}(B_{\text{init}} + BT)$ .  $\square$

695 **Corollary 4.** Suppose that assumptions of Corollary 3 hold, batch size  $B \leq \min \left\{ \frac{\sigma}{p_a \sqrt{\varepsilon n}}, \frac{L_\sigma^2}{\mathbb{1}_{p_a}^2 \hat{L}^2} \right\}$ ,  
696 we take RandK compressors with  $K = \Theta \left( \frac{Bd\sqrt{\varepsilon n}}{\sigma} \right)$ . Then the communication complexity equals  
697  $\mathcal{O} \left( \frac{d\sigma}{\sqrt{p_a} \sqrt{n\varepsilon}} + \frac{L_\sigma \Delta_0 d}{p_a \sqrt{n\varepsilon}} \right)$ , and the expected number of stochastic gradient calculations per node equals  
698  $\mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a} n \varepsilon} + \frac{L_\sigma \Delta_0 \sigma}{p_a \varepsilon^{3/2} n} \right)$ .

699 *Proof.* The communication complexity equals

$$\mathcal{O}(d + KT) = \mathcal{O} \left( d + \frac{\Delta_0}{\varepsilon} \left[ KL + K \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + K \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right] + K \frac{\sigma^2}{\sqrt{p_a} n \varepsilon B} \right).$$

700 Due to  $B \leq \frac{L_\sigma^2}{\mathbb{1}_{p_a}^2 \hat{L}^2}$ , we have  $\mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \leq \frac{2L_\sigma}{\sqrt{B}}$  and

$$\mathcal{O}(d + KT) = \mathcal{O} \left( d + \frac{\Delta_0}{\varepsilon} \left[ KL + K \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + K \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \frac{L_\sigma}{\sqrt{B}} \right] + K \frac{\sigma^2}{\sqrt{p_a n \varepsilon B}} \right).$$

701 From Theorem 6, we have  $\omega + 1 = \frac{d}{K}$ . Since  $K = \Theta \left( \frac{Bd\sqrt{\varepsilon n}}{\sigma} \right) = \mathcal{O} \left( \frac{d}{p_a \sqrt{n}} \right)$ , the communication  
702 complexity equals

$$\begin{aligned} \mathcal{O}(d + KT) &= \mathcal{O} \left( d + \frac{\Delta_0}{\varepsilon} \left[ \frac{d}{p_a \sqrt{n}} L + \frac{d}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{d}{p_a \sqrt{n}} L_\sigma \right] + \frac{d\sigma}{\sqrt{p_a} \sqrt{n \varepsilon}} \right) \\ &= \mathcal{O} \left( \frac{d\sigma}{\sqrt{p_a} \sqrt{n \varepsilon}} + \frac{L_\sigma \Delta_0 d}{p_a \sqrt{n \varepsilon}} \right) \end{aligned}$$

703 And the expected number of stochastic gradient calculations per node equals

$$\begin{aligned} &\mathcal{O}(B_{\text{init}} + BT) \\ &= \mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} + \frac{B\omega}{\sqrt{p_a}} \sqrt{\frac{\sigma^2}{n \varepsilon B}} + \frac{\Delta_0}{\varepsilon} \left[ BL + B \frac{\omega}{p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + B \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \left( \mathbb{1}_{p_a} \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right] + B \frac{\sigma^2}{\sqrt{p_a n \varepsilon B}} \right) \\ &= \mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} + \frac{Bd}{K \sqrt{p_a}} \sqrt{\frac{\sigma^2}{n \varepsilon B}} + \frac{\Delta_0}{\varepsilon} \left[ BL + B \frac{d}{K p_a \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + B \sqrt{\frac{\sigma^2}{p_a^2 \varepsilon n^2 B}} \frac{L_\sigma}{\sqrt{B}} \right] + \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} \right) \\ &= \mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} + \frac{\sigma^2}{\sqrt{p_a n \varepsilon} \sqrt{B}} + \frac{\Delta_0}{\varepsilon} \left[ \frac{\sigma}{p_a \sqrt{\varepsilon n}} L + \frac{\sigma}{p_a \sqrt{\varepsilon n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon n}} L_\sigma \right] \right) \\ &= \mathcal{O} \left( \frac{\sigma^2}{\sqrt{p_a n \varepsilon}} + \frac{L_\sigma \Delta_0 \sigma}{p_a \varepsilon^{3/2} n} \right). \end{aligned}$$

704

□

## F Analysis of DASHA-PP under Polyak-Łojasiewicz Condition

In this section, we provide the theoretical convergence rates of DASHA-PP under Polyak-Łojasiewicz Condition.

**Assumption 9.** The function  $f$  satisfy (Polyak-Łojasiewicz) PL-condition:

$$\|\nabla f(x)\|^2 \geq 2\mu(f(x) - f^*), \quad \forall x \in \mathbb{R}, \quad (27)$$

where  $f^* = \inf_{x \in \mathbb{R}^d} f(x) > -\infty$ .

Under Polyak-Łojasiewicz condition, a (random) point  $\hat{x}$  is  $\varepsilon$ -solution, if  $\mathbb{E}[f(\hat{x})] - f^* \leq \varepsilon$ .

We now provide the convergence rates of DASHA-PP under PL-condition.

### F.1 Gradient Setting

**Theorem 8.** Suppose that Assumption 1, 2, 3, 7, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_a}{2-p_a}$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega+1)}{np_a^2} + \frac{48}{np_a^2} \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}} \right)^{-1}, \frac{a}{4\mu} \right\},$$

and  $h_i^0 = g_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP), then  $\mathbb{E}[f(x^T)] - f^* \leq (1 - \gamma\mu)^T \Delta_0$ .

Let us provide bounds up to logarithmic factors and use  $\tilde{\mathcal{O}}(\cdot)$  notation. The provided theorem states that to get  $\varepsilon$ -solution DASHA-PP have to run

$$\tilde{\mathcal{O}} \left( \frac{\omega+1}{p_a} + \frac{L}{\mu} + \frac{\omega \hat{L}}{p_a \mu \sqrt{n}} + \frac{\hat{L}}{p_a \mu \sqrt{n}} \right),$$

communication rounds. The method DASHA from (Tyurin and Richtárik, 2023), have to run

$$\tilde{\mathcal{O}} \left( \omega + \frac{L}{\mu} + \frac{\omega \hat{L}}{\mu \sqrt{n}} \right),$$

communication rounds to get  $\varepsilon$ -solution. The difference is the same as in the general nonconvex case (see Section 6.1). Up to Lipschitz constants factors, we get the degeneration up to  $1/p_a$  factor due to the partial participation.

### F.2 Finite-Sum Setting

**Theorem 9.** Suppose that Assumption 1, 2, 3, 7, 4, 8, and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ , probability  $p_{page} = \frac{B}{m+B}$ ,  $b = \frac{p_{page} p_a}{2-p_a}$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{(1-p_{page})L_{\max}^2}{B} \right) + \frac{48}{np_a^2 p_{page}} \left( \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2 + \frac{(1-p_{page})L_{\max}^2}{B} \right)} \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

and  $h_i^0 = g_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-PAGE), then  $\mathbb{E}[f(x^T)] - f^* \leq (1 - \gamma\mu)^T \Delta_0$ .

The provided theorem states that to get  $\varepsilon$ -solution DASHA-PP have to run

$$\tilde{\mathcal{O}} \left( \frac{\omega+1}{p_a} + \frac{m}{p_a B} + \frac{L}{\mu} + \frac{\omega}{p_a \mu \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{\sqrt{m}}{p_a \mu \sqrt{n} B} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) \right),$$

communication rounds. The method DASHA-PAGE from (Tyurin and Richtárik, 2023), have to run

$$\tilde{\mathcal{O}} \left( \omega + \frac{m}{B} + \frac{L}{\mu} + \frac{\omega}{\mu \sqrt{n}} \left( \hat{L} + \frac{L_{\max}}{\sqrt{B}} \right) + \frac{\sqrt{m}}{\mu \sqrt{n} B} \left( \frac{L_{\max}}{\sqrt{B}} \right) \right),$$

communication rounds to get  $\varepsilon$ -solution. We can guarantee the degeneration up to  $1/p_a$  factor due to the partial participation only if  $B = \mathcal{O} \left( \frac{L_{\max}^2}{L^2} \right)$ . The same conclusion we have in Section 6.2.



### 728 F.3 Stochastic Setting

**Theorem 10.** Suppose that Assumption 1, 2, 3, 7, 5, 6, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  
 $b \in \left(0, \frac{p_a}{2-p_a}\right]$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega+1)}{np_a^2} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \hat{L}^2 \right)} + \frac{40}{np_a b} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2 \right) \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

729 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-MVR), then

$$\begin{aligned} & \mathbb{E} [f(x^T) - f^*] \\ & \leq (1 - \gamma\mu)^T \left( \Delta_0 + \frac{2\gamma}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{40\gamma b\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \\ & \quad + \frac{1}{\mu} \left( \frac{100b^2\omega(2\omega+1)}{p_a^2} + \frac{20b}{p_a} \right) \frac{\sigma^2}{nB}. \end{aligned}$$

730 The provided theorems states that to get  $\varepsilon$ -solution DASHA-PP have to run

$$\tilde{\mathcal{O}} \left( \underbrace{\frac{\omega+1}{p_a} + \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n \varepsilon B}}}_{\mathcal{P}_2} + \frac{\sigma^2}{p_a \mu n \varepsilon B} + \frac{L}{\mu} + \frac{\omega}{p_a \mu \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \underbrace{\frac{\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon B}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right)}_{\mathcal{P}_1} \right) \quad (28)$$

731 communication rounds. We take  $b = \Theta \left( \min \left\{ \frac{p_a}{\omega} \sqrt{\frac{\mu n \varepsilon B}{\sigma^2}}, \frac{p_a \mu n \varepsilon B}{\sigma^2} \right\} \right) \geq$   
 732  $\Theta \left( \min \left\{ \frac{p_a}{\omega^2}, \frac{p_a \mu n \varepsilon B}{\sigma^2} \right\} \right).$

733 The method DASHA-SYNC-MVR from (Tyurin and Richtárik, 2023), have to run

$$\tilde{\mathcal{O}} \left( \omega + \frac{\sigma^2}{\mu n \varepsilon B} + \frac{L}{\mu} + \frac{\omega}{\mu \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{n \mu^{3/2} \sqrt{\varepsilon B}} \left( \frac{L_\sigma}{\sqrt{B}} \right) \right) \quad (29)$$

734 communication rounds to get  $\varepsilon$ -solution<sup>8</sup>.

735 In the stochastic setting, the comparison is a little bit more complicated. As in the finite-sum setting,  
 736 we have to take  $B = \mathcal{O} \left( \frac{L_\sigma^2}{\hat{L}^2} \right)$  to guarantee the degeneration up to  $1/p_a$  of the term  $\mathcal{P}_1$  from (28).  
 737 However, DASHA-PP-MVR has also suboptimal term  $\mathcal{P}_2$ . This suboptimality is tightly connected with  
 738 the suboptimality of  $B_{\text{init}}$  in the general nonconvex case, which we discuss in Section 6.3, and it also  
 739 appears in the analysis of DASHA-MVR (Tyurin and Richtárik, 2023). Let us provide the counterpart  
 740 of Corollary 4. The corollary reveals that we can escape regimes when  $\mathcal{P}_2$  is the bottleneck by  
 741 choosing the parameters of the compressors.

742 **Corollary 5.** Suppose that assumptions of Theorem 10 hold, batch size  $B \leq \min \left\{ \frac{\sigma}{p_a \sqrt{\mu \varepsilon n}}, \frac{L_\sigma^2}{\hat{L}^2} \right\}$ ,  
 743 we take RandK compressors with  $K = \Theta \left( \frac{B d \sqrt{\mu \varepsilon n}}{\sigma} \right)$ . Then the communication complexity equals

$$\tilde{\mathcal{O}} \left( \frac{d\sigma}{p_a \sqrt{\mu \varepsilon n}} + \frac{dL_\sigma}{p_a \mu \sqrt{n}} \right),$$

744 and the expected number of stochastic gradient calculations per node equals

$$\tilde{\mathcal{O}} \left( \frac{\sigma^2}{p_a \mu n \varepsilon} + \frac{\sigma L_\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon}} \right).$$

745 Up to Lipschitz constants, DASHA-PP-MVR has the state-of-the-art oracle complexity under PL-  
 746 condition (see (Li et al., 2021a)). Moreover, DASHA-PP-MVR has the state-of-the-art communication  
 747 complexity of DASHA for a small enough  $\mu$ .

<sup>8</sup>For simplicity, we omitted  $\frac{d}{\zeta_C}$  term from the complexity in the stochastic setting, where  $\zeta_C$  is defined in Definition 12. For instance, for the RandK compressor (see Definition 5 and Theorem 6),  $\zeta_C = K$  and  $\frac{d}{\zeta_C} = \Theta(\omega)$ .

#### 748 **F.4 Proofs of Theorems**

749 The following proofs almost repeat the proofs from Section E. And one of the main changes is that  
750 instead of Lemma 3, we use the following lemma.

##### 751 **F.4.1 Standard Lemma under Polyak-Łojasiewicz Condition**

752 **Lemma 11.** *Suppose that Assumptions 1 and 9 hold and*

$$\mathbb{E} [f(x^{t+1})] + \gamma \Psi^{t+1} \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] + (1 - \gamma\mu)\gamma\Psi^t + \gamma C,$$

753 *where  $\Psi^t$  is a sequence of numbers,  $\Psi^t \geq 0$  for all  $t \in [T]$ , constant  $C \geq 0$ , constant  $\mu > 0$ , and*  
754 *constant  $\gamma \in (0, 1/\mu)$ . Then*

$$\mathbb{E} [f(x^T) - f^*] \leq (1 - \gamma\mu)^T ((f(x^0) - f^*) + \gamma\Psi^0) + \frac{C}{\mu}. \quad (30)$$

755 *Proof.* We subtract  $f^*$  and use PL-condition (27) to get

$$\begin{aligned} \mathbb{E} [f(x^{t+1}) - f^*] + \gamma\Psi^{t+1} &\leq \mathbb{E} [f(x^t) - f^*] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] + \gamma\Psi^t + \gamma C \\ &\leq (1 - \gamma\mu) \mathbb{E} [f(x^t) - f^*] + (1 - \gamma\mu)\gamma\Psi^t + \gamma C \\ &= (1 - \gamma\mu) (\mathbb{E} [f(x^t) - f^*] + \gamma\Psi^t) + \gamma C. \end{aligned}$$

756 Unrolling the inequality, we have

$$\begin{aligned} \mathbb{E} [f(x^{t+1}) - f^*] + \gamma\Psi^{t+1} &\leq (1 - \gamma\mu)^{t+1} ((f(x^0) - f^*) + \gamma\Psi^0) + \gamma C \sum_{i=0}^t (1 - \gamma\mu)^i \\ &\leq (1 - \gamma\mu)^{t+1} ((f(x^0) - f^*) + \gamma\Psi^0) + \frac{C}{\mu}. \end{aligned}$$

757 It is left to note that  $\Psi^t \geq 0$  for all  $t \in [T]$ . □

##### 758 **F.4.2 Generic Lemma**

759 We now provide the counterpart of Lemma 6.

760 **Lemma 12.** *Suppose that Assumptions 2, 7, 8 and 9 hold and let us take  $a = \frac{p_a}{2\omega+1}$ , then*

$$\begin{aligned} &\mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ &\leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ &\quad + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ &\quad + \frac{10\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right]. \end{aligned}$$

761 *Proof.* Let us fix some constants  $\kappa, \eta \in [0, \infty)$  that we will define later. Using the same reasoning as  
762 in Lemma 6, we can get

$$\begin{aligned} &\mathbb{E} [f(x^{t+1})] \\ &\quad + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ &\leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \end{aligned}$$

$$\begin{aligned}
& + \left( \gamma + \kappa (1 - a)^2 \right) \mathbb{E} \left[ \|g^t - h^t\|^2 \right] \\
& + \left( \frac{\kappa a^2 ((2\omega + 1) p_a - p_{aa})}{np_a^2} + \eta \left( \frac{a^2 (2\omega + 1 - p_a)}{p_a} + (1 - a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left( \frac{2\kappa\omega}{np_a} + \frac{2\eta\omega}{p_a} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

763 Let us take  $\kappa = \frac{2\gamma}{a}$ . One can show that  $\gamma + \kappa (1 - a)^2 \leq (1 - \frac{a}{2}) \kappa$ , and thus

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] \\
& + \frac{2\gamma}{a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma}{a} \mathbb{E} [\|g^t - h^t\|^2] \\
& + \left( \frac{2\gamma a ((2\omega + 1) p_a - p_{aa})}{np_a^2} + \eta \left( \frac{a^2 (2\omega + 1 - p_a)}{p_a} + (1 - a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left( \frac{4\gamma\omega}{anp_a} + \frac{2\eta\omega}{p_a} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

764 Considering the choice of  $a$ , one can show that  $\left( \frac{a^2 (2\omega + 1 - p_a)}{p_a} + (1 - a)^2 \right) \leq 1 - a$ . If we take  
765  $\eta = \frac{4\gamma((2\omega+1)p_a-p_{aa})}{np_a^2}$ , then  $\left( \frac{2\gamma a ((2\omega + 1) p_a - p_{aa})}{np_a^2} + \eta \left( \frac{a^2 (2\omega + 1 - p_a)}{p_a} + (1 - a)^2 \right) \right) \leq (1 - \frac{a}{2}) \eta$  and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] \\
& + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1) p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left( 1 - \frac{a}{2} \right) \frac{4\gamma((2\omega + 1) p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left( \frac{2\gamma(2\omega + 1)\omega}{np_a^2} + \frac{8\gamma((2\omega + 1) p_a - p_{aa})\omega}{np_a^3} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \\
& \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left( 1 - \frac{a}{2} \right) \frac{4\gamma((2\omega + 1) p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{10\gamma(2\omega + 1)\omega}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right].
\end{aligned}$$

766 It is left to consider that  $\gamma \leq \frac{a}{2\mu}$ , and therefore  $1 - \frac{a}{2} \leq 1 - \gamma\mu$ . □

767 **E.4.3 Proof for DASHA-PP under PL-condition**

**Theorem 8.** Suppose that Assumption 1, 2, 3, 7, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_a}{2-p_a}$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega+1)}{np_a^2} + \frac{48}{np_a^2} \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}} \right)^{-1}, \frac{a}{4\mu} \right\},$$

768 and  $h_i^0 = g_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP), then  $\mathbb{E}[f(x^T)] - f^* \leq$   
 769  $(1 - \gamma\mu)^T \Delta_0$ .

770 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 12, Lemma 7,  
 771 and the law of total expectation, we obtain

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \nu \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}\left[f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left(\frac{1}{2\gamma} - \frac{L}{2}\right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2\right] \\ & + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\ & + \frac{10\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E}\left[2\widehat{L}^2 \|x^{t+1} - x^t\|^2 + 2b^2 \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\ & + \nu \mathbb{E}\left[\frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \|x^{t+1} - x^t\|^2 + \frac{2b^2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2\right] \\ & + \rho \mathbb{E}\left[\frac{2(1-p_a)\widehat{L}^2}{p_a} \|x^{t+1} - x^t\|^2 + \left(\frac{2b^2(1-p_a)}{p_a} + (1-b)^2\right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right]. \end{aligned}$$

772 After rearranging the terms, we get

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \nu \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\ & + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\ & - \left(\frac{1}{2\gamma} - \frac{L}{2} - \frac{20\gamma\omega(2\omega+1)\widehat{L}^2}{np_a^2} - \nu \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} - \rho \frac{2(1-p_a)\widehat{L}^2}{p_a}\right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\ & + (\gamma + \nu(1-b)^2) \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] \\ & + \left(\frac{20b^2\gamma\omega(2\omega+1)}{np_a^2} + \nu \frac{2b^2(p_a - p_{aa})}{np_a^2} + \rho \left(\frac{2b^2(1-p_a)}{p_a} + (1-b)^2\right)\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right]. \end{aligned}$$

773 By taking  $\nu = \frac{2\gamma}{b}$ , one can show that  $(\gamma + \nu(1-b)^2) \leq (1 - \frac{b}{2})\nu$ , and

$$\mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right]$$

$$\begin{aligned}
& + \frac{2\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
\leq & \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{20\gamma\omega(2\omega + 1)\hat{L}^2}{np_a^2} - \frac{4\gamma(p_a - p_{aa})\hat{L}^2}{bnp_a^2} - \rho \frac{2(1 - p_a)\hat{L}^2}{p_a} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( 1 - \frac{b}{2} \right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{20b^2\gamma\omega(2\omega + 1)}{np_a^2} + \frac{4\gamma b(p_a - p_{aa})}{np_a^2} + \rho \left( \frac{2b^2(1 - p_a)}{p_a} + (1 - b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

774 Note that  $b = \frac{p_a}{2 - p_a}$ , thus

$$\begin{aligned}
& \left( \frac{20b^2\gamma\omega(2\omega + 1)}{np_a^2} + \frac{4\gamma b(p_a - p_{aa})}{np_a^2} + \rho \left( \frac{2b^2(1 - p_a)}{p_a} + (1 - b)^2 \right) \right) \\
& \leq \left( \frac{20b^2\gamma\omega(2\omega + 1)}{np_a^2} + \frac{4\gamma b(p_a - p_{aa})}{np_a^2} + \rho(1 - b) \right).
\end{aligned}$$

775 And if we take  $\rho = \frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}$ , then

$$\left( \frac{20b^2\gamma\omega(2\omega + 1)}{np_a^2} + \frac{4\gamma b(p_a - p_{aa})}{np_a^2} + \rho(1 - b) \right) \leq \left( 1 - \frac{b}{2} \right) \rho,$$

776 and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
\leq & \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{20\gamma\omega(2\omega + 1)\hat{L}^2}{np_a^2} - \frac{4\gamma(p_a - p_{aa})\hat{L}^2}{bnp_a^2} \right. \\
& \quad \left. - \frac{80b\gamma\omega(2\omega + 1)(1 - p_a)\hat{L}^2}{np_a^3} - \frac{16\gamma(p_a - p_{aa})(1 - p_a)\hat{L}^2}{np_a^3} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( 1 - \frac{b}{2} \right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( 1 - \frac{b}{2} \right) \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

777 Due to  $\frac{p_a}{2} \leq b \leq p_a$ , we have

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad - \left(\frac{1}{2\gamma} - \frac{L}{2} - \frac{100\gamma\omega(2\omega + 1)\hat{L}^2}{np_a^2} - \frac{24\gamma(p_a - p_{aa})\hat{L}^2}{np_a^3}\right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
&\quad + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left(\frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
\end{aligned}$$

778 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
&\mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
&\quad + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left(\frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
\end{aligned}$$

779 Note that  $\gamma \leq \frac{a}{4\mu} \leq \frac{p_a}{4\mu} \leq \frac{b}{2\mu}$ , thus  $1 - \frac{b}{2} \leq 1 - \gamma\mu$  and

$$\begin{aligned}
&\mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
&\quad + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad + (1 - \gamma\mu) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + (1 - \gamma\mu) \left(\frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
\end{aligned}$$

780 In the view of Lemma 11 with

$$\begin{aligned}
\Psi^t &= \frac{2(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{4((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad + \frac{2}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(\frac{40b\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right],
\end{aligned}$$

781 we can conclude the proof of the theorem.  $\square$

#### 782 F.4.4 Proof for DASHA-PP-PAGE under PL-condition

**Theorem 9.** Suppose that Assumption 1, 2, 3, 7, 4, 8, and 9 hold. Let us take  $a = \frac{p_a}{2\omega + 1}$ , probability

$$p_{\text{page}} = \frac{B}{m+B}, b = \frac{p_{\text{page}} p_a}{2-p_a},$$

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega + 1)}{np_a^2} \left( \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right)} + \frac{48}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

783 and  $h_i^0 = g_i^0 = \nabla f_i(x^0)$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-PAGE), then  $\mathbb{E}[f(x^T)] - f^* \leq$   
 784  $(1 - \gamma\mu)^T \Delta_0$ .

785 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 12, Lemma 8,  
 786 and the law of total expectation, we obtain

$$\begin{aligned}
 & \mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
 & + \nu \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
 & \leq \mathbb{E}\left[f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left(\frac{1}{2\gamma} - \frac{L}{2}\right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2\right] \\
 & + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
 & + \frac{10\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2\right] \\
 & + \nu \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
 & \leq \mathbb{E}\left[f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left(\frac{1}{2\gamma} - \frac{L}{2}\right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2\right] \\
 & + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
 & + \frac{10\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E}\left[\left(2\hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B}\right) \|x^{t+1} - x^t\|^2 + \frac{2b^2}{p_{\text{page}}} \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\
 & + \nu \mathbb{E}\left[\left(\frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} + \frac{(1 - p_{\text{page}})L_{\max}^2}{np_a B}\right) \|x^{t+1} - x^t\|^2\right. \\
 & \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{page}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left(p_{\text{page}} \left(1 - \frac{b}{p_{\text{page}}}\right)^2 + (1 - p_{\text{page}})\right) \|h^t - \nabla f(x^t)\|^2\right] \\
 & + \rho \mathbb{E}\left[\left(\frac{2(1 - p_a)\hat{L}^2}{p_a} + \frac{(1 - p_{\text{page}})L_{\max}^2}{p_a B}\right) \|x^{t+1} - x^t\|^2\right. \\
 & \quad \left. + \left(\frac{2(1 - p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left(1 - \frac{b}{p_{\text{page}}}\right)^2 + (1 - p_{\text{page}})\right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right].
 \end{aligned}$$

787 After rearranging the terms, we get

$$\begin{aligned}
 & \mathbb{E}[f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
 & + \nu \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
 & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
 & + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right]
 \end{aligned}$$

$$\begin{aligned}
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( 2\widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \nu \left( \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} + \frac{(1-p_{\text{page}})L_{\max}^2}{np_a B} \right) - \rho \left( \frac{2(1-p_a)\widehat{L}^2}{p_a} + \frac{(1-p_{\text{page}})L_{\max}^2}{p_a B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( \gamma + \nu \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \right) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{2\nu(p_a - p_{aa})b^2}{np_a^2 p_{\text{page}}} \right. \\
& \quad \left. + \rho \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Due to  $b = \frac{p_{\text{page}} p_a}{2-p_a} \leq p_{\text{page}}$ , one can show that  $\left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \leq 1-b$ . Thus, if we take  $\nu = \frac{2\gamma}{b}$ , then

$$\left( \gamma + \nu \left( p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \right) \leq \gamma + \nu(1-b) = \left( 1 - \frac{b}{2} \right) \nu,$$

788 therefore

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( 2\widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \frac{2\gamma}{bn p_a} \left( 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \widehat{L}^2 + \frac{(1-p_{\text{page}})L_{\max}^2}{B} \right) - \rho \left( \frac{2(1-p_a)\widehat{L}^2}{p_a} + \frac{(1-p_{\text{page}})L_{\max}^2}{p_a B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( 1 - \frac{b}{2} \right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{4\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{page}}} \right. \\
& \quad \left. + \rho \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Next, with the choice of  $b = \frac{p_{\text{page}} p_a}{2-p_a}$ , we ensure that

$$\left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \leq 1-b.$$

If we take  $\rho = \frac{40b^2\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}}$ , then

$$\left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{4\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{page}}} + \rho \left( \frac{2(1-p_a)b^2}{p_a p_{\text{page}}} + p_{\text{page}} \left( 1 - \frac{b}{p_{\text{page}}} \right)^2 + (1-p_{\text{page}}) \right) \right) \leq \left( 1 - \frac{b}{2} \right) \rho,$$



789 therefore

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( 2\hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \frac{2\gamma}{bnp_a} \left( 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^3 p_{\text{page}}} + \frac{8\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2(1 - p_a) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left( 1 - \frac{b}{2} \right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( 1 - \frac{b}{2} \right) \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

Let us simplify the inequality. First, due to  $b \geq \frac{p_{\text{page}} p_a}{2}$ , we have

$$\frac{2\gamma}{bnp_a} \left( 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \leq \frac{8\gamma}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right).$$

790 Second, due to  $b \leq p_a p_{\text{page}}$  and  $p_{aa} \leq p_a^2$ , we get

$$\begin{aligned}
& \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^3 p_{\text{page}}} + \frac{8\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2(1 - p_a) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \\
& \leq \left( \frac{40\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2 p_{\text{page}}} \right) \left( 2 \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \\
& \leq \frac{80\gamma\omega(2\omega+1)}{np_a^2} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \\
& \quad + \frac{16\gamma \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \\
& \leq \frac{80\gamma\omega(2\omega+1)}{np_a^2} \left( \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \\
& \quad + \frac{16\gamma}{np_a^2 p_{\text{page}}} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right).
\end{aligned}$$

791 Combining all bounds together, we obtain the following inequality:

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2]
\end{aligned}$$

$$\begin{aligned}
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{100\gamma\omega(2\omega + 1)}{np_a^2} \left( \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right. \\
& \quad \left. - \frac{24\gamma}{np_a^2 p_{\text{page}}} \left( \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2 + \frac{(1 - p_{\text{page}})L_{\max}^2}{B} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

792 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

793 Note that  $\gamma \leq \frac{b}{2\mu}$ , thus  $1 - \frac{b}{2} \leq 1 - \gamma\mu$  and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{4\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + (1 - \gamma\mu) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + (1 - \gamma\mu) \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right].
\end{aligned}$$

794 It is left to apply Lemma 11 with

$$\begin{aligned}
\Psi^t &= \frac{2(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{4((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{2}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left( \frac{40b\omega(2\omega + 1)}{np_a^2 p_{\text{page}}} + \frac{8(p_a - p_{aa})}{np_a^2 p_{\text{page}}} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]
\end{aligned}$$

795 to conclude the proof.  $\square$

796 **F.4.5 Proof for DASHA-PP-MVR under PL-condition**

**Theorem 10.** Suppose that Assumption 1, 2, 3, 7, 5, 6, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,

$$b \in \left(0, \frac{p_a}{2-p_a}\right],$$

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{200\omega(2\omega+1)}{np_a^2} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \widehat{L}^2 \right) + \frac{40}{np_a b} \left( \frac{(1-b)^2 L_\sigma^2}{B} + \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2 \right)} \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

797 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 1 (DASHA-PP-MVR), then

$$\begin{aligned} & \mathbb{E} [f(x^T) - f^*] \\ & \leq (1-\gamma\mu)^T \left( \Delta_0 + \frac{2\gamma}{b} \|h^0 - \nabla f(x^0)\|^2 + \left( \frac{40\gamma b\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) \\ & \quad + \frac{1}{\mu} \left( \frac{100b^2\omega(2\omega+1)}{p_a^2} + \frac{20b}{p_a} \right) \frac{\sigma^2}{nB}. \end{aligned}$$

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & \quad + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & \quad + \frac{10\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right]. \end{aligned}$$

798 *Proof.* Let us fix constants  $\nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 12, Lemma 10,  
799 and the law of total expectation, we obtain

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & \quad + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & \quad + \frac{10\gamma(2\omega+1)\omega}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|k_i^{t+1}\|^2 \right] \\ & \quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma \|h^t - \nabla f(x^t)\|^2 \right] \\ & \quad + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{10\gamma\omega(2\omega+1)}{np_a^2} \mathbb{E} \left[ \frac{2b^2\sigma^2}{B} + \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 + 2b^2 \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \nu \mathbb{E} \left( \frac{2b^2\sigma^2}{np_a B} + \left( \frac{2(1-b)^2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b)^2 \|h^t - \nabla f(x^t)\|^2 \right) \\
& + \rho \mathbb{E} \left( \frac{2b^2\sigma^2}{p_a B} + \left( \frac{2(1-b)^2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right. \\
& \quad \left. + \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right).
\end{aligned}$$

800 After rearranging the terms, we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \right. \\
& \quad \left. - \nu \left( \frac{2(1-b)^2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2(1-b)^2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + (\gamma + \nu(1-b)^2) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{2\nu(p_a - p_{aa})b^2}{np_a^2} + \rho \left( \frac{2(1-p_a)b^2}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2} + \nu \frac{2b^2}{np_a} + \rho \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

801 By taking  $\nu = \frac{2\gamma}{b}$ , one can show that  $(\gamma + \nu(1-b)^2) \leq (1 - \frac{b}{2})\nu$ , and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \right.
\end{aligned}$$

$$\begin{aligned}
& -\frac{2\gamma}{b} \left( \frac{2(1-b)^2 L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa}) \hat{L}^2}{np_a^2} \right) - \rho \left( \frac{2(1-b)^2 L_\sigma^2}{p_a B} + \frac{2(1-p_a) \hat{L}^2}{p_a} \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma (p_a - p_{aa}) b}{np_a^2} + \rho \left( \frac{2(1-p_a) b^2}{p_a} + (1-b)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma b}{np_a} + \rho \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

802 Note that  $b \leq \frac{p_a}{2-p_a}$ , thus

$$\begin{aligned}
& \left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma (p_a - p_{aa}) b}{np_a^2} + \rho \left( \frac{2(1-p_a) b^2}{p_a} + (1-b)^2 \right) \right) \\
& \leq \left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma (p_a - p_{aa}) b}{np_a^2} + \rho (1-b) \right).
\end{aligned}$$

803 And if we take  $\rho = \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a-p_{aa})}{np_a^2}$ , then

$$\left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma (p_a - p_{aa}) b}{np_a^2} + \rho (1-b) \right) \leq \rho,$$

804 and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a-p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{10\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2\hat{L}^2 \right) \right. \\
& \quad \left. - \frac{2\gamma}{np_a b} \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2 \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^3} + \frac{8\gamma \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a^2} \right) \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 2(1-p_a) \hat{L}^2 \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a-p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{20b^2 \gamma \omega (2\omega + 1)}{np_a^2} + \frac{4\gamma b}{np_a} + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a-p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

805 Let us simplify the inequality. First, due to  $b \leq p_a$  and  $(1-p_a) \leq \left(1 - \frac{p_{aa}}{p_a}\right)$ , we have

$$\left( \frac{40b\gamma\omega(2\omega+1)}{np_a^3} + \frac{2\gamma \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a^2} \right) \left( \frac{2(1-b)^2 L_\sigma^2}{B} + 8(1-p_a) \hat{L}^2 \right)$$

$$\begin{aligned}
&= \frac{40b\gamma\omega(2\omega+1)}{np_a^3} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2(1-p_a)\widehat{L}^2 \right) \\
&\quad + \frac{8\gamma \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a^2} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2(1-p_a)\widehat{L}^2 \right) \\
&\leq \frac{40\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \\
&\quad + \frac{8\gamma}{np_ab} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2 \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2 \right),
\end{aligned}$$

806 therefore

$$\begin{aligned}
&\mathbb{E}[f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\
&\quad + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{50\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2\widehat{L}^2 \right) \right. \\
&\quad \left. - \frac{10\gamma}{np_ab} \left( \frac{2(1-b)^2L_\sigma^2}{B} + 2 \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2 \right) \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
&\quad + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\
&\quad + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{4\gamma b}{np_a} + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B} \\
&\leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
&\quad + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{100\gamma\omega(2\omega+1)}{np_a^2} \left( \frac{(1-b)^2L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \frac{20\gamma}{np_ab} \left( \frac{(1-b)^2L_\sigma^2}{B} + \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2 \right) \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
&\quad + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\
&\quad + \left( \frac{20b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{4\gamma b}{np_a} + \left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

807 Also, we can simplify the last term:

$$\begin{aligned}
&\left( \frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \frac{2b^2}{p_a} \\
&= \frac{80b^3\gamma\omega(2\omega+1)}{np_a^3} + \frac{16b^2\gamma \left(1 - \frac{p_{aa}}{p_a}\right)}{np_a^2}
\end{aligned}$$

$$\leq \frac{80b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{16b\gamma}{np_a},$$

808 thus

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\ & + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\ & - \left(\frac{1}{2\gamma} - \frac{L}{2} - \frac{100\gamma\omega(2\omega+1)}{np_a^2} \left(\frac{(1-b)^2 L_\sigma^2}{B} + \widehat{L}^2\right)\right. \\ & \quad \left. - \frac{20\gamma}{np_a b} \left(\frac{(1-b)^2 L_\sigma^2}{B} + \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2\right)\right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\ & + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left(\frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\ & + \left(\frac{100b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{20\gamma b}{np_a}\right) \frac{\sigma^2}{B}. \end{aligned}$$

809 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\ & + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \\ & + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \left(\frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2\right] \\ & + \left(\frac{100b^2\gamma\omega(2\omega+1)}{np_a^2} + \frac{20\gamma b}{np_a}\right) \frac{\sigma^2}{B}. \end{aligned}$$

810 Note that  $\gamma \leq \frac{b}{2\mu}$ , thus  $1 - \frac{b}{2} \leq 1 - \gamma\mu$  and

$$\begin{aligned} & \mathbb{E}[f(x^{t+1})] + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2\right] \\ & + \frac{2\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \left(\frac{40b\gamma\omega(2\omega+1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2\right] \\ & \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\ & + (1-\gamma\mu) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + (1-\gamma\mu) \frac{4\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2\right] \end{aligned}$$

$$\begin{aligned}
& + (1 - \gamma\mu) \frac{2\gamma}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + (1 - \gamma\mu) \left( \frac{40b\gamma\omega(2\omega + 1)}{np_a^2} + \frac{8\gamma(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{100b^2\gamma\omega(2\omega + 1)}{np_a^2} + \frac{20\gamma b}{np_a} \right) \frac{\sigma^2}{B}.
\end{aligned}$$

811 It is left to apply Lemma 11 with

$$\begin{aligned}
\Psi^t &= \frac{2(2\omega + 1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{4((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&+ \frac{2}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \left( \frac{40b\omega(2\omega + 1)}{np_a^2} + \frac{8(p_a - p_{aa})}{np_a^2} \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]
\end{aligned}$$

812 and  $C = \left( \frac{100b^2\omega(2\omega+1)}{p_a^2} + \frac{20b}{p_a} \right) \frac{\sigma^2}{nB}$  to conclude the proof.  $\square$

813 **Corollary 5.** Suppose that assumptions of Theorem 10 hold, batch size  $B \leq \min \left\{ \frac{\sigma}{p_a\sqrt{\mu\varepsilon n}}, \frac{L_\sigma^2}{L^2} \right\}$ ,  
814 we take RandK compressors with  $K = \Theta \left( \frac{Bd\sqrt{\mu\varepsilon n}}{\sigma} \right)$ . Then the communication complexity equals

$$\tilde{\mathcal{O}} \left( \frac{d\sigma}{p_a\sqrt{\mu\varepsilon n}} + \frac{dL_\sigma}{p_a\mu\sqrt{n}} \right),$$

815 and the expected number of stochastic gradient calculations per node equals

$$\tilde{\mathcal{O}} \left( \frac{\sigma^2}{p_a\mu n\varepsilon} + \frac{\sigma L_\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon}} \right).$$

816 *Proof.* In the view of Theorem 10, DASHA-PP have to run

$$\tilde{\mathcal{O}} \left( \frac{\omega + 1}{p_a} + \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n\varepsilon B}} + \frac{\sigma^2}{p_a\mu n\varepsilon B} + \frac{L}{\mu} + \frac{\omega}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon B}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right)$$

817 communication rounds in the stochastic settings to get  $\varepsilon$ -solution. Note that  $K = \mathcal{O} \left( \frac{d}{p_a\sqrt{n}} \right)$ .

818 Moreover, we can skip the initialization procedure and initialize  $h_i^0$  and  $g_i^0$ , for instance, with zeros  
819 because the initialization error is under a logarithm. Considering Theorem 6, the communication  
820 complexity equals

$$\begin{aligned}
& \tilde{\mathcal{O}} \left( K \frac{\omega + 1}{p_a} + K \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n\varepsilon B}} + K \frac{\sigma^2}{p_a\mu n\varepsilon B} + K \frac{L}{\mu} + K \frac{\omega}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + K \frac{\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon B}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right) \\
&= \tilde{\mathcal{O}} \left( K \frac{\omega + 1}{p_a} + K \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n\varepsilon B}} + K \frac{\sigma^2}{p_a\mu n\varepsilon B} + K \frac{L}{\mu} + K \frac{\omega}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + K \frac{\sigma L_\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon B}} \right) \\
&= \tilde{\mathcal{O}} \left( \frac{d}{p_a} + \frac{d}{p_a} \sqrt{\frac{\sigma^2}{\mu n\varepsilon B}} + \frac{K\sigma^2}{p_a\mu n\varepsilon B} + \frac{dL}{p_a\mu\sqrt{n}} + \frac{d}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{K\sigma L_\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon B}} \right) \\
&= \tilde{\mathcal{O}} \left( \frac{d}{p_a} + \frac{d\sigma}{p_a\sqrt{\mu n\varepsilon B}} + \frac{d\sigma}{p_a\sqrt{\mu\varepsilon n}} + \frac{dL}{p_a\mu\sqrt{n}} + \frac{d}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{dL_\sigma}{p_a\mu\sqrt{n}} \right) \\
&= \tilde{\mathcal{O}} \left( \frac{d\sigma}{p_a\sqrt{\mu\varepsilon n}} + \frac{dL_\sigma}{p_a\mu\sqrt{n}} \right).
\end{aligned}$$

821 The expected number of stochastic gradient calculations per node equals

$$\tilde{\mathcal{O}} \left( B \frac{\omega + 1}{p_a} + B \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n\varepsilon B}} + B \frac{\sigma^2}{p_a\mu n\varepsilon B} + B \frac{L}{\mu} + B \frac{\omega}{p_a\mu\sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + B \frac{\sigma}{p_a n\mu^{3/2}\sqrt{\varepsilon B}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) \right)$$



$$\begin{aligned}
&= \tilde{\mathcal{O}} \left( B \frac{\omega + 1}{p_a} + B \frac{\omega}{p_a} \sqrt{\frac{\sigma^2}{\mu n \varepsilon B}} + B \frac{\sigma^2}{p_a \mu n \varepsilon B} + B \frac{L}{\mu} + B \frac{\omega}{p_a \mu \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + B \frac{\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon B}} \left( \frac{L_\sigma}{\sqrt{B}} \right) \right) \\
&= \tilde{\mathcal{O}} \left( \frac{Bd}{K p_a} + \frac{Bd}{K p_a} \sqrt{\frac{\sigma^2}{\mu n \varepsilon B}} + \frac{\sigma^2}{p_a \mu n \varepsilon} + B \frac{L}{\mu} + \frac{Bd}{K p_a \mu \sqrt{n}} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma L_\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon}} \right) \\
&= \tilde{\mathcal{O}} \left( \frac{\sigma}{p_a \sqrt{\mu \varepsilon n}} + \frac{\sigma^2}{p_a \mu \varepsilon n \sqrt{B}} + \frac{\sigma^2}{p_a \mu n \varepsilon} + \frac{\sigma L}{p_a \mu^{3/2} \sqrt{\varepsilon} n} + \frac{\sigma}{p_a \mu^{3/2} \sqrt{\varepsilon} n} \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma L_\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon}} \right) \\
&= \tilde{\mathcal{O}} \left( \frac{\sigma^2}{p_a \mu n \varepsilon} + \frac{\sigma L_\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon}} \right).
\end{aligned}$$

822

□

## 823 G Description of DASHA-PP-SYNC-MVR

824 By analogy to (Tyurin and Richtárik, 2023), we provide a “synchronized” version of the algorithm.  
 825 With a small probability, participating nodes calculate and send a mega batch without compression.  
 826 This helps us to resolve the suboptimality of DASHA-PP-MVR w.r.t.  $\omega$ . Note that this suboptimality is  
 827 not a problem. We show in Corollary 4 that DASHA-PP-MVR can have the optimal oracle complexity  
 828 and SOTA communication complexity with the particular choices of parameters of the compressors.

---

### Algorithm 8 DASHA-PP-SYNC-MVR

---

```

1: Input: starting point  $x^0 \in \mathbb{R}^d$ , stepsize  $\gamma > 0$ , momentum  $a \in (0, 1]$ , momentum  $b \in$ 
   (0, 1], probability  $p_{\text{mega}} \in (0, 1]$ , batch size  $B'$  and  $B$ , probability  $p_a \in (0, 1]$  that a node is
   participating(a), number of iterations  $T \geq 1$ .
2: Initialize  $g_i^0, h_i^0$  on the nodes and  $g^0 = \frac{1}{n} \sum_{i=1}^n g_i^0$  on the server
3: for  $t = 0, 1, \dots, T - 1$  do
4:    $x^{t+1} = x^t - \gamma g^t$ 
5:    $c^{t+1} = \begin{cases} 1, & \text{with probability } p_{\text{mega}}, \\ 0, & \text{with probability } 1 - p_{\text{mega}} \end{cases}$ 
6:   Broadcast  $x^{t+1}, x^t$  to all participating(a) nodes
7:   for  $i = 1, \dots, n$  in parallel do
8:     if  $i^{\text{th}}$  node is participating(a) then
9:       if  $c^{t+1} = 1$  then
10:        Generate i.i.d. samples  $\{\xi_{ik}^{t+1}\}_{k=1}^{B'}$  of size  $B'$  from  $\mathcal{D}_i$ .
11:         $k_i^{t+1} = \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) - \frac{b}{p_{\text{mega}}} \left( h_i^t - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) \right)$ 
12:         $m_i^{t+1} = \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t)$ 
13:       else
14:        Generate i.i.d. samples  $\{\xi_{ij}^{t+1}\}_{j=1}^B$  of size  $B$  from  $\mathcal{D}_i$ .
15:         $k_i^{t+1} = \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1})$ 
16:         $m_i^{t+1} = C_i \left( \frac{1}{p_a} k_i^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right)$ 
17:       end if
18:        $h_i^{t+1} = h_i^t + \frac{1}{p_a} k_i^{t+1}$ 
19:        $g_i^{t+1} = g_i^t + m_i^{t+1}$ 
20:       Send  $m_i^{t+1}$  to the server
21:     else
22:        $h_i^{t+1} = h_i^t$ 
23:        $m_i^{t+1} = 0$ 
24:        $g_i^{t+1} = g_i^t$ 
25:     end if
26:   end for
27:    $g^{t+1} = g^t + \frac{1}{n} \sum_{i=1}^n m_i^{t+1}$ 
28: end for
29: Output:  $\hat{x}^T$  chosen uniformly at random from  $\{x^t\}_{k=0}^{T-1}$ 
   (a): For the formal description see Section 2.2.

```

---

829 In the following theorem, we provide the convergence rate of DASHA-PP-SYNC-MVR.

**Theorem 11.** Suppose that Assumptions 1, 2, 3, 5, 6, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  
 $b = \frac{p_{\text{mega}} p_a}{2-p_a}$ , probability  $p_{\text{mega}} \in (0, 1]$ , batch size  $B' \geq B \geq 1$

$$\gamma \leq \left( L + \sqrt{\frac{8(2\omega+1)\omega}{np_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right) + \frac{16}{np_{\text{mega}} p_a^2} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{L_\sigma^2}{B} \right)} \right)^{-1},$$

830 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 8. Then

$$\begin{aligned} \mathbb{E} \left[ \|\nabla f(\hat{x}^T)\|^2 \right] &\leq \frac{1}{T} \left[ \frac{2\Delta_0}{\gamma} + \frac{4}{p_{\text{mega}} p_a} \|h^0 - \nabla f(x^0)\|^2 + \frac{4 \left(1 - \frac{p_{aa}}{p_a}\right)}{n p_{\text{mega}} p_a} \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right] \\ &\quad + \frac{12\sigma^2}{nB'}. \end{aligned}$$

831 First, we introduce the expected density of compressors (Gorbunov et al., 2021; Tyurin and Richtárik, 2023).

833 **Definition 12.** The expected density of the compressor  $\mathcal{C}_i$  is  $\zeta_{\mathcal{C}_i} := \sup_{x \in \mathbb{R}^d} \mathbb{E} [\|\mathcal{C}_i(x)\|_0]$ , where  
834  $\|x\|_0$  is the number of nonzero components of  $x \in \mathbb{R}^d$ . Let  $\zeta_{\mathcal{C}} = \max_{i \in [n]} \zeta_{\mathcal{C}_i}$ .

835 Note that  $\zeta_{\mathcal{C}}$  is finite and  $\zeta_{\mathcal{C}} \leq d$ .

836 In the next corollary, we choose particular algorithm parameters to reveal the communication and  
837 oracle complexity.

**Corollary 6.** Suppose that assumptions from Theorem 11 hold, probability  $p_{\text{mega}} = \min \left\{ \frac{\zeta_{\mathcal{C}}}{d}, \frac{n\varepsilon B}{\sigma^2} \right\}$ ,  
batch size  $B' = \Theta \left( \frac{\sigma^2}{n\varepsilon} \right)$ , and  $h_i^0 = g_i^0 = \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0)$  for all  $i \in [n]$ , initial batch size  
 $B_{\text{init}} = \Theta \left( \frac{B}{p_{\text{mega}} \sqrt{p_a}} \right) = \Theta \left( \max \left\{ \frac{Bd}{\sqrt{p_a} \zeta_{\mathcal{C}}}, \frac{\sigma^2}{\sqrt{p_a} n\varepsilon} \right\} \right)$ , then DASHA-PP-SYNC-MVR needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \left( \frac{\omega}{p_a \sqrt{n}} + \sqrt{\frac{d}{p_a^2 \zeta_{\mathcal{C}} n}} \right) \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon} n} \left( \frac{\hat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right] + \frac{\sigma^2}{\sqrt{p_a} n \varepsilon B} \right).$$

838 communication rounds to get an  $\varepsilon$ -solution, the expected communication complexity is equal to  
839  $\mathcal{O}(d + \zeta_{\mathcal{C}} T)$ , and the expected number of stochastic gradient calculations per node equals  $\mathcal{O}(B_{\text{init}} +$   
840  $BT)$ , where  $\zeta_{\mathcal{C}}$  is the expected density from Definition 12.

841 The main improvement of Corollary 6 over Corollary 3 is the size of the initial batch size  $B_{\text{init}}$ .  
842 However, Corollary 4 reveals that we can avoid regimes when DASHA-PP-MVR is suboptimal.

843 We also provide a theorem under PL-condition (see Assumption 9).

**Theorem 13.** Suppose that Assumptions 1, 2, 3, 5, 6, 7, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  
 $b = \frac{p_{\text{mega}} p_a}{2-p_a}$ , probability  $p_{\text{mega}} \in (0, 1]$ , batch size  $B' \geq B \geq 1$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{16(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \hat{L}^2 \right) + \left( \frac{48L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{24 \left(1 - \frac{p_{aa}}{p_a}\right) \hat{L}^2}{np_{\text{mega}} p_a^2} \right)} \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

844 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 8. Then

$$\begin{aligned} &\mathbb{E} [f(x^T) - f^*] \\ &\leq (1 - \gamma\mu)^T \left( \Delta_0 + \frac{2\gamma}{b} \|h^0 - \nabla f(x^0)\|^2 + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) + \frac{20\sigma^2}{\mu n B'}. \end{aligned}$$

845 Let us provide bounds up to logarithmic factors and use  $\tilde{\mathcal{O}}(\cdot)$  notation.

**Corollary 7.** Suppose that assumptions from Theorem 13 hold, probability  $p_{\text{mega}} =$   
 $\min \left\{ \frac{\zeta_{\mathcal{C}}}{d}, \frac{\mu n \varepsilon B}{\sigma^2} \right\}$ , batch size  $B' = \Theta \left( \frac{\sigma^2}{\mu n \varepsilon} \right)$  then DASHA-PP-SYNC-MVR needs

$$T := \tilde{\mathcal{O}} \left( \frac{\omega+1}{p_a} + \frac{d}{p_a \zeta_{\mathcal{C}}} + \frac{\sigma^2}{p_a \mu n \varepsilon B} + \frac{L}{\mu} + \frac{\omega}{p_a \mu \sqrt{n}} \left( \frac{L_\sigma}{\sqrt{B}} + \hat{L} \right) + \left( \frac{\sqrt{d}}{p_a \mu \sqrt{\zeta_{\mathcal{C}} n}} + \frac{\sigma}{p_a \mu^{3/2} \sqrt{\varepsilon B}} \right) \left( \frac{L_\sigma}{\sqrt{B}} + \hat{L} \right) \right).$$

communication rounds to get an  $\varepsilon$ -solution, the expected communication complexity is equal to  $\tilde{\mathcal{O}}(\zeta_c T)$ , and the expected number of stochastic gradient calculations per node equals  $\tilde{\mathcal{O}}(BT)$ , where  $\zeta_c$  is the expected density from Definition 12.

The proof of this corollary almost repeats the proof of Corollary 6. Note that we can skip the initialization procedure and initialize  $h_i^0$  and  $g_i^0$ , for instance, with zeros because the initialization error is under a logarithm.

Let us assume that  $\frac{d}{\zeta_c} = \Theta(\omega)$  (holds for the RandK compressor), then the convergence rate of DASHA-PP-SYNC-MVR is

$$\tilde{\mathcal{O}}\left(\frac{\omega+1}{p_a} + \frac{\sigma^2}{p_a \mu n \varepsilon B} + \frac{L}{\mu} + \frac{\omega}{p_a \mu \sqrt{n}} \left(\frac{L_\sigma}{\sqrt{B}} + \hat{L}\right) + \frac{\sigma}{p_a n \mu^{3/2} \sqrt{\varepsilon B}} \left(\frac{L_\sigma}{\sqrt{B}} + \hat{L}\right)\right). \quad (31)$$

Comparing (31) with the rate of DASHA-PP-MVR (28), one can see that DASHA-PP-SYNC-MVR improves the suboptimal term  $\mathcal{P}_2$  from (28). However, Corollary 5 reveals that we can escape these suboptimal regimes by choosing the parameter  $K$  of RandK compressors in a particular way.

### G.1 Proof for DASHA-PP-SYNC-MVR

In this section, we provide the proof of the convergence rate for DASHA-PP-SYNC-MVR. There are four different sources of randomness in Algorithm 8: the first one from random samples  $\xi_i^{t+1}$ , the second one from compressors  $\{\mathcal{C}_i\}_{i=1}^n$ , the third one from availability of nodes, and the fourth one from  $c^{t+1}$ . We define  $\mathbb{E}_k[\cdot]$ ,  $\mathbb{E}_c[\cdot]$ ,  $\mathbb{E}_{p_a}[\cdot]$  and  $\mathbb{E}_{p_{\text{mega}}}[\cdot]$  to be conditional expectations w.r.t.  $\xi_i^{t+1}$ ,  $\{\mathcal{C}_i\}_{i=1}^n$ , availability, and  $c^{t+1}$ , accordingly, conditioned on all previous randomness. Moreover, we define  $\mathbb{E}_{t+1}[\cdot]$  to be a conditional expectation w.r.t. all randomness in iteration  $t+1$  conditioned on all previous randomness.

Let us denote

$$\begin{aligned} k_{i,1}^{t+1} &:= \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) - \frac{b}{p_{\text{mega}}} \left( h_i^t - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) \right), \\ k_{i,2}^{t+1} &:= \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1}), \\ h_{i,1}^{t+1} &:= \begin{cases} h_i^t + \frac{1}{p_a} k_{i,1}^{t+1}, & i^{\text{th}} \text{ node is participating,} \\ h_i^t, & \text{otherwise,} \end{cases} \\ h_{i,2}^{t+1} &:= \begin{cases} h_i^t + \frac{1}{p_a} k_{i,2}^{t+1}, & i^{\text{th}} \text{ node is participating,} \\ h_i^t, & \text{otherwise,} \end{cases} \\ g_{i,1}^{t+1} &:= \begin{cases} g_i^t + \frac{1}{p_a} k_{i,1}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t), & i^{\text{th}} \text{ node is participating,} \\ g_i^t, & \text{otherwise,} \end{cases} \\ g_{i,2}^{t+1} &:= \begin{cases} g_i^t + \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right), & i^{\text{th}} \text{ node is participating,} \\ g_i^t, & \text{otherwise,} \end{cases} \end{aligned}$$

$h_1^{t+1} := \frac{1}{n} \sum_{i=1}^n h_{i,1}^{t+1}$ ,  $h_2^{t+1} := \frac{1}{n} \sum_{i=1}^n h_{i,2}^{t+1}$ ,  $g_1^{t+1} := \frac{1}{n} \sum_{i=1}^n g_{i,1}^{t+1}$ , and  $g_2^{t+1} := \frac{1}{n} \sum_{i=1}^n g_{i,2}^{t+1}$ . Note, that

$$h^{t+1} = \begin{cases} h_1^{t+1}, & c^{t+1} = 1, \\ h_2^{t+1}, & c^{t+1} = 0, \end{cases}$$

and

$$g^{t+1} = \begin{cases} g_1^{t+1}, & c^{t+1} = 1, \\ g_2^{t+1}, & c^{t+1} = 0 \end{cases}$$

First, we will prove two lemmas.

870 **Lemma 13.** Suppose that Assumptions 3, 5, 7 and 8 hold and let us consider sequences  $\{g_i^{t+1}\}_{i=1}^n$   
 871 and  $\{h_i^{t+1}\}_{i=1}^n$  from Algorithm 8, then

$$\begin{aligned} & \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \right] \\ & \leq \frac{2(1-p_{\text{mega}})\omega}{n^2 p_a} \sum_{i=1}^n \|k_{i,2}^{t+1}\|^2 + \left( \frac{(p_a - p_{aa})a^2}{n^2 p_a^2} + \frac{2(1-p_{\text{mega}})a^2\omega}{n^2 p_a} \right) \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \\ & \quad + (1-a)^2 \|g^t - h^t\|^2, \end{aligned}$$

872 and

$$\begin{aligned} & \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \right] \\ & \leq \frac{2(1-p_{\text{mega}})\omega}{p_a} \|k_{i,2}^{t+1}\|^2 + \left( \frac{(1-p_a)a^2}{p_a} + \frac{2(1-p_{\text{mega}})a^2\omega}{p_a} \right) \|g_i^t - h_i^t\|^2 \\ & \quad + (1-a)^2 \|g_i^t - h_i^t\|^2, \quad \forall i \in [n]. \end{aligned}$$

873 *Proof.* First, we get the bound for  $\mathbb{E}_{t+1} [\|g^{t+1} - h^{t+1}\|^2]$ :

$$\begin{aligned} & \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \right] \\ & = p_{\text{mega}} \mathbb{E}_{p_a} \left[ \|g_1^{t+1} - h_1^{t+1}\|^2 \right] + (1-p_{\text{mega}}) \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g_2^{t+1} - h_2^{t+1}\|^2 \right] \right]. \end{aligned}$$

874 Using

$$\mathbb{E}_{p_a} [g_{i,1}^{t+1} - h_{i,1}^{t+1}] = g_i^t + k_{i,1}^{t+1} - a(g_i^t - h_i^t) - h_i^t - k_{i,1}^{t+1} = (1-a)(g_i^t - h_i^t)$$

875 and

$$\mathbb{E}_{\mathcal{C}} [\mathbb{E}_{p_a} [g_{i,2}^{t+1} - h_{i,2}^{t+1}]] = g_i^t + k_{i,2}^{t+1} - a(g_i^t - h_i^t) - h_i^t - k_{i,2}^{t+1} = (1-a)(g_i^t - h_i^t),$$

876 we have

$$\begin{aligned} & \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \right] \\ & \stackrel{(13)}{=} p_{\text{mega}} \mathbb{E}_{p_a} \left[ \|g_1^{t+1} - h_1^{t+1} - \mathbb{E}_{p_a} [g_1^{t+1} - h_1^{t+1}]\|^2 \right] \\ & \quad + (1-p_{\text{mega}}) \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \|g_2^{t+1} - h_2^{t+1} - \mathbb{E}_{p_a} [g_2^{t+1} - h_2^{t+1}]\|^2 \right] \right] \\ & \quad + (1-a)^2 \|g^t - h^t\|^2. \end{aligned}$$

877 We can use Lemma 1 two times with i)  $r_i = g_i^t - h_i^t$  and  $s_i = -a(g_i^t - h_i^t)$  and ii)  $r_i = g_i^t - h_i^t$  and  
 878  $s_i = p_a \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - k_{i,2}^{t+1}$ , to obtain

$$\begin{aligned} & \mathbb{E}_{\mathcal{C}} \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \right] \\ & \leq \frac{p_{\text{mega}} a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \\ & \quad + (1-p_{\text{mega}}) \left( \frac{1}{n^2 p_a} \sum_{i=1}^n \mathbb{E}_{\mathcal{C}} \left[ \left\| p_a \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - (k_{i,2}^{t+1} - a(g_i^t - h_i^t)) \right\|^2 \right] \right) \\ & \quad + (1-p_{\text{mega}}) \left( \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right) \\ & \quad + (1-a)^2 \|g^t - h^t\|^2 \\ & = \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \end{aligned}$$

$$\begin{aligned}
& + (1 - p_{\text{mega}}) \left( \frac{p_a}{n^2} \sum_{i=1}^n \mathbb{E}_C \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right\|^2 \right] \right) \\
& + (1 - a)^2 \|g^t - h^t\|^2 \\
& \leq \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \\
& + \frac{(1 - p_{\text{mega}}) p_a \omega}{n^2} \sum_{i=1}^n \left\| \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right\|^2 \\
& + (1 - a)^2 \|g^t - h^t\|^2 \\
& = \frac{a^2 (p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \\
& + \frac{(1 - p_{\text{mega}}) \omega}{n^2 p_a} \sum_{i=1}^n \|k_{i,2}^{t+1} - a (g_i^t - h_i^t)\|^2 \\
& + (1 - a)^2 \|g^t - h^t\|^2.
\end{aligned}$$

879 In the last inequality, we use Assumption 7. Next, using (12), we have

$$\begin{aligned}
& \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] \right] \right] \\
& \leq \frac{2(1 - p_{\text{mega}}) \omega}{n^2 p_a} \sum_{i=1}^n \|k_{i,2}^{t+1}\|^2 + \left( \frac{(p_a - p_{aa}) a^2}{n^2 p_a^2} + \frac{2(1 - p_{\text{mega}}) \omega a^2}{n^2 p_a} \right) \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \\
& + (1 - a)^2 \|g^t - h^t\|^2.
\end{aligned}$$

880 The second inequality can be proved almost in the same way:

$$\begin{aligned}
& \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \right] \\
& = p_{\text{mega}} \mathbb{E}_{p_a} \left[ \|g_{i,1}^{t+1} - h_{i,1}^{t+1}\|^2 \right] + (1 - p_{\text{mega}}) \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g_{i,2}^{t+1} - h_{i,2}^{t+1}\|^2 \right] \right] \\
& \stackrel{(13)}{=} p_{\text{mega}} \mathbb{E}_{p_a} \left[ \|g_{i,1}^{t+1} - h_{i,1}^{t+1} - (1 - a)(g_i^t - h_i^t)\|^2 \right] + (1 - p_{\text{mega}}) \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g_{i,2}^{t+1} - h_{i,2}^{t+1}\|^2 \right] \right] \\
& + p_{\text{mega}} (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& = \frac{p_{\text{mega}} (1 - p_a) a^2}{p_a} \|g_i^t - h_i^t\|^2 + (1 - p_{\text{mega}}) \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g_{i,2}^{t+1} - h_{i,2}^{t+1}\|^2 \right] \right] \\
& + p_{\text{mega}} (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& \stackrel{(13)}{=} \frac{p_{\text{mega}} (1 - p_a) a^2}{p_a} \|g_i^t - h_i^t\|^2 + (1 - p_{\text{mega}}) \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \|g_{i,2}^{t+1} - h_{i,2}^{t+1} - (1 - a)(g_i^t - h_i^t)\|^2 \right] \right] \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& = \frac{p_{\text{mega}} (1 - p_a) a^2}{p_a} \|g_i^t - h_i^t\|^2 \\
& + (1 - p_{\text{mega}}) p_a \mathbb{E}_C \left[ \left\| g_i^t + \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( h_i^t + \frac{1}{p_a} k_{i,2}^{t+1} \right) - (1 - a)(g_i^t - h_i^t) \right\|^2 \right] \\
& + (1 - p_{\text{mega}}) (1 - p_a) \|g_i^t - h_i^t - (1 - a)(g_i^t - h_i^t)\|^2 \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& = \frac{p_{\text{mega}} (1 - p_a) a^2}{p_a} \|g_i^t - h_i^t\|^2 \\
& + (1 - p_{\text{mega}}) p_a \mathbb{E}_C \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_{i,2}^{t+1} - a(g_i^t - h_i^t) \right) \right\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
& + (1 - p_{\text{mega}}) (1 - p_a) a^2 \|g_i^t - h_i^t\|^2 \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& \stackrel{(13)}{=} \left( \frac{p_{\text{mega}}(1 - p_a)a^2}{p_a} + \frac{(1 - p_{\text{mega}})(1 - p_a)a^2}{p_a} \right) \|g_i^t - h_i^t\|^2 \\
& + (1 - p_{\text{mega}}) p_a \mathbb{E} \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right\|^2 \right] \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& = \frac{(1 - p_a)a^2}{p_a} \|g_i^t - h_i^t\|^2 \\
& + (1 - p_{\text{mega}}) p_a \mathbb{E} \left[ \left\| \mathcal{C}_i \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) - \left( \frac{1}{p_a} k_{i,2}^{t+1} - \frac{a}{p_a} (g_i^t - h_i^t) \right) \right\|^2 \right] \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& \leq \frac{(1 - p_a)a^2}{p_a} \|g_i^t - h_i^t\|^2 \\
& + \frac{(1 - p_{\text{mega}})\omega}{p_a} \|k_{i,2}^{t+1} - a(g_i^t - h_i^t)\|^2 \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2 \\
& \stackrel{(12)}{\leq} \frac{2(1 - p_{\text{mega}})\omega}{p_a} \|k_{i,2}^{t+1}\|^2 + \left( \frac{(1 - p_a)a^2}{p_a} + \frac{2(1 - p_{\text{mega}})a^2\omega}{p_a} \right) \|g_i^t - h_i^t\|^2 \\
& + (1 - a)^2 \|g_i^t - h_i^t\|^2.
\end{aligned}$$

881

□

882 **Lemma 14.** Suppose that Assumptions 3, 5, 6 and 8 hold and let us consider sequence  $\{h_i^{t+1}\}_{i=1}^n$   
883 from Algorithm 8, then

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\
& \leq \frac{2b^2\sigma^2}{np_{\text{mega}}p_a B'} + \left( \frac{2p_{\text{mega}}L_\sigma^2}{np_a B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1 - p_{\text{mega}})L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\hat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \\
& + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2,
\end{aligned}$$

884

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\
& \leq \frac{2b^2\sigma^2}{p_a p_{\text{mega}} B'} + \left( \frac{2p_{\text{mega}}L_\sigma^2}{p_a B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1 - p_{\text{mega}})L_\sigma^2}{p_a B} + \frac{2(1 - p_a)L_i^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \\
& + \frac{2(1 - p_a)b^2}{p_{\text{mega}}p_a} \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2, \quad \forall i \in [n],
\end{aligned}$$

885 and

$$\mathbb{E}_k \left[ \|k_{i,2}^{t+1}\|^2 \right] \leq \left( \frac{L_\sigma^2}{B} + L_i^2 \right) \|x^{t+1} - x^t\|^2, \quad \forall i \in [n],$$

886 *Proof.* First, we prove the bound for  $\mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right]$ . Using

$$\mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ h_{i,1}^{t+1} \right] \right]$$

$$\begin{aligned}
&= h_i^t + E_k \left[ \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) - \frac{b}{p_{\text{mega}}} \left( h_i^t - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) \right) \right] \\
&= h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{mega}}} (h_i^t - \nabla f_i(x^t))
\end{aligned}$$

887 and

$$\begin{aligned}
&E_k [E_{p_a} [h_{i,2}^{t+1}]] \\
&= h_i^t + E_k \left[ \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1}) \right] \\
&= h_i^t + \nabla f_i(x^{t+1}) - \nabla f_i(x^t),
\end{aligned}$$

888 we have

$$\begin{aligned}
&E_k [E_{p_a} [E_{p_{\text{mega}}} [\|h^{t+1} - \nabla f(x^{t+1})\|^2]]] \\
&= p_{\text{mega}} E_k [E_{p_a} [\|h_1^{t+1} - \nabla f(x^{t+1})\|^2]] + (1 - p_{\text{mega}}) E_k [E_{p_a} [\|h_2^{t+1} - \nabla f(x^{t+1})\|^2]] \\
&\stackrel{(13)}{=} p_{\text{mega}} E_k [E_{p_a} [\|h_1^{t+1} - E_k [E_{p_a} [h_1^{t+1}]]\|^2]] + (1 - p_{\text{mega}}) E_k [E_{p_a} [\|h_2^{t+1} - E_k [E_{p_a} [h_2^{t+1}]]\|^2]] \\
&\quad + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2.
\end{aligned}$$

889 We can use Lemma 1 two times with i)  $r_i = h_i^t$  and  $s_i = k_{i,1}^{t+1}$  and ii)  $r_i = h_i^t$  and  $s_i = k_{i,2}^{t+1}$ , to  
890 obtain

$$\begin{aligned}
&E_k [E_{p_a} [E_{p_{\text{mega}}} [\|h^{t+1} - \nabla f(x^{t+1})\|^2]]] \\
&\leq p_{\text{mega}} \left( \frac{1}{n^2 p_a} \sum_{i=1}^n E_k [\|k_{i,1}^{t+1} - E_k [k_{i,1}^{t+1}]\|^2] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{mega}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \right) \\
&\quad + (1 - p_{\text{mega}}) \left( \frac{1}{n^2 p_a} \sum_{i=1}^n E_k [\|k_{i,2}^{t+1} - E_k [k_{i,2}^{t+1}]\|^2] + \frac{p_a - p_{aa}}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \right) \\
&\quad + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2 \\
&\stackrel{(12)}{\leq} \frac{p_{\text{mega}}}{n^2 p_a} \sum_{i=1}^n E_k [\|k_{i,1}^{t+1} - E_k [k_{i,1}^{t+1}]\|^2] \\
&\quad + \frac{1 - p_{\text{mega}}}{n^2 p_a} \sum_{i=1}^n E_k [\|k_{i,2}^{t+1} - E_k [k_{i,2}^{t+1}]\|^2] \\
&\quad + \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
&\quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2.
\end{aligned} \tag{32}$$

891 Let us consider  $E_k [\|k_{i,1}^{t+1} - E_k [k_{i,1}^{t+1}]\|^2]$ .

$$\begin{aligned}
&E_k [\|k_{i,1}^{t+1} - E_k [k_{i,1}^{t+1}]\|^2] \\
&= E_k \left[ \left\| \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) - \frac{b}{p_{\text{mega}}} \left( h_i^t - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) \right) \right\|^2 \right]
\end{aligned}$$



$$\begin{aligned}
& - \left( \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{mega}}} (h_i^t - \nabla f_i(x^t)) \right) \Big\|^2 \Big] \\
& = \mathbb{E}_k \left[ \left\| \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) + \frac{b}{p_{\text{mega}}} \left( \frac{1}{B'} \sum_{k=1}^{B'} \nabla f_i(x^t; \xi_{ik}^{t+1}) \right) \right. \right. \\
& \quad \left. \left. - \left( \nabla f_i(x^{t+1}) - \nabla f_i(x^t) + \frac{b}{p_{\text{mega}}} (\nabla f_i(x^t)) \right) \right\|^2 \right] \\
& = \frac{1}{B'^2} \sum_{k=1}^{B'} \mathbb{E}_k \left[ \left\| \frac{b}{p_{\text{mega}}} (\nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \nabla f_i(x^{t+1})) \right. \right. \\
& \quad \left. \left. + \left( 1 - \frac{b}{p_{\text{mega}}} \right) (\nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \nabla f_i(x^t; \xi_{ik}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t))) \right\|^2 \right],
\end{aligned}$$

where we used independence of the mini-batch samples. Using (12), we get

$$\begin{aligned}
& \mathbb{E}_k \left[ \left\| k_{i,1}^{t+1} - \mathbb{E}_k [k_{i,1}^{t+1}] \right\|^2 \right] \\
& \leq \frac{2b^2}{B'^2 p_{\text{mega}}^2} \sum_{k=1}^{B'} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \nabla f_i(x^{t+1}) \right\|^2 \right] \\
& \quad + \frac{2}{B'^2} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 \sum_{k=1}^{B'} \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ik}^{t+1}) - \nabla f_i(x^t; \xi_{ik}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right].
\end{aligned}$$

Due to Assumptions 5 and 6, we have

$$\mathbb{E}_k \left[ \left\| k_{i,1}^{t+1} - \mathbb{E}_k [k_{i,1}^{t+1}] \right\|^2 \right] \leq \frac{2b^2 \sigma^2}{B' p_{\text{mega}}^2} + \frac{2L_\sigma^2}{B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 \|x^{t+1} - x^t\|^2. \quad (33)$$

Next, we estimate the bound for  $\mathbb{E}_k \left[ \left\| k_{i,2}^{t+1} - \mathbb{E}_k [k_{i,2}^{t+1}] \right\|^2 \right]$ .

$$\begin{aligned}
& \mathbb{E}_k \left[ \left\| k_{i,2}^{t+1} - \mathbb{E}_k [k_{i,2}^{t+1}] \right\|^2 \right] \\
& = \mathbb{E}_k \left[ \left\| \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \frac{1}{B} \sum_{j=1}^B \nabla f_i(x^t; \xi_{ij}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right] \\
& = \frac{1}{B^2} \sum_{j=1}^B \mathbb{E}_k \left[ \left\| \nabla f_i(x^{t+1}; \xi_{ij}^{t+1}) - \nabla f_i(x^t; \xi_{ij}^{t+1}) - (\nabla f_i(x^{t+1}) - \nabla f_i(x^t)) \right\|^2 \right].
\end{aligned}$$

Due to Assumptions 6, we have

$$\mathbb{E}_k \left[ \left\| k_{i,2}^{t+1} - \mathbb{E}_k [k_{i,2}^{t+1}] \right\|^2 \right] \leq \frac{L_\sigma^2}{B} \|x^{t+1} - x^t\|^2. \quad (34)$$

Plugging (33) and (34) into (32), we obtain

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\
& \leq \frac{p_{\text{mega}}}{np_a} \left( \frac{2b^2 \sigma^2}{B' p_{\text{mega}}^2} + \frac{2L_\sigma^2}{B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 \|x^{t+1} - x^t\|^2 \right) \\
& \quad + \frac{(1 - p_{\text{mega}}) L_\sigma^2}{np_a B} \|x^{t+1} - x^t\|^2 \\
& \quad + \frac{2(p_a - p_{aa})}{n^2 p_a^2} \sum_{i=1}^n \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2
\end{aligned}$$

$$+ \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2.$$

897 Using Assumption 3, we get

$$\begin{aligned} & \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] \right] \right] \\ & \leq \frac{2b^2 \sigma^2}{n p_{\text{mega}} p_a B'} + \left( \frac{2p_{\text{mega}} L_\sigma^2}{n p_a B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1 - p_{\text{mega}}) L_\sigma^2}{n p_a B} + \frac{2(p_a - p_{aa}) \widehat{L}^2}{n p_a^2} \right) \|x^{t+1} - x^t\|^2 \\ & \quad + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2. \end{aligned}$$

898 Using almost the same derivations, we can prove the second inequality:

$$\begin{aligned} & \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\ & = p_{\text{mega}} \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h_{i,1}^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] + (1 - p_{\text{mega}}) \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h_{i,2}^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \\ & \stackrel{(13)}{=} p_{\text{mega}} \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h_{i,1}^{t+1} - \mathbb{E}_k [h_{i,1}^{t+1}]\|^2 \right] \right] + (1 - p_{\text{mega}}) \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \|h_{i,2}^{t+1} - \mathbb{E}_k [h_{i,2}^{t+1}]\|^2 \right] \right] \\ & \quad + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\ & = p_{\text{mega}} p_a \mathbb{E}_k \left[ \left\| h_i^t + \frac{1}{p_a} k_{i,1}^{t+1} - (h_i^t + \mathbb{E}_k [k_{i,1}^{t+1}]) \right\|^2 \right] \\ & \quad + p_{\text{mega}} (1 - p_a) \|h_i^t - (h_i^t + \mathbb{E}_k [k_{i,1}^{t+1}])\|^2 \\ & \quad + (1 - p_{\text{mega}}) p_a \mathbb{E}_k \left[ \left\| h_i^t + \frac{1}{p_a} k_{i,2}^{t+1} - (h_i^t + \mathbb{E}_k [k_{i,2}^{t+1}]) \right\|^2 \right] \\ & \quad + (1 - p_{\text{mega}}) (1 - p_a) \|h_i^t - (h_i^t + \mathbb{E}_k [k_{i,2}^{t+1}])\|^2 \\ & \quad + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\ & = p_{\text{mega}} p_a \mathbb{E}_k \left[ \left\| \frac{1}{p_a} k_{i,1}^{t+1} - \mathbb{E}_k [k_{i,1}^{t+1}] \right\|^2 \right] \\ & \quad + p_{\text{mega}} (1 - p_a) \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{mega}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \quad + (1 - p_{\text{mega}}) p_a \mathbb{E}_k \left[ \left\| \frac{1}{p_a} k_{i,2}^{t+1} - \mathbb{E}_k [k_{i,2}^{t+1}] \right\|^2 \right] \\ & \quad + (1 - p_{\text{mega}}) (1 - p_a) \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\ & \quad + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\ & \stackrel{(13)}{=} \frac{p_{\text{mega}}}{p_a} \mathbb{E}_k \left[ \|k_{i,1}^{t+1} - \mathbb{E}_k [k_{i,1}^{t+1}]\|^2 \right] \\ & \quad + \frac{(1 - p_{\text{mega}})}{p_a} \mathbb{E}_k \left[ \|k_{i,2}^{t+1} - \mathbb{E}_k [k_{i,2}^{t+1}]\|^2 \right] \\ & \quad + \frac{p_{\text{mega}} (1 - p_a)}{p_a} \left\| \nabla f_i(x^{t+1}) - \nabla f_i(x^t) - \frac{b}{p_{\text{mega}}} (h_i^t - \nabla f_i(x^t)) \right\|^2 \\ & \quad + \frac{(1 - p_{\text{mega}}) (1 - p_a)}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \end{aligned}$$

$$\begin{aligned}
& + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2 \\
& \stackrel{(12)}{\leq} \frac{p_{\text{mega}}}{p_a} \mathbb{E}_k \left[ \|k_{i,1}^{t+1} - \mathbb{E}_k[k_{i,1}^{t+1}]\|^2 \right] \\
& + \frac{(1 - p_{\text{mega}})}{p_a} \mathbb{E}_k \left[ \|k_{i,2}^{t+1} - \mathbb{E}_k[k_{i,2}^{t+1}]\|^2 \right] \\
& + \frac{2(1 - p_a)}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
& + \frac{2(1 - p_a)b^2}{p_{\text{mega}}p_a} \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2.
\end{aligned}$$

899 Using (33) and (34), we get

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\
& \leq \frac{2b^2\sigma^2}{p_ap_{\text{mega}}B'} + \frac{2p_{\text{mega}}L_\sigma^2}{p_aB'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 \|x^{t+1} - x^t\|^2 \\
& + \frac{(1 - p_{\text{mega}})L_\sigma^2}{p_aB} \|x^{t+1} - x^t\|^2 \\
& + \frac{2(1 - p_a)}{p_a} \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
& + \frac{2(1 - p_a)b^2}{p_{\text{mega}}p_a} \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2.
\end{aligned}$$

900 Next, due to Assumption 3, we obtain

$$\begin{aligned}
& \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \\
& \leq \frac{2b^2\sigma^2}{p_ap_{\text{mega}}B'} + \left( \frac{2p_{\text{mega}}L_\sigma^2}{p_aB'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1 - p_{\text{mega}})L_\sigma^2}{p_aB} + \frac{2(1 - p_a)L_i^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \\
& + \frac{2(1 - p_a)b^2}{p_{\text{mega}}p_a} \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1 - p_{\text{mega}}) \right) \|h_i^t - \nabla f_i(x^t)\|^2.
\end{aligned}$$

901 The third inequality can be proved with the help of (34) and Assumption 3.

$$\begin{aligned}
& \mathbb{E}_k \left[ \|k_{i,2}^{t+1}\|^2 \right] \\
& \stackrel{(13)}{=} \mathbb{E}_k \left[ \|k_{i,2}^{t+1} - \mathbb{E}_k[k_{i,2}^{t+1}]\|^2 \right] + \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
& \leq \frac{L_\sigma^2}{B} \|x^{t+1} - x^t\|^2 + \|\nabla f_i(x^{t+1}) - \nabla f_i(x^t)\|^2 \\
& \leq \left( \frac{L_\sigma^2}{B} + L_i^2 \right) \|x^{t+1} - x^t\|^2.
\end{aligned}$$

902

□

**Theorem 11.** Suppose that Assumptions 1, 2, 3, 5, 6, 7 and 8 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_{\text{mega}}p_a}{2-p_a}$ , probability  $p_{\text{mega}} \in (0, 1]$ , batch size  $B' \geq B \geq 1$

$$\gamma \leq \left( L + \sqrt{\frac{8(2\omega+1)\omega}{np_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right) + \frac{16}{np_{\text{mega}}p_a^2} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{L_\sigma^2}{B} \right)} \right)^{-1},$$

903 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 8. Then

$$\mathbb{E} \left[ \|\nabla f(\hat{x}^T)\|^2 \right] \leq \frac{1}{T} \left[ \frac{2\Delta_0}{\gamma} + \frac{4}{p_{\text{mega}}p_a} \|h^0 - \nabla f(x^0)\|^2 + \frac{4 \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_{\text{mega}}p_a} \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right]$$

$$+ \frac{12\sigma^2}{nB'}.$$

904 *Proof.* Due to Lemma 2 and the update step from Line 4 in Algorithm 8, we have

$$\begin{aligned} & \mathbb{E}_{t+1} [f(x^{t+1})] \\ & \leq \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - \nabla f(x^t)\|^2 \right] \\ & = \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \frac{\gamma}{2} \|g^t - h^t + h^t - \nabla f(x^t)\|^2 \right] \\ & \stackrel{(13)}{\leq} \mathbb{E}_{t+1} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma (\|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2) \right]. \end{aligned}$$

905 Let us fix constants  $\kappa, \eta, \nu, \rho \in [0, \infty)$  that we will define later. Considering Lemma 13, Lemma 14,  
906 and the law of total expectation, we obtain

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & \quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma (\|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2) \right] \\ & \quad + \kappa \mathbb{E} \left[ \mathbb{E}_k \left[ \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} [\|g^{t+1} - h^{t+1}\|^2] \right] \right] \right] \right] \\ & \quad + \eta \mathbb{E} \left[ \mathbb{E}_k \left[ \mathbb{E}_C \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \right] \right] \right] \right] \\ & \quad + \nu \mathbb{E} \left[ \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] \right] \right] \right] \\ & \quad + \rho \mathbb{E} \left[ \mathbb{E}_k \left[ \mathbb{E}_{p_a} \left[ \mathbb{E}_{p_{\text{mega}}} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \right] \right] \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma (\|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2) \right] \\ & \quad + \kappa \mathbb{E} \left( \frac{2(1-p_{\text{mega}})\omega}{np_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \left( \frac{(p_a - p_{aa})a^2}{n^2 p_a^2} + \frac{2(1-p_{\text{mega}})a^2\omega}{n^2 p_a} \right) \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \right) \\ & \quad + \eta \mathbb{E} \left( \frac{2(1-p_{\text{mega}})\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \left( \frac{(1-p_a)a^2}{p_a} + \frac{2(1-p_{\text{mega}})a^2\omega}{p_a} \right) \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \right) \\ & \quad + \nu \mathbb{E} \left( \frac{2b^2\sigma^2}{np_{\text{mega}}p_a B'} + \left( \frac{2p_{\text{mega}}L_\sigma^2}{np_a B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1-p_{\text{mega}})L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2 p_a^2 p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1-p_{\text{mega}}) \right) \|h^t - \nabla f(x^t)\|^2 \right) \\ & \quad + \rho \mathbb{E} \left( \frac{2b^2\sigma^2}{p_a p_{\text{mega}} B'} + \left( \frac{2p_{\text{mega}}L_\sigma^2}{p_a B'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + \frac{(1-p_{\text{mega}})L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right) \end{aligned}$$

$$+ \frac{2(1-p_a)b^2}{np_{\text{mega}}p_a} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + \left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1-p_{\text{mega}}) \right) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \Bigg).$$

Let us simplify the last inequality. Since  $B' \geq B$  and  $b = \frac{p_{\text{mega}}p_a}{2-p_a} \leq p_{\text{mega}}$ , we have  $1 - p_{\text{mega}} \leq 1$ ,

$$\frac{2p_{\text{mega}}L_\sigma^2}{p_aB'} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 \leq \frac{2p_{\text{mega}}L_\sigma^2}{p_aB},$$

$$\left( p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1-p_{\text{mega}}) \right) \leq 1 - b,$$

and

$$\left( \frac{2(1-p_a)b^2}{p_{\text{mega}}p_a} + p_{\text{mega}} \left( 1 - \frac{b}{p_{\text{mega}}} \right)^2 + (1-p_{\text{mega}}) \right) \leq 1 - b.$$

907 Thus

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} \left[ f(x^t) - \frac{\gamma}{2} \|\nabla f(x^t)\|^2 - \left( \frac{1}{2\gamma} - \frac{L}{2} \right) \|x^{t+1} - x^t\|^2 + \gamma (\|g^t - h^t\|^2 + \|h^t - \nabla f(x^t)\|^2) \right] \\ & + \kappa \mathbb{E} \left( \frac{2\omega}{np_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \frac{((2\omega + 1)p_a - p_{aa})a^2}{n^2p_a^2} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \right) \\ & + \eta \mathbb{E} \left( \frac{2\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \frac{(2\omega + 1 - p_a)a^2}{p_a} \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 + (1-a)^2 \|g^t - h^t\|^2 \right) \\ & + \nu \mathbb{E} \left( \frac{2b^2\sigma^2}{np_{\text{mega}}p_aB'} + \left( \frac{2L_\sigma^2}{np_aB} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + \frac{2(p_a - p_{aa})b^2}{n^2p_a^2p_{\text{mega}}} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 + (1-b) \|h^t - \nabla f(x^t)\|^2 \right) \\ & + \rho \mathbb{E} \left( \frac{2b^2\sigma^2}{p_ap_{\text{mega}}B'} + \left( \frac{2L_\sigma^2}{p_aB} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \|x^{t+1} - x^t\|^2 \right. \\ & \quad \left. + (1-b) \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right). \end{aligned}$$

908 After rearranging the terms, we get

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{2\kappa\omega}{np_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \frac{2\eta\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
&\quad + (\gamma + \kappa(1-a)^2) \mathbb{E} [\|g^t - h^t\|^2] \\
&\quad + \left( \kappa \frac{((2\omega+1)p_a - p_{aa})a^2}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&\quad + (\gamma + \nu(1-b)) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
&\quad + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
&\quad + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'}.
\end{aligned}$$

909 Let us take  $\kappa = \frac{\gamma}{a}$ , thus  $\gamma + \kappa(1-a)^2 \leq \kappa$  and

$$\begin{aligned}
&\mathbb{E} [f(x^{t+1})] + \frac{\gamma}{a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
&\quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
&\leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{2\gamma\omega}{anp_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \frac{2\eta\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
&\quad + \frac{\gamma}{a} \mathbb{E} [\|g^t - h^t\|^2] \\
&\quad + \left( \frac{\gamma((2\omega+1)p_a - p_{aa})a}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&\quad + (\gamma + \nu(1-b)) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
&\quad + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
&\quad + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'}.
\end{aligned}$$

910 Next, since  $a = \frac{p_a}{2\omega+1}$ , we have  $\left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \leq 1-a$ . We the choice  $\eta =$

911  $\frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2}$ , we guarantee  $\frac{\gamma((2\omega+1)p_a - p_{aa})a}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \leq \eta$  and

$$\begin{aligned}
&\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
&\quad + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{2\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \frac{2\gamma((2\omega+1)p_a - p_{aa})\omega}{np_a^3} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
&\quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&\quad + (\gamma + \nu(1-b)) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
&\quad + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
&\quad + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'} \\
&\leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
&\quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&\quad + (\gamma + \nu(1-b)) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
&\quad + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
&\quad + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'},
\end{aligned}$$

912 where simplified the term using  $p_{aa} \geq 0$ . Let us take  $\nu = \frac{\gamma}{b}$  to obtain

$$\begin{aligned}
&\mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
&\quad + \frac{\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
&\leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
&\quad - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
&\quad \left. - \left( \frac{2\gamma L_\sigma^2}{bn p_a B} + \frac{2\gamma(p_a - p_{aa})\widehat{L}^2}{bn p_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
&\quad + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \frac{\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&\quad + \frac{\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2]
\end{aligned}$$

$$\begin{aligned}
& + \left( \frac{2\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{2\gamma b}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'}.
\end{aligned}$$

913 Next, we take  $\rho = \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}}$ , thus

$$\begin{aligned}
& \mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma(2\omega + 1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \hat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{2\gamma L_\sigma^2}{bn p_a B} + \frac{2\gamma(p_a - p_{aa})\hat{L}^2}{bn p_a^2} \right) - \left( \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \right) \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\hat{L}^2}{p_a} \right) \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{2\gamma b}{np_{\text{mega}} p_a} + \frac{4\gamma(p_a - p_{aa})b^2}{np_a^3 p_{\text{mega}}} \right) \frac{\sigma^2}{B'}.
\end{aligned}$$

914 Since  $\frac{p_{\text{mega}} p_a}{2} \leq b \leq p_{\text{mega}} p_a$  and  $1 - p_a \leq 1 - \frac{p_{aa}}{p_a} \leq 1$ , we get

$$\begin{aligned}
& \mathbb{E}[f(x^{t+1})] + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^{t+1} - h^{t+1}\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma(2\omega + 1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \hat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{4\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{4\gamma(p_a - p_{aa})\hat{L}^2}{np_{\text{mega}} p_a^3} \right) - \left( \frac{4\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{4\gamma(1-p_a)\hat{L}^2}{np_{\text{mega}} p_a^2} \right) \right) \mathbb{E}[\|x^{t+1} - x^t\|^2] \\
& + \frac{\gamma(2\omega + 1)}{p_a} \mathbb{E}[\|g^t - h^t\|^2] + \frac{\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E}[\|h^t - \nabla f(x^t)\|^2] + \frac{2\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{6\gamma\sigma^2}{nB'} \\
& \leq \mathbb{E}[f(x^t)] - \frac{\gamma}{2} \mathbb{E}[\|\nabla f(x^t)\|^2]
\end{aligned}$$



$$\begin{aligned}
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \left( \frac{8\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{8\gamma \left(1 - \frac{p_{\text{aa}}}{p_a}\right) \widehat{L}^2}{np_{\text{mega}} p_a^2} \right) \right) \mathbb{E} \left[ \|x^{t+1} - x^t\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{6\gamma\sigma^2}{nB'}.
\end{aligned}$$

915 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} \left[ \|\nabla f(x^t)\|^2 \right] \\
& + \frac{\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{\gamma((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \frac{\gamma}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \frac{2\gamma(p_a - p_{\text{aa}})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{6\gamma\sigma^2}{nB'}.
\end{aligned}$$

916 It is left to apply Lemma 3 with

$$\begin{aligned}
\Psi^t &= \frac{(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{((2\omega+1)p_a - p_{\text{aa}})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
&+ \frac{1}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \frac{2 \left(1 - \frac{p_{\text{aa}}}{p_a}\right)}{np_a p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]
\end{aligned}$$

917 and  $C = \frac{6\sigma^2}{nB'}$  to conclude the proof.  $\square$

**Corollary 6.** Suppose that assumptions from Theorem 11 hold, probability  $p_{\text{mega}} = \min \left\{ \frac{\zeta_c}{d}, \frac{n\varepsilon B}{\sigma^2} \right\}$ , batch size  $B' = \Theta \left( \frac{\sigma^2}{n\varepsilon} \right)$ , and  $h_i^0 = g_i^0 = \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0)$  for all  $i \in [n]$ , initial batch size  $B_{\text{init}} = \Theta \left( \frac{B}{p_{\text{mega}} \sqrt{p_a}} \right) = \Theta \left( \max \left\{ \frac{Bd}{\sqrt{p_a} \zeta_c}, \frac{\sigma^2}{\sqrt{p_a} n \varepsilon} \right\} \right)$ , then DASHA-PP-SYNC-MVR needs

$$T := \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \left( \frac{\omega}{p_a \sqrt{n}} + \sqrt{\frac{d}{p_a^2 \zeta_c n}} \right) \left( \widehat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon} n} \left( \frac{\widehat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right] + \frac{\sigma^2}{\sqrt{p_a} n \varepsilon B} \right).$$

918 communication rounds to get an  $\varepsilon$ -solution, the expected communication complexity is equal to  
919  $\mathcal{O}(d + \zeta_c T)$ , and the expected number of stochastic gradient calculations per node equals  $\mathcal{O}(B_{\text{init}} +$   
920  $BT)$ , where  $\zeta_c$  is the expected density from Definition 12.

921 *Proof.* Due to the choice of  $B'$ , we have

$$\begin{aligned} \mathbb{E} \left[ \|\nabla f(\hat{x}^T)\|^2 \right] &\leq \frac{1}{T} \left[ 2\Delta_0 \left( L + \sqrt{\frac{8(2\omega+1)\omega}{np_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right) + \frac{16}{np_{\text{mega}}p_a^2} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{L_\sigma^2}{B} \right)} \right) \right. \\ &\quad \left. + \frac{4}{p_{\text{mega}}p_a} \|h^0 - \nabla f(x^0)\|^2 + \frac{4 \left( 1 - \frac{p_{aa}}{p_a} \right)}{np_{\text{mega}}p_a} \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right] \\ &\quad + \frac{2\varepsilon}{3}. \end{aligned}$$

922 Using

$$\mathbb{E} \left[ \|h^0 - \nabla f(x^0)\|^2 \right] = \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0) - \nabla f(x^0) \right\|^2 \right] \leq \frac{\sigma^2}{nB_{\text{init}}}$$

923 and

$$\frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \|h_i^0 - \nabla f_i(x^0)\|^2 \right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left[ \left\| \frac{1}{B_{\text{init}}} \sum_{k=1}^{B_{\text{init}}} \nabla f_i(x^0; \xi_{ik}^0) - \nabla f_i(x^0) \right\|^2 \right] \leq \frac{\sigma^2}{nB_{\text{init}}},$$

924 we have

$$\begin{aligned} \mathbb{E} \left[ \|\nabla f(\hat{x}^T)\|^2 \right] &\leq \frac{1}{T} \left[ 2\Delta_0 \left( L + \sqrt{\frac{8(2\omega+1)\omega}{np_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right) + \frac{16}{np_{\text{mega}}p_a^2} \left( \left( 1 - \frac{p_{aa}}{p_a} \right) \hat{L}^2 + \frac{L_\sigma^2}{B} \right)} \right) \right. \\ &\quad \left. + \frac{8\sigma^2}{np_{\text{mega}}p_a B_{\text{init}}} \right] \\ &\quad + \frac{2\varepsilon}{3}. \end{aligned}$$

925 Therefore, we can take the following  $T$  to get  $\varepsilon$ -solution.

$$T = \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \sqrt{\frac{\omega^2}{np_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right) + \frac{1}{np_{\text{mega}}p_a^2} \left( \hat{L}^2 + \frac{L_\sigma^2}{B} \right)} \right) + \frac{\sigma^2}{np_{\text{mega}}p_a B_{\text{init}}} \right] \right)$$

926 Considering the choice of  $p_{\text{mega}}$  and  $B_{\text{init}}$ , we obtain

$$\begin{aligned} T &= \mathcal{O} \left( \frac{1}{\varepsilon} \left[ \Delta_0 \left( L + \left( \frac{\omega}{p_a \sqrt{n}} + \sqrt{\frac{d}{p_a^2 \zeta_C n}} \right) \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon n}} \left( \frac{\hat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right) + \frac{\sigma^2}{np_{\text{mega}}p_a B_{\text{init}}} \right] \right) \\ &= \mathcal{O} \left( \frac{\Delta_0}{\varepsilon} \left[ L + \left( \frac{\omega}{p_a \sqrt{n}} + \sqrt{\frac{d}{p_a^2 \zeta_C n}} \right) \left( \hat{L} + \frac{L_\sigma}{\sqrt{B}} \right) + \frac{\sigma}{p_a \sqrt{\varepsilon n}} \left( \frac{\hat{L}}{\sqrt{B}} + \frac{L_\sigma}{B} \right) \right] + \frac{\sigma^2}{\sqrt{p_a} n \varepsilon B} \right). \end{aligned}$$

927 The expected communication complexity equals  $\mathcal{O}(d + p_{\text{mega}}d + (1 - p_{\text{mega}})\zeta_C) =$   
 928  $\mathcal{O}(d + \zeta_C)$  and the expected number of stochastic gradient calculations per node equals  
 929  $\mathcal{O}(B_{\text{init}} + p_{\text{mega}}B' + (1 - p_{\text{mega}})B) = \mathcal{O}(B_{\text{init}} + B)$ .  $\square$

**Theorem 13.** Suppose that Assumptions 1, 2, 3, 5, 6, 7, 8 and 9 hold. Let us take  $a = \frac{p_a}{2\omega+1}$ ,  $b = \frac{p_{\text{mega}}p_a}{2-p_a}$ , probability  $p_{\text{mega}} \in (0, 1]$ , batch size  $B' \geq B \geq 1$ ,

$$\gamma \leq \min \left\{ \left( L + \sqrt{\frac{16(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) + \left( \frac{48L_\sigma^2}{np_{\text{mega}}p_a^2B} + \frac{24 \left( 1 - \frac{p_{aa}}{p_a} \right) \widehat{L}^2}{np_{\text{mega}}p_a^2} \right)} \right)^{-1}, \frac{a}{2\mu}, \frac{b}{2\mu} \right\},$$

930 and  $h_i^0 = g_i^0$  for all  $i \in [n]$  in Algorithm 8. Then

$$\begin{aligned} & \mathbb{E} [f(x^T) - f^*] \\ & \leq (1 - \gamma\mu)^T \left( \Delta_0 + \frac{2\gamma}{b} \|h^0 - \nabla f(x^0)\|^2 + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \frac{1}{n} \sum_{i=1}^n \|h_i^0 - \nabla f_i(x^0)\|^2 \right) + \frac{20\sigma^2}{\mu n B'}. \end{aligned}$$

931 *Proof.* Let us fix constants  $\kappa, \eta, \nu, \rho \in [0, \infty)$  that we will define later. As in the proof of Theorem 11,  
932 we can get

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \kappa \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\ & - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{2\kappa\omega}{np_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \frac{2\eta\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\ & \quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\ & + (\gamma + \kappa(1-a)^2) \mathbb{E} [\|g^t - h^t\|^2] \\ & + \left( \kappa \frac{((2\omega+1)p_a - p_{aa})a^2}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ & + (\gamma + \nu(1-b)) \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\ & + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\ & + \left( \frac{2\nu b^2}{np_{\text{mega}}p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'}. \end{aligned}$$

933 Let us take  $\kappa = \frac{2\gamma}{a}$ , thus  $\gamma + \kappa(1-a)^2 \leq (1 - \frac{a}{2})\kappa$  and

$$\begin{aligned} & \mathbb{E} [f(x^{t+1})] + \frac{2\gamma}{a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \eta \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\ & + \nu \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\ & \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\ & - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{4\gamma\omega}{anp_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \frac{2\eta\omega}{p_a} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \end{aligned}$$

$$\begin{aligned}
& -\nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \mathbb{E} \left[ \|x^{t+1} - x^t\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma}{a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] \\
& + \left( \frac{2\gamma((2\omega+1)p_a - p_{aa})a}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + (\gamma + \nu(1-b)) \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'}.
\end{aligned}$$

934 Next, since  $a = \frac{p_a}{2\omega+1}$ , we have  $\left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \leq 1-a$ . We the choice  $\eta =$   
935  $\frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2}$ , we guarantee  $\frac{\gamma((2\omega+1)p_a - p_{aa})a}{np_a^2} + \eta \left( \frac{(2\omega+1-p_a)a^2}{p_a} + (1-a)^2 \right) \leq (1-\frac{a}{2})\eta$  and

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \nu \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} \left[ \|\nabla f(x^t)\|^2 \right] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
& \quad \left. - \nu \left( \frac{2L_\sigma^2}{np_a B} + \frac{2(p_a - p_{aa})\widehat{L}^2}{np_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} \left[ \|x^{t+1} - x^t\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] \\
& + \left( 1 - \frac{a}{2} \right) \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + (\gamma + \nu(1-b)) \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] \\
& + \left( \nu \frac{2(p_a - p_{aa})b^2}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{2\nu b^2}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'},
\end{aligned}$$

936 where simplified the term using  $p_{aa} \geq 0$ . Let us take  $\nu = \frac{2\gamma}{b}$  to obtain

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} \left[ \|g^{t+1} - h^{t+1}\|^2 \right] + \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} \left[ \|h^{t+1} - \nabla f(x^{t+1})\|^2 \right] + \rho \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} \left[ \|\nabla f(x^t)\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{4\gamma L_\sigma^2}{bn p_a B} + \frac{4\gamma(p_a - p_{aa})\widehat{L}^2}{bn p_a^2} \right) - \rho \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{a}{2}\right) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left(1 - \frac{a}{2}\right) \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] \\
& + \left( \frac{4\gamma(p_a - p_{aa})b}{np_a^2 p_{\text{mega}}} + \rho(1-b) \right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{4\gamma b}{np_{\text{mega}} p_a} + \frac{2\rho b^2}{p_a p_{\text{mega}}} \right) \frac{\sigma^2}{B'},
\end{aligned}$$

937 Next, we take  $\rho = \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}}$ , thus

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{4\gamma L_\sigma^2}{bn p_a B} + \frac{4\gamma(p_a - p_{aa})\widehat{L}^2}{bn p_a^2} \right) - \left( \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \right) \left( \frac{2L_\sigma^2}{p_a B} + \frac{2(1-p_a)\widehat{L}^2}{p_a} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{a}{2}\right) \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left(1 - \frac{a}{2}\right) \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \left( \frac{4\gamma b}{np_{\text{mega}} p_a} + \frac{16\gamma(p_a - p_{aa})b^2}{np_a^3 p_{\text{mega}}^2} \right) \frac{\sigma^2}{B'},
\end{aligned}$$

938 Since  $\frac{p_{\text{mega}} p_a}{2} \leq b \leq p_{\text{mega}} p_a$  and  $1 - p_a \leq 1 - \frac{p_{aa}}{p_a} \leq 1$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega+1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{2\gamma((2\omega+1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma(2\omega+1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) \right. \\
& \quad \left. - \left( \frac{8\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{8\gamma(p_a - p_{aa})\widehat{L}^2}{np_{\text{mega}} p_a^3} \right) - \left( \frac{16\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{16\gamma(1-p_a)\widehat{L}^2}{np_{\text{mega}} p_a^2} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2]
\end{aligned}$$

$$\begin{aligned}
& + \left(1 - \frac{a}{2}\right) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left(1 - \frac{a}{2}\right) \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{20\gamma\sigma^2}{nB'} \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& - \left( \frac{1}{2\gamma} - \frac{L}{2} - \frac{8\gamma(2\omega + 1)\omega}{np_a^2} \left( \frac{L_\sigma^2}{B} + \widehat{L}^2 \right) - \left( \frac{24\gamma L_\sigma^2}{np_{\text{mega}} p_a^2 B} + \frac{24\gamma \left(1 - \frac{p_{aa}}{p_a}\right) \widehat{L}^2}{np_{\text{mega}} p_a^2} \right) \right) \mathbb{E} [\|x^{t+1} - x^t\|^2] \\
& + \left(1 - \frac{a}{2}\right) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left(1 - \frac{a}{2}\right) \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{20\gamma\sigma^2}{nB'}.
\end{aligned}$$

939 Using Lemma 4 and the assumption about  $\gamma$ , we get

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + \left(1 - \frac{a}{2}\right) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + \left(1 - \frac{a}{2}\right) \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + \left(1 - \frac{b}{2}\right) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + \left(1 - \frac{b}{2}\right) \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{20\gamma\sigma^2}{nB'}.
\end{aligned}$$

940 Due to  $\gamma \leq \frac{a}{2\mu}$  and  $\gamma \leq \frac{b}{2\mu}$ , we have

$$\begin{aligned}
& \mathbb{E} [f(x^{t+1})] + \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^{t+1} - h^{t+1}\|^2] + \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^{t+1} - h_i^{t+1}\|^2 \right] \\
& + \frac{2\gamma}{b} \mathbb{E} [\|h^{t+1} - \nabla f(x^{t+1})\|^2] + \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^{t+1} - \nabla f_i(x^{t+1})\|^2 \right] \\
& \leq \mathbb{E} [f(x^t)] - \frac{\gamma}{2} \mathbb{E} [\|\nabla f(x^t)\|^2] \\
& + (1 - \gamma\mu) \frac{2\gamma(2\omega + 1)}{p_a} \mathbb{E} [\|g^t - h^t\|^2] + (1 - \gamma\mu) \frac{2\gamma((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\
& + (1 - \gamma\mu) \frac{2\gamma}{b} \mathbb{E} [\|h^t - \nabla f(x^t)\|^2] + (1 - \gamma\mu) \frac{8\gamma(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right] \\
& + \frac{20\gamma\sigma^2}{nB'}.
\end{aligned}$$

941 It is left to apply Lemma 11 with

$$\begin{aligned}\Psi^t &= \frac{2(2\omega + 1)}{p_a} \mathbb{E} \left[ \|g^t - h^t\|^2 \right] + \frac{2((2\omega + 1)p_a - p_{aa})}{np_a^2} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|g_i^t - h_i^t\|^2 \right] \\ &+ \frac{2}{b} \mathbb{E} \left[ \|h^t - \nabla f(x^t)\|^2 \right] + \frac{8(p_a - p_{aa})}{np_a^2 p_{\text{mega}}} \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \|h_i^t - \nabla f_i(x^t)\|^2 \right]\end{aligned}$$

942 and  $C = \frac{20\sigma^2}{nB'}$  to conclude the proof. □