

P9185 Project 1: Protocol of a Phase II MATIK Trial

Kate Colvin, Chhiring Lama, Emily Carter

Contents

I. Introduction	1
II. Methods	1
III. Results	2
3.1 Primary Objective 1	2
GLMM	2
3.2 Primary Objective 2	7
3.3 Secondary Objective 1	9
3.4 Secondary Objective 2	21
IV. Discussion	23
V. Conclusion	23
VI. Appendix	24

I. Introduction

II. Methods


Notation:

k :subject, $k = 1, \dots, 180$
 j :sequence, $j = 0, 1, 2$
 i :period, $i = 1, 2, 3$
 h :week, $h = 1, 2, 3, 4$
 π :period effect
 τ :treatment effect
 α :grouped treatment effects
 β :demographic effects
 λ :sequence/carryover effect
 γ :week effect
 b :subject-specific intercepts
 μ :fixed-effect intercept
 ε :residual error

III. Results

3.1 Primary Objective 1

Proportion of Adverse Events per period



Patient Distribution: 30 people in per treatment sequence Initial Observation: Most patients do not face adverse events over the course of 4 weeks in each of the consecutive periods.

add demographic data

demographic summary

Table 1: Distributional Summary of Predictor variables in the Study

Characteristic	period1			period2			period3		
	No N = 164	Yes N = 16	p-value	No N = 168	Yes N = 12	p-value	No N = 163	Yes N = 17	p-value
age	31 (25, 37)	42 (40, 44)	<0.001	32 (25, 38)	40 (37, 44)	<0.001	32 (25, 38)	38 (32, 42)	0.012

¹ Median (Q1, Q3)

² Wilcoxon rank sum test

GLMM

Model Selection

Characteristic	OR	95% CI	p-value
(Intercept)	0.00	0.00, 0.00	<0.001
Treatment			
Pill A	—	—	
Gel B	1.20	0.55, 2.64	0.6
Gel C	0.84	0.37, 1.92	0.7
Period			
period1	—	—	
period2	0.71	0.31, 1.64	0.4
period3	1.13	0.52, 2.46	0.8
Treatment Sequence			
ABC or BAC	—	—	
CAB or ACB	0.61	0.23, 1.62	0.3
BCA or CBA	0.84	0.35, 2.02	0.7
age	1.19	1.11, 1.27	<0.001
gender			
Female	—	—	
Male	0.90	0.43, 1.89	0.8
race			
black	—	—	
others	1.55	0.61, 3.95	0.4
white	1.74	0.65, 4.68	0.3

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

Table 2: LRT Between Model w/ and w/o Interaction Between Tx and Period

term	npar	AIC	BIC	logLik	minus2logL	statistic	df	p.value
glmm2	12	278.67	330.17	-127.34	254.67	NA	NA	NA
glmm1	16	281.34	350.01	-124.67	249.34	5.33	4	0.25

Table 3: LRT Between Model w/ and w/o Race, Gender and Carryover Effect

term	npar	AIC	BIC	logLik	minus2logL	statistic	df	p.value
glmm3	7	271.25	301.29	-128.62	257.25	NA	NA	NA
glmm2	12	278.67	330.17	-127.34	254.67	2.57	5	0.77

Characteristic	OR	95% CI	p-value
(Intercept)	0.00	0.00, 0.00	<0.001
Treatment			
Pill A	—	—	
Gel B	1.18	0.54, 2.58	0.7
Gel C	0.85	0.37, 1.94	0.7
Period			
period1	—	—	
period2	0.72	0.31, 1.65	0.4
period3	1.11	0.51, 2.40	0.8
age	1.19	1.12, 1.27	<0.001

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

Model Result (with Carryover Effect)


Final Model Result (without Carryover Effect)

Notation:

$$\widehat{\text{logit}}(P(AE_{ijk} = 1)) = b_k + \mu + \tau_i + \pi_j + \beta a_k,$$


$$i = 1, 2, 3, j = 1, 2, 3, k = (1, 2, \dots, 180),$$

$$b_k \sim N(0, \sigma_b^2)$$




AE Prediction

Distribution of OR for Adverse Events




Predicted probabilities of adverse event



Model Diagnostics


Posterior Predictive Check

Model-predicted intervals should include observed data points




Binned Residuals

Points should be within error bounds




Influential Observations

Points should be inside the contour lines




Collinearity

High collinearity (VIF) may inflate parameter uncertainty




Distribution of Quantile Residuals

Dots should fall along the line




Normality of Random Effects (ptid)



Dots should be plotted along the line



3.2 Primary Objective 2

Individual Patient Adherence Over Time, by Treatment





Model

\$\$

$$Y_{hik} \sim \text{Binomial}(n = 7, p_{hik}), \quad (1)$$

$$\text{logit}(p_{hik}) = \mu + b_k + \pi_i + \tau_i + \lambda_j + \gamma * h, \quad (2)$$

$$b_k \sim \mathcal{N}(0, \sigma_b^2) \quad (3)$$

\$\$

Table 4: GLMM Results of Treatment on Adherence

Characteristic	OR	95% CI	p-value
Treatment			
Pill A	—	—	
Gel B	0.14	0.12, 0.17	<0.001
Gel C	0.21	0.17, 0.24	<0.001
Week	0.73	0.69, 0.76	<0.001
Period			
1	—	—	
2	1.15	1.00, 1.32	0.051
3	1.05	0.92, 1.20	0.5

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

Table 5: Table 1. Baseline Characteristics

Characteristic	Overall N = 180 ¹	ABC N = 30 ¹	ACB N = 30 ¹	BAC N = 30 ¹	BCA N = 30 ¹
Age (years)	32 (8) [18, 45]	33 (8) [19, 44]	31 (7) [18, 42]	31 (7) [20, 44]	33 (8) [19, 45]
Race					
Black	64 (36%)	12 (40%)	13 (43%)	12 (40%)	10 (33%)
White	52 (29%)	8 (27%)	9 (30%)	7 (23%)	7 (23%)
Other	64 (36%)	10 (33%)	8 (27%)	11 (37%)	13 (43%)
Female	81 (45%)	13 (43%)	11 (37%)	16 (53%)	14 (47%)

¹Mean (SD) [Min, Max]; n (%)


²Kruskal-Wallis rank sum test; Pearson's Chi-squared test

3.3 Secondary Objective 1

preliminary assessment and comparison of systemic and local Pharmacokinetics (PK) of Pill A, Gel B, and Gel C and the correlation of PK with adherence measures and the occurrence of adverse events

Blood PK

$$Y_{ik} = \mu + b_k + \pi_i + \tau_i + \lambda_i + \alpha_{ik} + \beta_{ik} + \varepsilon_{ik}, \quad b_k \sim N(0, \sigma_b^2), \quad \varepsilon_{ik} \sim N(0, \sigma^2)$$



Data: merged

```

## Models:
## model_blood: bloodVL_change ~ treatment + overall_adhere + overall_safety + period + (1 | ptid)
## model_large_blood: bloodVL_change ~ treatment + overall_adhere + overall_safety + sequence_ind + per
##
##          npar    AIC    BIC  logLik -2*log(L)  Chisq Df Pr(>Chisq)
## model_blood      9 4863.1 4901.7 -2422.6    4845.1
## model_large_blood 11 4866.5 4913.7 -2422.2    4844.5 0.6124  2    0.7362

## Linear mixed model fit by REML. t-tests use Satterthwaite's method [
## lmerModLmerTest]
## Formula: bloodVL_change ~ treatment + overall_adhere + overall_safety +
##          period + (1 | ptid)
##          Data: merged
##
## REML criterion at convergence: 4827.4
##
## Scaled residuals:
##          Min         1Q      Median         3Q          Max
## -3.5936 -0.5795  0.0153  0.6127  3.0873
##
## Random effects:
##      Groups      Name      Variance Std.Dev.
##      ptid      (Intercept)  23.09      4.805
##      Residual              445.60     21.109
## Number of obs: 540, groups:  ptid, 180
##
## Fixed effects:
##
##              Estimate Std. Error      df t value
## (Intercept)      73.4552      6.7058 419.7422  10.954
## treatmentGel B      -6.7916      2.3555 417.1354  -2.883
## treatmentGel C      -9.0880      2.2967 391.5595  -3.957
## overall_adhere        0.9365      0.1882 269.4894   4.975
## overall_safetyNo Adverse Event 26.4624      3.4675 521.9424   7.632
## periodperiod2      -2.3295      2.2271 356.0129  -1.046
## periodperiod3      -1.5433      2.2252 355.4545  -0.694
##
##              Pr(>|t|)
## (Intercept)      < 0.0000000000000002 ***
## treatmentGel B          0.00414 **
## treatmentGel C      0.000090106833765 ***
## overall_adhere      0.000001161748202 ***
## overall_safetyNo Adverse Event 0.000000000000111 ***
## periodperiod2          0.29629
## periodperiod3          0.48842
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) trtmGB trtmGC ovrll_ ov_NAE prdpr2
## treatmntG1B -0.437
## treatmntG1C -0.359  0.538
## overall_dhr -0.832  0.328  0.247
## ovrll_sfNAE -0.616  0.080  0.032  0.196
## periodperd2 -0.129 -0.009 -0.006 -0.026 -0.038
## periodperd3 -0.173  0.001  0.001  0.004  0.009  0.499
##
## treatment emmean SE df lower.CL upper.CL

```

Comparison	Beta	95% CI	p-value
Intercept (Pill A, Period 1, Sequence 0, No Adherence or AE)	73.5	60.3, 86.6	<0.001
Treatment			
Gel B vs Pill A	-6.8	-11.4, -2.2	0.004
Gel C vs Pill A	-9.1	-13.6, -4.6	<0.001
Additional Day of Adherence	0.9	0.6, 1.3	<0.001
Adverse Event			
No Adverse Event	26.5	19.7, 33.3	<0.001
Period			
Period 2 vs Period 1	-2.3	-6.7, 2.1	0.3
Period 3 vs Period 1	-1.5	-5.9, 2.8	0.5

Abbreviation: CI = Confidence Interval

```
## Pill A      107.9 2.27 525      103.4      112
## Gel B       101.1 2.13 523       96.9      105
## Gel C       98.8 2.18 526       94.5      103
##
## Results are averaged over the levels of: overall_safety, period
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## contrast      estimate    SE  df t.ratio p.value
## Pill A - Gel B      6.79 2.36 417   2.882 0.0115
## Pill A - Gel C      9.09 2.30 391   3.956 0.0003
## Gel B - Gel C       2.30 2.24 359   1.027 0.5602
##
## Results are averaged over the levels of: overall_safety, period
## Degrees-of-freedom method: kenward-roger
## P value adjustment: tukey method for comparing a family of 3 estimates







## overall_adhere emmean    SE  df lower.CL upper.CL
##              24      103 1.75 411    99.2    106
##
## Results are averaged over the levels of: treatment, overall_safety, period
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## overall_safety emmean    SE  df lower.CL upper.CL
## Adverse Event   89.4 3.34 500    82.8    95.9
## No Adverse Event 115.9 1.02 196   113.8   117.9
##
## Results are averaged over the levels of: treatment, period
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## period emmean    SE  df lower.CL upper.CL
## period1  104 2.16 527    99.7    108
## period2   102 2.21 527    97.2    106
## period3   102 2.14 527    98.2    107
##
## Results are averaged over the levels of: treatment, overall_safety
```


```
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## contrast      estimate    SE  df t.ratio p.value
## period1 - period2    2.330 2.23 355   1.046  0.5484
## period1 - period3    1.543 2.23 355   0.694  0.7674
## period2 - period3   -0.786 2.23 356  -0.353  0.9337
##
## Results are averaged over the levels of: treatment, overall_safety
## Degrees-of-freedom method: kenward-roger
## P value adjustment: tukey method for comparing a family of 3 estimates
```



Skin PK

Linear Model



Analysis of Variance Table

```
##
## Model 1: skinVL_change ~ treatment + overall_adhere + overall_safety +
##   period
## Model 2: skinVL_change ~ treatment + overall_adhere + overall_safety +
##   sequence_ind + period
##   Res.Df    RSS Df Sum of Sq    F Pr(>F)
## 1     533 571160
## 2     531 568220  2    2940.4 1.3739  0.254

##
## Call:
## lm(formula = skinVL_change ~ treatment + overall_adhere + overall_safety +
##   period, data = merged)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -112.388  -22.128    1.035   20.921  115.662
##
## Coefficients:
##              Estimate Std. Error t value      Pr(>|t|)
## (Intercept)      105.6003      9.9680  10.594 < 0.0000000000000002
## treatmentGel B      -24.4131      3.6314  -6.723  0.000000000000046
## treatmentGel C     -36.9951      3.5493 -10.423 < 0.0000000000000002
## overall_adhere       1.8154      0.2753   6.594  0.00000000000103
## overall_safetyNo Adverse Event  55.7940      5.2166  10.696 < 0.0000000000000002
## periodperiod2        0.8567      3.4535   0.248      0.804
## periodperiod3        1.6225      3.4507   0.470      0.638
##
## (Intercept)          ***
## treatmentGel B        ***
## treatmentGel C        ***
## overall_adhere        ***
## overall_safetyNo Adverse Event ***
## periodperiod2
## periodperiod3
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 32.74 on 533 degrees of freedom
## Multiple R-squared:  0.3581, Adjusted R-squared:  0.3508
## F-statistic: 49.55 on 6 and 533 DF, p-value: < 0.00000000000000022

##   treatment emmean   SE  df lower.CL upper.CL
## Pill A      178 3.42 533     171     185
## Gel B       154 3.20 533     147     160
## Gel C       141 3.29 533     135     147
##
## Results are averaged over the levels of: overall_safety, period
## Confidence level used: 0.95

##   contrast      estimate    SE  df t.ratio p.value
## Pill A - Gel B      24.4 3.63 533   6.723 <0.0001
## Pill A - Gel C      37.0 3.55 533  10.423 <0.0001
## Gel B - Gel C       12.6 3.47 533   3.631  0.0009
##
```

Comparison	Beta	95% CI	p-value
Intercept (Pill A, Period 1, Sequence 0, No Adherence or AE)	105.6	86.0, 125.2	<0.001
Treatment			
Gel B vs Pill A	-24.4	-31.5, -17.3	<0.001
Gel C vs Pill A	-37.0	-44.0, -30.0	<0.001
Additional Day of Adherence	1.8	1.3, 2.4	<0.001
Adverse Event			
No Adverse Event	55.8	45.5, 66.0	<0.001
Period			
Period 2 vs Period 1	0.9	-5.9, 7.6	0.8
Period 3 vs Period 1	1.6	-5.2, 8.4	0.6

Abbreviation: CI = Confidence Interval


```
## Results are averaged over the levels of: overall_safety, period
## P value adjustment: tukey method for comparing a family of 3 estimates

## overall_adhere emmean SE df lower.CL upper.CL
##          24      158 2.59 533      152      163
##
## Results are averaged over the levels of: treatment, overall_safety, period
## Confidence level used: 0.95

## overall_safety emmean SE df lower.CL upper.CL
## Adverse Event      130 4.99 533      120      139
## No Adverse Event    185 1.47 533      183      188
##
## Results are averaged over the levels of: treatment, period
## Confidence level used: 0.95

## period emmean SE df lower.CL upper.CL
## period1      157 3.25 533      150      163
## period2      158 3.33 533      151      164
## period3      158 3.23 533      152      165
##
## Results are averaged over the levels of: treatment, overall_safety
## Confidence level used: 0.95

## contrast estimate SE df t.ratio p.value
## period1 - period2 -0.857 3.45 533 -0.248 0.9666
## period1 - period3 -1.623 3.45 533 -0.470 0.8853
## period2 - period3 -0.766 3.45 533 -0.222 0.9733
##
## Results are averaged over the levels of: treatment, overall_safety
## P value adjustment: tukey method for comparing a family of 3 estimates
```

Linear Mixed Model

```
## Data: merged
## Models:
## model_skin: skinVL_change ~ treatment + overall_adhere + overall_safety + period + (1 | ptid)
## model_large_skin: skinVL_change ~ treatment + overall_adhere + overall_safety + sequence_ind + period
##               npar    AIC    BIC logLik -2*log(L)  Chisq Df Pr(>Chisq)
## model_skin      9 5310.9 5349.6 -2646.5   5292.9
## model_large_skin 11 5312.1 5359.4 -2645.1   5290.1 2.7872 2    0.2482

## Linear mixed model fit by REML. t-tests use Satterthwaite's method [
## lmerModLmerTest]
## Formula: skinVL_change ~ treatment + overall_adhere + overall_safety +
##         period + (1 | ptid)
## Data: merged
##
## REML criterion at convergence: 5269.4
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -3.4333 -0.6760  0.0316  0.6391  3.5333
##
## Random effects:
## Groups   Name                Variance Std.Dev.
## ptid     (Intercept)          0        0.00
## Residual                    1072      32.74
## Number of obs: 540, groups:  ptid, 180
##
```

Comparison	Beta	95% CI	p-value
Intercept (Pill A, Period 1, Sequence 0, No Adherence or AE)	105.6	86.0, 125.2	<0.001
Treatment			
Gel B vs Pill A	-24.4	-31.5, -17.3	<0.001
Gel C vs Pill A	-37.0	-44.0, -30.0	<0.001
Additional Day of Adherence	1.8	1.3, 2.4	<0.001
Adverse Event			
No Adverse Event	55.8	45.5, 66.0	<0.001
Period			
Period 2 vs Period 1	0.9	-5.9, 7.6	0.8
Period 3 vs Period 1	1.6	-5.2, 8.4	0.6

Abbreviation: CI = Confidence Interval

```
## Fixed effects:
##               Estimate Std. Error      df t value
## (Intercept)    105.6003     9.9680 533.0000   10.594
## treatmentGel B   -24.4131     3.6314 533.0000    -6.723
## treatmentGel C   -36.9951     3.5493 533.0000   -10.423
## overall_adhere     1.8154     0.2753 533.0000     6.594
## overall_safetyNo Adverse Event 55.7940     5.2166 533.0000   10.696
## periodperiod2     0.8567     3.4535 533.0000     0.248
## periodperiod3     1.6225     3.4507 533.0000     0.470
##               Pr(>|t|)
## (Intercept)      < 0.0000000000000002 ***
## treatmentGel B      0.0000000000046 ***
## treatmentGel C      < 0.0000000000000002 ***
## overall_adhere      0.000000000103 ***
## overall_safetyNo Adverse Event < 0.0000000000000002 ***
## periodperiod2           0.804
## periodperiod3           0.638
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##      (Intr) trtmGB trtmGC ovrll_ ov_NAE prdpr2
## treatmntG1B -0.429
## treatmntG1C -0.354  0.534
## overall_dhr -0.826  0.311  0.234
## ovrll_sfNAE -0.627  0.079  0.033  0.206
## periodperd2 -0.137 -0.008 -0.005 -0.025 -0.037
## periodperd3 -0.180  0.001  0.001  0.004  0.009  0.499
## optimizer (nloptwrap) convergence code: 0 (OK)
## boundary (singular) fit: see help('isSingular')
##
## treatment emmean   SE   df lower.CL upper.CL
## Pill A          178 3.43 525      171      185
## Gel B           154 3.21 525      147      160
## Gel C           141 3.29 527      135      147
##
## Results are averaged over the levels of: overall_safety, period
```

```

## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## contrast      estimate    SE  df t.ratio p.value
## Pill A - Gel B      24.4 3.63 412   6.721 <0.0001
## Pill A - Gel C      37.0 3.55 388  10.422 <0.0001
## Gel B - Gel C       12.6 3.47 359   3.631  0.0009
##
## Results are averaged over the levels of: overall_safety, period
## Degrees-of-freedom method: kenward-roger
## P value adjustment: tukey method for comparing a family of 3 estimates

## overall_adhere emmean  SE  df lower.CL upper.CL
##              24    158 2.6 401      152      163
##
## Results are averaged over the levels of: treatment, overall_safety, period
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## overall_safety  emmean    SE  df lower.CL upper.CL
## Adverse Event      130 5.00 485      120      139
## No Adverse Event    185 1.47 196      182      188
##
## Results are averaged over the levels of: treatment, period
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95







## period  emmean    SE  df lower.CL upper.CL
## period1  157 3.25 528      150      163
## period2  158 3.34 528      151      164
## period3  158 3.23 528      152      165
##
## Results are averaged over the levels of: treatment, overall_safety
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95

## contrast      estimate    SE  df t.ratio p.value
## period1 - period2  -0.857 3.45 356  -0.248  0.9666
## period1 - period3  -1.623 3.45 355  -0.470  0.8853
## period2 - period3  -0.766 3.45 356  -0.222  0.9733
##
## Results are averaged over the levels of: treatment, overall_safety
## Degrees-of-freedom method: kenward-roger
## P value adjustment: tukey method for comparing a family of 3 estimates

```


Comparison	Blood Viral Load			Skin	
	Beta	95% CI	p-value	Beta	95% CI
Intercept (Pill A, Period 1, Sequence 0, No Adherence or AE)	73.5	60.3, 86.6	<0.001	105.6	86.6, 124.6
Treatment					
Gel B vs Pill A	-6.8	-11.4, -2.2	0.004	-24.4	-31.1, -17.7
Gel C vs Pill A	-9.1	-13.6, -4.6	<0.001	-37.0	-44.1, -30.0
Additional Day of Adherence	0.9	0.6, 1.3	<0.001	1.8	1.3, 2.3
Adverse Event					
No Adverse Event	26.5	19.7, 33.3	<0.001	55.8	45.1, 66.5
Period					
Period 2 vs Period 1	-2.3	-6.7, 2.1	0.3	0.9	-5.1, 6.9
Period 3 vs Period 1	-1.5	-5.9, 2.8	0.5	1.6	-5.1, 8.3


Abbreviation: CI = Confidence Interval



Combined Table

3.4 Secondary Objective 2





Model

\$\$

$$Y_{hik} \sim \text{Binomial}(n = 7, p_{hik}), \quad (4)$$


$$\text{logit}(p_{hik}) = \mu + b_k + \pi_i + \lambda_j + \gamma * h + \beta_d * d_k + \beta_r * regimen + \beta_p * product \quad (5)$$

$$+ \beta_{dr} * d_k * regimen + \beta_{dp} * d_k * product \quad (6)$$

$$b_k \sim \mathcal{N}(0, \sigma_b^2) \quad (7)$$

\$\$

DHARMa residual



IV. Discussion

V. Conclusion

VI. Appendix

Table 6: Interaction Results of Demographic Variables on Adherence

Characteristic	OR	95% CI	p-value
Age	1.21	1.18, 1.24	<0.001
Product			
Gel	—	—	
Pill	4.35	2.98, 6.35	<0.001
Gender			
Male	—	—	
Female	1.36	0.95, 1.95	0.10
Race			
white	—	—	
black	1.02	0.65, 1.60	>0.9
others	0.97	0.62, 1.53	0.9
Regimen			
once	—	—	
three	0.76	0.58, 1.00	0.050
Period			
1	—	—	
2	1.14	0.99, 1.32	0.073
3	1.03	0.89, 1.18	0.7
Week	0.73	0.69, 0.76	<0.001
Age * Product			
Age * Pill	1.00	0.97, 1.03	>0.9
Product * Gender			
Pill * Female	1.00	0.71, 1.40	>0.9
Product * Race			
Pill * black	1.23	0.81, 1.86	0.3
Pill * others	1.05	0.69, 1.61	0.8
Age * Regimen			
Age * three	1.02	1.00, 1.04	0.074
Gender * Regimen			
Female * three	0.89	0.69, 1.15	0.4
Race * Regimen			
black * three	1.14	0.84, 1.56	0.4
others * three	1.02	0.74, 1.41	0.9

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

Table 7: Main Effect Results of Demographic Variables on Adherence

Characteristic	OR	95% CI	p-value
Age	1.22	1.19, 1.25	<0.001
Gender			
Male	—	—	
Female	1.29	0.92, 1.80	0.14
Race			
white	—	—	
black	1.11	0.73, 1.69	0.6
others	0.99	0.65, 1.50	>0.9
Regimen			
once	—	—	
three	0.71	0.63, 0.80	<0.001
Product			
Gel	—	—	
Pill	4.84	4.12, 5.70	<0.001
Period			
1	—	—	
2	1.14	0.99, 1.32	0.065
3	1.04	0.91, 1.20	0.6
Week	0.73	0.69, 0.76	<0.001

Abbreviations: CI = Confidence Interval, OR = Odds Ratio