Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Oldest Child

Age reference: 10 October 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,181	0.005 [0.026] N=1,278	0.012 [0.026] N=1,179	-0.001 [0.028] N=1,056	-0.004 [0.029] N=907	-0.015 [0.034] N=749
Vegetables consumed at least weekly	0.896 P=1,142	0.013 [0.031] N=1,278	0.002 [0.032] N=1,179	-0.016 [0.032] N=1,056	-0.006 [0.033] N=907	-0.031 [0.039] N=749
Meat fish or poultry consumed at least weekly	0.767 P=986	0.065 [0.052] N=1,278	0.078 [0.054] N=1,179	0.069 [0.059] N=1,056	0.068 [0.066] N=907	0.057 [0.071] N=749
Legumes consumed at least weekly	0.770 P=989	-0.036 [0.046] N=1,278	-0.060 [0.049] N=1,179	-0.058 [0.049] N=1,056	-0.048 [0.055] N=907	-0.049 [0.065] N=749
Animal products consumed at least weekly	0.904 P=1,158	0.026 [0.037] N=1,278	0.022 [0.040] N=1,179	-0.008 [0.043] N=1,056	-0.001 [0.047] N=907	-0.025 [0.052] N=749
Sweets consumed at least weekly	0.556 P=707	-0.036 [0.049] N=1,278	-0.034 [0.051] N=1,179	-0.022 [0.055] N=1,056	-0.039 [0.060] N=907	-0.005 [0.068] N=749
Snack foods consumed at least weekly	0.460 P=581	-0.015 [0.045] N=1,278	-0.018 [0.047] N=1,179	0.004 [0.051] N=1,056	0.005 [0.055] N=907	0.006 [0.057] N=749
Packaged soft drinks consumed at least weekly	0.264 P=329	-0.029 [0.047] N=1,278	-0.049 [0.049] N=1,179	-0.051 [0.051] N=1,056	-0.026 [0.054] N=907	-0.044 [0.062] N=749
Grains consumed at least weekly	0.865 P=1,107	0.032 [0.038] N=1,278	0.029 [0.039] N=1,179	0.020 [0.041] N=1,056	0.031 [0.045] N=907	0.024 [0.050] N=749
Child was ever breastfed	0.969 P=1,447	-0.016 [0.013] N=1,498	-0.015 [0.013] N=1,385	-0.019 [0.015] N=1,237	-0.020 [0.017] N=1,057	-0.023 [0.016] N=877

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.018 [0.040] N=1,082	0.031 [0.041] N=983	0.008 [0.046] N=860	-0.009 [0.053] N=711	-0.039 [0.071] N=553
Vegetables consumed at least weekly	0.028 [0.048] N=1,082	0.011 [0.051] N=983	-0.028 [0.054] N=860	-0.018 [0.063] N=711	-0.108 [0.090] N=553
Meat fish or poultry consumed at least weekly	0.042 [0.065] N=1,082	0.052 [0.068] N=983	0.026 [0.078] N=860	-0.022 [0.089] N=711	-0.098 [0.088] N=553
Legumes consumed at least weekly	-0.026 [0.069] N=1,082	-0.075 [0.074] N=983	-0.070 [0.074] N=860	-0.079 [0.088] N=711	-0.081 [0.125] N=553
Animal products consumed at least weekly	0.049 [0.051] N=1,082	0.046 [0.057] N=983	-0.014 [0.064] N=860	0.003 [0.078] N=711	-0.063 [0.104] N=553
Sweets consumed at least weekly	-0.067 [0.064] N=1,082	-0.067 [0.067] N=983	-0.054 [0.077] N=860	-0.130 [0.086] N=711	-0.061 [0.121] N=553
Snack foods consumed at least weekly	-0.043 [0.062] N=1,082	-0.053 [0.066] N=983	-0.022 [0.077] N=860	-0.040 [0.097] N=711	-0.080 [0.118] N=553
Packaged soft drinks consumed at least weekly	-0.035 [0.067] N=1,082	-0.072 [0.072] N=983	-0.079 [0.078] N=860	-0.048 [0.091] N=711	-0.096 [0.128] N=553
Grains consumed at least weekly	0.074 [0.058] N=1,082	0.073 [0.064] N=983	0.061 [0.071] N=860	0.099 [0.088] N=711	0.102 [0.125] N=553
Child was ever breastfed	-0.009 [0.019] N=1,265	-0.004 [0.022] N=1,152	-0.011 [0.027] N=1,004	-0.015 [0.035] N=824	-0.014 [0.041] N=644

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,181	0.005 [0.026] N=1,278	0.012 [0.026] N=1,179	-0.000 [0.028] N=1,056	-0.004 [0.029] N=907	-0.015 [0.034] N=749
Vegetables consumed at least weekly	0.896 P=1,142	0.013 [0.031] N=1,278	0.002 [0.031] N=1,179	-0.015 [0.032] N=1,056	-0.006 [0.033] N=907	-0.032 [0.038] N=749
Meat fish or poultry consumed at least weekly	0.767 P=986	0.065 [0.051] N=1,278	0.078 [0.054] N=1,179	0.069 [0.059] N=1,056	0.068 [0.066] N=907	0.057 [0.072] N=749
Legumes consumed at least weekly	0.770 P=989	-0.036 [0.047] N=1,278	-0.060 [0.049] N=1,179	-0.057 [0.049] N=1,056	-0.047 [0.055] N=907	-0.049 [0.066] N=749
Animal products consumed at least weekly	0.904 P=1,158	0.027 [0.037] N=1,278	0.022 [0.039] N=1,179	-0.006 [0.042] N=1,056	-0.000 [0.046] N=907	-0.025 [0.051] N=749
Sweets consumed at least weekly	0.556 P=707	-0.036 [0.049] N=1,278	-0.034 [0.051] N=1,179	-0.021 [0.055] N=1,056	-0.039 [0.060] N=907	-0.005 [0.068] N=749
Snack foods consumed at least weekly	0.460 P=581	-0.015 [0.045] N=1,278	-0.018 [0.048] N=1,179	0.006 [0.051] N=1,056	0.005 [0.055] N=907	0.005 [0.058] N=749
Packaged soft drinks consumed at least weekly	0.264 P=329	-0.029 [0.047] N=1,278	-0.049 [0.049] N=1,179	-0.051 [0.051] N=1,056	-0.026 [0.054] N=907	-0.044 [0.062] N=749
Grains consumed at least weekly	0.865 P=1,107	0.032 [0.038] N=1,278	0.029 [0.039] N=1,179	0.021 [0.041] N=1,056	0.031 [0.046] N=907	0.024 [0.051] N=749
Child was ever breastfed	0.969 P=1,447	-0.016 [0.013] N=1,498	-0.015 [0.013] N=1,385	-0.019 [0.015] N=1,237	-0.019 [0.017] N=1,057	-0.022 [0.016] N=877

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.018 [0.040] N=1,082	0.032 [0.041] N=983	0.009 [0.046] N=860	-0.009 [0.053] N=711	-0.039 [0.072] N=553
Vegetables consumed at least weekly	0.028 [0.048] N=1,082	0.012 [0.051] N=983	-0.026 [0.054] N=860	-0.017 [0.063] N=711	-0.107 [0.090] N=553
Meat fish or poultry consumed at least weekly	0.041 [0.065] N=1,082	0.052 [0.068] N=983	0.026 [0.078] N=860	-0.021 [0.089] N=711	-0.097 [0.087] N=553
Legumes consumed at least weekly	-0.026 [0.069] N=1,082	-0.074 [0.074] N=983	-0.067 [0.074] N=860	-0.077 [0.089] N=711	-0.080 [0.125] N=553
Animal products consumed at least weekly	0.052 [0.051] N=1,082	0.049 [0.056] N=983	-0.008 [0.064] N=860	0.007 [0.077] N=711	-0.061 [0.102] N=553
Sweets consumed at least weekly	-0.067 [0.064] N=1,082	-0.066 [0.067] N=983	-0.053 [0.077] N=860	-0.129 [0.086] N=711	-0.061 [0.121] N=553
Snack foods consumed at least weekly	-0.040 [0.063] N=1,082	-0.051 [0.067] N=983	-0.013 [0.077] N=860	-0.036 [0.097] N=711	-0.078 [0.117] N=553
Packaged soft drinks consumed at least weekly	-0.035 [0.067] N=1,082	-0.072 [0.072] N=983	-0.078 [0.078] N=860	-0.047 [0.091] N=711	-0.097 [0.129] N=553
Grains consumed at least weekly	0.075 [0.058] N=1,082	0.074 [0.064] N=983	0.065 [0.071] N=860	0.100 [0.088] N=711	0.103 [0.124] N=553
Child was ever breastfed	-0.009 [0.019] N=1,265	-0.004 [0.022] N=1,152	-0.011 [0.027] N=1,004	-0.016 [0.035] N=824	-0.014 [0.041] N=644

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,181	-0.002 [0.026] N=1,278	0.003 [0.027] N=1,179	-0.008 [0.028] N=1,056	-0.009 [0.030] N=907	-0.025 [0.035] N=749
Vegetables consumed at least weekly	0.896 P=1,142	0.008 [0.031] N=1,278	-0.005 [0.031] N=1,179	-0.023 [0.031] N=1,056	-0.011 [0.033] N=907	-0.041 [0.038] N=749
Meat fish or poultry consumed at least weekly	0.767 P=986	0.044 [0.050] N=1,278	0.052 [0.052] N=1,179	0.044 [0.058] N=1,056	0.052 [0.064] N=907	0.031 [0.069] N=749
Legumes consumed at least weekly	0.770 P=989	-0.042 [0.046] N=1,278	-0.067 [0.048] N=1,179	-0.063 [0.049] N=1,056	-0.050 [0.054] N=907	-0.052 [0.065] N=749
Animal products consumed at least weekly	0.904 P=1,158	0.022 [0.037] N=1,278	0.016 [0.039] N=1,179	-0.013 [0.043] N=1,056	-0.005 [0.047] N=907	-0.036 [0.052] N=749
Sweets consumed at least weekly	0.556 P=707	-0.038 [0.049] N=1,278	-0.037 [0.051] N=1,179	-0.023 [0.056] N=1,056	-0.040 [0.060] N=907	-0.013 [0.067] N=749
Snack foods consumed at least weekly	0.460 P=581	-0.011 [0.045] N=1,278	-0.018 [0.047] N=1,179	0.008 [0.051] N=1,056	0.008 [0.054] N=907	0.003 [0.056] N=749
Packaged soft drinks consumed at least weekly	0.264 P=329	-0.018 [0.046] N=1,278	-0.037 [0.048] N=1,179	-0.037 [0.050] N=1,056	-0.017 [0.053] N=907	-0.033 [0.060] N=749
Grains consumed at least weekly	0.865 P=1,107	0.027 [0.038] N=1,278	0.025 [0.040] N=1,179	0.016 [0.042] N=1,056	0.027 [0.046] N=907	0.021 [0.051] N=749
Child was ever breastfed	0.969 P=1,447	-0.018 [0.013] N=1,498	-0.017 [0.014] N=1,385	-0.021 [0.015] N=1,237	-0.020 [0.017] N=1,057	-0.025 [0.017] N=877

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.016 [0.039] N=1,082	0.029 [0.041] N=983	0.010 [0.045] N=860	0.010 [0.054] N=711	-0.028 [0.073] N=553
Vegetables consumed at least weekly	0.028 [0.047] N=1,082	0.009 [0.050] N=983	-0.027 [0.053] N=860	-0.002 [0.063] N=711	-0.092 [0.090] N=553
Meat fish or poultry consumed at least weekly	0.031 [0.064] N=1,082	0.036 [0.066] N=983	0.012 [0.079] N=860	-0.003 [0.090] N=711	-0.088 [0.088] N=553
Legumes consumed at least weekly	-0.027 [0.069] N=1,082	-0.078 [0.073] N=983	-0.067 [0.074] N=860	-0.065 [0.089] N=711	-0.068 [0.128] N=553
Animal products consumed at least weekly	0.053 [0.049] N=1,082	0.047 [0.055] N=983	-0.009 [0.063] N=860	0.020 [0.078] N=711	-0.043 [0.102] N=553
Sweets consumed at least weekly	-0.064 [0.065] N=1,082	-0.063 [0.068] N=983	-0.042 [0.078] N=860	-0.113 [0.087] N=711	-0.056 [0.125] N=553
Snack foods consumed at least weekly	-0.034 [0.062] N=1,082	-0.049 [0.067] N=983	-0.004 [0.076] N=860	-0.020 [0.097] N=711	-0.071 [0.115] N=553
Packaged soft drinks consumed at least weekly	-0.024 [0.065] N=1,082	-0.057 [0.071] N=983	-0.059 [0.076] N=860	-0.035 [0.089] N=711	-0.087 [0.124] N=553
Grains consumed at least weekly	0.071 [0.057] N=1,082	0.072 [0.064] N=983	0.063 [0.071] N=860	0.107 [0.088] N=711	0.104 [0.124] N=553
Child was ever breastfed	-0.006 [0.019] N=1,265	-0.003 [0.022] N=1,152	-0.007 [0.026] N=1,004	-0.003 [0.034] N=824	-0.012 [0.040] N=644

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,181	-0.006 [0.027] N=1,278	-0.000 [0.027] N=1,179	-0.016 [0.028] N=1,056	-0.018 [0.031] N=907	-0.036 [0.035] N=749
Vegetables consumed at least weekly	0.896 P=1,142	0.006 [0.031] N=1,278	-0.005 [0.031] N=1,179	-0.023 [0.031] N=1,056	-0.013 [0.033] N=907	-0.041 [0.039] N=749
Meat fish or poultry consumed at least weekly	0.767 P=986	0.036 [0.051] N=1,278	0.046 [0.053] N=1,179	0.033 [0.059] N=1,056	0.034 [0.065] N=907	0.005 [0.071] N=749
Legumes consumed at least weekly	0.770 P=989	-0.046 [0.047] N=1,278	-0.071 [0.050] N=1,179	-0.071 [0.050] N=1,056	-0.058 [0.057] N=907	-0.066 [0.068] N=749
Animal products consumed at least weekly	0.904 P=1,158	0.018 [0.038] N=1,278	0.013 [0.040] N=1,179	-0.017 [0.043] N=1,056	-0.012 [0.047] N=907	-0.045 [0.051] N=749
Sweets consumed at least weekly	0.556 P=707	-0.042 [0.049] N=1,278	-0.043 [0.051] N=1,179	-0.034 [0.055] N=1,056	-0.048 [0.060] N=907	-0.025 [0.068] N=749
Snack foods consumed at least weekly	0.460 P=581	-0.017 [0.044] N=1,278	-0.023 [0.047] N=1,179	-0.004 [0.051] N=1,056	-0.004 [0.055] N=907	-0.011 [0.057] N=749
Packaged soft drinks consumed at least weekly	0.264 P=329	-0.018 [0.046] N=1,278	-0.041 [0.048] N=1,179	-0.042 [0.050] N=1,056	-0.020 [0.053] N=907	-0.035 [0.060] N=749
Grains consumed at least weekly	0.865 P=1,107	0.028 [0.038] N=1,278	0.025 [0.040] N=1,179	0.016 [0.042] N=1,056	0.026 [0.046] N=907	0.021 [0.053] N=749
Child was ever breastfed	0.969 P=1,447	-0.021 [0.014] N=1,498	-0.020 [0.015] N=1,385	-0.025 [0.016] N=1,237	-0.024 [0.018] N=1,057	-0.029 [0.018] N=877

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.016 [0.040] N=1,082	0.032 [0.042] N=983	0.008 [0.047] N=860	-0.004 [0.054] N=711	-0.056 [0.075] N=553
Vegetables consumed at least weekly	0.025 [0.048] N=1,082	0.009 [0.051] N=983	-0.033 [0.054] N=860	-0.022 [0.062] N=711	-0.130 [0.092] N=553
Meat fish or poultry consumed at least weekly	0.027 [0.063] N=1,082	0.038 [0.065] N=983	0.009 [0.079] N=860	-0.007 [0.087] N=711	-0.127 [0.093] N=553
Legumes consumed at least weekly	-0.027 [0.069] N=1,082	-0.075 [0.073] N=983	-0.066 [0.073] N=860	-0.056 [0.089] N=711	-0.066 [0.136] N=553
Animal products consumed at least weekly	0.053 [0.050] N=1,082	0.050 [0.056] N=983	-0.012 [0.063] N=860	0.013 [0.077] N=711	-0.064 [0.108] N=553
Sweets consumed at least weekly	-0.054 [0.064] N=1,082	-0.058 [0.066] N=983	-0.038 [0.076] N=860	-0.097 [0.087] N=711	-0.051 [0.124] N=553
Snack foods consumed at least weekly	-0.026 [0.061] N=1,082	-0.042 [0.065] N=983	0.002 [0.073] N=860	-0.010 [0.094] N=711	-0.072 [0.110] N=553
Packaged soft drinks consumed at least weekly	-0.017 [0.063] N=1,082	-0.057 [0.069] N=983	-0.059 [0.074] N=860	-0.030 [0.086] N=711	-0.082 [0.122] N=553
Grains consumed at least weekly	0.075 [0.057] N=1,082	0.074 [0.063] N=983	0.065 [0.071] N=860	0.106 [0.087] N=711	0.084 [0.121] N=553
Child was ever breastfed	-0.006 [0.019] N=1,265	-0.003 [0.022] N=1,152	-0.010 [0.027] N=1,004	-0.007 [0.037] N=824	-0.011 [0.043] N=644

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,087	-0.002 [0.028] N=1,175	-0.002 [0.028] N=1,089	-0.013 [0.029] N=975	-0.018 [0.031] N=839	-0.036 [0.033] N=697
Vegetables consumed at least weekly	0.899 P=1,053	0.004 [0.033] N=1,175	-0.011 [0.033] N=1,089	-0.031 [0.033] N=975	-0.016 [0.035] N=839	-0.041 [0.039] N=697
Meat fish or poultry consumed at least weekly	0.766 P=904	0.055 [0.048] N=1,175	0.052 [0.050] N=1,089	0.040 [0.054] N=975	0.031 [0.060] N=839	0.013 [0.064] N=697
Legumes consumed at least weekly	0.765 P=903	-0.060 [0.048] N=1,175	**-0.093* [0.051] N=1,089**	**-0.094* [0.050] N=975**	-0.087 [0.057] N=839	-0.089 [0.066] N=697
Animal products consumed at least weekly	0.906 P=1,067	0.028 [0.039] N=1,175	0.020 [0.042] N=1,089	-0.011 [0.044] N=975	-0.013 [0.048] N=839	-0.039 [0.053] N=697
Sweets consumed at least weekly	0.557 P=649	-0.014 [0.051] N=1,175	-0.019 [0.054] N=1,089	-0.010 [0.058] N=975	-0.024 [0.063] N=839	0.012 [0.071] N=697
Snack foods consumed at least weekly	0.463 P=536	-0.015 [0.048] N=1,175	-0.019 [0.051] N=1,089	0.003 [0.055] N=975	-0.001 [0.060] N=839	0.014 [0.064] N=697
Packaged soft drinks consumed at least weekly	0.267 P=306	0.004 [0.046] N=1,175	-0.018 [0.048] N=1,089	-0.032 [0.050] N=975	-0.009 [0.051] N=839	-0.015 [0.058] N=697
Grains consumed at least weekly	0.867 P=1,020	0.023 [0.040] N=1,175	0.016 [0.041] N=1,089	-0.004 [0.043] N=975	0.007 [0.048] N=839	-0.000 [0.052] N=697
Child was ever breastfed	0.969 P=1,326	-0.021 [0.013] N=1,372	-0.018 [0.015] N=1,273	-0.022 [0.016] N=1,134	-0.018 [0.018] N=974	-0.025 [0.018] N=813

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.018 [0.043] N=992	0.019 [0.045] N=906	-0.005 [0.050] N=792	-0.025 [0.059] N=656	-0.064 [0.076] N=514
Vegetables consumed at least weekly	0.024 [0.052] N=992	-0.002 [0.056] N=906	-0.046 [0.058] N=792	-0.026 [0.069] N=656	-0.100 [0.088] N=514
Meat fish or poultry consumed at least weekly	0.055 [0.066] N=992	0.037 [0.068] N=906	0.012 [0.076] N=792	-0.041 [0.090] N=656	-0.104 [0.086] N=514
Legumes consumed at least weekly	-0.035 [0.072] N=992	-0.102 [0.077] N=906	-0.100 [0.074] N=792	-0.116 [0.090] N=656	-0.106 [0.118] N=514
Animal products consumed at least weekly	0.062 [0.055] N=992	0.050 [0.062] N=906	-0.011 [0.068] N=792	-0.006 [0.085] N=656	-0.069 [0.110] N=514
Sweets consumed at least weekly	-0.048 [0.066] N=992	-0.063 [0.072] N=906	-0.060 [0.082] N=792	-0.136 [0.092] N=656	-0.072 [0.130] N=514
Snack foods consumed at least weekly	-0.045 [0.065] N=992	-0.059 [0.070] N=906	-0.027 [0.079] N=792	-0.062 [0.098] N=656	-0.083 [0.117] N=514
Packaged soft drinks consumed at least weekly	-0.007 [0.066] N=992	-0.053 [0.071] N=906	-0.082 [0.077] N=792	-0.059 [0.087] N=656	-0.081 [0.122] N=514
Grains consumed at least weekly	0.076 [0.061] N=992	0.067 [0.068] N=906	0.034 [0.076] N=792	0.072 [0.094] N=656	0.082 [0.128] N=514
Child was ever breastfed	-0.017 [0.021] N=1,155	-0.009 [0.024] N=1,056	-0.016 [0.029] N=917	-0.014 [0.037] N=757	-0.023 [0.044] N=596

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,087	-0.004 [0.027] N=1,175	-0.004 [0.027] N=1,089	-0.013 [0.028] N=975	-0.017 [0.030] N=839	-0.036 [0.033] N=697
Vegetables consumed at least weekly	0.899 P=1,053	0.003 [0.033] N=1,175	-0.012 [0.033] N=1,089	-0.033 [0.033] N=975	-0.018 [0.035] N=839	-0.044 [0.039] N=697
Meat fish or poultry consumed at least weekly	0.766 P=904	0.052 [0.049] N=1,175	0.050 [0.050] N=1,089	0.039 [0.054] N=975	0.034 [0.060] N=839	0.019 [0.064] N=697
Legumes consumed at least weekly	0.765 P=903	-0.060 [0.048] N=1,175	**-0.093* [0.051] N=1,089**	**-0.095* [0.050] N=975**	-0.090 [0.057] N=839	-0.095 [0.067] N=697
Animal products consumed at least weekly	0.906 P=1,067	0.027 [0.039] N=1,175	0.019 [0.041] N=1,089	-0.011 [0.044] N=975	-0.011 [0.048] N=839	-0.038 [0.052] N=697
Sweets consumed at least weekly	0.557 P=649	-0.014 [0.050] N=1,175	-0.020 [0.054] N=1,089	-0.010 [0.057] N=975	-0.024 [0.062] N=839	0.011 [0.070] N=697
Snack foods consumed at least weekly	0.463 P=536	-0.016 [0.048] N=1,175	-0.020 [0.051] N=1,089	0.002 [0.055] N=975	-0.002 [0.060] N=839	0.012 [0.064] N=697
Packaged soft drinks consumed at least weekly	0.267 P=306	0.004 [0.046] N=1,175	-0.018 [0.049] N=1,089	-0.031 [0.051] N=975	-0.006 [0.052] N=839	-0.013 [0.058] N=697
Grains consumed at least weekly	0.867 P=1,020	0.023 [0.039] N=1,175	0.016 [0.041] N=1,089	-0.003 [0.043] N=975	0.008 [0.048] N=839	0.001 [0.051] N=697
Child was ever breastfed	0.969 P=1,326	-0.021 [0.013] N=1,372	-0.017 [0.015] N=1,273	-0.021 [0.016] N=1,134	-0.019 [0.018] N=974	-0.025 [0.018] N=813

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.011 [0.042] N=992	0.012 [0.044] N=906	-0.006 [0.049] N=792	-0.022 [0.057] N=656	-0.065 [0.072] N=514
Vegetables consumed at least weekly	0.019 [0.051] N=992	-0.006 [0.055] N=906	-0.047 [0.058] N=792	-0.030 [0.067] N=656	-0.114 [0.086] N=514
Meat fish or poultry consumed at least weekly	0.050 [0.067] N=992	0.034 [0.068] N=906	0.011 [0.077] N=792	-0.040 [0.089] N=656	-0.108 [0.087] N=514
Legumes consumed at least weekly	-0.039 [0.071] N=992	-0.106 [0.076] N=906	-0.101 [0.074] N=792	-0.117 [0.090] N=656	-0.117 [0.118] N=514
Animal products consumed at least weekly	0.056 [0.054] N=992	0.044 [0.060] N=906	-0.012 [0.067] N=792	-0.005 [0.082] N=656	-0.073 [0.106] N=514
Sweets consumed at least weekly	-0.056 [0.066] N=992	-0.074 [0.072] N=906	-0.071 [0.082] N=792	-0.142 [0.090] N=656	-0.085 [0.128] N=514
Snack foods consumed at least weekly	-0.053 [0.064] N=992	-0.069 [0.069] N=906	-0.035 [0.079] N=792	-0.069 [0.099] N=656	-0.094 [0.119] N=514
Packaged soft drinks consumed at least weekly	-0.012 [0.067] N=992	-0.058 [0.072] N=906	-0.085 [0.078] N=792	-0.057 [0.088] N=656	-0.077 [0.123] N=514
Grains consumed at least weekly	0.073 [0.060] N=992	0.066 [0.067] N=906	0.036 [0.074] N=792	0.076 [0.093] N=656	0.083 [0.127] N=514
Child was ever breastfed	-0.015 [0.020] N=1,155	-0.007 [0.024] N=1,056	-0.015 [0.028] N=917	-0.014 [0.037] N=757	-0.025 [0.044] N=596

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,025	0.051 [0.032] N=1,103	0.054 [0.033] N=998	0.022 [0.036] N=869	0.004 [0.039] N=727	0.012 [0.045] N=587
Vegetables consumed at least weekly	0.898 P=988	0.036 [0.037] N=1,103	0.044 [0.038] N=998	0.039 [0.042] N=869	0.016 [0.045] N=727	0.024 [0.047] N=587
Meat fish or poultry consumed at least weekly	0.762 P=850	**0.097* [0.055] N=1,103**	**0.100* [0.057] N=998**	0.087 [0.064] N=869	**0.119* [0.067] N=727**	0.109 [0.077] N=587
Legumes consumed at least weekly	0.768 P=857	-0.010 [0.046] N=1,103	0.033 [0.048] N=998	0.018 [0.053] N=869	-0.012 [0.058] N=727	0.020 [0.066] N=587
Animal products consumed at least weekly	0.905 P=1,001	0.014 [0.041] N=1,103	0.015 [0.043] N=998	-0.009 [0.050] N=869	-0.023 [0.054] N=727	-0.018 [0.062] N=587
Sweets consumed at least weekly	0.563 P=622	-0.048 [0.061] N=1,103	-0.056 [0.064] N=998	-0.076 [0.069] N=869	-0.074 [0.073] N=727	-0.135 [0.083] N=587
Snack foods consumed at least weekly	0.463 P=509	-0.041 [0.057] N=1,103	-0.040 [0.060] N=998	-0.006 [0.063] N=869	-0.019 [0.068] N=727	-0.072 [0.077] N=587
Packaged soft drinks consumed at least weekly	0.255 P=276	-0.065 [0.049] N=1,103	-0.074 [0.051] N=998	-0.022 [0.055] N=869	-0.047 [0.060] N=727	-0.075 [0.067] N=587
Grains consumed at least weekly	0.856 P=949	0.044 [0.044] N=1,103	0.053 [0.046] N=998	0.050 [0.049] N=869	0.059 [0.054] N=727	0.040 [0.065] N=587
Child was ever breastfed	0.968 P=1,250	-0.002 [0.023] N=1,293	-0.006 [0.023] N=1,171	0.005 [0.022] N=1,017	0.005 [0.022] N=856	0.012 [0.021] N=688

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.055 [0.051] N=961	0.058 [0.053] N=856	-0.001 [0.061] N=727	-0.011 [0.071] N=585	0.007 [0.093] N=445
Vegetables consumed at least weekly	-0.005 [0.060] N=961	0.003 [0.064] N=856	-0.019 [0.073] N=727	-0.058 [0.085] N=585	-0.058 [0.108] N=445
Meat fish or poultry consumed at least weekly	0.033 [0.074] N=961	0.025 [0.078] N=856	-0.037 [0.089] N=727	0.013 [0.098] N=585	-0.051 [0.130] N=445
Legumes consumed at least weekly	-0.053 [0.057] N=961	0.011 [0.061] N=856	-0.016 [0.072] N=727	-0.065 [0.083] N=585	0.015 [0.095] N=445
Animal products consumed at least weekly	-0.011 [0.053] N=961	-0.011 [0.058] N=856	-0.067 [0.070] N=727	-0.093 [0.073] N=585	-0.107 [0.090] N=445
Sweets consumed at least weekly	-0.028 [0.082] N=961	-0.039 [0.087] N=856	-0.085 [0.097] N=727	-0.083 [0.113] N=585	**-0.273* [0.141] N=445**
Snack foods consumed at least weekly	-0.058 [0.080] N=961	-0.061 [0.084] N=856	-0.007 [0.093] N=727	-0.016 [0.109] N=585	-0.169 [0.133] N=445
Packaged soft drinks consumed at least weekly	-0.089 [0.075] N=961	-0.100 [0.080] N=856	-0.008 [0.094] N=727	-0.046 [0.109] N=585	-0.101 [0.134] N=445
Grains consumed at least weekly	-0.045 [0.051] N=961	-0.049 [0.056] N=856	-0.079 [0.058] N=727	-0.110 [0.069] N=585	**-0.253*** [0.092] N=445**
Child was ever breastfed	0.013 [0.042] N=1,127	0.006 [0.044] N=1,005	0.026 [0.047] N=851	0.028 [0.055] N=690	0.058 [0.068] N=522

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,025	0.052 [0.032] N=1,103	0.054 [0.033] N=998	0.022 [0.037] N=869	0.004 [0.040] N=727	0.012 [0.045] N=587
Vegetables consumed at least weekly	0.898 P=988	0.037 [0.037] N=1,103	0.045 [0.038] N=998	0.039 [0.042] N=869	0.017 [0.045] N=727	0.024 [0.046] N=587
Meat fish or poultry consumed at least weekly	0.762 P=850	**0.098* [0.055] N=1,103**	**0.100* [0.057] N=998**	0.087 [0.064] N=869	**0.118* [0.067] N=727**	0.109 [0.077] N=587
Legumes consumed at least weekly	0.768 P=857	-0.010 [0.046] N=1,103	0.033 [0.048] N=998	0.018 [0.053] N=869	-0.012 [0.058] N=727	0.020 [0.066] N=587
Animal products consumed at least weekly	0.905 P=1,001	0.016 [0.040] N=1,103	0.016 [0.043] N=998	-0.008 [0.050] N=869	-0.021 [0.054] N=727	-0.018 [0.062] N=587
Sweets consumed at least weekly	0.563 P=622	-0.045 [0.061] N=1,103	-0.054 [0.064] N=998	-0.075 [0.069] N=869	-0.074 [0.073] N=727	-0.135 [0.083] N=587
Snack foods consumed at least weekly	0.463 P=509	-0.038 [0.056] N=1,103	-0.038 [0.059] N=998	-0.004 [0.062] N=869	-0.018 [0.067] N=727	-0.072 [0.075] N=587
Packaged soft drinks consumed at least weekly	0.255 P=276	-0.064 [0.049] N=1,103	-0.073 [0.051] N=998	-0.021 [0.054] N=869	-0.047 [0.060] N=727	-0.075 [0.067] N=587
Grains consumed at least weekly	0.856 P=949	0.045 [0.044] N=1,103	0.054 [0.046] N=998	0.051 [0.049] N=869	0.059 [0.055] N=727	0.040 [0.065] N=587
Child was ever breastfed	0.968 P=1,250	-0.001 [0.023] N=1,293	-0.005 [0.023] N=1,171	0.005 [0.022] N=1,017	0.005 [0.023] N=856	0.012 [0.021] N=688

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.056 [0.050] N=961	0.059 [0.053] N=856	-0.001 [0.061] N=727	-0.012 [0.072] N=585	0.005 [0.096] N=445
Vegetables consumed at least weekly	-0.003 [0.059] N=961	0.004 [0.063] N=856	-0.017 [0.073] N=727	-0.053 [0.084] N=585	-0.055 [0.107] N=445
Meat fish or poultry consumed at least weekly	0.034 [0.073] N=961	0.027 [0.077] N=856	-0.035 [0.088] N=727	0.015 [0.097] N=585	-0.048 [0.128] N=445
Legumes consumed at least weekly	-0.053 [0.056] N=961	0.012 [0.061] N=856	-0.016 [0.072] N=727	-0.068 [0.083] N=585	0.003 [0.095] N=445
Animal products consumed at least weekly	-0.006 [0.050] N=961	-0.006 [0.055] N=856	-0.062 [0.068] N=727	-0.085 [0.071] N=585	-0.095 [0.085] N=445
Sweets consumed at least weekly	-0.023 [0.081] N=961	-0.034 [0.086] N=856	-0.080 [0.097] N=727	-0.083 [0.114] N=585	**-0.273* [0.142] N=445**
Snack foods consumed at least weekly	-0.050 [0.078] N=961	-0.055 [0.083] N=856	-0.001 [0.091] N=727	-0.011 [0.108] N=585	-0.161 [0.130] N=445
Packaged soft drinks consumed at least weekly	-0.087 [0.074] N=961	-0.099 [0.079] N=856	-0.006 [0.093] N=727	-0.045 [0.109] N=585	-0.094 [0.131] N=445
Grains consumed at least weekly	-0.044 [0.051] N=961	-0.048 [0.056] N=856	-0.078 [0.059] N=727	-0.114 [0.069] N=585	**-0.259*** [0.095] N=445**
Child was ever breastfed	0.014 [0.043] N=1,127	0.007 [0.045] N=1,005	0.027 [0.048] N=851	0.028 [0.055] N=690	0.061 [0.069] N=522

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,025	0.046 [0.033] N=1,103	0.047 [0.034] N=998	0.019 [0.037] N=869	0.001 [0.041] N=727	0.010 [0.047] N=587
Vegetables consumed at least weekly	0.898 P=988	0.032 [0.037] N=1,103	0.040 [0.038] N=998	0.035 [0.042] N=869	0.013 [0.045] N=727	0.021 [0.048] N=587
Meat fish or poultry consumed at least weekly	0.762 P=850	0.078 [0.055] N=1,103	0.080 [0.057] N=998	0.080 [0.064] N=869	0.112 [0.067] N=727	0.105 [0.078] N=587
Legumes consumed at least weekly	0.768 P=857	-0.017 [0.047] N=1,103	0.026 [0.049] N=998	0.015 [0.054] N=869	-0.015 [0.059] N=727	0.021 [0.068] N=587
Animal products consumed at least weekly	0.905 P=1,001	0.009 [0.041] N=1,103	0.008 [0.044] N=998	-0.013 [0.050] N=869	-0.025 [0.055] N=727	-0.022 [0.063] N=587
Sweets consumed at least weekly	0.563 P=622	-0.049 [0.062] N=1,103	-0.059 [0.064] N=998	-0.080 [0.069] N=869	-0.077 [0.074] N=727	-0.140 [0.085] N=587
Snack foods consumed at least weekly	0.463 P=509	-0.039 [0.056] N=1,103	-0.041 [0.060] N=998	-0.010 [0.062] N=869	-0.022 [0.068] N=727	-0.078 [0.077] N=587
Packaged soft drinks consumed at least weekly	0.255 P=276	-0.058 [0.049] N=1,103	-0.069 [0.051] N=998	-0.022 [0.054] N=869	-0.044 [0.060] N=727	-0.075 [0.067] N=587
Grains consumed at least weekly	0.856 P=949	0.042 [0.044] N=1,103	0.051 [0.047] N=998	0.050 [0.050] N=869	0.058 [0.057] N=727	0.041 [0.067] N=587
Child was ever breastfed	0.968 P=1,250	-0.004 [0.023] N=1,293	-0.008 [0.024] N=1,171	0.005 [0.023] N=1,017	0.004 [0.023] N=856	0.011 [0.021] N=688

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.045 [0.050] N=961	0.047 [0.052] N=856	-0.004 [0.060] N=727	-0.018 [0.073] N=585	-0.004 [0.096] N=445
Vegetables consumed at least weekly	-0.011 [0.059] N=961	-0.004 [0.063] N=856	-0.023 [0.071] N=727	-0.060 [0.083] N=585	-0.062 [0.106] N=445
Meat fish or poultry consumed at least weekly	0.001 [0.072] N=961	-0.010 [0.076] N=856	-0.051 [0.087] N=727	-0.008 [0.101] N=585	-0.085 [0.129] N=445
Legumes consumed at least weekly	-0.061 [0.056] N=961	0.003 [0.062] N=856	-0.020 [0.071] N=727	-0.075 [0.083] N=585	0.005 [0.097] N=445
Animal products consumed at least weekly	-0.013 [0.050] N=961	-0.015 [0.055] N=856	-0.064 [0.067] N=727	-0.087 [0.072] N=585	-0.097 [0.086] N=445
Sweets consumed at least weekly	-0.020 [0.079] N=961	-0.033 [0.084] N=856	-0.078 [0.094] N=727	-0.075 [0.113] N=585	**-0.259* [0.136] N=445**
Snack foods consumed at least weekly	-0.044 [0.077] N=961	-0.051 [0.082] N=856	-0.001 [0.089] N=727	-0.009 [0.106] N=585	-0.155 [0.128] N=445
Packaged soft drinks consumed at least weekly	-0.068 [0.074] N=961	-0.079 [0.079] N=856	0.005 [0.092] N=727	-0.029 [0.108] N=585	-0.071 [0.131] N=445
Grains consumed at least weekly	-0.051 [0.052] N=961	-0.053 [0.057] N=856	-0.079 [0.059] N=727	-0.115 [0.070] N=585	**-0.256*** [0.094] N=445**
Child was ever breastfed	0.010 [0.041] N=1,127	0.001 [0.044] N=1,005	0.026 [0.047] N=851	0.027 [0.053] N=690	0.056 [0.066] N=522

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,025	0.042 [0.033] N=1,103	0.044 [0.033] N=998	0.016 [0.037] N=869	-0.008 [0.040] N=727	-0.002 [0.045] N=587
Vegetables consumed at least weekly	0.898 P=988	0.026 [0.037] N=1,103	0.035 [0.039] N=998	0.030 [0.042] N=869	0.002 [0.044] N=727	0.010 [0.045] N=587
Meat fish or poultry consumed at least weekly	0.762 P=850	0.076 [0.055] N=1,103	0.077 [0.058] N=998	0.067 [0.065] N=869	0.088 [0.067] N=727	0.070 [0.075] N=587
Legumes consumed at least weekly	0.768 P=857	-0.012 [0.047] N=1,103	0.030 [0.050] N=998	0.020 [0.053] N=869	-0.016 [0.057] N=727	0.013 [0.065] N=587
Animal products consumed at least weekly	0.905 P=1,001	0.015 [0.040] N=1,103	0.015 [0.043] N=998	-0.002 [0.050] N=869	-0.019 [0.054] N=727	-0.021 [0.061] N=587
Sweets consumed at least weekly	0.563 P=622	-0.037 [0.063] N=1,103	-0.047 [0.066] N=998	-0.071 [0.072] N=869	-0.072 [0.075] N=727	-0.138 [0.087] N=587
Snack foods consumed at least weekly	0.463 P=509	-0.029 [0.057] N=1,103	-0.030 [0.060] N=998	0.002 [0.064] N=869	-0.016 [0.068] N=727	-0.071 [0.077] N=587
Packaged soft drinks consumed at least weekly	0.255 P=276	-0.047 [0.048] N=1,103	-0.060 [0.050] N=998	-0.015 [0.054] N=869	-0.039 [0.060] N=727	-0.065 [0.069] N=587
Grains consumed at least weekly	0.856 P=949	0.038 [0.044] N=1,103	0.046 [0.047] N=998	0.043 [0.050] N=869	0.050 [0.056] N=727	0.029 [0.066] N=587
Child was ever breastfed	0.968 P=1,250	-0.001 [0.022] N=1,293	-0.006 [0.022] N=1,171	0.007 [0.021] N=1,017	0.005 [0.022] N=856	0.012 [0.021] N=688

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.044 [0.050] N=961	0.046 [0.052] N=856	-0.004 [0.061] N=727	-0.030 [0.074] N=585	-0.033 [0.094] N=445
Vegetables consumed at least weekly	-0.018 [0.060] N=961	-0.012 [0.064] N=856	-0.032 [0.073] N=727	-0.081 [0.082] N=585	-0.103 [0.104] N=445
Meat fish or poultry consumed at least weekly	0.005 [0.075] N=961	-0.007 [0.080] N=856	-0.061 [0.093] N=727	-0.039 [0.109] N=585	-0.149 [0.133] N=445
Legumes consumed at least weekly	-0.053 [0.058] N=961	0.009 [0.064] N=856	-0.002 [0.075] N=727	-0.064 [0.087] N=585	-0.005 [0.100] N=445
Animal products consumed at least weekly	-0.007 [0.047] N=961	-0.007 [0.052] N=856	-0.049 [0.065] N=727	-0.077 [0.070] N=585	-0.098 [0.081] N=445
Sweets consumed at least weekly	-0.002 [0.080] N=961	-0.014 [0.085] N=856	-0.060 [0.095] N=727	-0.059 [0.113] N=585	**-0.250* [0.137] N=445**
Snack foods consumed at least weekly	-0.033 [0.077] N=961	-0.038 [0.083] N=856	0.018 [0.090] N=727	0.002 [0.104] N=585	-0.147 [0.123] N=445
Packaged soft drinks consumed at least weekly	-0.060 [0.072] N=961	-0.074 [0.078] N=856	0.005 [0.091] N=727	-0.025 [0.107] N=585	-0.051 [0.133] N=445
Grains consumed at least weekly	-0.049 [0.051] N=961	-0.052 [0.056] N=856	-0.082 [0.059] N=727	**-0.124* [0.071] N=585**	**-0.274*** [0.092] N=445**
Child was ever breastfed	0.013 [0.041] N=1,127	0.004 [0.043] N=1,005	0.030 [0.045] N=851	0.028 [0.051] N=690	0.054 [0.064] N=522

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=940	0.048 [0.035] N=1,009	0.048 [0.035] N=910	0.033 [0.039] N=799	0.005 [0.043] N=668	0.014 [0.047] N=544
Vegetables consumed at least weekly	0.902 P=909	0.030 [0.037] N=1,009	0.035 [0.039] N=910	0.047 [0.042] N=799	0.018 [0.047] N=668	0.011 [0.049] N=544
Meat fish or poultry consumed at least weekly	0.762 P=777	0.085 [0.055] N=1,009	0.080 [0.057] N=910	0.080 [0.063] N=799	0.090 [0.068] N=668	0.084 [0.077] N=544
Legumes consumed at least weekly	0.765 P=782	-0.010 [0.051] N=1,009	0.034 [0.054] N=910	0.023 [0.058] N=799	-0.014 [0.064] N=668	0.017 [0.075] N=544
Animal products consumed at least weekly	0.907 P=919	0.014 [0.044] N=1,009	0.018 [0.047] N=910	0.006 [0.054] N=799	-0.011 [0.059] N=668	-0.011 [0.067] N=544
Sweets consumed at least weekly	0.574 P=579	-0.045 [0.062] N=1,009	-0.052 [0.065] N=910	-0.078 [0.071] N=799	-0.087 [0.075] N=668	-0.145 [0.088] N=544
Snack foods consumed at least weekly	0.473 P=475	-0.024 [0.058] N=1,009	-0.025 [0.062] N=910	-0.001 [0.065] N=799	-0.020 [0.070] N=668	-0.066 [0.080] N=544
Packaged soft drinks consumed at least weekly	0.264 P=262	-0.072 [0.049] N=1,009	**-0.086* [0.051] N=910**	-0.032 [0.055] N=799	-0.065 [0.062] N=668	-0.105 [0.071] N=544
Grains consumed at least weekly	0.859 P=871	0.065 [0.046] N=1,009	0.073 [0.048] N=910	0.073 [0.051] N=799	0.076 [0.058] N=668	0.051 [0.067] N=544
Child was ever breastfed	0.970 P=1,145	-0.010 [0.022] N=1,181	-0.014 [0.023] N=1,068	-0.002 [0.020] N=934	-0.006 [0.021] N=787	0.003 [0.020] N=637

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.039 [0.051] N=877	0.039 [0.053] N=778	0.009 [0.061] N=667	-0.034 [0.073] N=536	-0.028 [0.083] N=412
Vegetables consumed at least weekly	-0.017 [0.052] N=877	-0.016 [0.058] N=778	-0.010 [0.063] N=667	-0.066 [0.076] N=536	-0.111 [0.089] N=412
Meat fish or poultry consumed at least weekly	0.038 [0.080] N=877	0.022 [0.084] N=778	-0.010 [0.094] N=667	-0.018 [0.103] N=536	-0.077 [0.131] N=412
Legumes consumed at least weekly	-0.048 [0.061] N=877	0.018 [0.066] N=778	0.011 [0.075] N=667	-0.038 [0.090] N=536	0.028 [0.103] N=412
Animal products consumed at least weekly	-0.023 [0.051] N=877	-0.020 [0.057] N=778	-0.050 [0.067] N=667	-0.090 [0.073] N=536	-0.115 [0.085] N=412
Sweets consumed at least weekly	-0.024 [0.083] N=877	-0.036 [0.088] N=778	-0.102 [0.099] N=667	-0.129 [0.113] N=536	**-0.318** [0.148] N=412**
Snack foods consumed at least weekly	-0.039 [0.081] N=877	-0.045 [0.088] N=778	-0.015 [0.094] N=667	-0.051 [0.108] N=536	-0.178 [0.138] N=412
Packaged soft drinks consumed at least weekly	-0.098 [0.075] N=877	-0.115 [0.081] N=778	-0.026 [0.095] N=667	-0.087 [0.115] N=536	-0.202 [0.143] N=412
Grains consumed at least weekly	-0.019 [0.056] N=877	-0.024 [0.062] N=778	-0.045 [0.066] N=667	-0.104 [0.079] N=536	**-0.286*** [0.102] N=412**
Child was ever breastfed	0.002 [0.039] N=1,027	-0.005 [0.042] N=914	0.015 [0.042] N=780	0.006 [0.050] N=633	0.034 [0.061] N=483

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=940	0.049 [0.034] N=1,009	0.050 [0.035] N=910	0.034 [0.039] N=799	0.005 [0.042] N=668	0.013 [0.046] N=544
Vegetables consumed at least weekly	0.902 P=909	0.033 [0.037] N=1,009	0.038 [0.039] N=910	0.051 [0.041] N=799	0.019 [0.046] N=668	0.011 [0.048] N=544
Meat fish or poultry consumed at least weekly	0.762 P=777	**0.094* [0.054] N=1,009**	0.089 [0.056] N=910	0.087 [0.062] N=799	0.091 [0.065] N=668	0.084 [0.074] N=544
Legumes consumed at least weekly	0.765 P=782	-0.011 [0.050] N=1,009	0.033 [0.053] N=910	0.024 [0.057] N=799	-0.018 [0.063] N=668	0.015 [0.074] N=544
Animal products consumed at least weekly	0.907 P=919	0.013 [0.043] N=1,009	0.017 [0.047] N=910	0.005 [0.053] N=799	-0.012 [0.058] N=668	-0.010 [0.066] N=544
Sweets consumed at least weekly	0.574 P=579	-0.047 [0.062] N=1,009	-0.056 [0.065] N=910	-0.083 [0.071] N=799	-0.093 [0.076] N=668	-0.148 [0.089] N=544
Snack foods consumed at least weekly	0.473 P=475	-0.024 [0.057] N=1,009	-0.026 [0.061] N=910	-0.001 [0.064] N=799	-0.020 [0.069] N=668	-0.064 [0.079] N=544
Packaged soft drinks consumed at least weekly	0.264 P=262	-0.074 [0.049] N=1,009	**-0.089* [0.051] N=910**	-0.036 [0.054] N=799	-0.069 [0.060] N=668	-0.107 [0.069] N=544
Grains consumed at least weekly	0.859 P=871	0.065 [0.045] N=1,009	0.073 [0.048] N=910	0.072 [0.050] N=799	0.076 [0.057] N=668	0.050 [0.067] N=544
Child was ever breastfed	0.970 P=1,145	-0.009 [0.022] N=1,181	-0.013 [0.023] N=1,068	0.000 [0.021] N=934	-0.003 [0.022] N=787	0.006 [0.020] N=637

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.044 [0.051] N=877	0.046 [0.054] N=778	0.016 [0.062] N=667	-0.030 [0.074] N=536	-0.021 [0.090] N=412
Vegetables consumed at least weekly	-0.015 [0.053] N=877	-0.014 [0.058] N=778	-0.005 [0.063] N=667	-0.069 [0.077] N=536	-0.118 [0.093] N=412
Meat fish or poultry consumed at least weekly	0.048 [0.080] N=877	0.030 [0.085] N=778	-0.006 [0.094] N=667	-0.021 [0.102] N=536	-0.078 [0.132] N=412
Legumes consumed at least weekly	-0.048 [0.061] N=877	0.019 [0.067] N=778	0.011 [0.075] N=667	-0.049 [0.090] N=536	0.033 [0.101] N=412
Animal products consumed at least weekly	-0.023 [0.051] N=877	-0.018 [0.057] N=778	-0.049 [0.067] N=667	-0.091 [0.074] N=536	-0.117 [0.089] N=412
Sweets consumed at least weekly	-0.022 [0.083] N=877	-0.037 [0.089] N=778	-0.101 [0.100] N=667	-0.128 [0.115] N=536	**-0.320** [0.147] N=412**
Snack foods consumed at least weekly	-0.038 [0.081] N=877	-0.046 [0.087] N=778	-0.010 [0.094] N=667	-0.036 [0.109] N=536	-0.172 [0.134] N=412
Packaged soft drinks consumed at least weekly	-0.098 [0.077] N=877	-0.115 [0.082] N=778	-0.020 [0.095] N=667	-0.072 [0.113] N=536	-0.168 [0.142] N=412
Grains consumed at least weekly	-0.020 [0.056] N=877	-0.025 [0.061] N=778	-0.047 [0.064] N=667	-0.110 [0.077] N=536	**-0.291*** [0.099] N=412**
Child was ever breastfed	0.002 [0.039] N=1,027	-0.006 [0.042] N=914	0.018 [0.043] N=780	0.012 [0.052] N=633	0.039 [0.063] N=483

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.