Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Oldest Child

Age reference: 10 October 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.053	0.179 [0.251] N=1,284	0.291 [0.264] N=1,191	0.362 [0.287] N=1,062	0.400 [0.291] N=912	0.042 [0.315] N=761
WHO weight-for-age z score	0.314	-0.104 [0.133] N=1,483	-0.074 [0.149] N=1,371	-0.029 [0.154] N=1,225	-0.005 [0.155] N=1,047	-0.099 [0.173] N=868
WHO weight-for-length/height z score, 0-60 months	0.317	0.346 [0.218] N=1,280	0.339 [0.228] N=1,186	0.343 [0.243] N=1,062	0.348 [0.253] N=911	0.435 [0.268] N=757
WHO BMI-for-age z score	0.313	0.271 [0.245] N=1,253	0.326 [0.261] N=1,164	0.316 [0.284] N=1,039	0.198 [0.300] N=887	0.268 [0.324] N=739
Stunted: HAZ below -2	0.200 P=249	-0.044 [0.048] N=1,284	-0.054 [0.050] N=1,191	-0.045 [0.053] N=1,062	-0.032 [0.055] N=912	-0.000 [0.063] N=761
Underweight: WAZ below -2	0.057 P=88	**0.037* [0.021] N=1,483**	0.028 [0.023] N=1,371	0.028 [0.024] N=1,225	0.016 [0.024] N=1,047	0.010 [0.032] N=868
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=150	-0.047 [0.035] N=1,280	-0.037 [0.037] N=1,186	-0.053 [0.040] N=1,062	-0.033 [0.042] N=911	-0.048 [0.044] N=757
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=246	0.039 [0.046] N=1,280	0.043 [0.050] N=1,186	0.028 [0.052] N=1,062	0.039 [0.059] N=911	0.044 [0.066] N=757
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=113	0.024 [0.034] N=1,280	0.029 [0.036] N=1,186	0.038 [0.038] N=1,062	0.042 [0.042] N=911	0.050 [0.046] N=757
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=327	-0.064 [0.052] N=1,222	-0.049 [0.054] N=1,133	-0.049 [0.058] N=1,013	-0.031 [0.062] N=869	-0.011 [0.071] N=724
BMI-for-age z score below -2	0.124 P=153	-0.012 [0.039] N=1,253	-0.004 [0.042] N=1,164	-0.006 [0.047] N=1,039	0.045 [0.049] N=887	0.045 [0.053] N=739

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.191 [0.390] N=1,076	0.397 [0.441] N=983	0.655 [0.501] N=854	**0.966* [0.523] N=704**	0.278 [0.724] N=553
WHO weight-for-age z score	**-0.353* [0.191] N=1,252**	-0.340 [0.231] N=1,140	-0.298 [0.240] N=994	-0.344 [0.278] N=816	**-0.799* [0.410] N=637**
WHO weight-for-length/height z score, 0-60 months	0.164 [0.310] N=1,074	0.128 [0.342] N=980	0.035 [0.369] N=856	-0.006 [0.430] N=705	-0.047 [0.508] N=551
WHO BMI-for-age z score	0.204 [0.306] N=1,051	0.287 [0.339] N=962	0.167 [0.379] N=837	-0.142 [0.421] N=685	-0.224 [0.495] N=537
Stunted: HAZ below -2	-0.042 [0.067] N=1,076	-0.058 [0.073] N=983	-0.055 [0.080] N=854	-0.037 [0.093] N=704	0.051 [0.139] N=553
Underweight: WAZ below -2	**0.072** [0.028] N=1,252**	**0.065** [0.033] N=1,140**	**0.073** [0.033] N=994**	**0.065* [0.035] N=816**	0.082 [0.061] N=637
Wasted: WLZ/WHZ below -2, 0-60 months	-0.053 [0.053] N=1,074	-0.036 [0.059] N=980	-0.065 [0.068] N=856	-0.039 [0.083] N=705	-0.080 [0.099] N=551
Overweight: WLZ/WHZ above +2, 0-60 months	-0.004 [0.056] N=1,074	-0.001 [0.064] N=980	-0.045 [0.065] N=856	-0.041 [0.084] N=705	-0.088 [0.104] N=551
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.041] N=1,074	0.008 [0.047] N=980	0.020 [0.049] N=856	0.030 [0.058] N=705	0.041 [0.073] N=551
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.050 [0.074] N=1,022	-0.017 [0.081] N=933	-0.017 [0.091] N=813	0.018 [0.111] N=669	0.108 [0.161] N=524
BMI-for-age z score below -2	-0.053 [0.050] N=1,051	-0.045 [0.056] N=962	-0.050 [0.068] N=837	0.033 [0.078] N=685	0.040 [0.091] N=537

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.053	0.180 [0.251] N=1,284	0.291 [0.264] N=1,191	0.362 [0.288] N=1,062	0.406 [0.292] N=912	0.055 [0.318] N=761
WHO weight-for-age z score	0.314	-0.105 [0.132] N=1,483	-0.079 [0.148] N=1,371	-0.031 [0.153] N=1,225	-0.009 [0.155] N=1,047	-0.107 [0.172] N=868
WHO weight-for-length/height z score, 0-60 months	0.317	0.343 [0.218] N=1,280	0.336 [0.229] N=1,186	0.342 [0.244] N=1,062	0.345 [0.255] N=911	0.427 [0.271] N=757
WHO BMI-for-age z score	0.313	0.270 [0.246] N=1,253	0.323 [0.262] N=1,164	0.314 [0.285] N=1,039	0.196 [0.301] N=887	0.265 [0.325] N=739
Stunted: HAZ below -2	0.200 P=249	-0.044 [0.048] N=1,284	-0.053 [0.050] N=1,191	-0.045 [0.053] N=1,062	-0.031 [0.056] N=912	0.001 [0.064] N=761
Underweight: WAZ below -2	0.057 P=88	**0.037* [0.021] N=1,483**	0.029 [0.023] N=1,371	0.028 [0.024] N=1,225	0.016 [0.025] N=1,047	0.011 [0.032] N=868
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=150	-0.045 [0.035] N=1,280	-0.035 [0.037] N=1,186	-0.051 [0.040] N=1,062	-0.030 [0.043] N=911	-0.044 [0.045] N=757
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=246	0.039 [0.046] N=1,280	0.043 [0.050] N=1,186	0.028 [0.052] N=1,062	0.040 [0.059] N=911	0.046 [0.065] N=757
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=113	0.024 [0.034] N=1,280	0.030 [0.036] N=1,186	0.039 [0.038] N=1,062	0.043 [0.042] N=911	0.052 [0.046] N=757
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=327	-0.062 [0.051] N=1,222	-0.046 [0.053] N=1,133	-0.047 [0.057] N=1,013	-0.028 [0.061] N=869	-0.005 [0.070] N=724
BMI-for-age z score below -2	0.124 P=153	-0.012 [0.039] N=1,253	-0.003 [0.042] N=1,164	-0.005 [0.048] N=1,039	0.047 [0.049] N=887	0.048 [0.054] N=739

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.184 [0.387] N=1,076	0.392 [0.439] N=983	0.643 [0.497] N=854	**0.936* [0.514] N=704**	0.259 [0.713] N=553
WHO weight-for-age z score	**-0.342* [0.190] N=1,252**	-0.333 [0.231] N=1,140	-0.287 [0.242] N=994	-0.338 [0.280] N=816	**-0.796* [0.409] N=637**
WHO weight-for-length/height z score, 0-60 months	0.166 [0.309] N=1,074	0.128 [0.341] N=980	0.039 [0.369] N=856	-0.003 [0.431] N=705	-0.050 [0.513] N=551
WHO BMI-for-age z score	0.211 [0.305] N=1,051	0.291 [0.339] N=962	0.178 [0.379] N=837	-0.136 [0.424] N=685	-0.224 [0.500] N=537
Stunted: HAZ below -2	-0.043 [0.067] N=1,076	-0.059 [0.073] N=983	-0.056 [0.080] N=854	-0.036 [0.093] N=704	0.052 [0.138] N=553
Underweight: WAZ below -2	**0.071** [0.028] N=1,252**	**0.065** [0.032] N=1,140**	**0.072** [0.033] N=994**	**0.065* [0.035] N=816**	0.082 [0.060] N=637
Wasted: WLZ/WHZ below -2, 0-60 months	-0.054 [0.052] N=1,074	-0.036 [0.059] N=980	-0.068 [0.067] N=856	-0.040 [0.083] N=705	-0.079 [0.099] N=551
Overweight: WLZ/WHZ above +2, 0-60 months	-0.004 [0.056] N=1,074	-0.001 [0.064] N=980	-0.045 [0.065] N=856	-0.041 [0.084] N=705	-0.088 [0.104] N=551
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.041] N=1,074	0.008 [0.048] N=980	0.019 [0.049] N=856	0.029 [0.058] N=705	0.041 [0.073] N=551
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.053 [0.073] N=1,022	-0.021 [0.080] N=933	-0.027 [0.089] N=813	0.010 [0.110] N=669	0.103 [0.160] N=524
BMI-for-age z score below -2	-0.054 [0.050] N=1,051	-0.046 [0.056] N=962	-0.053 [0.068] N=837	0.031 [0.079] N=685	0.040 [0.093] N=537

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.053	0.153 [0.251] N=1,284	0.253 [0.265] N=1,191	0.303 [0.286] N=1,062	0.382 [0.291] N=912	0.006 [0.315] N=761
WHO weight-for-age z score	0.314	-0.138 [0.126] N=1,483	-0.115 [0.142] N=1,371	-0.070 [0.146] N=1,225	-0.027 [0.150] N=1,047	-0.143 [0.165] N=868
WHO weight-for-length/height z score, 0-60 months	0.317	0.339 [0.221] N=1,280	0.337 [0.234] N=1,186	0.357 [0.249] N=1,062	0.357 [0.259] N=911	0.426 [0.280] N=757
WHO BMI-for-age z score	0.313	0.275 [0.248] N=1,253	0.332 [0.266] N=1,164	0.334 [0.289] N=1,039	0.210 [0.305] N=887	0.280 [0.334] N=739
Stunted: HAZ below -2	0.200 P=249	-0.040 [0.048] N=1,284	-0.046 [0.051] N=1,191	-0.035 [0.053] N=1,062	-0.028 [0.056] N=912	0.010 [0.065] N=761
Underweight: WAZ below -2	0.057 P=88	**0.039* [0.020] N=1,483**	0.033 [0.022] N=1,371	0.032 [0.023] N=1,225	0.018 [0.024] N=1,047	0.017 [0.031] N=868
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=150	-0.043 [0.035] N=1,280	-0.032 [0.038] N=1,186	-0.050 [0.040] N=1,062	-0.030 [0.043] N=911	-0.042 [0.046] N=757
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=246	0.041 [0.046] N=1,280	0.046 [0.050] N=1,186	0.032 [0.053] N=1,062	0.042 [0.059] N=911	0.048 [0.066] N=757
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=113	0.025 [0.034] N=1,280	0.032 [0.037] N=1,186	0.040 [0.039] N=1,062	0.044 [0.042] N=911	0.052 [0.046] N=757
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=327	-0.055 [0.052] N=1,222	-0.034 [0.054] N=1,133	-0.037 [0.057] N=1,013	-0.024 [0.061] N=869	0.012 [0.072] N=724
BMI-for-age z score below -2	0.124 P=153	-0.012 [0.039] N=1,253	-0.002 [0.042] N=1,164	-0.006 [0.048] N=1,039	0.046 [0.050] N=887	0.046 [0.055] N=739

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.170 [0.385] N=1,076	0.355 [0.432] N=983	0.557 [0.485] N=854	**0.913* [0.496] N=704**	0.244 [0.682] N=553
WHO weight-for-age z score	**-0.355* [0.185] N=1,252**	-0.353 [0.226] N=1,140	-0.312 [0.232] N=994	-0.324 [0.282] N=816	**-0.798* [0.404] N=637**
WHO weight-for-length/height z score, 0-60 months	0.172 [0.309] N=1,074	0.137 [0.345] N=980	0.074 [0.369] N=856	-0.003 [0.423] N=705	-0.074 [0.511] N=551
WHO BMI-for-age z score	0.216 [0.305] N=1,051	0.296 [0.341] N=962	0.205 [0.381] N=837	-0.163 [0.413] N=685	-0.230 [0.502] N=537
Stunted: HAZ below -2	-0.042 [0.066] N=1,076	-0.054 [0.073] N=983	-0.049 [0.078] N=854	-0.045 [0.090] N=704	0.047 [0.134] N=553
Underweight: WAZ below -2	**0.071** [0.028] N=1,252**	**0.066** [0.032] N=1,140**	**0.072** [0.033] N=994**	0.058 [0.037] N=816	0.079 [0.062] N=637
Wasted: WLZ/WHZ below -2, 0-60 months	-0.054 [0.053] N=1,074	-0.035 [0.060] N=980	-0.072 [0.068] N=856	-0.044 [0.083] N=705	-0.074 [0.101] N=551
Overweight: WLZ/WHZ above +2, 0-60 months	-0.004 [0.056] N=1,074	0.001 [0.065] N=980	-0.043 [0.065] N=856	-0.048 [0.085] N=705	-0.086 [0.106] N=551
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.041] N=1,074	0.008 [0.049] N=980	0.018 [0.050] N=856	0.026 [0.059] N=705	0.040 [0.073] N=551
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.054 [0.074] N=1,022	-0.017 [0.081] N=933	-0.029 [0.089] N=813	-0.012 [0.109] N=669	0.105 [0.159] N=524
BMI-for-age z score below -2	-0.055 [0.050] N=1,051	-0.044 [0.056] N=962	-0.056 [0.069] N=837	0.033 [0.077] N=685	0.045 [0.093] N=537

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.053	0.126 [0.253] N=1,284	0.235 [0.267] N=1,191	0.286 [0.290] N=1,062	0.374 [0.296] N=912	-0.014 [0.327] N=761
WHO weight-for-age z score	0.314	-0.155 [0.126] N=1,483	-0.129 [0.143] N=1,371	-0.086 [0.148] N=1,225	-0.058 [0.151] N=1,047	-0.190 [0.170] N=868
WHO weight-for-length/height z score, 0-60 months	0.317	0.337 [0.222] N=1,280	0.331 [0.234] N=1,186	0.338 [0.249] N=1,062	0.319 [0.262] N=911	0.377 [0.280] N=757
WHO BMI-for-age z score	0.313	0.279 [0.248] N=1,253	0.337 [0.264] N=1,164	0.333 [0.287] N=1,039	0.186 [0.307] N=887	0.229 [0.333] N=739
Stunted: HAZ below -2	0.200 P=249	-0.033 [0.048] N=1,284	-0.039 [0.051] N=1,191	-0.025 [0.054] N=1,062	-0.018 [0.057] N=912	0.022 [0.066] N=761
Underweight: WAZ below -2	0.057 P=88	**0.043** [0.020] N=1,483**	**0.037* [0.022] N=1,371**	**0.038* [0.022] N=1,225**	0.025 [0.023] N=1,047	0.027 [0.030] N=868
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=150	-0.042 [0.035] N=1,280	-0.030 [0.037] N=1,186	-0.045 [0.040] N=1,062	-0.025 [0.043] N=911	-0.039 [0.045] N=757
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=246	0.042 [0.047] N=1,280	0.048 [0.051] N=1,186	0.035 [0.053] N=1,062	0.041 [0.059] N=911	0.044 [0.067] N=757
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=113	0.024 [0.034] N=1,280	0.029 [0.037] N=1,186	0.039 [0.038] N=1,062	0.040 [0.042] N=911	0.044 [0.045] N=757
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=327	-0.046 [0.052] N=1,222	-0.024 [0.054] N=1,133	-0.020 [0.058] N=1,013	-0.005 [0.063] N=869	0.034 [0.074] N=724
BMI-for-age z score below -2	0.124 P=153	-0.010 [0.039] N=1,253	-0.001 [0.042] N=1,164	-0.001 [0.048] N=1,039	0.051 [0.050] N=887	0.050 [0.055] N=739

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.138 [0.392] N=1,076	0.336 [0.436] N=983	0.532 [0.496] N=854	**0.924* [0.523] N=704**	0.186 [0.709] N=553
WHO weight-for-age z score	**-0.372* [0.189] N=1,252**	-0.357 [0.228] N=1,140	-0.330 [0.236] N=994	-0.302 [0.287] N=816	**-0.756* [0.405] N=637**
WHO weight-for-length/height z score, 0-60 months	0.189 [0.309] N=1,074	0.160 [0.340] N=980	0.081 [0.373] N=856	0.059 [0.435] N=705	0.029 [0.521] N=551
WHO BMI-for-age z score	0.216 [0.305] N=1,051	0.314 [0.337] N=962	0.213 [0.385] N=837	-0.080 [0.429] N=685	-0.105 [0.509] N=537
Stunted: HAZ below -2	-0.039 [0.067] N=1,076	-0.052 [0.073] N=983	-0.048 [0.080] N=854	-0.050 [0.092] N=704	0.051 [0.137] N=553
Underweight: WAZ below -2	**0.071** [0.028] N=1,252**	**0.064** [0.032] N=1,140**	**0.072** [0.033] N=994**	0.053 [0.038] N=816	0.080 [0.063] N=637
Wasted: WLZ/WHZ below -2, 0-60 months	-0.055 [0.053] N=1,074	-0.039 [0.059] N=980	-0.072 [0.068] N=856	-0.044 [0.085] N=705	-0.078 [0.103] N=551
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.056] N=1,074	0.004 [0.064] N=980	-0.039 [0.066] N=856	-0.033 [0.085] N=705	-0.052 [0.102] N=551
Obese: WLZ/WHZ above +3, 0-60 months	0.004 [0.042] N=1,074	0.015 [0.048] N=980	0.026 [0.050] N=856	0.036 [0.058] N=705	0.051 [0.071] N=551
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.051 [0.076] N=1,022	-0.019 [0.083] N=933	-0.025 [0.091] N=813	-0.013 [0.112] N=669	0.109 [0.166] N=524
BMI-for-age z score below -2	-0.054 [0.050] N=1,051	-0.050 [0.056] N=962	-0.056 [0.068] N=837	0.028 [0.080] N=685	0.032 [0.095] N=537

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.056	0.151 [0.246] N=1,179	0.219 [0.263] N=1,098	0.281 [0.290] N=976	0.252 [0.291] N=844	-0.134 [0.307] N=706
WHO weight-for-age z score	0.322	-0.026 [0.134] N=1,358	-0.008 [0.147] N=1,260	0.053 [0.153] N=1,123	0.076 [0.152] N=965	-0.001 [0.173] N=805
WHO weight-for-length/height z score, 0-60 months	0.335	**0.491** [0.217] N=1,179**	**0.480** [0.226] N=1,098**	**0.485** [0.242] N=982**	**0.512** [0.245] N=849**	**0.638** [0.260] N=708**
WHO BMI-for-age z score	0.334	**0.428* [0.249] N=1,152**	**0.457* [0.262] N=1,075**	0.465 [0.284] N=959	0.383 [0.292] N=825	0.494 [0.320] N=690
Stunted: HAZ below -2	0.200 P=228	-0.033 [0.049] N=1,179	-0.037 [0.052] N=1,098	-0.026 [0.055] N=976	-0.005 [0.058] N=844	0.036 [0.067] N=706
Underweight: WAZ below -2	0.057 P=81	0.032 [0.022] N=1,358	0.025 [0.024] N=1,260	0.024 [0.025] N=1,123	0.013 [0.026] N=965	0.010 [0.034] N=805
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=137	**-0.069* [0.036] N=1,179**	-0.056 [0.037] N=1,098	**-0.070* [0.040] N=982**	-0.052 [0.042] N=849	-0.069 [0.044] N=708
Overweight: WLZ/WHZ above +2, 0-60 months	0.201 P=229	0.058 [0.047] N=1,179	0.062 [0.050] N=1,098	0.049 [0.053] N=982	0.059 [0.058] N=849	0.077 [0.065] N=708
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=108	0.022 [0.036] N=1,179	0.030 [0.038] N=1,098	0.042 [0.040] N=982	0.054 [0.043] N=849	0.069 [0.046] N=708
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=300	-0.068 [0.054] N=1,124	-0.047 [0.056] N=1,047	-0.051 [0.060] N=934	-0.035 [0.063] N=808	-0.012 [0.075] N=675
BMI-for-age z score below -2	0.120 P=136	-0.033 [0.038] N=1,152	-0.023 [0.040] N=1,075	-0.029 [0.046] N=959	0.017 [0.048] N=825	0.007 [0.053] N=690

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.367 [0.391] N=985	0.549 [0.448] N=904	0.837 [0.510] N=782	**1.005* [0.525] N=650**	0.334 [0.736] N=512
WHO weight-for-age z score	-0.296 [0.193] N=1,143	-0.307 [0.228] N=1,045	-0.254 [0.235] N=908	-0.322 [0.261] N=750	**-0.745* [0.406] N=590**
WHO weight-for-length/height z score, 0-60 months	0.181 [0.323] N=986	0.097 [0.347] N=905	-0.031 [0.377] N=789	-0.016 [0.433] N=656	-0.009 [0.515] N=515
WHO BMI-for-age z score	0.225 [0.324] N=963	0.220 [0.347] N=886	0.106 [0.380] N=770	-0.113 [0.416] N=636	-0.193 [0.516] N=501
Stunted: HAZ below -2	-0.060 [0.070] N=985	-0.077 [0.076] N=904	-0.076 [0.083] N=782	-0.052 [0.095] N=650	0.034 [0.150] N=512
Underweight: WAZ below -2	**0.069** [0.029] N=1,143**	**0.065* [0.034] N=1,045**	**0.073** [0.034] N=908**	**0.070* [0.036] N=750**	0.098 [0.064] N=590
Wasted: WLZ/WHZ below -2, 0-60 months	-0.083 [0.055] N=986	-0.059 [0.060] N=905	-0.087 [0.068] N=789	-0.076 [0.082] N=656	-0.145 [0.097] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.007 [0.057] N=986	-0.012 [0.064] N=905	-0.058 [0.066] N=789	-0.061 [0.086] N=656	-0.097 [0.107] N=515
Obese: WLZ/WHZ above +3, 0-60 months	-0.023 [0.044] N=986	-0.015 [0.051] N=905	-0.001 [0.053] N=789	0.023 [0.062] N=656	0.040 [0.077] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.076 [0.076] N=937	-0.040 [0.084] N=860	-0.051 [0.093] N=747	-0.039 [0.110] N=621	0.018 [0.167] N=488
BMI-for-age z score below -2	-0.052 [0.051] N=963	-0.033 [0.056] N=886	-0.041 [0.068] N=770	0.032 [0.079] N=636	0.023 [0.095] N=501

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.056	0.152 [0.247] N=1,179	0.216 [0.264] N=1,098	0.276 [0.290] N=976	0.245 [0.288] N=844	-0.151 [0.302] N=706
WHO weight-for-age z score	0.322	-0.009 [0.134] N=1,358	0.005 [0.147] N=1,260	0.060 [0.152] N=1,123	0.069 [0.154] N=965	-0.008 [0.178] N=805
WHO weight-for-length/height z score, 0-60 months	0.335	**0.493** [0.214] N=1,179**	**0.482** [0.222] N=1,098**	**0.481** [0.238] N=982**	**0.501** [0.242] N=849**	**0.626** [0.255] N=708**
WHO BMI-for-age z score	0.334	**0.429* [0.244] N=1,152**	**0.457* [0.256] N=1,075**	0.462 [0.279] N=959	0.374 [0.288] N=825	0.477 [0.314] N=690
Stunted: HAZ below -2	0.200 P=228	-0.035 [0.050] N=1,179	-0.039 [0.052] N=1,098	-0.027 [0.055] N=976	-0.004 [0.058] N=844	0.037 [0.067] N=706
Underweight: WAZ below -2	0.057 P=81	0.029 [0.022] N=1,358	0.023 [0.023] N=1,260	0.023 [0.025] N=1,123	0.014 [0.026] N=965	0.009 [0.034] N=805
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=137	**-0.068* [0.035] N=1,179**	-0.055 [0.037] N=1,098	**-0.070* [0.040] N=982**	-0.051 [0.042] N=849	-0.070 [0.044] N=708
Overweight: WLZ/WHZ above +2, 0-60 months	0.201 P=229	0.059 [0.046] N=1,179	0.063 [0.049] N=1,098	0.048 [0.052] N=982	0.055 [0.057] N=849	0.072 [0.064] N=708
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=108	0.022 [0.035] N=1,179	0.030 [0.038] N=1,098	0.042 [0.039] N=982	0.052 [0.042] N=849	0.067 [0.046] N=708
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=300	-0.069 [0.054] N=1,124	-0.047 [0.056] N=1,047	-0.050 [0.061] N=934	-0.030 [0.064] N=808	-0.008 [0.076] N=675
BMI-for-age z score below -2	0.120 P=136	-0.032 [0.037] N=1,152	-0.022 [0.040] N=1,075	-0.029 [0.045] N=959	0.017 [0.048] N=825	0.007 [0.052] N=690

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.368 [0.395] N=985	0.540 [0.453] N=904	0.826 [0.516] N=782	**0.992* [0.527] N=650**	0.328 [0.729] N=512
WHO weight-for-age z score	-0.262 [0.190] N=1,143	-0.276 [0.227] N=1,045	-0.231 [0.233] N=908	-0.327 [0.263] N=750	**-0.743* [0.416] N=590**
WHO weight-for-length/height z score, 0-60 months	0.199 [0.318] N=986	0.122 [0.346] N=905	-0.011 [0.378] N=789	-0.029 [0.430] N=656	-0.042 [0.510] N=515
WHO BMI-for-age z score	0.237 [0.319] N=963	0.238 [0.344] N=886	0.121 [0.379] N=770	-0.128 [0.409] N=636	-0.237 [0.497] N=501
Stunted: HAZ below -2	-0.065 [0.070] N=985	-0.081 [0.077] N=904	-0.079 [0.083] N=782	-0.053 [0.095] N=650	0.028 [0.147] N=512
Underweight: WAZ below -2	**0.063** [0.030] N=1,143**	**0.060* [0.034] N=1,045**	**0.070** [0.035] N=908**	**0.071* [0.037] N=750**	0.090 [0.067] N=590
Wasted: WLZ/WHZ below -2, 0-60 months	-0.081 [0.054] N=986	-0.058 [0.060] N=905	-0.087 [0.068] N=789	-0.075 [0.082] N=656	-0.144 [0.096] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.057] N=986	-0.005 [0.064] N=905	-0.053 [0.066] N=789	-0.062 [0.086] N=656	-0.098 [0.106] N=515
Obese: WLZ/WHZ above +3, 0-60 months	-0.021 [0.044] N=986	-0.013 [0.051] N=905	0.000 [0.053] N=789	0.023 [0.062] N=656	0.039 [0.076] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.078 [0.076] N=937	-0.044 [0.084] N=860	-0.055 [0.093] N=747	-0.039 [0.111] N=621	0.013 [0.168] N=488
BMI-for-age z score below -2	-0.049 [0.051] N=963	-0.032 [0.056] N=886	-0.041 [0.068] N=770	0.034 [0.079] N=636	0.027 [0.095] N=501

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146	-0.066 [0.342] N=1,115	-0.175 [0.346] N=1,002	-0.009 [0.373] N=870	-0.062 [0.424] N=738	-0.044 [0.481] N=597
WHO weight-for-age z score	0.318	0.036 [0.160] N=1,281	0.036 [0.170] N=1,160	-0.028 [0.180] N=1,006	-0.024 [0.192] N=849	0.065 [0.209] N=682
WHO weight-for-length/height z score, 0-60 months	0.343	0.089 [0.269] N=1,115	0.069 [0.277] N=1,007	0.008 [0.303] N=874	0.070 [0.340] N=738	0.003 [0.378] N=595
WHO BMI-for-age z score	0.329	0.132 [0.271] N=1,090	0.171 [0.278] N=984	0.094 [0.312] N=853	0.220 [0.352] N=721	0.268 [0.392] N=585
Stunted: HAZ below -2	0.180 P=201	-0.033 [0.052] N=1,115	-0.017 [0.054] N=1,002	-0.058 [0.056] N=870	-0.043 [0.064] N=738	-0.094 [0.071] N=597
Underweight: WAZ below -2	0.053 P=71	-0.004 [0.028] N=1,281	-0.005 [0.028] N=1,160	0.013 [0.029] N=1,006	0.025 [0.029] N=849	0.015 [0.029] N=682
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=129	0.008 [0.041] N=1,115	0.018 [0.043] N=1,007	0.035 [0.048] N=874	0.004 [0.048] N=738	0.019 [0.052] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=219	0.052 [0.049] N=1,115	0.055 [0.051] N=1,007	0.026 [0.053] N=874	0.025 [0.058] N=738	0.032 [0.065] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=104	0.034 [0.040] N=1,115	0.028 [0.041] N=1,007	0.016 [0.043] N=874	0.006 [0.048] N=738	-0.019 [0.048] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=277	0.015 [0.049] N=1,064	0.027 [0.052] N=959	0.008 [0.053] N=832	-0.009 [0.055] N=704	-0.036 [0.059] N=568
BMI-for-age z score below -2	0.124 P=134	0.030 [0.046] N=1,090	0.038 [0.049] N=984	0.037 [0.056] N=853	0.019 [0.059] N=721	0.043 [0.067] N=585

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.162 [0.459] N=968	-0.320 [0.479] N=855	-0.098 [0.550] N=723	-0.356 [0.666] N=591	-0.307 [0.766] N=450
WHO weight-for-age z score	0.013 [0.225] N=1,115	0.009 [0.247] N=994	-0.148 [0.269] N=840	-0.199 [0.303] N=683	-0.075 [0.349] N=516
WHO weight-for-length/height z score, 0-60 months	-0.102 [0.372] N=970	-0.172 [0.396] N=862	-0.335 [0.472] N=729	-0.279 [0.539] N=593	-0.800 [0.633] N=450
WHO BMI-for-age z score	-0.263 [0.361] N=950	-0.244 [0.381] N=844	-0.489 [0.469] N=713	-0.371 [0.530] N=581	-0.575 [0.592] N=445
Stunted: HAZ below -2	0.037 [0.075] N=968	0.062 [0.081] N=855	0.009 [0.086] N=723	0.078 [0.110] N=591	-0.073 [0.127] N=450
Underweight: WAZ below -2	-0.023 [0.048] N=1,115	-0.026 [0.051] N=994	0.007 [0.057] N=840	0.024 [0.063] N=683	-0.003 [0.069] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.043 [0.058] N=970	0.067 [0.062] N=862	0.106 [0.076] N=729	0.046 [0.077] N=593	0.125 [0.090] N=450
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.076] N=970	0.024 [0.081] N=862	-0.035 [0.088] N=729	-0.052 [0.102] N=593	-0.121 [0.125] N=450
Obese: WLZ/WHZ above +3, 0-60 months	0.057 [0.056] N=970	0.039 [0.059] N=862	0.019 [0.067] N=729	-0.007 [0.084] N=593	-0.133 [0.098] N=450
CIAF: any stunting, underweight, or wasting, 0-60 months	0.080 [0.079] N=923	0.101 [0.085] N=818	0.088 [0.094] N=691	0.080 [0.105] N=563	-0.003 [0.127] N=427
BMI-for-age z score below -2	0.057 [0.050] N=950	0.073 [0.054] N=844	0.078 [0.071] N=713	0.033 [0.067] N=581	0.109 [0.079] N=445

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146	-0.078 [0.341] N=1,115	-0.191 [0.345] N=1,002	-0.033 [0.370] N=870	-0.085 [0.420] N=738	-0.061 [0.475] N=597
WHO weight-for-age z score	0.318	0.044 [0.160] N=1,281	0.042 [0.169] N=1,160	-0.024 [0.179] N=1,006	-0.017 [0.191] N=849	0.070 [0.208] N=682
WHO weight-for-length/height z score, 0-60 months	0.343	0.092 [0.269] N=1,115	0.069 [0.277] N=1,007	0.008 [0.302] N=874	0.074 [0.338] N=738	0.013 [0.373] N=595
WHO BMI-for-age z score	0.329	0.132 [0.271] N=1,090	0.167 [0.278] N=984	0.090 [0.313] N=853	0.220 [0.352] N=721	0.269 [0.391] N=585
Stunted: HAZ below -2	0.180 P=201	-0.035 [0.052] N=1,115	-0.018 [0.054] N=1,002	-0.057 [0.056] N=870	-0.041 [0.065] N=738	-0.093 [0.071] N=597
Underweight: WAZ below -2	0.053 P=71	-0.006 [0.028] N=1,281	-0.006 [0.028] N=1,160	0.013 [0.029] N=1,006	0.025 [0.029] N=849	0.015 [0.029] N=682
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=129	0.004 [0.041] N=1,115	0.013 [0.043] N=1,007	0.031 [0.048] N=874	0.001 [0.047] N=738	0.015 [0.051] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=219	0.049 [0.050] N=1,115	0.051 [0.052] N=1,007	0.022 [0.053] N=874	0.022 [0.059] N=738	0.030 [0.066] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=104	0.031 [0.040] N=1,115	0.024 [0.041] N=1,007	0.012 [0.042] N=874	0.003 [0.048] N=738	-0.022 [0.049] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=277	0.007 [0.050] N=1,064	0.019 [0.052] N=959	0.002 [0.054] N=832	-0.012 [0.055] N=704	-0.037 [0.059] N=568
BMI-for-age z score below -2	0.124 P=134	0.027 [0.046] N=1,090	0.035 [0.049] N=984	0.034 [0.056] N=853	0.017 [0.058] N=721	0.041 [0.066] N=585

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.168 [0.462] N=968	-0.340 [0.483] N=855	-0.143 [0.549] N=723	-0.427 [0.669] N=591	-0.366 [0.770] N=450
WHO weight-for-age z score	0.032 [0.222] N=1,115	0.026 [0.244] N=994	-0.134 [0.266] N=840	-0.162 [0.298] N=683	-0.028 [0.349] N=516
WHO weight-for-length/height z score, 0-60 months	-0.098 [0.376] N=970	-0.176 [0.401] N=862	-0.342 [0.478] N=729	-0.260 [0.548] N=593	-0.729 [0.658] N=450
WHO BMI-for-age z score	-0.258 [0.366] N=950	-0.246 [0.386] N=844	-0.495 [0.473] N=713	-0.353 [0.538] N=581	-0.522 [0.617] N=445
Stunted: HAZ below -2	0.030 [0.075] N=968	0.055 [0.081] N=855	0.006 [0.086] N=723	0.079 [0.113] N=591	-0.071 [0.130] N=450
Underweight: WAZ below -2	-0.026 [0.048] N=1,115	-0.029 [0.051] N=994	0.006 [0.057] N=840	0.023 [0.064] N=683	-0.003 [0.071] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.059] N=970	0.056 [0.064] N=862	0.094 [0.078] N=729	0.028 [0.077] N=593	0.094 [0.093] N=450
Overweight: WLZ/WHZ above +2, 0-60 months	0.029 [0.076] N=970	0.015 [0.081] N=862	-0.048 [0.088] N=729	-0.070 [0.103] N=593	-0.135 [0.126] N=450
Obese: WLZ/WHZ above +3, 0-60 months	0.051 [0.056] N=970	0.031 [0.059] N=862	0.010 [0.067] N=729	-0.021 [0.084] N=593	-0.151 [0.100] N=450
CIAF: any stunting, underweight, or wasting, 0-60 months	0.057 [0.079] N=923	0.075 [0.085] N=818	0.060 [0.094] N=691	0.049 [0.105] N=563	-0.043 [0.123] N=427
BMI-for-age z score below -2	0.048 [0.051] N=950	0.061 [0.055] N=844	0.067 [0.072] N=713	0.017 [0.067] N=581	0.088 [0.083] N=445

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146	-0.096 [0.329] N=1,115	-0.184 [0.332] N=1,002	-0.012 [0.359] N=870	-0.077 [0.408] N=738	-0.057 [0.462] N=597
WHO weight-for-age z score	0.318	0.037 [0.156] N=1,281	0.039 [0.165] N=1,160	-0.011 [0.175] N=1,006	-0.010 [0.189] N=849	0.073 [0.205] N=682
WHO weight-for-length/height z score, 0-60 months	0.343	0.092 [0.268] N=1,115	0.063 [0.276] N=1,007	0.013 [0.300] N=874	0.080 [0.334] N=738	0.009 [0.371] N=595
WHO BMI-for-age z score	0.329	0.129 [0.270] N=1,090	0.154 [0.277] N=984	0.080 [0.306] N=853	0.214 [0.342] N=721	0.241 [0.381] N=585
Stunted: HAZ below -2	0.180 P=201	-0.030 [0.051] N=1,115	-0.017 [0.053] N=1,002	-0.062 [0.054] N=870	-0.044 [0.063] N=738	-0.100 [0.067] N=597
Underweight: WAZ below -2	0.053 P=71	-0.004 [0.027] N=1,281	-0.005 [0.028] N=1,160	0.012 [0.029] N=1,006	0.024 [0.028] N=849	0.014 [0.029] N=682
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=129	0.005 [0.042] N=1,115	0.014 [0.044] N=1,007	0.030 [0.049] N=874	0.000 [0.049] N=738	0.017 [0.053] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=219	0.051 [0.050] N=1,115	0.051 [0.052] N=1,007	0.021 [0.053] N=874	0.019 [0.058] N=738	0.030 [0.066] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=104	0.032 [0.040] N=1,115	0.024 [0.041] N=1,007	0.012 [0.043] N=874	0.003 [0.048] N=738	-0.022 [0.049] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=277	0.013 [0.051] N=1,064	0.021 [0.053] N=959	-0.004 [0.055] N=832	-0.019 [0.057] N=704	-0.047 [0.059] N=568
BMI-for-age z score below -2	0.124 P=134	0.029 [0.047] N=1,090	0.037 [0.049] N=984	0.036 [0.057] N=853	0.019 [0.059] N=721	0.047 [0.067] N=585

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.259 [0.465] N=968	-0.405 [0.484] N=855	-0.178 [0.544] N=723	-0.499 [0.659] N=591	-0.494 [0.773] N=450
WHO weight-for-age z score	-0.026 [0.223] N=1,115	-0.029 [0.245] N=994	-0.140 [0.267] N=840	-0.183 [0.301] N=683	-0.087 [0.355] N=516
WHO weight-for-length/height z score, 0-60 months	-0.113 [0.378] N=970	-0.194 [0.403] N=862	-0.336 [0.475] N=729	-0.247 [0.544] N=593	-0.741 [0.663] N=450
WHO BMI-for-age z score	-0.262 [0.369] N=950	-0.262 [0.390] N=844	-0.486 [0.470] N=713	-0.329 [0.535] N=581	-0.504 [0.626] N=445
Stunted: HAZ below -2	0.042 [0.076] N=968	0.063 [0.081] N=855	0.012 [0.086] N=723	0.094 [0.112] N=591	-0.052 [0.131] N=450
Underweight: WAZ below -2	-0.020 [0.047] N=1,115	-0.022 [0.050] N=994	0.008 [0.056] N=840	0.026 [0.062] N=683	0.005 [0.072] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.041 [0.061] N=970	0.064 [0.065] N=862	0.097 [0.079] N=729	0.031 [0.078] N=593	0.107 [0.096] N=450
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.076] N=970	0.020 [0.082] N=862	-0.046 [0.088] N=729	-0.065 [0.103] N=593	-0.132 [0.126] N=450
Obese: WLZ/WHZ above +3, 0-60 months	0.053 [0.056] N=970	0.034 [0.060] N=862	0.012 [0.067] N=729	-0.017 [0.084] N=593	-0.147 [0.098] N=450
CIAF: any stunting, underweight, or wasting, 0-60 months	0.077 [0.080] N=923	0.090 [0.086] N=818	0.068 [0.096] N=691	0.060 [0.110] N=563	-0.018 [0.128] N=427
BMI-for-age z score below -2	0.048 [0.053] N=950	0.064 [0.056] N=844	0.068 [0.072] N=713	0.018 [0.067] N=581	0.093 [0.084] N=445

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146	-0.097 [0.328] N=1,115	-0.181 [0.332] N=1,002	0.009 [0.359] N=870	-0.070 [0.403] N=738	-0.081 [0.458] N=597
WHO weight-for-age z score	0.318	0.023 [0.156] N=1,281	0.024 [0.166] N=1,160	-0.021 [0.177] N=1,006	-0.019 [0.189] N=849	0.062 [0.207] N=682
WHO weight-for-length/height z score, 0-60 months	0.343	0.108 [0.263] N=1,115	0.079 [0.270] N=1,007	0.020 [0.292] N=874	0.108 [0.326] N=738	0.070 [0.365] N=595
WHO BMI-for-age z score	0.329	0.139 [0.269] N=1,090	0.169 [0.276] N=984	0.094 [0.304] N=853	0.245 [0.342] N=721	0.321 [0.385] N=585
Stunted: HAZ below -2	0.180 P=201	-0.033 [0.051] N=1,115	-0.020 [0.053] N=1,002	-0.065 [0.055] N=870	-0.044 [0.063] N=738	-0.092 [0.068] N=597
Underweight: WAZ below -2	0.053 P=71	-0.005 [0.027] N=1,281	-0.006 [0.028] N=1,160	0.012 [0.029] N=1,006	0.025 [0.029] N=849	0.016 [0.031] N=682
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=129	0.003 [0.041] N=1,115	0.013 [0.043] N=1,007	0.032 [0.047] N=874	-0.001 [0.046] N=738	0.011 [0.048] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=219	0.052 [0.049] N=1,115	0.053 [0.051] N=1,007	0.025 [0.052] N=874	0.029 [0.057] N=738	0.041 [0.065] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=104	0.037 [0.039] N=1,115	0.030 [0.040] N=1,007	0.016 [0.042] N=874	0.009 [0.046] N=738	-0.015 [0.048] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=277	0.012 [0.051] N=1,064	0.023 [0.053] N=959	0.000 [0.055] N=832	-0.011 [0.056] N=704	-0.028 [0.059] N=568
BMI-for-age z score below -2	0.124 P=134	0.024 [0.046] N=1,090	0.033 [0.048] N=984	0.033 [0.055] N=853	0.015 [0.057] N=721	0.033 [0.064] N=585

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.237 [0.447] N=968	-0.369 [0.470] N=855	-0.061 [0.536] N=723	-0.388 [0.659] N=591	-0.414 [0.759] N=450
WHO weight-for-age z score	-0.009 [0.217] N=1,115	-0.012 [0.238] N=994	-0.117 [0.266] N=840	-0.153 [0.292] N=683	-0.043 [0.343] N=516
WHO weight-for-length/height z score, 0-60 months	-0.087 [0.372] N=970	-0.170 [0.396] N=862	-0.346 [0.463] N=729	-0.235 [0.530] N=593	-0.676 [0.653] N=450
WHO BMI-for-age z score	-0.255 [0.365] N=950	-0.247 [0.386] N=844	-0.503 [0.466] N=713	-0.321 [0.526] N=581	-0.457 [0.613] N=445
Stunted: HAZ below -2	0.040 [0.074] N=968	0.058 [0.079] N=855	-0.004 [0.085] N=723	0.077 [0.113] N=591	-0.052 [0.128] N=450
Underweight: WAZ below -2	-0.024 [0.046] N=1,115	-0.028 [0.050] N=994	0.002 [0.057] N=840	0.021 [0.063] N=683	-0.003 [0.073] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.037 [0.060] N=970	0.061 [0.064] N=862	0.102 [0.076] N=729	0.038 [0.076] N=593	0.107 [0.091] N=450
Overweight: WLZ/WHZ above +2, 0-60 months	0.032 [0.076] N=970	0.017 [0.081] N=862	-0.049 [0.086] N=729	-0.067 [0.101] N=593	-0.124 [0.128] N=450
Obese: WLZ/WHZ above +3, 0-60 months	0.058 [0.055] N=970	0.038 [0.058] N=862	0.012 [0.064] N=729	-0.012 [0.081] N=593	-0.137 [0.098] N=450
CIAF: any stunting, underweight, or wasting, 0-60 months	0.070 [0.080] N=923	0.085 [0.086] N=818	0.060 [0.096] N=691	0.052 [0.108] N=563	-0.007 [0.127] N=427
BMI-for-age z score below -2	0.049 [0.052] N=950	0.064 [0.056] N=844	0.073 [0.071] N=713	0.021 [0.065] N=581	0.086 [0.080] N=445

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.160	-0.172 [0.341] N=1,020	-0.296 [0.346] N=916	-0.160 [0.367] N=801	-0.248 [0.404] N=677	-0.267 [0.466] N=550
WHO weight-for-age z score	0.319	0.060 [0.167] N=1,170	0.079 [0.176] N=1,058	-0.015 [0.184] N=924	-0.031 [0.191] N=781	0.020 [0.206] N=632
WHO weight-for-length/height z score, 0-60 months	0.353	0.189 [0.287] N=1,024	0.188 [0.300] N=925	0.094 [0.326] N=807	0.121 [0.358] N=681	0.027 [0.388] N=553
WHO BMI-for-age z score	0.341	0.188 [0.293] N=1,000	0.250 [0.304] N=903	0.119 [0.342] N=788	0.218 [0.382] N=666	0.211 [0.421] N=544
Stunted: HAZ below -2	0.178 P=182	-0.025 [0.054] N=1,020	-0.004 [0.056] N=916	-0.047 [0.056] N=801	-0.033 [0.062] N=677	-0.078 [0.070] N=550
Underweight: WAZ below -2	0.054 P=66	0.001 [0.030] N=1,170	-0.002 [0.031] N=1,058	0.016 [0.032] N=924	0.034 [0.032] N=781	0.027 [0.032] N=632
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=118	0.004 [0.043] N=1,024	0.010 [0.045] N=925	0.029 [0.050] N=807	0.002 [0.048] N=681	0.020 [0.052] N=553
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=204	0.058 [0.053] N=1,024	0.062 [0.055] N=925	0.026 [0.056] N=807	0.014 [0.060] N=681	0.015 [0.065] N=553
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=98	0.031 [0.040] N=1,024	0.023 [0.042] N=925	0.013 [0.043] N=807	0.003 [0.047] N=681	-0.016 [0.046] N=553
CIAF: any stunting, underweight, or wasting, 0-60 months	0.258 P=254	0.020 [0.054] N=976	0.027 [0.056] N=880	0.007 [0.055] N=768	-0.013 [0.056] N=648	-0.026 [0.062] N=526
BMI-for-age z score below -2	0.121 P=119	0.011 [0.048] N=1,000	0.014 [0.051] N=903	0.014 [0.058] N=788	0.007 [0.061] N=666	0.040 [0.068] N=544

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.061 [0.488] N=884	-0.245 [0.506] N=780	-0.067 [0.576] N=665	-0.372 [0.693] N=541	-0.157 [0.822] N=414
WHO weight-for-age z score	0.095 [0.233] N=1,016	0.110 [0.254] N=904	-0.105 [0.281] N=770	-0.134 [0.311] N=627	0.046 [0.378] N=478
WHO weight-for-length/height z score, 0-60 months	-0.025 [0.408] N=888	-0.074 [0.437] N=789	-0.314 [0.515] N=671	-0.285 [0.604] N=545	-0.938 [0.723] N=417
WHO BMI-for-age z score	-0.204 [0.385] N=868	-0.158 [0.407] N=771	-0.497 [0.497] N=656	-0.374 [0.573] N=534	-0.756 [0.661] N=412
Stunted: HAZ below -2	0.020 [0.083] N=884	0.049 [0.090] N=780	-0.003 [0.094] N=665	0.053 [0.120] N=541	-0.119 [0.145] N=414
Underweight: WAZ below -2	-0.031 [0.051] N=1,016	-0.037 [0.055] N=904	-0.005 [0.062] N=770	0.022 [0.073] N=627	-0.013 [0.082] N=478
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.063] N=888	0.052 [0.068] N=789	0.086 [0.082] N=671	0.012 [0.082] N=545	0.085 [0.095] N=417
Overweight: WLZ/WHZ above +2, 0-60 months	0.033 [0.084] N=888	0.020 [0.090] N=789	-0.057 [0.097] N=671	-0.109 [0.116] N=545	-0.219 [0.137] N=417
Obese: WLZ/WHZ above +3, 0-60 months	0.034 [0.060] N=888	0.011 [0.064] N=789	-0.008 [0.072] N=671	-0.047 [0.092] N=545	-0.183 [0.113] N=417
CIAF: any stunting, underweight, or wasting, 0-60 months	0.064 [0.089] N=844	0.077 [0.097] N=748	0.055 [0.102] N=636	0.006 [0.116] N=516	-0.121 [0.141] N=394
BMI-for-age z score below -2	0.037 [0.056] N=868	0.045 [0.061] N=771	0.044 [0.079] N=656	-0.010 [0.078] N=534	0.093 [0.092] N=412

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.160	-0.196 [0.346] N=1,020	-0.320 [0.352] N=916	-0.179 [0.372] N=801	-0.271 [0.407] N=677	-0.269 [0.469] N=550
WHO weight-for-age z score	0.319	0.057 [0.164] N=1,170	0.068 [0.174] N=1,058	-0.018 [0.182] N=924	-0.032 [0.189] N=781	0.023 [0.202] N=632
WHO weight-for-length/height z score, 0-60 months	0.353	0.191 [0.291] N=1,024	0.186 [0.303] N=925	0.095 [0.330] N=807	0.123 [0.361] N=681	0.021 [0.396] N=553
WHO BMI-for-age z score	0.341	0.186 [0.296] N=1,000	0.249 [0.307] N=903	0.117 [0.345] N=788	0.219 [0.384] N=666	0.210 [0.426] N=544
Stunted: HAZ below -2	0.178 P=182	-0.024 [0.054] N=1,020	-0.004 [0.056] N=916	-0.047 [0.056] N=801	-0.033 [0.062] N=677	-0.079 [0.069] N=550
Underweight: WAZ below -2	0.054 P=66	-0.003 [0.029] N=1,170	-0.004 [0.030] N=1,058	0.012 [0.031] N=924	0.029 [0.031] N=781	0.023 [0.031] N=632
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=118	-0.000 [0.043] N=1,024	0.006 [0.045] N=925	0.024 [0.050] N=807	-0.006 [0.048] N=681	0.013 [0.051] N=553
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=204	0.058 [0.052] N=1,024	0.061 [0.054] N=925	0.025 [0.055] N=807	0.012 [0.060] N=681	0.012 [0.065] N=553
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=98	0.030 [0.040] N=1,024	0.021 [0.041] N=925	0.012 [0.042] N=807	0.002 [0.046] N=681	-0.020 [0.045] N=553
CIAF: any stunting, underweight, or wasting, 0-60 months	0.258 P=254	0.015 [0.053] N=976	0.021 [0.055] N=880	0.000 [0.054] N=768	-0.023 [0.055] N=648	-0.038 [0.057] N=526
BMI-for-age z score below -2	0.121 P=119	0.010 [0.049] N=1,000	0.013 [0.051] N=903	0.013 [0.059] N=788	0.003 [0.060] N=666	0.037 [0.067] N=544

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.083 [0.486] N=884	-0.259 [0.504] N=780	-0.064 [0.568] N=665	-0.367 [0.672] N=541	-0.185 [0.784] N=414
WHO weight-for-age z score	0.084 [0.232] N=1,016	0.093 [0.255] N=904	-0.108 [0.280] N=770	-0.143 [0.308] N=627	0.010 [0.364] N=478
WHO weight-for-length/height z score, 0-60 months	-0.044 [0.403] N=888	-0.100 [0.431] N=789	-0.342 [0.509] N=671	-0.329 [0.581] N=545	-0.954 [0.682] N=417
WHO BMI-for-age z score	-0.213 [0.382] N=868	-0.169 [0.403] N=771	-0.508 [0.493] N=656	-0.379 [0.560] N=534	-0.716 [0.627] N=412
Stunted: HAZ below -2	0.020 [0.083] N=884	0.048 [0.090] N=780	-0.007 [0.093] N=665	0.048 [0.118] N=541	-0.112 [0.140] N=414
Underweight: WAZ below -2	-0.033 [0.052] N=1,016	-0.037 [0.056] N=904	-0.007 [0.062] N=770	0.021 [0.072] N=627	-0.005 [0.079] N=478
Wasted: WLZ/WHZ below -2, 0-60 months	0.037 [0.063] N=888	0.054 [0.068] N=789	0.087 [0.083] N=671	0.014 [0.081] N=545	0.087 [0.094] N=417
Overweight: WLZ/WHZ above +2, 0-60 months	0.033 [0.082] N=888	0.019 [0.088] N=789	-0.060 [0.094] N=671	-0.116 [0.111] N=545	-0.210 [0.131] N=417
Obese: WLZ/WHZ above +3, 0-60 months	0.033 [0.059] N=888	0.010 [0.062] N=789	-0.009 [0.071] N=671	-0.051 [0.089] N=545	**-0.183* [0.104] N=417**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.065 [0.088] N=844	0.077 [0.096] N=748	0.054 [0.101] N=636	0.006 [0.116] N=516	-0.087 [0.138] N=394
BMI-for-age z score below -2	0.040 [0.056] N=868	0.048 [0.061] N=771	0.047 [0.079] N=656	-0.005 [0.075] N=534	0.095 [0.089] N=412

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.