Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,106	0.005 [0.029] N=1,197	0.013 [0.029] N=1,106	0.005 [0.032] N=990	0.001 [0.033] N=852	-0.008 [0.037] N=700
Vegetables consumed at least weekly	0.896 P=1,070	-0.001 [0.033] N=1,197	-0.013 [0.033] N=1,106	-0.022 [0.034] N=990	-0.015 [0.036] N=852	-0.046 [0.042] N=700
Meat fish or poultry consumed at least weekly	0.762 P=918	0.057 [0.057] N=1,197	0.072 [0.059] N=1,106	0.077 [0.064] N=990	0.058 [0.071] N=852	0.047 [0.079] N=700
Legumes consumed at least weekly	0.759 P=914	-0.038 [0.048] N=1,197	-0.067 [0.050] N=1,106	-0.049 [0.050] N=990	-0.046 [0.054] N=852	-0.047 [0.063] N=700
Animal products consumed at least weekly	0.904 P=1,085	0.017 [0.039] N=1,197	0.012 [0.041] N=1,106	-0.012 [0.045] N=990	-0.002 [0.049] N=852	-0.014 [0.055] N=700
Sweets consumed at least weekly	0.533 P=635	-0.021 [0.052] N=1,197	-0.017 [0.054] N=1,106	-0.005 [0.058] N=990	-0.024 [0.065] N=852	0.016 [0.075] N=700
Snack foods consumed at least weekly	0.439 P=519	0.001 [0.046] N=1,197	-0.007 [0.048] N=1,106	0.017 [0.051] N=990	0.020 [0.057] N=852	0.025 [0.061] N=700
Packaged soft drinks consumed at least weekly	0.248 P=291	-0.020 [0.046] N=1,197	-0.043 [0.049] N=1,106	-0.042 [0.050] N=990	-0.037 [0.052] N=852	-0.062 [0.059] N=700
Grains consumed at least weekly	0.860 P=1,031	0.028 [0.038] N=1,197	0.024 [0.040] N=1,106	0.018 [0.042] N=990	0.030 [0.044] N=852	0.015 [0.051] N=700
Child was ever breastfed	0.968 P=1,373	-0.017 [0.013] N=1,422	-0.016 [0.014] N=1,317	-0.019 [0.015] N=1,176	-0.018 [0.017] N=1,006	-0.020 [0.016] N=830

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.046] N=1,015	0.019 [0.048] N=924	-0.002 [0.055] N=808	-0.016 [0.065] N=670	-0.052 [0.083] N=518
Vegetables consumed at least weekly	0.008 [0.050] N=1,015	-0.011 [0.052] N=924	-0.033 [0.057] N=808	-0.026 [0.066] N=670	-0.132 [0.092] N=518
Meat fish or poultry consumed at least weekly	0.007 [0.065] N=1,015	0.019 [0.069] N=924	0.014 [0.079] N=808	-0.084 [0.084] N=670	**-0.198** [0.089] N=518**
Legumes consumed at least weekly	-0.045 [0.073] N=1,015	-0.106 [0.076] N=924	-0.075 [0.077] N=808	-0.093 [0.090] N=670	-0.109 [0.128] N=518
Animal products consumed at least weekly	0.017 [0.051] N=1,015	0.011 [0.058] N=924	-0.044 [0.065] N=808	-0.017 [0.076] N=670	-0.068 [0.094] N=518
Sweets consumed at least weekly	-0.048 [0.069] N=1,015	-0.046 [0.072] N=924	-0.032 [0.079] N=808	-0.110 [0.091] N=670	-0.018 [0.124] N=518
Snack foods consumed at least weekly	-0.017 [0.062] N=1,015	-0.038 [0.066] N=924	0.001 [0.074] N=808	-0.012 [0.097] N=670	-0.008 [0.119] N=518
Packaged soft drinks consumed at least weekly	-0.006 [0.069] N=1,015	-0.048 [0.076] N=924	-0.043 [0.080] N=808	-0.044 [0.091] N=670	-0.130 [0.122] N=518
Grains consumed at least weekly	0.062 [0.058] N=1,015	0.059 [0.064] N=924	0.053 [0.070] N=808	0.086 [0.084] N=670	0.049 [0.120] N=518
Child was ever breastfed	-0.010 [0.020] N=1,203	-0.006 [0.023] N=1,098	-0.011 [0.029] N=957	-0.013 [0.036] N=787	-0.007 [0.042] N=611

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,106	0.005 [0.029] N=1,197	0.013 [0.029] N=1,106	0.006 [0.032] N=990	0.002 [0.033] N=852	-0.009 [0.036] N=700
Vegetables consumed at least weekly	0.896 P=1,070	-0.001 [0.032] N=1,197	-0.013 [0.032] N=1,106	-0.021 [0.034] N=990	-0.015 [0.036] N=852	-0.047 [0.042] N=700
Meat fish or poultry consumed at least weekly	0.762 P=918	0.057 [0.056] N=1,197	0.072 [0.059] N=1,106	0.077 [0.064] N=990	0.058 [0.071] N=852	0.046 [0.080] N=700
Legumes consumed at least weekly	0.759 P=914	-0.037 [0.049] N=1,197	-0.067 [0.050] N=1,106	-0.048 [0.050] N=990	-0.046 [0.055] N=852	-0.048 [0.063] N=700
Animal products consumed at least weekly	0.904 P=1,085	0.017 [0.038] N=1,197	0.012 [0.041] N=1,106	-0.011 [0.044] N=990	-0.002 [0.048] N=852	-0.015 [0.055] N=700
Sweets consumed at least weekly	0.533 P=635	-0.021 [0.052] N=1,197	-0.017 [0.054] N=1,106	-0.004 [0.058] N=990	-0.024 [0.066] N=852	0.016 [0.076] N=700
Snack foods consumed at least weekly	0.439 P=519	0.001 [0.046] N=1,197	-0.007 [0.049] N=1,106	0.019 [0.052] N=990	0.020 [0.058] N=852	0.024 [0.061] N=700
Packaged soft drinks consumed at least weekly	0.248 P=291	-0.020 [0.046] N=1,197	-0.043 [0.049] N=1,106	-0.041 [0.050] N=990	-0.036 [0.052] N=852	-0.062 [0.059] N=700
Grains consumed at least weekly	0.860 P=1,031	0.028 [0.038] N=1,197	0.024 [0.040] N=1,106	0.019 [0.042] N=990	0.031 [0.044] N=852	0.014 [0.051] N=700
Child was ever breastfed	0.968 P=1,373	-0.017 [0.013] N=1,422	-0.016 [0.014] N=1,317	-0.019 [0.015] N=1,176	-0.017 [0.017] N=1,006	-0.019 [0.016] N=830

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.046] N=1,015	0.019 [0.048] N=924	-0.000 [0.055] N=808	-0.016 [0.066] N=670	-0.052 [0.082] N=518
Vegetables consumed at least weekly	0.008 [0.050] N=1,015	-0.011 [0.052] N=924	-0.032 [0.058] N=808	-0.025 [0.067] N=670	-0.132 [0.092] N=518
Meat fish or poultry consumed at least weekly	0.007 [0.066] N=1,015	0.019 [0.070] N=924	0.015 [0.079] N=808	-0.084 [0.085] N=670	**-0.198** [0.089] N=518**
Legumes consumed at least weekly	-0.044 [0.073] N=1,015	-0.106 [0.076] N=924	-0.072 [0.077] N=808	-0.092 [0.090] N=670	-0.109 [0.127] N=518
Animal products consumed at least weekly	0.019 [0.051] N=1,015	0.011 [0.057] N=924	-0.040 [0.064] N=808	-0.016 [0.076] N=670	-0.068 [0.093] N=518
Sweets consumed at least weekly	-0.048 [0.069] N=1,015	-0.046 [0.071] N=924	-0.031 [0.079] N=808	-0.109 [0.091] N=670	-0.018 [0.123] N=518
Snack foods consumed at least weekly	-0.015 [0.062] N=1,015	-0.037 [0.066] N=924	0.007 [0.073] N=808	-0.009 [0.097] N=670	-0.008 [0.117] N=518
Packaged soft drinks consumed at least weekly	-0.006 [0.069] N=1,015	-0.048 [0.076] N=924	-0.042 [0.080] N=808	-0.044 [0.091] N=670	-0.130 [0.123] N=518
Grains consumed at least weekly	0.063 [0.058] N=1,015	0.059 [0.064] N=924	0.056 [0.069] N=808	0.088 [0.084] N=670	0.049 [0.119] N=518
Child was ever breastfed	-0.010 [0.020] N=1,203	-0.007 [0.023] N=1,098	-0.011 [0.029] N=957	-0.014 [0.036] N=787	-0.006 [0.042] N=611

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,106	-0.005 [0.029] N=1,197	0.003 [0.030] N=1,106	-0.004 [0.032] N=990	-0.005 [0.034] N=852	-0.021 [0.037] N=700
Vegetables consumed at least weekly	0.896 P=1,070	-0.007 [0.033] N=1,197	-0.022 [0.032] N=1,106	-0.031 [0.034] N=990	-0.022 [0.035] N=852	-0.060 [0.041] N=700
Meat fish or poultry consumed at least weekly	0.762 P=918	0.033 [0.055] N=1,197	0.044 [0.058] N=1,106	0.050 [0.063] N=990	0.039 [0.069] N=852	0.017 [0.077] N=700
Legumes consumed at least weekly	0.759 P=914	-0.044 [0.049] N=1,197	-0.075 [0.050] N=1,106	-0.056 [0.050] N=990	-0.049 [0.054] N=852	-0.053 [0.062] N=700
Animal products consumed at least weekly	0.904 P=1,085	0.011 [0.038] N=1,197	0.004 [0.041] N=1,106	-0.019 [0.045] N=990	-0.009 [0.048] N=852	-0.028 [0.055] N=700
Sweets consumed at least weekly	0.533 P=635	-0.021 [0.052] N=1,197	-0.019 [0.055] N=1,106	-0.005 [0.058] N=990	-0.024 [0.065] N=852	0.007 [0.074] N=700
Snack foods consumed at least weekly	0.439 P=519	0.006 [0.045] N=1,197	-0.007 [0.048] N=1,106	0.020 [0.051] N=990	0.024 [0.056] N=852	0.023 [0.060] N=700
Packaged soft drinks consumed at least weekly	0.248 P=291	-0.007 [0.045] N=1,197	-0.029 [0.049] N=1,106	-0.027 [0.049] N=990	-0.025 [0.051] N=852	-0.049 [0.059] N=700
Grains consumed at least weekly	0.860 P=1,031	0.021 [0.039] N=1,197	0.018 [0.040] N=1,106	0.012 [0.042] N=990	0.025 [0.044] N=852	0.009 [0.051] N=700
Child was ever breastfed	0.968 P=1,373	-0.018 [0.013] N=1,422	-0.019 [0.014] N=1,317	-0.022 [0.015] N=1,176	-0.019 [0.017] N=1,006	-0.022 [0.017] N=830

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.045] N=1,015	0.018 [0.047] N=924	0.002 [0.054] N=808	0.007 [0.066] N=670	-0.036 [0.085] N=518
Vegetables consumed at least weekly	0.009 [0.049] N=1,015	-0.011 [0.051] N=924	-0.030 [0.057] N=808	-0.005 [0.066] N=670	-0.108 [0.091] N=518
Meat fish or poultry consumed at least weekly	-0.002 [0.063] N=1,015	0.008 [0.067] N=924	0.003 [0.079] N=808	-0.065 [0.084] N=670	**-0.186** [0.088] N=518**
Legumes consumed at least weekly	-0.045 [0.073] N=1,015	-0.107 [0.076] N=924	-0.069 [0.077] N=808	-0.074 [0.090] N=670	-0.092 [0.129] N=518
Animal products consumed at least weekly	0.022 [0.050] N=1,015	0.013 [0.057] N=924	-0.039 [0.064] N=808	0.002 [0.075] N=670	-0.040 [0.092] N=518
Sweets consumed at least weekly	-0.044 [0.069] N=1,015	-0.043 [0.071] N=924	-0.022 [0.079] N=808	-0.094 [0.090] N=670	-0.009 [0.127] N=518
Snack foods consumed at least weekly	-0.010 [0.060] N=1,015	-0.036 [0.065] N=924	0.015 [0.070] N=808	0.005 [0.094] N=670	-0.003 [0.115] N=518
Packaged soft drinks consumed at least weekly	0.009 [0.067] N=1,015	-0.032 [0.075] N=924	-0.022 [0.077] N=808	-0.029 [0.088] N=670	-0.123 [0.118] N=518
Grains consumed at least weekly	0.059 [0.058] N=1,015	0.058 [0.063] N=924	0.053 [0.069] N=808	0.098 [0.083] N=670	0.052 [0.118] N=518
Child was ever breastfed	-0.007 [0.020] N=1,203	-0.005 [0.023] N=1,098	-0.008 [0.028] N=957	-0.001 [0.036] N=787	-0.005 [0.041] N=611

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,106	-0.009 [0.029] N=1,197	0.000 [0.030] N=1,106	-0.012 [0.032] N=990	-0.014 [0.033] N=852	-0.031 [0.037] N=700
Vegetables consumed at least weekly	0.896 P=1,070	-0.011 [0.033] N=1,197	-0.022 [0.033] N=1,106	-0.033 [0.034] N=990	-0.024 [0.035] N=852	-0.060 [0.042] N=700
Meat fish or poultry consumed at least weekly	0.762 P=918	0.026 [0.055] N=1,197	0.042 [0.058] N=1,106	0.043 [0.063] N=990	0.027 [0.069] N=852	0.002 [0.078] N=700
Legumes consumed at least weekly	0.759 P=914	-0.049 [0.050] N=1,197	-0.079 [0.052] N=1,106	-0.065 [0.052] N=990	-0.058 [0.057] N=852	-0.064 [0.066] N=700
Animal products consumed at least weekly	0.904 P=1,085	0.007 [0.039] N=1,197	0.001 [0.041] N=1,106	-0.023 [0.044] N=990	-0.014 [0.048] N=852	-0.034 [0.054] N=700
Sweets consumed at least weekly	0.533 P=635	-0.025 [0.053] N=1,197	-0.024 [0.055] N=1,106	-0.016 [0.059] N=990	-0.032 [0.066] N=852	-0.003 [0.076] N=700
Snack foods consumed at least weekly	0.439 P=519	0.000 [0.046] N=1,197	-0.010 [0.049] N=1,106	0.009 [0.051] N=990	0.012 [0.057] N=852	0.010 [0.061] N=700
Packaged soft drinks consumed at least weekly	0.248 P=291	-0.008 [0.045] N=1,197	-0.034 [0.049] N=1,106	-0.033 [0.050] N=990	-0.030 [0.051] N=852	-0.052 [0.058] N=700
Grains consumed at least weekly	0.860 P=1,031	0.021 [0.040] N=1,197	0.018 [0.041] N=1,106	0.011 [0.043] N=990	0.024 [0.045] N=852	0.010 [0.055] N=700
Child was ever breastfed	0.968 P=1,373	-0.021 [0.014] N=1,422	-0.020 [0.015] N=1,317	-0.025 [0.016] N=1,176	-0.023 [0.018] N=1,006	-0.027 [0.018] N=830

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.004 [0.045] N=1,015	0.022 [0.048] N=924	0.000 [0.054] N=808	-0.005 [0.065] N=670	-0.064 [0.083] N=518
Vegetables consumed at least weekly	0.007 [0.049] N=1,015	-0.011 [0.051] N=924	-0.036 [0.057] N=808	-0.020 [0.064] N=670	-0.143 [0.091] N=518
Meat fish or poultry consumed at least weekly	-0.006 [0.064] N=1,015	0.012 [0.069] N=924	0.001 [0.081] N=808	-0.070 [0.086] N=670	**-0.227** [0.097] N=518**
Legumes consumed at least weekly	-0.045 [0.073] N=1,015	-0.105 [0.077] N=924	-0.073 [0.077] N=808	-0.072 [0.090] N=670	-0.097 [0.135] N=518
Animal products consumed at least weekly	0.020 [0.051] N=1,015	0.013 [0.057] N=924	-0.042 [0.064] N=808	-0.005 [0.074] N=670	-0.061 [0.097] N=518
Sweets consumed at least weekly	-0.035 [0.069] N=1,015	-0.038 [0.071] N=924	-0.018 [0.078] N=808	-0.079 [0.091] N=670	-0.004 [0.127] N=518
Snack foods consumed at least weekly	-0.002 [0.061] N=1,015	-0.027 [0.066] N=924	0.022 [0.071] N=808	0.015 [0.094] N=670	0.002 [0.114] N=518
Packaged soft drinks consumed at least weekly	0.011 [0.066] N=1,015	-0.034 [0.073] N=924	-0.026 [0.076] N=808	-0.026 [0.085] N=670	-0.117 [0.113] N=518
Grains consumed at least weekly	0.064 [0.058] N=1,015	0.060 [0.064] N=924	0.056 [0.069] N=808	0.099 [0.083] N=670	0.039 [0.116] N=518
Child was ever breastfed	-0.008 [0.020] N=1,203	-0.005 [0.023] N=1,098	-0.011 [0.029] N=957	-0.004 [0.039] N=787	-0.003 [0.043] N=611

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,015	0.003 [0.030] N=1,098	0.005 [0.030] N=1,018	-0.003 [0.032] N=911	-0.005 [0.034] N=786	-0.026 [0.037] N=650
Vegetables consumed at least weekly	0.897 P=982	-0.007 [0.035] N=1,098	-0.026 [0.035] N=1,018	-0.037 [0.036] N=911	-0.023 [0.038] N=786	-0.057 [0.042] N=650
Meat fish or poultry consumed at least weekly	0.762 P=840	0.050 [0.053] N=1,098	0.048 [0.055] N=1,018	0.047 [0.058] N=911	0.021 [0.065] N=786	-0.004 [0.072] N=650
Legumes consumed at least weekly	0.755 P=832	-0.061 [0.051] N=1,098	**-0.100* [0.052] N=1,018**	-0.084 [0.051] N=911	-0.081 [0.056] N=786	-0.088 [0.064] N=650
Animal products consumed at least weekly	0.906 P=997	0.020 [0.040] N=1,098	0.010 [0.043] N=1,018	-0.015 [0.046] N=911	-0.012 [0.050] N=786	-0.032 [0.056] N=650
Sweets consumed at least weekly	0.535 P=582	-0.001 [0.055] N=1,098	-0.007 [0.058] N=1,018	0.005 [0.061] N=911	-0.011 [0.070] N=786	0.026 [0.081] N=650
Snack foods consumed at least weekly	0.440 P=475	-0.005 [0.049] N=1,098	-0.015 [0.053] N=1,018	0.010 [0.055] N=911	0.011 [0.062] N=786	0.028 [0.067] N=650
Packaged soft drinks consumed at least weekly	0.250 P=268	0.010 [0.045] N=1,098	-0.018 [0.048] N=1,018	-0.028 [0.049] N=911	-0.028 [0.050] N=786	-0.041 [0.057] N=650
Grains consumed at least weekly	0.861 P=946	0.018 [0.040] N=1,098	0.009 [0.041] N=1,018	-0.004 [0.043] N=911	0.013 [0.046] N=786	-0.009 [0.051] N=650
Child was ever breastfed	0.969 P=1,257	**-0.024* [0.014] N=1,301**	-0.021 [0.015] N=1,208	-0.023 [0.016] N=1,076	-0.017 [0.018] N=926	-0.020 [0.017] N=769

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.013 [0.048] N=929	0.016 [0.051] N=849	-0.007 [0.057] N=742	-0.018 [0.069] N=617	-0.074 [0.086] N=481
Vegetables consumed at least weekly	0.011 [0.053] N=929	-0.021 [0.056] N=849	-0.049 [0.061] N=742	-0.024 [0.072] N=617	-0.118 [0.090] N=481
Meat fish or poultry consumed at least weekly	0.026 [0.067] N=929	0.011 [0.070] N=849	0.003 [0.078] N=742	-0.097 [0.086] N=617	**-0.217** [0.095] N=481**
Legumes consumed at least weekly	-0.050 [0.077] N=929	-0.128 [0.080] N=849	-0.101 [0.077] N=742	-0.115 [0.092] N=617	-0.143 [0.125] N=481
Animal products consumed at least weekly	0.035 [0.055] N=929	0.017 [0.061] N=849	-0.040 [0.067] N=742	-0.019 [0.080] N=617	-0.074 [0.099] N=481
Sweets consumed at least weekly	-0.022 [0.073] N=929	-0.038 [0.078] N=849	-0.032 [0.086] N=742	-0.110 [0.098] N=617	-0.032 [0.132] N=481
Snack foods consumed at least weekly	-0.020 [0.066] N=929	-0.046 [0.072] N=849	-0.003 [0.078] N=742	-0.029 [0.101] N=617	-0.012 [0.120] N=481
Packaged soft drinks consumed at least weekly	0.032 [0.068] N=929	-0.020 [0.075] N=849	-0.036 [0.079] N=742	-0.049 [0.088] N=617	-0.098 [0.120] N=481
Grains consumed at least weekly	0.065 [0.060] N=929	0.052 [0.067] N=849	0.029 [0.072] N=742	0.077 [0.086] N=617	0.026 [0.120] N=481
Child was ever breastfed	-0.022 [0.022] N=1,098	-0.015 [0.025] N=1,005	-0.020 [0.031] N=873	-0.014 [0.039] N=723	-0.013 [0.045] N=566

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,015	-0.001 [0.029] N=1,098	0.000 [0.029] N=1,018	-0.006 [0.032] N=911	-0.008 [0.034] N=786	-0.029 [0.036] N=650
Vegetables consumed at least weekly	0.897 P=982	-0.010 [0.034] N=1,098	-0.028 [0.034] N=1,018	-0.039 [0.036] N=911	-0.025 [0.037] N=786	-0.059 [0.042] N=650
Meat fish or poultry consumed at least weekly	0.762 P=840	0.043 [0.053] N=1,098	0.043 [0.055] N=1,018	0.045 [0.058] N=911	0.021 [0.065] N=786	0.001 [0.072] N=650
Legumes consumed at least weekly	0.755 P=832	-0.062 [0.051] N=1,098	**-0.101* [0.052] N=1,018**	**-0.087* [0.051] N=911**	-0.086 [0.056] N=786	-0.096 [0.065] N=650
Animal products consumed at least weekly	0.906 P=997	0.018 [0.039] N=1,098	0.008 [0.042] N=1,018	-0.017 [0.045] N=911	-0.011 [0.049] N=786	-0.031 [0.055] N=650
Sweets consumed at least weekly	0.535 P=582	0.003 [0.054] N=1,098	-0.003 [0.058] N=1,018	0.007 [0.061] N=911	-0.009 [0.069] N=786	0.026 [0.079] N=650
Snack foods consumed at least weekly	0.440 P=475	-0.005 [0.049] N=1,098	-0.016 [0.052] N=1,018	0.011 [0.054] N=911	0.012 [0.062] N=786	0.028 [0.066] N=650
Packaged soft drinks consumed at least weekly	0.250 P=268	0.012 [0.045] N=1,098	-0.017 [0.048] N=1,018	-0.026 [0.049] N=911	-0.023 [0.050] N=786	-0.040 [0.056] N=650
Grains consumed at least weekly	0.861 P=946	0.017 [0.039] N=1,098	0.009 [0.041] N=1,018	-0.005 [0.043] N=911	0.013 [0.045] N=786	-0.010 [0.050] N=650
Child was ever breastfed	0.969 P=1,257	**-0.023* [0.014] N=1,301**	-0.019 [0.015] N=1,208	-0.022 [0.016] N=1,076	-0.016 [0.018] N=926	-0.020 [0.017] N=769

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.005 [0.047] N=929	0.008 [0.050] N=849	-0.009 [0.056] N=742	-0.019 [0.068] N=617	-0.078 [0.082] N=481
Vegetables consumed at least weekly	0.006 [0.052] N=929	-0.025 [0.056] N=849	-0.052 [0.060] N=742	-0.028 [0.070] N=617	-0.130 [0.087] N=481
Meat fish or poultry consumed at least weekly	0.017 [0.068] N=929	0.006 [0.071] N=849	0.002 [0.079] N=742	-0.099 [0.087] N=617	**-0.222** [0.096] N=481**
Legumes consumed at least weekly	-0.055 [0.076] N=929	**-0.133* [0.079] N=849**	-0.102 [0.077] N=742	-0.117 [0.092] N=617	-0.150 [0.125] N=481
Animal products consumed at least weekly	0.028 [0.053] N=929	0.010 [0.060] N=849	-0.043 [0.066] N=742	-0.019 [0.077] N=617	-0.079 [0.096] N=481
Sweets consumed at least weekly	-0.026 [0.072] N=929	-0.044 [0.077] N=849	-0.038 [0.085] N=742	-0.114 [0.096] N=617	-0.043 [0.129] N=481
Snack foods consumed at least weekly	-0.027 [0.066] N=929	-0.054 [0.071] N=849	-0.008 [0.077] N=742	-0.033 [0.101] N=617	-0.020 [0.119] N=481
Packaged soft drinks consumed at least weekly	0.026 [0.070] N=929	-0.026 [0.076] N=849	-0.040 [0.080] N=742	-0.046 [0.090] N=617	-0.097 [0.122] N=481
Grains consumed at least weekly	0.061 [0.059] N=929	0.050 [0.065] N=849	0.030 [0.071] N=742	0.078 [0.086] N=617	0.025 [0.119] N=481
Child was ever breastfed	-0.019 [0.021] N=1,098	-0.012 [0.025] N=1,005	-0.017 [0.030] N=873	-0.012 [0.039] N=723	-0.015 [0.045] N=566

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.931 P=946	**0.074* [0.038] N=1,018**	**0.074* [0.039] N=917**	0.053 [0.043] N=805	0.040 [0.045] N=673	0.053 [0.054] N=548
Vegetables consumed at least weekly	0.896 P=911	0.061 [0.042] N=1,018	0.067 [0.044] N=917	0.071 [0.047] N=805	0.053 [0.051] N=673	0.069 [0.054] N=548
Meat fish or poultry consumed at least weekly	0.757 P=779	0.097 [0.063] N=1,018	0.093 [0.064] N=917	0.085 [0.071] N=805	**0.146* [0.075] N=673**	0.145 [0.088] N=548
Legumes consumed at least weekly	0.754 P=778	-0.004 [0.048] N=1,018	0.036 [0.050] N=917	0.030 [0.057] N=805	-0.002 [0.060] N=673	0.035 [0.070] N=548
Animal products consumed at least weekly	0.904 P=923	0.014 [0.047] N=1,018	0.010 [0.050] N=917	0.001 [0.058] N=805	-0.013 [0.063] N=673	-0.005 [0.073] N=548
Sweets consumed at least weekly	0.533 P=544	-0.063 [0.064] N=1,018	-0.078 [0.067] N=917	-0.087 [0.071] N=805	-0.087 [0.073] N=673	**-0.154* [0.083] N=548**
Snack foods consumed at least weekly	0.439 P=445	-0.030 [0.057] N=1,018	-0.032 [0.060] N=917	-0.008 [0.063] N=805	-0.027 [0.067] N=673	-0.080 [0.075] N=548
Packaged soft drinks consumed at least weekly	0.233 P=234	-0.053 [0.048] N=1,018	-0.065 [0.050] N=917	-0.002 [0.052] N=805	-0.019 [0.055] N=673	-0.015 [0.060] N=548
Grains consumed at least weekly	0.849 P=869	0.059 [0.047] N=1,018	0.066 [0.051] N=917	0.067 [0.053] N=805	0.069 [0.059] N=673	0.054 [0.071] N=548
Child was ever breastfed	0.967 P=1,172	0.001 [0.023] N=1,214	-0.004 [0.024] N=1,096	0.009 [0.023] N=958	0.010 [0.024] N=807	0.014 [0.024] N=653

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.054] N=887	0.057 [0.055] N=786	0.026 [0.066] N=674	0.018 [0.074] N=542	0.048 [0.108] N=417
Vegetables consumed at least weekly	0.005 [0.063] N=887	0.006 [0.068] N=786	-0.001 [0.080] N=674	-0.036 [0.095] N=542	-0.019 [0.119] N=417
Meat fish or poultry consumed at least weekly	0.012 [0.075] N=887	-0.004 [0.077] N=786	-0.056 [0.087] N=674	0.039 [0.094] N=542	-0.008 [0.129] N=417
Legumes consumed at least weekly	-0.048 [0.060] N=887	0.006 [0.065] N=786	0.000 [0.076] N=674	-0.044 [0.084] N=542	0.035 [0.103] N=417
Animal products consumed at least weekly	-0.024 [0.058] N=887	-0.032 [0.064] N=786	-0.049 [0.078] N=674	-0.076 [0.080] N=542	-0.088 [0.102] N=417
Sweets consumed at least weekly	-0.033 [0.090] N=887	-0.054 [0.096] N=786	-0.076 [0.106] N=674	-0.072 [0.124] N=542	-0.247 [0.162] N=417
Snack foods consumed at least weekly	-0.037 [0.082] N=887	-0.041 [0.089] N=786	0.004 [0.096] N=674	-0.018 [0.112] N=542	-0.176 [0.142] N=417
Packaged soft drinks consumed at least weekly	-0.074 [0.075] N=887	-0.098 [0.080] N=786	0.004 [0.093] N=674	-0.028 [0.108] N=542	-0.041 [0.127] N=417
Grains consumed at least weekly	-0.052 [0.051] N=887	-0.061 [0.057] N=786	-0.081 [0.058] N=674	**-0.147** [0.062] N=542**	**-0.313*** [0.080] N=417**
Child was ever breastfed	0.009 [0.043] N=1,058	0.002 [0.045] N=940	0.027 [0.047] N=802	0.031 [0.053] N=651	0.050 [0.067] N=497

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.931 P=946	**0.074* [0.038] N=1,018**	**0.074* [0.039] N=917**	0.053 [0.043] N=805	0.039 [0.046] N=673	0.053 [0.055] N=548
Vegetables consumed at least weekly	0.896 P=911	0.061 [0.041] N=1,018	0.067 [0.043] N=917	0.071 [0.047] N=805	0.054 [0.050] N=673	0.069 [0.053] N=548
Meat fish or poultry consumed at least weekly	0.757 P=779	0.097 [0.063] N=1,018	0.094 [0.064] N=917	0.085 [0.071] N=805	**0.146* [0.075] N=673**	0.144 [0.088] N=548
Legumes consumed at least weekly	0.754 P=778	-0.004 [0.048] N=1,018	0.036 [0.050] N=917	0.030 [0.057] N=805	-0.002 [0.060] N=673	0.035 [0.071] N=548
Animal products consumed at least weekly	0.904 P=923	0.014 [0.045] N=1,018	0.010 [0.049] N=917	0.002 [0.057] N=805	-0.012 [0.062] N=673	-0.005 [0.071] N=548
Sweets consumed at least weekly	0.533 P=544	-0.063 [0.064] N=1,018	-0.078 [0.067] N=917	-0.086 [0.071] N=805	-0.086 [0.073] N=673	**-0.154* [0.083] N=548**
Snack foods consumed at least weekly	0.439 P=445	-0.030 [0.056] N=1,018	-0.032 [0.060] N=917	-0.007 [0.062] N=805	-0.026 [0.066] N=673	-0.080 [0.074] N=548
Packaged soft drinks consumed at least weekly	0.233 P=234	-0.053 [0.048] N=1,018	-0.065 [0.050] N=917	-0.002 [0.052] N=805	-0.019 [0.055] N=673	-0.015 [0.060] N=548
Grains consumed at least weekly	0.849 P=869	0.059 [0.047] N=1,018	0.067 [0.050] N=917	0.067 [0.053] N=805	0.069 [0.059] N=673	0.054 [0.071] N=548
Child was ever breastfed	0.967 P=1,172	0.001 [0.024] N=1,214	-0.004 [0.024] N=1,096	0.009 [0.023] N=958	0.009 [0.024] N=807	0.013 [0.024] N=653

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.059 [0.053] N=887	0.058 [0.055] N=786	0.026 [0.066] N=674	0.017 [0.075] N=542	0.047 [0.110] N=417
Vegetables consumed at least weekly	0.005 [0.062] N=887	0.006 [0.067] N=786	-0.001 [0.079] N=674	-0.033 [0.094] N=542	-0.018 [0.119] N=417
Meat fish or poultry consumed at least weekly	0.012 [0.075] N=887	-0.004 [0.076] N=786	-0.054 [0.086] N=674	0.040 [0.094] N=542	-0.008 [0.129] N=417
Legumes consumed at least weekly	-0.048 [0.060] N=887	0.007 [0.064] N=786	0.001 [0.075] N=674	-0.045 [0.085] N=542	0.030 [0.105] N=417
Animal products consumed at least weekly	-0.024 [0.054] N=887	-0.031 [0.061] N=786	-0.046 [0.075] N=674	-0.069 [0.076] N=542	-0.081 [0.094] N=417
Sweets consumed at least weekly	-0.033 [0.088] N=887	-0.053 [0.094] N=786	-0.074 [0.105] N=674	-0.072 [0.124] N=542	-0.247 [0.162] N=417
Snack foods consumed at least weekly	-0.037 [0.080] N=887	-0.040 [0.086] N=786	0.007 [0.094] N=674	-0.014 [0.111] N=542	-0.173 [0.140] N=417
Packaged soft drinks consumed at least weekly	-0.075 [0.074] N=887	-0.098 [0.080] N=786	0.005 [0.092] N=674	-0.028 [0.108] N=542	-0.039 [0.126] N=417
Grains consumed at least weekly	-0.052 [0.051] N=887	-0.061 [0.057] N=786	-0.080 [0.058] N=674	**-0.150** [0.062] N=542**	**-0.315*** [0.082] N=417**
Child was ever breastfed	0.009 [0.043] N=1,058	0.002 [0.045] N=940	0.026 [0.047] N=802	0.029 [0.053] N=651	0.049 [0.067] N=497

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.931 P=946	**0.071* [0.038] N=1,018**	**0.071* [0.039] N=917**	0.053 [0.043] N=805	0.040 [0.047] N=673	0.054 [0.056] N=548
Vegetables consumed at least weekly	0.896 P=911	0.059 [0.041] N=1,018	0.065 [0.043] N=917	0.071 [0.047] N=805	0.054 [0.051] N=673	0.067 [0.054] N=548
Meat fish or poultry consumed at least weekly	0.757 P=779	0.089 [0.060] N=1,018	0.086 [0.061] N=917	0.088 [0.068] N=805	**0.150** [0.072] N=673**	**0.149* [0.084] N=548**
Legumes consumed at least weekly	0.754 P=778	-0.008 [0.049] N=1,018	0.033 [0.051] N=917	0.028 [0.057] N=805	-0.003 [0.061] N=673	0.035 [0.073] N=548
Animal products consumed at least weekly	0.904 P=923	0.011 [0.045] N=1,018	0.005 [0.049] N=917	-0.001 [0.056] N=805	-0.013 [0.062] N=673	-0.005 [0.071] N=548
Sweets consumed at least weekly	0.533 P=544	-0.065 [0.064] N=1,018	-0.083 [0.068] N=917	-0.090 [0.071] N=805	-0.089 [0.073] N=673	**-0.159* [0.084] N=548**
Snack foods consumed at least weekly	0.439 P=445	-0.029 [0.056] N=1,018	-0.031 [0.060] N=917	-0.009 [0.062] N=805	-0.027 [0.067] N=673	-0.083 [0.074] N=548
Packaged soft drinks consumed at least weekly	0.233 P=234	-0.050 [0.048] N=1,018	-0.065 [0.050] N=917	-0.008 [0.053] N=805	-0.023 [0.056] N=673	-0.022 [0.062] N=548
Grains consumed at least weekly	0.849 P=869	0.056 [0.047] N=1,018	0.064 [0.050] N=917	0.067 [0.053] N=805	0.070 [0.060] N=673	0.054 [0.072] N=548
Child was ever breastfed	0.967 P=1,172	-0.000 [0.024] N=1,214	-0.005 [0.024] N=1,096	0.010 [0.024] N=958	0.010 [0.024] N=807	0.012 [0.024] N=653

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.052 [0.053] N=887	0.049 [0.054] N=786	0.026 [0.065] N=674	0.013 [0.076] N=542	0.038 [0.112] N=417
Vegetables consumed at least weekly	0.000 [0.061] N=887	0.000 [0.066] N=786	-0.003 [0.077] N=674	-0.038 [0.093] N=542	-0.031 [0.119] N=417
Meat fish or poultry consumed at least weekly	-0.008 [0.074] N=887	-0.023 [0.076] N=786	-0.055 [0.083] N=674	0.028 [0.096] N=542	-0.031 [0.129] N=417
Legumes consumed at least weekly	-0.050 [0.060] N=887	0.006 [0.065] N=786	0.001 [0.074] N=674	-0.050 [0.085] N=542	0.028 [0.108] N=417
Animal products consumed at least weekly	-0.027 [0.054] N=887	-0.036 [0.060] N=786	-0.046 [0.073] N=674	-0.072 [0.077] N=542	-0.086 [0.095] N=417
Sweets consumed at least weekly	-0.031 [0.086] N=887	-0.054 [0.093] N=786	-0.072 [0.102] N=674	-0.067 [0.123] N=542	-0.236 [0.159] N=417
Snack foods consumed at least weekly	-0.032 [0.079] N=887	-0.034 [0.086] N=786	0.006 [0.094] N=674	-0.012 [0.110] N=542	-0.171 [0.139] N=417
Packaged soft drinks consumed at least weekly	-0.060 [0.073] N=887	-0.084 [0.079] N=786	0.009 [0.092] N=674	-0.017 [0.106] N=542	-0.020 [0.123] N=417
Grains consumed at least weekly	-0.058 [0.051] N=887	-0.066 [0.057] N=786	-0.079 [0.057] N=674	**-0.151** [0.061] N=542**	**-0.315*** [0.080] N=417**
Child was ever breastfed	0.007 [0.042] N=1,058	-0.002 [0.044] N=940	0.026 [0.046] N=802	0.027 [0.051] N=651	0.043 [0.064] N=497

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.931 P=946	**0.067* [0.038] N=1,018**	**0.068* [0.039] N=917**	0.051 [0.043] N=805	0.030 [0.046] N=673	0.041 [0.055] N=548
Vegetables consumed at least weekly	0.896 P=911	0.053 [0.042] N=1,018	0.060 [0.044] N=917	0.064 [0.047] N=805	0.041 [0.049] N=673	0.055 [0.051] N=548
Meat fish or poultry consumed at least weekly	0.757 P=779	0.082 [0.060] N=1,018	0.077 [0.062] N=917	0.073 [0.069] N=805	**0.121* [0.072] N=673**	0.107 [0.082] N=548
Legumes consumed at least weekly	0.754 P=778	-0.006 [0.049] N=1,018	0.034 [0.051] N=917	0.034 [0.056] N=805	-0.005 [0.060] N=673	0.028 [0.069] N=548
Animal products consumed at least weekly	0.904 P=923	0.015 [0.044] N=1,018	0.012 [0.048] N=917	0.010 [0.055] N=805	-0.008 [0.061] N=673	-0.004 [0.070] N=548
Sweets consumed at least weekly	0.533 P=544	-0.055 [0.066] N=1,018	-0.073 [0.069] N=917	-0.082 [0.073] N=805	-0.085 [0.075] N=673	**-0.155* [0.085] N=548**
Snack foods consumed at least weekly	0.439 P=445	-0.021 [0.057] N=1,018	-0.024 [0.061] N=917	0.001 [0.064] N=805	-0.022 [0.067] N=673	-0.072 [0.076] N=548
Packaged soft drinks consumed at least weekly	0.233 P=234	-0.040 [0.048] N=1,018	-0.057 [0.050] N=917	0.005 [0.053] N=805	-0.012 [0.056] N=673	0.002 [0.063] N=548
Grains consumed at least weekly	0.849 P=869	0.054 [0.047] N=1,018	0.060 [0.051] N=917	0.063 [0.053] N=805	0.062 [0.060] N=673	0.045 [0.071] N=548
Child was ever breastfed	0.967 P=1,172	0.001 [0.023] N=1,214	-0.004 [0.023] N=1,096	0.010 [0.022] N=958	0.010 [0.023] N=807	0.013 [0.023] N=653

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.050 [0.052] N=887	0.047 [0.054] N=786	0.026 [0.065] N=674	-0.003 [0.076] N=542	0.009 [0.108] N=417
Vegetables consumed at least weekly	-0.008 [0.062] N=887	-0.008 [0.066] N=786	-0.011 [0.078] N=674	-0.060 [0.091] N=542	-0.066 [0.114] N=417
Meat fish or poultry consumed at least weekly	-0.009 [0.076] N=887	-0.027 [0.078] N=786	-0.069 [0.086] N=674	-0.012 [0.101] N=542	-0.106 [0.126] N=417
Legumes consumed at least weekly	-0.051 [0.062] N=887	0.002 [0.067] N=786	0.012 [0.076] N=674	-0.048 [0.086] N=542	0.013 [0.106] N=417
Animal products consumed at least weekly	-0.026 [0.051] N=887	-0.033 [0.057] N=786	-0.033 [0.071] N=674	-0.068 [0.075] N=542	-0.092 [0.090] N=417
Sweets consumed at least weekly	-0.019 [0.087] N=887	-0.043 [0.093] N=786	-0.057 [0.103] N=674	-0.052 [0.122] N=542	-0.219 [0.157] N=417
Snack foods consumed at least weekly	-0.029 [0.080] N=887	-0.033 [0.087] N=786	0.020 [0.095] N=674	-0.007 [0.109] N=542	-0.157 [0.138] N=417
Packaged soft drinks consumed at least weekly	-0.061 [0.073] N=887	-0.090 [0.079] N=786	0.009 [0.090] N=674	-0.013 [0.105] N=542	0.013 [0.125] N=417
Grains consumed at least weekly	-0.057 [0.051] N=887	-0.068 [0.056] N=786	-0.081 [0.058] N=674	**-0.163** [0.063] N=542**	**-0.334*** [0.080] N=417**
Child was ever breastfed	0.008 [0.041] N=1,058	0.001 [0.043] N=940	0.030 [0.044] N=802	0.028 [0.049] N=651	0.039 [0.062] N=497

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=865	**0.070* [0.041] N=929**	0.066 [0.041] N=834	0.054 [0.046] N=739	0.024 [0.050] N=617	0.031 [0.060] N=507
Vegetables consumed at least weekly	0.899 P=834	0.049 [0.042] N=929	0.051 [0.045] N=834	0.073 [0.047] N=739	0.040 [0.051] N=617	0.030 [0.055] N=507
Meat fish or poultry consumed at least weekly	0.757 P=710	0.075 [0.065] N=929	0.061 [0.066] N=834	0.065 [0.074] N=739	0.106 [0.080] N=617	0.106 [0.094] N=507
Legumes consumed at least weekly	0.750 P=707	-0.014 [0.055] N=929	0.027 [0.058] N=834	0.021 [0.064] N=739	-0.027 [0.069] N=617	0.011 [0.081] N=507
Animal products consumed at least weekly	0.907 P=845	0.009 [0.049] N=929	0.007 [0.054] N=834	0.008 [0.062] N=739	-0.023 [0.069] N=617	-0.027 [0.080] N=507
Sweets consumed at least weekly	0.544 P=506	-0.056 [0.066] N=929	-0.070 [0.069] N=834	-0.089 [0.074] N=739	-0.105 [0.078] N=617	**-0.174* [0.089] N=507**
Snack foods consumed at least weekly	0.447 P=412	-0.026 [0.059] N=929	-0.029 [0.063] N=834	-0.017 [0.065] N=739	-0.047 [0.070] N=617	-0.088 [0.079] N=507
Packaged soft drinks consumed at least weekly	0.239 P=220	-0.060 [0.049] N=929	-0.080 [0.050] N=834	-0.016 [0.053] N=739	-0.041 [0.058] N=617	-0.050 [0.064] N=507
Grains consumed at least weekly	0.850 P=794	0.072 [0.050] N=929	0.079 [0.054] N=834	0.086 [0.057] N=739	0.076 [0.065] N=617	0.048 [0.077] N=507
Child was ever breastfed	0.969 P=1,072	-0.010 [0.022] N=1,107	-0.014 [0.023] N=998	0.002 [0.021] N=879	-0.003 [0.022] N=741	0.006 [0.022] N=604

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.054 [0.057] N=808	0.049 [0.057] N=713	0.034 [0.067] N=618	-0.008 [0.077] N=496	-0.008 [0.108] N=386
Vegetables consumed at least weekly	-0.010 [0.057] N=808	-0.015 [0.063] N=713	0.009 [0.069] N=618	-0.052 [0.081] N=496	-0.100 [0.101] N=386
Meat fish or poultry consumed at least weekly	0.008 [0.083] N=808	-0.017 [0.085] N=713	-0.048 [0.094] N=618	-0.004 [0.106] N=496	-0.068 [0.141] N=386
Legumes consumed at least weekly	-0.052 [0.067] N=808	0.005 [0.071] N=713	0.008 [0.080] N=618	-0.056 [0.091] N=496	0.008 [0.111] N=386
Animal products consumed at least weekly	-0.037 [0.058] N=808	-0.039 [0.065] N=713	-0.037 [0.077] N=618	-0.093 [0.085] N=496	-0.141 [0.106] N=386
Sweets consumed at least weekly	-0.022 [0.092] N=808	-0.041 [0.100] N=713	-0.092 [0.108] N=618	-0.126 [0.126] N=496	**-0.324* [0.166] N=386**
Snack foods consumed at least weekly	-0.033 [0.085] N=808	-0.037 [0.094] N=713	-0.020 [0.100] N=618	-0.079 [0.114] N=496	-0.211 [0.147] N=386
Packaged soft drinks consumed at least weekly	-0.084 [0.076] N=808	-0.120 [0.082] N=713	-0.024 [0.096] N=618	-0.080 [0.114] N=496	-0.161 [0.135] N=386
Grains consumed at least weekly	-0.035 [0.060] N=808	-0.042 [0.068] N=713	-0.047 [0.071] N=618	**-0.143* [0.081] N=496**	**-0.364*** [0.102] N=386**
Child was ever breastfed	-0.007 [0.040] N=963	-0.016 [0.043] N=854	0.014 [0.042] N=735	0.006 [0.048] N=597	0.030 [0.060] N=460

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=865	**0.073* [0.040] N=929**	**0.071* [0.041] N=834**	0.060 [0.046] N=739	0.025 [0.049] N=617	0.032 [0.059] N=507
Vegetables consumed at least weekly	0.899 P=834	0.054 [0.041] N=929	0.055 [0.044] N=834	**0.078* [0.046] N=739**	0.040 [0.050] N=617	0.030 [0.054] N=507
Meat fish or poultry consumed at least weekly	0.757 P=710	0.087 [0.063] N=929	0.074 [0.065] N=834	0.074 [0.072] N=739	0.103 [0.076] N=617	0.102 [0.090] N=507
Legumes consumed at least weekly	0.750 P=707	-0.015 [0.054] N=929	0.027 [0.057] N=834	0.024 [0.063] N=739	-0.031 [0.067] N=617	0.009 [0.079] N=507
Animal products consumed at least weekly	0.907 P=845	0.008 [0.049] N=929	0.007 [0.053] N=834	0.008 [0.061] N=739	-0.022 [0.067] N=617	-0.025 [0.078] N=507
Sweets consumed at least weekly	0.544 P=506	-0.060 [0.065] N=929	-0.073 [0.069] N=834	-0.095 [0.073] N=739	-0.112 [0.078] N=617	**-0.178* [0.090] N=507**
Snack foods consumed at least weekly	0.447 P=412	-0.021 [0.058] N=929	-0.025 [0.063] N=834	-0.012 [0.064] N=739	-0.042 [0.069] N=617	-0.082 [0.078] N=507
Packaged soft drinks consumed at least weekly	0.239 P=220	-0.062 [0.049] N=929	-0.081 [0.051] N=834	-0.018 [0.053] N=739	-0.043 [0.057] N=617	-0.052 [0.063] N=507
Grains consumed at least weekly	0.850 P=794	0.073 [0.050] N=929	0.080 [0.054] N=834	0.085 [0.056] N=739	0.074 [0.065] N=617	0.047 [0.077] N=507
Child was ever breastfed	0.969 P=1,072	-0.009 [0.023] N=1,107	-0.014 [0.023] N=998	0.004 [0.022] N=879	0.001 [0.022] N=741	0.010 [0.022] N=604

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.056] N=808	0.058 [0.057] N=713	0.048 [0.068] N=618	-0.008 [0.078] N=496	-0.001 [0.114] N=386
Vegetables consumed at least weekly	-0.006 [0.057] N=808	-0.011 [0.063] N=713	0.017 [0.070] N=618	-0.058 [0.081] N=496	-0.108 [0.104] N=386
Meat fish or poultry consumed at least weekly	0.022 [0.082] N=808	-0.005 [0.084] N=713	-0.038 [0.093] N=618	-0.012 [0.103] N=496	-0.070 [0.143] N=386
Legumes consumed at least weekly	-0.050 [0.067] N=808	0.009 [0.071] N=713	0.014 [0.080] N=618	-0.070 [0.090] N=496	0.012 [0.109] N=386
Animal products consumed at least weekly	-0.036 [0.058] N=808	-0.037 [0.065] N=713	-0.035 [0.077] N=618	-0.096 [0.084] N=496	-0.147 [0.109] N=386
Sweets consumed at least weekly	-0.021 [0.092] N=808	-0.041 [0.100] N=713	-0.091 [0.110] N=618	-0.125 [0.128] N=496	**-0.326* [0.170] N=386**
Snack foods consumed at least weekly	-0.030 [0.085] N=808	-0.036 [0.094] N=713	-0.012 [0.101] N=618	-0.061 [0.117] N=496	-0.205 [0.151] N=386
Packaged soft drinks consumed at least weekly	-0.085 [0.078] N=808	-0.118 [0.084] N=713	-0.013 [0.095] N=618	-0.066 [0.112] N=496	-0.131 [0.133] N=386
Grains consumed at least weekly	-0.034 [0.060] N=808	-0.041 [0.067] N=713	-0.049 [0.068] N=618	**-0.150* [0.078] N=496**	**-0.369*** [0.099] N=386**
Child was ever breastfed	-0.008 [0.041] N=963	-0.018 [0.043] N=854	0.015 [0.043] N=735	0.011 [0.050] N=597	0.032 [0.063] N=460

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.