Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.733	0.113 [0.266] N=1,220	0.054 [0.284] N=1,133	0.135 [0.310] N=1,011	0.191 [0.327] N=869	0.037 [0.355] N=721
WHO weight-for-age z score	-0.259	-0.122 [0.136] N=1,407	-0.104 [0.149] N=1,304	-0.078 [0.155] N=1,165	-0.007 [0.154] N=997	-0.089 [0.164] N=823
WHO weight-for-length/height z score, 0-60 months	0.306	0.322 [0.235] N=1,216	0.340 [0.245] N=1,127	0.293 [0.259] N=1,009	0.347 [0.273] N=865	0.405 [0.298] N=714
WHO BMI-for-age z score	0.303	0.389 [0.269] N=1,196	0.429 [0.283] N=1,110	0.390 [0.304] N=991	0.305 [0.331] N=849	0.332 [0.367] N=704
Stunted: HAZ below -2	0.301 P=359	-0.031 [0.047] N=1,220	-0.028 [0.050] N=1,133	-0.022 [0.054] N=1,011	-0.032 [0.058] N=869	-0.003 [0.061] N=721
Underweight: WAZ below -2	0.109 P=150	**0.083*** [0.027] N=1,407**	**0.069** [0.028] N=1,304**	**0.058* [0.030] N=1,165**	**0.052* [0.030] N=997**	**0.060* [0.034] N=823**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=147	-0.045 [0.039] N=1,216	-0.037 [0.041] N=1,127	-0.042 [0.043] N=1,009	-0.031 [0.047] N=865	-0.049 [0.048] N=714
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=236	0.048 [0.048] N=1,216	0.054 [0.051] N=1,127	0.034 [0.054] N=1,009	0.054 [0.060] N=865	0.044 [0.066] N=714
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=107	0.006 [0.035] N=1,216	0.011 [0.038] N=1,127	0.017 [0.040] N=1,009	0.037 [0.043] N=865	0.044 [0.048] N=714
CIAF: any stunting, underweight, or wasting, 0-60 months	0.398 P=456	-0.061 [0.048] N=1,165	-0.053 [0.051] N=1,080	-0.044 [0.054] N=966	-0.058 [0.060] N=830	-0.032 [0.066] N=687
BMI-for-age z score below -2	0.138 P=160	-0.037 [0.043] N=1,196	-0.030 [0.045] N=1,110	-0.026 [0.049] N=991	0.016 [0.053] N=849	0.017 [0.057] N=704

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146 [0.381] N=1,028	0.025 [0.432] N=941	0.263 [0.511] N=819	0.540 [0.569] N=677	0.490 [0.722] N=529
WHO weight-for-age z score	**-0.352* [0.191] N=1,190**	-0.352 [0.227] N=1,087	-0.344 [0.245] N=948	-0.274 [0.289] N=780	-0.577 [0.373] N=606
WHO weight-for-length/height z score, 0-60 months	0.190 [0.341] N=1,024	0.196 [0.371] N=935	0.022 [0.403] N=817	0.087 [0.460] N=673	-0.072 [0.560] N=522
WHO BMI-for-age z score	0.298 [0.347] N=1,004	0.325 [0.371] N=918	0.160 [0.415] N=799	-0.126 [0.473] N=657	-0.445 [0.566] N=512
Stunted: HAZ below -2	-0.050 [0.063] N=1,028	-0.048 [0.071] N=941	-0.051 [0.082] N=819	-0.095 [0.095] N=677	-0.098 [0.117] N=529
Underweight: WAZ below -2	**0.114*** [0.040] N=1,190**	**0.093** [0.045] N=1,087**	0.076 [0.050] N=948	0.073 [0.060] N=780	0.105 [0.081] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.052 [0.057] N=1,024	-0.037 [0.063] N=935	-0.043 [0.071] N=817	-0.026 [0.087] N=673	-0.061 [0.107] N=522
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.062] N=1,024	0.021 [0.069] N=935	-0.030 [0.070] N=817	-0.001 [0.090] N=673	-0.098 [0.109] N=522
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.043] N=1,024	-0.016 [0.049] N=935	-0.016 [0.051] N=817	0.028 [0.060] N=673	0.024 [0.080] N=522
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.071 [0.071] N=981	-0.058 [0.078] N=896	-0.055 [0.088] N=782	-0.105 [0.111] N=646	-0.097 [0.151] N=503
BMI-for-age z score below -2	-0.069 [0.059] N=1,004	-0.057 [0.063] N=918	-0.054 [0.073] N=799	0.030 [0.088] N=657	0.041 [0.107] N=512

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.733	0.111 [0.266] N=1,220	0.050 [0.283] N=1,133	0.132 [0.309] N=1,011	0.189 [0.327] N=869	0.036 [0.354] N=721
WHO weight-for-age z score	-0.259	-0.125 [0.136] N=1,407	-0.111 [0.150] N=1,304	-0.083 [0.155] N=1,165	-0.014 [0.154] N=997	-0.105 [0.162] N=823
WHO weight-for-length/height z score, 0-60 months	0.306	0.318 [0.236] N=1,216	0.338 [0.246] N=1,127	0.290 [0.260] N=1,009	0.346 [0.274] N=865	0.397 [0.301] N=714
WHO BMI-for-age z score	0.303	0.390 [0.268] N=1,196	0.430 [0.282] N=1,110	0.390 [0.304] N=991	0.306 [0.331] N=849	0.332 [0.368] N=704
Stunted: HAZ below -2	0.301 P=359	-0.030 [0.047] N=1,220	-0.027 [0.050] N=1,133	-0.021 [0.054] N=1,011	-0.031 [0.058] N=869	-0.000 [0.061] N=721
Underweight: WAZ below -2	0.109 P=150	**0.083*** [0.027] N=1,407**	**0.070** [0.028] N=1,304**	**0.059** [0.030] N=1,165**	**0.053* [0.030] N=997**	**0.063* [0.033] N=823**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=147	-0.043 [0.039] N=1,216	-0.036 [0.041] N=1,127	-0.040 [0.043] N=1,009	-0.029 [0.047] N=865	-0.045 [0.049] N=714
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=236	0.049 [0.048] N=1,216	0.054 [0.051] N=1,127	0.035 [0.054] N=1,009	0.056 [0.060] N=865	0.047 [0.066] N=714
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=107	0.006 [0.035] N=1,216	0.012 [0.037] N=1,127	0.018 [0.040] N=1,009	0.038 [0.043] N=865	0.046 [0.048] N=714
CIAF: any stunting, underweight, or wasting, 0-60 months	0.398 P=456	-0.059 [0.048] N=1,165	-0.050 [0.050] N=1,080	-0.043 [0.054] N=966	-0.056 [0.060] N=830	-0.024 [0.065] N=687
BMI-for-age z score below -2	0.138 P=160	-0.036 [0.044] N=1,196	-0.029 [0.045] N=1,110	-0.025 [0.050] N=991	0.017 [0.054] N=849	0.020 [0.058] N=704

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.146 [0.381] N=1,028	0.028 [0.432] N=941	0.267 [0.510] N=819	0.535 [0.570] N=677	0.491 [0.723] N=529
WHO weight-for-age z score	**-0.343* [0.191] N=1,190**	-0.349 [0.230] N=1,087	-0.335 [0.249] N=948	-0.270 [0.293] N=780	-0.594 [0.376] N=606
WHO weight-for-length/height z score, 0-60 months	0.190 [0.341] N=1,024	0.195 [0.371] N=935	0.023 [0.403] N=817	0.089 [0.461] N=673	-0.084 [0.559] N=522
WHO BMI-for-age z score	0.298 [0.347] N=1,004	0.325 [0.371] N=918	0.164 [0.414] N=799	-0.125 [0.474] N=657	-0.456 [0.565] N=512
Stunted: HAZ below -2	-0.051 [0.063] N=1,028	-0.049 [0.071] N=941	-0.054 [0.082] N=819	-0.096 [0.095] N=677	-0.098 [0.117] N=529
Underweight: WAZ below -2	**0.113*** [0.040] N=1,190**	**0.093** [0.045] N=1,087**	0.075 [0.051] N=948	0.073 [0.061] N=780	0.107 [0.082] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.051 [0.057] N=1,024	-0.036 [0.063] N=935	-0.044 [0.071] N=817	-0.027 [0.087] N=673	-0.057 [0.106] N=522
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.062] N=1,024	0.021 [0.069] N=935	-0.030 [0.070] N=817	-0.001 [0.091] N=673	-0.097 [0.109] N=522
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.043] N=1,024	-0.016 [0.049] N=935	-0.016 [0.051] N=817	0.028 [0.060] N=673	0.025 [0.080] N=522
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.074 [0.072] N=981	-0.062 [0.079] N=896	-0.064 [0.089] N=782	-0.113 [0.114] N=646	-0.098 [0.154] N=503
BMI-for-age z score below -2	-0.069 [0.059] N=1,004	-0.057 [0.063] N=918	-0.055 [0.073] N=799	0.030 [0.089] N=657	0.045 [0.108] N=512

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.733	0.079 [0.268] N=1,220	0.009 [0.285] N=1,133	0.082 [0.310] N=1,011	0.158 [0.327] N=869	-0.021 [0.360] N=721
WHO weight-for-age z score	-0.259	-0.163 [0.129] N=1,407	-0.151 [0.143] N=1,304	-0.127 [0.148] N=1,165	-0.038 [0.148] N=997	-0.151 [0.157] N=823
WHO weight-for-length/height z score, 0-60 months	0.306	0.318 [0.238] N=1,216	0.341 [0.251] N=1,127	0.308 [0.264] N=1,009	0.362 [0.277] N=865	0.403 [0.314] N=714
WHO BMI-for-age z score	0.303	0.398 [0.272] N=1,196	0.439 [0.288] N=1,110	0.411 [0.310] N=991	0.329 [0.337] N=849	0.349 [0.385] N=704
Stunted: HAZ below -2	0.301 P=359	-0.023 [0.047] N=1,220	-0.016 [0.050] N=1,133	-0.010 [0.054] N=1,011	-0.024 [0.057] N=869	0.016 [0.061] N=721
Underweight: WAZ below -2	0.109 P=150	**0.089*** [0.026] N=1,407**	**0.077*** [0.028] N=1,304**	**0.067** [0.029] N=1,165**	**0.057* [0.029] N=997**	**0.073** [0.032] N=823**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=147	-0.042 [0.039] N=1,216	-0.034 [0.041] N=1,127	-0.040 [0.043] N=1,009	-0.030 [0.047] N=865	-0.044 [0.049] N=714
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=236	0.050 [0.048] N=1,216	0.057 [0.051] N=1,127	0.038 [0.054] N=1,009	0.058 [0.059] N=865	0.049 [0.067] N=714
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=107	0.007 [0.036] N=1,216	0.014 [0.038] N=1,127	0.020 [0.040] N=1,009	0.040 [0.043] N=865	0.045 [0.049] N=714
CIAF: any stunting, underweight, or wasting, 0-60 months	0.398 P=456	-0.049 [0.048] N=1,165	-0.034 [0.050] N=1,080	-0.030 [0.053] N=966	-0.046 [0.059] N=830	0.000 [0.065] N=687
BMI-for-age z score below -2	0.138 P=160	-0.036 [0.044] N=1,196	-0.026 [0.046] N=1,110	-0.025 [0.051] N=991	0.015 [0.055] N=849	0.019 [0.060] N=704

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.132 [0.380] N=1,028	-0.001 [0.426] N=941	0.219 [0.498] N=819	0.526 [0.551] N=677	0.503 [0.709] N=529
WHO weight-for-age z score	**-0.347* [0.186] N=1,190**	-0.358 [0.226] N=1,087	-0.347 [0.242] N=948	-0.238 [0.293] N=780	-0.578 [0.376] N=606
WHO weight-for-length/height z score, 0-60 months	0.189 [0.338] N=1,024	0.197 [0.373] N=935	0.047 [0.397] N=817	0.079 [0.449] N=673	-0.139 [0.558] N=522
WHO BMI-for-age z score	0.297 [0.347] N=1,004	0.315 [0.374] N=918	0.170 [0.411] N=799	-0.159 [0.461] N=657	-0.482 [0.567] N=512
Stunted: HAZ below -2	-0.052 [0.063] N=1,028	-0.046 [0.071] N=941	-0.051 [0.081] N=819	-0.109 [0.093] N=677	-0.107 [0.114] N=529
Underweight: WAZ below -2	**0.111*** [0.041] N=1,190**	**0.094** [0.045] N=1,087**	0.075 [0.051] N=948	0.060 [0.062] N=780	0.102 [0.083] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.051 [0.057] N=1,024	-0.036 [0.063] N=935	-0.049 [0.071] N=817	-0.031 [0.086] N=673	-0.053 [0.107] N=522
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.062] N=1,024	0.020 [0.070] N=935	-0.033 [0.070] N=817	-0.012 [0.091] N=673	-0.103 [0.111] N=522
Obese: WLZ/WHZ above +3, 0-60 months	-0.024 [0.043] N=1,024	-0.017 [0.051] N=935	-0.019 [0.051] N=817	0.021 [0.061] N=673	0.021 [0.080] N=522
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.080 [0.072] N=981	-0.064 [0.080] N=896	-0.075 [0.088] N=782	-0.143 [0.113] N=646	-0.114 [0.154] N=503
BMI-for-age z score below -2	-0.070 [0.059] N=1,004	-0.054 [0.063] N=918	-0.059 [0.074] N=799	0.023 [0.088] N=657	0.048 [0.110] N=512

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.733	0.062 [0.267] N=1,220	0.009 [0.284] N=1,133	0.084 [0.311] N=1,011	0.174 [0.329] N=869	0.010 [0.361] N=721
WHO weight-for-age z score	-0.259	-0.180 [0.130] N=1,407	-0.161 [0.144] N=1,304	-0.140 [0.151] N=1,165	-0.074 [0.150] N=997	-0.209 [0.163] N=823
WHO weight-for-length/height z score, 0-60 months	0.306	0.313 [0.239] N=1,216	0.336 [0.251] N=1,127	0.291 [0.265] N=1,009	0.320 [0.279] N=865	0.344 [0.312] N=714
WHO BMI-for-age z score	0.303	0.396 [0.272] N=1,196	0.443 [0.286] N=1,110	0.410 [0.308] N=991	0.296 [0.335] N=849	0.284 [0.380] N=704
Stunted: HAZ below -2	0.301 P=359	-0.017 [0.047] N=1,220	-0.013 [0.050] N=1,133	-0.005 [0.054] N=1,011	-0.020 [0.057] N=869	0.016 [0.061] N=721
Underweight: WAZ below -2	0.109 P=150	**0.094*** [0.026] N=1,407**	**0.081*** [0.027] N=1,304**	**0.073** [0.029] N=1,165**	**0.067** [0.029] N=997**	**0.086*** [0.032] N=823**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=147	-0.040 [0.039] N=1,216	-0.031 [0.041] N=1,127	-0.034 [0.044] N=1,009	-0.024 [0.047] N=865	-0.040 [0.049] N=714
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=236	0.052 [0.049] N=1,216	0.060 [0.052] N=1,127	0.043 [0.055] N=1,009	0.059 [0.060] N=865	0.045 [0.068] N=714
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=107	0.007 [0.035] N=1,216	0.012 [0.038] N=1,127	0.018 [0.040] N=1,009	0.036 [0.043] N=865	0.036 [0.048] N=714
CIAF: any stunting, underweight, or wasting, 0-60 months	0.398 P=456	-0.038 [0.049] N=1,165	-0.025 [0.051] N=1,080	-0.014 [0.055] N=966	-0.028 [0.061] N=830	0.014 [0.068] N=687
BMI-for-age z score below -2	0.138 P=160	-0.032 [0.044] N=1,196	-0.022 [0.046] N=1,110	-0.017 [0.051] N=991	0.027 [0.055] N=849	0.032 [0.060] N=704

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.103 [0.383] N=1,028	-0.014 [0.430] N=941	0.209 [0.509] N=819	0.533 [0.572] N=677	0.493 [0.717] N=529
WHO weight-for-age z score	**-0.366* [0.189] N=1,190**	-0.358 [0.225] N=1,087	-0.357 [0.244] N=948	-0.229 [0.298] N=780	-0.557 [0.378] N=606
WHO weight-for-length/height z score, 0-60 months	0.218 [0.337] N=1,024	0.232 [0.367] N=935	0.065 [0.401] N=817	0.153 [0.454] N=673	-0.041 [0.547] N=522
WHO BMI-for-age z score	0.312 [0.346] N=1,004	0.348 [0.368] N=918	0.197 [0.414] N=799	-0.063 [0.470] N=657	-0.360 [0.552] N=512
Stunted: HAZ below -2	-0.041 [0.064] N=1,028	-0.040 [0.071] N=941	-0.045 [0.082] N=819	-0.106 [0.096] N=677	-0.091 [0.114] N=529
Underweight: WAZ below -2	**0.115*** [0.040] N=1,190**	**0.093** [0.045] N=1,087**	0.077 [0.050] N=948	0.059 [0.061] N=780	0.107 [0.083] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.054 [0.057] N=1,024	-0.041 [0.063] N=935	-0.050 [0.071] N=817	-0.035 [0.087] N=673	-0.059 [0.106] N=522
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.062] N=1,024	0.023 [0.069] N=935	-0.029 [0.070] N=817	0.001 [0.090] N=673	-0.077 [0.104] N=522
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.043] N=1,024	-0.010 [0.049] N=935	-0.010 [0.051] N=817	0.034 [0.059] N=673	0.032 [0.077] N=522
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.070 [0.074] N=981	-0.059 [0.081] N=896	-0.063 [0.090] N=782	-0.134 [0.116] N=646	-0.099 [0.159] N=503
BMI-for-age z score below -2	-0.073 [0.059] N=1,004	-0.062 [0.062] N=918	-0.064 [0.073] N=799	0.011 [0.088] N=657	0.032 [0.107] N=512

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.742	0.043 [0.276] N=1,123	-0.014 [0.297] N=1,046	0.079 [0.326] N=931	0.098 [0.339] N=807	-0.049 [0.357] N=672
WHO weight-for-age z score	-0.253	-0.033 [0.136] N=1,287	-0.030 [0.148] N=1,196	0.004 [0.152] N=1,066	0.069 [0.149] N=918	-0.004 [0.158] N=763
WHO weight-for-length/height z score, 0-60 months	0.323	**0.484** [0.230] N=1,119**	**0.505** [0.239] N=1,041**	**0.450* [0.253] N=931**	**0.525** [0.262] N=805**	**0.622** [0.286] N=667**
WHO BMI-for-age z score	0.326	**0.551** [0.269] N=1,101**	**0.560** [0.279] N=1,024**	**0.526* [0.300] N=914**	0.477 [0.321] N=790	0.563 [0.358] N=658
Stunted: HAZ below -2	0.303 P=331	-0.020 [0.051] N=1,123	-0.018 [0.054] N=1,046	-0.011 [0.058] N=931	-0.016 [0.063] N=807	0.021 [0.066] N=672
Underweight: WAZ below -2	0.111 P=139	**0.076*** [0.028] N=1,287**	**0.064** [0.029] N=1,196**	**0.053* [0.031] N=1,066**	0.047 [0.031] N=918	**0.059* [0.034] N=763**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=134	**-0.070* [0.038] N=1,119**	-0.059 [0.040] N=1,041	-0.060 [0.042] N=931	-0.051 [0.045] N=805	-0.070 [0.047] N=667
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=219	0.068 [0.048] N=1,119	0.074 [0.051] N=1,041	0.057 [0.054] N=931	0.076 [0.060] N=805	0.081 [0.067] N=667
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=102	0.009 [0.037] N=1,119	0.019 [0.040] N=1,041	0.028 [0.042] N=931	0.054 [0.045] N=805	0.066 [0.050] N=667
CIAF: any stunting, underweight, or wasting, 0-60 months	0.399 P=419	-0.069 [0.053] N=1,072	-0.057 [0.055] N=997	-0.050 [0.059] N=890	-0.064 [0.067] N=772	-0.030 [0.074] N=641
BMI-for-age z score below -2	0.137 P=144	-0.060 [0.044] N=1,101	-0.046 [0.045] N=1,024	-0.045 [0.050] N=914	-0.005 [0.055] N=790	-0.013 [0.059] N=658

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.178 [0.407] N=944	0.092 [0.469] N=867	0.389 [0.554] N=752	0.601 [0.620] N=628	0.644 [0.786] N=493
WHO weight-for-age z score	-0.272 [0.192] N=1,086	-0.302 [0.225] N=995	-0.288 [0.241] N=865	-0.249 [0.275] N=717	-0.555 [0.360] N=562
WHO weight-for-length/height z score, 0-60 months	0.254 [0.347] N=940	0.223 [0.375] N=862	0.003 [0.408] N=752	0.114 [0.465] N=626	-0.032 [0.571] N=488
WHO BMI-for-age z score	0.317 [0.367] N=922	0.250 [0.387] N=845	0.058 [0.420] N=735	-0.171 [0.473] N=611	-0.485 [0.579] N=479
Stunted: HAZ below -2	-0.069 [0.067] N=944	-0.077 [0.075] N=867	-0.090 [0.087] N=752	-0.135 [0.102] N=628	-0.143 [0.124] N=493
Underweight: WAZ below -2	**0.106** [0.042] N=1,086**	**0.089* [0.047] N=995**	0.071 [0.052] N=865	0.075 [0.061] N=717	0.120 [0.083] N=562
Wasted: WLZ/WHZ below -2, 0-60 months	-0.088 [0.058] N=940	-0.067 [0.063] N=862	-0.069 [0.071] N=752	-0.070 [0.086] N=626	-0.125 [0.105] N=488
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.062] N=940	0.017 [0.068] N=862	-0.037 [0.069] N=752	-0.015 [0.091] N=626	-0.098 [0.111] N=488
Obese: WLZ/WHZ above +3, 0-60 months	-0.035 [0.046] N=940	-0.026 [0.053] N=862	-0.021 [0.056] N=752	0.032 [0.065] N=626	0.034 [0.086] N=488
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.114 [0.075] N=901	-0.103 [0.083] N=826	-0.115 [0.093] N=719	-0.191 [0.116] N=601	-0.203 [0.152] N=470
BMI-for-age z score below -2	-0.091 [0.061] N=922	-0.068 [0.063] N=845	-0.066 [0.073] N=735	0.003 [0.089] N=611	-0.010 [0.112] N=479

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.742	0.059 [0.276] N=1,123	0.004 [0.296] N=1,046	0.085 [0.325] N=931	0.097 [0.338] N=807	-0.061 [0.356] N=672
WHO weight-for-age z score	-0.253	-0.023 [0.135] N=1,287	-0.021 [0.148] N=1,196	0.008 [0.151] N=1,066	0.059 [0.151] N=918	-0.006 [0.164] N=763
WHO weight-for-length/height z score, 0-60 months	0.323	**0.471** [0.226] N=1,119**	**0.490** [0.235] N=1,041**	**0.433* [0.248] N=931**	**0.497* [0.257] N=805**	**0.588** [0.282] N=667**
WHO BMI-for-age z score	0.326	**0.539** [0.265] N=1,101**	**0.545** [0.275] N=1,024**	**0.514* [0.296] N=914**	0.456 [0.318] N=790	0.530 [0.354] N=658
Stunted: HAZ below -2	0.303 P=331	-0.019 [0.050] N=1,123	-0.017 [0.053] N=1,046	-0.010 [0.057] N=931	-0.017 [0.061] N=807	0.017 [0.065] N=672
Underweight: WAZ below -2	0.111 P=139	**0.075*** [0.028] N=1,287**	**0.063** [0.030] N=1,196**	**0.053* [0.031] N=1,066**	0.050 [0.032] N=918	0.057 [0.035] N=763
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=134	**-0.067* [0.038] N=1,119**	-0.057 [0.040] N=1,041	-0.059 [0.042] N=931	-0.049 [0.045] N=805	-0.070 [0.047] N=667
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=219	0.067 [0.048] N=1,119	0.072 [0.051] N=1,041	0.053 [0.054] N=931	0.070 [0.060] N=805	0.073 [0.066] N=667
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=102	0.009 [0.037] N=1,119	0.019 [0.040] N=1,041	0.028 [0.042] N=931	0.052 [0.044] N=805	0.063 [0.049] N=667
CIAF: any stunting, underweight, or wasting, 0-60 months	0.399 P=419	-0.066 [0.052] N=1,072	-0.055 [0.055] N=997	-0.048 [0.059] N=890	-0.066 [0.066] N=772	-0.034 [0.072] N=641
BMI-for-age z score below -2	0.137 P=144	-0.056 [0.044] N=1,101	-0.044 [0.045] N=1,024	-0.043 [0.050] N=914	-0.004 [0.055] N=790	-0.011 [0.059] N=658

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.210 [0.408] N=944	0.126 [0.470] N=867	0.408 [0.556] N=752	0.604 [0.624] N=628	0.663 [0.787] N=493
WHO weight-for-age z score	-0.239 [0.190] N=1,086	-0.270 [0.223] N=995	-0.256 [0.238] N=865	-0.242 [0.278] N=717	-0.510 [0.377] N=562
WHO weight-for-length/height z score, 0-60 months	0.259 [0.343] N=940	0.236 [0.374] N=862	0.016 [0.407] N=752	0.097 [0.461] N=626	-0.062 [0.569] N=488
WHO BMI-for-age z score	0.320 [0.361] N=922	0.250 [0.381] N=845	0.069 [0.417] N=735	-0.186 [0.467] N=611	-0.527 [0.568] N=479
Stunted: HAZ below -2	-0.062 [0.067] N=944	-0.069 [0.075] N=867	-0.083 [0.086] N=752	-0.127 [0.102] N=628	-0.147 [0.123] N=493
Underweight: WAZ below -2	**0.101** [0.043] N=1,086**	**0.084* [0.047] N=995**	0.066 [0.053] N=865	0.074 [0.063] N=717	0.104 [0.087] N=562
Wasted: WLZ/WHZ below -2, 0-60 months	-0.085 [0.058] N=940	-0.065 [0.063] N=862	-0.069 [0.070] N=752	-0.068 [0.085] N=626	-0.124 [0.103] N=488
Overweight: WLZ/WHZ above +2, 0-60 months	0.021 [0.061] N=940	0.022 [0.068] N=862	-0.033 [0.069] N=752	-0.016 [0.090] N=626	-0.098 [0.110] N=488
Obese: WLZ/WHZ above +3, 0-60 months	-0.035 [0.045] N=940	-0.026 [0.053] N=862	-0.020 [0.055] N=752	0.033 [0.064] N=626	0.035 [0.084] N=488
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.102 [0.075] N=901	-0.094 [0.082] N=826	-0.109 [0.092] N=719	-0.186 [0.116] N=601	-0.212 [0.153] N=470
BMI-for-age z score below -2	-0.084 [0.060] N=922	-0.063 [0.062] N=845	-0.064 [0.073] N=735	0.005 [0.089] N=611	-0.007 [0.111] N=479

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.087 [0.327] N=1,045	-0.196 [0.338] N=935	-0.028 [0.364] N=816	-0.109 [0.389] N=692	-0.211 [0.444] N=563
WHO weight-for-age z score	-0.267	-0.050 [0.155] N=1,203	-0.038 [0.166] N=1,085	-0.089 [0.174] N=947	-0.033 [0.182] N=800	-0.002 [0.198] N=648
WHO weight-for-length/height z score, 0-60 months	0.321	-0.043 [0.301] N=1,046	-0.098 [0.309] N=941	-0.199 [0.335] N=821	-0.020 [0.367] N=696	-0.141 [0.412] N=565
WHO BMI-for-age z score	0.289	0.034 [0.285] N=1,028	0.067 [0.291] N=924	-0.004 [0.324] N=804	0.235 [0.357] N=682	0.154 [0.389] N=558
Stunted: HAZ below -2	0.284 P=294	0.059 [0.069] N=1,045	0.064 [0.071] N=935	0.039 [0.075] N=816	0.043 [0.083] N=692	0.019 [0.091] N=563
Underweight: WAZ below -2	0.113 P=132	0.016 [0.037] N=1,203	0.019 [0.040] N=1,085	0.042 [0.041] N=947	0.037 [0.044] N=800	0.020 [0.046] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=126	0.017 [0.047] N=1,046	0.035 [0.049] N=941	0.066 [0.054] N=821	0.015 [0.053] N=696	0.022 [0.062] N=565
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=204	0.025 [0.055] N=1,046	0.026 [0.057] N=941	-0.000 [0.060] N=821	0.011 [0.065] N=696	0.003 [0.073] N=565
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=96	0.017 [0.042] N=1,046	0.006 [0.043] N=941	0.008 [0.046] N=821	-0.007 [0.051] N=696	-0.057 [0.051] N=565
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=385	0.100 [0.061] N=1,000	**0.109* [0.063] N=896**	0.101 [0.067] N=781	0.051 [0.072] N=662	0.028 [0.077] N=537
BMI-for-age z score below -2	0.139 P=141	-0.005 [0.048] N=1,028	0.004 [0.050] N=924	0.003 [0.057] N=804	-0.045 [0.058] N=682	-0.031 [0.066] N=558

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.245 [0.489] N=909	-0.431 [0.516] N=799	-0.139 [0.603] N=680	-0.413 [0.691] N=556	-0.709 [0.860] N=427
WHO weight-for-age z score	-0.069 [0.237] N=1,047	-0.090 [0.257] N=929	-0.213 [0.285] N=791	-0.184 [0.315] N=644	-0.051 [0.359] N=492
WHO weight-for-length/height z score, 0-60 months	-0.130 [0.376] N=909	-0.252 [0.397] N=804	-0.420 [0.460] N=684	-0.106 [0.525] N=559	-0.478 [0.641] N=428
WHO BMI-for-age z score	-0.234 [0.374] N=898	-0.213 [0.389] N=794	-0.332 [0.475] N=674	0.107 [0.551] N=552	-0.108 [0.596] N=428
Stunted: HAZ below -2	0.102 [0.099] N=909	0.115 [0.103] N=799	0.073 [0.117] N=680	0.107 [0.141] N=556	0.068 [0.166] N=427
Underweight: WAZ below -2	-0.008 [0.059] N=1,047	0.003 [0.065] N=929	0.045 [0.073] N=791	0.045 [0.083] N=644	0.002 [0.094] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.053 [0.063] N=909	0.085 [0.067] N=804	**0.142* [0.078] N=684**	0.048 [0.079] N=559	0.090 [0.104] N=428
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.074] N=909	0.012 [0.079] N=804	-0.031 [0.087] N=684	-0.002 [0.102] N=559	-0.047 [0.127] N=428
Obese: WLZ/WHZ above +3, 0-60 months	0.055 [0.055] N=909	0.030 [0.057] N=804	0.040 [0.068] N=684	0.008 [0.084] N=559	-0.132 [0.093] N=428
CIAF: any stunting, underweight, or wasting, 0-60 months	0.158 [0.098] N=868	**0.184* [0.104] N=764**	0.169 [0.120] N=649	0.090 [0.142] N=530	0.040 [0.169] N=405
BMI-for-age z score below -2	0.015 [0.062] N=898	0.032 [0.066] N=794	0.017 [0.083] N=674	-0.098 [0.079] N=552	-0.088 [0.100] N=428

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.082 [0.327] N=1,045	-0.193 [0.338] N=935	-0.027 [0.363] N=816	-0.109 [0.389] N=692	-0.212 [0.444] N=563
WHO weight-for-age z score	-0.267	-0.041 [0.153] N=1,203	-0.030 [0.163] N=1,085	-0.080 [0.171] N=947	-0.020 [0.178] N=800	0.013 [0.195] N=648
WHO weight-for-length/height z score, 0-60 months	0.321	-0.040 [0.300] N=1,046	-0.096 [0.309] N=941	-0.199 [0.334] N=821	-0.015 [0.365] N=696	-0.121 [0.407] N=565
WHO BMI-for-age z score	0.289	0.031 [0.286] N=1,028	0.064 [0.291] N=924	-0.010 [0.324] N=804	0.231 [0.358] N=682	0.156 [0.391] N=558
Stunted: HAZ below -2	0.284 P=294	0.057 [0.069] N=1,045	0.062 [0.071] N=935	0.037 [0.075] N=816	0.042 [0.083] N=692	0.018 [0.092] N=563
Underweight: WAZ below -2	0.113 P=132	0.014 [0.037] N=1,203	0.017 [0.039] N=1,085	0.039 [0.041] N=947	0.035 [0.044] N=800	0.017 [0.046] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=126	0.015 [0.047] N=1,046	0.032 [0.049] N=941	0.063 [0.054] N=821	0.011 [0.053] N=696	0.016 [0.061] N=565
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=204	0.024 [0.055] N=1,046	0.024 [0.057] N=941	-0.004 [0.060] N=821	0.008 [0.067] N=696	0.001 [0.074] N=565
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=96	0.016 [0.042] N=1,046	0.004 [0.043] N=941	0.005 [0.046] N=821	-0.009 [0.051] N=696	-0.060 [0.052] N=565
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=385	0.096 [0.061] N=1,000	0.104 [0.063] N=896	0.095 [0.067] N=781	0.048 [0.072] N=662	0.022 [0.078] N=537
BMI-for-age z score below -2	0.139 P=141	-0.006 [0.048] N=1,028	0.002 [0.050] N=924	0.003 [0.057] N=804	-0.045 [0.059] N=682	-0.032 [0.066] N=558

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.230 [0.489] N=909	-0.425 [0.517] N=799	-0.140 [0.600] N=680	-0.423 [0.690] N=556	-0.712 [0.859] N=427
WHO weight-for-age z score	-0.056 [0.232] N=1,047	-0.076 [0.252] N=929	-0.196 [0.279] N=791	-0.145 [0.308] N=644	0.001 [0.359] N=492
WHO weight-for-length/height z score, 0-60 months	-0.129 [0.378] N=909	-0.254 [0.398] N=804	-0.428 [0.462] N=684	-0.091 [0.532] N=559	-0.410 [0.661] N=428
WHO BMI-for-age z score	-0.237 [0.376] N=898	-0.211 [0.392] N=794	-0.337 [0.477] N=674	0.119 [0.560] N=552	-0.050 [0.623] N=428
Stunted: HAZ below -2	0.098 [0.098] N=909	0.112 [0.103] N=799	0.071 [0.116] N=680	0.111 [0.142] N=556	0.071 [0.167] N=427
Underweight: WAZ below -2	-0.010 [0.059] N=1,047	0.000 [0.065] N=929	0.041 [0.072] N=791	0.040 [0.083] N=644	-0.005 [0.094] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.049 [0.064] N=909	0.079 [0.069] N=804	**0.134* [0.080] N=684**	0.034 [0.080] N=559	0.066 [0.108] N=428
Overweight: WLZ/WHZ above +2, 0-60 months	0.021 [0.073] N=909	0.008 [0.079] N=804	-0.039 [0.086] N=684	-0.013 [0.103] N=559	-0.053 [0.127] N=428
Obese: WLZ/WHZ above +3, 0-60 months	0.053 [0.054] N=909	0.027 [0.057] N=804	0.035 [0.068] N=684	0.001 [0.085] N=559	-0.141 [0.094] N=428
CIAF: any stunting, underweight, or wasting, 0-60 months	0.146 [0.096] N=868	0.168 [0.102] N=764	0.149 [0.117] N=649	0.068 [0.140] N=530	0.003 [0.164] N=405
BMI-for-age z score below -2	0.012 [0.062] N=898	0.027 [0.067] N=794	0.014 [0.084] N=674	-0.104 [0.081] N=552	-0.099 [0.105] N=428

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.083 [0.323] N=1,045	-0.176 [0.334] N=935	0.005 [0.363] N=816	-0.083 [0.389] N=692	-0.190 [0.443] N=563
WHO weight-for-age z score	-0.267	-0.040 [0.146] N=1,203	-0.024 [0.156] N=1,085	-0.054 [0.163] N=947	-0.003 [0.170] N=800	0.026 [0.183] N=648
WHO weight-for-length/height z score, 0-60 months	0.321	-0.038 [0.300] N=1,046	-0.100 [0.308] N=941	-0.175 [0.332] N=821	0.005 [0.362] N=696	-0.106 [0.409] N=565
WHO BMI-for-age z score	0.289	0.027 [0.286] N=1,028	0.053 [0.292] N=924	-0.001 [0.319] N=804	0.243 [0.349] N=682	0.153 [0.386] N=558
Stunted: HAZ below -2	0.284 P=294	0.056 [0.068] N=1,045	0.057 [0.070] N=935	0.027 [0.074] N=816	0.031 [0.082] N=692	0.009 [0.089] N=563
Underweight: WAZ below -2	0.113 P=132	0.015 [0.035] N=1,203	0.018 [0.037] N=1,085	0.035 [0.038] N=947	0.032 [0.040] N=800	0.016 [0.039] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=126	0.015 [0.048] N=1,046	0.031 [0.049] N=941	0.060 [0.055] N=821	0.009 [0.054] N=696	0.016 [0.063] N=565
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=204	0.024 [0.055] N=1,046	0.024 [0.058] N=941	-0.006 [0.061] N=821	0.004 [0.067] N=696	0.002 [0.075] N=565
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=96	0.016 [0.042] N=1,046	0.003 [0.043] N=941	0.004 [0.047] N=821	-0.010 [0.052] N=696	-0.059 [0.053] N=565
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=385	0.097 [0.061] N=1,000	0.101 [0.063] N=896	0.084 [0.068] N=781	0.034 [0.072] N=662	0.012 [0.076] N=537
BMI-for-age z score below -2	0.139 P=141	-0.005 [0.048] N=1,028	0.005 [0.050] N=924	-0.000 [0.057] N=804	-0.048 [0.059] N=682	-0.032 [0.067] N=558

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.270 [0.486] N=909	-0.427 [0.509] N=799	-0.142 [0.597] N=680	-0.442 [0.684] N=556	-0.776 [0.877] N=427
WHO weight-for-age z score	-0.098 [0.228] N=1,047	-0.113 [0.247] N=929	-0.186 [0.276] N=791	-0.169 [0.308] N=644	-0.082 [0.359] N=492
WHO weight-for-length/height z score, 0-60 months	-0.154 [0.378] N=909	-0.288 [0.399] N=804	-0.414 [0.454] N=684	-0.085 [0.527] N=559	-0.413 [0.678] N=428
WHO BMI-for-age z score	-0.256 [0.378] N=898	-0.244 [0.396] N=794	-0.328 [0.472] N=674	0.125 [0.554] N=552	-0.053 [0.630] N=428
Stunted: HAZ below -2	0.104 [0.098] N=909	0.111 [0.102] N=799	0.070 [0.114] N=680	0.117 [0.140] N=556	0.090 [0.166] N=427
Underweight: WAZ below -2	-0.003 [0.057] N=1,047	0.009 [0.062] N=929	0.039 [0.070] N=791	0.045 [0.081] N=644	0.009 [0.087] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.054 [0.065] N=909	0.087 [0.069] N=804	0.133 [0.081] N=684	0.034 [0.081] N=559	0.072 [0.110] N=428
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.074] N=909	0.011 [0.080] N=804	-0.040 [0.087] N=684	-0.012 [0.104] N=559	-0.053 [0.129] N=428
Obese: WLZ/WHZ above +3, 0-60 months	0.053 [0.055] N=909	0.028 [0.058] N=804	0.035 [0.068] N=684	0.002 [0.085] N=559	-0.140 [0.093] N=428
CIAF: any stunting, underweight, or wasting, 0-60 months	0.160 [0.098] N=868	**0.178* [0.104] N=764**	0.147 [0.120] N=649	0.075 [0.144] N=530	0.027 [0.170] N=405
BMI-for-age z score below -2	0.018 [0.062] N=898	0.037 [0.066] N=794	0.013 [0.083] N=674	-0.104 [0.080] N=552	-0.093 [0.105] N=428

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.091 [0.317] N=1,045	-0.173 [0.328] N=935	0.012 [0.351] N=816	-0.084 [0.376] N=692	-0.248 [0.429] N=563
WHO weight-for-age z score	-0.267	-0.047 [0.144] N=1,203	-0.033 [0.153] N=1,085	-0.063 [0.161] N=947	-0.012 [0.165] N=800	0.004 [0.179] N=648
WHO weight-for-length/height z score, 0-60 months	0.321	-0.021 [0.297] N=1,046	-0.087 [0.305] N=941	-0.177 [0.327] N=821	0.026 [0.356] N=696	-0.052 [0.401] N=565
WHO BMI-for-age z score	0.289	0.042 [0.285] N=1,028	0.072 [0.293] N=924	0.012 [0.320] N=804	0.278 [0.350] N=682	0.236 [0.388] N=558
Stunted: HAZ below -2	0.284 P=294	0.057 [0.068] N=1,045	0.056 [0.069] N=935	0.029 [0.074] N=816	0.038 [0.081] N=692	0.022 [0.089] N=563
Underweight: WAZ below -2	0.113 P=132	0.015 [0.035] N=1,203	0.019 [0.037] N=1,085	0.038 [0.039] N=947	0.036 [0.041] N=800	0.022 [0.042] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=126	0.012 [0.048] N=1,046	0.031 [0.050] N=941	0.063 [0.054] N=821	0.009 [0.052] N=696	0.011 [0.060] N=565
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=204	0.024 [0.054] N=1,046	0.022 [0.057] N=941	-0.003 [0.060] N=821	0.011 [0.065] N=696	0.010 [0.074] N=565
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=96	0.021 [0.041] N=1,046	0.008 [0.042] N=941	0.008 [0.045] N=821	-0.004 [0.050] N=696	-0.052 [0.051] N=565
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=385	0.094 [0.061] N=1,000	0.099 [0.063] N=896	0.087 [0.067] N=781	0.044 [0.072] N=662	0.027 [0.077] N=537
BMI-for-age z score below -2	0.139 P=141	-0.010 [0.048] N=1,028	-0.001 [0.051] N=924	-0.003 [0.056] N=804	-0.051 [0.057] N=682	-0.043 [0.065] N=558

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263 [0.471] N=909	-0.416 [0.497] N=799	-0.046 [0.572] N=680	-0.325 [0.657] N=556	-0.725 [0.828] N=427
WHO weight-for-age z score	-0.075 [0.226] N=1,047	-0.092 [0.244] N=929	-0.161 [0.279] N=791	-0.134 [0.302] N=644	-0.046 [0.356] N=492
WHO weight-for-length/height z score, 0-60 months	-0.123 [0.377] N=909	-0.264 [0.396] N=804	-0.436 [0.449] N=684	-0.095 [0.517] N=559	-0.398 [0.658] N=428
WHO BMI-for-age z score	-0.230 [0.376] N=898	-0.213 [0.395] N=794	-0.336 [0.475] N=674	0.136 [0.552] N=552	-0.021 [0.620] N=428
Stunted: HAZ below -2	0.106 [0.095] N=909	0.111 [0.099] N=799	0.059 [0.112] N=680	0.104 [0.139] N=556	0.090 [0.165] N=427
Underweight: WAZ below -2	-0.008 [0.056] N=1,047	0.004 [0.062] N=929	0.033 [0.071] N=791	0.038 [0.081] N=644	0.000 [0.090] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.050 [0.065] N=909	0.082 [0.069] N=804	**0.136* [0.079] N=684**	0.039 [0.079] N=559	0.072 [0.106] N=428
Overweight: WLZ/WHZ above +2, 0-60 months	0.021 [0.073] N=909	0.006 [0.079] N=804	-0.047 [0.085] N=684	-0.018 [0.101] N=559	-0.049 [0.127] N=428
Obese: WLZ/WHZ above +3, 0-60 months	0.056 [0.054] N=909	0.028 [0.057] N=804	0.033 [0.066] N=684	0.004 [0.083] N=559	-0.134 [0.092] N=428
CIAF: any stunting, underweight, or wasting, 0-60 months	0.152 [0.097] N=868	0.168 [0.104] N=764	0.136 [0.120] N=649	0.065 [0.145] N=530	0.031 [0.170] N=405
BMI-for-age z score below -2	0.014 [0.063] N=898	0.029 [0.067] N=794	0.013 [0.084] N=674	-0.102 [0.081] N=552	-0.101 [0.104] N=428

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.088 [0.349] N=958	-0.207 [0.367] N=857	-0.007 [0.389] N=754	-0.112 [0.404] N=637	-0.224 [0.457] N=521
WHO weight-for-age z score	-0.269	-0.016 [0.163] N=1,097	0.016 [0.175] N=988	-0.072 [0.182] N=869	-0.037 [0.188] N=735	-0.050 [0.200] N=600
WHO weight-for-length/height z score, 0-60 months	0.329	0.068 [0.317] N=959	0.019 [0.331] N=863	-0.137 [0.353] N=757	0.007 [0.380] N=641	-0.153 [0.416] N=524
WHO BMI-for-age z score	0.302	0.112 [0.303] N=943	0.168 [0.315] N=848	0.025 [0.347] N=743	0.243 [0.384] N=630	0.116 [0.418] N=519
Stunted: HAZ below -2	0.284 P=268	0.056 [0.075] N=958	0.064 [0.078] N=857	0.025 [0.081] N=754	0.024 [0.087] N=637	0.001 [0.096] N=521
Underweight: WAZ below -2	0.116 P=124	0.011 [0.040] N=1,097	0.013 [0.043] N=988	0.036 [0.044] N=869	0.039 [0.047] N=735	0.027 [0.050] N=600
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=115	0.012 [0.050] N=959	0.028 [0.052] N=863	0.062 [0.057] N=757	0.011 [0.055] N=641	0.028 [0.063] N=524
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=189	0.030 [0.056] N=959	0.031 [0.059] N=863	-0.008 [0.061] N=757	-0.012 [0.066] N=641	-0.029 [0.073] N=524
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=90	0.015 [0.044] N=959	0.000 [0.044] N=863	-0.000 [0.047] N=757	-0.021 [0.050] N=641	-0.069 [0.050] N=524
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=352	0.096 [0.067] N=917	0.101 [0.070] N=822	0.079 [0.073] N=721	0.018 [0.076] N=609	0.009 [0.082] N=497
BMI-for-age z score below -2	0.138 P=126	-0.002 [0.050] N=943	0.006 [0.053] N=848	0.005 [0.059] N=743	-0.036 [0.061] N=630	-0.009 [0.070] N=519

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.234 [0.534] N=831	-0.445 [0.568] N=730	-0.155 [0.657] N=627	-0.550 [0.726] N=510	-0.817 [0.902] N=394
WHO weight-for-age z score	0.010 [0.238] N=953	0.032 [0.259] N=844	-0.156 [0.290] N=725	-0.105 [0.320] N=591	0.045 [0.365] N=456
WHO weight-for-length/height z score, 0-60 months	-0.017 [0.409] N=832	-0.110 [0.439] N=736	-0.365 [0.508] N=630	-0.059 [0.589] N=514	-0.478 [0.730] N=397
WHO BMI-for-age z score	-0.138 [0.381] N=821	-0.058 [0.402] N=726	-0.264 [0.490] N=621	0.210 [0.572] N=508	-0.031 [0.667] N=397
Stunted: HAZ below -2	0.109 [0.110] N=831	0.128 [0.118] N=730	0.070 [0.130] N=627	0.092 [0.154] N=510	0.037 [0.181] N=394
Underweight: WAZ below -2	-0.024 [0.063] N=953	-0.018 [0.070] N=844	0.021 [0.076] N=725	0.029 [0.090] N=591	-0.031 [0.100] N=456
Wasted: WLZ/WHZ below -2, 0-60 months	0.041 [0.067] N=832	0.067 [0.074] N=736	0.119 [0.087] N=630	0.012 [0.085] N=514	0.062 [0.113] N=397
Overweight: WLZ/WHZ above +2, 0-60 months	0.029 [0.080] N=832	0.022 [0.087] N=736	-0.039 [0.096] N=630	-0.046 [0.114] N=514	-0.118 [0.141] N=397
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.061] N=832	0.013 [0.064] N=736	0.021 [0.076] N=630	-0.028 [0.094] N=514	-0.175 [0.108] N=397
CIAF: any stunting, underweight, or wasting, 0-60 months	0.157 [0.110] N=794	0.172 [0.119] N=699	0.127 [0.134] N=598	-0.001 [0.156] N=486	-0.063 [0.183] N=374
BMI-for-age z score below -2	0.021 [0.063] N=821	0.033 [0.070] N=726	0.014 [0.087] N=621	-0.097 [0.085] N=508	-0.063 [0.114] N=397

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.616	-0.116 [0.352] N=958	-0.232 [0.370] N=857	-0.026 [0.391] N=754	-0.132 [0.406] N=637	-0.242 [0.464] N=521
WHO weight-for-age z score	-0.269	-0.017 [0.160] N=1,097	0.006 [0.173] N=988	-0.074 [0.180] N=869	-0.035 [0.184] N=735	-0.044 [0.195] N=600
WHO weight-for-length/height z score, 0-60 months	0.329	0.082 [0.317] N=959	0.030 [0.331] N=863	-0.125 [0.352] N=757	0.013 [0.380] N=641	-0.145 [0.416] N=524
WHO BMI-for-age z score	0.302	0.118 [0.303] N=943	0.175 [0.314] N=848	0.031 [0.346] N=743	0.251 [0.382] N=630	0.128 [0.414] N=519
Stunted: HAZ below -2	0.284 P=268	0.057 [0.074] N=958	0.063 [0.077] N=857	0.023 [0.081] N=754	0.020 [0.086] N=637	-0.005 [0.095] N=521
Underweight: WAZ below -2	0.116 P=124	0.008 [0.039] N=1,097	0.011 [0.042] N=988	0.033 [0.043] N=869	0.035 [0.046] N=735	0.023 [0.048] N=600
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=115	0.008 [0.050] N=959	0.024 [0.052] N=863	0.055 [0.057] N=757	0.004 [0.055] N=641	0.020 [0.063] N=524
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=189	0.032 [0.056] N=959	0.031 [0.059] N=863	-0.009 [0.060] N=757	-0.014 [0.066] N=641	-0.030 [0.072] N=524
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=90	0.015 [0.043] N=959	0.000 [0.044] N=863	0.000 [0.046] N=757	-0.022 [0.049] N=641	-0.069 [0.048] N=524
CIAF: any stunting, underweight, or wasting, 0-60 months	0.387 P=352	0.095 [0.066] N=917	0.097 [0.069] N=822	0.072 [0.072] N=721	0.008 [0.076] N=609	-0.005 [0.083] N=497
BMI-for-age z score below -2	0.138 P=126	-0.004 [0.051] N=943	0.003 [0.053] N=848	0.002 [0.059] N=743	-0.040 [0.060] N=630	-0.015 [0.068] N=519

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.258 [0.532] N=831	-0.456 [0.565] N=730	-0.140 [0.649] N=627	-0.538 [0.713] N=510	-0.839 [0.884] N=394
WHO weight-for-age z score	0.001 [0.236] N=953	0.015 [0.260] N=844	-0.159 [0.289] N=725	-0.111 [0.318] N=591	0.026 [0.362] N=456
WHO weight-for-length/height z score, 0-60 months	-0.031 [0.406] N=832	-0.130 [0.437] N=736	-0.389 [0.503] N=630	-0.101 [0.572] N=514	-0.469 [0.704] N=397
WHO BMI-for-age z score	-0.147 [0.380] N=821	-0.067 [0.401] N=726	-0.281 [0.487] N=621	0.203 [0.567] N=508	0.002 [0.649] N=397
Stunted: HAZ below -2	0.109 [0.110] N=831	0.126 [0.117] N=730	0.063 [0.128] N=627	0.087 [0.152] N=510	0.032 [0.178] N=394
Underweight: WAZ below -2	-0.027 [0.063] N=953	-0.019 [0.070] N=844	0.019 [0.077] N=725	0.028 [0.090] N=591	-0.017 [0.101] N=456
Wasted: WLZ/WHZ below -2, 0-60 months	0.046 [0.068] N=832	0.072 [0.074] N=736	0.123 [0.088] N=630	0.016 [0.085] N=514	0.063 [0.113] N=397
Overweight: WLZ/WHZ above +2, 0-60 months	0.032 [0.079] N=832	0.022 [0.085] N=736	-0.044 [0.094] N=630	-0.051 [0.111] N=514	-0.106 [0.137] N=397
Obese: WLZ/WHZ above +3, 0-60 months	0.041 [0.059] N=832	0.013 [0.063] N=736	0.020 [0.074] N=630	-0.030 [0.091] N=514	**-0.171* [0.101] N=397**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.164 [0.109] N=794	0.176 [0.118] N=699	0.128 [0.133] N=598	-0.001 [0.157] N=486	-0.065 [0.187] N=374
BMI-for-age z score below -2	0.025 [0.064] N=821	0.035 [0.070] N=726	0.017 [0.089] N=621	-0.093 [0.086] N=508	-0.065 [0.115] N=397

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.