Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Oldest Child

Age reference: 12 November 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,153	-0.003 [0.027] N=1,245	0.001 [0.027] N=1,148	-0.011 [0.028] N=1,029	-0.018 [0.029] N=883	-0.031 [0.032] N=727
Vegetables consumed at least weekly	0.898 P=1,115	-0.002 [0.032] N=1,245	-0.017 [0.032] N=1,148	-0.033 [0.033] N=1,029	-0.030 [0.034] N=883	-0.059 [0.039] N=727
Meat fish or poultry consumed at least weekly	0.764 P=958	0.062 [0.055] N=1,245	0.069 [0.058] N=1,148	0.063 [0.063] N=1,029	0.056 [0.071] N=883	0.042 [0.077] N=727
Legumes consumed at least weekly	0.766 P=959	-0.045 [0.047] N=1,245	-0.074 [0.049] N=1,148	-0.065 [0.050] N=1,029	-0.065 [0.056] N=883	-0.068 [0.067] N=727
Animal products consumed at least weekly	0.907 P=1,132	0.016 [0.038] N=1,245	0.012 [0.040] N=1,148	-0.014 [0.043] N=1,029	-0.010 [0.047] N=883	-0.030 [0.054] N=727
Sweets consumed at least weekly	0.546 P=677	-0.041 [0.050] N=1,245	-0.037 [0.052] N=1,148	-0.029 [0.057] N=1,029	-0.045 [0.062] N=883	-0.016 [0.069] N=727
Snack foods consumed at least weekly	0.452 P=556	-0.010 [0.046] N=1,245	-0.018 [0.048] N=1,148	-0.001 [0.052] N=1,029	-0.006 [0.056] N=883	-0.012 [0.060] N=727
Packaged soft drinks consumed at least weekly	0.260 P=316	-0.013 [0.046] N=1,245	-0.039 [0.048] N=1,148	-0.045 [0.050] N=1,029	-0.030 [0.053] N=883	-0.054 [0.060] N=727
Grains consumed at least weekly	0.865 P=1,078	0.021 [0.036] N=1,245	0.017 [0.038] N=1,148	0.008 [0.040] N=1,029	0.016 [0.044] N=883	0.002 [0.050] N=727
Child was ever breastfed	0.968 P=1,416	-0.018 [0.013] N=1,467	-0.015 [0.014] N=1,356	-0.018 [0.015] N=1,212	-0.020 [0.017] N=1,034	-0.022 [0.016] N=855

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.005 [0.041] N=1,054	0.016 [0.042] N=957	-0.010 [0.047] N=838	-0.035 [0.054] N=692	-0.082 [0.072] N=536
Vegetables consumed at least weekly	0.014 [0.049] N=1,054	-0.008 [0.052] N=957	-0.042 [0.057] N=838	-0.045 [0.066] N=692	-0.154 [0.095] N=536
Meat fish or poultry consumed at least weekly	0.035 [0.068] N=1,054	0.039 [0.072] N=957	0.020 [0.082] N=838	-0.050 [0.091] N=692	-0.147 [0.093] N=536
Legumes consumed at least weekly	-0.039 [0.070] N=1,054	-0.094 [0.074] N=957	-0.072 [0.074] N=838	-0.103 [0.089] N=692	-0.123 [0.124] N=536
Animal products consumed at least weekly	0.034 [0.052] N=1,054	0.031 [0.057] N=957	-0.020 [0.064] N=838	-0.004 [0.077] N=692	-0.050 [0.102] N=536
Sweets consumed at least weekly	-0.074 [0.065] N=1,054	-0.070 [0.068] N=957	-0.061 [0.079] N=838	-0.141 [0.088] N=692	-0.081 [0.121] N=536
Snack foods consumed at least weekly	-0.036 [0.062] N=1,054	-0.058 [0.066] N=957	-0.036 [0.077] N=838	-0.076 [0.094] N=692	-0.136 [0.116] N=536
Packaged soft drinks consumed at least weekly	-0.020 [0.069] N=1,054	-0.070 [0.075] N=957	-0.087 [0.081] N=838	-0.080 [0.096] N=692	-0.164 [0.132] N=536
Grains consumed at least weekly	0.056 [0.054] N=1,054	0.051 [0.059] N=957	0.036 [0.064] N=838	0.060 [0.076] N=692	0.021 [0.105] N=536
Child was ever breastfed	-0.010 [0.019] N=1,239	-0.004 [0.022] N=1,128	-0.008 [0.028] N=984	-0.013 [0.035] N=806	-0.000 [0.040] N=627

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,153	-0.002 [0.026] N=1,245	0.001 [0.026] N=1,148	-0.010 [0.028] N=1,029	-0.018 [0.029] N=883	-0.031 [0.032] N=727
Vegetables consumed at least weekly	0.898 P=1,115	-0.002 [0.032] N=1,245	-0.017 [0.032] N=1,148	-0.032 [0.032] N=1,029	-0.030 [0.034] N=883	-0.060 [0.039] N=727
Meat fish or poultry consumed at least weekly	0.764 P=958	0.062 [0.055] N=1,245	0.069 [0.058] N=1,148	0.064 [0.063] N=1,029	0.056 [0.071] N=883	0.042 [0.078] N=727
Legumes consumed at least weekly	0.766 P=959	-0.045 [0.047] N=1,245	-0.074 [0.049] N=1,148	-0.063 [0.050] N=1,029	-0.065 [0.057] N=883	-0.069 [0.068] N=727
Animal products consumed at least weekly	0.907 P=1,132	0.017 [0.037] N=1,245	0.012 [0.039] N=1,148	-0.012 [0.042] N=1,029	-0.009 [0.047] N=883	-0.030 [0.053] N=727
Sweets consumed at least weekly	0.546 P=677	-0.041 [0.050] N=1,245	-0.037 [0.052] N=1,148	-0.028 [0.057] N=1,029	-0.045 [0.062] N=883	-0.016 [0.070] N=727
Snack foods consumed at least weekly	0.452 P=556	-0.009 [0.046] N=1,245	-0.017 [0.049] N=1,148	0.001 [0.053] N=1,029	-0.006 [0.057] N=883	-0.012 [0.061] N=727
Packaged soft drinks consumed at least weekly	0.260 P=316	-0.013 [0.046] N=1,245	-0.039 [0.048] N=1,148	-0.044 [0.050] N=1,029	-0.030 [0.054] N=883	-0.054 [0.060] N=727
Grains consumed at least weekly	0.865 P=1,078	0.021 [0.036] N=1,245	0.017 [0.038] N=1,148	0.009 [0.040] N=1,029	0.016 [0.044] N=883	0.002 [0.050] N=727
Child was ever breastfed	0.968 P=1,416	-0.018 [0.013] N=1,467	-0.015 [0.014] N=1,356	-0.018 [0.015] N=1,212	-0.020 [0.017] N=1,034	-0.021 [0.016] N=855

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.006 [0.041] N=1,054	0.016 [0.042] N=957	-0.009 [0.048] N=838	-0.034 [0.055] N=692	-0.082 [0.072] N=536
Vegetables consumed at least weekly	0.015 [0.049] N=1,054	-0.007 [0.052] N=957	-0.040 [0.057] N=838	-0.043 [0.067] N=692	-0.154 [0.095] N=536
Meat fish or poultry consumed at least weekly	0.035 [0.068] N=1,054	0.038 [0.072] N=957	0.020 [0.082] N=838	-0.049 [0.091] N=692	-0.147 [0.092] N=536
Legumes consumed at least weekly	-0.037 [0.070] N=1,054	-0.093 [0.074] N=957	-0.069 [0.074] N=838	-0.101 [0.089] N=692	-0.123 [0.123] N=536
Animal products consumed at least weekly	0.037 [0.051] N=1,054	0.033 [0.057] N=957	-0.015 [0.063] N=838	-0.001 [0.077] N=692	-0.050 [0.100] N=536
Sweets consumed at least weekly	-0.074 [0.065] N=1,054	-0.069 [0.068] N=957	-0.060 [0.079] N=838	-0.141 [0.088] N=692	-0.081 [0.121] N=536
Snack foods consumed at least weekly	-0.033 [0.062] N=1,054	-0.056 [0.066] N=957	-0.028 [0.077] N=838	-0.073 [0.095] N=692	-0.136 [0.115] N=536
Packaged soft drinks consumed at least weekly	-0.021 [0.069] N=1,054	-0.071 [0.075] N=957	-0.087 [0.081] N=838	-0.080 [0.096] N=692	-0.164 [0.133] N=536
Grains consumed at least weekly	0.057 [0.053] N=1,054	0.053 [0.058] N=957	0.041 [0.062] N=838	0.062 [0.075] N=692	0.021 [0.104] N=536
Child was ever breastfed	-0.010 [0.020] N=1,239	-0.003 [0.023] N=1,128	-0.008 [0.028] N=984	-0.013 [0.035] N=806	-0.000 [0.040] N=627

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,153	-0.010 [0.027] N=1,245	-0.007 [0.027] N=1,148	-0.019 [0.028] N=1,029	-0.023 [0.030] N=883	-0.041 [0.033] N=727
Vegetables consumed at least weekly	0.898 P=1,115	-0.007 [0.032] N=1,245	-0.025 [0.031] N=1,148	-0.040 [0.032] N=1,029	-0.034 [0.034] N=883	**-0.069* [0.039] N=727**
Meat fish or poultry consumed at least weekly	0.764 P=958	0.043 [0.053] N=1,245	0.043 [0.056] N=1,148	0.039 [0.061] N=1,029	0.042 [0.068] N=883	0.018 [0.074] N=727
Legumes consumed at least weekly	0.766 P=959	-0.051 [0.047] N=1,245	**-0.081* [0.049] N=1,148**	-0.071 [0.049] N=1,029	-0.068 [0.055] N=883	-0.072 [0.067] N=727
Animal products consumed at least weekly	0.907 P=1,132	0.012 [0.037] N=1,245	0.005 [0.039] N=1,148	-0.019 [0.043] N=1,029	-0.014 [0.047] N=883	-0.041 [0.053] N=727
Sweets consumed at least weekly	0.546 P=677	-0.042 [0.050] N=1,245	-0.041 [0.052] N=1,148	-0.031 [0.057] N=1,029	-0.046 [0.061] N=883	-0.023 [0.068] N=727
Snack foods consumed at least weekly	0.452 P=556	-0.004 [0.045] N=1,245	-0.016 [0.048] N=1,148	0.003 [0.051] N=1,029	-0.004 [0.056] N=883	-0.014 [0.059] N=727
Packaged soft drinks consumed at least weekly	0.260 P=316	-0.003 [0.045] N=1,245	-0.027 [0.047] N=1,148	-0.031 [0.048] N=1,029	-0.023 [0.052] N=883	-0.044 [0.058] N=727
Grains consumed at least weekly	0.865 P=1,078	0.016 [0.037] N=1,245	0.012 [0.038] N=1,148	0.005 [0.040] N=1,029	0.013 [0.044] N=883	-0.001 [0.050] N=727
Child was ever breastfed	0.968 P=1,416	-0.019 [0.013] N=1,467	-0.018 [0.014] N=1,356	-0.021 [0.016] N=1,212	-0.021 [0.017] N=1,034	-0.024 [0.017] N=855

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.004 [0.040] N=1,054	0.013 [0.041] N=957	-0.009 [0.046] N=838	-0.017 [0.055] N=692	-0.074 [0.073] N=536
Vegetables consumed at least weekly	0.015 [0.048] N=1,054	-0.010 [0.051] N=957	-0.041 [0.055] N=838	-0.028 [0.066] N=692	-0.139 [0.094] N=536
Meat fish or poultry consumed at least weekly	0.026 [0.065] N=1,054	0.024 [0.069] N=957	0.007 [0.081] N=838	-0.028 [0.090] N=692	-0.136 [0.091] N=536
Legumes consumed at least weekly	-0.040 [0.069] N=1,054	-0.096 [0.073] N=957	-0.070 [0.074] N=838	-0.089 [0.089] N=692	-0.113 [0.125] N=536
Animal products consumed at least weekly	0.039 [0.049] N=1,054	0.032 [0.055] N=957	-0.014 [0.063] N=838	0.014 [0.076] N=692	-0.032 [0.097] N=536
Sweets consumed at least weekly	-0.071 [0.065] N=1,054	-0.067 [0.069] N=957	-0.051 [0.080] N=838	-0.130 [0.089] N=692	-0.080 [0.125] N=536
Snack foods consumed at least weekly	-0.027 [0.061] N=1,054	-0.053 [0.066] N=957	-0.019 [0.075] N=838	-0.063 [0.093] N=692	-0.135 [0.112] N=536
Packaged soft drinks consumed at least weekly	-0.008 [0.067] N=1,054	-0.055 [0.073] N=957	-0.068 [0.079] N=838	-0.072 [0.093] N=692	-0.161 [0.128] N=536
Grains consumed at least weekly	0.053 [0.053] N=1,054	0.050 [0.058] N=957	0.038 [0.063] N=838	0.069 [0.076] N=692	0.019 [0.103] N=536
Child was ever breastfed	-0.007 [0.019] N=1,239	-0.003 [0.022] N=1,128	-0.006 [0.027] N=984	-0.004 [0.035] N=806	0.000 [0.040] N=627

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,153	-0.013 [0.027] N=1,245	-0.010 [0.027] N=1,148	-0.026 [0.029] N=1,029	-0.032 [0.030] N=883	-0.053 [0.034] N=727
Vegetables consumed at least weekly	0.898 P=1,115	-0.010 [0.032] N=1,245	-0.025 [0.032] N=1,148	-0.041 [0.032] N=1,029	-0.037 [0.034] N=883	**-0.070* [0.040] N=727**
Meat fish or poultry consumed at least weekly	0.764 P=958	0.035 [0.053] N=1,245	0.039 [0.056] N=1,148	0.029 [0.062] N=1,029	0.026 [0.069] N=883	-0.006 [0.076] N=727
Legumes consumed at least weekly	0.766 P=959	-0.055 [0.048] N=1,245	**-0.086* [0.051] N=1,148**	-0.079 [0.051] N=1,029	-0.076 [0.058] N=883	-0.086 [0.070] N=727
Animal products consumed at least weekly	0.907 P=1,132	0.009 [0.037] N=1,245	0.002 [0.039] N=1,148	-0.024 [0.042] N=1,029	-0.020 [0.047] N=883	-0.050 [0.052] N=727
Sweets consumed at least weekly	0.546 P=677	-0.045 [0.050] N=1,245	-0.046 [0.052] N=1,148	-0.041 [0.057] N=1,029	-0.053 [0.061] N=883	-0.032 [0.069] N=727
Snack foods consumed at least weekly	0.452 P=556	-0.011 [0.046] N=1,245	-0.021 [0.049] N=1,148	-0.009 [0.052] N=1,029	-0.015 [0.056] N=883	-0.025 [0.060] N=727
Packaged soft drinks consumed at least weekly	0.260 P=316	-0.003 [0.045] N=1,245	-0.032 [0.047] N=1,148	-0.038 [0.049] N=1,029	-0.027 [0.052] N=883	-0.045 [0.059] N=727
Grains consumed at least weekly	0.865 P=1,078	0.017 [0.037] N=1,245	0.013 [0.039] N=1,148	0.004 [0.040] N=1,029	0.011 [0.045] N=883	-0.003 [0.053] N=727
Child was ever breastfed	0.968 P=1,416	-0.022 [0.014] N=1,467	-0.020 [0.015] N=1,356	-0.025 [0.016] N=1,212	-0.025 [0.018] N=1,034	-0.028 [0.018] N=855

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.006 [0.041] N=1,054	0.017 [0.042] N=957	-0.010 [0.047] N=838	-0.026 [0.054] N=692	-0.099 [0.073] N=536
Vegetables consumed at least weekly	0.014 [0.049] N=1,054	-0.009 [0.052] N=957	-0.046 [0.056] N=838	-0.043 [0.065] N=692	**-0.173* [0.095] N=536**
Meat fish or poultry consumed at least weekly	0.025 [0.065] N=1,054	0.029 [0.069] N=957	0.006 [0.081] N=838	-0.032 [0.088] N=692	**-0.181* [0.095] N=536**
Legumes consumed at least weekly	-0.037 [0.070] N=1,054	-0.093 [0.074] N=957	-0.070 [0.074] N=838	-0.080 [0.089] N=692	-0.109 [0.131] N=536
Animal products consumed at least weekly	0.039 [0.050] N=1,054	0.034 [0.056] N=957	-0.017 [0.063] N=838	0.010 [0.075] N=692	-0.045 [0.104] N=536
Sweets consumed at least weekly	-0.062 [0.065] N=1,054	-0.062 [0.068] N=957	-0.048 [0.078] N=838	-0.115 [0.089] N=692	-0.076 [0.125] N=536
Snack foods consumed at least weekly	-0.020 [0.060] N=1,054	-0.047 [0.065] N=957	-0.015 [0.074] N=838	-0.056 [0.092] N=692	-0.135 [0.111] N=536
Packaged soft drinks consumed at least weekly	-0.004 [0.066] N=1,054	-0.057 [0.071] N=957	-0.072 [0.077] N=838	-0.069 [0.091] N=692	-0.155 [0.127] N=536
Grains consumed at least weekly	0.059 [0.053] N=1,054	0.053 [0.058] N=957	0.041 [0.063] N=838	0.071 [0.075] N=692	0.005 [0.102] N=536
Child was ever breastfed	-0.007 [0.020] N=1,239	-0.002 [0.023] N=1,128	-0.008 [0.028] N=984	-0.006 [0.037] N=806	0.001 [0.043] N=627

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=1,060	-0.006 [0.027] N=1,144	-0.009 [0.027] N=1,060	-0.019 [0.029] N=950	-0.027 [0.030] N=817	-0.048 [0.032] N=677
Vegetables consumed at least weekly	0.900 P=1,026	-0.011 [0.033] N=1,144	-0.030 [0.033] N=1,060	-0.046 [0.034] N=950	-0.036 [0.035] N=817	**-0.067* [0.039] N=677**
Meat fish or poultry consumed at least weekly	0.763 P=878	0.054 [0.051] N=1,144	0.045 [0.053] N=1,060	0.036 [0.057] N=950	0.020 [0.064] N=817	-0.002 [0.069] N=677
Legumes consumed at least weekly	0.761 P=875	-0.066 [0.049] N=1,144	**-0.103** [0.051] N=1,060**	**-0.096* [0.051] N=950**	**-0.100* [0.059] N=817**	-0.104 [0.069] N=677
Animal products consumed at least weekly	0.908 P=1,042	0.020 [0.039] N=1,144	0.011 [0.041] N=1,060	-0.015 [0.044] N=950	-0.020 [0.048] N=817	-0.044 [0.054] N=677
Sweets consumed at least weekly	0.548 P=622	-0.018 [0.053] N=1,144	-0.021 [0.056] N=1,060	-0.015 [0.060] N=950	-0.029 [0.066] N=817	-0.000 [0.074] N=677
Snack foods consumed at least weekly	0.454 P=511	-0.012 [0.050] N=1,144	-0.020 [0.053] N=1,060	-0.002 [0.057] N=950	-0.013 [0.062] N=817	-0.004 [0.068] N=677
Packaged soft drinks consumed at least weekly	0.262 P=293	0.016 [0.045] N=1,144	-0.011 [0.047] N=1,060	-0.029 [0.049] N=950	-0.016 [0.051] N=817	-0.024 [0.056] N=677
Grains consumed at least weekly	0.866 P=991	0.013 [0.037] N=1,144	0.005 [0.039] N=1,060	-0.011 [0.040] N=950	-0.003 [0.045] N=817	-0.020 [0.050] N=677
Child was ever breastfed	0.969 P=1,298	**-0.025* [0.014] N=1,344**	-0.020 [0.015] N=1,247	-0.023 [0.016] N=1,112	-0.020 [0.018] N=954	-0.024 [0.017] N=794

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.043] N=966	0.008 [0.045] N=882	-0.016 [0.050] N=772	-0.043 [0.059] N=639	-0.103 [0.075] N=499
Vegetables consumed at least weekly	0.013 [0.053] N=966	-0.017 [0.056] N=882	-0.053 [0.061] N=772	-0.041 [0.071] N=639	-0.138 [0.091] N=499
Meat fish or poultry consumed at least weekly	0.053 [0.069] N=966	0.029 [0.072] N=882	0.010 [0.081] N=772	-0.065 [0.093] N=639	**-0.156* [0.093] N=499**
Legumes consumed at least weekly	-0.041 [0.073] N=966	-0.113 [0.077] N=882	-0.095 [0.075] N=772	-0.133 [0.091] N=639	-0.150 [0.118] N=499
Animal products consumed at least weekly	0.052 [0.055] N=966	0.039 [0.061] N=882	-0.011 [0.067] N=772	-0.007 [0.082] N=639	-0.053 [0.106] N=499
Sweets consumed at least weekly	-0.047 [0.068] N=966	-0.056 [0.075] N=882	-0.057 [0.086] N=772	-0.137 [0.096] N=639	-0.085 [0.132] N=499
Snack foods consumed at least weekly	-0.038 [0.066] N=966	-0.061 [0.072] N=882	-0.035 [0.081] N=772	-0.093 [0.099] N=639	-0.132 [0.118] N=499
Packaged soft drinks consumed at least weekly	0.006 [0.070] N=966	-0.050 [0.074] N=882	-0.090 [0.081] N=772	-0.092 [0.094] N=639	-0.139 [0.132] N=499
Grains consumed at least weekly	0.060 [0.055] N=966	0.049 [0.060] N=882	0.017 [0.064] N=772	0.042 [0.076] N=639	0.000 [0.104] N=499
Child was ever breastfed	-0.020 [0.021] N=1,132	-0.011 [0.025] N=1,035	-0.016 [0.030] N=900	-0.013 [0.038] N=742	-0.008 [0.043] N=582

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=1,060	-0.009 [0.027] N=1,144	-0.011 [0.027] N=1,060	-0.020 [0.028] N=950	-0.027 [0.030] N=817	-0.049 [0.031] N=677
Vegetables consumed at least weekly	0.900 P=1,026	-0.012 [0.033] N=1,144	-0.031 [0.033] N=1,060	-0.047 [0.034] N=950	-0.038 [0.035] N=817	**-0.070* [0.039] N=677**
Meat fish or poultry consumed at least weekly	0.763 P=878	0.050 [0.052] N=1,144	0.042 [0.053] N=1,060	0.035 [0.057] N=950	0.023 [0.064] N=817	0.003 [0.069] N=677
Legumes consumed at least weekly	0.761 P=875	-0.066 [0.048] N=1,144	**-0.104** [0.051] N=1,060**	**-0.098* [0.051] N=950**	**-0.103* [0.059] N=817**	-0.111 [0.070] N=677
Animal products consumed at least weekly	0.908 P=1,042	0.019 [0.039] N=1,144	0.010 [0.041] N=1,060	-0.015 [0.043] N=950	-0.018 [0.048] N=817	-0.043 [0.053] N=677
Sweets consumed at least weekly	0.548 P=622	-0.017 [0.052] N=1,144	-0.021 [0.055] N=1,060	-0.014 [0.060] N=950	-0.029 [0.065] N=817	-0.001 [0.073] N=677
Snack foods consumed at least weekly	0.454 P=511	-0.013 [0.049] N=1,144	-0.022 [0.053] N=1,060	-0.002 [0.056] N=950	-0.013 [0.062] N=817	-0.006 [0.067] N=677
Packaged soft drinks consumed at least weekly	0.262 P=293	0.016 [0.046] N=1,144	-0.011 [0.048] N=1,060	-0.026 [0.049] N=950	-0.012 [0.051] N=817	-0.022 [0.056] N=677
Grains consumed at least weekly	0.866 P=991	0.013 [0.037] N=1,144	0.006 [0.039] N=1,060	-0.011 [0.040] N=950	-0.002 [0.044] N=817	-0.020 [0.049] N=677
Child was ever breastfed	0.969 P=1,298	**-0.024* [0.014] N=1,344**	-0.020 [0.015] N=1,247	-0.023 [0.016] N=1,112	-0.020 [0.018] N=954	-0.024 [0.017] N=794

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.042] N=966	0.002 [0.044] N=882	-0.017 [0.050] N=772	-0.041 [0.058] N=639	-0.104 [0.071] N=499
Vegetables consumed at least weekly	0.010 [0.052] N=966	-0.019 [0.056] N=882	-0.055 [0.060] N=772	-0.046 [0.070] N=639	-0.150 [0.091] N=499
Meat fish or poultry consumed at least weekly	0.047 [0.070] N=966	0.026 [0.073] N=882	0.009 [0.081] N=772	-0.065 [0.093] N=639	-0.158 [0.096] N=499
Legumes consumed at least weekly	-0.045 [0.072] N=966	-0.117 [0.077] N=882	-0.096 [0.075] N=772	-0.135 [0.092] N=639	-0.160 [0.118] N=499
Animal products consumed at least weekly	0.047 [0.054] N=966	0.034 [0.059] N=882	-0.011 [0.066] N=772	-0.005 [0.080] N=639	-0.056 [0.103] N=499
Sweets consumed at least weekly	-0.055 [0.068] N=966	-0.066 [0.074] N=882	-0.066 [0.086] N=772	-0.141 [0.095] N=639	-0.097 [0.130] N=499
Snack foods consumed at least weekly	-0.047 [0.066] N=966	-0.071 [0.071] N=882	-0.043 [0.081] N=772	-0.099 [0.099] N=639	-0.142 [0.118] N=499
Packaged soft drinks consumed at least weekly	-0.000 [0.071] N=966	-0.057 [0.076] N=882	-0.093 [0.082] N=772	-0.088 [0.096] N=639	-0.136 [0.133] N=499
Grains consumed at least weekly	0.058 [0.054] N=966	0.049 [0.059] N=882	0.019 [0.063] N=772	0.045 [0.076] N=639	0.002 [0.103] N=499
Child was ever breastfed	-0.018 [0.021] N=1,132	-0.009 [0.024] N=1,035	-0.015 [0.030] N=900	-0.013 [0.037] N=742	-0.011 [0.043] N=582

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=996	**0.055* [0.033] N=1,068**	**0.058* [0.034] N=965**	0.029 [0.037] N=841	0.010 [0.039] N=702	0.014 [0.045] N=573
Vegetables consumed at least weekly	0.898 P=958	0.043 [0.038] N=1,068	0.053 [0.039] N=965	0.045 [0.043] N=841	0.019 [0.046] N=702	0.028 [0.048] N=573
Meat fish or poultry consumed at least weekly	0.759 P=821	**0.105* [0.055] N=1,068**	**0.106* [0.056] N=965**	0.096 [0.064] N=841	**0.136** [0.067] N=702**	0.117 [0.077] N=573
Legumes consumed at least weekly	0.762 P=825	-0.008 [0.047] N=1,068	0.034 [0.049] N=965	0.022 [0.055] N=841	-0.012 [0.059] N=702	0.020 [0.067] N=573
Animal products consumed at least weekly	0.906 P=971	0.016 [0.041] N=1,068	0.014 [0.044] N=965	-0.004 [0.051] N=841	-0.017 [0.055] N=702	-0.015 [0.064] N=573
Sweets consumed at least weekly	0.550 P=589	-0.050 [0.062] N=1,068	-0.057 [0.065] N=965	-0.068 [0.070] N=841	-0.059 [0.072] N=702	-0.118 [0.085] N=573
Snack foods consumed at least weekly	0.455 P=484	-0.028 [0.058] N=1,068	-0.026 [0.062] N=965	0.007 [0.065] N=841	-0.004 [0.071] N=702	-0.052 [0.081] N=573
Packaged soft drinks consumed at least weekly	0.247 P=259	-0.070 [0.047] N=1,068	-0.079 [0.049] N=965	-0.025 [0.054] N=841	-0.052 [0.058] N=702	-0.078 [0.065] N=573
Grains consumed at least weekly	0.855 P=917	0.051 [0.044] N=1,068	0.060 [0.048] N=965	0.052 [0.051] N=841	0.061 [0.056] N=702	0.040 [0.067] N=573
Child was ever breastfed	0.967 P=1,219	-0.002 [0.023] N=1,262	-0.007 [0.023] N=1,142	0.007 [0.022] N=992	0.007 [0.022] N=834	0.013 [0.021] N=676

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.055 [0.051] N=929	0.058 [0.054] N=826	0.005 [0.061] N=702	-0.004 [0.069] N=563	0.004 [0.093] N=434
Vegetables consumed at least weekly	0.002 [0.061] N=929	0.012 [0.066] N=826	-0.016 [0.076] N=702	-0.064 [0.087] N=563	-0.064 [0.109] N=434
Meat fish or poultry consumed at least weekly	0.044 [0.074] N=929	0.034 [0.076] N=826	-0.015 [0.088] N=702	0.057 [0.095] N=563	-0.012 [0.125] N=434
Legumes consumed at least weekly	-0.055 [0.058] N=929	0.007 [0.062] N=826	-0.016 [0.075] N=702	-0.080 [0.085] N=563	0.002 [0.099] N=434
Animal products consumed at least weekly	-0.020 [0.053] N=929	-0.024 [0.058] N=826	-0.063 [0.071] N=702	-0.087 [0.072] N=563	-0.102 [0.092] N=434
Sweets consumed at least weekly	-0.037 [0.083] N=929	-0.044 [0.089] N=826	-0.071 [0.101] N=702	-0.056 [0.115] N=563	-0.235 [0.150] N=434
Snack foods consumed at least weekly	-0.054 [0.079] N=929	-0.053 [0.084] N=826	-0.006 [0.094] N=702	-0.016 [0.110] N=563	-0.171 [0.132] N=434
Packaged soft drinks consumed at least weekly	-0.084 [0.072] N=929	-0.095 [0.077] N=826	0.006 [0.092] N=702	-0.034 [0.106] N=563	-0.074 [0.126] N=434
Grains consumed at least weekly	-0.042 [0.051] N=929	-0.044 [0.057] N=826	-0.086 [0.060] N=702	**-0.126* [0.069] N=563**	**-0.276*** [0.088] N=434**
Child was ever breastfed	0.012 [0.041] N=1,098	0.005 [0.043] N=978	0.029 [0.046] N=828	0.033 [0.053] N=670	0.065 [0.064] N=512

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=996	**0.055* [0.033] N=1,068**	**0.058* [0.034] N=965**	0.029 [0.037] N=841	0.010 [0.040] N=702	0.014 [0.045] N=573
Vegetables consumed at least weekly	0.898 P=958	0.044 [0.037] N=1,068	0.053 [0.039] N=965	0.046 [0.043] N=841	0.020 [0.046] N=702	0.027 [0.048] N=573
Meat fish or poultry consumed at least weekly	0.759 P=821	**0.106* [0.055] N=1,068**	**0.107* [0.056] N=965**	0.097 [0.064] N=841	**0.137** [0.067] N=702**	0.117 [0.077] N=573
Legumes consumed at least weekly	0.762 P=825	-0.008 [0.047] N=1,068	0.035 [0.049] N=965	0.022 [0.055] N=841	-0.012 [0.059] N=702	0.020 [0.067] N=573
Animal products consumed at least weekly	0.906 P=971	0.017 [0.040] N=1,068	0.015 [0.043] N=965	-0.003 [0.051] N=841	-0.016 [0.056] N=702	-0.015 [0.064] N=573
Sweets consumed at least weekly	0.550 P=589	-0.048 [0.062] N=1,068	-0.055 [0.065] N=965	-0.067 [0.070] N=841	-0.059 [0.072] N=702	-0.118 [0.085] N=573
Snack foods consumed at least weekly	0.455 P=484	-0.025 [0.057] N=1,068	-0.024 [0.061] N=965	0.009 [0.064] N=841	-0.002 [0.070] N=702	-0.052 [0.080] N=573
Packaged soft drinks consumed at least weekly	0.247 P=259	-0.070 [0.047] N=1,068	-0.079 [0.049] N=965	-0.024 [0.053] N=841	-0.052 [0.058] N=702	-0.078 [0.064] N=573
Grains consumed at least weekly	0.855 P=917	0.051 [0.044] N=1,068	0.061 [0.047] N=965	0.053 [0.050] N=841	0.061 [0.056] N=702	0.040 [0.067] N=573
Child was ever breastfed	0.967 P=1,219	-0.002 [0.023] N=1,262	-0.007 [0.023] N=1,142	0.006 [0.022] N=992	0.006 [0.023] N=834	0.013 [0.021] N=676

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.056 [0.051] N=929	0.059 [0.053] N=826	0.006 [0.061] N=702	-0.005 [0.070] N=563	0.003 [0.095] N=434
Vegetables consumed at least weekly	0.003 [0.060] N=929	0.013 [0.065] N=826	-0.014 [0.075] N=702	-0.059 [0.086] N=563	-0.062 [0.108] N=434
Meat fish or poultry consumed at least weekly	0.045 [0.073] N=929	0.037 [0.075] N=826	-0.013 [0.086] N=702	0.061 [0.094] N=563	-0.011 [0.125] N=434
Legumes consumed at least weekly	-0.054 [0.058] N=929	0.007 [0.062] N=826	-0.015 [0.075] N=702	-0.081 [0.085] N=563	-0.004 [0.098] N=434
Animal products consumed at least weekly	-0.017 [0.050] N=929	-0.021 [0.055] N=826	-0.059 [0.068] N=702	-0.079 [0.070] N=563	-0.094 [0.086] N=434
Sweets consumed at least weekly	-0.035 [0.082] N=929	-0.041 [0.087] N=826	-0.067 [0.100] N=702	-0.056 [0.116] N=563	-0.235 [0.149] N=434
Snack foods consumed at least weekly	-0.049 [0.077] N=929	-0.049 [0.082] N=826	-0.000 [0.092] N=702	-0.010 [0.108] N=563	-0.166 [0.129] N=434
Packaged soft drinks consumed at least weekly	-0.083 [0.071] N=929	-0.094 [0.077] N=826	0.007 [0.091] N=702	-0.033 [0.106] N=563	-0.071 [0.124] N=434
Grains consumed at least weekly	-0.040 [0.051] N=929	-0.043 [0.057] N=826	-0.085 [0.060] N=702	**-0.129* [0.069] N=563**	**-0.278*** [0.090] N=434**
Child was ever breastfed	0.012 [0.041] N=1,098	0.005 [0.044] N=978	0.028 [0.046] N=828	0.032 [0.053] N=670	0.065 [0.065] N=512

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=996	0.050 [0.033] N=1,068	0.052 [0.034] N=965	0.026 [0.038] N=841	0.008 [0.041] N=702	0.013 [0.047] N=573
Vegetables consumed at least weekly	0.898 P=958	0.039 [0.038] N=1,068	0.048 [0.039] N=965	0.042 [0.043] N=841	0.016 [0.047] N=702	0.023 [0.049] N=573
Meat fish or poultry consumed at least weekly	0.759 P=821	0.089 [0.055] N=1,068	0.088 [0.057] N=965	0.091 [0.064] N=841	**0.132** [0.066] N=702**	0.115 [0.077] N=573
Legumes consumed at least weekly	0.762 P=825	-0.014 [0.048] N=1,068	0.028 [0.050] N=965	0.019 [0.056] N=841	-0.015 [0.061] N=702	0.021 [0.071] N=573
Animal products consumed at least weekly	0.906 P=971	0.011 [0.041] N=1,068	0.006 [0.044] N=965	-0.008 [0.051] N=841	-0.019 [0.057] N=702	-0.019 [0.065] N=573
Sweets consumed at least weekly	0.550 P=589	-0.050 [0.062] N=1,068	-0.060 [0.065] N=965	-0.072 [0.070] N=841	-0.061 [0.073] N=702	-0.123 [0.086] N=573
Snack foods consumed at least weekly	0.455 P=484	-0.026 [0.057] N=1,068	-0.026 [0.061] N=965	0.003 [0.065] N=841	-0.007 [0.071] N=702	-0.060 [0.082] N=573
Packaged soft drinks consumed at least weekly	0.247 P=259	-0.064 [0.047] N=1,068	-0.075 [0.049] N=965	-0.026 [0.053] N=841	-0.049 [0.058] N=702	-0.079 [0.065] N=573
Grains consumed at least weekly	0.855 P=917	0.048 [0.045] N=1,068	0.058 [0.048] N=965	0.052 [0.052] N=841	0.062 [0.058] N=702	0.041 [0.069] N=573
Child was ever breastfed	0.967 P=1,219	-0.004 [0.023] N=1,262	-0.009 [0.024] N=1,142	0.006 [0.023] N=992	0.006 [0.023] N=834	0.012 [0.022] N=676

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.047 [0.050] N=929	0.049 [0.052] N=826	0.004 [0.060] N=702	-0.008 [0.071] N=563	-0.003 [0.097] N=434
Vegetables consumed at least weekly	-0.004 [0.060] N=929	0.005 [0.065] N=826	-0.019 [0.074] N=702	-0.066 [0.086] N=563	-0.074 [0.109] N=434
Meat fish or poultry consumed at least weekly	0.017 [0.072] N=929	0.007 [0.074] N=826	-0.022 [0.085] N=702	0.045 [0.097] N=563	-0.038 [0.124] N=434
Legumes consumed at least weekly	-0.061 [0.058] N=929	0.001 [0.063] N=826	-0.019 [0.074] N=702	-0.088 [0.085] N=563	-0.004 [0.100] N=434
Animal products consumed at least weekly	-0.023 [0.050] N=929	-0.029 [0.055] N=826	-0.060 [0.067] N=702	-0.080 [0.070] N=563	-0.097 [0.088] N=434
Sweets consumed at least weekly	-0.030 [0.079] N=929	-0.039 [0.085] N=826	-0.065 [0.096] N=702	-0.049 [0.113] N=563	-0.219 [0.144] N=434
Snack foods consumed at least weekly	-0.042 [0.076] N=929	-0.042 [0.081] N=826	-0.001 [0.091] N=702	-0.007 [0.107] N=563	-0.161 [0.127] N=434
Packaged soft drinks consumed at least weekly	-0.066 [0.071] N=929	-0.077 [0.076] N=826	0.015 [0.090] N=702	-0.019 [0.103] N=563	-0.050 [0.122] N=434
Grains consumed at least weekly	-0.047 [0.051] N=929	-0.048 [0.057] N=826	-0.085 [0.060] N=702	**-0.128* [0.069] N=563**	**-0.275*** [0.088] N=434**
Child was ever breastfed	0.009 [0.040] N=1,098	-0.000 [0.043] N=978	0.028 [0.045] N=828	0.030 [0.051] N=670	0.059 [0.062] N=512

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=996	0.046 [0.033] N=1,068	0.049 [0.034] N=965	0.025 [0.038] N=841	-0.000 [0.040] N=702	0.001 [0.046] N=573
Vegetables consumed at least weekly	0.898 P=958	0.034 [0.038] N=1,068	0.043 [0.040] N=965	0.036 [0.043] N=841	0.006 [0.045] N=702	0.011 [0.047] N=573
Meat fish or poultry consumed at least weekly	0.759 P=821	0.084 [0.055] N=1,068	0.084 [0.057] N=965	0.078 [0.064] N=841	0.106 [0.066] N=702	0.075 [0.074] N=573
Legumes consumed at least weekly	0.762 P=825	-0.010 [0.048] N=1,068	0.031 [0.051] N=965	0.025 [0.056] N=841	-0.016 [0.059] N=702	0.010 [0.067] N=573
Animal products consumed at least weekly	0.906 P=971	0.017 [0.040] N=1,068	0.015 [0.043] N=965	0.004 [0.051] N=841	-0.013 [0.056] N=702	-0.017 [0.064] N=573
Sweets consumed at least weekly	0.550 P=589	-0.038 [0.063] N=1,068	-0.047 [0.066] N=965	-0.060 [0.072] N=841	-0.054 [0.075] N=702	-0.117 [0.087] N=573
Snack foods consumed at least weekly	0.455 P=484	-0.015 [0.059] N=1,068	-0.014 [0.062] N=965	0.018 [0.067] N=841	0.003 [0.072] N=702	-0.046 [0.083] N=573
Packaged soft drinks consumed at least weekly	0.247 P=259	-0.052 [0.047] N=1,068	-0.064 [0.049] N=965	-0.015 [0.054] N=841	-0.040 [0.058] N=702	-0.062 [0.066] N=573
Grains consumed at least weekly	0.855 P=917	0.047 [0.045] N=1,068	0.056 [0.048] N=965	0.050 [0.051] N=841	0.057 [0.057] N=702	0.034 [0.067] N=573
Child was ever breastfed	0.967 P=1,219	-0.002 [0.022] N=1,262	-0.007 [0.022] N=1,142	0.008 [0.021] N=992	0.006 [0.022] N=834	0.012 [0.020] N=676

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.046 [0.050] N=929	0.047 [0.052] N=826	0.005 [0.061] N=702	-0.023 [0.072] N=563	-0.033 [0.095] N=434
Vegetables consumed at least weekly	-0.012 [0.061] N=929	-0.004 [0.066] N=826	-0.028 [0.075] N=702	-0.088 [0.085] N=563	-0.114 [0.107] N=434
Meat fish or poultry consumed at least weekly	0.021 [0.075] N=929	0.009 [0.079] N=826	-0.032 [0.091] N=702	0.010 [0.106] N=563	-0.112 [0.127] N=434
Legumes consumed at least weekly	-0.054 [0.059] N=929	0.006 [0.065] N=826	0.003 [0.079] N=702	-0.078 [0.088] N=563	-0.015 [0.103] N=434
Animal products consumed at least weekly	-0.016 [0.047] N=929	-0.019 [0.052] N=826	-0.043 [0.066] N=702	-0.071 [0.070] N=563	-0.099 [0.085] N=434
Sweets consumed at least weekly	-0.013 [0.081] N=929	-0.020 [0.086] N=826	-0.043 [0.098] N=702	-0.030 [0.114] N=563	-0.204 [0.144] N=434
Snack foods consumed at least weekly	-0.033 [0.076] N=929	-0.033 [0.082] N=826	0.020 [0.093] N=702	0.003 [0.105] N=563	-0.143 [0.125] N=434
Packaged soft drinks consumed at least weekly	-0.061 [0.070] N=929	-0.073 [0.075] N=826	0.018 [0.090] N=702	-0.014 [0.104] N=563	-0.021 [0.126] N=434
Grains consumed at least weekly	-0.042 [0.051] N=929	-0.044 [0.057] N=826	-0.083 [0.061] N=702	**-0.134* [0.070] N=563**	**-0.286*** [0.087] N=434**
Child was ever breastfed	0.011 [0.040] N=1,098	0.002 [0.042] N=978	0.032 [0.043] N=828	0.031 [0.049] N=670	0.055 [0.059] N=512

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=913	0.053 [0.035] N=977	0.053 [0.036] N=880	0.039 [0.040] N=774	0.008 [0.043] N=645	0.008 [0.049] N=531
Vegetables consumed at least weekly	0.902 P=880	0.033 [0.038] N=977	0.038 [0.041] N=880	0.052 [0.043] N=774	0.016 [0.049] N=645	0.002 [0.052] N=531
Meat fish or poultry consumed at least weekly	0.759 P=750	**0.096* [0.056] N=977**	0.088 [0.058] N=880	0.090 [0.064] N=774	0.109 [0.069] N=645	0.088 [0.079] N=531
Legumes consumed at least weekly	0.759 P=752	-0.008 [0.052] N=977	0.036 [0.055] N=880	0.026 [0.060] N=774	-0.021 [0.066] N=645	0.006 [0.078] N=531
Animal products consumed at least weekly	0.909 P=891	0.016 [0.044] N=977	0.015 [0.047] N=880	0.010 [0.055] N=774	-0.010 [0.061] N=645	-0.019 [0.069] N=531
Sweets consumed at least weekly	0.562 P=549	-0.044 [0.064] N=977	-0.049 [0.066] N=880	-0.072 [0.072] N=774	-0.076 [0.077] N=645	-0.138 [0.090] N=531
Snack foods consumed at least weekly	0.463 P=450	-0.015 [0.060] N=977	-0.013 [0.064] N=880	0.008 [0.067] N=774	-0.008 [0.074] N=645	-0.054 [0.084] N=531
Packaged soft drinks consumed at least weekly	0.254 P=245	-0.079 [0.048] N=977	**-0.093* [0.050] N=880**	-0.038 [0.054] N=774	-0.072 [0.060] N=645	-0.110 [0.069] N=531
Grains consumed at least weekly	0.856 P=840	0.067 [0.047] N=977	0.074 [0.050] N=880	0.072 [0.053] N=774	0.073 [0.061] N=645	0.040 [0.071] N=531
Child was ever breastfed	0.970 P=1,117	-0.011 [0.021] N=1,153	-0.016 [0.022] N=1,042	-0.002 [0.020] N=912	-0.007 [0.021] N=767	0.006 [0.020] N=626

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.041 [0.052] N=848	0.041 [0.054] N=751	0.015 [0.062] N=645	-0.035 [0.072] N=516	-0.057 [0.090] N=402
Vegetables consumed at least weekly	-0.020 [0.055] N=848	-0.018 [0.060] N=751	-0.008 [0.066] N=645	-0.083 [0.080] N=516	-0.148 [0.096] N=402
Meat fish or poultry consumed at least weekly	0.058 [0.082] N=848	0.043 [0.084] N=751	0.022 [0.094] N=645	0.037 [0.104] N=516	-0.045 [0.129] N=402
Legumes consumed at least weekly	-0.045 [0.064] N=848	0.018 [0.068] N=751	0.016 [0.078] N=645	-0.059 [0.093] N=516	0.008 [0.108] N=402
Animal products consumed at least weekly	-0.033 [0.051] N=848	-0.035 [0.056] N=751	-0.047 [0.067] N=645	-0.091 [0.074] N=516	-0.142 [0.093] N=402
Sweets consumed at least weekly	-0.028 [0.085] N=848	-0.033 [0.091] N=751	-0.091 [0.103] N=645	-0.111 [0.118] N=516	**-0.301* [0.156] N=402**
Snack foods consumed at least weekly	-0.042 [0.081] N=848	-0.040 [0.087] N=751	-0.019 [0.094] N=645	-0.051 [0.108] N=516	-0.183 [0.135] N=402
Packaged soft drinks consumed at least weekly	-0.096 [0.073] N=848	-0.115 [0.079] N=751	-0.020 [0.094] N=645	-0.085 [0.114] N=516	-0.193 [0.138] N=402
Grains consumed at least weekly	-0.023 [0.057] N=848	-0.027 [0.064] N=751	-0.049 [0.068] N=645	-0.126 [0.082] N=516	**-0.325*** [0.099] N=402**
Child was ever breastfed	-0.002 [0.038] N=1,001	-0.011 [0.041] N=890	0.014 [0.040] N=760	0.005 [0.048] N=615	0.040 [0.056] N=474

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=913	0.054 [0.035] N=977	0.056 [0.036] N=880	0.040 [0.039] N=774	0.008 [0.042] N=645	0.007 [0.048] N=531
Vegetables consumed at least weekly	0.902 P=880	0.037 [0.038] N=977	0.041 [0.040] N=880	0.056 [0.042] N=774	0.017 [0.048] N=645	0.004 [0.051] N=531
Meat fish or poultry consumed at least weekly	0.759 P=750	**0.105* [0.054] N=977**	**0.099* [0.056] N=880**	0.098 [0.062] N=774	0.108 [0.066] N=645	0.085 [0.076] N=531
Legumes consumed at least weekly	0.759 P=752	-0.008 [0.051] N=977	0.035 [0.054] N=880	0.026 [0.059] N=774	-0.025 [0.065] N=645	0.002 [0.076] N=531
Animal products consumed at least weekly	0.909 P=891	0.014 [0.043] N=977	0.015 [0.047] N=880	0.010 [0.054] N=774	-0.010 [0.060] N=645	-0.017 [0.068] N=531
Sweets consumed at least weekly	0.562 P=549	-0.047 [0.063] N=977	-0.052 [0.066] N=880	-0.077 [0.072] N=774	-0.082 [0.076] N=645	-0.140 [0.091] N=531
Snack foods consumed at least weekly	0.463 P=450	-0.013 [0.059] N=977	-0.012 [0.063] N=880	0.010 [0.066] N=774	-0.006 [0.072] N=645	-0.049 [0.083] N=531
Packaged soft drinks consumed at least weekly	0.254 P=245	**-0.081* [0.048] N=977**	**-0.095* [0.050] N=880**	-0.042 [0.053] N=774	-0.076 [0.059] N=645	**-0.114* [0.068] N=531**
Grains consumed at least weekly	0.856 P=840	0.067 [0.047] N=977	0.074 [0.050] N=880	0.072 [0.052] N=774	0.073 [0.060] N=645	0.040 [0.071] N=531
Child was ever breastfed	0.970 P=1,117	-0.010 [0.022] N=1,153	-0.016 [0.023] N=1,042	0.000 [0.021] N=912	-0.003 [0.022] N=767	0.008 [0.020] N=626

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.046 [0.051] N=848	0.047 [0.054] N=751	0.022 [0.062] N=645	-0.032 [0.073] N=516	-0.049 [0.095] N=402
Vegetables consumed at least weekly	-0.017 [0.055] N=848	-0.016 [0.061] N=751	-0.003 [0.067] N=645	-0.087 [0.081] N=516	-0.152 [0.098] N=402
Meat fish or poultry consumed at least weekly	0.069 [0.081] N=848	0.052 [0.084] N=751	0.026 [0.093] N=645	0.031 [0.102] N=516	-0.049 [0.129] N=402
Legumes consumed at least weekly	-0.045 [0.064] N=848	0.020 [0.068] N=751	0.017 [0.078] N=645	-0.069 [0.092] N=516	0.011 [0.106] N=402
Animal products consumed at least weekly	-0.033 [0.051] N=848	-0.034 [0.056] N=751	-0.046 [0.068] N=645	-0.093 [0.075] N=516	-0.145 [0.096] N=402
Sweets consumed at least weekly	-0.027 [0.086] N=848	-0.033 [0.092] N=751	-0.090 [0.104] N=645	-0.110 [0.119] N=516	**-0.305* [0.155] N=402**
Snack foods consumed at least weekly	-0.039 [0.081] N=848	-0.040 [0.087] N=751	-0.012 [0.095] N=645	-0.035 [0.110] N=516	-0.177 [0.134] N=402
Packaged soft drinks consumed at least weekly	-0.096 [0.074] N=848	-0.114 [0.080] N=751	-0.013 [0.093] N=645	-0.070 [0.111] N=516	-0.162 [0.135] N=402
Grains consumed at least weekly	-0.023 [0.057] N=848	-0.028 [0.063] N=751	-0.051 [0.066] N=645	-0.132 [0.080] N=516	**-0.329*** [0.097] N=402**
Child was ever breastfed	-0.002 [0.038] N=1,001	-0.012 [0.041] N=890	0.017 [0.041] N=760	0.011 [0.050] N=615	0.045 [0.059] N=474

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.