Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Oldest Child

Age reference: 12 November 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	0.011 [0.250] N=1,257	0.038 [0.261] N=1,164	0.073 [0.289] N=1,039	0.111 [0.306] N=892	-0.101 [0.337] N=741
WHO weight-for-age z score	0.010	-0.140 [0.136] N=1,451	-0.105 [0.150] N=1,342	-0.076 [0.155] N=1,200	-0.042 [0.154] N=1,024	-0.093 [0.169] N=846
WHO weight-for-length/height z score, 0-60 months	0.312	0.315 [0.227] N=1,255	0.316 [0.238] N=1,163	0.317 [0.254] N=1,042	0.377 [0.260] N=891	**0.467* [0.278] N=738**
WHO BMI-for-age z score	0.301	0.286 [0.252] N=1,231	0.298 [0.264] N=1,142	0.284 [0.290] N=1,020	0.239 [0.307] N=871	0.301 [0.334] N=723
Stunted: HAZ below -2	0.245 P=296	-0.016 [0.048] N=1,257	-0.020 [0.051] N=1,164	-0.019 [0.054] N=1,039	-0.034 [0.058] N=892	-0.012 [0.061] N=741
Underweight: WAZ below -2	0.083 P=120	**0.071*** [0.026] N=1,451**	**0.056** [0.027] N=1,342**	**0.050* [0.029] N=1,200**	0.046 [0.031] N=1,024	0.040 [0.034] N=846
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=152	-0.033 [0.037] N=1,255	-0.022 [0.039] N=1,163	-0.032 [0.042] N=1,042	-0.020 [0.044] N=891	-0.034 [0.046] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=244	0.052 [0.047] N=1,255	0.058 [0.050] N=1,163	0.047 [0.053] N=1,042	0.064 [0.059] N=891	0.059 [0.066] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=112	0.012 [0.035] N=1,255	0.020 [0.037] N=1,163	0.030 [0.038] N=1,042	0.042 [0.042] N=891	0.056 [0.046] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=393	-0.031 [0.051] N=1,202	-0.022 [0.053] N=1,113	-0.022 [0.057] N=996	-0.032 [0.062] N=853	-0.024 [0.068] N=708
BMI-for-age z score below -2	0.131 P=160	-0.021 [0.041] N=1,231	-0.015 [0.043] N=1,142	-0.012 [0.048] N=1,020	0.021 [0.051] N=871	0.019 [0.055] N=723

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.018 [0.372] N=1,056	0.078 [0.400] N=963	0.239 [0.470] N=838	0.500 [0.514] N=691	0.224 [0.676] N=540
WHO weight-for-age z score	**-0.392** [0.193] N=1,225**	-0.366 [0.232] N=1,116	-0.350 [0.241] N=974	-0.384 [0.279] N=798	**-0.701* [0.395] N=620**
WHO weight-for-length/height z score, 0-60 months	0.145 [0.332] N=1,054	0.101 [0.360] N=962	0.016 [0.396] N=841	0.116 [0.445] N=690	0.062 [0.527] N=537
WHO BMI-for-age z score	0.196 [0.332] N=1,032	0.161 [0.354] N=943	0.042 [0.407] N=821	-0.144 [0.443] N=672	-0.327 [0.510] N=524
Stunted: HAZ below -2	-0.009 [0.069] N=1,056	-0.021 [0.076] N=963	-0.031 [0.085] N=838	-0.080 [0.101] N=691	-0.071 [0.135] N=540
Underweight: WAZ below -2	**0.113*** [0.036] N=1,225**	**0.089** [0.040] N=1,116**	**0.087** [0.042] N=974**	**0.099** [0.050] N=798**	0.108 [0.069] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	-0.041 [0.055] N=1,054	-0.018 [0.061] N=962	-0.035 [0.070] N=841	-0.024 [0.085] N=690	-0.055 [0.099] N=537
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.058] N=1,054	0.014 [0.064] N=962	-0.020 [0.065] N=841	0.002 [0.083] N=690	-0.080 [0.100] N=537
Obese: WLZ/WHZ above +3, 0-60 months	-0.020 [0.042] N=1,054	-0.011 [0.049] N=962	-0.001 [0.050] N=841	0.022 [0.058] N=690	0.033 [0.075] N=537
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.004 [0.077] N=1,007	0.012 [0.084] N=918	0.010 [0.093] N=801	-0.022 [0.119] N=658	-0.010 [0.157] N=513
BMI-for-age z score below -2	-0.042 [0.056] N=1,032	-0.030 [0.061] N=943	-0.025 [0.072] N=821	0.035 [0.084] N=672	0.057 [0.098] N=524

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	0.011 [0.251] N=1,257	0.038 [0.261] N=1,164	0.073 [0.288] N=1,039	0.112 [0.307] N=892	-0.095 [0.338] N=741
WHO weight-for-age z score	0.010	-0.140 [0.135] N=1,451	-0.108 [0.150] N=1,342	-0.078 [0.154] N=1,200	-0.046 [0.154] N=1,024	-0.101 [0.168] N=846
WHO weight-for-length/height z score, 0-60 months	0.312	0.314 [0.228] N=1,255	0.315 [0.238] N=1,163	0.316 [0.254] N=1,042	0.376 [0.261] N=891	0.462 [0.281] N=738
WHO BMI-for-age z score	0.301	0.286 [0.252] N=1,231	0.299 [0.264] N=1,142	0.284 [0.290] N=1,020	0.240 [0.308] N=871	0.301 [0.334] N=723
Stunted: HAZ below -2	0.245 P=296	-0.016 [0.048] N=1,257	-0.019 [0.050] N=1,164	-0.019 [0.053] N=1,039	-0.033 [0.058] N=892	-0.009 [0.061] N=741
Underweight: WAZ below -2	0.083 P=120	**0.071*** [0.026] N=1,451**	**0.057** [0.027] N=1,342**	**0.051* [0.029] N=1,200**	0.046 [0.031] N=1,024	0.042 [0.034] N=846
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=152	-0.032 [0.037] N=1,255	-0.021 [0.038] N=1,163	-0.031 [0.042] N=1,042	-0.018 [0.045] N=891	-0.031 [0.047] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=244	0.052 [0.047] N=1,255	0.058 [0.050] N=1,163	0.048 [0.053] N=1,042	0.065 [0.059] N=891	0.061 [0.065] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=112	0.012 [0.035] N=1,255	0.021 [0.037] N=1,163	0.031 [0.038] N=1,042	0.042 [0.042] N=891	0.058 [0.046] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=393	-0.031 [0.050] N=1,202	-0.022 [0.052] N=1,113	-0.024 [0.056] N=996	-0.032 [0.062] N=853	-0.018 [0.069] N=708
BMI-for-age z score below -2	0.131 P=160	-0.021 [0.041] N=1,231	-0.015 [0.043] N=1,142	-0.012 [0.048] N=1,020	0.021 [0.052] N=871	0.020 [0.055] N=723

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.011 [0.370] N=1,056	0.076 [0.399] N=963	0.236 [0.467] N=838	0.480 [0.512] N=691	0.228 [0.674] N=540
WHO weight-for-age z score	**-0.382** [0.193] N=1,225**	-0.359 [0.233] N=1,116	-0.340 [0.244] N=974	-0.379 [0.281] N=798	**-0.710* [0.397] N=620**
WHO weight-for-length/height z score, 0-60 months	0.148 [0.331] N=1,054	0.103 [0.360] N=962	0.020 [0.396] N=841	0.118 [0.446] N=690	0.050 [0.529] N=537
WHO BMI-for-age z score	0.198 [0.332] N=1,032	0.163 [0.354] N=943	0.050 [0.407] N=821	-0.142 [0.444] N=672	-0.337 [0.510] N=524
Stunted: HAZ below -2	-0.011 [0.069] N=1,056	-0.024 [0.076] N=963	-0.036 [0.084] N=838	-0.082 [0.101] N=691	-0.071 [0.135] N=540
Underweight: WAZ below -2	**0.112*** [0.036] N=1,225**	**0.089** [0.040] N=1,116**	**0.086** [0.043] N=974**	**0.099* [0.050] N=798**	0.108 [0.069] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	-0.042 [0.055] N=1,054	-0.019 [0.060] N=962	-0.038 [0.070] N=841	-0.025 [0.085] N=690	-0.051 [0.098] N=537
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.058] N=1,054	0.014 [0.064] N=962	-0.021 [0.065] N=841	0.002 [0.084] N=690	-0.078 [0.100] N=537
Obese: WLZ/WHZ above +3, 0-60 months	-0.020 [0.043] N=1,054	-0.011 [0.049] N=962	-0.002 [0.050] N=841	0.022 [0.058] N=690	0.035 [0.075] N=537
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.010 [0.077] N=1,007	0.004 [0.084] N=918	-0.004 [0.093] N=801	-0.032 [0.118] N=658	-0.008 [0.160] N=513
BMI-for-age z score below -2	-0.044 [0.056] N=1,032	-0.031 [0.061] N=943	-0.028 [0.072] N=821	0.034 [0.084] N=672	0.061 [0.099] N=524

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	-0.002 [0.252] N=1,257	0.017 [0.263] N=1,164	0.038 [0.289] N=1,039	0.101 [0.306] N=892	-0.124 [0.335] N=741
WHO weight-for-age z score	0.010	-0.169 [0.129] N=1,451	-0.142 [0.144] N=1,342	-0.116 [0.148] N=1,200	-0.067 [0.147] N=1,024	-0.140 [0.161] N=846
WHO weight-for-length/height z score, 0-60 months	0.312	0.310 [0.230] N=1,255	0.315 [0.243] N=1,163	0.327 [0.259] N=1,042	0.382 [0.263] N=891	0.457 [0.290] N=738
WHO BMI-for-age z score	0.301	0.289 [0.255] N=1,231	0.304 [0.269] N=1,142	0.299 [0.295] N=1,020	0.252 [0.312] N=871	0.306 [0.346] N=723
Stunted: HAZ below -2	0.245 P=296	-0.013 [0.048] N=1,257	-0.012 [0.051] N=1,164	-0.011 [0.053] N=1,039	-0.030 [0.057] N=892	0.003 [0.061] N=741
Underweight: WAZ below -2	0.083 P=120	**0.074*** [0.026] N=1,451**	**0.061** [0.027] N=1,342**	**0.056* [0.029] N=1,200**	0.049 [0.031] N=1,024	0.049 [0.034] N=846
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=152	-0.030 [0.037] N=1,255	-0.018 [0.039] N=1,163	-0.030 [0.042] N=1,042	-0.018 [0.045] N=891	-0.028 [0.047] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=244	0.053 [0.047] N=1,255	0.061 [0.050] N=1,163	0.051 [0.053] N=1,042	0.066 [0.058] N=891	0.063 [0.066] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=112	0.012 [0.035] N=1,255	0.022 [0.038] N=1,163	0.032 [0.039] N=1,042	0.043 [0.042] N=891	0.057 [0.046] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=393	-0.024 [0.050] N=1,202	-0.008 [0.053] N=1,113	-0.012 [0.056] N=996	-0.025 [0.061] N=853	0.003 [0.069] N=708
BMI-for-age z score below -2	0.131 P=160	-0.021 [0.042] N=1,231	-0.013 [0.044] N=1,142	-0.011 [0.049] N=1,020	0.021 [0.052] N=871	0.020 [0.056] N=723

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.006 [0.370] N=1,056	0.054 [0.396] N=963	0.182 [0.460] N=838	0.473 [0.496] N=691	0.245 [0.653] N=540
WHO weight-for-age z score	**-0.389** [0.190] N=1,225**	-0.376 [0.230] N=1,116	-0.360 [0.239] N=974	-0.360 [0.283] N=798	**-0.707* [0.393] N=620**
WHO weight-for-length/height z score, 0-60 months	0.152 [0.330] N=1,054	0.108 [0.362] N=962	0.050 [0.394] N=841	0.111 [0.437] N=690	0.015 [0.529] N=537
WHO BMI-for-age z score	0.198 [0.333] N=1,032	0.165 [0.357] N=943	0.071 [0.407] N=821	-0.171 [0.433] N=672	-0.348 [0.511] N=524
Stunted: HAZ below -2	-0.013 [0.068] N=1,056	-0.020 [0.075] N=963	-0.032 [0.083] N=838	-0.095 [0.096] N=691	-0.080 [0.127] N=540
Underweight: WAZ below -2	**0.110*** [0.036] N=1,225**	**0.090** [0.041] N=1,116**	**0.087** [0.043] N=974**	**0.091* [0.051] N=798**	0.107 [0.069] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	-0.043 [0.054] N=1,054	-0.019 [0.061] N=962	-0.042 [0.070] N=841	-0.030 [0.084] N=690	-0.048 [0.099] N=537
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.059] N=1,054	0.014 [0.065] N=962	-0.020 [0.066] N=841	-0.006 [0.085] N=690	-0.078 [0.103] N=537
Obese: WLZ/WHZ above +3, 0-60 months	-0.021 [0.042] N=1,054	-0.012 [0.050] N=962	-0.003 [0.050] N=841	0.019 [0.059] N=690	0.033 [0.075] N=537
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.015 [0.077] N=1,007	0.007 [0.085] N=918	-0.008 [0.092] N=801	-0.057 [0.116] N=658	-0.013 [0.154] N=513
BMI-for-age z score below -2	-0.045 [0.056] N=1,032	-0.028 [0.061] N=943	-0.030 [0.073] N=821	0.033 [0.083] N=672	0.062 [0.100] N=524

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	-0.022 [0.252] N=1,257	0.015 [0.262] N=1,164	0.039 [0.290] N=1,039	0.114 [0.307] N=892	-0.107 [0.337] N=741
WHO weight-for-age z score	0.010	-0.185 [0.130] N=1,451	-0.154 [0.145] N=1,342	-0.129 [0.151] N=1,200	-0.100 [0.149] N=1,024	-0.192 [0.167] N=846
WHO weight-for-length/height z score, 0-60 months	0.312	0.306 [0.231] N=1,255	0.303 [0.243] N=1,163	0.305 [0.259] N=1,042	0.343 [0.267] N=891	0.401 [0.291] N=738
WHO BMI-for-age z score	0.301	0.287 [0.254] N=1,231	0.302 [0.266] N=1,142	0.293 [0.293] N=1,020	0.222 [0.312] N=871	0.250 [0.344] N=723
Stunted: HAZ below -2	0.245 P=296	-0.007 [0.049] N=1,257	-0.008 [0.051] N=1,164	-0.003 [0.054] N=1,039	-0.021 [0.058] N=892	0.012 [0.062] N=741
Underweight: WAZ below -2	0.083 P=120	**0.078*** [0.026] N=1,451**	**0.065** [0.027] N=1,342**	**0.061** [0.029] N=1,200**	**0.058* [0.031] N=1,024**	**0.059* [0.034] N=846**
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=152	-0.028 [0.037] N=1,255	-0.015 [0.039] N=1,163	-0.024 [0.042] N=1,042	-0.012 [0.045] N=891	-0.024 [0.047] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=244	0.054 [0.048] N=1,255	0.063 [0.051] N=1,163	0.054 [0.053] N=1,042	0.067 [0.059] N=891	0.060 [0.067] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=112	0.011 [0.035] N=1,255	0.019 [0.037] N=1,163	0.029 [0.038] N=1,042	0.038 [0.042] N=891	0.048 [0.046] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=393	-0.013 [0.051] N=1,202	0.003 [0.054] N=1,113	0.008 [0.058] N=996	-0.002 [0.064] N=853	0.024 [0.072] N=708
BMI-for-age z score below -2	0.131 P=160	-0.018 [0.042] N=1,231	-0.010 [0.043] N=1,142	-0.005 [0.049] N=1,020	0.028 [0.052] N=871	0.028 [0.056] N=723

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.031 [0.377] N=1,056	0.037 [0.400] N=963	0.156 [0.469] N=838	0.465 [0.512] N=691	0.196 [0.660] N=540
WHO weight-for-age z score	**-0.411** [0.194] N=1,225**	-0.382 [0.232] N=1,116	-0.375 [0.242] N=974	-0.341 [0.287] N=798	**-0.669* [0.391] N=620**
WHO weight-for-length/height z score, 0-60 months	0.173 [0.329] N=1,054	0.133 [0.357] N=962	0.058 [0.399] N=841	0.178 [0.446] N=690	0.107 [0.532] N=537
WHO BMI-for-age z score	0.208 [0.329] N=1,032	0.185 [0.349] N=943	0.086 [0.409] N=821	-0.086 [0.445] N=672	-0.240 [0.507] N=524
Stunted: HAZ below -2	-0.006 [0.070] N=1,056	-0.018 [0.076] N=963	-0.026 [0.085] N=838	-0.094 [0.097] N=691	-0.071 [0.129] N=540
Underweight: WAZ below -2	**0.112*** [0.037] N=1,225**	**0.089** [0.041] N=1,116**	**0.088** [0.043] N=974**	**0.087* [0.051] N=798**	0.110 [0.069] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	-0.044 [0.055] N=1,054	-0.022 [0.061] N=962	-0.042 [0.070] N=841	-0.031 [0.085] N=690	-0.050 [0.101] N=537
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.058] N=1,054	0.016 [0.064] N=962	-0.018 [0.066] N=841	0.006 [0.084] N=690	-0.054 [0.098] N=537
Obese: WLZ/WHZ above +3, 0-60 months	-0.015 [0.043] N=1,054	-0.006 [0.049] N=962	0.004 [0.050] N=841	0.028 [0.058] N=690	0.042 [0.073] N=537
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.010 [0.079] N=1,007	0.006 [0.086] N=918	-0.001 [0.094] N=801	-0.055 [0.118] N=658	-0.005 [0.162] N=513
BMI-for-age z score below -2	-0.046 [0.056] N=1,032	-0.035 [0.060] N=943	-0.034 [0.073] N=821	0.023 [0.085] N=672	0.045 [0.099] N=524

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.352	-0.021 [0.246] N=1,155	-0.020 [0.264] N=1,075	0.010 [0.292] N=957	-0.006 [0.303] N=828	-0.242 [0.327] N=690
WHO weight-for-age z score	0.016	-0.051 [0.135] N=1,329	-0.031 [0.147] N=1,234	0.010 [0.152] N=1,101	0.038 [0.150] N=945	0.008 [0.167] N=786
WHO weight-for-length/height z score, 0-60 months	0.327	**0.485** [0.224] N=1,156**	**0.479** [0.233] N=1,077**	**0.477* [0.248] N=964**	**0.547** [0.247] N=831**	**0.668** [0.266] N=691**
WHO BMI-for-age z score	0.324	**0.441* [0.250] N=1,133**	0.429 [0.260] N=1,056	0.431 [0.283] N=943	0.419 [0.292] N=812	0.527 [0.319] N=677
Stunted: HAZ below -2	0.247 P=274	-0.007 [0.050] N=1,155	-0.010 [0.053] N=1,075	-0.009 [0.056] N=957	-0.016 [0.060] N=828	0.019 [0.064] N=690
Underweight: WAZ below -2	0.083 P=110	**0.066** [0.028] N=1,329**	**0.053* [0.030] N=1,234**	0.046 [0.031] N=1,101	0.042 [0.034] N=945	0.036 [0.038] N=786
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=139	-0.058 [0.036] N=1,156	-0.043 [0.037] N=1,077	-0.052 [0.040] N=964	-0.040 [0.042] N=831	-0.054 [0.044] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=226	0.072 [0.048] N=1,156	0.079 [0.050] N=1,077	0.071 [0.054] N=964	0.085 [0.059] N=831	0.091 [0.066] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.014 [0.036] N=1,156	0.026 [0.039] N=1,077	0.039 [0.041] N=964	0.054 [0.044] N=831	0.073 [0.047] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.338 P=363	-0.043 [0.053] N=1,105	-0.028 [0.056] N=1,029	-0.034 [0.060] N=919	-0.042 [0.066] N=794	-0.027 [0.073] N=661
BMI-for-age z score below -2	0.129 P=143	-0.032 [0.041] N=1,133	-0.021 [0.043] N=1,056	-0.022 [0.048] N=943	0.007 [0.051] N=812	-0.004 [0.056] N=677

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.151 [0.374] N=967	0.198 [0.420] N=887	0.401 [0.496] N=769	0.568 [0.541] N=640	0.353 [0.728] N=502
WHO weight-for-age z score	**-0.319* [0.190] N=1,119**	-0.324 [0.224] N=1,024	-0.301 [0.230] N=891	-0.351 [0.260] N=735	-0.632 [0.385] N=576
WHO weight-for-length/height z score, 0-60 months	0.211 [0.345] N=968	0.127 [0.370] N=889	0.001 [0.409] N=776	0.144 [0.450] N=643	0.160 [0.547] N=503
WHO BMI-for-age z score	0.208 [0.352] N=947	0.099 [0.369] N=870	-0.030 [0.414] N=757	-0.146 [0.438] N=626	-0.315 [0.526] N=491
Stunted: HAZ below -2	-0.040 [0.072] N=967	-0.060 [0.079] N=887	-0.081 [0.088] N=769	-0.130 [0.103] N=640	-0.119 [0.142] N=502
Underweight: WAZ below -2	**0.108*** [0.038] N=1,119**	**0.089** [0.042] N=1,024**	**0.087* [0.044] N=891**	**0.103** [0.051] N=735**	0.113 [0.075] N=576
Wasted: WLZ/WHZ below -2, 0-60 months	-0.075 [0.056] N=968	-0.044 [0.061] N=889	-0.061 [0.069] N=776	-0.062 [0.083] N=643	-0.119 [0.096] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.059] N=968	0.013 [0.065] N=889	-0.024 [0.068] N=776	-0.008 [0.085] N=643	-0.073 [0.102] N=503
Obese: WLZ/WHZ above +3, 0-60 months	-0.034 [0.046] N=968	-0.020 [0.054] N=889	-0.006 [0.055] N=776	0.027 [0.063] N=643	0.047 [0.079] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.058 [0.079] N=923	-0.041 [0.087] N=847	-0.064 [0.096] N=737	-0.120 [0.118] N=612	-0.125 [0.160] N=479
BMI-for-age z score below -2	-0.040 [0.057] N=947	-0.015 [0.060] N=870	-0.015 [0.072] N=757	0.036 [0.084] N=626	0.037 [0.099] N=491

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.352	-0.012 [0.246] N=1,155	-0.012 [0.263] N=1,075	0.011 [0.291] N=957	-0.011 [0.302] N=828	-0.261 [0.324] N=690
WHO weight-for-age z score	0.016	-0.035 [0.135] N=1,329	-0.018 [0.148] N=1,234	0.016 [0.152] N=1,101	0.031 [0.152] N=945	0.004 [0.172] N=786
WHO weight-for-length/height z score, 0-60 months	0.327	**0.483** [0.221] N=1,156**	**0.476** [0.230] N=1,077**	**0.471* [0.245] N=964**	**0.536** [0.244] N=831**	**0.655** [0.262] N=691**
WHO BMI-for-age z score	0.324	**0.442* [0.247] N=1,133**	**0.426* [0.256] N=1,056**	0.427 [0.280] N=943	0.410 [0.288] N=812	0.509 [0.313] N=677
Stunted: HAZ below -2	0.247 P=274	-0.007 [0.049] N=1,155	-0.009 [0.053] N=1,075	-0.009 [0.056] N=957	-0.017 [0.060] N=828	0.019 [0.063] N=690
Underweight: WAZ below -2	0.083 P=110	**0.063** [0.027] N=1,329**	**0.051* [0.029] N=1,234**	0.045 [0.030] N=1,101	0.043 [0.033] N=945	0.034 [0.037] N=786
Wasted: WLZ/WHZ below -2, 0-60 months	0.120 P=139	-0.057 [0.036] N=1,156	-0.042 [0.037] N=1,077	-0.052 [0.040] N=964	-0.040 [0.042] N=831	-0.056 [0.044] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=226	0.073 [0.047] N=1,156	0.079 [0.050] N=1,077	0.069 [0.053] N=964	0.081 [0.058] N=831	0.086 [0.065] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.014 [0.036] N=1,156	0.026 [0.039] N=1,077	0.039 [0.040] N=964	0.053 [0.043] N=831	0.071 [0.047] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.338 P=363	-0.040 [0.053] N=1,105	-0.025 [0.056] N=1,029	-0.033 [0.060] N=919	-0.041 [0.066] N=794	-0.024 [0.073] N=661
BMI-for-age z score below -2	0.129 P=143	-0.030 [0.041] N=1,133	-0.020 [0.043] N=1,056	-0.022 [0.048] N=943	0.008 [0.051] N=812	-0.002 [0.055] N=677

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.166 [0.376] N=967	0.208 [0.423] N=887	0.402 [0.499] N=769	0.558 [0.546] N=640	0.336 [0.728] N=502
WHO weight-for-age z score	-0.283 [0.188] N=1,119	-0.291 [0.224] N=1,024	-0.278 [0.229] N=891	-0.354 [0.261] N=735	-0.617 [0.397] N=576
WHO weight-for-length/height z score, 0-60 months	0.225 [0.341] N=968	0.146 [0.369] N=889	0.015 [0.409] N=776	0.134 [0.448] N=643	0.125 [0.542] N=503
WHO BMI-for-age z score	0.223 [0.349] N=947	0.106 [0.367] N=870	-0.021 [0.414] N=757	-0.154 [0.433] N=626	-0.352 [0.514] N=491
Stunted: HAZ below -2	-0.037 [0.072] N=967	-0.055 [0.079] N=887	-0.078 [0.088] N=769	-0.127 [0.104] N=640	-0.124 [0.141] N=502
Underweight: WAZ below -2	**0.103*** [0.037] N=1,119**	**0.082** [0.041] N=1,024**	**0.082* [0.043] N=891**	**0.103** [0.051] N=735**	0.102 [0.074] N=576
Wasted: WLZ/WHZ below -2, 0-60 months	-0.072 [0.056] N=968	-0.042 [0.060] N=889	-0.060 [0.069] N=776	-0.061 [0.083] N=643	-0.117 [0.094] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	0.019 [0.059] N=968	0.019 [0.064] N=889	-0.020 [0.068] N=776	-0.008 [0.084] N=643	-0.072 [0.101] N=503
Obese: WLZ/WHZ above +3, 0-60 months	-0.032 [0.046] N=968	-0.019 [0.053] N=889	-0.004 [0.055] N=776	0.027 [0.062] N=643	0.046 [0.078] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.050 [0.079] N=923	-0.034 [0.087] N=847	-0.061 [0.096] N=737	-0.118 [0.118] N=612	-0.132 [0.161] N=479
BMI-for-age z score below -2	-0.036 [0.056] N=947	-0.013 [0.060] N=870	-0.014 [0.072] N=757	0.036 [0.084] N=626	0.038 [0.099] N=491

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.108 [0.316] N=1,088	-0.213 [0.321] N=978	-0.003 [0.342] N=849	0.004 [0.373] N=718	-0.026 [0.416] N=587
WHO weight-for-age z score	0.018	0.016 [0.165] N=1,250	0.034 [0.176] N=1,131	-0.043 [0.184] N=981	0.004 [0.201] N=827	0.062 [0.220] N=670
WHO weight-for-length/height z score, 0-60 months	0.338	0.098 [0.267] N=1,089	0.090 [0.276] N=983	-0.009 [0.298] N=854	0.116 [0.327] N=721	0.012 [0.371] N=586
WHO BMI-for-age z score	0.312	0.137 [0.261] N=1,067	0.195 [0.267] N=961	0.106 [0.300] N=834	0.318 [0.331] N=705	0.261 [0.370] N=576
Stunted: HAZ below -2	0.228 P=244	0.039 [0.058] N=1,088	0.042 [0.059] N=978	0.002 [0.062] N=849	-0.006 [0.070] N=718	-0.051 [0.077] N=587
Underweight: WAZ below -2	0.081 P=101	-0.009 [0.036] N=1,250	-0.005 [0.038] N=1,131	0.007 [0.039] N=981	0.021 [0.042] N=827	0.018 [0.045] N=670
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=130	0.002 [0.043] N=1,089	0.011 [0.045] N=983	0.034 [0.050] N=854	-0.008 [0.049] N=721	0.008 [0.056] N=586
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=216	0.043 [0.049] N=1,089	0.046 [0.051] N=983	0.016 [0.052] N=854	0.016 [0.055] N=721	0.021 [0.062] N=586
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=102	0.034 [0.041] N=1,089	0.027 [0.041] N=983	0.016 [0.043] N=854	0.010 [0.048] N=721	-0.024 [0.047] N=586
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=332	0.055 [0.054] N=1,041	0.055 [0.056] N=938	0.047 [0.059] N=813	0.006 [0.063] N=687	-0.011 [0.068] N=560
BMI-for-age z score below -2	0.130 P=138	0.018 [0.047] N=1,067	0.027 [0.049] N=961	0.020 [0.057] N=834	-0.014 [0.057] N=705	0.007 [0.065] N=576

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.279 [0.458] N=944	-0.433 [0.480] N=834	-0.115 [0.543] N=705	-0.195 [0.613] N=574	-0.277 [0.756] N=443
WHO weight-for-age z score	-0.023 [0.232] N=1,086	-0.004 [0.255] N=967	-0.181 [0.278] N=817	-0.153 [0.317] N=663	-0.108 [0.370] N=506
WHO weight-for-length/height z score, 0-60 months	-0.088 [0.386] N=945	-0.134 [0.411] N=839	-0.363 [0.482] N=710	-0.218 [0.539] N=577	-0.782 [0.645] N=442
WHO BMI-for-age z score	-0.222 [0.373] N=929	-0.154 [0.392] N=823	-0.353 [0.484] N=696	-0.014 [0.546] N=567	-0.376 [0.582] N=438
Stunted: HAZ below -2	0.107 [0.078] N=944	0.109 [0.082] N=834	0.055 [0.086] N=705	0.064 [0.107] N=574	-0.052 [0.123] N=443
Underweight: WAZ below -2	-0.010 [0.060] N=1,086	-0.004 [0.066] N=967	0.024 [0.073] N=817	0.060 [0.086] N=663	0.067 [0.103] N=506
Wasted: WLZ/WHZ below -2, 0-60 months	0.043 [0.060] N=945	0.064 [0.065] N=839	0.113 [0.078] N=710	0.039 [0.076] N=577	0.105 [0.092] N=442
Overweight: WLZ/WHZ above +2, 0-60 months	0.030 [0.078] N=945	0.023 [0.082] N=839	-0.038 [0.090] N=710	-0.048 [0.103] N=577	-0.121 [0.126] N=442
Obese: WLZ/WHZ above +3, 0-60 months	0.058 [0.058] N=945	0.041 [0.061] N=839	0.024 [0.069] N=710	-0.000 [0.087] N=577	-0.159 [0.101] N=442
CIAF: any stunting, underweight, or wasting, 0-60 months	0.120 [0.082] N=901	0.124 [0.089] N=798	0.123 [0.096] N=673	0.065 [0.108] N=547	0.017 [0.131] N=420
BMI-for-age z score below -2	0.060 [0.055] N=929	0.074 [0.059] N=823	0.068 [0.075] N=696	-0.020 [0.068] N=567	0.017 [0.084] N=438

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.110 [0.316] N=1,088	-0.218 [0.321] N=978	-0.015 [0.340] N=849	-0.011 [0.371] N=718	-0.038 [0.414] N=587
WHO weight-for-age z score	0.018	0.023 [0.165] N=1,250	0.039 [0.175] N=1,131	-0.038 [0.183] N=981	0.014 [0.200] N=827	0.068 [0.220] N=670
WHO weight-for-length/height z score, 0-60 months	0.338	0.099 [0.267] N=1,089	0.087 [0.276] N=983	-0.012 [0.298] N=854	0.118 [0.326] N=721	0.020 [0.368] N=586
WHO BMI-for-age z score	0.312	0.129 [0.261] N=1,067	0.185 [0.267] N=961	0.096 [0.300] N=834	0.311 [0.331] N=705	0.258 [0.371] N=576
Stunted: HAZ below -2	0.228 P=244	0.035 [0.058] N=1,088	0.038 [0.059] N=978	-0.000 [0.062] N=849	-0.007 [0.070] N=718	-0.051 [0.077] N=587
Underweight: WAZ below -2	0.081 P=101	-0.010 [0.036] N=1,250	-0.007 [0.039] N=1,131	0.004 [0.039] N=981	0.018 [0.042] N=827	0.017 [0.045] N=670
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=130	-0.001 [0.043] N=1,089	0.007 [0.045] N=983	0.031 [0.050] N=854	-0.011 [0.048] N=721	0.004 [0.055] N=586
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=216	0.041 [0.050] N=1,089	0.042 [0.051] N=983	0.012 [0.052] N=854	0.012 [0.056] N=721	0.019 [0.063] N=586
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=102	0.032 [0.041] N=1,089	0.024 [0.041] N=983	0.013 [0.043] N=854	0.008 [0.048] N=721	-0.027 [0.047] N=586
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=332	0.047 [0.055] N=1,041	0.046 [0.057] N=938	0.039 [0.061] N=813	0.001 [0.064] N=687	-0.016 [0.070] N=560
BMI-for-age z score below -2	0.130 P=138	0.017 [0.047] N=1,067	0.025 [0.049] N=961	0.019 [0.057] N=834	-0.015 [0.057] N=705	0.007 [0.065] N=576

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.268 [0.461] N=944	-0.431 [0.483] N=834	-0.129 [0.543] N=705	-0.232 [0.618] N=574	-0.299 [0.768] N=443
WHO weight-for-age z score	-0.009 [0.230] N=1,086	0.008 [0.253] N=967	-0.167 [0.275] N=817	-0.113 [0.312] N=663	-0.068 [0.372] N=506
WHO weight-for-length/height z score, 0-60 months	-0.089 [0.389] N=945	-0.143 [0.415] N=839	-0.376 [0.487] N=710	-0.206 [0.549] N=577	-0.736 [0.670] N=442
WHO BMI-for-age z score	-0.232 [0.376] N=929	-0.164 [0.395] N=823	-0.370 [0.485] N=696	-0.009 [0.553] N=567	-0.348 [0.608] N=438
Stunted: HAZ below -2	0.097 [0.078] N=944	0.099 [0.082] N=834	0.047 [0.086] N=705	0.062 [0.108] N=574	-0.053 [0.125] N=443
Underweight: WAZ below -2	-0.013 [0.060] N=1,086	-0.007 [0.066] N=967	0.019 [0.073] N=817	0.052 [0.086] N=663	0.061 [0.104] N=506
Wasted: WLZ/WHZ below -2, 0-60 months	0.036 [0.062] N=945	0.055 [0.067] N=839	0.104 [0.080] N=710	0.023 [0.077] N=577	0.080 [0.095] N=442
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.078] N=945	0.015 [0.082] N=839	-0.049 [0.090] N=710	-0.064 [0.104] N=577	-0.135 [0.127] N=442
Obese: WLZ/WHZ above +3, 0-60 months	0.054 [0.058] N=945	0.035 [0.061] N=839	0.016 [0.070] N=710	-0.010 [0.088] N=577	**-0.174* [0.103] N=442**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.099 [0.082] N=901	0.098 [0.089] N=798	0.096 [0.097] N=673	0.034 [0.109] N=547	-0.024 [0.128] N=420
BMI-for-age z score below -2	0.055 [0.056] N=929	0.067 [0.061] N=823	0.061 [0.076] N=696	-0.032 [0.071] N=567	0.004 [0.090] N=438

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.119 [0.309] N=1,088	-0.207 [0.314] N=978	-0.003 [0.337] N=849	-0.010 [0.369] N=718	-0.045 [0.413] N=587
WHO weight-for-age z score	0.018	0.016 [0.162] N=1,250	0.034 [0.172] N=1,131	-0.025 [0.181] N=981	0.020 [0.199] N=827	0.071 [0.219] N=670
WHO weight-for-length/height z score, 0-60 months	0.338	0.102 [0.266] N=1,089	0.084 [0.276] N=983	0.005 [0.296] N=854	0.133 [0.324] N=721	0.026 [0.368] N=586
WHO BMI-for-age z score	0.312	0.127 [0.259] N=1,067	0.176 [0.266] N=961	0.101 [0.295] N=834	0.319 [0.322] N=705	0.249 [0.365] N=576
Stunted: HAZ below -2	0.228 P=244	0.039 [0.057] N=1,088	0.038 [0.058] N=978	-0.003 [0.061] N=849	-0.008 [0.069] N=718	-0.052 [0.075] N=587
Underweight: WAZ below -2	0.081 P=101	-0.007 [0.035] N=1,250	-0.003 [0.037] N=1,131	0.005 [0.038] N=981	0.019 [0.040] N=827	0.019 [0.043] N=670
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=130	-0.000 [0.044] N=1,089	0.007 [0.046] N=983	0.028 [0.051] N=854	-0.013 [0.050] N=721	0.004 [0.057] N=586
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=216	0.042 [0.050] N=1,089	0.043 [0.051] N=983	0.011 [0.052] N=854	0.010 [0.056] N=721	0.021 [0.062] N=586
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=102	0.033 [0.041] N=1,089	0.025 [0.042] N=983	0.013 [0.044] N=854	0.008 [0.048] N=721	-0.027 [0.048] N=586
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=332	0.055 [0.056] N=1,041	0.050 [0.058] N=938	0.034 [0.062] N=813	-0.004 [0.065] N=687	-0.021 [0.069] N=560
BMI-for-age z score below -2	0.130 P=138	0.020 [0.047] N=1,067	0.028 [0.050] N=961	0.020 [0.058] N=834	-0.015 [0.058] N=705	0.011 [0.066] N=576

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.323 [0.461] N=944	-0.456 [0.480] N=834	-0.160 [0.537] N=705	-0.290 [0.610] N=574	-0.394 [0.777] N=443
WHO weight-for-age z score	-0.060 [0.230] N=1,086	-0.042 [0.252] N=967	-0.168 [0.277] N=817	-0.134 [0.318] N=663	-0.122 [0.383] N=506
WHO weight-for-length/height z score, 0-60 months	-0.107 [0.389] N=945	-0.164 [0.416] N=839	-0.353 [0.482] N=710	-0.185 [0.541] N=577	-0.732 [0.673] N=442
WHO BMI-for-age z score	-0.235 [0.377] N=929	-0.180 [0.398] N=823	-0.350 [0.479] N=696	0.011 [0.542] N=567	-0.324 [0.614] N=438
Stunted: HAZ below -2	0.109 [0.078] N=944	0.105 [0.081] N=834	0.054 [0.084] N=705	0.076 [0.107] N=574	-0.027 [0.125] N=443
Underweight: WAZ below -2	-0.006 [0.059] N=1,086	0.001 [0.064] N=967	0.020 [0.071] N=817	0.055 [0.084] N=663	0.070 [0.101] N=506
Wasted: WLZ/WHZ below -2, 0-60 months	0.043 [0.063] N=945	0.064 [0.068] N=839	0.104 [0.081] N=710	0.025 [0.078] N=577	0.092 [0.098] N=442
Overweight: WLZ/WHZ above +2, 0-60 months	0.028 [0.078] N=945	0.018 [0.083] N=839	-0.046 [0.089] N=710	-0.060 [0.104] N=577	-0.129 [0.126] N=442
Obese: WLZ/WHZ above +3, 0-60 months	0.056 [0.057] N=945	0.038 [0.061] N=839	0.019 [0.070] N=710	-0.006 [0.087] N=577	-0.168 [0.101] N=442
CIAF: any stunting, underweight, or wasting, 0-60 months	0.122 [0.083] N=901	0.118 [0.090] N=798	0.101 [0.099] N=673	0.045 [0.112] N=547	0.006 [0.128] N=420
BMI-for-age z score below -2	0.058 [0.057] N=929	0.072 [0.062] N=823	0.061 [0.076] N=696	-0.032 [0.071] N=567	0.006 [0.092] N=438

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.134 [0.308] N=1,088	-0.216 [0.314] N=978	0.013 [0.333] N=849	-0.002 [0.361] N=718	-0.067 [0.404] N=587
WHO weight-for-age z score	0.018	0.003 [0.162] N=1,250	0.023 [0.173] N=1,131	-0.031 [0.183] N=981	0.014 [0.200] N=827	0.059 [0.223] N=670
WHO weight-for-length/height z score, 0-60 months	0.338	0.118 [0.263] N=1,089	0.102 [0.272] N=983	0.015 [0.289] N=854	0.162 [0.318] N=721	0.080 [0.364] N=586
WHO BMI-for-age z score	0.312	0.142 [0.259] N=1,067	0.194 [0.266] N=961	0.114 [0.293] N=834	0.345 [0.323] N=705	0.315 [0.368] N=576
Stunted: HAZ below -2	0.228 P=244	0.040 [0.057] N=1,088	0.038 [0.058] N=978	-0.005 [0.061] N=849	-0.009 [0.069] N=718	-0.048 [0.076] N=587
Underweight: WAZ below -2	0.081 P=101	-0.010 [0.035] N=1,250	-0.006 [0.038] N=1,131	0.002 [0.039] N=981	0.019 [0.042] N=827	0.018 [0.046] N=670
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=130	-0.002 [0.043] N=1,089	0.007 [0.046] N=983	0.030 [0.050] N=854	-0.014 [0.047] N=721	-0.001 [0.053] N=586
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=216	0.043 [0.049] N=1,089	0.043 [0.050] N=983	0.015 [0.051] N=854	0.018 [0.055] N=721	0.027 [0.062] N=586
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=102	0.038 [0.039] N=1,089	0.030 [0.040] N=983	0.019 [0.042] N=854	0.016 [0.046] N=721	-0.018 [0.046] N=586
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=332	0.053 [0.056] N=1,041	0.051 [0.058] N=938	0.036 [0.061] N=813	0.000 [0.064] N=687	-0.008 [0.070] N=560
BMI-for-age z score below -2	0.130 P=138	0.014 [0.047] N=1,067	0.022 [0.050] N=961	0.016 [0.056] N=834	-0.020 [0.056] N=705	-0.004 [0.063] N=576

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.333 [0.446] N=944	-0.462 [0.468] N=834	-0.068 [0.524] N=705	-0.195 [0.601] N=574	-0.326 [0.758] N=443
WHO weight-for-age z score	-0.041 [0.227] N=1,086	-0.020 [0.250] N=967	-0.133 [0.281] N=817	-0.093 [0.314] N=663	-0.079 [0.382] N=506
WHO weight-for-length/height z score, 0-60 months	-0.074 [0.386] N=945	-0.132 [0.411] N=839	-0.356 [0.476] N=710	-0.166 [0.535] N=577	-0.685 [0.671] N=442
WHO BMI-for-age z score	-0.219 [0.373] N=929	-0.154 [0.393] N=823	-0.357 [0.479] N=696	0.030 [0.544] N=567	-0.298 [0.611] N=438
Stunted: HAZ below -2	0.110 [0.077] N=944	0.106 [0.080] N=834	0.041 [0.084] N=705	0.060 [0.108] N=574	-0.031 [0.126] N=443
Underweight: WAZ below -2	-0.012 [0.058] N=1,086	-0.006 [0.064] N=967	0.012 [0.072] N=817	0.050 [0.084] N=663	0.060 [0.104] N=506
Wasted: WLZ/WHZ below -2, 0-60 months	0.038 [0.062] N=945	0.059 [0.067] N=839	0.107 [0.078] N=710	0.027 [0.076] N=577	0.091 [0.094] N=442
Overweight: WLZ/WHZ above +2, 0-60 months	0.028 [0.077] N=945	0.016 [0.082] N=839	-0.050 [0.088] N=710	-0.062 [0.104] N=577	-0.126 [0.129] N=442
Obese: WLZ/WHZ above +3, 0-60 months	0.060 [0.056] N=945	0.041 [0.060] N=839	0.020 [0.067] N=710	0.001 [0.085] N=577	-0.159 [0.102] N=442
CIAF: any stunting, underweight, or wasting, 0-60 months	0.111 [0.082] N=901	0.107 [0.089] N=798	0.086 [0.098] N=673	0.028 [0.111] N=547	0.008 [0.131] N=420
BMI-for-age z score below -2	0.057 [0.057] N=929	0.069 [0.061] N=823	0.063 [0.076] N=696	-0.030 [0.070] N=567	-0.001 [0.090] N=438

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.244	-0.166 [0.323] N=996	-0.288 [0.331] N=895	-0.056 [0.346] N=783	-0.073 [0.366] N=660	-0.134 [0.421] N=542
WHO weight-for-age z score	0.017	0.037 [0.170] N=1,142	0.074 [0.180] N=1,032	-0.029 [0.185] N=902	-0.006 [0.198] N=761	0.005 [0.215] N=621
WHO weight-for-length/height z score, 0-60 months	0.345	0.186 [0.284] N=1,000	0.192 [0.297] N=903	0.050 [0.318] N=789	0.128 [0.344] N=665	-0.005 [0.376] N=544
WHO BMI-for-age z score	0.325	0.196 [0.278] N=980	0.278 [0.290] N=883	0.130 [0.323] N=772	0.317 [0.356] N=652	0.214 [0.393] N=536
Stunted: HAZ below -2	0.228 P=223	0.051 [0.062] N=996	0.058 [0.064] N=895	0.015 [0.066] N=783	0.000 [0.071] N=660	-0.040 [0.079] N=542
Underweight: WAZ below -2	0.082 P=93	-0.006 [0.037] N=1,142	-0.004 [0.040] N=1,032	0.005 [0.040] N=902	0.025 [0.043] N=761	0.030 [0.045] N=621
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=119	-0.004 [0.045] N=1,000	0.001 [0.048] N=903	0.027 [0.052] N=789	-0.011 [0.049] N=665	0.012 [0.055] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=200	0.046 [0.052] N=1,000	0.048 [0.053] N=903	0.007 [0.054] N=789	-0.008 [0.057] N=665	-0.012 [0.062] N=544
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=96	0.031 [0.041] N=1,000	0.022 [0.042] N=903	0.008 [0.042] N=789	-0.002 [0.046] N=665	-0.033 [0.045] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	0.324 P=305	0.067 [0.059] N=954	0.062 [0.061] N=860	0.054 [0.062] N=750	0.005 [0.064] N=632	0.007 [0.070] N=518
BMI-for-age z score below -2	0.128 P=123	0.016 [0.049] N=980	0.021 [0.052] N=883	0.017 [0.059] N=772	-0.007 [0.059] N=652	0.026 [0.068] N=536

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.185 [0.499] N=862	-0.364 [0.524] N=761	-0.024 [0.581] N=649	-0.120 [0.647] N=526	-0.054 [0.825] N=408
WHO weight-for-age z score	0.049 [0.238] N=990	0.099 [0.260] N=880	-0.138 [0.289] N=750	-0.086 [0.328] N=609	-0.054 [0.401] N=469
WHO weight-for-length/height z score, 0-60 months	-0.010 [0.421] N=866	-0.030 [0.453] N=769	-0.351 [0.522] N=655	-0.240 [0.601] N=531	-0.988 [0.715] N=410
WHO BMI-for-age z score	-0.173 [0.390] N=850	-0.066 [0.416] N=753	-0.372 [0.502] N=642	-0.030 [0.586] N=522	-0.587 [0.637] N=406
Stunted: HAZ below -2	0.099 [0.089] N=862	0.107 [0.095] N=761	0.049 [0.098] N=649	0.039 [0.119] N=526	-0.099 [0.147] N=408
Underweight: WAZ below -2	-0.016 [0.062] N=990	-0.012 [0.068] N=880	0.008 [0.075] N=750	0.053 [0.092] N=609	0.058 [0.113] N=469
Wasted: WLZ/WHZ below -2, 0-60 months	0.032 [0.065] N=866	0.044 [0.072] N=769	0.089 [0.084] N=655	0.001 [0.081] N=531	0.083 [0.096] N=410
Overweight: WLZ/WHZ above +2, 0-60 months	0.030 [0.085] N=866	0.022 [0.090] N=769	-0.059 [0.098] N=655	-0.111 [0.116] N=531	**-0.225* [0.135] N=410**
Obese: WLZ/WHZ above +3, 0-60 months	0.044 [0.063] N=866	0.023 [0.068] N=769	-0.001 [0.076] N=655	-0.038 [0.096] N=531	**-0.206* [0.114] N=410**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.106 [0.093] N=824	0.099 [0.101] N=730	0.089 [0.108] N=620	-0.021 [0.122] N=502	-0.081 [0.151] N=388
BMI-for-age z score below -2	0.056 [0.060] N=850	0.064 [0.066] N=753	0.056 [0.081] N=642	-0.028 [0.079] N=522	0.026 [0.096] N=406

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.244	-0.188 [0.325] N=996	-0.309 [0.333] N=895	-0.070 [0.347] N=783	-0.091 [0.366] N=660	-0.141 [0.421] N=542
WHO weight-for-age z score	0.017	0.037 [0.167] N=1,142	0.065 [0.179] N=1,032	-0.031 [0.184] N=902	-0.006 [0.196] N=761	0.009 [0.213] N=621
WHO weight-for-length/height z score, 0-60 months	0.345	0.188 [0.286] N=1,000	0.190 [0.298] N=903	0.048 [0.319] N=789	0.125 [0.345] N=665	-0.012 [0.382] N=544
WHO BMI-for-age z score	0.325	0.193 [0.281] N=980	0.276 [0.291] N=883	0.125 [0.324] N=772	0.316 [0.358] N=652	0.212 [0.397] N=536
Stunted: HAZ below -2	0.228 P=223	0.049 [0.061] N=996	0.056 [0.063] N=895	0.012 [0.065] N=783	-0.003 [0.070] N=660	-0.043 [0.078] N=542
Underweight: WAZ below -2	0.082 P=93	-0.010 [0.036] N=1,142	-0.007 [0.039] N=1,032	0.001 [0.039] N=902	0.021 [0.041] N=761	0.027 [0.044] N=621
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=119	-0.008 [0.045] N=1,000	-0.003 [0.048] N=903	0.022 [0.052] N=789	-0.018 [0.049] N=665	0.005 [0.056] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=200	0.046 [0.051] N=1,000	0.047 [0.053] N=903	0.006 [0.053] N=789	-0.010 [0.057] N=665	-0.014 [0.062] N=544
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=96	0.031 [0.041] N=1,000	0.021 [0.041] N=903	0.007 [0.041] N=789	-0.004 [0.045] N=665	-0.037 [0.043] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	0.324 P=305	0.060 [0.059] N=954	0.054 [0.061] N=860	0.046 [0.063] N=750	-0.007 [0.064] N=632	-0.006 [0.070] N=518
BMI-for-age z score below -2	0.128 P=123	0.015 [0.050] N=980	0.019 [0.052] N=883	0.015 [0.059] N=772	-0.012 [0.059] N=652	0.021 [0.066] N=536

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.209 [0.496] N=862	-0.380 [0.521] N=761	-0.024 [0.572] N=649	-0.132 [0.628] N=526	-0.125 [0.796] N=408
WHO weight-for-age z score	0.042 [0.237] N=990	0.084 [0.262] N=880	-0.138 [0.289] N=750	-0.092 [0.327] N=609	-0.062 [0.392] N=469
WHO weight-for-length/height z score, 0-60 months	-0.030 [0.417] N=866	-0.056 [0.449] N=769	-0.378 [0.517] N=655	-0.284 [0.585] N=531	-0.983 [0.686] N=410
WHO BMI-for-age z score	-0.186 [0.388] N=850	-0.078 [0.413] N=753	-0.388 [0.499] N=642	-0.032 [0.577] N=522	-0.553 [0.612] N=406
Stunted: HAZ below -2	0.098 [0.088] N=862	0.104 [0.093] N=761	0.047 [0.096] N=649	0.038 [0.116] N=526	-0.082 [0.141] N=408
Underweight: WAZ below -2	-0.020 [0.062] N=990	-0.016 [0.068] N=880	0.003 [0.075] N=750	0.050 [0.091] N=609	0.068 [0.110] N=469
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.065] N=866	0.047 [0.072] N=769	0.092 [0.085] N=655	0.004 [0.080] N=531	0.085 [0.097] N=410
Overweight: WLZ/WHZ above +2, 0-60 months	0.031 [0.083] N=866	0.020 [0.089] N=769	-0.063 [0.095] N=655	-0.118 [0.112] N=531	-0.215 [0.130] N=410
Obese: WLZ/WHZ above +3, 0-60 months	0.044 [0.062] N=866	0.023 [0.067] N=769	-0.002 [0.074] N=655	-0.040 [0.093] N=531	**-0.204* [0.107] N=410**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.108 [0.092] N=824	0.100 [0.101] N=730	0.092 [0.108] N=620	-0.016 [0.124] N=502	-0.047 [0.149] N=388
BMI-for-age z score below -2	0.060 [0.060] N=850	0.068 [0.066] N=753	0.060 [0.082] N=642	-0.025 [0.077] N=522	0.038 [0.094] N=406

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.