Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,143	-0.000 [0.028] N=1,238	0.012 [0.027] N=1,140	0.002 [0.029] N=1,020	-0.003 [0.031] N=881	-0.012 [0.036] N=729
Vegetables consumed at least weekly	0.897 P=1,107	0.013 [0.032] N=1,238	0.005 [0.031] N=1,140	-0.013 [0.032] N=1,020	-0.014 [0.034] N=881	-0.032 [0.041] N=729
Meat fish or poultry consumed at least weekly	0.760 P=947	0.051 [0.056] N=1,238	0.066 [0.059] N=1,140	0.065 [0.065] N=1,020	0.054 [0.072] N=881	0.055 [0.081] N=729
Legumes consumed at least weekly	0.763 P=949	-0.049 [0.047] N=1,238	-0.072 [0.049] N=1,140	-0.064 [0.051] N=1,020	-0.073 [0.056] N=881	-0.069 [0.068] N=729
Animal products consumed at least weekly	0.900 P=1,118	0.026 [0.038] N=1,238	0.018 [0.040] N=1,140	-0.008 [0.043] N=1,020	-0.001 [0.047] N=881	-0.021 [0.053] N=729
Sweets consumed at least weekly	0.530 P=654	-0.044 [0.046] N=1,238	-0.046 [0.048] N=1,140	-0.034 [0.052] N=1,020	-0.049 [0.055] N=881	-0.004 [0.064] N=729
Snack foods consumed at least weekly	0.439 P=538	-0.021 [0.047] N=1,238	-0.026 [0.048] N=1,140	-0.008 [0.052] N=1,020	-0.003 [0.054] N=881	0.011 [0.057] N=729
Packaged soft drinks consumed at least weekly	0.248 P=299	-0.027 [0.046] N=1,238	-0.044 [0.048] N=1,140	-0.051 [0.050] N=1,020	-0.035 [0.055] N=881	-0.043 [0.063] N=729
Grains consumed at least weekly	0.858 P=1,065	0.032 [0.038] N=1,238	0.036 [0.040] N=1,140	0.018 [0.042] N=1,020	0.025 [0.046] N=881	0.013 [0.053] N=729
Child was ever breastfed	0.970 P=1,447	-0.010 [0.012] N=1,497	-0.008 [0.013] N=1,384	-0.012 [0.015] N=1,236	-0.018 [0.016] N=1,055	-0.019 [0.016] N=875

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.005 [0.043] N=1,044	0.029 [0.044] N=946	0.011 [0.049] N=826	-0.008 [0.059] N=687	-0.034 [0.082] N=535
Vegetables consumed at least weekly	0.024 [0.049] N=1,044	0.010 [0.050] N=946	-0.027 [0.053] N=826	-0.044 [0.063] N=687	-0.126 [0.094] N=535
Meat fish or poultry consumed at least weekly	0.013 [0.067] N=1,044	0.026 [0.069] N=946	0.009 [0.080] N=826	-0.068 [0.087] N=687	-0.124 [0.094] N=535
Legumes consumed at least weekly	-0.036 [0.070] N=1,044	-0.081 [0.074] N=946	-0.060 [0.077] N=826	-0.109 [0.090] N=687	-0.106 [0.135] N=535
Animal products consumed at least weekly	0.044 [0.051] N=1,044	0.032 [0.056] N=946	-0.024 [0.064] N=826	-0.014 [0.077] N=687	-0.079 [0.105] N=535
Sweets consumed at least weekly	-0.064 [0.064] N=1,044	-0.071 [0.067] N=946	-0.056 [0.080] N=826	-0.121 [0.085] N=687	-0.027 [0.123] N=535
Snack foods consumed at least weekly	-0.043 [0.066] N=1,044	-0.058 [0.071] N=946	-0.032 [0.084] N=826	-0.045 [0.099] N=687	-0.045 [0.125] N=535
Packaged soft drinks consumed at least weekly	-0.017 [0.062] N=1,044	-0.047 [0.066] N=946	-0.060 [0.071] N=826	-0.042 [0.086] N=687	-0.075 [0.122] N=535
Grains consumed at least weekly	0.075 [0.060] N=1,044	0.087 [0.065] N=946	0.062 [0.073] N=826	0.095 [0.091] N=687	0.090 [0.126] N=535
Child was ever breastfed	0.000 [0.019] N=1,265	0.006 [0.022] N=1,152	0.001 [0.026] N=1,004	-0.018 [0.033] N=823	-0.007 [0.038] N=643

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,143	-0.001 [0.028] N=1,238	0.011 [0.027] N=1,140	0.002 [0.029] N=1,020	-0.003 [0.031] N=881	-0.013 [0.036] N=729
Vegetables consumed at least weekly	0.897 P=1,107	0.013 [0.032] N=1,238	0.004 [0.031] N=1,140	-0.013 [0.031] N=1,020	-0.015 [0.034] N=881	-0.034 [0.040] N=729
Meat fish or poultry consumed at least weekly	0.760 P=947	0.051 [0.056] N=1,238	0.066 [0.059] N=1,140	0.065 [0.065] N=1,020	0.053 [0.073] N=881	0.053 [0.081] N=729
Legumes consumed at least weekly	0.763 P=949	-0.049 [0.047] N=1,238	-0.072 [0.049] N=1,140	-0.064 [0.051] N=1,020	-0.074 [0.057] N=881	-0.070 [0.069] N=729
Animal products consumed at least weekly	0.900 P=1,118	0.025 [0.038] N=1,238	0.016 [0.040] N=1,140	-0.009 [0.043] N=1,020	-0.002 [0.047] N=881	-0.023 [0.053] N=729
Sweets consumed at least weekly	0.530 P=654	-0.044 [0.046] N=1,238	-0.047 [0.048] N=1,140	-0.034 [0.052] N=1,020	-0.049 [0.056] N=881	-0.005 [0.064] N=729
Snack foods consumed at least weekly	0.439 P=538	-0.022 [0.047] N=1,238	-0.028 [0.049] N=1,140	-0.009 [0.053] N=1,020	-0.004 [0.055] N=881	0.010 [0.057] N=729
Packaged soft drinks consumed at least weekly	0.248 P=299	-0.027 [0.046] N=1,238	-0.044 [0.048] N=1,140	-0.051 [0.050] N=1,020	-0.035 [0.055] N=881	-0.044 [0.064] N=729
Grains consumed at least weekly	0.858 P=1,065	0.032 [0.038] N=1,238	0.035 [0.040] N=1,140	0.018 [0.042] N=1,020	0.025 [0.047] N=881	0.012 [0.054] N=729
Child was ever breastfed	0.970 P=1,447	-0.010 [0.012] N=1,497	-0.008 [0.013] N=1,384	-0.011 [0.015] N=1,236	-0.017 [0.016] N=1,055	-0.018 [0.016] N=875

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.006 [0.043] N=1,044	0.029 [0.043] N=946	0.012 [0.049] N=826	-0.007 [0.059] N=687	-0.033 [0.082] N=535
Vegetables consumed at least weekly	0.023 [0.049] N=1,044	0.010 [0.050] N=946	-0.027 [0.053] N=826	-0.043 [0.063] N=687	-0.125 [0.094] N=535
Meat fish or poultry consumed at least weekly	0.012 [0.067] N=1,044	0.026 [0.069] N=946	0.009 [0.080] N=826	-0.067 [0.087] N=687	-0.121 [0.092] N=535
Legumes consumed at least weekly	-0.035 [0.071] N=1,044	-0.081 [0.074] N=946	-0.058 [0.077] N=826	-0.108 [0.091] N=687	-0.104 [0.135] N=535
Animal products consumed at least weekly	0.046 [0.051] N=1,044	0.033 [0.056] N=946	-0.019 [0.064] N=826	-0.011 [0.077] N=687	-0.075 [0.104] N=535
Sweets consumed at least weekly	-0.064 [0.064] N=1,044	-0.071 [0.067] N=946	-0.055 [0.080] N=826	-0.121 [0.085] N=687	-0.026 [0.122] N=535
Snack foods consumed at least weekly	-0.041 [0.066] N=1,044	-0.056 [0.071] N=946	-0.026 [0.084] N=826	-0.042 [0.100] N=687	-0.042 [0.124] N=535
Packaged soft drinks consumed at least weekly	-0.018 [0.062] N=1,044	-0.047 [0.066] N=946	-0.060 [0.071] N=826	-0.041 [0.086] N=687	-0.075 [0.122] N=535
Grains consumed at least weekly	0.076 [0.059] N=1,044	0.088 [0.065] N=946	0.066 [0.072] N=826	0.096 [0.090] N=687	0.091 [0.125] N=535
Child was ever breastfed	0.000 [0.019] N=1,265	0.006 [0.022] N=1,152	0.000 [0.026] N=1,004	-0.019 [0.033] N=823	-0.009 [0.038] N=643

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,143	-0.008 [0.028] N=1,238	0.003 [0.027] N=1,140	-0.006 [0.029] N=1,020	-0.008 [0.032] N=881	-0.022 [0.037] N=729
Vegetables consumed at least weekly	0.897 P=1,107	0.008 [0.032] N=1,238	-0.004 [0.031] N=1,140	-0.021 [0.031] N=1,020	-0.019 [0.034] N=881	-0.043 [0.040] N=729
Meat fish or poultry consumed at least weekly	0.760 P=947	0.031 [0.053] N=1,238	0.042 [0.056] N=1,140	0.042 [0.063] N=1,020	0.040 [0.070] N=881	0.028 [0.078] N=729
Legumes consumed at least weekly	0.763 P=949	-0.055 [0.047] N=1,238	-0.079 [0.048] N=1,140	-0.069 [0.050] N=1,020	-0.075 [0.056] N=881	-0.073 [0.068] N=729
Animal products consumed at least weekly	0.900 P=1,118	0.021 [0.038] N=1,238	0.011 [0.040] N=1,140	-0.014 [0.043] N=1,020	-0.006 [0.048] N=881	-0.032 [0.053] N=729
Sweets consumed at least weekly	0.530 P=654	-0.045 [0.047] N=1,238	-0.049 [0.048] N=1,140	-0.036 [0.053] N=1,020	-0.048 [0.056] N=881	-0.010 [0.063] N=729
Snack foods consumed at least weekly	0.439 P=538	-0.017 [0.046] N=1,238	-0.026 [0.048] N=1,140	-0.007 [0.052] N=1,020	-0.001 [0.054] N=881	0.012 [0.056] N=729
Packaged soft drinks consumed at least weekly	0.248 P=299	-0.016 [0.044] N=1,238	-0.032 [0.046] N=1,140	-0.037 [0.049] N=1,020	-0.025 [0.053] N=881	-0.032 [0.061] N=729
Grains consumed at least weekly	0.858 P=1,065	0.029 [0.039] N=1,238	0.032 [0.040] N=1,140	0.016 [0.042] N=1,020	0.023 [0.046] N=881	0.012 [0.053] N=729
Child was ever breastfed	0.970 P=1,447	-0.011 [0.013] N=1,497	-0.010 [0.014] N=1,384	-0.013 [0.015] N=1,236	-0.017 [0.017] N=1,055	-0.020 [0.017] N=875

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.004 [0.042] N=1,044	0.025 [0.042] N=946	0.013 [0.048] N=826	0.012 [0.058] N=687	-0.023 [0.081] N=535
Vegetables consumed at least weekly	0.024 [0.049] N=1,044	0.008 [0.049] N=946	-0.025 [0.053] N=826	-0.022 [0.064] N=687	-0.110 [0.094] N=535
Meat fish or poultry consumed at least weekly	0.001 [0.064] N=1,044	0.009 [0.066] N=946	-0.006 [0.078] N=826	-0.055 [0.085] N=687	-0.124 [0.089] N=535
Legumes consumed at least weekly	-0.039 [0.070] N=1,044	-0.085 [0.073] N=946	-0.057 [0.076] N=826	-0.097 [0.091] N=687	-0.093 [0.137] N=535
Animal products consumed at least weekly	0.047 [0.049] N=1,044	0.033 [0.055] N=946	-0.018 [0.064] N=826	0.005 [0.078] N=687	-0.052 [0.103] N=535
Sweets consumed at least weekly	-0.061 [0.065] N=1,044	-0.069 [0.067] N=946	-0.046 [0.080] N=826	-0.109 [0.086] N=687	-0.021 [0.126] N=535
Snack foods consumed at least weekly	-0.033 [0.066] N=1,044	-0.053 [0.071] N=946	-0.016 [0.083] N=826	-0.030 [0.097] N=687	-0.036 [0.122] N=535
Packaged soft drinks consumed at least weekly	-0.006 [0.060] N=1,044	-0.032 [0.064] N=946	-0.039 [0.070] N=826	-0.032 [0.084] N=687	-0.062 [0.118] N=535
Grains consumed at least weekly	0.073 [0.059] N=1,044	0.087 [0.065] N=946	0.066 [0.072] N=826	0.104 [0.090] N=687	0.097 [0.125] N=535
Child was ever breastfed	0.003 [0.019] N=1,265	0.007 [0.021] N=1,152	0.003 [0.026] N=1,004	-0.008 [0.033] N=823	-0.007 [0.038] N=643

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,143	-0.010 [0.028] N=1,238	0.002 [0.028] N=1,140	-0.011 [0.030] N=1,020	-0.015 [0.032] N=881	-0.033 [0.038] N=729
Vegetables consumed at least weekly	0.897 P=1,107	0.006 [0.032] N=1,238	-0.002 [0.031] N=1,140	-0.021 [0.031] N=1,020	-0.022 [0.034] N=881	-0.043 [0.041] N=729
Meat fish or poultry consumed at least weekly	0.760 P=947	0.025 [0.054] N=1,238	0.037 [0.056] N=1,140	0.030 [0.063] N=1,020	0.024 [0.070] N=881	0.001 [0.080] N=729
Legumes consumed at least weekly	0.763 P=949	-0.058 [0.048] N=1,238	-0.083 [0.050] N=1,140	-0.078 [0.052] N=1,020	-0.084 [0.058] N=881	-0.090 [0.071] N=729
Animal products consumed at least weekly	0.900 P=1,118	0.017 [0.038] N=1,238	0.008 [0.040] N=1,140	-0.019 [0.043] N=1,020	-0.012 [0.047] N=881	-0.042 [0.053] N=729
Sweets consumed at least weekly	0.530 P=654	-0.047 [0.047] N=1,238	-0.054 [0.048] N=1,140	-0.044 [0.053] N=1,020	-0.054 [0.056] N=881	-0.018 [0.064] N=729
Snack foods consumed at least weekly	0.439 P=538	-0.021 [0.047] N=1,238	-0.028 [0.049] N=1,140	-0.013 [0.053] N=1,020	-0.008 [0.055] N=881	0.006 [0.057] N=729
Packaged soft drinks consumed at least weekly	0.248 P=299	-0.017 [0.044] N=1,238	-0.037 [0.047] N=1,140	-0.042 [0.050] N=1,020	-0.027 [0.053] N=881	-0.032 [0.061] N=729
Grains consumed at least weekly	0.858 P=1,065	0.032 [0.039] N=1,238	0.036 [0.040] N=1,140	0.020 [0.042] N=1,020	0.025 [0.047] N=881	0.016 [0.056] N=729
Child was ever breastfed	0.970 P=1,447	-0.014 [0.013] N=1,497	-0.012 [0.015] N=1,384	-0.016 [0.016] N=1,236	-0.021 [0.017] N=1,055	-0.024 [0.018] N=875

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.005 [0.043] N=1,044	0.030 [0.043] N=946	0.011 [0.049] N=826	0.001 [0.059] N=687	-0.051 [0.084] N=535
Vegetables consumed at least weekly	0.023 [0.049] N=1,044	0.011 [0.050] N=946	-0.031 [0.053] N=826	-0.041 [0.063] N=687	-0.148 [0.096] N=535
Meat fish or poultry consumed at least weekly	0.005 [0.064] N=1,044	0.021 [0.064] N=946	0.002 [0.077] N=826	-0.037 [0.084] N=687	-0.142 [0.095] N=535
Legumes consumed at least weekly	-0.034 [0.071] N=1,044	-0.079 [0.074] N=946	-0.054 [0.076] N=826	-0.088 [0.091] N=687	-0.094 [0.142] N=535
Animal products consumed at least weekly	0.048 [0.051] N=1,044	0.036 [0.056] N=946	-0.021 [0.064] N=826	-0.008 [0.077] N=687	-0.085 [0.109] N=535
Sweets consumed at least weekly	-0.052 [0.064] N=1,044	-0.063 [0.066] N=946	-0.041 [0.079] N=826	-0.096 [0.086] N=687	-0.021 [0.126] N=535
Snack foods consumed at least weekly	-0.030 [0.065] N=1,044	-0.050 [0.070] N=946	-0.014 [0.081] N=826	-0.031 [0.096] N=687	-0.046 [0.119] N=535
Packaged soft drinks consumed at least weekly	-0.003 [0.059] N=1,044	-0.037 [0.063] N=946	-0.045 [0.068] N=826	-0.035 [0.081] N=687	-0.058 [0.118] N=535
Grains consumed at least weekly	0.075 [0.059] N=1,044	0.088 [0.065] N=946	0.065 [0.072] N=826	0.099 [0.090] N=687	0.070 [0.123] N=535
Child was ever breastfed	0.002 [0.020] N=1,265	0.007 [0.022] N=1,152	0.000 [0.027] N=1,004	-0.013 [0.034] N=823	-0.006 [0.040] N=643

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,049	-0.011 [0.028] N=1,135	-0.005 [0.028] N=1,050	-0.012 [0.030] N=939	-0.018 [0.032] N=813	-0.036 [0.036] N=677
Vegetables consumed at least weekly	0.899 P=1,017	-0.000 [0.033] N=1,135	-0.014 [0.033] N=1,050	-0.032 [0.033] N=939	-0.028 [0.036] N=813	-0.045 [0.041] N=677
Meat fish or poultry consumed at least weekly	0.761 P=868	0.042 [0.051] N=1,135	0.042 [0.054] N=1,050	0.036 [0.059] N=939	0.014 [0.066] N=813	0.004 [0.073] N=677
Legumes consumed at least weekly	0.759 P=864	-0.076 [0.048] N=1,135	**-0.109** [0.050] N=1,050**	**-0.105** [0.051] N=939**	**-0.120** [0.058] N=813**	**-0.117* [0.070] N=677**
Animal products consumed at least weekly	0.903 P=1,028	0.028 [0.040] N=1,135	0.015 [0.042] N=1,050	-0.013 [0.045] N=939	-0.015 [0.049] N=813	-0.043 [0.055] N=677
Sweets consumed at least weekly	0.531 P=598	-0.024 [0.047] N=1,135	-0.036 [0.050] N=1,050	-0.027 [0.054] N=939	-0.040 [0.058] N=813	0.010 [0.065] N=677
Snack foods consumed at least weekly	0.442 P=495	-0.024 [0.049] N=1,135	-0.031 [0.052] N=1,050	-0.014 [0.056] N=939	-0.015 [0.060] N=813	0.017 [0.062] N=677
Packaged soft drinks consumed at least weekly	0.251 P=278	0.006 [0.046] N=1,135	-0.012 [0.047] N=1,050	-0.033 [0.050] N=939	-0.022 [0.053] N=813	-0.021 [0.060] N=677
Grains consumed at least weekly	0.860 P=978	0.020 [0.040] N=1,135	0.019 [0.042] N=1,050	-0.008 [0.043] N=939	0.001 [0.049] N=813	-0.015 [0.055] N=677
Child was ever breastfed	0.970 P=1,326	-0.015 [0.013] N=1,371	-0.011 [0.014] N=1,272	-0.013 [0.016] N=1,133	-0.017 [0.018] N=972	-0.020 [0.018] N=811

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.045] N=954	0.015 [0.047] N=869	-0.004 [0.054] N=758	-0.024 [0.066] N=632	-0.064 [0.088] N=496
Vegetables consumed at least weekly	0.013 [0.052] N=954	-0.008 [0.054] N=869	-0.049 [0.057] N=758	-0.056 [0.068] N=632	-0.129 [0.092] N=496
Meat fish or poultry consumed at least weekly	0.030 [0.067] N=954	0.016 [0.069] N=869	-0.002 [0.080] N=758	-0.092 [0.089] N=632	-0.145 [0.093] N=496
Legumes consumed at least weekly	-0.048 [0.072] N=954	-0.112 [0.076] N=869	-0.094 [0.077] N=758	**-0.157* [0.093] N=632**	-0.151 [0.133] N=496
Animal products consumed at least weekly	0.059 [0.056] N=954	0.037 [0.061] N=869	-0.019 [0.068] N=758	-0.025 [0.084] N=632	-0.102 [0.111] N=496
Sweets consumed at least weekly	-0.044 [0.066] N=954	-0.071 [0.071] N=869	-0.067 [0.085] N=758	-0.138 [0.092] N=632	-0.048 [0.130] N=496
Snack foods consumed at least weekly	-0.048 [0.069] N=954	-0.066 [0.074] N=869	-0.042 [0.087] N=758	-0.077 [0.103] N=632	-0.060 [0.126] N=496
Packaged soft drinks consumed at least weekly	0.014 [0.064] N=954	-0.023 [0.068] N=869	-0.061 [0.074] N=758	-0.056 [0.088] N=632	-0.070 [0.124] N=496
Grains consumed at least weekly	0.070 [0.062] N=954	0.075 [0.069] N=869	0.030 [0.076] N=758	0.067 [0.095] N=632	0.060 [0.129] N=496
Child was ever breastfed	-0.008 [0.021] N=1,155	0.001 [0.024] N=1,056	-0.003 [0.029] N=917	-0.017 [0.035] N=756	-0.015 [0.042] N=595

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=1,049	-0.012 [0.028] N=1,135	-0.006 [0.027] N=1,050	-0.012 [0.029] N=939	-0.017 [0.032] N=813	-0.034 [0.035] N=677
Vegetables consumed at least weekly	0.899 P=1,017	-0.001 [0.032] N=1,135	-0.015 [0.032] N=1,050	-0.033 [0.033] N=939	-0.028 [0.036] N=813	-0.047 [0.041] N=677
Meat fish or poultry consumed at least weekly	0.761 P=868	0.041 [0.052] N=1,135	0.041 [0.054] N=1,050	0.036 [0.060] N=939	0.016 [0.066] N=813	0.006 [0.073] N=677
Legumes consumed at least weekly	0.759 P=864	-0.076 [0.048] N=1,135	**-0.109** [0.050] N=1,050**	**-0.105** [0.051] N=939**	**-0.121** [0.058] N=813**	**-0.121* [0.070] N=677**
Animal products consumed at least weekly	0.903 P=1,028	0.027 [0.040] N=1,135	0.014 [0.042] N=1,050	-0.012 [0.044] N=939	-0.013 [0.048] N=813	-0.040 [0.054] N=677
Sweets consumed at least weekly	0.531 P=598	-0.024 [0.047] N=1,135	-0.037 [0.050] N=1,050	-0.026 [0.054] N=939	-0.038 [0.056] N=813	0.012 [0.063] N=677
Snack foods consumed at least weekly	0.442 P=495	-0.025 [0.049] N=1,135	-0.032 [0.052] N=1,050	-0.014 [0.056] N=939	-0.014 [0.060] N=813	0.016 [0.062] N=677
Packaged soft drinks consumed at least weekly	0.251 P=278	0.006 [0.046] N=1,135	-0.012 [0.048] N=1,050	-0.030 [0.051] N=939	-0.018 [0.053] N=813	-0.016 [0.061] N=677
Grains consumed at least weekly	0.860 P=978	0.020 [0.039] N=1,135	0.019 [0.042] N=1,050	-0.008 [0.043] N=939	0.002 [0.049] N=813	-0.013 [0.055] N=677
Child was ever breastfed	0.970 P=1,326	-0.015 [0.013] N=1,371	-0.011 [0.014] N=1,272	-0.013 [0.016] N=1,133	-0.017 [0.017] N=972	-0.020 [0.018] N=811

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.044] N=954	0.010 [0.046] N=869	-0.003 [0.053] N=758	-0.020 [0.064] N=632	-0.061 [0.083] N=496
Vegetables consumed at least weekly	0.008 [0.051] N=954	-0.013 [0.053] N=869	-0.051 [0.056] N=758	-0.057 [0.067] N=632	-0.139 [0.089] N=496
Meat fish or poultry consumed at least weekly	0.027 [0.068] N=954	0.016 [0.070] N=869	-0.001 [0.080] N=758	-0.091 [0.089] N=632	-0.154 [0.094] N=496
Legumes consumed at least weekly	-0.052 [0.072] N=954	-0.116 [0.076] N=869	-0.094 [0.078] N=758	**-0.156* [0.093] N=632**	-0.158 [0.132] N=496
Animal products consumed at least weekly	0.054 [0.054] N=954	0.032 [0.059] N=869	-0.019 [0.067] N=758	-0.022 [0.081] N=632	-0.098 [0.107] N=496
Sweets consumed at least weekly	-0.053 [0.066] N=954	-0.084 [0.071] N=869	-0.077 [0.085] N=758	-0.142 [0.090] N=632	-0.056 [0.128] N=496
Snack foods consumed at least weekly	-0.056 [0.069] N=954	-0.077 [0.074] N=869	-0.050 [0.088] N=758	-0.083 [0.103] N=632	-0.068 [0.128] N=496
Packaged soft drinks consumed at least weekly	0.008 [0.065] N=954	-0.030 [0.069] N=869	-0.064 [0.075] N=758	-0.053 [0.088] N=632	-0.061 [0.125] N=496
Grains consumed at least weekly	0.068 [0.061] N=954	0.074 [0.068] N=869	0.033 [0.075] N=758	0.071 [0.095] N=632	0.064 [0.129] N=496
Child was ever breastfed	-0.005 [0.020] N=1,155	0.004 [0.023] N=1,056	-0.002 [0.028] N=917	-0.016 [0.035] N=756	-0.017 [0.041] N=595

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,002	0.054 [0.033] N=1,080	**0.057* [0.034] N=976**	0.030 [0.037] N=848	0.007 [0.039] N=709	0.010 [0.046] N=573
Vegetables consumed at least weekly	0.900 P=970	0.037 [0.037] N=1,080	0.045 [0.038] N=976	0.033 [0.042] N=848	0.002 [0.046] N=709	-0.002 [0.049] N=573
Meat fish or poultry consumed at least weekly	0.758 P=829	**0.103* [0.054] N=1,080**	**0.104* [0.056] N=976**	0.094 [0.063] N=848	**0.124* [0.067] N=709**	0.113 [0.076] N=573
Legumes consumed at least weekly	0.761 P=832	-0.000 [0.043] N=1,080	0.040 [0.046] N=976	0.017 [0.051] N=848	-0.022 [0.055] N=709	-0.002 [0.061] N=573
Animal products consumed at least weekly	0.902 P=979	0.018 [0.040] N=1,080	0.018 [0.042] N=976	-0.003 [0.048] N=848	-0.013 [0.053] N=709	-0.012 [0.061] N=573
Sweets consumed at least weekly	0.537 P=584	-0.057 [0.062] N=1,080	-0.063 [0.064] N=976	-0.080 [0.070] N=848	-0.088 [0.073] N=709	**-0.144* [0.083] N=573**
Snack foods consumed at least weekly	0.440 P=476	-0.045 [0.056] N=1,080	-0.040 [0.060] N=976	-0.008 [0.063] N=848	-0.030 [0.068] N=709	-0.080 [0.076] N=573
Packaged soft drinks consumed at least weekly	0.242 P=256	-0.055 [0.051] N=1,080	-0.065 [0.053] N=976	-0.017 [0.057] N=848	-0.056 [0.061] N=709	-0.092 [0.068] N=573
Grains consumed at least weekly	0.851 P=926	0.045 [0.044] N=1,080	0.056 [0.047] N=976	0.049 [0.050] N=848	0.057 [0.057] N=709	0.034 [0.068] N=573
Child was ever breastfed	0.968 P=1,250	-0.004 [0.023] N=1,293	-0.008 [0.023] N=1,171	0.003 [0.022] N=1,017	0.005 [0.022] N=855	0.018 [0.023] N=688

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.059 [0.051] N=942	0.062 [0.053] N=838	0.014 [0.061] N=710	-0.002 [0.070] N=571	-0.001 [0.094] N=435
Vegetables consumed at least weekly	0.017 [0.059] N=942	0.025 [0.063] N=838	-0.005 [0.072] N=710	-0.059 [0.086] N=571	-0.089 [0.107] N=435
Meat fish or poultry consumed at least weekly	0.037 [0.071] N=942	0.031 [0.075] N=838	-0.022 [0.087] N=710	0.011 [0.095] N=571	-0.064 [0.125] N=435
Legumes consumed at least weekly	-0.016 [0.057] N=942	0.049 [0.061] N=838	0.016 [0.075] N=710	-0.045 [0.085] N=571	0.013 [0.094] N=435
Animal products consumed at least weekly	-0.006 [0.053] N=942	-0.006 [0.058] N=838	-0.060 [0.070] N=710	-0.083 [0.075] N=571	-0.114 [0.095] N=435
Sweets consumed at least weekly	-0.042 [0.084] N=942	-0.049 [0.089] N=838	-0.083 [0.101] N=710	-0.104 [0.118] N=571	**-0.288* [0.145] N=435**
Snack foods consumed at least weekly	-0.064 [0.078] N=942	-0.059 [0.083] N=838	-0.003 [0.093] N=710	-0.037 [0.109] N=571	-0.184 [0.128] N=435
Packaged soft drinks consumed at least weekly	-0.072 [0.075] N=942	-0.080 [0.080] N=838	0.010 [0.095] N=710	-0.055 [0.110] N=571	-0.129 [0.132] N=435
Grains consumed at least weekly	-0.039 [0.051] N=942	-0.038 [0.058] N=838	-0.075 [0.062] N=710	-0.115 [0.072] N=571	**-0.269*** [0.095] N=435**
Child was ever breastfed	0.009 [0.042] N=1,127	0.002 [0.044] N=1,005	0.022 [0.047] N=851	0.027 [0.054] N=689	0.072 [0.070] N=522

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,002	**0.055* [0.033] N=1,080**	**0.057* [0.033] N=976**	0.030 [0.037] N=848	0.007 [0.040] N=709	0.010 [0.046] N=573
Vegetables consumed at least weekly	0.900 P=970	0.038 [0.037] N=1,080	0.046 [0.038] N=976	0.033 [0.042] N=848	0.003 [0.046] N=709	-0.002 [0.048] N=573
Meat fish or poultry consumed at least weekly	0.758 P=829	**0.104* [0.054] N=1,080**	**0.105* [0.056] N=976**	0.095 [0.063] N=848	**0.124* [0.067] N=709**	0.114 [0.076] N=573
Legumes consumed at least weekly	0.761 P=832	-0.000 [0.043] N=1,080	0.040 [0.046] N=976	0.017 [0.051] N=848	-0.023 [0.055] N=709	-0.003 [0.061] N=573
Animal products consumed at least weekly	0.902 P=979	0.021 [0.040] N=1,080	0.021 [0.042] N=976	-0.002 [0.048] N=848	-0.011 [0.053] N=709	-0.011 [0.061] N=573
Sweets consumed at least weekly	0.537 P=584	-0.053 [0.062] N=1,080	-0.060 [0.064] N=976	-0.077 [0.069] N=848	-0.087 [0.073] N=709	**-0.143* [0.083] N=573**
Snack foods consumed at least weekly	0.440 P=476	-0.041 [0.055] N=1,080	-0.037 [0.058] N=976	-0.005 [0.062] N=848	-0.029 [0.067] N=709	-0.079 [0.074] N=573
Packaged soft drinks consumed at least weekly	0.242 P=256	-0.055 [0.051] N=1,080	-0.064 [0.053] N=976	-0.016 [0.056] N=848	-0.056 [0.061] N=709	-0.091 [0.067] N=573
Grains consumed at least weekly	0.851 P=926	0.046 [0.044] N=1,080	0.057 [0.047] N=976	0.050 [0.050] N=848	0.056 [0.057] N=709	0.034 [0.069] N=573
Child was ever breastfed	0.968 P=1,250	-0.004 [0.023] N=1,293	-0.008 [0.023] N=1,171	0.003 [0.022] N=1,017	0.004 [0.023] N=855	0.017 [0.023] N=688

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.050] N=942	0.064 [0.052] N=838	0.015 [0.061] N=710	-0.002 [0.071] N=571	-0.002 [0.097] N=435
Vegetables consumed at least weekly	0.018 [0.059] N=942	0.026 [0.063] N=838	-0.003 [0.072] N=710	-0.053 [0.084] N=571	-0.087 [0.107] N=435
Meat fish or poultry consumed at least weekly	0.039 [0.070] N=942	0.035 [0.073] N=838	-0.017 [0.086] N=710	0.017 [0.093] N=571	-0.057 [0.122] N=435
Legumes consumed at least weekly	-0.016 [0.056] N=942	0.050 [0.061] N=838	0.015 [0.075] N=710	-0.046 [0.086] N=571	0.000 [0.095] N=435
Animal products consumed at least weekly	-0.001 [0.050] N=942	0.000 [0.055] N=838	-0.055 [0.068] N=710	-0.074 [0.072] N=571	-0.101 [0.090] N=435
Sweets consumed at least weekly	-0.035 [0.082] N=942	-0.042 [0.087] N=838	-0.076 [0.100] N=710	-0.101 [0.118] N=571	**-0.281* [0.144] N=435**
Snack foods consumed at least weekly	-0.056 [0.076] N=942	-0.051 [0.081] N=838	0.005 [0.091] N=710	-0.030 [0.107] N=571	-0.172 [0.124] N=435
Packaged soft drinks consumed at least weekly	-0.071 [0.075] N=942	-0.079 [0.079] N=838	0.014 [0.093] N=710	-0.053 [0.109] N=571	-0.118 [0.127] N=435
Grains consumed at least weekly	-0.037 [0.052] N=942	-0.036 [0.058] N=838	-0.075 [0.063] N=710	-0.119 [0.074] N=571	**-0.274*** [0.097] N=435**
Child was ever breastfed	0.010 [0.043] N=1,127	0.002 [0.045] N=1,005	0.022 [0.048] N=851	0.027 [0.055] N=689	0.072 [0.071] N=522

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,002	0.049 [0.033] N=1,080	0.050 [0.034] N=976	0.026 [0.038] N=848	0.004 [0.041] N=709	0.008 [0.048] N=573
Vegetables consumed at least weekly	0.900 P=970	0.031 [0.037] N=1,080	0.038 [0.039] N=976	0.028 [0.043] N=848	-0.001 [0.047] N=709	-0.006 [0.050] N=573
Meat fish or poultry consumed at least weekly	0.758 P=829	0.084 [0.055] N=1,080	0.083 [0.057] N=976	0.088 [0.064] N=848	**0.117* [0.067] N=709**	0.109 [0.077] N=573
Legumes consumed at least weekly	0.761 P=832	-0.008 [0.045] N=1,080	0.032 [0.048] N=976	0.013 [0.053] N=848	-0.026 [0.058] N=709	-0.001 [0.065] N=573
Animal products consumed at least weekly	0.902 P=979	0.014 [0.040] N=1,080	0.011 [0.043] N=976	-0.008 [0.049] N=848	-0.016 [0.054] N=709	-0.015 [0.063] N=573
Sweets consumed at least weekly	0.537 P=584	-0.055 [0.062] N=1,080	-0.063 [0.065] N=976	-0.082 [0.070] N=848	-0.090 [0.074] N=709	**-0.150* [0.084] N=573**
Snack foods consumed at least weekly	0.440 P=476	-0.041 [0.056] N=1,080	-0.038 [0.059] N=976	-0.011 [0.063] N=848	-0.033 [0.068] N=709	-0.086 [0.075] N=573
Packaged soft drinks consumed at least weekly	0.242 P=256	-0.046 [0.051] N=1,080	-0.057 [0.052] N=976	-0.015 [0.056] N=848	-0.053 [0.061] N=709	-0.091 [0.068] N=573
Grains consumed at least weekly	0.851 P=926	0.043 [0.045] N=1,080	0.054 [0.048] N=976	0.049 [0.052] N=848	0.056 [0.059] N=709	0.035 [0.071] N=573
Child was ever breastfed	0.968 P=1,250	-0.006 [0.023] N=1,293	-0.010 [0.024] N=1,171	0.003 [0.023] N=1,017	0.005 [0.024] N=855	0.017 [0.024] N=688

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.051 [0.050] N=942	0.053 [0.051] N=838	0.013 [0.059] N=710	-0.006 [0.071] N=571	-0.010 [0.096] N=435
Vegetables consumed at least weekly	0.009 [0.057] N=942	0.015 [0.062] N=838	-0.009 [0.070] N=710	-0.059 [0.083] N=571	-0.096 [0.103] N=435
Meat fish or poultry consumed at least weekly	0.008 [0.069] N=942	-0.001 [0.072] N=838	-0.031 [0.085] N=710	-0.006 [0.096] N=571	-0.099 [0.120] N=435
Legumes consumed at least weekly	-0.023 [0.057] N=942	0.041 [0.063] N=838	0.014 [0.074] N=710	-0.051 [0.087] N=571	0.004 [0.096] N=435
Animal products consumed at least weekly	-0.006 [0.050] N=942	-0.008 [0.054] N=838	-0.056 [0.067] N=710	-0.074 [0.072] N=571	-0.104 [0.089] N=435
Sweets consumed at least weekly	-0.031 [0.080] N=942	-0.039 [0.086] N=838	-0.072 [0.097] N=710	-0.093 [0.115] N=571	**-0.268* [0.136] N=435**
Snack foods consumed at least weekly	-0.047 [0.075] N=942	-0.044 [0.080] N=838	0.006 [0.089] N=710	-0.027 [0.105] N=571	-0.166 [0.119] N=435
Packaged soft drinks consumed at least weekly	-0.050 [0.075] N=942	-0.058 [0.080] N=838	0.025 [0.093] N=710	-0.037 [0.108] N=571	-0.094 [0.128] N=435
Grains consumed at least weekly	-0.041 [0.053] N=942	-0.040 [0.059] N=838	-0.074 [0.063] N=710	-0.119 [0.074] N=571	**-0.271*** [0.097] N=435**
Child was ever breastfed	0.006 [0.042] N=1,127	-0.003 [0.044] N=1,005	0.022 [0.047] N=851	0.025 [0.053] N=689	0.069 [0.068] N=522

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.929 P=1,002	0.043 [0.033] N=1,080	0.045 [0.033] N=976	0.022 [0.037] N=848	-0.008 [0.040] N=709	-0.009 [0.046] N=573
Vegetables consumed at least weekly	0.900 P=970	0.029 [0.037] N=1,080	0.036 [0.039] N=976	0.024 [0.043] N=848	-0.011 [0.045] N=709	-0.016 [0.048] N=573
Meat fish or poultry consumed at least weekly	0.758 P=829	0.084 [0.055] N=1,080	0.082 [0.057] N=976	0.077 [0.064] N=848	0.094 [0.067] N=709	0.073 [0.074] N=573
Legumes consumed at least weekly	0.761 P=832	-0.003 [0.045] N=1,080	0.036 [0.048] N=976	0.018 [0.052] N=848	-0.028 [0.056] N=709	-0.013 [0.061] N=573
Animal products consumed at least weekly	0.902 P=979	0.020 [0.039] N=1,080	0.019 [0.041] N=976	0.004 [0.048] N=848	-0.010 [0.053] N=709	-0.015 [0.060] N=573
Sweets consumed at least weekly	0.537 P=584	-0.045 [0.063] N=1,080	-0.052 [0.066] N=976	-0.073 [0.071] N=848	-0.084 [0.075] N=709	**-0.144* [0.086] N=573**
Snack foods consumed at least weekly	0.440 P=476	-0.034 [0.056] N=1,080	-0.029 [0.059] N=976	-0.001 [0.063] N=848	-0.028 [0.067] N=709	-0.080 [0.076] N=573
Packaged soft drinks consumed at least weekly	0.242 P=256	-0.038 [0.050] N=1,080	-0.050 [0.052] N=976	-0.009 [0.056] N=848	-0.047 [0.061] N=709	-0.080 [0.069] N=573
Grains consumed at least weekly	0.851 P=926	0.039 [0.045] N=1,080	0.049 [0.048] N=976	0.041 [0.052] N=848	0.046 [0.058] N=709	0.021 [0.069] N=573
Child was ever breastfed	0.968 P=1,250	-0.004 [0.022] N=1,293	-0.009 [0.023] N=1,171	0.003 [0.021] N=1,017	0.003 [0.022] N=855	0.016 [0.022] N=688

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.047 [0.050] N=942	0.050 [0.051] N=838	0.009 [0.061] N=710	-0.025 [0.072] N=571	-0.047 [0.094] N=435
Vegetables consumed at least weekly	0.005 [0.059] N=942	0.012 [0.063] N=838	-0.015 [0.072] N=710	-0.077 [0.084] N=571	-0.132 [0.103] N=435
Meat fish or poultry consumed at least weekly	0.017 [0.073] N=942	0.007 [0.077] N=838	-0.035 [0.091] N=710	-0.033 [0.106] N=571	-0.153 [0.126] N=435
Legumes consumed at least weekly	-0.015 [0.057] N=942	0.048 [0.064] N=838	0.029 [0.078] N=710	-0.045 [0.089] N=571	-0.010 [0.099] N=435
Animal products consumed at least weekly	-0.000 [0.047] N=942	0.001 [0.052] N=838	-0.041 [0.066] N=710	-0.070 [0.071] N=571	-0.106 [0.085] N=435
Sweets consumed at least weekly	-0.015 [0.080] N=942	-0.021 [0.086] N=838	-0.055 [0.098] N=710	-0.078 [0.115] N=571	**-0.253* [0.140] N=435**
Snack foods consumed at least weekly	-0.041 [0.075] N=942	-0.035 [0.080] N=838	0.019 [0.090] N=710	-0.022 [0.104] N=571	-0.162 [0.119] N=435
Packaged soft drinks consumed at least weekly	-0.047 [0.073] N=942	-0.057 [0.078] N=838	0.021 [0.092] N=710	-0.034 [0.107] N=571	-0.073 [0.130] N=435
Grains consumed at least weekly	-0.041 [0.052] N=942	-0.042 [0.058] N=838	-0.082 [0.064] N=710	**-0.134* [0.075] N=571**	**-0.295*** [0.094] N=435**
Child was ever breastfed	0.008 [0.041] N=1,127	-0.002 [0.043] N=1,005	0.023 [0.045] N=851	0.022 [0.052] N=689	0.063 [0.066] N=522

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=917	0.053 [0.035] N=986	0.054 [0.036] N=888	0.043 [0.039] N=778	0.008 [0.043] N=650	0.011 [0.049] N=530
Vegetables consumed at least weekly	0.904 P=890	0.027 [0.037] N=986	0.031 [0.039] N=888	0.043 [0.042] N=778	0.004 [0.049] N=650	-0.015 [0.051] N=530
Meat fish or poultry consumed at least weekly	0.760 P=757	**0.095* [0.054] N=986**	0.087 [0.056] N=888	0.091 [0.062] N=778	0.097 [0.066] N=650	0.090 [0.074] N=530
Legumes consumed at least weekly	0.759 P=758	-0.001 [0.048] N=986	0.039 [0.051] N=888	0.023 [0.055] N=778	-0.026 [0.062] N=650	-0.008 [0.070] N=530
Animal products consumed at least weekly	0.906 P=898	0.020 [0.043] N=986	0.023 [0.045] N=888	0.013 [0.052] N=778	-0.001 [0.058] N=650	-0.002 [0.066] N=530
Sweets consumed at least weekly	0.549 P=543	-0.052 [0.062] N=986	-0.057 [0.065] N=888	-0.081 [0.071] N=778	-0.099 [0.074] N=650	**-0.147* [0.087] N=530**
Snack foods consumed at least weekly	0.450 P=443	-0.031 [0.058] N=986	-0.028 [0.062] N=888	-0.002 [0.065] N=778	-0.029 [0.069] N=650	-0.066 [0.078] N=530
Packaged soft drinks consumed at least weekly	0.249 P=242	-0.065 [0.052] N=986	-0.078 [0.053] N=888	-0.028 [0.057] N=778	-0.077 [0.063] N=650	**-0.122* [0.071] N=530**
Grains consumed at least weekly	0.854 P=848	0.063 [0.046] N=986	0.073 [0.049] N=888	0.077 [0.052] N=778	0.077 [0.060] N=650	0.046 [0.071] N=530
Child was ever breastfed	0.970 P=1,145	-0.012 [0.022] N=1,181	-0.016 [0.023] N=1,068	-0.005 [0.021] N=934	-0.008 [0.022] N=786	0.009 [0.021] N=637

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.047 [0.050] N=858	0.047 [0.051] N=760	0.029 [0.059] N=650	-0.026 [0.068] N=522	-0.044 [0.080] N=402
Vegetables consumed at least weekly	-0.004 [0.051] N=858	-0.004 [0.056] N=760	0.006 [0.061] N=650	-0.075 [0.077] N=522	**-0.164* [0.085] N=402**
Meat fish or poultry consumed at least weekly	0.047 [0.077] N=858	0.032 [0.080] N=760	0.010 [0.091] N=650	-0.024 [0.099] N=522	-0.101 [0.122] N=402
Legumes consumed at least weekly	-0.012 [0.061] N=858	0.053 [0.067] N=760	0.045 [0.077] N=650	-0.018 [0.094] N=522	0.024 [0.104] N=402
Animal products consumed at least weekly	-0.018 [0.051] N=858	-0.014 [0.056] N=760	-0.041 [0.067] N=650	-0.077 [0.074] N=522	-0.121 [0.089] N=402
Sweets consumed at least weekly	-0.035 [0.084] N=858	-0.041 [0.090] N=760	-0.096 [0.102] N=650	-0.150 [0.116] N=522	**-0.332** [0.151] N=402**
Snack foods consumed at least weekly	-0.050 [0.080] N=858	-0.046 [0.087] N=760	-0.009 [0.093] N=650	-0.070 [0.106] N=522	-0.182 [0.132] N=402
Packaged soft drinks consumed at least weekly	-0.084 [0.077] N=858	-0.095 [0.082] N=760	-0.009 [0.096] N=650	-0.098 [0.115] N=522	-0.223 [0.140] N=402
Grains consumed at least weekly	-0.020 [0.056] N=858	-0.021 [0.063] N=760	-0.034 [0.069] N=650	-0.105 [0.081] N=522	**-0.302*** [0.103] N=402**
Child was ever breastfed	-0.001 [0.040] N=1,027	-0.010 [0.042] N=914	0.010 [0.043] N=780	0.003 [0.050] N=632	0.043 [0.063] N=483

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=917	0.053 [0.035] N=986	0.055 [0.035] N=888	0.044 [0.039] N=778	0.007 [0.042] N=650	0.011 [0.048] N=530
Vegetables consumed at least weekly	0.904 P=890	0.030 [0.037] N=986	0.034 [0.039] N=888	0.046 [0.041] N=778	0.004 [0.047] N=650	-0.016 [0.049] N=530
Meat fish or poultry consumed at least weekly	0.760 P=757	**0.101* [0.052] N=986**	**0.094* [0.055] N=888**	0.096 [0.060] N=778	0.096 [0.064] N=650	0.090 [0.071] N=530
Legumes consumed at least weekly	0.759 P=758	-0.001 [0.048] N=986	0.040 [0.051] N=888	0.024 [0.054] N=778	-0.029 [0.061] N=650	-0.011 [0.069] N=530
Animal products consumed at least weekly	0.906 P=898	0.020 [0.042] N=986	0.023 [0.045] N=888	0.013 [0.051] N=778	-0.001 [0.057] N=650	-0.002 [0.065] N=530
Sweets consumed at least weekly	0.549 P=543	-0.055 [0.062] N=986	-0.060 [0.065] N=888	-0.085 [0.071] N=778	-0.105 [0.074] N=650	**-0.151* [0.087] N=530**
Snack foods consumed at least weekly	0.450 P=443	-0.029 [0.057] N=986	-0.027 [0.061] N=888	-0.001 [0.064] N=778	-0.028 [0.068] N=650	-0.064 [0.077] N=530
Packaged soft drinks consumed at least weekly	0.249 P=242	-0.065 [0.051] N=986	-0.078 [0.053] N=888	-0.030 [0.056] N=778	-0.079 [0.061] N=650	**-0.124* [0.069] N=530**
Grains consumed at least weekly	0.854 P=848	0.063 [0.046] N=986	0.073 [0.049] N=888	0.076 [0.052] N=778	0.076 [0.060] N=650	0.046 [0.071] N=530
Child was ever breastfed	0.970 P=1,145	-0.011 [0.022] N=1,181	-0.015 [0.023] N=1,068	-0.003 [0.021] N=934	-0.005 [0.022] N=786	0.011 [0.022] N=637

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.050 [0.051] N=858	0.052 [0.052] N=760	0.034 [0.060] N=650	-0.023 [0.071] N=522	-0.035 [0.089] N=402
Vegetables consumed at least weekly	-0.001 [0.051] N=858	-0.000 [0.057] N=760	0.012 [0.061] N=650	-0.076 [0.076] N=522	**-0.158* [0.089] N=402**
Meat fish or poultry consumed at least weekly	0.051 [0.076] N=858	0.036 [0.080] N=760	0.011 [0.091] N=650	-0.027 [0.098] N=522	-0.089 [0.124] N=402
Legumes consumed at least weekly	-0.009 [0.061] N=858	0.058 [0.066] N=760	0.049 [0.076] N=650	-0.025 [0.093] N=522	0.035 [0.103] N=402
Animal products consumed at least weekly	-0.017 [0.051] N=858	-0.013 [0.057] N=760	-0.040 [0.067] N=650	-0.079 [0.074] N=522	-0.125 [0.093] N=402
Sweets consumed at least weekly	-0.032 [0.085] N=858	-0.040 [0.090] N=760	-0.092 [0.103] N=650	-0.143 [0.118] N=522	**-0.323** [0.152] N=402**
Snack foods consumed at least weekly	-0.046 [0.080] N=858	-0.044 [0.086] N=760	0.000 [0.093] N=650	-0.049 [0.108] N=522	-0.167 [0.130] N=402
Packaged soft drinks consumed at least weekly	-0.081 [0.078] N=858	-0.092 [0.083] N=760	0.000 [0.095] N=650	-0.080 [0.113] N=522	-0.192 [0.139] N=402
Grains consumed at least weekly	-0.020 [0.056] N=858	-0.023 [0.062] N=760	-0.038 [0.068] N=650	-0.112 [0.080] N=522	**-0.307*** [0.101] N=402**
Child was ever breastfed	-0.002 [0.040] N=1,027	-0.011 [0.042] N=914	0.013 [0.044] N=780	0.009 [0.052] N=632	0.048 [0.065] N=483

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.