Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.063	0.237 [0.236] N=1,273	0.314 [0.244] N=1,179	0.374 [0.270] N=1,053	**0.516* [0.276] N=906**	0.233 [0.304] N=756
WHO weight-for-age z score	0.300	-0.167 [0.131] N=1,479	-0.150 [0.147] N=1,368	-0.099 [0.154] N=1,222	-0.049 [0.153] N=1,043	-0.133 [0.173] N=864
WHO weight-for-length/height z score, 0-60 months	0.313	0.280 [0.211] N=1,264	0.275 [0.220] N=1,169	0.336 [0.228] N=1,047	0.218 [0.238] N=900	0.265 [0.252] N=748
WHO BMI-for-age z score	0.298	0.242 [0.231] N=1,245	0.302 [0.246] N=1,154	0.347 [0.261] N=1,031	0.150 [0.268] N=881	0.174 [0.287] N=733
Stunted: HAZ below -2	0.195 P=240	-0.051 [0.047] N=1,273	-0.051 [0.049] N=1,179	-0.041 [0.053] N=1,053	-0.047 [0.055] N=906	-0.026 [0.064] N=756
Underweight: WAZ below -2	0.061 P=93	**0.046** [0.023] N=1,479**	0.041 [0.025] N=1,368	**0.043* [0.026] N=1,222**	0.023 [0.027] N=1,043	0.012 [0.035] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=147	-0.048 [0.035] N=1,264	-0.035 [0.037] N=1,169	-0.055 [0.039] N=1,047	-0.033 [0.042] N=900	-0.048 [0.044] N=748
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=244	0.023 [0.045] N=1,264	0.027 [0.048] N=1,169	0.026 [0.050] N=1,047	0.016 [0.056] N=900	0.009 [0.064] N=748
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=113	0.002 [0.034] N=1,264	0.006 [0.037] N=1,169	0.023 [0.039] N=1,047	0.013 [0.043] N=900	0.020 [0.048] N=748
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=315	-0.070 [0.051] N=1,206	-0.041 [0.053] N=1,116	-0.036 [0.056] N=998	-0.044 [0.060] N=859	-0.026 [0.071] N=715
BMI-for-age z score below -2	0.125 P=153	-0.027 [0.038] N=1,245	-0.019 [0.041] N=1,154	-0.024 [0.047] N=1,031	0.021 [0.048] N=881	0.026 [0.053] N=733

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.128 [0.365] N=1,067	0.246 [0.406] N=973	0.438 [0.463] N=847	**0.874* [0.486] N=700**	0.335 [0.664] N=550
WHO weight-for-age z score	**-0.417** [0.187] N=1,249**	**-0.422* [0.229] N=1,138**	-0.373 [0.240] N=992	-0.352 [0.267] N=813	**-0.703* [0.406] N=634**
WHO weight-for-length/height z score, 0-60 months	0.230 [0.304] N=1,060	0.219 [0.334] N=965	0.251 [0.350] N=843	0.033 [0.412] N=696	-0.012 [0.487] N=544
WHO BMI-for-age z score	0.308 [0.306] N=1,044	0.425 [0.340] N=953	0.421 [0.368] N=830	0.068 [0.412] N=680	-0.004 [0.496] N=532
Stunted: HAZ below -2	-0.031 [0.067] N=1,067	-0.024 [0.072] N=973	-0.013 [0.080] N=847	-0.018 [0.093] N=700	0.043 [0.133] N=550
Underweight: WAZ below -2	**0.084*** [0.032] N=1,249**	**0.082** [0.037] N=1,138**	**0.096** [0.039] N=992**	**0.074* [0.043] N=813**	0.075 [0.075] N=634
Wasted: WLZ/WHZ below -2, 0-60 months	-0.068 [0.049] N=1,060	-0.046 [0.055] N=965	-0.085 [0.061] N=843	-0.060 [0.074] N=696	-0.117 [0.086] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	-0.013 [0.056] N=1,060	-0.009 [0.063] N=965	-0.030 [0.062] N=843	-0.059 [0.081] N=696	-0.149 [0.101] N=544
Obese: WLZ/WHZ above +3, 0-60 months	-0.028 [0.042] N=1,060	-0.025 [0.049] N=965	-0.000 [0.049] N=843	-0.023 [0.061] N=696	-0.042 [0.078] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.051 [0.071] N=1,008	0.008 [0.077] N=918	0.019 [0.084] N=800	0.010 [0.101] N=661	0.084 [0.140] N=517
BMI-for-age z score below -2	**-0.082* [0.045] N=1,044**	-0.078 [0.049] N=953	-0.092 [0.060] N=830	-0.028 [0.067] N=680	-0.034 [0.075] N=532

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.063	0.238 [0.236] N=1,273	0.314 [0.245] N=1,179	0.376 [0.271] N=1,053	**0.524* [0.276] N=906**	0.249 [0.305] N=756
WHO weight-for-age z score	0.300	-0.172 [0.130] N=1,479	-0.159 [0.147] N=1,368	-0.106 [0.152] N=1,222	-0.055 [0.153] N=1,043	-0.142 [0.172] N=864
WHO weight-for-length/height z score, 0-60 months	0.313	0.278 [0.211] N=1,264	0.274 [0.221] N=1,169	0.334 [0.229] N=1,047	0.213 [0.239] N=900	0.256 [0.254] N=748
WHO BMI-for-age z score	0.298	0.240 [0.231] N=1,245	0.299 [0.246] N=1,154	0.345 [0.261] N=1,031	0.149 [0.269] N=881	0.172 [0.287] N=733
Stunted: HAZ below -2	0.195 P=240	-0.050 [0.047] N=1,273	-0.049 [0.049] N=1,179	-0.040 [0.053] N=1,053	-0.046 [0.055] N=906	-0.024 [0.064] N=756
Underweight: WAZ below -2	0.061 P=93	**0.047** [0.023] N=1,479**	**0.042* [0.025] N=1,368**	**0.044* [0.026] N=1,222**	0.023 [0.027] N=1,043	0.013 [0.035] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=147	-0.046 [0.035] N=1,264	-0.032 [0.037] N=1,169	-0.052 [0.040] N=1,047	-0.029 [0.042] N=900	-0.043 [0.045] N=748
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=244	0.024 [0.045] N=1,264	0.028 [0.048] N=1,169	0.027 [0.050] N=1,047	0.018 [0.056] N=900	0.011 [0.064] N=748
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=113	0.003 [0.034] N=1,264	0.008 [0.037] N=1,169	0.024 [0.039] N=1,047	0.015 [0.043] N=900	0.022 [0.048] N=748
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=315	-0.066 [0.050] N=1,206	-0.036 [0.051] N=1,116	-0.032 [0.055] N=998	-0.040 [0.059] N=859	-0.019 [0.070] N=715
BMI-for-age z score below -2	0.125 P=153	-0.026 [0.039] N=1,245	-0.017 [0.042] N=1,154	-0.022 [0.047] N=1,031	0.023 [0.049] N=881	0.029 [0.054] N=733

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.126 [0.364] N=1,067	0.245 [0.405] N=973	0.429 [0.460] N=847	**0.843* [0.480] N=700**	0.281 [0.654] N=550
WHO weight-for-age z score	**-0.410** [0.186] N=1,249**	**-0.420* [0.229] N=1,138**	-0.365 [0.241] N=992	-0.347 [0.268] N=813	**-0.694* [0.405] N=634**
WHO weight-for-length/height z score, 0-60 months	0.231 [0.304] N=1,060	0.219 [0.334] N=965	0.254 [0.350] N=843	0.036 [0.414] N=696	-0.004 [0.492] N=544
WHO BMI-for-age z score	0.311 [0.306] N=1,044	0.426 [0.340] N=953	0.429 [0.368] N=830	0.076 [0.416] N=680	0.008 [0.503] N=532
Stunted: HAZ below -2	-0.032 [0.067] N=1,067	-0.026 [0.072] N=973	-0.015 [0.079] N=847	-0.018 [0.093] N=700	0.045 [0.132] N=550
Underweight: WAZ below -2	**0.084*** [0.032] N=1,249**	**0.082** [0.037] N=1,138**	**0.096** [0.039] N=992**	**0.074* [0.043] N=813**	0.074 [0.074] N=634
Wasted: WLZ/WHZ below -2, 0-60 months	-0.068 [0.049] N=1,060	-0.046 [0.055] N=965	-0.088 [0.061] N=843	-0.061 [0.075] N=696	-0.120 [0.086] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	-0.013 [0.056] N=1,060	-0.009 [0.063] N=965	-0.030 [0.061] N=843	-0.059 [0.081] N=696	-0.150 [0.101] N=544
Obese: WLZ/WHZ above +3, 0-60 months	-0.029 [0.042] N=1,060	-0.025 [0.048] N=965	-0.002 [0.049] N=843	-0.024 [0.061] N=696	-0.043 [0.077] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.053 [0.070] N=1,008	0.005 [0.075] N=918	0.011 [0.081] N=800	0.000 [0.099] N=661	0.071 [0.139] N=517
BMI-for-age z score below -2	**-0.083* [0.045] N=1,044**	-0.078 [0.049] N=953	-0.095 [0.060] N=830	-0.031 [0.069] N=680	-0.038 [0.078] N=532

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.063	0.209 [0.236] N=1,273	0.274 [0.244] N=1,179	0.317 [0.270] N=1,053	**0.498* [0.273] N=906**	0.202 [0.300] N=756
WHO weight-for-age z score	0.300	-0.204 [0.125] N=1,479	-0.192 [0.142] N=1,368	-0.140 [0.147] N=1,222	-0.069 [0.149] N=1,043	-0.169 [0.168] N=864
WHO weight-for-length/height z score, 0-60 months	0.313	0.276 [0.213] N=1,264	0.277 [0.224] N=1,169	0.348 [0.233] N=1,047	0.221 [0.242] N=900	0.258 [0.261] N=748
WHO BMI-for-age z score	0.298	0.243 [0.233] N=1,245	0.303 [0.249] N=1,154	0.359 [0.265] N=1,031	0.159 [0.273] N=881	0.180 [0.296] N=733
Stunted: HAZ below -2	0.195 P=240	-0.046 [0.048] N=1,273	-0.042 [0.049] N=1,179	-0.033 [0.053] N=1,053	-0.043 [0.055] N=906	-0.018 [0.065] N=756
Underweight: WAZ below -2	0.061 P=93	**0.049** [0.022] N=1,479**	**0.045* [0.024] N=1,368**	**0.047* [0.025] N=1,222**	0.024 [0.026] N=1,043	0.018 [0.035] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=147	-0.045 [0.035] N=1,264	-0.030 [0.037] N=1,169	-0.053 [0.040] N=1,047	-0.030 [0.042] N=900	-0.041 [0.045] N=748
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=244	0.025 [0.045] N=1,264	0.030 [0.048] N=1,169	0.029 [0.050] N=1,047	0.018 [0.056] N=900	0.014 [0.064] N=748
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=113	0.003 [0.035] N=1,264	0.009 [0.038] N=1,169	0.025 [0.039] N=1,047	0.016 [0.043] N=900	0.023 [0.048] N=748
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=315	-0.060 [0.050] N=1,206	-0.027 [0.052] N=1,116	-0.026 [0.055] N=998	-0.039 [0.060] N=859	-0.008 [0.070] N=715
BMI-for-age z score below -2	0.125 P=153	-0.026 [0.039] N=1,245	-0.016 [0.042] N=1,154	-0.022 [0.048] N=1,031	0.023 [0.049] N=881	0.029 [0.055] N=733

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.100 [0.362] N=1,067	0.199 [0.399] N=973	0.333 [0.453] N=847	**0.805* [0.466] N=700**	0.262 [0.625] N=550
WHO weight-for-age z score	**-0.427** [0.183] N=1,249**	**-0.441* [0.226] N=1,138**	**-0.388* [0.234] N=992**	-0.332 [0.273] N=813	**-0.687* [0.405] N=634**
WHO weight-for-length/height z score, 0-60 months	0.232 [0.302] N=1,060	0.225 [0.334] N=965	0.275 [0.345] N=843	0.023 [0.403] N=696	-0.032 [0.489] N=544
WHO BMI-for-age z score	0.317 [0.306] N=1,044	0.431 [0.341] N=953	0.457 [0.366] N=830	0.069 [0.402] N=680	-0.003 [0.499] N=532
Stunted: HAZ below -2	-0.030 [0.066] N=1,067	-0.020 [0.071] N=973	-0.008 [0.078] N=847	-0.023 [0.090] N=700	0.041 [0.129] N=550
Underweight: WAZ below -2	**0.084*** [0.031] N=1,249**	**0.083** [0.037] N=1,138**	**0.095** [0.038] N=992**	0.068 [0.044] N=813	0.071 [0.075] N=634
Wasted: WLZ/WHZ below -2, 0-60 months	-0.069 [0.050] N=1,060	-0.046 [0.056] N=965	-0.095 [0.062] N=843	-0.070 [0.077] N=696	-0.119 [0.089] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	-0.013 [0.056] N=1,060	-0.008 [0.064] N=965	-0.033 [0.062] N=843	-0.072 [0.080] N=696	-0.149 [0.102] N=544
Obese: WLZ/WHZ above +3, 0-60 months	-0.030 [0.042] N=1,060	-0.026 [0.049] N=965	-0.005 [0.050] N=843	-0.031 [0.062] N=696	-0.047 [0.078] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.054 [0.071] N=1,008	0.007 [0.077] N=918	0.004 [0.081] N=800	-0.026 [0.100] N=661	0.065 [0.139] N=517
BMI-for-age z score below -2	**-0.084* [0.046] N=1,044**	-0.077 [0.050] N=953	-0.101 [0.062] N=830	-0.037 [0.069] N=680	-0.035 [0.079] N=532

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.063	0.196 [0.236] N=1,273	0.276 [0.244] N=1,179	0.318 [0.270] N=1,053	**0.510* [0.276] N=906**	0.213 [0.307] N=756
WHO weight-for-age z score	0.300	**-0.224* [0.124] N=1,479**	-0.216 [0.140] N=1,368	-0.170 [0.147] N=1,222	-0.102 [0.148] N=1,043	-0.223 [0.170] N=864
WHO weight-for-length/height z score, 0-60 months	0.313	0.270 [0.214] N=1,264	0.262 [0.224] N=1,169	0.326 [0.233] N=1,047	0.190 [0.244] N=900	0.201 [0.261] N=748
WHO BMI-for-age z score	0.298	0.237 [0.233] N=1,245	0.293 [0.247] N=1,154	0.344 [0.263] N=1,031	0.128 [0.273] N=881	0.115 [0.294] N=733
Stunted: HAZ below -2	0.195 P=240	-0.042 [0.048] N=1,273	-0.038 [0.049] N=1,179	-0.026 [0.053] N=1,053	-0.037 [0.055] N=906	-0.008 [0.064] N=756
Underweight: WAZ below -2	0.061 P=93	**0.053** [0.022] N=1,479**	**0.050** [0.024] N=1,368**	**0.053** [0.024] N=1,222**	0.031 [0.025] N=1,043	0.029 [0.034] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=147	-0.044 [0.035] N=1,264	-0.028 [0.036] N=1,169	-0.047 [0.039] N=1,047	-0.024 [0.042] N=900	-0.037 [0.044] N=748
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=244	0.025 [0.045] N=1,264	0.031 [0.049] N=1,169	0.031 [0.051] N=1,047	0.019 [0.057] N=900	0.008 [0.065] N=748
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=113	0.002 [0.034] N=1,264	0.006 [0.037] N=1,169	0.023 [0.039] N=1,047	0.012 [0.043] N=900	0.014 [0.047] N=748
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=315	-0.052 [0.050] N=1,206	-0.016 [0.052] N=1,116	-0.007 [0.055] N=998	-0.020 [0.059] N=859	0.017 [0.070] N=715
BMI-for-age z score below -2	0.125 P=153	-0.024 [0.039] N=1,245	-0.014 [0.042] N=1,154	-0.017 [0.048] N=1,031	0.030 [0.050] N=881	0.037 [0.055] N=733

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062 [0.369] N=1,067	0.186 [0.402] N=973	0.304 [0.456] N=847	0.811 [0.494] N=700	0.198 [0.653] N=550
WHO weight-for-age z score	**-0.428** [0.183] N=1,249**	**-0.434* [0.222] N=1,138**	**-0.396* [0.232] N=992**	-0.296 [0.275] N=813	-0.639 [0.400] N=634
WHO weight-for-length/height z score, 0-60 months	0.255 [0.303] N=1,060	0.247 [0.332] N=965	0.290 [0.353] N=843	0.080 [0.417] N=696	0.045 [0.498] N=544
WHO BMI-for-age z score	0.333 [0.307] N=1,044	0.452 [0.338] N=953	0.471 [0.374] N=830	0.141 [0.417] N=680	0.102 [0.510] N=532
Stunted: HAZ below -2	-0.027 [0.067] N=1,067	-0.021 [0.072] N=973	-0.007 [0.080] N=847	-0.027 [0.092] N=700	0.040 [0.131] N=550
Underweight: WAZ below -2	**0.082*** [0.031] N=1,249**	**0.080** [0.036] N=1,138**	**0.094** [0.038] N=992**	0.061 [0.044] N=813	0.072 [0.074] N=634
Wasted: WLZ/WHZ below -2, 0-60 months	-0.068 [0.050] N=1,060	-0.048 [0.055] N=965	-0.091 [0.062] N=843	-0.064 [0.077] N=696	-0.116 [0.091] N=544
Overweight: WLZ/WHZ above +2, 0-60 months	-0.010 [0.056] N=1,060	-0.006 [0.063] N=965	-0.027 [0.062] N=843	-0.058 [0.080] N=696	-0.122 [0.099] N=544
Obese: WLZ/WHZ above +3, 0-60 months	-0.023 [0.043] N=1,060	-0.019 [0.049] N=965	0.004 [0.050] N=843	-0.023 [0.061] N=696	-0.041 [0.074] N=544
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.053 [0.071] N=1,008	0.004 [0.077] N=918	0.007 [0.082] N=800	-0.030 [0.102] N=661	0.061 [0.143] N=517
BMI-for-age z score below -2	**-0.084* [0.045] N=1,044**	-0.082 [0.050] N=953	-0.100 [0.061] N=830	-0.042 [0.071] N=680	-0.045 [0.081] N=532

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.073	0.241 [0.236] N=1,169	0.263 [0.247] N=1,087	0.300 [0.275] N=968	0.368 [0.280] N=839	0.026 [0.304] N=702
WHO weight-for-age z score	0.308	-0.092 [0.136] N=1,354	-0.089 [0.150] N=1,257	-0.023 [0.156] N=1,120	0.027 [0.155] N=961	-0.060 [0.177] N=801
WHO weight-for-length/height z score, 0-60 months	0.327	**0.407* [0.211] N=1,165**	**0.403* [0.217] N=1,083**	**0.477** [0.224] N=969**	**0.381* [0.229] N=840**	**0.476* [0.240] N=701**
WHO BMI-for-age z score	0.315	0.386 [0.235] N=1,146	**0.423* [0.245] N=1,067**	**0.497* [0.258] N=953**	0.331 [0.259] N=821	0.389 [0.279] N=686
Stunted: HAZ below -2	0.193 P=217	-0.045 [0.050] N=1,169	-0.038 [0.051] N=1,087	-0.024 [0.055] N=968	-0.021 [0.058] N=839	0.013 [0.068] N=702
Underweight: WAZ below -2	0.060 P=85	**0.041* [0.023] N=1,354**	0.036 [0.025] N=1,257	0.040 [0.027] N=1,120	0.021 [0.028] N=961	0.017 [0.036] N=801
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=134	**-0.071** [0.035] N=1,165**	-0.054 [0.036] N=1,083	**-0.072* [0.039] N=969**	-0.050 [0.041] N=840	-0.065 [0.043] N=701
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=227	0.039 [0.045] N=1,165	0.045 [0.048] N=1,083	0.049 [0.050] N=969	0.042 [0.055] N=840	0.048 [0.063] N=701
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=108	-0.002 [0.036] N=1,165	0.005 [0.039] N=1,083	0.028 [0.041] N=969	0.028 [0.044] N=840	0.044 [0.049] N=701
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=287	-0.077 [0.053] N=1,110	-0.041 [0.055] N=1,032	-0.036 [0.058] N=921	-0.048 [0.061] N=800	-0.025 [0.074] N=668
BMI-for-age z score below -2	0.121 P=136	-0.050 [0.037] N=1,146	-0.040 [0.039] N=1,067	-0.047 [0.045] N=953	-0.007 [0.046] N=821	-0.012 [0.051] N=686

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.377 [0.380] N=977	0.461 [0.427] N=895	0.649 [0.488] N=776	**0.928* [0.516] N=647**	0.312 [0.711] N=510
WHO weight-for-age z score	**-0.353* [0.195] N=1,140**	-0.383 [0.232] N=1,043	-0.319 [0.243] N=906	-0.314 [0.261] N=747	-0.656 [0.407] N=587
WHO weight-for-length/height z score, 0-60 months	0.218 [0.324] N=974	0.163 [0.347] N=892	0.189 [0.366] N=778	0.048 [0.433] N=649	0.096 [0.507] N=510
WHO BMI-for-age z score	0.305 [0.336] N=958	0.337 [0.362] N=879	0.364 [0.386] N=765	0.107 [0.433] N=633	0.034 [0.534] N=498
Stunted: HAZ below -2	-0.061 [0.070] N=977	-0.055 [0.075] N=895	-0.039 [0.083] N=776	-0.035 [0.095] N=647	0.030 [0.144] N=510
Underweight: WAZ below -2	**0.078** [0.032] N=1,140**	**0.078** [0.038] N=1,043**	**0.094** [0.040] N=906**	**0.077* [0.045] N=747**	0.093 [0.074] N=587
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.100* [0.051] N=974**	-0.072 [0.055] N=892	**-0.110* [0.060] N=778**	-0.100 [0.073] N=649	**-0.182** [0.082] N=510**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.023 [0.059] N=974	-0.028 [0.065] N=892	-0.046 [0.065] N=778	-0.074 [0.086] N=649	-0.142 [0.109] N=510
Obese: WLZ/WHZ above +3, 0-60 months	-0.058 [0.046] N=974	-0.054 [0.053] N=892	-0.025 [0.055] N=778	-0.028 [0.067] N=649	-0.035 [0.084] N=510
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.086 [0.073] N=925	-0.024 [0.080] N=847	-0.018 [0.086] N=736	-0.052 [0.100] N=615	-0.015 [0.150] N=483
BMI-for-age z score below -2	**-0.084* [0.047] N=958**	-0.070 [0.051] N=879	-0.088 [0.061] N=765	-0.036 [0.070] N=633	-0.057 [0.079] N=498

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.073	0.235 [0.238] N=1,169	0.253 [0.249] N=1,087	0.294 [0.277] N=968	0.369 [0.278] N=839	0.028 [0.299] N=702
WHO weight-for-age z score	0.308	-0.077 [0.135] N=1,354	-0.075 [0.150] N=1,257	-0.014 [0.155] N=1,120	0.025 [0.155] N=961	-0.055 [0.179] N=801
WHO weight-for-length/height z score, 0-60 months	0.327	**0.416** [0.208] N=1,165**	**0.412* [0.214] N=1,083**	**0.477** [0.220] N=969**	0.368 [0.226] N=840	**0.456* [0.237] N=701**
WHO BMI-for-age z score	0.315	**0.391* [0.231] N=1,146**	**0.428* [0.240] N=1,067**	**0.495* [0.253] N=953**	0.322 [0.254] N=821	0.378 [0.272] N=686
Stunted: HAZ below -2	0.193 P=217	-0.046 [0.050] N=1,169	-0.039 [0.051] N=1,087	-0.024 [0.055] N=968	-0.019 [0.057] N=839	0.013 [0.067] N=702
Underweight: WAZ below -2	0.060 P=85	0.038 [0.023] N=1,354	0.034 [0.025] N=1,257	0.038 [0.026] N=1,120	0.021 [0.028] N=961	0.015 [0.036] N=801
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=134	**-0.070** [0.035] N=1,165**	-0.053 [0.036] N=1,083	**-0.071* [0.039] N=969**	-0.049 [0.041] N=840	-0.066 [0.043] N=701
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=227	0.042 [0.045] N=1,165	0.048 [0.047] N=1,083	0.049 [0.049] N=969	0.035 [0.055] N=840	0.039 [0.062] N=701
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=108	-0.000 [0.036] N=1,165	0.007 [0.039] N=1,083	0.028 [0.040] N=969	0.026 [0.044] N=840	0.038 [0.048] N=701
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=287	-0.077 [0.053] N=1,110	-0.041 [0.055] N=1,032	-0.035 [0.058] N=921	-0.044 [0.062] N=800	-0.022 [0.074] N=668
BMI-for-age z score below -2	0.121 P=136	-0.049 [0.037] N=1,146	-0.039 [0.039] N=1,067	-0.047 [0.045] N=953	-0.007 [0.046] N=821	-0.011 [0.051] N=686

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.357 [0.383] N=977	0.426 [0.433] N=895	0.622 [0.497] N=776	**0.917* [0.520] N=647**	0.340 [0.706] N=510
WHO weight-for-age z score	**-0.323* [0.190] N=1,140**	-0.352 [0.229] N=1,043	-0.295 [0.237] N=906	-0.315 [0.260] N=747	-0.637 [0.414] N=587
WHO weight-for-length/height z score, 0-60 months	0.252 [0.319] N=974	0.208 [0.345] N=892	0.221 [0.366] N=778	0.025 [0.431] N=649	0.027 [0.506] N=510
WHO BMI-for-age z score	0.330 [0.332] N=958	0.369 [0.359] N=879	0.388 [0.386] N=765	0.099 [0.427] N=633	0.014 [0.515] N=498
Stunted: HAZ below -2	-0.064 [0.071] N=977	-0.057 [0.076] N=895	-0.042 [0.083] N=776	-0.036 [0.095] N=647	0.024 [0.141] N=510
Underweight: WAZ below -2	**0.072** [0.033] N=1,140**	**0.072* [0.038] N=1,043**	**0.090** [0.040] N=906**	**0.077* [0.045] N=747**	0.086 [0.076] N=587
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.098* [0.050] N=974**	-0.071 [0.055] N=892	**-0.111* [0.060] N=778**	-0.099 [0.074] N=649	**-0.182** [0.082] N=510**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.011 [0.059] N=974	-0.014 [0.065] N=892	-0.036 [0.065] N=778	-0.078 [0.085] N=649	-0.155 [0.107] N=510
Obese: WLZ/WHZ above +3, 0-60 months	-0.054 [0.046] N=974	-0.050 [0.053] N=892	-0.021 [0.054] N=778	-0.031 [0.067] N=649	-0.048 [0.084] N=510
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.087 [0.073] N=925	-0.027 [0.081] N=847	-0.020 [0.086] N=736	-0.052 [0.100] N=615	-0.020 [0.148] N=483
BMI-for-age z score below -2	**-0.081* [0.047] N=958**	-0.069 [0.051] N=879	-0.088 [0.061] N=765	-0.035 [0.070] N=633	-0.055 [0.079] N=498

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.122	-0.079 [0.344] N=1,113	-0.171 [0.350] N=1,002	-0.046 [0.375] N=871	0.003 [0.433] N=736	0.037 [0.491] N=594
WHO weight-for-age z score	0.307	0.018 [0.159] N=1,279	0.025 [0.168] N=1,158	-0.042 [0.176] N=1,004	-0.008 [0.191] N=846	0.063 [0.213] N=680
WHO weight-for-length/height z score, 0-60 months	0.357	0.055 [0.272] N=1,111	0.056 [0.281] N=1,004	0.010 [0.308] N=872	0.068 [0.347] N=734	0.016 [0.391] N=591
WHO BMI-for-age z score	0.331	0.138 [0.271] N=1,090	0.172 [0.280] N=985	0.118 [0.315] N=855	0.202 [0.353] N=720	0.238 [0.396] N=582
Stunted: HAZ below -2	0.180 P=201	-0.048 [0.054] N=1,113	-0.034 [0.055] N=1,002	-0.077 [0.057] N=871	-0.068 [0.067] N=736	-0.121 [0.074] N=594
Underweight: WAZ below -2	0.056 P=74	-0.003 [0.028] N=1,279	-0.006 [0.029] N=1,158	0.007 [0.029] N=1,004	0.016 [0.028] N=846	0.006 [0.031] N=680
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=127	0.007 [0.042] N=1,111	0.016 [0.044] N=1,004	0.024 [0.049] N=872	-0.005 [0.050] N=734	0.006 [0.055] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=222	0.042 [0.050] N=1,111	0.054 [0.052] N=1,004	0.025 [0.053] N=872	0.023 [0.060] N=734	0.024 [0.068] N=591
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.028 [0.041] N=1,111	0.026 [0.042] N=1,004	0.015 [0.043] N=872	0.005 [0.049] N=734	-0.019 [0.050] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	0.260 P=275	-0.010 [0.049] N=1,059	0.003 [0.051] N=955	-0.027 [0.053] N=829	-0.043 [0.055] N=699	-0.070 [0.059] N=562
BMI-for-age z score below -2	0.126 P=136	0.034 [0.047] N=1,090	0.043 [0.049] N=985	0.033 [0.057] N=855	0.017 [0.060] N=720	0.043 [0.068] N=582

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.212 [0.463] N=968	-0.367 [0.489] N=857	-0.238 [0.564] N=726	-0.287 [0.703] N=591	-0.214 [0.802] N=449
WHO weight-for-age z score	-0.012 [0.225] N=1,115	-0.003 [0.246] N=994	-0.176 [0.265] N=840	-0.162 [0.300] N=682	-0.103 [0.350] N=516
WHO weight-for-length/height z score, 0-60 months	-0.161 [0.378] N=968	-0.187 [0.403] N=861	-0.334 [0.473] N=729	-0.274 [0.549] N=591	-0.771 [0.633] N=448
WHO BMI-for-age z score	-0.221 [0.363] N=951	-0.190 [0.384] N=846	-0.375 [0.465] N=716	-0.309 [0.531] N=581	-0.560 [0.578] N=443
Stunted: HAZ below -2	0.022 [0.078] N=968	0.047 [0.083] N=857	-0.006 [0.090] N=726	0.049 [0.116] N=591	-0.103 [0.126] N=449
Underweight: WAZ below -2	-0.015 [0.048] N=1,115	-0.020 [0.052] N=994	0.008 [0.057] N=840	0.019 [0.061] N=682	-0.007 [0.069] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.045 [0.059] N=968	0.066 [0.062] N=861	0.089 [0.075] N=729	0.039 [0.077] N=591	0.107 [0.090] N=448
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.075] N=968	0.032 [0.081] N=861	-0.023 [0.086] N=729	-0.035 [0.102] N=591	-0.109 [0.124] N=448
Obese: WLZ/WHZ above +3, 0-60 months	0.047 [0.057] N=968	0.038 [0.060] N=861	0.020 [0.067] N=729	-0.003 [0.085] N=591	-0.115 [0.098] N=448
CIAF: any stunting, underweight, or wasting, 0-60 months	0.054 [0.081] N=921	0.079 [0.085] N=817	0.049 [0.095] N=691	0.045 [0.108] N=561	-0.035 [0.123] N=424
BMI-for-age z score below -2	0.057 [0.051] N=951	0.073 [0.055] N=846	0.062 [0.072] N=716	0.019 [0.069] N=581	0.095 [0.081] N=443

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.122	-0.095 [0.342] N=1,113	-0.194 [0.348] N=1,002	-0.084 [0.371] N=871	-0.031 [0.428] N=736	0.004 [0.483] N=594
WHO weight-for-age z score	0.307	0.026 [0.159] N=1,279	0.033 [0.168] N=1,158	-0.040 [0.176] N=1,004	-0.003 [0.191] N=846	0.066 [0.213] N=680
WHO weight-for-length/height z score, 0-60 months	0.357	0.059 [0.271] N=1,111	0.055 [0.282] N=1,004	0.009 [0.309] N=872	0.070 [0.346] N=734	0.025 [0.388] N=591
WHO BMI-for-age z score	0.331	0.140 [0.272] N=1,090	0.171 [0.281] N=985	0.115 [0.315] N=855	0.201 [0.354] N=720	0.242 [0.396] N=582
Stunted: HAZ below -2	0.180 P=201	-0.050 [0.054] N=1,113	-0.035 [0.056] N=1,002	-0.077 [0.058] N=871	-0.067 [0.068] N=736	-0.119 [0.074] N=594
Underweight: WAZ below -2	0.056 P=74	-0.004 [0.028] N=1,279	-0.007 [0.029] N=1,158	0.007 [0.029] N=1,004	0.016 [0.029] N=846	0.006 [0.031] N=680
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=127	0.002 [0.042] N=1,111	0.011 [0.044] N=1,004	0.019 [0.049] N=872	-0.009 [0.049] N=734	0.002 [0.053] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=222	0.039 [0.050] N=1,111	0.049 [0.053] N=1,004	0.019 [0.054] N=872	0.018 [0.061] N=734	0.022 [0.070] N=591
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.024 [0.041] N=1,111	0.022 [0.042] N=1,004	0.010 [0.043] N=872	0.001 [0.049] N=734	-0.023 [0.050] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	0.260 P=275	-0.020 [0.050] N=1,059	-0.006 [0.051] N=955	-0.035 [0.053] N=829	-0.048 [0.056] N=699	-0.073 [0.059] N=562
BMI-for-age z score below -2	0.126 P=136	0.030 [0.046] N=1,090	0.039 [0.049] N=985	0.030 [0.057] N=855	0.014 [0.059] N=720	0.040 [0.068] N=582

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.225 [0.465] N=968	-0.400 [0.492] N=857	-0.313 [0.563] N=726	-0.380 [0.704] N=591	-0.328 [0.813] N=449
WHO weight-for-age z score	0.005 [0.224] N=1,115	0.015 [0.245] N=994	-0.167 [0.264] N=840	-0.138 [0.299] N=682	-0.074 [0.350] N=516
WHO weight-for-length/height z score, 0-60 months	-0.154 [0.383] N=968	-0.190 [0.410] N=861	-0.339 [0.480] N=729	-0.267 [0.559] N=591	-0.724 [0.656] N=448
WHO BMI-for-age z score	-0.208 [0.369] N=951	-0.184 [0.391] N=846	-0.371 [0.471] N=716	-0.293 [0.538] N=581	-0.502 [0.601] N=443
Stunted: HAZ below -2	0.014 [0.079] N=968	0.041 [0.084] N=857	-0.009 [0.091] N=726	0.048 [0.118] N=591	-0.102 [0.129] N=449
Underweight: WAZ below -2	-0.017 [0.049] N=1,115	-0.022 [0.052] N=994	0.008 [0.058] N=840	0.019 [0.063] N=682	-0.005 [0.072] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.061] N=968	0.055 [0.064] N=861	0.078 [0.077] N=729	0.025 [0.078] N=591	0.081 [0.094] N=448
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.075] N=968	0.021 [0.081] N=861	-0.039 [0.086] N=729	-0.056 [0.102] N=591	-0.127 [0.124] N=448
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.057] N=968	0.029 [0.060] N=861	0.009 [0.066] N=729	-0.017 [0.085] N=591	-0.134 [0.098] N=448
CIAF: any stunting, underweight, or wasting, 0-60 months	0.030 [0.080] N=921	0.053 [0.085] N=817	0.022 [0.094] N=691	0.016 [0.108] N=561	-0.071 [0.121] N=424
BMI-for-age z score below -2	0.047 [0.052] N=951	0.061 [0.057] N=846	0.050 [0.073] N=716	0.004 [0.070] N=581	0.076 [0.084] N=443

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.122	-0.118 [0.330] N=1,113	-0.197 [0.336] N=1,002	-0.070 [0.362] N=871	-0.030 [0.416] N=736	0.003 [0.472] N=594
WHO weight-for-age z score	0.307	0.018 [0.156] N=1,279	0.025 [0.164] N=1,158	-0.032 [0.174] N=1,004	-0.000 [0.190] N=846	0.063 [0.211] N=680
WHO weight-for-length/height z score, 0-60 months	0.357	0.064 [0.271] N=1,111	0.053 [0.282] N=1,004	0.014 [0.307] N=872	0.071 [0.342] N=734	0.012 [0.385] N=591
WHO BMI-for-age z score	0.331	0.139 [0.271] N=1,090	0.159 [0.280] N=985	0.111 [0.309] N=855	0.198 [0.344] N=720	0.212 [0.385] N=582
Stunted: HAZ below -2	0.180 P=201	-0.046 [0.052] N=1,113	-0.033 [0.055] N=1,002	-0.080 [0.056] N=871	-0.068 [0.065] N=736	**-0.127* [0.071] N=594**
Underweight: WAZ below -2	0.056 P=74	-0.003 [0.028] N=1,279	-0.005 [0.029] N=1,158	0.007 [0.029] N=1,004	0.015 [0.029] N=846	0.005 [0.031] N=680
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=127	0.003 [0.043] N=1,111	0.012 [0.045] N=1,004	0.018 [0.050] N=872	-0.010 [0.050] N=734	0.003 [0.055] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=222	0.041 [0.050] N=1,111	0.050 [0.053] N=1,004	0.018 [0.053] N=872	0.015 [0.060] N=734	0.020 [0.068] N=591
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.026 [0.041] N=1,111	0.022 [0.042] N=1,004	0.009 [0.044] N=872	-0.000 [0.050] N=734	-0.026 [0.050] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	0.260 P=275	-0.014 [0.051] N=1,059	-0.003 [0.052] N=955	-0.040 [0.055] N=829	-0.055 [0.058] N=699	-0.086 [0.059] N=562
BMI-for-age z score below -2	0.126 P=136	0.032 [0.047] N=1,090	0.041 [0.050] N=985	0.029 [0.058] N=855	0.013 [0.060] N=720	0.043 [0.069] N=582

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.322 [0.466] N=968	-0.475 [0.491] N=857	-0.338 [0.558] N=726	-0.452 [0.693] N=591	-0.454 [0.819] N=449
WHO weight-for-age z score	-0.048 [0.223] N=1,115	-0.038 [0.243] N=994	-0.174 [0.261] N=840	-0.163 [0.297] N=682	-0.140 [0.349] N=516
WHO weight-for-length/height z score, 0-60 months	-0.163 [0.385] N=968	-0.202 [0.412] N=861	-0.336 [0.478] N=729	-0.259 [0.556] N=591	-0.727 [0.667] N=448
WHO BMI-for-age z score	-0.212 [0.372] N=951	-0.199 [0.394] N=846	-0.358 [0.467] N=716	-0.263 [0.535] N=581	-0.475 [0.610] N=443
Stunted: HAZ below -2	0.026 [0.078] N=968	0.048 [0.083] N=857	-0.004 [0.089] N=726	0.062 [0.116] N=591	-0.084 [0.129] N=449
Underweight: WAZ below -2	-0.012 [0.048] N=1,115	-0.017 [0.051] N=994	0.010 [0.057] N=840	0.022 [0.062] N=682	0.005 [0.072] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.041 [0.062] N=968	0.064 [0.065] N=861	0.080 [0.078] N=729	0.026 [0.079] N=591	0.089 [0.096] N=448
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.076] N=968	0.027 [0.082] N=861	-0.038 [0.086] N=729	-0.052 [0.102] N=591	-0.121 [0.124] N=448
Obese: WLZ/WHZ above +3, 0-60 months	0.042 [0.057] N=968	0.033 [0.060] N=861	0.010 [0.067] N=729	-0.014 [0.084] N=591	-0.130 [0.098] N=448
CIAF: any stunting, underweight, or wasting, 0-60 months	0.048 [0.081] N=921	0.068 [0.085] N=817	0.027 [0.097] N=691	0.026 [0.112] N=561	-0.053 [0.122] N=424
BMI-for-age z score below -2	0.049 [0.054] N=951	0.066 [0.058] N=846	0.050 [0.073] N=716	0.004 [0.070] N=581	0.077 [0.086] N=443

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.122	-0.108 [0.330] N=1,113	-0.185 [0.337] N=1,002	-0.040 [0.362] N=871	-0.016 [0.412] N=736	-0.003 [0.469] N=594
WHO weight-for-age z score	0.307	0.013 [0.155] N=1,279	0.025 [0.164] N=1,158	-0.025 [0.175] N=1,004	0.002 [0.191] N=846	0.071 [0.216] N=680
WHO weight-for-length/height z score, 0-60 months	0.357	0.074 [0.268] N=1,111	0.068 [0.278] N=1,004	0.023 [0.300] N=872	0.103 [0.336] N=734	0.085 [0.382] N=591
WHO BMI-for-age z score	0.331	0.150 [0.271] N=1,090	0.177 [0.280] N=985	0.123 [0.308] N=855	0.224 [0.344] N=720	0.290 [0.393] N=582
Stunted: HAZ below -2	0.180 P=201	-0.050 [0.053] N=1,113	-0.038 [0.055] N=1,002	-0.085 [0.057] N=871	-0.070 [0.066] N=736	**-0.121* [0.072] N=594**
Underweight: WAZ below -2	0.056 P=74	-0.005 [0.028] N=1,279	-0.007 [0.028] N=1,158	0.005 [0.029] N=1,004	0.017 [0.029] N=846	0.007 [0.032] N=680
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=127	0.001 [0.042] N=1,111	0.011 [0.044] N=1,004	0.021 [0.048] N=872	-0.011 [0.048] N=734	-0.003 [0.051] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=222	0.041 [0.049] N=1,111	0.050 [0.052] N=1,004	0.022 [0.053] N=872	0.024 [0.060] N=734	0.031 [0.069] N=591
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=107	0.030 [0.040] N=1,111	0.028 [0.041] N=1,004	0.014 [0.043] N=872	0.007 [0.048] N=734	-0.015 [0.049] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	0.260 P=275	-0.017 [0.050] N=1,059	-0.004 [0.052] N=955	-0.039 [0.054] N=829	-0.050 [0.056] N=699	-0.069 [0.058] N=562
BMI-for-age z score below -2	0.126 P=136	0.028 [0.047] N=1,090	0.038 [0.049] N=985	0.029 [0.056] N=855	0.012 [0.058] N=720	0.035 [0.065] N=582

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.276 [0.446] N=968	-0.418 [0.476] N=857	-0.216 [0.551] N=726	-0.340 [0.690] N=591	-0.366 [0.804] N=449
WHO weight-for-age z score	-0.020 [0.215] N=1,115	-0.002 [0.235] N=994	-0.120 [0.260] N=840	-0.110 [0.290] N=682	-0.054 [0.346] N=516
WHO weight-for-length/height z score, 0-60 months	-0.159 [0.381] N=968	-0.196 [0.407] N=861	-0.352 [0.471] N=729	-0.258 [0.546] N=591	-0.676 [0.656] N=448
WHO BMI-for-age z score	-0.208 [0.368] N=951	-0.184 [0.392] N=846	-0.369 [0.467] N=716	-0.260 [0.529] N=581	-0.438 [0.601] N=443
Stunted: HAZ below -2	0.020 [0.076] N=968	0.041 [0.082] N=857	-0.022 [0.089] N=726	0.043 [0.117] N=591	-0.089 [0.127] N=449
Underweight: WAZ below -2	-0.019 [0.047] N=1,115	-0.024 [0.050] N=994	0.001 [0.057] N=840	0.016 [0.062] N=682	-0.006 [0.073] N=516
Wasted: WLZ/WHZ below -2, 0-60 months	0.041 [0.061] N=968	0.063 [0.065] N=861	0.088 [0.076] N=729	0.034 [0.077] N=591	0.093 [0.090] N=448
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.075] N=968	0.020 [0.081] N=861	-0.044 [0.085] N=729	-0.058 [0.102] N=591	-0.118 [0.127] N=448
Obese: WLZ/WHZ above +3, 0-60 months	0.045 [0.056] N=968	0.035 [0.059] N=861	0.012 [0.064] N=729	-0.009 [0.082] N=591	-0.119 [0.098] N=448
CIAF: any stunting, underweight, or wasting, 0-60 months	0.041 [0.080] N=921	0.061 [0.085] N=817	0.015 [0.095] N=691	0.012 [0.109] N=561	-0.048 [0.120] N=424
BMI-for-age z score below -2	0.050 [0.053] N=951	0.065 [0.058] N=846	0.055 [0.073] N=716	0.008 [0.069] N=581	0.077 [0.082] N=443

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.135	-0.211 [0.340] N=1,019	-0.324 [0.348] N=917	-0.211 [0.365] N=802	-0.187 [0.409] N=675	-0.200 [0.472] N=547
WHO weight-for-age z score	0.310	0.031 [0.166] N=1,168	0.055 [0.174] N=1,056	-0.035 [0.181] N=922	-0.018 [0.192] N=778	0.021 [0.210] N=630
WHO weight-for-length/height z score, 0-60 months	0.364	0.175 [0.287] N=1,021	0.197 [0.303] N=923	0.112 [0.329] N=805	0.133 [0.363] N=677	0.048 [0.401] N=549
WHO BMI-for-age z score	0.339	0.198 [0.293] N=1,001	0.258 [0.307] N=905	0.142 [0.345] N=790	0.197 [0.383] N=665	0.172 [0.428] N=541
Stunted: HAZ below -2	0.177 P=182	-0.042 [0.056] N=1,019	-0.024 [0.058] N=917	-0.070 [0.058] N=802	-0.063 [0.065] N=675	-0.110 [0.074] N=547
Underweight: WAZ below -2	0.056 P=68	0.001 [0.030] N=1,168	-0.002 [0.031] N=1,056	0.011 [0.032] N=922	0.025 [0.032] N=778	0.018 [0.034] N=630
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=116	0.001 [0.044] N=1,021	0.008 [0.046] N=923	0.016 [0.050] N=805	-0.010 [0.050] N=677	0.005 [0.054] N=549
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=207	0.053 [0.053] N=1,021	0.066 [0.056] N=923	0.027 [0.057] N=805	0.015 [0.063] N=677	0.007 [0.069] N=549
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=101	0.029 [0.042] N=1,021	0.027 [0.043] N=923	0.017 [0.044] N=805	0.005 [0.050] N=677	-0.016 [0.048] N=549
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=252	-0.010 [0.053] N=972	0.001 [0.055] N=877	-0.034 [0.054] N=765	-0.055 [0.058] N=643	-0.071 [0.064] N=520
BMI-for-age z score below -2	0.123 P=121	0.014 [0.049] N=1,001	0.019 [0.052] N=905	0.009 [0.060] N=790	0.001 [0.063] N=665	0.036 [0.071] N=541

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.152 [0.487] N=885	-0.347 [0.514] N=783	-0.229 [0.580] N=668	-0.296 [0.717] N=541	-0.113 [0.858] N=413
WHO weight-for-age z score	0.046 [0.233] N=1,016	0.069 [0.252] N=904	-0.140 [0.278] N=770	-0.099 [0.312] N=626	-0.003 [0.372] N=478
WHO weight-for-length/height z score, 0-60 months	-0.051 [0.413] N=887	-0.054 [0.445] N=789	-0.277 [0.515] N=671	-0.258 [0.606] N=543	-0.899 [0.717] N=415
WHO BMI-for-age z score	-0.151 [0.387] N=870	-0.086 [0.413] N=774	-0.374 [0.493] N=659	-0.317 [0.563] N=534	-0.721 [0.630] N=410
Stunted: HAZ below -2	-0.000 [0.086] N=885	0.032 [0.093] N=783	-0.024 [0.096] N=668	0.011 [0.125] N=541	-0.169 [0.148] N=413
Underweight: WAZ below -2	-0.021 [0.052] N=1,016	-0.027 [0.055] N=904	-0.002 [0.062] N=770	0.016 [0.071] N=626	-0.015 [0.082] N=478
Wasted: WLZ/WHZ below -2, 0-60 months	0.034 [0.064] N=887	0.051 [0.068] N=789	0.067 [0.082] N=671	0.005 [0.083] N=543	0.061 [0.094] N=415
Overweight: WLZ/WHZ above +2, 0-60 months	0.032 [0.083] N=887	0.038 [0.090] N=789	-0.037 [0.095] N=671	-0.082 [0.115] N=543	-0.197 [0.136] N=415
Obese: WLZ/WHZ above +3, 0-60 months	0.031 [0.060] N=887	0.020 [0.064] N=789	0.002 [0.071] N=671	-0.034 [0.092] N=543	-0.162 [0.111] N=415
CIAF: any stunting, underweight, or wasting, 0-60 months	0.033 [0.089] N=843	0.052 [0.096] N=748	0.012 [0.101] N=636	-0.037 [0.119] N=514	-0.174 [0.143] N=391
BMI-for-age z score below -2	0.036 [0.058] N=870	0.045 [0.063] N=774	0.027 [0.081] N=659	-0.024 [0.080] N=534	0.068 [0.094] N=410

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.135	-0.227 [0.346] N=1,019	-0.339 [0.354] N=917	-0.228 [0.370] N=802	-0.205 [0.411] N=675	-0.202 [0.474] N=547
WHO weight-for-age z score	0.310	0.038 [0.163] N=1,168	0.054 [0.172] N=1,056	-0.031 [0.179] N=922	-0.012 [0.189] N=778	0.024 [0.206] N=630
WHO weight-for-length/height z score, 0-60 months	0.364	0.161 [0.291] N=1,021	0.182 [0.306] N=923	0.102 [0.332] N=805	0.128 [0.365] N=677	0.045 [0.410] N=549
WHO BMI-for-age z score	0.339	0.200 [0.296] N=1,001	0.262 [0.310] N=905	0.145 [0.348] N=790	0.201 [0.385] N=665	0.175 [0.434] N=541
Stunted: HAZ below -2	0.177 P=182	-0.041 [0.056] N=1,019	-0.023 [0.058] N=917	-0.068 [0.058] N=802	-0.062 [0.065] N=675	-0.111 [0.073] N=547
Underweight: WAZ below -2	0.056 P=68	-0.003 [0.030] N=1,168	-0.005 [0.031] N=1,056	0.007 [0.031] N=922	0.020 [0.031] N=778	0.013 [0.033] N=630
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=116	-0.004 [0.043] N=1,021	0.004 [0.046] N=923	0.011 [0.050] N=805	-0.016 [0.049] N=677	-0.001 [0.054] N=549
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=207	0.049 [0.053] N=1,021	0.062 [0.056] N=923	0.023 [0.056] N=805	0.010 [0.062] N=677	0.004 [0.069] N=549
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=101	0.024 [0.041] N=1,021	0.022 [0.042] N=923	0.013 [0.043] N=805	0.002 [0.048] N=677	-0.019 [0.046] N=549
CIAF: any stunting, underweight, or wasting, 0-60 months	0.259 P=252	-0.015 [0.052] N=972	-0.005 [0.054] N=877	-0.040 [0.053] N=765	-0.064 [0.055] N=643	-0.082 [0.059] N=520
BMI-for-age z score below -2	0.123 P=121	0.013 [0.049] N=1,001	0.017 [0.052] N=905	0.007 [0.060] N=790	-0.002 [0.062] N=665	0.033 [0.070] N=541

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.152 [0.485] N=885	-0.341 [0.512] N=783	-0.220 [0.574] N=668	-0.285 [0.697] N=541	-0.132 [0.830] N=413
WHO weight-for-age z score	0.050 [0.232] N=1,016	0.068 [0.253] N=904	-0.132 [0.276] N=770	-0.103 [0.307] N=626	-0.048 [0.361] N=478
WHO weight-for-length/height z score, 0-60 months	-0.105 [0.410] N=887	-0.111 [0.441] N=789	-0.335 [0.510] N=671	-0.325 [0.585] N=543	-0.908 [0.684] N=415
WHO BMI-for-age z score	-0.156 [0.386] N=870	-0.089 [0.411] N=774	-0.377 [0.491] N=659	-0.313 [0.551] N=534	-0.672 [0.604] N=410
Stunted: HAZ below -2	-0.001 [0.086] N=885	0.030 [0.092] N=783	-0.025 [0.095] N=668	0.007 [0.123] N=541	-0.162 [0.142] N=413
Underweight: WAZ below -2	-0.023 [0.052] N=1,016	-0.026 [0.056] N=904	-0.004 [0.063] N=770	0.016 [0.071] N=626	-0.005 [0.080] N=478
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.063] N=887	0.054 [0.068] N=789	0.070 [0.082] N=671	0.010 [0.081] N=543	0.070 [0.094] N=415
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.082] N=887	0.029 [0.088] N=789	-0.048 [0.093] N=671	-0.094 [0.110] N=543	-0.192 [0.130] N=415
Obese: WLZ/WHZ above +3, 0-60 months	0.022 [0.060] N=887	0.011 [0.063] N=789	-0.007 [0.070] N=671	-0.043 [0.089] N=543	-0.159 [0.102] N=415
CIAF: any stunting, underweight, or wasting, 0-60 months	0.032 [0.089] N=843	0.052 [0.095] N=748	0.012 [0.100] N=636	-0.036 [0.117] N=514	-0.136 [0.137] N=391
BMI-for-age z score below -2	0.039 [0.058] N=870	0.048 [0.063] N=774	0.030 [0.081] N=659	-0.019 [0.077] N=534	0.071 [0.091] N=410

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.