Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Youngest Child

Age reference: 12 November 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,112	-0.005 [0.028] N=1,204	0.006 [0.028] N=1,108	-0.006 [0.029] N=992	-0.013 [0.031] N=856	-0.026 [0.035] N=706
Vegetables consumed at least weekly	0.896 P=1,076	0.001 [0.032] N=1,204	-0.010 [0.031] N=1,108	-0.027 [0.032] N=992	-0.033 [0.035] N=856	-0.058 [0.041] N=706
Meat fish or poultry consumed at least weekly	0.756 P=917	0.052 [0.059] N=1,204	0.062 [0.062] N=1,108	0.062 [0.069] N=992	0.045 [0.077] N=856	0.043 [0.086] N=706
Legumes consumed at least weekly	0.758 P=918	-0.053 [0.047] N=1,204	-0.080 [0.049] N=1,108	-0.066 [0.051] N=992	-0.084 [0.057] N=856	-0.083 [0.070] N=706
Animal products consumed at least weekly	0.902 P=1,090	0.015 [0.039] N=1,204	0.008 [0.040] N=1,108	-0.016 [0.043] N=992	-0.012 [0.048] N=856	-0.024 [0.055] N=706
Sweets consumed at least weekly	0.520 P=625	-0.042 [0.047] N=1,204	-0.045 [0.048] N=1,108	-0.042 [0.053] N=992	-0.058 [0.056] N=856	-0.017 [0.063] N=706
Snack foods consumed at least weekly	0.431 P=514	-0.013 [0.047] N=1,204	-0.023 [0.049] N=1,108	-0.014 [0.053] N=992	-0.013 [0.055] N=856	-0.009 [0.058] N=706
Packaged soft drinks consumed at least weekly	0.244 P=288	-0.014 [0.046] N=1,204	-0.037 [0.047] N=1,108	-0.049 [0.049] N=992	-0.038 [0.054] N=856	-0.058 [0.061] N=706
Grains consumed at least weekly	0.856 P=1,033	0.022 [0.037] N=1,204	0.026 [0.038] N=1,108	0.006 [0.040] N=992	0.013 [0.045] N=856	-0.005 [0.052] N=706
Child was ever breastfed	0.969 P=1,416	-0.012 [0.013] N=1,466	-0.009 [0.014] N=1,355	-0.012 [0.015] N=1,211	-0.019 [0.016] N=1,032	-0.020 [0.016] N=853

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.000 [0.045] N=1,015	0.023 [0.045] N=919	0.000 [0.051] N=803	-0.021 [0.061] N=667	-0.064 [0.084] N=517
Vegetables consumed at least weekly	0.019 [0.050] N=1,015	0.003 [0.051] N=919	-0.030 [0.055] N=803	-0.052 [0.066] N=667	-0.157 [0.099] N=517
Meat fish or poultry consumed at least weekly	0.015 [0.068] N=1,015	0.024 [0.071] N=919	0.017 [0.081] N=803	-0.077 [0.086] N=667	-0.148 [0.095] N=517
Legumes consumed at least weekly	-0.037 [0.072] N=1,015	-0.085 [0.075] N=919	-0.045 [0.078] N=803	-0.107 [0.091] N=667	-0.121 [0.131] N=517
Animal products consumed at least weekly	0.028 [0.052] N=1,015	0.020 [0.058] N=919	-0.029 [0.065] N=803	-0.021 [0.078] N=667	-0.058 [0.104] N=517
Sweets consumed at least weekly	-0.059 [0.064] N=1,015	-0.067 [0.067] N=919	-0.067 [0.081] N=803	**-0.142* [0.084] N=667**	-0.053 [0.119] N=517
Snack foods consumed at least weekly	-0.036 [0.065] N=1,015	-0.063 [0.070] N=919	-0.054 [0.082] N=803	-0.084 [0.095] N=667	-0.118 [0.117] N=517
Packaged soft drinks consumed at least weekly	-0.008 [0.065] N=1,015	-0.054 [0.070] N=919	-0.079 [0.076] N=803	-0.074 [0.091] N=667	-0.158 [0.126] N=517
Grains consumed at least weekly	0.057 [0.055] N=1,015	0.069 [0.059] N=919	0.036 [0.064] N=803	0.060 [0.076] N=667	0.021 [0.106] N=517
Child was ever breastfed	-0.001 [0.020] N=1,239	0.005 [0.023] N=1,128	0.003 [0.027] N=984	-0.017 [0.034] N=805	-0.004 [0.040] N=626

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,112	-0.006 [0.028] N=1,204	0.005 [0.027] N=1,108	-0.006 [0.029] N=992	-0.014 [0.031] N=856	-0.027 [0.035] N=706
Vegetables consumed at least weekly	0.896 P=1,076	0.001 [0.032] N=1,204	-0.011 [0.031] N=1,108	-0.027 [0.032] N=992	-0.034 [0.035] N=856	-0.060 [0.041] N=706
Meat fish or poultry consumed at least weekly	0.756 P=917	0.052 [0.059] N=1,204	0.061 [0.062] N=1,108	0.062 [0.069] N=992	0.044 [0.078] N=856	0.042 [0.087] N=706
Legumes consumed at least weekly	0.758 P=918	-0.053 [0.048] N=1,204	-0.081 [0.050] N=1,108	-0.067 [0.052] N=992	-0.085 [0.058] N=856	-0.084 [0.070] N=706
Animal products consumed at least weekly	0.902 P=1,090	0.014 [0.038] N=1,204	0.006 [0.040] N=1,108	-0.017 [0.043] N=992	-0.013 [0.047] N=856	-0.026 [0.054] N=706
Sweets consumed at least weekly	0.520 P=625	-0.042 [0.047] N=1,204	-0.045 [0.048] N=1,108	-0.042 [0.053] N=992	-0.058 [0.056] N=856	-0.018 [0.064] N=706
Snack foods consumed at least weekly	0.431 P=514	-0.013 [0.047] N=1,204	-0.025 [0.049] N=1,108	-0.014 [0.053] N=992	-0.014 [0.055] N=856	-0.010 [0.058] N=706
Packaged soft drinks consumed at least weekly	0.244 P=288	-0.014 [0.046] N=1,204	-0.037 [0.047] N=1,108	-0.049 [0.049] N=992	-0.039 [0.054] N=856	-0.059 [0.061] N=706
Grains consumed at least weekly	0.856 P=1,033	0.022 [0.037] N=1,204	0.024 [0.039] N=1,108	0.006 [0.040] N=992	0.012 [0.046] N=856	-0.007 [0.053] N=706
Child was ever breastfed	0.969 P=1,416	-0.012 [0.013] N=1,466	-0.009 [0.014] N=1,355	-0.011 [0.015] N=1,211	-0.018 [0.016] N=1,032	-0.018 [0.016] N=853

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.001 [0.045] N=1,015	0.024 [0.045] N=919	0.001 [0.051] N=803	-0.020 [0.061] N=667	-0.064 [0.084] N=517
Vegetables consumed at least weekly	0.019 [0.051] N=1,015	0.003 [0.050] N=919	-0.030 [0.055] N=803	-0.051 [0.066] N=667	-0.157 [0.099] N=517
Meat fish or poultry consumed at least weekly	0.015 [0.069] N=1,015	0.024 [0.071] N=919	0.017 [0.081] N=803	-0.077 [0.086] N=667	-0.147 [0.094] N=517
Legumes consumed at least weekly	-0.036 [0.072] N=1,015	-0.085 [0.075] N=919	-0.043 [0.078] N=803	-0.106 [0.092] N=667	-0.120 [0.131] N=517
Animal products consumed at least weekly	0.030 [0.052] N=1,015	0.020 [0.057] N=919	-0.026 [0.064] N=803	-0.019 [0.077] N=667	-0.057 [0.103] N=517
Sweets consumed at least weekly	-0.059 [0.064] N=1,015	-0.067 [0.067] N=919	-0.066 [0.081] N=803	**-0.142* [0.084] N=667**	-0.053 [0.119] N=517
Snack foods consumed at least weekly	-0.035 [0.065] N=1,015	-0.062 [0.070] N=919	-0.050 [0.082] N=803	-0.082 [0.095] N=667	-0.117 [0.117] N=517
Packaged soft drinks consumed at least weekly	-0.009 [0.065] N=1,015	-0.054 [0.070] N=919	-0.079 [0.075] N=803	-0.074 [0.090] N=667	-0.158 [0.126] N=517
Grains consumed at least weekly	0.058 [0.054] N=1,015	0.070 [0.058] N=919	0.039 [0.062] N=803	0.061 [0.075] N=667	0.022 [0.104] N=517
Child was ever breastfed	-0.001 [0.020] N=1,239	0.005 [0.023] N=1,128	0.002 [0.027] N=984	-0.017 [0.034] N=805	-0.004 [0.040] N=626

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,112	-0.013 [0.028] N=1,204	-0.004 [0.028] N=1,108	-0.015 [0.029] N=992	-0.019 [0.032] N=856	-0.036 [0.036] N=706
Vegetables consumed at least weekly	0.896 P=1,076	-0.005 [0.032] N=1,204	-0.019 [0.030] N=1,108	-0.035 [0.031] N=992	-0.038 [0.034] N=856	**-0.069* [0.040] N=706**
Meat fish or poultry consumed at least weekly	0.756 P=917	0.034 [0.056] N=1,204	0.038 [0.059] N=1,108	0.040 [0.066] N=992	0.032 [0.074] N=856	0.020 [0.083] N=706
Legumes consumed at least weekly	0.758 P=918	-0.059 [0.047] N=1,204	**-0.088* [0.049] N=1,108**	-0.072 [0.051] N=992	-0.086 [0.057] N=856	-0.087 [0.069] N=706
Animal products consumed at least weekly	0.902 P=1,090	0.010 [0.038] N=1,204	0.001 [0.040] N=1,108	-0.022 [0.043] N=992	-0.016 [0.048] N=856	-0.034 [0.054] N=706
Sweets consumed at least weekly	0.520 P=625	-0.042 [0.047] N=1,204	-0.048 [0.049] N=1,108	-0.044 [0.054] N=992	-0.058 [0.056] N=856	-0.023 [0.063] N=706
Snack foods consumed at least weekly	0.431 P=514	-0.008 [0.046] N=1,204	-0.022 [0.048] N=1,108	-0.013 [0.052] N=992	-0.011 [0.054] N=856	-0.009 [0.057] N=706
Packaged soft drinks consumed at least weekly	0.244 P=288	-0.003 [0.044] N=1,204	-0.025 [0.046] N=1,108	-0.035 [0.048] N=992	-0.031 [0.052] N=856	-0.048 [0.059] N=706
Grains consumed at least weekly	0.856 P=1,033	0.018 [0.037] N=1,204	0.021 [0.039] N=1,108	0.004 [0.040] N=992	0.011 [0.045] N=856	-0.007 [0.052] N=706
Child was ever breastfed	0.969 P=1,416	-0.013 [0.013] N=1,466	-0.012 [0.014] N=1,355	-0.014 [0.016] N=1,211	-0.019 [0.017] N=1,032	-0.021 [0.017] N=853

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.044] N=1,015	0.019 [0.044] N=919	0.000 [0.049] N=803	-0.002 [0.060] N=667	-0.054 [0.083] N=517
Vegetables consumed at least weekly	0.019 [0.050] N=1,015	0.000 [0.049] N=919	-0.029 [0.054] N=803	-0.030 [0.065] N=667	-0.140 [0.098] N=517
Meat fish or poultry consumed at least weekly	0.006 [0.064] N=1,015	0.009 [0.066] N=919	0.004 [0.078] N=803	-0.059 [0.082] N=667	-0.141 [0.089] N=517
Legumes consumed at least weekly	-0.040 [0.071] N=1,015	-0.089 [0.074] N=919	-0.044 [0.077] N=803	-0.095 [0.092] N=667	-0.110 [0.133] N=517
Animal products consumed at least weekly	0.032 [0.051] N=1,015	0.021 [0.056] N=919	-0.023 [0.064] N=803	-0.001 [0.078] N=667	-0.033 [0.101] N=517
Sweets consumed at least weekly	-0.056 [0.065] N=1,015	-0.065 [0.068] N=919	-0.059 [0.081] N=803	-0.135 [0.085] N=667	-0.052 [0.123] N=517
Snack foods consumed at least weekly	-0.027 [0.064] N=1,015	-0.057 [0.069] N=919	-0.040 [0.080] N=803	-0.075 [0.092] N=667	-0.117 [0.114] N=517
Packaged soft drinks consumed at least weekly	0.005 [0.063] N=1,015	-0.036 [0.068] N=919	-0.057 [0.075] N=803	-0.067 [0.090] N=667	-0.151 [0.124] N=517
Grains consumed at least weekly	0.055 [0.054] N=1,015	0.068 [0.058] N=919	0.038 [0.063] N=803	0.068 [0.075] N=667	0.021 [0.103] N=517
Child was ever breastfed	0.001 [0.020] N=1,239	0.006 [0.022] N=1,128	0.003 [0.027] N=984	-0.009 [0.034] N=805	-0.004 [0.040] N=626

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.926 P=1,112	-0.015 [0.028] N=1,204	-0.004 [0.028] N=1,108	-0.020 [0.030] N=992	-0.026 [0.032] N=856	-0.047 [0.037] N=706
Vegetables consumed at least weekly	0.896 P=1,076	-0.006 [0.032] N=1,204	-0.017 [0.031] N=1,108	-0.035 [0.032] N=992	-0.040 [0.035] N=856	-0.069 [0.042] N=706
Meat fish or poultry consumed at least weekly	0.756 P=917	0.028 [0.056] N=1,204	0.034 [0.059] N=1,108	0.029 [0.066] N=992	0.017 [0.074] N=856	-0.006 [0.084] N=706
Legumes consumed at least weekly	0.758 P=918	-0.063 [0.049] N=1,204	**-0.093* [0.051] N=1,108**	-0.082 [0.053] N=992	-0.096 [0.059] N=856	-0.105 [0.073] N=706
Animal products consumed at least weekly	0.902 P=1,090	0.006 [0.038] N=1,204	-0.003 [0.040] N=1,108	-0.027 [0.043] N=992	-0.023 [0.047] N=856	-0.045 [0.053] N=706
Sweets consumed at least weekly	0.520 P=625	-0.045 [0.047] N=1,204	-0.053 [0.049] N=1,108	-0.053 [0.054] N=992	-0.063 [0.056] N=856	-0.029 [0.064] N=706
Snack foods consumed at least weekly	0.431 P=514	-0.013 [0.047] N=1,204	-0.026 [0.049] N=1,108	-0.021 [0.053] N=992	-0.019 [0.055] N=856	-0.015 [0.058] N=706
Packaged soft drinks consumed at least weekly	0.244 P=288	-0.005 [0.044] N=1,204	-0.033 [0.046] N=1,108	-0.044 [0.048] N=992	-0.035 [0.052] N=856	-0.049 [0.059] N=706
Grains consumed at least weekly	0.856 P=1,033	0.021 [0.038] N=1,204	0.024 [0.039] N=1,108	0.007 [0.041] N=992	0.012 [0.046] N=856	-0.004 [0.055] N=706
Child was ever breastfed	0.969 P=1,416	-0.015 [0.014] N=1,466	-0.013 [0.015] N=1,355	-0.016 [0.016] N=1,211	-0.022 [0.017] N=1,032	-0.025 [0.018] N=853

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.001 [0.044] N=1,015	0.026 [0.045] N=919	0.000 [0.050] N=803	-0.010 [0.060] N=667	-0.078 [0.083] N=517
Vegetables consumed at least weekly	0.018 [0.050] N=1,015	0.003 [0.050] N=919	-0.034 [0.054] N=803	-0.045 [0.064] N=667	**-0.175* [0.098] N=517**
Meat fish or poultry consumed at least weekly	0.013 [0.065] N=1,015	0.022 [0.066] N=919	0.011 [0.077] N=803	-0.044 [0.082] N=667	**-0.162* [0.093] N=517**
Legumes consumed at least weekly	-0.034 [0.072] N=1,015	-0.083 [0.075] N=919	-0.041 [0.077] N=803	-0.084 [0.092] N=667	-0.107 [0.136] N=517
Animal products consumed at least weekly	0.032 [0.052] N=1,015	0.023 [0.057] N=919	-0.027 [0.064] N=803	-0.011 [0.077] N=667	-0.057 [0.107] N=517
Sweets consumed at least weekly	-0.047 [0.065] N=1,015	-0.061 [0.067] N=919	-0.056 [0.080] N=803	-0.124 [0.086] N=667	-0.053 [0.124] N=517
Snack foods consumed at least weekly	-0.024 [0.064] N=1,015	-0.056 [0.069] N=919	-0.040 [0.080] N=803	-0.079 [0.093] N=667	-0.125 [0.116] N=517
Packaged soft drinks consumed at least weekly	0.005 [0.062] N=1,015	-0.045 [0.066] N=919	-0.067 [0.072] N=803	-0.073 [0.086] N=667	-0.149 [0.122] N=517
Grains consumed at least weekly	0.058 [0.055] N=1,015	0.070 [0.058] N=919	0.039 [0.063] N=803	0.066 [0.076] N=667	0.005 [0.103] N=517
Child was ever breastfed	0.001 [0.020] N=1,239	0.007 [0.023] N=1,128	0.001 [0.027] N=984	-0.012 [0.035] N=805	-0.002 [0.042] N=626

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,019	-0.012 [0.028] N=1,103	-0.006 [0.028] N=1,020	-0.015 [0.030] N=913	-0.024 [0.032] N=790	-0.045 [0.035] N=656
Vegetables consumed at least weekly	0.898 P=987	-0.010 [0.033] N=1,103	-0.025 [0.032] N=1,020	-0.043 [0.033] N=913	-0.043 [0.036] N=790	-0.068 [0.041] N=656
Meat fish or poultry consumed at least weekly	0.758 P=840	0.046 [0.054] N=1,103	0.042 [0.057] N=1,020	0.037 [0.063] N=913	0.008 [0.071] N=790	-0.008 [0.079] N=656
Legumes consumed at least weekly	0.754 P=835	-0.076 [0.049] N=1,103	**-0.112** [0.051] N=1,020**	**-0.102* [0.052] N=913**	**-0.126** [0.060] N=790**	**-0.129* [0.072] N=656**
Animal products consumed at least weekly	0.905 P=1,001	0.019 [0.040] N=1,103	0.006 [0.042] N=1,020	-0.019 [0.044] N=913	-0.025 [0.049] N=790	-0.048 [0.056] N=656
Sweets consumed at least weekly	0.521 P=571	-0.021 [0.049] N=1,103	-0.033 [0.052] N=1,020	-0.033 [0.056] N=913	-0.048 [0.060] N=790	-0.004 [0.068] N=656
Snack foods consumed at least weekly	0.433 P=471	-0.017 [0.050] N=1,103	-0.029 [0.053] N=1,020	-0.021 [0.057] N=913	-0.025 [0.060] N=790	-0.005 [0.064] N=656
Packaged soft drinks consumed at least weekly	0.248 P=267	0.017 [0.045] N=1,103	-0.007 [0.047] N=1,020	-0.032 [0.049] N=913	-0.027 [0.051] N=790	-0.034 [0.058] N=656
Grains consumed at least weekly	0.858 P=947	0.013 [0.037] N=1,103	0.012 [0.039] N=1,020	-0.016 [0.040] N=913	-0.009 [0.046] N=790	-0.032 [0.053] N=656
Child was ever breastfed	0.969 P=1,298	-0.018 [0.014] N=1,343	-0.014 [0.015] N=1,246	-0.016 [0.016] N=1,111	-0.018 [0.018] N=952	-0.021 [0.018] N=792

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.046] N=927	0.015 [0.048] N=844	-0.008 [0.054] N=737	-0.030 [0.066] N=614	-0.090 [0.089] N=480
Vegetables consumed at least weekly	0.015 [0.053] N=927	-0.008 [0.054] N=844	-0.044 [0.058] N=737	-0.053 [0.070] N=614	-0.148 [0.097] N=480
Meat fish or poultry consumed at least weekly	0.038 [0.068] N=927	0.022 [0.071] N=844	0.013 [0.081] N=737	-0.095 [0.090] N=614	**-0.171* [0.097] N=480**
Legumes consumed at least weekly	-0.041 [0.074] N=927	-0.106 [0.077] N=844	-0.071 [0.078] N=737	-0.147 [0.095] N=614	-0.169 [0.131] N=480
Animal products consumed at least weekly	0.048 [0.056] N=927	0.029 [0.061] N=844	-0.021 [0.067] N=737	-0.028 [0.083] N=614	-0.085 [0.109] N=480
Sweets consumed at least weekly	-0.034 [0.068] N=927	-0.057 [0.073] N=844	-0.070 [0.087] N=737	-0.149 [0.093] N=614	-0.068 [0.129] N=480
Snack foods consumed at least weekly	-0.041 [0.069] N=927	-0.069 [0.074] N=844	-0.063 [0.086] N=737	-0.116 [0.100] N=614	-0.131 [0.122] N=480
Packaged soft drinks consumed at least weekly	0.021 [0.068] N=927	-0.027 [0.072] N=844	-0.079 [0.079] N=737	-0.089 [0.095] N=614	-0.150 [0.135] N=480
Grains consumed at least weekly	0.058 [0.054] N=927	0.063 [0.059] N=844	0.009 [0.062] N=737	0.035 [0.076] N=614	-0.018 [0.104] N=480
Child was ever breastfed	-0.010 [0.022] N=1,132	-0.001 [0.025] N=1,035	-0.004 [0.029] N=900	-0.015 [0.035] N=741	-0.009 [0.043] N=581

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.927 P=1,019	-0.014 [0.028] N=1,103	-0.008 [0.028] N=1,020	-0.016 [0.029] N=913	-0.024 [0.032] N=790	-0.045 [0.034] N=656
Vegetables consumed at least weekly	0.898 P=987	-0.011 [0.033] N=1,103	-0.026 [0.032] N=1,020	-0.043 [0.033] N=913	-0.043 [0.036] N=790	**-0.070* [0.041] N=656**
Meat fish or poultry consumed at least weekly	0.758 P=840	0.043 [0.054] N=1,103	0.040 [0.058] N=1,020	0.036 [0.063] N=913	0.009 [0.071] N=790	-0.006 [0.079] N=656
Legumes consumed at least weekly	0.754 P=835	-0.076 [0.049] N=1,103	**-0.113** [0.051] N=1,020**	**-0.103* [0.052] N=913**	**-0.128** [0.060] N=790**	**-0.134* [0.072] N=656**
Animal products consumed at least weekly	0.905 P=1,001	0.018 [0.040] N=1,103	0.005 [0.041] N=1,020	-0.019 [0.044] N=913	-0.023 [0.048] N=790	-0.046 [0.055] N=656
Sweets consumed at least weekly	0.521 P=571	-0.020 [0.049] N=1,103	-0.033 [0.051] N=1,020	-0.031 [0.056] N=913	-0.045 [0.059] N=790	-0.003 [0.066] N=656
Snack foods consumed at least weekly	0.433 P=471	-0.018 [0.050] N=1,103	-0.030 [0.053] N=1,020	-0.020 [0.057] N=913	-0.025 [0.060] N=790	-0.005 [0.064] N=656
Packaged soft drinks consumed at least weekly	0.248 P=267	0.018 [0.046] N=1,103	-0.006 [0.047] N=1,020	-0.029 [0.049] N=913	-0.022 [0.052] N=790	-0.029 [0.058] N=656
Grains consumed at least weekly	0.858 P=947	0.013 [0.037] N=1,103	0.013 [0.039] N=1,020	-0.016 [0.040] N=913	-0.009 [0.046] N=790	-0.032 [0.053] N=656
Child was ever breastfed	0.969 P=1,298	-0.017 [0.014] N=1,343	-0.013 [0.015] N=1,246	-0.015 [0.016] N=1,111	-0.018 [0.017] N=952	-0.021 [0.018] N=792

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.045] N=927	0.010 [0.047] N=844	-0.007 [0.054] N=737	-0.026 [0.065] N=614	-0.087 [0.085] N=480
Vegetables consumed at least weekly	0.010 [0.052] N=927	-0.012 [0.054] N=844	-0.046 [0.057] N=737	-0.055 [0.069] N=614	-0.157 [0.096] N=480
Meat fish or poultry consumed at least weekly	0.035 [0.069] N=927	0.022 [0.072] N=844	0.014 [0.081] N=737	-0.094 [0.091] N=614	**-0.179* [0.099] N=480**
Legumes consumed at least weekly	-0.045 [0.073] N=927	-0.110 [0.077] N=844	-0.071 [0.079] N=737	-0.147 [0.095] N=614	-0.175 [0.130] N=480
Animal products consumed at least weekly	0.043 [0.055] N=927	0.025 [0.060] N=844	-0.020 [0.066] N=737	-0.024 [0.080] N=614	-0.081 [0.105] N=480
Sweets consumed at least weekly	-0.042 [0.068] N=927	-0.069 [0.073] N=844	-0.078 [0.087] N=737	**-0.152* [0.091] N=614**	-0.076 [0.127] N=480
Snack foods consumed at least weekly	-0.049 [0.069] N=927	-0.080 [0.074] N=844	-0.070 [0.087] N=737	-0.119 [0.101] N=614	-0.139 [0.123] N=480
Packaged soft drinks consumed at least weekly	0.015 [0.069] N=927	-0.034 [0.073] N=844	-0.082 [0.081] N=737	-0.084 [0.097] N=614	-0.142 [0.136] N=480
Grains consumed at least weekly	0.057 [0.054] N=927	0.064 [0.058] N=844	0.012 [0.061] N=737	0.038 [0.075] N=614	-0.014 [0.104] N=480
Child was ever breastfed	-0.008 [0.021] N=1,132	0.001 [0.024] N=1,035	-0.002 [0.029] N=900	-0.014 [0.035] N=741	-0.012 [0.043] N=581

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=972	**0.058* [0.033] N=1,046**	**0.061* [0.034] N=944**	0.037 [0.037] N=821	0.011 [0.040] N=685	0.016 [0.046] N=559
Vegetables consumed at least weekly	0.899 P=939	0.040 [0.038] N=1,046	0.048 [0.039] N=944	0.042 [0.043] N=821	0.008 [0.047] N=685	0.006 [0.049] N=559
Meat fish or poultry consumed at least weekly	0.753 P=799	**0.113** [0.054] N=1,046**	**0.114** [0.055] N=944**	0.102 [0.063] N=821	**0.133** [0.066] N=685**	0.121 [0.077] N=559
Legumes consumed at least weekly	0.755 P=801	-0.001 [0.044] N=1,046	0.040 [0.047] N=944	0.020 [0.053] N=821	-0.022 [0.057] N=685	-0.000 [0.064] N=559
Animal products consumed at least weekly	0.903 P=949	0.026 [0.041] N=1,046	0.024 [0.043] N=944	0.003 [0.050] N=821	-0.010 [0.054] N=685	-0.007 [0.063] N=559
Sweets consumed at least weekly	0.526 P=554	-0.051 [0.062] N=1,046	-0.057 [0.065] N=944	-0.069 [0.070] N=821	-0.074 [0.072] N=685	-0.128 [0.084] N=559
Snack foods consumed at least weekly	0.432 P=453	-0.033 [0.059] N=1,046	-0.028 [0.063] N=944	0.001 [0.067] N=821	-0.021 [0.072] N=685	-0.063 [0.082] N=559
Packaged soft drinks consumed at least weekly	0.234 P=241	-0.065 [0.049] N=1,046	-0.076 [0.051] N=944	-0.027 [0.055] N=821	-0.071 [0.059] N=685	-0.104 [0.065] N=559
Grains consumed at least weekly	0.849 P=894	0.052 [0.045] N=1,046	0.065 [0.048] N=944	0.055 [0.052] N=821	0.062 [0.058] N=685	0.038 [0.070] N=559
Child was ever breastfed	0.967 P=1,220	-0.005 [0.023] N=1,263	-0.009 [0.023] N=1,143	0.003 [0.022] N=993	0.003 [0.023] N=834	0.016 [0.022] N=676

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.059 [0.052] N=911	0.063 [0.054] N=809	0.019 [0.061] N=686	-0.002 [0.068] N=550	0.007 [0.093] N=424
Vegetables consumed at least weekly	0.015 [0.060] N=911	0.024 [0.065] N=809	0.002 [0.075] N=686	-0.060 [0.087] N=550	-0.083 [0.106] N=424
Meat fish or poultry consumed at least weekly	0.052 [0.072] N=911	0.048 [0.074] N=809	-0.000 [0.087] N=686	0.042 [0.094] N=550	-0.021 [0.126] N=424
Legumes consumed at least weekly	-0.022 [0.057] N=911	0.041 [0.062] N=809	0.014 [0.077] N=686	-0.058 [0.087] N=550	0.011 [0.101] N=424
Animal products consumed at least weekly	-0.011 [0.055] N=911	-0.015 [0.059] N=809	-0.058 [0.071] N=686	-0.085 [0.074] N=550	-0.110 [0.094] N=424
Sweets consumed at least weekly	-0.037 [0.085] N=911	-0.041 [0.090] N=809	-0.061 [0.104] N=686	-0.078 [0.120] N=550	**-0.255* [0.151] N=424**
Snack foods consumed at least weekly	-0.062 [0.080] N=911	-0.056 [0.084] N=809	-0.008 [0.097] N=686	-0.049 [0.112] N=550	-0.195 [0.130] N=424
Packaged soft drinks consumed at least weekly	-0.074 [0.075] N=911	-0.084 [0.080] N=809	0.013 [0.094] N=686	-0.061 [0.108] N=550	-0.121 [0.128] N=424
Grains consumed at least weekly	-0.038 [0.051] N=911	-0.034 [0.059] N=809	-0.075 [0.063] N=686	**-0.122* [0.071] N=550**	**-0.283*** [0.089] N=424**
Child was ever breastfed	0.009 [0.041] N=1,099	0.002 [0.043] N=979	0.024 [0.045] N=829	0.029 [0.052] N=670	0.076 [0.066] N=512

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=972	**0.058* [0.033] N=1,046**	**0.061* [0.034] N=944**	0.037 [0.038] N=821	0.011 [0.040] N=685	0.016 [0.046] N=559
Vegetables consumed at least weekly	0.899 P=939	0.040 [0.038] N=1,046	0.049 [0.039] N=944	0.042 [0.043] N=821	0.009 [0.046] N=685	0.006 [0.049] N=559
Meat fish or poultry consumed at least weekly	0.753 P=799	**0.113** [0.054] N=1,046**	**0.115** [0.055] N=944**	0.103 [0.063] N=821	**0.134** [0.066] N=685**	0.121 [0.077] N=559
Legumes consumed at least weekly	0.755 P=801	-0.000 [0.045] N=1,046	0.041 [0.047] N=944	0.020 [0.053] N=821	-0.022 [0.057] N=685	-0.001 [0.063] N=559
Animal products consumed at least weekly	0.903 P=949	0.027 [0.040] N=1,046	0.026 [0.043] N=944	0.004 [0.050] N=821	-0.009 [0.054] N=685	-0.006 [0.063] N=559
Sweets consumed at least weekly	0.526 P=554	-0.049 [0.062] N=1,046	-0.055 [0.065] N=944	-0.066 [0.070] N=821	-0.073 [0.072] N=685	-0.128 [0.084] N=559
Snack foods consumed at least weekly	0.432 P=453	-0.031 [0.058] N=1,046	-0.026 [0.062] N=944	0.004 [0.066] N=821	-0.019 [0.071] N=685	-0.062 [0.080] N=559
Packaged soft drinks consumed at least weekly	0.234 P=241	-0.064 [0.049] N=1,046	-0.076 [0.051] N=944	-0.026 [0.055] N=821	-0.071 [0.059] N=685	-0.103 [0.065] N=559
Grains consumed at least weekly	0.849 P=894	0.052 [0.045] N=1,046	0.066 [0.048] N=944	0.056 [0.052] N=821	0.062 [0.058] N=685	0.038 [0.070] N=559
Child was ever breastfed	0.967 P=1,220	-0.005 [0.023] N=1,263	-0.009 [0.023] N=1,143	0.002 [0.022] N=993	0.003 [0.023] N=834	0.015 [0.022] N=676

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.051] N=911	0.064 [0.053] N=809	0.019 [0.062] N=686	-0.002 [0.069] N=550	0.006 [0.095] N=424
Vegetables consumed at least weekly	0.016 [0.060] N=911	0.025 [0.065] N=809	0.002 [0.075] N=686	-0.055 [0.086] N=550	-0.081 [0.106] N=424
Meat fish or poultry consumed at least weekly	0.054 [0.071] N=911	0.052 [0.073] N=809	0.005 [0.085] N=686	0.050 [0.092] N=550	-0.015 [0.124] N=424
Legumes consumed at least weekly	-0.022 [0.057] N=911	0.043 [0.062] N=809	0.014 [0.077] N=686	-0.058 [0.087] N=550	0.003 [0.100] N=424
Animal products consumed at least weekly	-0.008 [0.051] N=911	-0.010 [0.056] N=809	-0.054 [0.069] N=686	-0.076 [0.071] N=550	-0.099 [0.089] N=424
Sweets consumed at least weekly	-0.033 [0.083] N=911	-0.035 [0.089] N=809	-0.055 [0.102] N=686	-0.076 [0.120] N=550	-0.250 [0.150] N=424
Snack foods consumed at least weekly	-0.057 [0.078] N=911	-0.049 [0.082] N=809	-0.000 [0.094] N=686	-0.041 [0.111] N=550	-0.185 [0.127] N=424
Packaged soft drinks consumed at least weekly	-0.072 [0.074] N=911	-0.083 [0.079] N=809	0.016 [0.093] N=686	-0.059 [0.108] N=550	-0.114 [0.125] N=424
Grains consumed at least weekly	-0.036 [0.051] N=911	-0.032 [0.059] N=809	-0.074 [0.064] N=686	**-0.125* [0.072] N=550**	**-0.286*** [0.091] N=424**
Child was ever breastfed	0.009 [0.041] N=1,099	0.002 [0.043] N=979	0.023 [0.046] N=829	0.029 [0.053] N=670	0.076 [0.066] N=512

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=972	0.052 [0.034] N=1,046	0.054 [0.034] N=944	0.034 [0.038] N=821	0.009 [0.041] N=685	0.014 [0.048] N=559
Vegetables consumed at least weekly	0.899 P=939	0.034 [0.038] N=1,046	0.041 [0.039] N=944	0.037 [0.044] N=821	0.004 [0.048] N=685	0.001 [0.051] N=559
Meat fish or poultry consumed at least weekly	0.753 P=799	**0.097* [0.054] N=1,046**	**0.096* [0.055] N=944**	0.097 [0.063] N=821	**0.129* [0.066] N=685**	0.117 [0.076] N=559
Legumes consumed at least weekly	0.755 P=801	-0.007 [0.046] N=1,046	0.033 [0.049] N=944	0.016 [0.056] N=821	-0.026 [0.060] N=685	-0.000 [0.068] N=559
Animal products consumed at least weekly	0.903 P=949	0.020 [0.041] N=1,046	0.016 [0.043] N=944	-0.002 [0.050] N=821	-0.013 [0.055] N=685	-0.011 [0.064] N=559
Sweets consumed at least weekly	0.526 P=554	-0.050 [0.062] N=1,046	-0.058 [0.065] N=944	-0.071 [0.070] N=821	-0.076 [0.073] N=685	-0.135 [0.085] N=559
Snack foods consumed at least weekly	0.432 P=453	-0.030 [0.058] N=1,046	-0.027 [0.062] N=944	-0.002 [0.066] N=821	-0.024 [0.072] N=685	-0.072 [0.082] N=559
Packaged soft drinks consumed at least weekly	0.234 P=241	-0.058 [0.048] N=1,046	-0.071 [0.050] N=944	-0.027 [0.054] N=821	-0.068 [0.058] N=685	-0.104 [0.065] N=559
Grains consumed at least weekly	0.849 P=894	0.050 [0.045] N=1,046	0.062 [0.049] N=944	0.054 [0.053] N=821	0.061 [0.060] N=685	0.038 [0.072] N=559
Child was ever breastfed	0.967 P=1,220	-0.007 [0.023] N=1,263	-0.011 [0.024] N=1,143	0.002 [0.023] N=993	0.003 [0.024] N=834	0.015 [0.023] N=676

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.051 [0.050] N=911	0.053 [0.052] N=809	0.019 [0.060] N=686	-0.004 [0.070] N=550	0.001 [0.097] N=424
Vegetables consumed at least weekly	0.009 [0.059] N=911	0.015 [0.063] N=809	-0.002 [0.072] N=686	-0.060 [0.085] N=550	-0.093 [0.105] N=424
Meat fish or poultry consumed at least weekly	0.029 [0.069] N=911	0.022 [0.071] N=809	-0.003 [0.083] N=686	0.035 [0.095] N=550	-0.046 [0.122] N=424
Legumes consumed at least weekly	-0.028 [0.058] N=911	0.035 [0.064] N=809	0.013 [0.077] N=686	-0.062 [0.088] N=550	0.007 [0.103] N=424
Animal products consumed at least weekly	-0.013 [0.051] N=911	-0.019 [0.056] N=809	-0.054 [0.068] N=686	-0.076 [0.071] N=550	-0.102 [0.090] N=424
Sweets consumed at least weekly	-0.028 [0.081] N=911	-0.033 [0.087] N=809	-0.051 [0.099] N=686	-0.068 [0.117] N=550	-0.234 [0.144] N=424
Snack foods consumed at least weekly	-0.048 [0.076] N=911	-0.043 [0.081] N=809	0.000 [0.092] N=686	-0.038 [0.108] N=550	-0.180 [0.122] N=424
Packaged soft drinks consumed at least weekly	-0.054 [0.073] N=911	-0.065 [0.079] N=809	0.024 [0.092] N=686	-0.045 [0.105] N=550	-0.092 [0.124] N=424
Grains consumed at least weekly	-0.039 [0.052] N=911	-0.036 [0.059] N=809	-0.073 [0.064] N=686	**-0.125* [0.072] N=550**	**-0.284*** [0.090] N=424**
Child was ever breastfed	0.006 [0.040] N=1,099	-0.002 [0.043] N=979	0.023 [0.045] N=829	0.027 [0.051] N=670	0.071 [0.064] N=512

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=972	0.047 [0.033] N=1,046	0.050 [0.034] N=944	0.031 [0.038] N=821	-0.002 [0.040] N=685	-0.001 [0.046] N=559
Vegetables consumed at least weekly	0.899 P=939	0.031 [0.038] N=1,046	0.039 [0.040] N=944	0.034 [0.044] N=821	-0.005 [0.046] N=685	-0.009 [0.049] N=559
Meat fish or poultry consumed at least weekly	0.753 P=799	**0.095* [0.054] N=1,046**	**0.093* [0.056] N=944**	0.087 [0.063] N=821	0.105 [0.066] N=685	0.079 [0.074] N=559
Legumes consumed at least weekly	0.755 P=801	-0.003 [0.046] N=1,046	0.037 [0.049] N=944	0.023 [0.055] N=821	-0.026 [0.057] N=685	-0.010 [0.064] N=559
Animal products consumed at least weekly	0.903 P=949	0.027 [0.039] N=1,046	0.025 [0.042] N=944	0.011 [0.049] N=821	-0.007 [0.054] N=685	-0.011 [0.062] N=559
Sweets consumed at least weekly	0.526 P=554	-0.040 [0.063] N=1,046	-0.046 [0.066] N=944	-0.060 [0.072] N=821	-0.068 [0.074] N=685	-0.126 [0.086] N=559
Snack foods consumed at least weekly	0.432 P=453	-0.021 [0.059] N=1,046	-0.016 [0.063] N=944	0.012 [0.068] N=821	-0.014 [0.072] N=685	-0.058 [0.082] N=559
Packaged soft drinks consumed at least weekly	0.234 P=241	-0.047 [0.048] N=1,046	-0.060 [0.050] N=944	-0.017 [0.055] N=821	-0.058 [0.059] N=685	-0.085 [0.067] N=559
Grains consumed at least weekly	0.849 P=894	0.048 [0.045] N=1,046	0.060 [0.049] N=944	0.051 [0.053] N=821	0.056 [0.059] N=685	0.031 [0.070] N=559
Child was ever breastfed	0.967 P=1,220	-0.006 [0.022] N=1,263	-0.011 [0.022] N=1,143	0.003 [0.021] N=993	0.002 [0.022] N=834	0.014 [0.021] N=676

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.048 [0.050] N=911	0.051 [0.052] N=809	0.017 [0.061] N=686	-0.024 [0.070] N=550	-0.036 [0.094] N=424
Vegetables consumed at least weekly	0.003 [0.060] N=911	0.009 [0.065] N=809	-0.008 [0.074] N=686	-0.079 [0.085] N=550	-0.128 [0.105] N=424
Meat fish or poultry consumed at least weekly	0.036 [0.073] N=911	0.029 [0.076] N=809	-0.005 [0.091] N=686	0.006 [0.106] N=550	-0.110 [0.129] N=424
Legumes consumed at least weekly	-0.022 [0.058] N=911	0.041 [0.065] N=809	0.034 [0.081] N=686	-0.054 [0.090] N=550	-0.005 [0.106] N=424
Animal products consumed at least weekly	-0.007 [0.049] N=911	-0.008 [0.054] N=809	-0.036 [0.067] N=686	-0.070 [0.071] N=550	-0.104 [0.089] N=424
Sweets consumed at least weekly	-0.014 [0.082] N=911	-0.015 [0.088] N=809	-0.032 [0.100] N=686	-0.053 [0.117] N=550	-0.219 [0.146] N=424
Snack foods consumed at least weekly	-0.042 [0.076] N=911	-0.034 [0.083] N=809	0.018 [0.094] N=686	-0.031 [0.108] N=550	-0.167 [0.123] N=424
Packaged soft drinks consumed at least weekly	-0.052 [0.072] N=911	-0.063 [0.077] N=809	0.024 [0.092] N=686	-0.040 [0.106] N=550	-0.061 [0.128] N=424
Grains consumed at least weekly	-0.037 [0.051] N=911	-0.035 [0.059] N=809	-0.075 [0.065] N=686	**-0.134* [0.073] N=550**	**-0.299*** [0.089] N=424**
Child was ever breastfed	0.007 [0.040] N=1,099	-0.002 [0.041] N=979	0.025 [0.043] N=829	0.024 [0.049] N=670	0.064 [0.060] N=512

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=889	0.058 [0.036] N=955	0.060 [0.036] N=859	0.049 [0.040] N=754	0.008 [0.044] N=628	0.009 [0.051] N=517
Vegetables consumed at least weekly	0.903 P=861	0.030 [0.038] N=955	0.036 [0.041] N=859	0.051 [0.044] N=754	0.004 [0.050] N=628	-0.021 [0.053] N=517
Meat fish or poultry consumed at least weekly	0.755 P=729	**0.106* [0.054] N=955**	**0.098* [0.056] N=859**	0.100 [0.063] N=754	0.108 [0.068] N=628	0.093 [0.078] N=517
Legumes consumed at least weekly	0.753 P=729	-0.001 [0.050] N=955	0.040 [0.053] N=859	0.025 [0.058] N=754	-0.032 [0.064] N=628	-0.017 [0.074] N=517
Animal products consumed at least weekly	0.907 P=870	0.023 [0.043] N=955	0.023 [0.046] N=859	0.018 [0.053] N=754	-0.004 [0.059] N=628	-0.011 [0.068] N=517
Sweets consumed at least weekly	0.537 P=515	-0.045 [0.064] N=955	-0.048 [0.067] N=859	-0.071 [0.073] N=754	-0.088 [0.077] N=628	-0.141 [0.089] N=517
Snack foods consumed at least weekly	0.441 P=420	-0.022 [0.061] N=955	-0.017 [0.065] N=859	0.003 [0.068] N=754	-0.023 [0.074] N=628	-0.057 [0.084] N=517
Packaged soft drinks consumed at least weekly	0.241 P=227	-0.075 [0.050] N=955	**-0.090* [0.052] N=859**	-0.042 [0.055] N=754	-0.095 [0.061] N=628	**-0.139** [0.069] N=517**
Grains consumed at least weekly	0.851 P=818	0.069 [0.047] N=955	0.079 [0.051] N=859	0.080 [0.054] N=754	0.076 [0.063] N=628	0.040 [0.074] N=517
Child was ever breastfed	0.969 P=1,118	-0.014 [0.022] N=1,154	-0.019 [0.022] N=1,043	-0.006 [0.020] N=913	-0.011 [0.021] N=767	0.007 [0.021] N=626

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.048 [0.052] N=830	0.050 [0.053] N=734	0.032 [0.061] N=629	-0.033 [0.069] N=503	-0.061 [0.086] N=392
Vegetables consumed at least weekly	-0.006 [0.053] N=830	-0.004 [0.059] N=734	0.010 [0.065] N=629	-0.087 [0.080] N=503	**-0.186** [0.091] N=392**
Meat fish or poultry consumed at least weekly	0.069 [0.079] N=830	0.057 [0.081] N=734	0.040 [0.092] N=629	0.014 [0.103] N=503	-0.073 [0.129] N=392
Legumes consumed at least weekly	-0.014 [0.064] N=830	0.051 [0.068] N=734	0.047 [0.080] N=629	-0.036 [0.097] N=503	0.012 [0.113] N=392
Animal products consumed at least weekly	-0.026 [0.052] N=830	-0.028 [0.057] N=734	-0.041 [0.069] N=629	-0.090 [0.076] N=503	-0.147 [0.095] N=392
Sweets consumed at least weekly	-0.028 [0.086] N=830	-0.028 [0.093] N=734	-0.079 [0.105] N=629	-0.135 [0.123] N=503	**-0.326** [0.159] N=392**
Snack foods consumed at least weekly	-0.052 [0.082] N=830	-0.045 [0.088] N=734	-0.021 [0.096] N=629	-0.086 [0.110] N=503	-0.203 [0.134] N=392
Packaged soft drinks consumed at least weekly	-0.088 [0.076] N=830	-0.103 [0.082] N=734	-0.015 [0.096] N=629	-0.118 [0.115] N=503	**-0.239* [0.141] N=392**
Grains consumed at least weekly	-0.016 [0.057] N=830	-0.015 [0.064] N=734	-0.031 [0.070] N=629	-0.114 [0.083] N=503	**-0.325*** [0.100] N=392**
Child was ever breastfed	-0.004 [0.038] N=1,002	-0.013 [0.040] N=891	0.010 [0.040] N=761	0.000 [0.047] N=615	0.049 [0.057] N=474

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=889	**0.059* [0.035] N=955**	**0.062* [0.036] N=859**	0.050 [0.040] N=754	0.008 [0.043] N=628	0.008 [0.050] N=517
Vegetables consumed at least weekly	0.903 P=861	0.034 [0.038] N=955	0.039 [0.040] N=859	0.054 [0.043] N=754	0.003 [0.048] N=628	-0.021 [0.051] N=517
Meat fish or poultry consumed at least weekly	0.755 P=729	**0.112** [0.053] N=955**	**0.106* [0.055] N=859**	**0.105* [0.061] N=754**	0.104 [0.065] N=628	0.090 [0.075] N=517
Legumes consumed at least weekly	0.753 P=729	-0.001 [0.049] N=955	0.040 [0.052] N=859	0.026 [0.057] N=754	-0.036 [0.062] N=628	-0.022 [0.072] N=517
Animal products consumed at least weekly	0.907 P=870	0.023 [0.042] N=955	0.023 [0.045] N=859	0.018 [0.052] N=754	-0.003 [0.058] N=628	-0.009 [0.067] N=517
Sweets consumed at least weekly	0.537 P=515	-0.048 [0.063] N=955	-0.051 [0.066] N=859	-0.075 [0.072] N=754	-0.094 [0.076] N=628	-0.145 [0.090] N=517
Snack foods consumed at least weekly	0.441 P=420	-0.019 [0.060] N=955	-0.015 [0.064] N=859	0.006 [0.068] N=754	-0.022 [0.073] N=628	-0.055 [0.083] N=517
Packaged soft drinks consumed at least weekly	0.241 P=227	-0.075 [0.050] N=955	**-0.090* [0.052] N=859**	-0.044 [0.055] N=754	-0.098 [0.060] N=628	**-0.142** [0.068] N=517**
Grains consumed at least weekly	0.851 P=818	0.069 [0.047] N=955	0.079 [0.051] N=859	0.080 [0.054] N=754	0.076 [0.062] N=628	0.039 [0.074] N=517
Child was ever breastfed	0.969 P=1,118	-0.013 [0.022] N=1,154	-0.018 [0.023] N=1,043	-0.004 [0.021] N=913	-0.008 [0.022] N=767	0.010 [0.021] N=626

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.052 [0.051] N=830	0.055 [0.054] N=734	0.037 [0.062] N=629	-0.032 [0.071] N=503	-0.052 [0.092] N=392
Vegetables consumed at least weekly	-0.003 [0.053] N=830	-0.000 [0.059] N=734	0.016 [0.065] N=629	-0.088 [0.079] N=503	**-0.180* [0.092] N=392**
Meat fish or poultry consumed at least weekly	0.075 [0.078] N=830	0.062 [0.081] N=734	0.041 [0.091] N=629	0.009 [0.101] N=503	-0.062 [0.129] N=392
Legumes consumed at least weekly	-0.011 [0.063] N=830	0.056 [0.068] N=734	0.051 [0.079] N=629	-0.044 [0.095] N=503	0.020 [0.111] N=392
Animal products consumed at least weekly	-0.026 [0.052] N=830	-0.026 [0.057] N=734	-0.040 [0.069] N=629	-0.092 [0.076] N=503	-0.153 [0.098] N=392
Sweets consumed at least weekly	-0.026 [0.087] N=830	-0.027 [0.093] N=734	-0.076 [0.106] N=629	-0.128 [0.124] N=503	**-0.318** [0.159] N=392**
Snack foods consumed at least weekly	-0.048 [0.081] N=830	-0.042 [0.088] N=734	-0.010 [0.097] N=629	-0.065 [0.113] N=503	-0.189 [0.134] N=392
Packaged soft drinks consumed at least weekly	-0.084 [0.078] N=830	-0.099 [0.083] N=734	-0.005 [0.095] N=629	-0.100 [0.113] N=503	-0.211 [0.138] N=392
Grains consumed at least weekly	-0.017 [0.056] N=830	-0.017 [0.064] N=734	-0.035 [0.069] N=629	-0.122 [0.082] N=503	**-0.333*** [0.099] N=392**
Child was ever breastfed	-0.004 [0.038] N=1,002	-0.014 [0.041] N=891	0.012 [0.041] N=761	0.006 [0.049] N=615	0.053 [0.059] N=474

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.