Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Youngest Child

Age reference: 12 November 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.093 [0.239] N=1,245	0.089 [0.246] N=1,151	0.122 [0.275] N=1,027	0.258 [0.286] N=883	0.070 [0.319] N=735
WHO weight-for-age z score	-0.023	-0.191 [0.135] N=1,449	-0.164 [0.150] N=1,340	-0.125 [0.155] N=1,198	-0.087 [0.155] N=1,021	-0.153 [0.174] N=843
WHO weight-for-length/height z score, 0-60 months	0.305	0.246 [0.224] N=1,240	0.252 [0.231] N=1,147	0.310 [0.240] N=1,026	0.237 [0.248] N=879	0.288 [0.268] N=728
WHO BMI-for-age z score	0.283	0.199 [0.244] N=1,222	0.227 [0.253] N=1,132	0.272 [0.271] N=1,010	0.129 [0.281] N=863	0.140 [0.306] N=715
Stunted: HAZ below -2	0.246 P=295	-0.040 [0.047] N=1,245	-0.039 [0.050] N=1,151	-0.034 [0.052] N=1,027	-0.069 [0.056] N=883	-0.052 [0.059] N=735
Underweight: WAZ below -2	0.090 P=130	**0.077*** [0.027] N=1,449**	**0.063** [0.028] N=1,340**	**0.059** [0.029] N=1,198**	0.052 [0.032] N=1,021	0.045 [0.036] N=843
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=148	-0.040 [0.037] N=1,240	-0.027 [0.038] N=1,147	-0.041 [0.041] N=1,026	-0.025 [0.045] N=879	-0.033 [0.047] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=241	0.033 [0.047] N=1,240	0.041 [0.049] N=1,147	0.043 [0.051] N=1,026	0.037 [0.057] N=879	0.023 [0.064] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=111	-0.012 [0.035] N=1,240	-0.005 [0.038] N=1,147	0.014 [0.039] N=1,026	0.010 [0.043] N=879	0.025 [0.048] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	0.339 P=392	-0.059 [0.049] N=1,189	-0.045 [0.052] N=1,099	-0.044 [0.055] N=982	-0.068 [0.061] N=843	-0.055 [0.066] N=700
BMI-for-age z score below -2	0.131 P=159	-0.031 [0.040] N=1,222	-0.028 [0.042] N=1,132	-0.029 [0.047] N=1,010	-0.001 [0.050] N=863	0.006 [0.053] N=715

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.004 [0.360] N=1,046	-0.011 [0.384] N=952	0.124 [0.453] N=828	0.520 [0.484] N=684	0.243 [0.616] N=536
WHO weight-for-age z score	**-0.450** [0.191] N=1,224**	**-0.429* [0.229] N=1,115**	-0.389 [0.237] N=973	-0.389 [0.267] N=796	**-0.688* [0.397] N=618**
WHO weight-for-length/height z score, 0-60 months	0.190 [0.329] N=1,041	0.181 [0.353] N=948	0.234 [0.379] N=827	0.131 [0.435] N=680	0.074 [0.507] N=529
WHO BMI-for-age z score	0.240 [0.337] N=1,025	0.269 [0.360] N=935	0.292 [0.402] N=813	0.017 [0.442] N=666	-0.138 [0.518] N=518
Stunted: HAZ below -2	-0.019 [0.071] N=1,046	-0.019 [0.078] N=952	-0.020 [0.087] N=828	-0.098 [0.104] N=684	-0.104 [0.132] N=536
Underweight: WAZ below -2	**0.121*** [0.037] N=1,224**	**0.098** [0.041] N=1,115**	**0.097** [0.043] N=973**	**0.102** [0.051] N=796**	0.098 [0.073] N=618
Wasted: WLZ/WHZ below -2, 0-60 months	-0.063 [0.051] N=1,041	-0.039 [0.056] N=948	-0.071 [0.063] N=827	-0.058 [0.076] N=680	-0.097 [0.087] N=529
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.059] N=1,041	0.002 [0.064] N=948	-0.011 [0.064] N=827	-0.028 [0.081] N=680	-0.143 [0.095] N=529
Obese: WLZ/WHZ above +3, 0-60 months	-0.055 [0.044] N=1,041	-0.050 [0.051] N=948	-0.028 [0.051] N=827	-0.043 [0.063] N=680	-0.052 [0.080] N=529
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.031 [0.075] N=996	-0.007 [0.080] N=906	-0.009 [0.087] N=789	-0.063 [0.109] N=650	-0.058 [0.136] N=507
BMI-for-age z score below -2	-0.070 [0.051] N=1,025	-0.068 [0.054] N=935	-0.079 [0.064] N=813	-0.044 [0.073] N=666	-0.035 [0.084] N=518

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.093 [0.240] N=1,245	0.087 [0.246] N=1,151	0.122 [0.275] N=1,027	0.263 [0.287] N=883	0.079 [0.320] N=735
WHO weight-for-age z score	-0.023	-0.197 [0.134] N=1,449	-0.173 [0.149] N=1,340	-0.131 [0.154] N=1,198	-0.092 [0.154] N=1,021	-0.161 [0.173] N=843
WHO weight-for-length/height z score, 0-60 months	0.305	0.244 [0.224] N=1,240	0.250 [0.231] N=1,147	0.308 [0.241] N=1,026	0.233 [0.248] N=879	0.281 [0.270] N=728
WHO BMI-for-age z score	0.283	0.200 [0.244] N=1,222	0.227 [0.253] N=1,132	0.272 [0.271] N=1,010	0.129 [0.281] N=863	0.139 [0.306] N=715
Stunted: HAZ below -2	0.246 P=295	-0.038 [0.047] N=1,245	-0.037 [0.049] N=1,151	-0.032 [0.052] N=1,027	-0.067 [0.056] N=883	-0.049 [0.059] N=735
Underweight: WAZ below -2	0.090 P=130	**0.078*** [0.027] N=1,449**	**0.064** [0.028] N=1,340**	**0.059** [0.029] N=1,198**	**0.053* [0.032] N=1,021**	0.046 [0.036] N=843
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=148	-0.038 [0.036] N=1,240	-0.024 [0.038] N=1,147	-0.039 [0.041] N=1,026	-0.022 [0.045] N=879	-0.029 [0.047] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=241	0.034 [0.047] N=1,240	0.042 [0.049] N=1,147	0.044 [0.051] N=1,026	0.039 [0.056] N=879	0.025 [0.064] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=111	-0.011 [0.035] N=1,240	-0.003 [0.038] N=1,147	0.014 [0.039] N=1,026	0.012 [0.043] N=879	0.027 [0.048] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	0.339 P=392	-0.056 [0.048] N=1,189	-0.041 [0.050] N=1,099	-0.042 [0.054] N=982	-0.066 [0.060] N=843	-0.049 [0.066] N=700
BMI-for-age z score below -2	0.131 P=159	-0.030 [0.040] N=1,222	-0.026 [0.042] N=1,132	-0.028 [0.047] N=1,010	0.001 [0.050] N=863	0.008 [0.054] N=715

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.000 [0.359] N=1,046	-0.012 [0.384] N=952	0.119 [0.451] N=828	0.500 [0.484] N=684	0.226 [0.615] N=536
WHO weight-for-age z score	**-0.446** [0.190] N=1,224**	**-0.429* [0.229] N=1,115**	-0.384 [0.238] N=973	-0.387 [0.269] N=796	**-0.688* [0.397] N=618**
WHO weight-for-length/height z score, 0-60 months	0.190 [0.329] N=1,041	0.180 [0.353] N=948	0.237 [0.379] N=827	0.131 [0.436] N=680	0.072 [0.509] N=529
WHO BMI-for-age z score	0.241 [0.337] N=1,025	0.269 [0.360] N=935	0.297 [0.402] N=813	0.019 [0.443] N=666	-0.139 [0.519] N=518
Stunted: HAZ below -2	-0.021 [0.070] N=1,046	-0.022 [0.077] N=952	-0.025 [0.085] N=828	-0.101 [0.103] N=684	-0.105 [0.132] N=536
Underweight: WAZ below -2	**0.120*** [0.037] N=1,224**	**0.098** [0.041] N=1,115**	**0.097** [0.043] N=973**	**0.103** [0.051] N=796**	0.098 [0.073] N=618
Wasted: WLZ/WHZ below -2, 0-60 months	-0.063 [0.051] N=1,041	-0.039 [0.055] N=948	-0.073 [0.062] N=827	-0.058 [0.076] N=680	-0.095 [0.086] N=529
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.059] N=1,041	0.002 [0.064] N=948	-0.012 [0.064] N=827	-0.028 [0.081] N=680	-0.143 [0.095] N=529
Obese: WLZ/WHZ above +3, 0-60 months	-0.055 [0.044] N=1,041	-0.049 [0.050] N=948	-0.029 [0.051] N=827	-0.043 [0.063] N=680	-0.051 [0.080] N=529
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.036 [0.073] N=996	-0.011 [0.079] N=906	-0.020 [0.085] N=789	-0.074 [0.108] N=650	-0.065 [0.137] N=507
BMI-for-age z score below -2	-0.071 [0.051] N=1,025	-0.068 [0.054] N=935	-0.081 [0.064] N=813	-0.045 [0.073] N=666	-0.035 [0.085] N=518

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.075 [0.241] N=1,245	0.061 [0.247] N=1,151	0.081 [0.276] N=1,027	0.243 [0.285] N=883	0.043 [0.315] N=735
WHO weight-for-age z score	-0.023	**-0.223* [0.130] N=1,449**	-0.203 [0.145] N=1,340	-0.165 [0.149] N=1,198	-0.110 [0.150] N=1,021	-0.189 [0.168] N=843
WHO weight-for-length/height z score, 0-60 months	0.305	0.240 [0.225] N=1,240	0.252 [0.234] N=1,147	0.317 [0.243] N=1,026	0.238 [0.250] N=879	0.282 [0.275] N=728
WHO BMI-for-age z score	0.283	0.202 [0.246] N=1,222	0.232 [0.256] N=1,132	0.283 [0.275] N=1,010	0.136 [0.285] N=863	0.145 [0.315] N=715
Stunted: HAZ below -2	0.246 P=295	-0.035 [0.047] N=1,245	-0.030 [0.049] N=1,151	-0.026 [0.052] N=1,027	-0.065 [0.055] N=883	-0.039 [0.058] N=735
Underweight: WAZ below -2	0.090 P=130	**0.081*** [0.026] N=1,449**	**0.068** [0.028] N=1,340**	**0.064** [0.029] N=1,198**	**0.056* [0.031] N=1,021**	0.052 [0.036] N=843
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=148	-0.036 [0.036] N=1,240	-0.022 [0.038] N=1,147	-0.038 [0.041] N=1,026	-0.021 [0.045] N=879	-0.027 [0.047] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=241	0.035 [0.047] N=1,240	0.044 [0.049] N=1,147	0.046 [0.051] N=1,026	0.040 [0.056] N=879	0.029 [0.064] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=111	-0.011 [0.036] N=1,240	-0.002 [0.038] N=1,147	0.015 [0.040] N=1,026	0.012 [0.043] N=879	0.027 [0.048] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	0.339 P=392	-0.050 [0.048] N=1,189	-0.031 [0.050] N=1,099	-0.033 [0.053] N=982	-0.062 [0.059] N=843	-0.034 [0.066] N=700
BMI-for-age z score below -2	0.131 P=159	-0.030 [0.040] N=1,222	-0.025 [0.042] N=1,132	-0.028 [0.048] N=1,010	0.001 [0.050] N=863	0.008 [0.054] N=715

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.012 [0.359] N=1,046	-0.041 [0.380] N=952	0.051 [0.445] N=828	0.479 [0.468] N=684	0.235 [0.598] N=536
WHO weight-for-age z score	**-0.455** [0.187] N=1,224**	**-0.446* [0.226] N=1,115**	**-0.407* [0.234] N=973**	-0.373 [0.274] N=796	**-0.678* [0.398] N=618**
WHO weight-for-length/height z score, 0-60 months	0.187 [0.326] N=1,041	0.182 [0.353] N=948	0.248 [0.373] N=827	0.105 [0.426] N=680	0.028 [0.504] N=529
WHO BMI-for-age z score	0.244 [0.338] N=1,025	0.271 [0.361] N=935	0.311 [0.399] N=813	-0.007 [0.428] N=666	-0.160 [0.510] N=518
Stunted: HAZ below -2	-0.022 [0.070] N=1,046	-0.017 [0.076] N=952	-0.020 [0.084] N=828	-0.113 [0.098] N=684	-0.111 [0.123] N=536
Underweight: WAZ below -2	**0.119*** [0.037] N=1,224**	**0.099** [0.041] N=1,115**	**0.097** [0.043] N=973**	**0.094* [0.052] N=796**	0.097 [0.074] N=618
Wasted: WLZ/WHZ below -2, 0-60 months	-0.063 [0.051] N=1,041	-0.038 [0.056] N=948	-0.078 [0.063] N=827	-0.064 [0.077] N=680	-0.093 [0.088] N=529
Overweight: WLZ/WHZ above +2, 0-60 months	-0.007 [0.059] N=1,041	0.002 [0.065] N=948	-0.015 [0.064] N=827	-0.039 [0.081] N=680	-0.142 [0.096] N=529
Obese: WLZ/WHZ above +3, 0-60 months	-0.056 [0.044] N=1,041	-0.051 [0.051] N=948	-0.032 [0.051] N=827	-0.049 [0.064] N=680	-0.056 [0.080] N=529
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.039 [0.073] N=996	-0.009 [0.080] N=906	-0.024 [0.085] N=789	-0.100 [0.105] N=650	-0.074 [0.132] N=507
BMI-for-age z score below -2	-0.073 [0.051] N=1,025	-0.066 [0.055] N=935	-0.085 [0.065] N=813	-0.049 [0.074] N=666	-0.033 [0.085] N=518

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.065 [0.240] N=1,245	0.073 [0.245] N=1,151	0.095 [0.275] N=1,027	0.273 [0.286] N=883	0.097 [0.316] N=735
WHO weight-for-age z score	-0.023	**-0.244* [0.128] N=1,449**	-0.227 [0.142] N=1,340	-0.193 [0.149] N=1,198	-0.146 [0.148] N=1,021	-0.249 [0.171] N=843
WHO weight-for-length/height z score, 0-60 months	0.305	0.235 [0.227] N=1,240	0.237 [0.234] N=1,147	0.297 [0.244] N=1,026	0.207 [0.252] N=879	0.219 [0.276] N=728
WHO BMI-for-age z score	0.283	0.194 [0.246] N=1,222	0.218 [0.254] N=1,132	0.269 [0.273] N=1,010	0.109 [0.284] N=863	0.082 [0.314] N=715
Stunted: HAZ below -2	0.246 P=295	-0.030 [0.048] N=1,245	-0.028 [0.049] N=1,151	-0.020 [0.052] N=1,027	-0.059 [0.056] N=883	-0.034 [0.058] N=735
Underweight: WAZ below -2	0.090 P=130	**0.085*** [0.026] N=1,449**	**0.073*** [0.028] N=1,340**	**0.070** [0.029] N=1,198**	**0.065** [0.031] N=1,021**	**0.065* [0.036] N=843**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=148	-0.036 [0.036] N=1,240	-0.021 [0.038] N=1,147	-0.034 [0.041] N=1,026	-0.018 [0.044] N=879	-0.024 [0.046] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=241	0.035 [0.047] N=1,240	0.044 [0.049] N=1,147	0.049 [0.051] N=1,026	0.041 [0.057] N=879	0.023 [0.065] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=111	-0.012 [0.035] N=1,240	-0.006 [0.038] N=1,147	0.012 [0.039] N=1,026	0.008 [0.043] N=879	0.017 [0.048] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	0.339 P=392	-0.041 [0.049] N=1,189	-0.021 [0.051] N=1,099	-0.015 [0.054] N=982	-0.042 [0.061] N=843	-0.014 [0.068] N=700
BMI-for-age z score below -2	0.131 P=159	-0.027 [0.040] N=1,222	-0.022 [0.042] N=1,132	-0.022 [0.047] N=1,010	0.008 [0.050] N=863	0.017 [0.054] N=715

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.051 [0.364] N=1,046	-0.049 [0.384] N=952	0.035 [0.451] N=828	0.488 [0.488] N=684	0.206 [0.613] N=536
WHO weight-for-age z score	**-0.463** [0.188] N=1,224**	**-0.447** [0.224] N=1,115**	**-0.412* [0.233] N=973**	-0.331 [0.272] N=796	-0.617 [0.388] N=618
WHO weight-for-length/height z score, 0-60 months	0.214 [0.327] N=1,041	0.207 [0.351] N=948	0.265 [0.381] N=827	0.159 [0.437] N=680	0.086 [0.511] N=529
WHO BMI-for-age z score	0.265 [0.337] N=1,025	0.293 [0.356] N=935	0.335 [0.406] N=813	0.064 [0.442] N=666	-0.085 [0.517] N=518
Stunted: HAZ below -2	-0.014 [0.070] N=1,046	-0.017 [0.076] N=952	-0.015 [0.085] N=828	-0.112 [0.099] N=684	-0.108 [0.125] N=536
Underweight: WAZ below -2	**0.120*** [0.037] N=1,224**	**0.098** [0.041] N=1,115**	**0.098** [0.043] N=973**	**0.088* [0.050] N=796**	0.097 [0.070] N=618
Wasted: WLZ/WHZ below -2, 0-60 months	-0.063 [0.051] N=1,041	-0.042 [0.056] N=948	-0.076 [0.063] N=827	-0.059 [0.078] N=680	-0.089 [0.090] N=529
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.059] N=1,041	0.003 [0.064] N=948	-0.011 [0.064] N=827	-0.030 [0.081] N=680	-0.127 [0.094] N=529
Obese: WLZ/WHZ above +3, 0-60 months	-0.050 [0.044] N=1,041	-0.045 [0.050] N=948	-0.025 [0.051] N=827	-0.042 [0.062] N=680	-0.052 [0.076] N=529
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.033 [0.075] N=996	-0.009 [0.081] N=906	-0.016 [0.086] N=789	-0.101 [0.107] N=650	-0.074 [0.138] N=507
BMI-for-age z score below -2	-0.075 [0.051] N=1,025	-0.073 [0.054] N=935	-0.088 [0.066] N=813	-0.057 [0.075] N=666	-0.045 [0.085] N=518

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.098 [0.241] N=1,144	0.056 [0.255] N=1,063	0.064 [0.284] N=946	0.147 [0.292] N=820	-0.097 [0.317] N=685
WHO weight-for-age z score	-0.016	-0.105 [0.136] N=1,327	-0.096 [0.149] N=1,232	-0.041 [0.155] N=1,099	0.001 [0.153] N=942	-0.049 [0.171] N=783
WHO weight-for-length/height z score, 0-60 months	0.317	**0.396* [0.223] N=1,143**	**0.399* [0.228] N=1,063**	**0.474** [0.234] N=950**	**0.414* [0.237] N=821**	**0.507** [0.253] N=683**
WHO BMI-for-age z score	0.303	0.343 [0.246] N=1,126	0.351 [0.251] N=1,048	0.426 [0.264] N=935	0.308 [0.269] N=806	0.363 [0.291] N=671
Stunted: HAZ below -2	0.247 P=271	-0.037 [0.050] N=1,144	-0.034 [0.053] N=1,063	-0.025 [0.056] N=946	-0.052 [0.060] N=820	-0.020 [0.063] N=685
Underweight: WAZ below -2	0.089 P=119	**0.070** [0.028] N=1,327**	**0.058* [0.030] N=1,232**	**0.053* [0.031] N=1,099**	0.047 [0.035] N=942	0.043 [0.039] N=783
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=135	**-0.066* [0.036] N=1,143**	-0.049 [0.037] N=1,063	-0.061 [0.040] N=950	-0.044 [0.042] N=821	-0.052 [0.044] N=683
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=223	0.051 [0.047] N=1,143	0.060 [0.049] N=1,063	0.068 [0.051] N=950	0.064 [0.057] N=821	0.063 [0.065] N=683
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=106	-0.012 [0.038] N=1,143	-0.001 [0.040] N=1,063	0.023 [0.042] N=950	0.025 [0.045] N=821	0.047 [0.049] N=683
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=361	-0.076 [0.053] N=1,094	-0.054 [0.055] N=1,017	-0.055 [0.059] N=907	-0.080 [0.065] N=786	-0.059 [0.071] N=655
BMI-for-age z score below -2	0.129 P=142	-0.044 [0.040] N=1,126	-0.036 [0.042] N=1,048	-0.040 [0.047] N=935	-0.014 [0.049] N=806	-0.016 [0.053] N=671

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.214 [0.377] N=958	0.179 [0.421] N=877	0.319 [0.497] N=760	0.619 [0.539] N=634	0.339 [0.700] N=499
WHO weight-for-age z score	**-0.378** [0.191] N=1,118**	**-0.395* [0.225] N=1,023**	-0.339 [0.233] N=890	-0.337 [0.257] N=733	-0.580 [0.381] N=574
WHO weight-for-length/height z score, 0-60 months	0.228 [0.352] N=957	0.177 [0.373] N=877	0.228 [0.402] N=764	0.185 [0.460] N=635	0.249 [0.537] N=497
WHO BMI-for-age z score	0.231 [0.368] N=942	0.188 [0.385] N=864	0.231 [0.420] N=751	0.019 [0.459] N=622	-0.123 [0.543] N=487
Stunted: HAZ below -2	-0.065 [0.075] N=958	-0.073 [0.082] N=877	-0.079 [0.091] N=760	-0.158 [0.109] N=634	-0.157 [0.143] N=499
Underweight: WAZ below -2	**0.112*** [0.039] N=1,118**	**0.093** [0.043] N=1,023**	**0.095** [0.045] N=890**	**0.105* [0.053] N=733**	0.106 [0.077] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.099* [0.053] N=957**	-0.067 [0.056] N=877	-0.099 [0.062] N=764	-0.099 [0.074] N=635	**-0.164** [0.081] N=497**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.009 [0.061] N=957	-0.007 [0.066] N=877	-0.016 [0.068] N=764	-0.030 [0.086] N=635	-0.121 [0.104] N=497
Obese: WLZ/WHZ above +3, 0-60 months	-0.075 [0.048] N=957	-0.067 [0.056] N=877	-0.036 [0.057] N=764	-0.035 [0.069] N=635	-0.031 [0.086] N=497
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.098 [0.079] N=914	-0.070 [0.086] N=837	-0.089 [0.093] N=727	-0.171 [0.112] N=606	-0.189 [0.147] N=475
BMI-for-age z score below -2	-0.073 [0.053] N=942	-0.060 [0.055] N=864	-0.075 [0.065] N=751	-0.050 [0.074] N=622	-0.066 [0.082] N=487

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.100 [0.240] N=1,144	0.057 [0.253] N=1,063	0.064 [0.283] N=946	0.149 [0.290] N=820	-0.098 [0.313] N=685
WHO weight-for-age z score	-0.016	-0.086 [0.136] N=1,327	-0.080 [0.150] N=1,232	-0.031 [0.154] N=1,099	-0.008 [0.155] N=942	-0.060 [0.176] N=783
WHO weight-for-length/height z score, 0-60 months	0.317	**0.403* [0.221] N=1,143**	**0.405* [0.226] N=1,063**	**0.472** [0.231] N=950**	**0.399* [0.235] N=821**	**0.486* [0.252] N=683**
WHO BMI-for-age z score	0.303	0.349 [0.243] N=1,126	0.352 [0.248] N=1,048	0.423 [0.261] N=935	0.299 [0.265] N=806	0.353 [0.285] N=671
Stunted: HAZ below -2	0.247 P=271	-0.036 [0.050] N=1,144	-0.032 [0.053] N=1,063	-0.025 [0.055] N=946	-0.053 [0.059] N=820	-0.020 [0.062] N=685
Underweight: WAZ below -2	0.089 P=119	**0.067** [0.028] N=1,327**	**0.055* [0.029] N=1,232**	**0.052* [0.031] N=1,099**	0.049 [0.034] N=942	0.042 [0.039] N=783
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=135	**-0.065* [0.036] N=1,143**	-0.047 [0.037] N=1,063	-0.061 [0.040] N=950	-0.045 [0.042] N=821	-0.054 [0.044] N=683
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=223	0.054 [0.047] N=1,143	0.062 [0.049] N=1,063	0.067 [0.051] N=950	0.056 [0.057] N=821	0.053 [0.064] N=683
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=106	-0.010 [0.037] N=1,143	0.000 [0.040] N=1,063	0.023 [0.041] N=950	0.023 [0.045] N=821	0.041 [0.049] N=683
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=361	-0.072 [0.052] N=1,094	-0.051 [0.055] N=1,017	-0.054 [0.058] N=907	-0.079 [0.065] N=786	-0.056 [0.070] N=655
BMI-for-age z score below -2	0.129 P=142	-0.041 [0.040] N=1,126	-0.034 [0.042] N=1,048	-0.040 [0.047] N=935	-0.014 [0.049] N=806	-0.015 [0.052] N=671

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.209 [0.378] N=958	0.164 [0.424] N=877	0.306 [0.502] N=760	0.609 [0.547] N=634	0.345 [0.699] N=499
WHO weight-for-age z score	**-0.338* [0.189] N=1,118**	-0.354 [0.225] N=1,023	-0.312 [0.231] N=890	-0.350 [0.260] N=733	-0.600 [0.401] N=574
WHO weight-for-length/height z score, 0-60 months	0.262 [0.348] N=957	0.219 [0.371] N=877	0.256 [0.403] N=764	0.165 [0.459] N=635	0.180 [0.537] N=497
WHO BMI-for-age z score	0.264 [0.366] N=942	0.214 [0.384] N=864	0.252 [0.423] N=751	0.020 [0.457] N=622	-0.125 [0.533] N=487
Stunted: HAZ below -2	-0.060 [0.075] N=958	-0.066 [0.082] N=877	-0.075 [0.091] N=760	-0.155 [0.109] N=634	-0.162 [0.141] N=499
Underweight: WAZ below -2	**0.106*** [0.038] N=1,118**	**0.087** [0.042] N=1,023**	**0.090** [0.044] N=890**	**0.106** [0.053] N=733**	0.098 [0.076] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.097* [0.052] N=957**	-0.066 [0.056] N=877	-0.099 [0.062] N=764	-0.098 [0.075] N=635	**-0.163** [0.081] N=497**
Overweight: WLZ/WHZ above +2, 0-60 months	0.003 [0.061] N=957	0.006 [0.066] N=877	-0.007 [0.068] N=764	-0.033 [0.085] N=635	-0.132 [0.101] N=497
Obese: WLZ/WHZ above +3, 0-60 months	-0.070 [0.048] N=957	-0.061 [0.055] N=877	-0.033 [0.057] N=764	-0.038 [0.069] N=635	-0.044 [0.086] N=497
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.087 [0.078] N=914	-0.061 [0.085] N=837	-0.085 [0.093] N=727	-0.168 [0.112] N=606	-0.194 [0.144] N=475
BMI-for-age z score below -2	-0.068 [0.053] N=942	-0.055 [0.055] N=864	-0.073 [0.065] N=751	-0.049 [0.075] N=622	-0.066 [0.082] N=487

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.303	-0.106 [0.318] N=1,084	-0.183 [0.325] N=976	0.011 [0.347] N=848	0.105 [0.382] N=715	0.084 [0.428] N=583
WHO weight-for-age z score	-0.006	-0.010 [0.166] N=1,250	0.021 [0.176] N=1,131	-0.059 [0.182] N=981	0.017 [0.200] N=826	0.086 [0.226] N=669
WHO weight-for-length/height z score, 0-60 months	0.356	0.056 [0.269] N=1,084	0.060 [0.280] N=979	-0.017 [0.302] N=851	0.119 [0.333] N=716	0.048 [0.383] N=581
WHO BMI-for-age z score	0.319	0.119 [0.264] N=1,065	0.178 [0.271] N=961	0.117 [0.304] N=835	0.307 [0.333] N=703	0.256 [0.375] N=572
Stunted: HAZ below -2	0.232 P=248	0.030 [0.059] N=1,084	0.031 [0.061] N=976	-0.004 [0.063] N=848	-0.017 [0.072] N=715	-0.061 [0.078] N=583
Underweight: WAZ below -2	0.086 P=107	-0.006 [0.036] N=1,250	-0.005 [0.038] N=1,131	0.007 [0.039] N=981	0.018 [0.042] N=826	0.013 [0.046] N=669
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=127	-0.001 [0.044] N=1,084	0.008 [0.046] N=979	0.021 [0.051] N=851	-0.022 [0.050] N=716	-0.011 [0.058] N=581
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=219	0.033 [0.050] N=1,084	0.043 [0.052] N=979	0.012 [0.052] N=851	0.015 [0.057] N=716	0.018 [0.065] N=581
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=105	0.027 [0.042] N=1,084	0.025 [0.042] N=979	0.015 [0.043] N=851	0.011 [0.048] N=716	-0.018 [0.048] N=581
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=334	0.036 [0.055] N=1,037	0.037 [0.056] N=935	0.023 [0.059] N=811	-0.018 [0.062] N=683	-0.037 [0.067] N=555
BMI-for-age z score below -2	0.130 P=138	0.024 [0.047] N=1,065	0.033 [0.050] N=961	0.018 [0.057] N=835	-0.019 [0.057] N=703	0.002 [0.065] N=572

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.304 [0.476] N=942	-0.431 [0.501] N=834	-0.141 [0.572] N=706	-0.043 [0.659] N=573	-0.092 [0.826] N=441
WHO weight-for-age z score	-0.080 [0.239] N=1,088	-0.038 [0.260] N=969	-0.218 [0.281] N=819	-0.134 [0.315] N=664	-0.093 [0.366] N=507
WHO weight-for-length/height z score, 0-60 months	-0.160 [0.392] N=942	-0.181 [0.418] N=837	-0.379 [0.483] N=709	-0.197 [0.546] N=574	-0.648 [0.647] N=439
WHO BMI-for-age z score	-0.203 [0.378] N=928	-0.127 [0.396] N=824	-0.271 [0.480] N=698	0.040 [0.546] N=566	-0.300 [0.577] N=435
Stunted: HAZ below -2	0.088 [0.084] N=942	0.089 [0.087] N=834	0.041 [0.093] N=706	0.039 [0.113] N=573	-0.080 [0.129] N=441
Underweight: WAZ below -2	0.001 [0.060] N=1,088	0.005 [0.066] N=969	0.034 [0.073] N=819	0.062 [0.084] N=664	0.060 [0.101] N=507
Wasted: WLZ/WHZ below -2, 0-60 months	0.043 [0.061] N=942	0.064 [0.065] N=837	0.097 [0.075] N=709	0.029 [0.076] N=574	0.082 [0.092] N=439
Overweight: WLZ/WHZ above +2, 0-60 months	0.021 [0.077] N=942	0.029 [0.082] N=837	-0.029 [0.088] N=709	-0.027 [0.102] N=574	-0.085 [0.125] N=439
Obese: WLZ/WHZ above +3, 0-60 months	0.045 [0.058] N=942	0.035 [0.062] N=837	0.019 [0.068] N=709	0.004 [0.086] N=574	-0.127 [0.102] N=439
CIAF: any stunting, underweight, or wasting, 0-60 months	0.095 [0.086] N=900	0.100 [0.092] N=798	0.094 [0.099] N=674	0.038 [0.112] N=546	-0.018 [0.134] N=418
BMI-for-age z score below -2	0.062 [0.056] N=928	0.075 [0.060] N=824	0.056 [0.076] N=698	-0.030 [0.070] N=566	0.003 [0.086] N=435

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.303	-0.110 [0.318] N=1,084	-0.193 [0.325] N=976	-0.011 [0.345] N=848	0.085 [0.380] N=715	0.060 [0.425] N=583
WHO weight-for-age z score	-0.006	-0.003 [0.166] N=1,250	0.027 [0.176] N=1,131	-0.057 [0.183] N=981	0.023 [0.200] N=826	0.090 [0.227] N=669
WHO weight-for-length/height z score, 0-60 months	0.356	0.059 [0.269] N=1,084	0.058 [0.280] N=979	-0.019 [0.303] N=851	0.118 [0.333] N=716	0.055 [0.380] N=581
WHO BMI-for-age z score	0.319	0.114 [0.264] N=1,065	0.172 [0.272] N=961	0.109 [0.304] N=835	0.300 [0.333] N=703	0.255 [0.376] N=572
Stunted: HAZ below -2	0.232 P=248	0.026 [0.060] N=1,084	0.027 [0.061] N=976	-0.007 [0.064] N=848	-0.019 [0.072] N=715	-0.061 [0.079] N=583
Underweight: WAZ below -2	0.086 P=107	-0.008 [0.037] N=1,250	-0.007 [0.039] N=1,131	0.006 [0.039] N=981	0.016 [0.042] N=826	0.011 [0.046] N=669
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=127	-0.005 [0.044] N=1,084	0.004 [0.046] N=979	0.017 [0.051] N=851	-0.025 [0.049] N=716	-0.016 [0.056] N=581
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=219	0.030 [0.050] N=1,084	0.038 [0.052] N=979	0.007 [0.053] N=851	0.010 [0.058] N=716	0.015 [0.066] N=581
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=105	0.025 [0.041] N=1,084	0.021 [0.042] N=979	0.011 [0.043] N=851	0.007 [0.048] N=716	-0.022 [0.048] N=581
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=334	0.026 [0.056] N=1,037	0.026 [0.057] N=935	0.014 [0.060] N=811	-0.024 [0.063] N=683	-0.044 [0.069] N=555
BMI-for-age z score below -2	0.130 P=138	0.022 [0.047] N=1,065	0.030 [0.050] N=961	0.016 [0.057] N=835	-0.020 [0.058] N=703	0.001 [0.066] N=572

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.300 [0.477] N=942	-0.442 [0.504] N=834	-0.188 [0.572] N=706	-0.096 [0.664] N=573	-0.164 [0.842] N=441
WHO weight-for-age z score	-0.068 [0.238] N=1,088	-0.027 [0.260] N=969	-0.210 [0.281] N=819	-0.112 [0.316] N=664	-0.068 [0.370] N=507
WHO weight-for-length/height z score, 0-60 months	-0.154 [0.396] N=942	-0.184 [0.423] N=837	-0.383 [0.490] N=709	-0.190 [0.556] N=574	-0.606 [0.672] N=439
WHO BMI-for-age z score	-0.203 [0.381] N=928	-0.125 [0.401] N=824	-0.270 [0.485] N=698	0.054 [0.553] N=566	-0.246 [0.605] N=435
Stunted: HAZ below -2	0.078 [0.084] N=942	0.079 [0.088] N=834	0.035 [0.093] N=706	0.036 [0.115] N=573	-0.078 [0.131] N=441
Underweight: WAZ below -2	-0.001 [0.061] N=1,088	0.002 [0.067] N=969	0.030 [0.073] N=819	0.056 [0.085] N=664	0.056 [0.101] N=507
Wasted: WLZ/WHZ below -2, 0-60 months	0.035 [0.063] N=942	0.054 [0.067] N=837	0.086 [0.078] N=709	0.013 [0.077] N=574	0.058 [0.096] N=439
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.077] N=942	0.019 [0.082] N=837	-0.042 [0.088] N=709	-0.046 [0.102] N=574	-0.103 [0.125] N=439
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.058] N=942	0.028 [0.062] N=837	0.010 [0.068] N=709	-0.007 [0.087] N=574	-0.144 [0.102] N=439
CIAF: any stunting, underweight, or wasting, 0-60 months	0.074 [0.086] N=900	0.075 [0.091] N=798	0.069 [0.099] N=674	0.010 [0.112] N=546	-0.055 [0.132] N=418
BMI-for-age z score below -2	0.055 [0.057] N=928	0.066 [0.062] N=824	0.046 [0.077] N=698	-0.042 [0.073] N=566	-0.013 [0.093] N=435

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.303	-0.125 [0.311] N=1,084	-0.191 [0.318] N=976	-0.004 [0.340] N=848	0.081 [0.376] N=715	0.049 [0.422] N=583
WHO weight-for-age z score	-0.006	-0.010 [0.163] N=1,250	0.020 [0.172] N=1,131	-0.049 [0.181] N=981	0.026 [0.199] N=826	0.089 [0.224] N=669
WHO weight-for-length/height z score, 0-60 months	0.356	0.065 [0.269] N=1,084	0.058 [0.281] N=979	-0.006 [0.302] N=851	0.125 [0.331] N=716	0.048 [0.379] N=581
WHO BMI-for-age z score	0.319	0.115 [0.263] N=1,065	0.162 [0.271] N=961	0.116 [0.299] N=835	0.307 [0.326] N=703	0.243 [0.371] N=572
Stunted: HAZ below -2	0.232 P=248	0.030 [0.058] N=1,084	0.028 [0.060] N=976	-0.009 [0.062] N=848	-0.020 [0.070] N=715	-0.063 [0.076] N=583
Underweight: WAZ below -2	0.086 P=107	-0.004 [0.036] N=1,250	-0.002 [0.038] N=1,131	0.007 [0.038] N=981	0.017 [0.041] N=826	0.013 [0.044] N=669
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=127	-0.005 [0.044] N=1,084	0.004 [0.046] N=979	0.014 [0.052] N=851	-0.026 [0.050] N=716	-0.016 [0.058] N=581
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=219	0.032 [0.050] N=1,084	0.040 [0.052] N=979	0.006 [0.053] N=851	0.006 [0.058] N=716	0.015 [0.065] N=581
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=105	0.026 [0.041] N=1,084	0.022 [0.042] N=979	0.010 [0.044] N=851	0.006 [0.049] N=716	-0.025 [0.049] N=581
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=334	0.033 [0.056] N=1,037	0.031 [0.058] N=935	0.009 [0.061] N=811	-0.029 [0.064] N=683	-0.051 [0.067] N=555
BMI-for-age z score below -2	0.130 P=138	0.024 [0.047] N=1,065	0.033 [0.050] N=961	0.015 [0.058] N=835	-0.021 [0.058] N=703	0.003 [0.066] N=572

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366 [0.477] N=942	-0.481 [0.501] N=834	-0.214 [0.565] N=706	-0.156 [0.656] N=573	-0.263 [0.855] N=441
WHO weight-for-age z score	-0.120 [0.237] N=1,088	-0.079 [0.257] N=969	-0.218 [0.279] N=819	-0.142 [0.317] N=664	-0.138 [0.374] N=507
WHO weight-for-length/height z score, 0-60 months	-0.165 [0.396] N=942	-0.197 [0.424] N=837	-0.368 [0.486] N=709	-0.179 [0.553] N=574	-0.603 [0.683] N=439
WHO BMI-for-age z score	-0.202 [0.383] N=928	-0.140 [0.404] N=824	-0.251 [0.479] N=698	0.073 [0.545] N=566	-0.225 [0.615] N=435
Stunted: HAZ below -2	0.092 [0.084] N=942	0.086 [0.087] N=834	0.041 [0.091] N=706	0.050 [0.113] N=573	-0.054 [0.132] N=441
Underweight: WAZ below -2	0.006 [0.059] N=1,088	0.011 [0.065] N=969	0.032 [0.071] N=819	0.060 [0.083] N=664	0.066 [0.099] N=507
Wasted: WLZ/WHZ below -2, 0-60 months	0.041 [0.064] N=942	0.062 [0.068] N=837	0.087 [0.079] N=709	0.016 [0.078] N=574	0.065 [0.098] N=439
Overweight: WLZ/WHZ above +2, 0-60 months	0.020 [0.078] N=942	0.024 [0.083] N=837	-0.041 [0.088] N=709	-0.043 [0.103] N=574	-0.096 [0.127] N=439
Obese: WLZ/WHZ above +3, 0-60 months	0.043 [0.058] N=942	0.032 [0.062] N=837	0.012 [0.068] N=709	-0.004 [0.086] N=574	-0.139 [0.103] N=439
CIAF: any stunting, underweight, or wasting, 0-60 months	0.098 [0.086] N=900	0.097 [0.092] N=798	0.074 [0.100] N=674	0.023 [0.114] N=546	-0.029 [0.131] N=418
BMI-for-age z score below -2	0.057 [0.059] N=928	0.073 [0.063] N=824	0.046 [0.077] N=698	-0.042 [0.073] N=566	-0.011 [0.094] N=435

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.303	-0.131 [0.310] N=1,084	-0.192 [0.318] N=976	0.020 [0.338] N=848	0.092 [0.370] N=715	0.037 [0.415] N=583
WHO weight-for-age z score	-0.006	-0.016 [0.163] N=1,250	0.019 [0.173] N=1,131	-0.038 [0.183] N=981	0.031 [0.202] N=826	0.090 [0.231] N=669
WHO weight-for-length/height z score, 0-60 months	0.356	0.075 [0.266] N=1,084	0.072 [0.278] N=979	0.005 [0.296] N=851	0.157 [0.326] N=716	0.115 [0.378] N=581
WHO BMI-for-age z score	0.319	0.128 [0.264] N=1,065	0.182 [0.272] N=961	0.127 [0.299] N=835	0.329 [0.328] N=703	0.306 [0.376] N=572
Stunted: HAZ below -2	0.232 P=248	0.030 [0.058] N=1,084	0.027 [0.060] N=976	-0.012 [0.063] N=848	-0.020 [0.070] N=715	-0.058 [0.077] N=583
Underweight: WAZ below -2	0.086 P=107	-0.008 [0.035] N=1,250	-0.007 [0.038] N=1,131	0.002 [0.039] N=981	0.016 [0.042] N=826	0.012 [0.047] N=669
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=127	-0.005 [0.044] N=1,084	0.004 [0.046] N=979	0.017 [0.050] N=851	-0.027 [0.048] N=716	-0.021 [0.055] N=581
Overweight: WLZ/WHZ above +2, 0-60 months	0.204 P=219	0.031 [0.049] N=1,084	0.039 [0.051] N=979	0.009 [0.052] N=851	0.014 [0.057] N=716	0.021 [0.065] N=581
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=105	0.031 [0.040] N=1,084	0.027 [0.041] N=979	0.017 [0.042] N=851	0.015 [0.047] N=716	-0.013 [0.047] N=581
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=334	0.032 [0.055] N=1,037	0.031 [0.057] N=935	0.010 [0.059] N=811	-0.025 [0.062] N=683	-0.039 [0.068] N=555
BMI-for-age z score below -2	0.130 P=138	0.020 [0.047] N=1,065	0.028 [0.050] N=961	0.013 [0.057] N=835	-0.024 [0.057] N=703	-0.008 [0.064] N=572

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.357 [0.459] N=942	-0.471 [0.487] N=834	-0.115 [0.553] N=706	-0.062 [0.646] N=573	-0.188 [0.833] N=441
WHO weight-for-age z score	-0.088 [0.230] N=1,088	-0.040 [0.251] N=969	-0.148 [0.281] N=819	-0.068 [0.313] N=664	-0.042 [0.380] N=507
WHO weight-for-length/height z score, 0-60 months	-0.155 [0.395] N=942	-0.187 [0.423] N=837	-0.378 [0.483] N=709	-0.171 [0.548] N=574	-0.565 [0.677] N=439
WHO BMI-for-age z score	-0.191 [0.380] N=928	-0.116 [0.401] N=824	-0.253 [0.482] N=698	0.089 [0.547] N=566	-0.200 [0.613] N=435
Stunted: HAZ below -2	0.089 [0.082] N=942	0.085 [0.086] N=834	0.026 [0.091] N=706	0.033 [0.114] N=573	-0.060 [0.132] N=441
Underweight: WAZ below -2	-0.003 [0.058] N=1,088	0.001 [0.064] N=969	0.020 [0.072] N=819	0.052 [0.084] N=664	0.052 [0.102] N=507
Wasted: WLZ/WHZ below -2, 0-60 months	0.039 [0.063] N=942	0.060 [0.067] N=837	0.092 [0.077] N=709	0.020 [0.076] N=574	0.069 [0.093] N=439
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.077] N=942	0.018 [0.082] N=837	-0.046 [0.087] N=709	-0.049 [0.104] N=574	-0.096 [0.129] N=439
Obese: WLZ/WHZ above +3, 0-60 months	0.045 [0.057] N=942	0.033 [0.060] N=837	0.014 [0.066] N=709	0.003 [0.085] N=574	-0.129 [0.102] N=439
CIAF: any stunting, underweight, or wasting, 0-60 months	0.086 [0.084] N=900	0.085 [0.090] N=798	0.054 [0.098] N=674	0.003 [0.111] N=546	-0.031 [0.130] N=418
BMI-for-age z score below -2	0.058 [0.058] N=928	0.069 [0.063] N=824	0.047 [0.077] N=698	-0.041 [0.073] N=566	-0.014 [0.093] N=435

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.292	-0.175 [0.327] N=993	-0.269 [0.338] N=894	-0.050 [0.351] N=782	0.037 [0.375] N=657	-0.016 [0.432] N=538
WHO weight-for-age z score	-0.005	0.008 [0.171] N=1,142	0.054 [0.180] N=1,032	-0.041 [0.185] N=902	0.008 [0.200] N=760	0.033 [0.222] N=620
WHO weight-for-length/height z score, 0-60 months	0.361	0.154 [0.284] N=996	0.174 [0.299] N=900	0.053 [0.320] N=786	0.141 [0.347] N=660	0.036 [0.385] N=539
WHO BMI-for-age z score	0.329	0.167 [0.280] N=979	0.252 [0.293] N=884	0.128 [0.325] N=773	0.293 [0.357] N=650	0.191 [0.399] N=532
Stunted: HAZ below -2	0.232 P=227	0.040 [0.064] N=993	0.044 [0.066] N=894	0.006 [0.067] N=782	-0.014 [0.073] N=657	-0.051 [0.082] N=538
Underweight: WAZ below -2	0.086 P=98	-0.001 [0.037] N=1,142	-0.000 [0.040] N=1,032	0.005 [0.040] N=902	0.023 [0.043] N=760	0.025 [0.047] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=116	-0.007 [0.045] N=996	-0.001 [0.047] N=900	0.013 [0.052] N=786	-0.026 [0.050] N=660	-0.008 [0.056] N=539
Overweight: WLZ/WHZ above +2, 0-60 months	0.205 P=203	0.039 [0.052] N=996	0.049 [0.054] N=900	0.006 [0.054] N=786	-0.007 [0.059] N=660	-0.014 [0.065] N=539
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=99	0.028 [0.042] N=996	0.024 [0.043] N=900	0.011 [0.043] N=786	0.002 [0.047] N=660	-0.026 [0.047] N=539
CIAF: any stunting, underweight, or wasting, 0-60 months	0.328 P=307	0.043 [0.060] N=951	0.041 [0.062] N=858	0.027 [0.062] N=748	-0.022 [0.064] N=628	-0.024 [0.072] N=513
BMI-for-age z score below -2	0.128 P=123	0.022 [0.049] N=979	0.028 [0.053] N=884	0.014 [0.060] N=773	-0.012 [0.060] N=650	0.020 [0.069] N=532

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.240 [0.514] N=861	-0.397 [0.546] N=762	-0.085 [0.604] N=650	0.041 [0.687] N=525	0.112 [0.896] N=406
WHO weight-for-age z score	-0.001 [0.244] N=992	0.064 [0.265] N=882	-0.160 [0.293] N=752	-0.060 [0.330] N=610	-0.022 [0.399] N=470
WHO weight-for-length/height z score, 0-60 months	-0.053 [0.424] N=864	-0.043 [0.458] N=768	-0.320 [0.520] N=654	-0.175 [0.601] N=528	-0.768 [0.715] N=407
WHO BMI-for-age z score	-0.152 [0.393] N=850	-0.028 [0.420] N=755	-0.277 [0.499] N=644	0.036 [0.575] N=521	-0.432 [0.626] N=403
Stunted: HAZ below -2	0.077 [0.095] N=861	0.082 [0.100] N=762	0.032 [0.103] N=650	0.005 [0.125] N=525	-0.136 [0.157] N=406
Underweight: WAZ below -2	-0.003 [0.062] N=992	0.001 [0.068] N=882	0.018 [0.075] N=752	0.056 [0.091] N=610	0.048 [0.111] N=470
Wasted: WLZ/WHZ below -2, 0-60 months	0.030 [0.065] N=864	0.044 [0.071] N=768	0.071 [0.082] N=654	-0.012 [0.081] N=528	0.046 [0.096] N=407
Overweight: WLZ/WHZ above +2, 0-60 months	0.029 [0.084] N=864	0.036 [0.090] N=768	-0.041 [0.096] N=654	-0.079 [0.114] N=528	-0.173 [0.136] N=407
Obese: WLZ/WHZ above +3, 0-60 months	0.037 [0.064] N=864	0.025 [0.067] N=768	0.004 [0.073] N=654	-0.024 [0.094] N=528	-0.166 [0.113] N=407
CIAF: any stunting, underweight, or wasting, 0-60 months	0.077 [0.097] N=824	0.073 [0.104] N=731	0.059 [0.109] N=621	-0.050 [0.126] N=501	-0.131 [0.163] N=386
BMI-for-age z score below -2	0.057 [0.061] N=850	0.065 [0.067] N=755	0.042 [0.083] N=644	-0.041 [0.080] N=521	-0.002 [0.098] N=403

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.292	-0.188 [0.328] N=993	-0.281 [0.339] N=894	-0.061 [0.351] N=782	0.021 [0.373] N=657	-0.026 [0.430] N=538
WHO weight-for-age z score	-0.005	-0.001 [0.169] N=1,142	0.037 [0.179] N=1,032	-0.051 [0.183] N=902	0.005 [0.197] N=760	0.037 [0.218] N=620
WHO weight-for-length/height z score, 0-60 months	0.361	0.141 [0.286] N=996	0.159 [0.301] N=900	0.039 [0.321] N=786	0.132 [0.348] N=660	0.031 [0.391] N=539
WHO BMI-for-age z score	0.329	0.170 [0.283] N=979	0.256 [0.294] N=884	0.131 [0.327] N=773	0.297 [0.359] N=650	0.192 [0.404] N=532
Stunted: HAZ below -2	0.232 P=227	0.039 [0.063] N=993	0.042 [0.065] N=894	0.005 [0.067] N=782	-0.017 [0.072] N=657	-0.053 [0.080] N=538
Underweight: WAZ below -2	0.086 P=98	-0.005 [0.037] N=1,142	-0.002 [0.039] N=1,032	0.002 [0.039] N=902	0.020 [0.042] N=760	0.022 [0.045] N=620
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=116	-0.011 [0.045] N=996	-0.005 [0.048] N=900	0.009 [0.052] N=786	-0.032 [0.050] N=660	-0.015 [0.057] N=539
Overweight: WLZ/WHZ above +2, 0-60 months	0.205 P=203	0.034 [0.052] N=996	0.044 [0.054] N=900	0.002 [0.054] N=786	-0.012 [0.058] N=660	-0.017 [0.065] N=539
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=99	0.023 [0.042] N=996	0.019 [0.042] N=900	0.006 [0.042] N=786	-0.002 [0.046] N=660	-0.029 [0.045] N=539
CIAF: any stunting, underweight, or wasting, 0-60 months	0.328 P=307	0.038 [0.059] N=951	0.035 [0.062] N=858	0.021 [0.062] N=748	-0.033 [0.064] N=628	-0.035 [0.071] N=513
BMI-for-age z score below -2	0.128 P=123	0.020 [0.049] N=979	0.025 [0.053] N=884	0.012 [0.060] N=773	-0.017 [0.059] N=650	0.015 [0.067] N=532

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.242 [0.509] N=861	-0.394 [0.542] N=762	-0.078 [0.596] N=650	0.034 [0.670] N=525	0.045 [0.866] N=406
WHO weight-for-age z score	-0.030 [0.245] N=992	0.031 [0.269] N=882	-0.183 [0.293] N=752	-0.087 [0.327] N=610	-0.047 [0.385] N=470
WHO weight-for-length/height z score, 0-60 months	-0.106 [0.421] N=864	-0.098 [0.455] N=768	-0.379 [0.516] N=654	-0.241 [0.586] N=528	-0.766 [0.692] N=407
WHO BMI-for-age z score	-0.156 [0.392] N=850	-0.031 [0.419] N=755	-0.281 [0.498] N=644	0.048 [0.570] N=521	-0.394 [0.605] N=403
Stunted: HAZ below -2	0.077 [0.094] N=861	0.081 [0.099] N=762	0.032 [0.102] N=650	0.007 [0.123] N=525	-0.117 [0.149] N=406
Underweight: WAZ below -2	-0.005 [0.062] N=992	-0.001 [0.069] N=882	0.016 [0.075] N=752	0.055 [0.090] N=610	0.062 [0.107] N=470
Wasted: WLZ/WHZ below -2, 0-60 months	0.033 [0.065] N=864	0.047 [0.071] N=768	0.074 [0.083] N=654	-0.008 [0.080] N=528	0.054 [0.096] N=407
Overweight: WLZ/WHZ above +2, 0-60 months	0.020 [0.083] N=864	0.026 [0.089] N=768	-0.053 [0.094] N=654	-0.091 [0.110] N=528	-0.168 [0.131] N=407
Obese: WLZ/WHZ above +3, 0-60 months	0.027 [0.063] N=864	0.017 [0.067] N=768	-0.005 [0.072] N=654	-0.032 [0.091] N=528	-0.163 [0.107] N=407
CIAF: any stunting, underweight, or wasting, 0-60 months	0.080 [0.096] N=824	0.076 [0.103] N=731	0.064 [0.109] N=621	-0.043 [0.127] N=501	-0.090 [0.157] N=386
BMI-for-age z score below -2	0.059 [0.061] N=850	0.068 [0.067] N=755	0.044 [0.083] N=644	-0.039 [0.079] N=521	0.009 [0.097] N=403

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.