Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-2: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.924 P=1,068	0.003 [0.030] N=1,159	0.018 [0.029] N=1,069	0.010 [0.032] N=956	0.004 [0.034] N=829	-0.005 [0.039] N=682
Vegetables consumed at least weekly	0.894 P=1,034	0.003 [0.033] N=1,159	-0.003 [0.032] N=1,069	-0.015 [0.033] N=956	-0.017 [0.036] N=829	-0.046 [0.044] N=682
Meat fish or poultry consumed at least weekly	0.755 P=881	0.051 [0.061] N=1,159	0.070 [0.064] N=1,069	0.079 [0.070] N=956	0.058 [0.078] N=829	0.053 [0.089] N=682
Legumes consumed at least weekly	0.753 P=877	-0.046 [0.049] N=1,159	-0.072 [0.050] N=1,069	-0.050 [0.051] N=956	-0.062 [0.056] N=829	-0.058 [0.066] N=682
Animal products consumed at least weekly	0.900 P=1,047	0.010 [0.039] N=1,159	0.007 [0.042] N=1,069	-0.016 [0.045] N=956	-0.006 [0.049] N=829	-0.013 [0.056] N=682
Sweets consumed at least weekly	0.508 P=589	-0.025 [0.049] N=1,159	-0.028 [0.050] N=1,069	-0.021 [0.055] N=956	-0.039 [0.060] N=829	0.013 [0.069] N=682
Snack foods consumed at least weekly	0.419 P=482	-0.005 [0.047] N=1,159	-0.016 [0.049] N=1,069	0.001 [0.052] N=956	0.008 [0.056] N=829	0.028 [0.059] N=682
Packaged soft drinks consumed at least weekly	0.235 P=268	-0.016 [0.045] N=1,159	-0.038 [0.047] N=1,069	-0.044 [0.049] N=956	-0.044 [0.052] N=829	-0.064 [0.059] N=682
Grains consumed at least weekly	0.852 P=990	0.028 [0.039] N=1,159	0.034 [0.040] N=1,069	0.018 [0.042] N=956	0.027 [0.045] N=829	0.007 [0.053] N=682
Child was ever breastfed	0.969 P=1,373	-0.012 [0.013] N=1,421	-0.010 [0.014] N=1,316	-0.012 [0.015] N=1,175	-0.017 [0.017] N=1,005	-0.017 [0.017] N=829

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.049] N=979	0.024 [0.049] N=889	0.007 [0.056] N=776	-0.011 [0.066] N=649	-0.043 [0.086] N=502
Vegetables consumed at least weekly	0.012 [0.052] N=979	0.002 [0.052] N=889	-0.019 [0.056] N=776	-0.032 [0.065] N=649	-0.126 [0.093] N=502
Meat fish or poultry consumed at least weekly	0.000 [0.067] N=979	0.020 [0.068] N=889	0.028 [0.079] N=776	-0.073 [0.083] N=649	**-0.166* [0.093] N=502**
Legumes consumed at least weekly	-0.043 [0.074] N=979	-0.093 [0.076] N=889	-0.045 [0.080] N=776	-0.089 [0.093] N=649	-0.090 [0.137] N=502
Animal products consumed at least weekly	0.006 [0.052] N=979	-0.001 [0.057] N=889	-0.053 [0.064] N=776	-0.036 [0.074] N=649	-0.080 [0.092] N=502
Sweets consumed at least weekly	-0.039 [0.068] N=979	-0.052 [0.072] N=889	-0.045 [0.082] N=776	-0.121 [0.088] N=649	-0.001 [0.122] N=502
Snack foods consumed at least weekly	-0.026 [0.066] N=979	-0.056 [0.071] N=889	-0.032 [0.082] N=776	-0.042 [0.100] N=649	0.001 [0.126] N=502
Packaged soft drinks consumed at least weekly	0.014 [0.065] N=979	-0.025 [0.071] N=889	-0.029 [0.076] N=776	-0.038 [0.089] N=649	-0.114 [0.120] N=502
Grains consumed at least weekly	0.064 [0.060] N=979	0.080 [0.064] N=889	0.059 [0.070] N=776	0.085 [0.083] N=649	0.050 [0.118] N=502
Child was ever breastfed	-0.004 [0.020] N=1,203	0.002 [0.023] N=1,098	-0.000 [0.028] N=957	-0.017 [0.034] N=787	-0.003 [0.042] N=611

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.924 P=1,068	0.002 [0.030] N=1,159	0.017 [0.029] N=1,069	0.010 [0.031] N=956	0.004 [0.033] N=829	-0.006 [0.038] N=682
Vegetables consumed at least weekly	0.894 P=1,034	0.003 [0.033] N=1,159	-0.004 [0.032] N=1,069	-0.015 [0.033] N=956	-0.017 [0.036] N=829	-0.047 [0.043] N=682
Meat fish or poultry consumed at least weekly	0.755 P=881	0.051 [0.061] N=1,159	0.070 [0.064] N=1,069	0.079 [0.070] N=956	0.057 [0.078] N=829	0.051 [0.090] N=682
Legumes consumed at least weekly	0.753 P=877	-0.047 [0.049] N=1,159	-0.073 [0.050] N=1,069	-0.051 [0.051] N=956	-0.064 [0.056] N=829	-0.060 [0.066] N=682
Animal products consumed at least weekly	0.900 P=1,047	0.010 [0.039] N=1,159	0.005 [0.041] N=1,069	-0.017 [0.045] N=956	-0.007 [0.048] N=829	-0.015 [0.056] N=682
Sweets consumed at least weekly	0.508 P=589	-0.025 [0.049] N=1,159	-0.029 [0.050] N=1,069	-0.021 [0.055] N=956	-0.040 [0.060] N=829	0.012 [0.069] N=682
Snack foods consumed at least weekly	0.419 P=482	-0.006 [0.047] N=1,159	-0.018 [0.049] N=1,069	0.000 [0.052] N=956	0.007 [0.057] N=829	0.026 [0.059] N=682
Packaged soft drinks consumed at least weekly	0.235 P=268	-0.016 [0.045] N=1,159	-0.039 [0.047] N=1,069	-0.044 [0.049] N=956	-0.045 [0.052] N=829	-0.065 [0.059] N=682
Grains consumed at least weekly	0.852 P=990	0.028 [0.039] N=1,159	0.033 [0.040] N=1,069	0.017 [0.042] N=956	0.026 [0.045] N=829	0.006 [0.053] N=682
Child was ever breastfed	0.969 P=1,373	-0.012 [0.013] N=1,421	-0.010 [0.014] N=1,316	-0.012 [0.015] N=1,175	-0.016 [0.016] N=1,005	-0.015 [0.016] N=829

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.049] N=979	0.024 [0.049] N=889	0.008 [0.056] N=776	-0.011 [0.066] N=649	-0.042 [0.085] N=502
Vegetables consumed at least weekly	0.012 [0.052] N=979	0.002 [0.052] N=889	-0.018 [0.056] N=776	-0.031 [0.065] N=649	-0.126 [0.093] N=502
Meat fish or poultry consumed at least weekly	-0.000 [0.067] N=979	0.020 [0.069] N=889	0.028 [0.079] N=776	-0.073 [0.083] N=649	**-0.164* [0.093] N=502**
Legumes consumed at least weekly	-0.042 [0.074] N=979	-0.093 [0.076] N=889	-0.043 [0.080] N=776	-0.088 [0.093] N=649	-0.088 [0.137] N=502
Animal products consumed at least weekly	0.008 [0.052] N=979	-0.001 [0.057] N=889	-0.050 [0.064] N=776	-0.035 [0.074] N=649	-0.078 [0.092] N=502
Sweets consumed at least weekly	-0.039 [0.068] N=979	-0.052 [0.071] N=889	-0.044 [0.082] N=776	-0.120 [0.087] N=649	0.000 [0.121] N=502
Snack foods consumed at least weekly	-0.025 [0.066] N=979	-0.056 [0.071] N=889	-0.027 [0.081] N=776	-0.040 [0.100] N=649	0.004 [0.124] N=502
Packaged soft drinks consumed at least weekly	0.013 [0.065] N=979	-0.025 [0.071] N=889	-0.029 [0.076] N=776	-0.037 [0.088] N=649	-0.114 [0.119] N=502
Grains consumed at least weekly	0.065 [0.060] N=979	0.080 [0.064] N=889	0.062 [0.069] N=776	0.086 [0.082] N=649	0.051 [0.117] N=502
Child was ever breastfed	-0.004 [0.020] N=1,203	0.002 [0.023] N=1,098	-0.001 [0.028] N=957	-0.018 [0.035] N=787	-0.003 [0.042] N=611

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.924 P=1,068	-0.007 [0.030] N=1,159	0.006 [0.030] N=1,069	-0.001 [0.032] N=956	-0.002 [0.035] N=829	-0.017 [0.039] N=682
Vegetables consumed at least weekly	0.894 P=1,034	-0.004 [0.034] N=1,159	-0.014 [0.032] N=1,069	-0.025 [0.033] N=956	-0.024 [0.036] N=829	-0.060 [0.043] N=682
Meat fish or poultry consumed at least weekly	0.755 P=881	0.028 [0.058] N=1,159	0.044 [0.062] N=1,069	0.054 [0.068] N=956	0.041 [0.075] N=829	0.025 [0.085] N=682
Legumes consumed at least weekly	0.753 P=877	-0.053 [0.049] N=1,159	-0.080 [0.050] N=1,069	-0.056 [0.051] N=956	-0.065 [0.055] N=829	-0.063 [0.065] N=682
Animal products consumed at least weekly	0.900 P=1,047	0.004 [0.039] N=1,159	-0.002 [0.042] N=1,069	-0.024 [0.045] N=956	-0.012 [0.049] N=829	-0.025 [0.056] N=682
Sweets consumed at least weekly	0.508 P=589	-0.025 [0.049] N=1,159	-0.031 [0.051] N=1,069	-0.023 [0.055] N=956	-0.038 [0.060] N=829	0.007 [0.068] N=682
Snack foods consumed at least weekly	0.419 P=482	-0.000 [0.045] N=1,159	-0.016 [0.048] N=1,069	0.001 [0.051] N=956	0.011 [0.056] N=829	0.028 [0.059] N=682
Packaged soft drinks consumed at least weekly	0.235 P=268	-0.002 [0.044] N=1,159	-0.024 [0.047] N=1,069	-0.029 [0.049] N=956	-0.033 [0.051] N=829	-0.051 [0.058] N=682
Grains consumed at least weekly	0.852 P=990	0.023 [0.039] N=1,159	0.029 [0.040] N=1,069	0.015 [0.042] N=956	0.024 [0.045] N=829	0.005 [0.053] N=682
Child was ever breastfed	0.969 P=1,373	-0.013 [0.013] N=1,421	-0.013 [0.014] N=1,316	-0.015 [0.016] N=1,175	-0.017 [0.017] N=1,005	-0.018 [0.017] N=829

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.048] N=979	0.021 [0.048] N=889	0.009 [0.055] N=776	0.012 [0.067] N=649	-0.024 [0.087] N=502
Vegetables consumed at least weekly	0.013 [0.051] N=979	0.002 [0.050] N=889	-0.015 [0.056] N=776	-0.006 [0.064] N=649	-0.101 [0.092] N=502
Meat fish or poultry consumed at least weekly	-0.008 [0.064] N=979	0.009 [0.066] N=889	0.016 [0.077] N=776	-0.058 [0.082] N=649	**-0.156* [0.092] N=502**
Legumes consumed at least weekly	-0.044 [0.075] N=979	-0.094 [0.076] N=889	-0.040 [0.079] N=776	-0.071 [0.093] N=649	-0.068 [0.138] N=502
Animal products consumed at least weekly	0.012 [0.051] N=979	0.001 [0.057] N=889	-0.047 [0.065] N=776	-0.014 [0.075] N=649	-0.045 [0.092] N=502
Sweets consumed at least weekly	-0.036 [0.068] N=979	-0.050 [0.071] N=889	-0.038 [0.081] N=776	-0.107 [0.087] N=649	0.009 [0.126] N=502
Snack foods consumed at least weekly	-0.018 [0.063] N=979	-0.053 [0.069] N=889	-0.019 [0.079] N=776	-0.028 [0.097] N=649	0.009 [0.123] N=502
Packaged soft drinks consumed at least weekly	0.030 [0.063] N=979	-0.007 [0.069] N=889	-0.006 [0.075] N=776	-0.025 [0.086] N=649	-0.104 [0.115] N=502
Grains consumed at least weekly	0.062 [0.059] N=979	0.080 [0.063] N=889	0.062 [0.069] N=776	0.098 [0.082] N=649	0.059 [0.115] N=502
Child was ever breastfed	-0.001 [0.020] N=1,203	0.003 [0.022] N=1,098	0.001 [0.027] N=957	-0.007 [0.034] N=787	-0.002 [0.041] N=611

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.924 P=1,068	-0.010 [0.031] N=1,159	0.006 [0.030] N=1,069	-0.006 [0.032] N=956	-0.010 [0.034] N=829	-0.026 [0.039] N=682
Vegetables consumed at least weekly	0.894 P=1,034	-0.007 [0.034] N=1,159	-0.012 [0.032] N=1,069	-0.025 [0.033] N=956	-0.026 [0.036] N=829	-0.058 [0.044] N=682
Meat fish or poultry consumed at least weekly	0.755 P=881	0.021 [0.058] N=1,159	0.042 [0.061] N=1,069	0.045 [0.067] N=956	0.028 [0.074] N=829	0.006 [0.085] N=682
Legumes consumed at least weekly	0.753 P=877	-0.058 [0.051] N=1,159	-0.085 [0.052] N=1,069	-0.067 [0.053] N=956	-0.076 [0.058] N=829	-0.080 [0.069] N=682
Animal products consumed at least weekly	0.900 P=1,047	0.000 [0.039] N=1,159	-0.005 [0.042] N=1,069	-0.028 [0.045] N=956	-0.018 [0.049] N=829	-0.033 [0.055] N=682
Sweets consumed at least weekly	0.508 P=589	-0.029 [0.050] N=1,159	-0.036 [0.052] N=1,069	-0.032 [0.056] N=956	-0.045 [0.061] N=829	-0.002 [0.070] N=682
Snack foods consumed at least weekly	0.419 P=482	-0.005 [0.047] N=1,159	-0.019 [0.050] N=1,069	-0.007 [0.053] N=956	0.002 [0.057] N=829	0.019 [0.059] N=682
Packaged soft drinks consumed at least weekly	0.235 P=268	-0.005 [0.044] N=1,159	-0.031 [0.047] N=1,069	-0.037 [0.049] N=956	-0.037 [0.051] N=829	-0.053 [0.058] N=682
Grains consumed at least weekly	0.852 P=990	0.025 [0.040] N=1,159	0.032 [0.041] N=1,069	0.016 [0.043] N=956	0.024 [0.046] N=829	0.008 [0.056] N=682
Child was ever breastfed	0.969 P=1,373	-0.016 [0.014] N=1,421	-0.013 [0.015] N=1,316	-0.017 [0.016] N=1,175	-0.021 [0.018] N=1,005	-0.022 [0.018] N=829

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.048] N=979	0.027 [0.048] N=889	0.007 [0.055] N=776	-0.000 [0.066] N=649	-0.052 [0.086] N=502
Vegetables consumed at least weekly	0.011 [0.052] N=979	0.004 [0.051] N=889	-0.022 [0.055] N=776	-0.024 [0.063] N=649	-0.140 [0.092] N=502
Meat fish or poultry consumed at least weekly	-0.007 [0.064] N=979	0.019 [0.066] N=889	0.021 [0.077] N=776	-0.048 [0.082] N=649	**-0.182* [0.096] N=502**
Legumes consumed at least weekly	-0.041 [0.075] N=979	-0.090 [0.077] N=889	-0.041 [0.079] N=776	-0.065 [0.093] N=649	-0.070 [0.143] N=502
Animal products consumed at least weekly	0.009 [0.051] N=979	0.001 [0.057] N=889	-0.051 [0.064] N=776	-0.029 [0.073] N=649	-0.079 [0.097] N=502
Sweets consumed at least weekly	-0.027 [0.069] N=979	-0.044 [0.071] N=889	-0.033 [0.081] N=776	-0.096 [0.089] N=649	0.011 [0.128] N=502
Snack foods consumed at least weekly	-0.013 [0.065] N=979	-0.048 [0.070] N=889	-0.015 [0.080] N=776	-0.026 [0.098] N=649	0.007 [0.123] N=502
Packaged soft drinks consumed at least weekly	0.029 [0.062] N=979	-0.016 [0.069] N=889	-0.017 [0.073] N=776	-0.032 [0.083] N=649	-0.106 [0.111] N=502
Grains consumed at least weekly	0.064 [0.060] N=979	0.080 [0.063] N=889	0.061 [0.069] N=776	0.094 [0.082] N=649	0.039 [0.115] N=502
Child was ever breastfed	-0.002 [0.020] N=1,203	0.004 [0.023] N=1,098	-0.002 [0.028] N=957	-0.011 [0.036] N=787	0.000 [0.043] N=611

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.925 P=977	-0.001 [0.030] N=1,060	0.007 [0.030] N=981	0.001 [0.032] N=877	-0.004 [0.035] N=763	-0.025 [0.039] N=632
Vegetables consumed at least weekly	0.895 P=946	-0.005 [0.035] N=1,060	-0.019 [0.034] N=981	-0.033 [0.035] N=877	-0.029 [0.038] N=763	-0.060 [0.044] N=632
Meat fish or poultry consumed at least weekly	0.758 P=806	0.046 [0.056] N=1,060	0.050 [0.059] N=981	0.051 [0.064] N=877	0.021 [0.071] N=763	-0.001 [0.083] N=632
Legumes consumed at least weekly	0.748 P=796	-0.071 [0.051] N=1,060	**-0.107** [0.052] N=981**	**-0.090* [0.052] N=877**	**-0.105* [0.058] N=763**	-0.107 [0.067] N=632
Animal products consumed at least weekly	0.903 P=960	0.015 [0.041] N=1,060	0.005 [0.043] N=981	-0.020 [0.046] N=877	-0.019 [0.051] N=763	-0.038 [0.058] N=632
Sweets consumed at least weekly	0.509 P=537	-0.009 [0.051] N=1,060	-0.023 [0.054] N=981	-0.017 [0.058] N=877	-0.032 [0.064] N=763	0.019 [0.073] N=632
Snack foods consumed at least weekly	0.421 P=440	-0.012 [0.049] N=1,060	-0.028 [0.052] N=981	-0.013 [0.055] N=877	-0.008 [0.061] N=763	0.025 [0.064] N=632
Packaged soft drinks consumed at least weekly	0.237 P=247	0.015 [0.045] N=1,060	-0.012 [0.047] N=981	-0.031 [0.048] N=877	-0.038 [0.050] N=763	-0.049 [0.057] N=632
Grains consumed at least weekly	0.854 P=906	0.018 [0.040] N=1,060	0.019 [0.041] N=981	-0.005 [0.043] N=877	0.006 [0.047] N=763	-0.020 [0.054] N=632
Child was ever breastfed	0.970 P=1,257	-0.019 [0.014] N=1,300	-0.015 [0.015] N=1,207	-0.016 [0.017] N=1,075	-0.016 [0.018] N=925	-0.017 [0.018] N=768

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.006 [0.050] N=893	0.019 [0.052] N=814	0.000 [0.058] N=710	-0.014 [0.071] N=596	-0.068 [0.090] N=465
Vegetables consumed at least weekly	0.013 [0.055] N=893	-0.010 [0.056] N=814	-0.038 [0.059] N=710	-0.035 [0.070] N=596	-0.119 [0.092] N=465
Meat fish or poultry consumed at least weekly	0.025 [0.066] N=893	0.021 [0.069] N=814	0.023 [0.078] N=710	-0.085 [0.086] N=596	**-0.192* [0.099] N=465**
Legumes consumed at least weekly	-0.047 [0.077] N=893	-0.115 [0.079] N=814	-0.072 [0.079] N=710	-0.117 [0.096] N=596	-0.131 [0.138] N=465
Animal products consumed at least weekly	0.028 [0.055] N=893	0.008 [0.061] N=814	-0.046 [0.066] N=710	-0.040 [0.078] N=596	-0.102 [0.097] N=465
Sweets consumed at least weekly	-0.017 [0.072] N=893	-0.049 [0.077] N=814	-0.052 [0.088] N=710	-0.130 [0.095] N=596	-0.024 [0.128] N=465
Snack foods consumed at least weekly	-0.030 [0.069] N=893	-0.068 [0.075] N=814	-0.045 [0.086] N=710	-0.074 [0.104] N=596	-0.014 [0.126] N=465
Packaged soft drinks consumed at least weekly	0.056 [0.067] N=893	0.007 [0.073] N=814	-0.023 [0.080] N=710	-0.044 [0.091] N=596	-0.093 [0.127] N=465
Grains consumed at least weekly	0.065 [0.061] N=893	0.072 [0.067] N=814	0.032 [0.072] N=710	0.068 [0.085] N=596	0.017 [0.120] N=465
Child was ever breastfed	-0.015 [0.022] N=1,098	-0.006 [0.025] N=1,005	-0.008 [0.030] N=873	-0.018 [0.036] N=723	-0.009 [0.045] N=566

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.925 P=977	-0.005 [0.030] N=1,060	0.003 [0.029] N=981	-0.002 [0.032] N=877	-0.007 [0.034] N=763	-0.027 [0.038] N=632
Vegetables consumed at least weekly	0.895 P=946	-0.007 [0.034] N=1,060	-0.021 [0.034] N=981	-0.034 [0.035] N=877	-0.028 [0.038] N=763	-0.061 [0.043] N=632
Meat fish or poultry consumed at least weekly	0.758 P=806	0.040 [0.057] N=1,060	0.046 [0.059] N=981	0.049 [0.064] N=877	0.020 [0.071] N=763	-0.000 [0.082] N=632
Legumes consumed at least weekly	0.748 P=796	-0.073 [0.051] N=1,060	**-0.109** [0.051] N=981**	**-0.093* [0.052] N=877**	**-0.109* [0.058] N=763**	**-0.113* [0.067] N=632**
Animal products consumed at least weekly	0.903 P=960	0.014 [0.040] N=1,060	0.003 [0.043] N=981	-0.022 [0.046] N=877	-0.017 [0.050] N=763	-0.037 [0.057] N=632
Sweets consumed at least weekly	0.509 P=537	-0.003 [0.050] N=1,060	-0.019 [0.053] N=981	-0.013 [0.057] N=877	-0.028 [0.062] N=763	0.021 [0.071] N=632
Snack foods consumed at least weekly	0.421 P=440	-0.012 [0.049] N=1,060	-0.028 [0.052] N=981	-0.011 [0.055] N=877	-0.006 [0.060] N=763	0.025 [0.063] N=632
Packaged soft drinks consumed at least weekly	0.237 P=247	0.017 [0.045] N=1,060	-0.011 [0.047] N=981	-0.028 [0.048] N=877	-0.033 [0.050] N=763	-0.046 [0.056] N=632
Grains consumed at least weekly	0.854 P=906	0.018 [0.039] N=1,060	0.019 [0.041] N=981	-0.007 [0.043] N=877	0.006 [0.046] N=763	-0.022 [0.054] N=632
Child was ever breastfed	0.970 P=1,257	-0.017 [0.014] N=1,300	-0.014 [0.015] N=1,207	-0.015 [0.016] N=1,075	-0.015 [0.018] N=925	-0.017 [0.018] N=768

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.001 [0.049] N=893	0.012 [0.051] N=814	-0.001 [0.058] N=710	-0.014 [0.070] N=596	-0.071 [0.086] N=465
Vegetables consumed at least weekly	0.007 [0.054] N=893	-0.015 [0.055] N=814	-0.040 [0.059] N=710	-0.037 [0.069] N=596	-0.130 [0.088] N=465
Meat fish or poultry consumed at least weekly	0.019 [0.067] N=893	0.019 [0.070] N=814	0.023 [0.079] N=710	-0.087 [0.087] N=596	**-0.202** [0.101] N=465**
Legumes consumed at least weekly	-0.051 [0.077] N=893	-0.119 [0.078] N=814	-0.073 [0.080] N=710	-0.117 [0.096] N=596	-0.137 [0.137] N=465
Animal products consumed at least weekly	0.023 [0.054] N=893	0.003 [0.059] N=814	-0.047 [0.065] N=710	-0.039 [0.076] N=596	-0.102 [0.094] N=465
Sweets consumed at least weekly	-0.021 [0.072] N=893	-0.057 [0.077] N=814	-0.058 [0.088] N=710	-0.133 [0.093] N=596	-0.033 [0.126] N=465
Snack foods consumed at least weekly	-0.037 [0.069] N=893	-0.077 [0.075] N=814	-0.049 [0.086] N=710	-0.076 [0.104] N=596	-0.023 [0.127] N=465
Packaged soft drinks consumed at least weekly	0.050 [0.069] N=893	0.000 [0.074] N=814	-0.025 [0.081] N=710	-0.041 [0.093] N=596	-0.088 [0.128] N=465
Grains consumed at least weekly	0.063 [0.060] N=893	0.071 [0.065] N=814	0.034 [0.071] N=710	0.070 [0.085] N=596	0.018 [0.119] N=465
Child was ever breastfed	-0.011 [0.021] N=1,098	-0.003 [0.025] N=1,005	-0.006 [0.030] N=873	-0.017 [0.036] N=723	-0.012 [0.045] N=566

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=925	**0.074* [0.038] N=998**	**0.077** [0.039] N=898**	0.059 [0.043] N=787	0.039 [0.046] N=658	0.055 [0.056] N=536
Vegetables consumed at least weekly	0.897 P=894	0.056 [0.041] N=998	0.062 [0.043] N=898	0.067 [0.047] N=787	0.042 [0.051] N=658	0.048 [0.054] N=536
Meat fish or poultry consumed at least weekly	0.752 P=759	0.099 [0.062] N=998	0.096 [0.064] N=898	0.087 [0.072] N=787	**0.139* [0.077] N=658**	0.144 [0.091] N=536
Legumes consumed at least weekly	0.747 P=756	-0.001 [0.045] N=998	0.038 [0.048] N=898	0.026 [0.055] N=787	-0.014 [0.058] N=658	0.014 [0.066] N=536
Animal products consumed at least weekly	0.901 P=903	0.019 [0.046] N=998	0.015 [0.050] N=898	0.003 [0.057] N=787	-0.010 [0.062] N=658	0.000 [0.072] N=536
Sweets consumed at least weekly	0.509 P=513	-0.067 [0.065] N=998	-0.076 [0.068] N=898	-0.085 [0.073] N=787	-0.099 [0.074] N=658	**-0.164* [0.084] N=536**
Snack foods consumed at least weekly	0.417 P=417	-0.035 [0.057] N=998	-0.030 [0.061] N=898	-0.007 [0.065] N=787	-0.037 [0.068] N=658	-0.084 [0.076] N=536
Packaged soft drinks consumed at least weekly	0.222 P=219	-0.044 [0.050] N=998	-0.055 [0.052] N=898	0.003 [0.055] N=787	-0.033 [0.057] N=658	-0.045 [0.062] N=536
Grains consumed at least weekly	0.843 P=848	0.059 [0.048] N=998	0.069 [0.052] N=898	0.067 [0.055] N=787	0.068 [0.061] N=658	0.053 [0.074] N=536
Child was ever breastfed	0.966 P=1,173	-0.003 [0.024] N=1,215	-0.008 [0.024] N=1,097	0.003 [0.024] N=959	0.004 [0.024] N=807	0.013 [0.024] N=653

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.054] N=871	0.062 [0.056] N=771	0.036 [0.067] N=660	0.020 [0.075] N=531	0.052 [0.110] N=409
Vegetables consumed at least weekly	0.016 [0.062] N=871	0.016 [0.066] N=771	0.016 [0.078] N=660	-0.025 [0.094] N=531	-0.029 [0.116] N=409
Meat fish or poultry consumed at least weekly	0.014 [0.073] N=871	0.001 [0.074] N=771	-0.045 [0.085] N=660	0.019 [0.095] N=531	-0.023 [0.131] N=409
Legumes consumed at least weekly	-0.021 [0.060] N=871	0.035 [0.065] N=771	0.026 [0.078] N=660	-0.027 [0.086] N=531	0.039 [0.104] N=409
Animal products consumed at least weekly	-0.020 [0.058] N=871	-0.027 [0.065] N=771	-0.049 [0.079] N=660	-0.075 [0.082] N=531	-0.092 [0.104] N=409
Sweets consumed at least weekly	-0.035 [0.092] N=871	-0.047 [0.098] N=771	-0.060 [0.110] N=660	-0.090 [0.127] N=531	-0.268 [0.162] N=409
Snack foods consumed at least weekly	-0.044 [0.082] N=871	-0.036 [0.088] N=771	0.016 [0.097] N=660	-0.033 [0.112] N=531	-0.175 [0.137] N=409
Packaged soft drinks consumed at least weekly	-0.060 [0.077] N=871	-0.076 [0.083] N=771	0.028 [0.095] N=660	-0.046 [0.109] N=531	-0.090 [0.129] N=409
Grains consumed at least weekly	-0.048 [0.051] N=871	-0.049 [0.058] N=771	-0.069 [0.062] N=660	**-0.139** [0.063] N=531**	**-0.313*** [0.077] N=409**
Child was ever breastfed	0.006 [0.043] N=1,059	-0.001 [0.045] N=941	0.019 [0.047] N=803	0.023 [0.053] N=651	0.056 [0.068] N=497

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=925	**0.074* [0.038] N=998**	**0.077** [0.039] N=898**	0.059 [0.044] N=787	0.039 [0.046] N=658	0.055 [0.057] N=536
Vegetables consumed at least weekly	0.897 P=894	0.057 [0.041] N=998	0.062 [0.043] N=898	0.066 [0.047] N=787	0.042 [0.051] N=658	0.048 [0.054] N=536
Meat fish or poultry consumed at least weekly	0.752 P=759	0.099 [0.062] N=998	0.096 [0.064] N=898	0.088 [0.072] N=787	**0.139* [0.077] N=658**	0.144 [0.091] N=536
Legumes consumed at least weekly	0.747 P=756	-0.001 [0.045] N=998	0.038 [0.048] N=898	0.027 [0.055] N=787	-0.013 [0.058] N=658	0.014 [0.066] N=536
Animal products consumed at least weekly	0.901 P=903	0.020 [0.045] N=998	0.016 [0.049] N=898	0.004 [0.056] N=787	-0.009 [0.061] N=658	0.000 [0.071] N=536
Sweets consumed at least weekly	0.509 P=513	-0.066 [0.065] N=998	-0.075 [0.068] N=898	-0.083 [0.073] N=787	-0.098 [0.074] N=658	**-0.164* [0.084] N=536**
Snack foods consumed at least weekly	0.417 P=417	-0.034 [0.056] N=998	-0.029 [0.060] N=898	-0.004 [0.064] N=787	-0.036 [0.067] N=658	-0.084 [0.075] N=536
Packaged soft drinks consumed at least weekly	0.222 P=219	-0.043 [0.050] N=998	-0.054 [0.052] N=898	0.005 [0.054] N=787	-0.032 [0.057] N=658	-0.045 [0.062] N=536
Grains consumed at least weekly	0.843 P=848	0.059 [0.047] N=998	0.070 [0.051] N=898	0.067 [0.054] N=787	0.068 [0.061] N=658	0.053 [0.074] N=536
Child was ever breastfed	0.966 P=1,173	-0.003 [0.024] N=1,215	-0.008 [0.024] N=1,097	0.003 [0.024] N=959	0.004 [0.024] N=807	0.013 [0.024] N=653

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.060 [0.054] N=871	0.062 [0.056] N=771	0.036 [0.067] N=660	0.020 [0.076] N=531	0.051 [0.112] N=409
Vegetables consumed at least weekly	0.016 [0.061] N=871	0.017 [0.066] N=771	0.016 [0.078] N=660	-0.023 [0.093] N=531	-0.029 [0.116] N=409
Meat fish or poultry consumed at least weekly	0.015 [0.072] N=871	0.003 [0.073] N=771	-0.042 [0.083] N=660	0.024 [0.093] N=531	-0.019 [0.129] N=409
Legumes consumed at least weekly	-0.021 [0.060] N=871	0.036 [0.064] N=771	0.027 [0.077] N=660	-0.025 [0.086] N=531	0.034 [0.106] N=409
Animal products consumed at least weekly	-0.019 [0.056] N=871	-0.025 [0.062] N=771	-0.046 [0.077] N=660	-0.068 [0.078] N=531	-0.085 [0.097] N=409
Sweets consumed at least weekly	-0.034 [0.090] N=871	-0.044 [0.096] N=771	-0.055 [0.108] N=660	-0.087 [0.127] N=531	-0.265 [0.161] N=409
Snack foods consumed at least weekly	-0.043 [0.080] N=871	-0.033 [0.086] N=771	0.021 [0.095] N=660	-0.027 [0.111] N=531	-0.169 [0.135] N=409
Packaged soft drinks consumed at least weekly	-0.059 [0.076] N=871	-0.075 [0.082] N=771	0.030 [0.094] N=660	-0.044 [0.109] N=531	-0.086 [0.127] N=409
Grains consumed at least weekly	-0.048 [0.051] N=871	-0.048 [0.059] N=771	-0.069 [0.062] N=660	**-0.142** [0.064] N=531**	**-0.316*** [0.080] N=409**
Child was ever breastfed	0.006 [0.043] N=1,059	-0.001 [0.045] N=941	0.019 [0.047] N=803	0.023 [0.054] N=651	0.056 [0.068] N=497

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=925	**0.071* [0.038] N=998**	**0.073* [0.039] N=898**	0.059 [0.044] N=787	0.039 [0.047] N=658	0.054 [0.057] N=536
Vegetables consumed at least weekly	0.897 P=894	0.054 [0.041] N=998	0.058 [0.043] N=898	0.066 [0.047] N=787	0.042 [0.051] N=658	0.045 [0.055] N=536
Meat fish or poultry consumed at least weekly	0.752 P=759	0.092 [0.060] N=998	0.089 [0.062] N=898	0.091 [0.069] N=787	**0.145* [0.074] N=658**	**0.147* [0.087] N=536**
Legumes consumed at least weekly	0.747 P=756	-0.004 [0.046] N=998	0.034 [0.049] N=898	0.025 [0.056] N=787	-0.014 [0.060] N=658	0.013 [0.070] N=536
Animal products consumed at least weekly	0.901 P=903	0.017 [0.045] N=998	0.011 [0.049] N=898	0.001 [0.056] N=787	-0.011 [0.061] N=658	-0.002 [0.071] N=536
Sweets consumed at least weekly	0.509 P=513	-0.067 [0.065] N=998	-0.079 [0.068] N=898	-0.086 [0.073] N=787	-0.102 [0.075] N=658	**-0.172** [0.085] N=536**
Snack foods consumed at least weekly	0.417 P=417	-0.032 [0.057] N=998	-0.028 [0.061] N=898	-0.007 [0.064] N=787	-0.039 [0.068] N=658	-0.091 [0.075] N=536
Packaged soft drinks consumed at least weekly	0.222 P=219	-0.040 [0.050] N=998	-0.054 [0.052] N=898	-0.001 [0.055] N=787	-0.038 [0.058] N=658	-0.052 [0.063] N=536
Grains consumed at least weekly	0.843 P=848	0.057 [0.047] N=998	0.067 [0.051] N=898	0.067 [0.055] N=787	0.068 [0.063] N=658	0.053 [0.075] N=536
Child was ever breastfed	0.966 P=1,173	-0.004 [0.024] N=1,215	-0.009 [0.025] N=1,097	0.004 [0.024] N=959	0.005 [0.025] N=807	0.012 [0.024] N=653

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.055 [0.053] N=871	0.055 [0.054] N=771	0.037 [0.066] N=660	0.016 [0.077] N=531	0.043 [0.113] N=409
Vegetables consumed at least weekly	0.012 [0.060] N=871	0.009 [0.064] N=771	0.015 [0.075] N=660	-0.028 [0.091] N=531	-0.042 [0.115] N=409
Meat fish or poultry consumed at least weekly	-0.003 [0.071] N=871	-0.017 [0.072] N=771	-0.043 [0.080] N=660	0.012 [0.095] N=531	-0.051 [0.129] N=409
Legumes consumed at least weekly	-0.022 [0.060] N=871	0.035 [0.065] N=771	0.030 [0.076] N=660	-0.029 [0.086] N=531	0.036 [0.109] N=409
Animal products consumed at least weekly	-0.020 [0.055] N=871	-0.030 [0.061] N=771	-0.045 [0.074] N=660	-0.070 [0.078] N=531	-0.094 [0.096] N=409
Sweets consumed at least weekly	-0.030 [0.087] N=871	-0.044 [0.094] N=771	-0.053 [0.105] N=660	-0.084 [0.125] N=531	-0.259 [0.157] N=409
Snack foods consumed at least weekly	-0.037 [0.078] N=871	-0.027 [0.085] N=771	0.020 [0.094] N=660	-0.026 [0.109] N=531	-0.170 [0.132] N=409
Packaged soft drinks consumed at least weekly	-0.044 [0.075] N=871	-0.062 [0.081] N=771	0.034 [0.094] N=660	-0.034 [0.108] N=531	-0.066 [0.124] N=409
Grains consumed at least weekly	-0.050 [0.052] N=871	-0.051 [0.059] N=771	-0.067 [0.062] N=660	**-0.144** [0.063] N=531**	**-0.317*** [0.079] N=409**
Child was ever breastfed	0.004 [0.042] N=1,059	-0.004 [0.044] N=941	0.019 [0.047] N=803	0.020 [0.053] N=651	0.051 [0.065] N=497

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.928 P=925	**0.066* [0.038] N=998**	**0.069* [0.039] N=898**	0.055 [0.044] N=787	0.028 [0.046] N=658	0.039 [0.056] N=536
Vegetables consumed at least weekly	0.897 P=894	0.049 [0.041] N=998	0.055 [0.043] N=898	0.061 [0.047] N=787	0.031 [0.050] N=658	0.035 [0.053] N=536
Meat fish or poultry consumed at least weekly	0.752 P=759	0.087 [0.061] N=998	0.081 [0.062] N=898	0.078 [0.069] N=787	0.118 [0.075] N=658	0.108 [0.085] N=536
Legumes consumed at least weekly	0.747 P=756	-0.002 [0.046] N=998	0.036 [0.049] N=898	0.030 [0.055] N=787	-0.014 [0.058] N=658	0.008 [0.066] N=536
Animal products consumed at least weekly	0.901 P=903	0.021 [0.044] N=998	0.018 [0.047] N=898	0.013 [0.055] N=787	-0.005 [0.060] N=658	0.000 [0.070] N=536
Sweets consumed at least weekly	0.509 P=513	-0.058 [0.066] N=998	-0.070 [0.069] N=898	-0.079 [0.074] N=787	-0.097 [0.076] N=658	**-0.165* [0.086] N=536**
Snack foods consumed at least weekly	0.417 P=417	-0.026 [0.057] N=998	-0.022 [0.062] N=898	0.002 [0.065] N=787	-0.032 [0.068] N=658	-0.078 [0.077] N=536
Packaged soft drinks consumed at least weekly	0.222 P=219	-0.032 [0.049] N=998	-0.047 [0.052] N=898	0.008 [0.055] N=787	-0.027 [0.059] N=658	-0.028 [0.065] N=536
Grains consumed at least weekly	0.843 P=848	0.053 [0.048] N=998	0.062 [0.052] N=898	0.062 [0.055] N=787	0.060 [0.062] N=658	0.043 [0.074] N=536
Child was ever breastfed	0.966 P=1,173	-0.003 [0.023] N=1,215	-0.009 [0.024] N=1,097	0.003 [0.023] N=959	0.003 [0.023] N=807	0.011 [0.023] N=653

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.049 [0.053] N=871	0.049 [0.055] N=771	0.032 [0.066] N=660	-0.004 [0.076] N=531	0.008 [0.109] N=409
Vegetables consumed at least weekly	0.004 [0.061] N=871	0.002 [0.065] N=771	0.007 [0.076] N=660	-0.047 [0.090] N=531	-0.073 [0.112] N=409
Meat fish or poultry consumed at least weekly	-0.002 [0.073] N=871	-0.020 [0.076] N=771	-0.052 [0.085] N=660	-0.024 [0.102] N=531	-0.113 [0.129] N=409
Legumes consumed at least weekly	-0.025 [0.061] N=871	0.029 [0.067] N=771	0.037 [0.079] N=660	-0.029 [0.087] N=531	0.019 [0.109] N=409
Animal products consumed at least weekly	-0.021 [0.052] N=871	-0.028 [0.059] N=771	-0.033 [0.073] N=660	-0.069 [0.077] N=531	-0.100 [0.094] N=409
Sweets consumed at least weekly	-0.021 [0.089] N=871	-0.034 [0.095] N=771	-0.040 [0.106] N=660	-0.072 [0.124] N=531	-0.239 [0.158] N=409
Snack foods consumed at least weekly	-0.035 [0.079] N=871	-0.027 [0.087] N=771	0.030 [0.096] N=660	-0.023 [0.109] N=531	-0.155 [0.133] N=409
Packaged soft drinks consumed at least weekly	-0.047 [0.075] N=871	-0.069 [0.081] N=771	0.030 [0.092] N=660	-0.032 [0.108] N=531	-0.036 [0.127] N=409
Grains consumed at least weekly	-0.053 [0.052] N=871	-0.057 [0.059] N=771	-0.074 [0.063] N=660	**-0.159** [0.065] N=531**	**-0.337*** [0.079] N=409**
Child was ever breastfed	0.004 [0.042] N=1,059	-0.004 [0.044] N=941	0.020 [0.045] N=803	0.018 [0.050] N=651	0.043 [0.062] N=497

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=844	**0.072* [0.041] N=909**	**0.073* [0.042] N=815**	0.064 [0.047] N=721	0.024 [0.051] N=602	0.034 [0.063] N=495
Vegetables consumed at least weekly	0.900 P=817	0.046 [0.042] N=909	0.048 [0.045] N=815	0.071 [0.047] N=721	0.028 [0.052] N=602	0.008 [0.057] N=495
Meat fish or poultry consumed at least weekly	0.753 P=691	0.081 [0.064] N=909	0.065 [0.066] N=815	0.070 [0.074] N=721	0.099 [0.081] N=602	0.102 [0.095] N=495
Legumes consumed at least weekly	0.744 P=686	-0.010 [0.052] N=909	0.029 [0.055] N=815	0.018 [0.062] N=721	-0.038 [0.067] N=602	-0.010 [0.077] N=495
Animal products consumed at least weekly	0.905 P=826	0.014 [0.049] N=909	0.012 [0.053] N=815	0.012 [0.061] N=721	-0.020 [0.068] N=602	-0.022 [0.080] N=495
Sweets consumed at least weekly	0.520 P=476	-0.060 [0.066] N=909	-0.067 [0.070] N=815	-0.085 [0.075] N=721	-0.117 [0.080] N=602	**-0.180** [0.091] N=495**
Snack foods consumed at least weekly	0.424 P=385	-0.031 [0.060] N=909	-0.027 [0.064] N=815	-0.015 [0.067] N=721	-0.056 [0.071] N=602	-0.088 [0.079] N=495
Packaged soft drinks consumed at least weekly	0.227 P=205	-0.051 [0.051] N=909	-0.068 [0.053] N=815	-0.009 [0.055] N=721	-0.057 [0.060] N=602	-0.083 [0.065] N=495
Grains consumed at least weekly	0.845 P=774	0.074 [0.051] N=909	0.086 [0.056] N=815	0.093 [0.058] N=721	0.078 [0.068] N=602	0.051 [0.080] N=495
Child was ever breastfed	0.968 P=1,073	-0.013 [0.023] N=1,108	-0.019 [0.024] N=999	-0.004 [0.021] N=880	-0.009 [0.022] N=741	0.005 [0.021] N=604

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.057 [0.057] N=792	0.058 [0.058] N=698	0.050 [0.068] N=604	-0.006 [0.076] N=485	-0.009 [0.106] N=378
Vegetables consumed at least weekly	0.004 [0.056] N=792	0.000 [0.062] N=698	0.030 [0.068] N=604	-0.046 [0.081] N=485	-0.129 [0.095] N=378
Meat fish or poultry consumed at least weekly	0.017 [0.079] N=792	-0.009 [0.081] N=698	-0.032 [0.091] N=604	-0.028 [0.105] N=485	-0.099 [0.140] N=378
Legumes consumed at least weekly	-0.021 [0.067] N=792	0.037 [0.072] N=698	0.040 [0.083] N=604	-0.034 [0.095] N=485	0.017 [0.116] N=378
Animal products consumed at least weekly	-0.032 [0.060] N=792	-0.033 [0.067] N=698	-0.032 [0.080] N=604	-0.089 [0.086] N=485	-0.142 [0.108] N=378
Sweets consumed at least weekly	-0.024 [0.093] N=792	-0.032 [0.101] N=698	-0.072 [0.111] N=604	-0.146 [0.129] N=485	**-0.345** [0.166] N=378**
Snack foods consumed at least weekly	-0.042 [0.084] N=792	-0.033 [0.093] N=698	-0.006 [0.100] N=604	-0.094 [0.112] N=485	-0.202 [0.139] N=378
Packaged soft drinks consumed at least weekly	-0.071 [0.079] N=792	-0.096 [0.086] N=698	0.001 [0.098] N=604	-0.103 [0.117] N=485	-0.211 [0.138] N=378
Grains consumed at least weekly	-0.028 [0.059] N=792	-0.025 [0.068] N=698	-0.025 [0.073] N=604	**-0.134* [0.081] N=485**	**-0.362*** [0.099] N=378**
Child was ever breastfed	-0.009 [0.041] N=964	-0.019 [0.044] N=855	0.007 [0.043] N=736	-0.002 [0.048] N=597	0.035 [0.059] N=460

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.930 P=844	**0.075* [0.040] N=909**	**0.078* [0.042] N=815**	0.069 [0.047] N=721	0.024 [0.050] N=602	0.034 [0.061] N=495
Vegetables consumed at least weekly	0.900 P=817	0.050 [0.041] N=909	0.052 [0.044] N=815	**0.077* [0.046] N=721**	0.028 [0.050] N=602	0.007 [0.054] N=495
Meat fish or poultry consumed at least weekly	0.753 P=691	0.089 [0.062] N=909	0.075 [0.064] N=815	0.077 [0.071] N=721	0.093 [0.078] N=602	0.098 [0.092] N=495
Legumes consumed at least weekly	0.744 P=686	-0.009 [0.051] N=909	0.031 [0.054] N=815	0.023 [0.061] N=721	-0.043 [0.064] N=602	-0.014 [0.075] N=495
Animal products consumed at least weekly	0.905 P=826	0.014 [0.048] N=909	0.013 [0.053] N=815	0.012 [0.060] N=721	-0.019 [0.066] N=602	-0.020 [0.078] N=495
Sweets consumed at least weekly	0.520 P=476	-0.062 [0.065] N=909	-0.070 [0.069] N=815	-0.090 [0.074] N=721	-0.123 [0.078] N=602	**-0.185** [0.090] N=495**
Snack foods consumed at least weekly	0.424 P=385	-0.026 [0.059] N=909	-0.022 [0.064] N=815	-0.009 [0.066] N=721	-0.052 [0.070] N=602	-0.082 [0.078] N=495
Packaged soft drinks consumed at least weekly	0.227 P=205	-0.051 [0.051] N=909	-0.067 [0.053] N=815	-0.010 [0.055] N=721	-0.058 [0.059] N=602	-0.085 [0.064] N=495
Grains consumed at least weekly	0.845 P=774	0.074 [0.051] N=909	0.086 [0.055] N=815	0.091 [0.058] N=721	0.076 [0.067] N=602	0.050 [0.079] N=495
Child was ever breastfed	0.968 P=1,073	-0.012 [0.023] N=1,108	-0.018 [0.024] N=999	-0.002 [0.022] N=880	-0.005 [0.022] N=741	0.008 [0.022] N=604

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.062 [0.056] N=792	0.066 [0.058] N=698	0.061 [0.069] N=604	-0.007 [0.077] N=485	0.001 [0.113] N=378
Vegetables consumed at least weekly	0.008 [0.056] N=792	0.005 [0.062] N=698	0.040 [0.069] N=604	-0.051 [0.080] N=485	-0.126 [0.100] N=378
Meat fish or poultry consumed at least weekly	0.024 [0.078] N=792	-0.003 [0.080] N=698	-0.026 [0.090] N=604	-0.036 [0.103] N=485	-0.088 [0.145] N=378
Legumes consumed at least weekly	-0.018 [0.067] N=792	0.043 [0.071] N=698	0.048 [0.082] N=604	-0.045 [0.093] N=485	0.025 [0.115] N=378
Animal products consumed at least weekly	-0.030 [0.060] N=792	-0.031 [0.067] N=698	-0.032 [0.080] N=604	-0.094 [0.086] N=485	-0.152 [0.111] N=378
Sweets consumed at least weekly	-0.022 [0.093] N=792	-0.030 [0.101] N=698	-0.068 [0.112] N=604	-0.140 [0.131] N=485	**-0.337* [0.171] N=378**
Snack foods consumed at least weekly	-0.037 [0.084] N=792	-0.029 [0.093] N=698	0.005 [0.101] N=604	-0.072 [0.115] N=485	-0.189 [0.144] N=378
Packaged soft drinks consumed at least weekly	-0.068 [0.080] N=792	-0.091 [0.087] N=698	0.014 [0.097] N=604	-0.085 [0.114] N=485	-0.184 [0.136] N=378
Grains consumed at least weekly	-0.028 [0.059] N=792	-0.025 [0.067] N=698	-0.029 [0.070] N=604	**-0.142* [0.078] N=485**	**-0.367*** [0.097] N=378**
Child was ever breastfed	-0.010 [0.041] N=964	-0.021 [0.044] N=855	0.008 [0.044] N=736	0.004 [0.051] N=597	0.038 [0.062] N=460

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.