Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-2: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.765	0.185 [0.253] N=1,206	0.101 [0.267] N=1,119	0.169 [0.293] N=998	0.323 [0.302] N=859	0.199 [0.334] N=715
WHO weight-for-age z score	-0.299	-0.170 [0.139] N=1,405	-0.148 [0.153] N=1,302	-0.102 [0.160] N=1,163	-0.055 [0.159] N=996	-0.176 [0.174] N=822
WHO weight-for-length/height z score, 0-60 months	0.298	0.236 [0.226] N=1,200	0.251 [0.236] N=1,111	0.267 [0.241] N=993	0.181 [0.255] N=854	0.200 [0.283] N=705
WHO BMI-for-age z score	0.282	0.273 [0.258] N=1,186	0.313 [0.271] N=1,100	0.320 [0.287] N=981	0.173 [0.305] N=841	0.157 [0.339] N=696
Stunted: HAZ below -2	0.305 P=359	-0.059 [0.046] N=1,206	-0.056 [0.049] N=1,119	-0.048 [0.052] N=998	-0.070 [0.055] N=859	-0.040 [0.058] N=715
Underweight: WAZ below -2	0.118 P=161	**0.088*** [0.028] N=1,405**	**0.074** [0.029] N=1,302**	**0.057* [0.031] N=1,163**	**0.063** [0.031] N=996**	**0.069* [0.036] N=822**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=143	-0.052 [0.039] N=1,200	-0.042 [0.040] N=1,111	-0.053 [0.043] N=993	-0.037 [0.047] N=854	-0.048 [0.050] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=233	0.028 [0.048] N=1,200	0.034 [0.050] N=1,111	0.029 [0.052] N=993	0.024 [0.058] N=854	0.003 [0.064] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=106	-0.022 [0.036] N=1,200	-0.017 [0.039] N=1,111	-0.003 [0.041] N=993	0.003 [0.045] N=854	0.007 [0.051] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=458	**-0.086* [0.046] N=1,150**	-0.078 [0.048] N=1,065	-0.074 [0.051] N=951	-0.091 [0.056] N=820	-0.057 [0.061] N=680
BMI-for-age z score below -2	0.141 P=163	-0.053 [0.044] N=1,186	-0.045 [0.046] N=1,100	-0.045 [0.050] N=981	-0.008 [0.054] N=841	0.004 [0.059] N=696

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.099 [0.369] N=1,016	-0.085 [0.413] N=929	0.103 [0.490] N=808	0.493 [0.541] N=669	0.387 [0.676] N=525
WHO weight-for-age z score	**-0.403** [0.195] N=1,189**	**-0.389* [0.232] N=1,086**	-0.331 [0.254] N=947	-0.287 [0.285] N=780	**-0.670* [0.379] N=606**
WHO weight-for-length/height z score, 0-60 months	0.222 [0.330] N=1,010	0.247 [0.358] N=921	0.229 [0.378] N=803	0.076 [0.437] N=664	-0.083 [0.542] N=515
WHO BMI-for-age z score	0.311 [0.346] N=996	0.370 [0.370] N=910	0.317 [0.408] N=791	0.013 [0.458] N=651	-0.256 [0.545] N=506
Stunted: HAZ below -2	-0.067 [0.065] N=1,016	-0.063 [0.072] N=929	-0.064 [0.082] N=808	-0.121 [0.093] N=669	-0.117 [0.112] N=525
Underweight: WAZ below -2	**0.123*** [0.042] N=1,189**	**0.102** [0.046] N=1,086**	0.072 [0.052] N=947	0.095 [0.060] N=780	0.125 [0.081] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.073 [0.053] N=1,010	-0.056 [0.057] N=921	-0.080 [0.063] N=803	-0.060 [0.075] N=664	-0.099 [0.092] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.003 [0.061] N=1,010	0.006 [0.068] N=921	-0.018 [0.067] N=803	-0.031 [0.086] N=664	-0.166 [0.101] N=515
Obese: WLZ/WHZ above +3, 0-60 months	-0.060 [0.044] N=1,010	-0.058 [0.051] N=921	-0.042 [0.052] N=803	-0.036 [0.065] N=664	-0.068 [0.087] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.090 [0.066] N=968	-0.077 [0.072] N=883	-0.083 [0.078] N=769	-0.134 [0.092] N=638	-0.110 [0.116] N=498
BMI-for-age z score below -2	**-0.108* [0.056] N=996**	**-0.102* [0.058] N=910**	**-0.112* [0.067] N=791**	-0.060 [0.079] N=651	-0.060 [0.097] N=506

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.765	0.185 [0.253] N=1,206	0.098 [0.267] N=1,119	0.169 [0.293] N=998	0.327 [0.303] N=859	0.206 [0.333] N=715
WHO weight-for-age z score	-0.299	-0.176 [0.139] N=1,405	-0.157 [0.153] N=1,302	-0.109 [0.160] N=1,163	-0.061 [0.159] N=996	-0.188 [0.173] N=822
WHO weight-for-length/height z score, 0-60 months	0.298	0.231 [0.226] N=1,200	0.248 [0.236] N=1,111	0.263 [0.242] N=993	0.177 [0.256] N=854	0.190 [0.284] N=705
WHO BMI-for-age z score	0.282	0.274 [0.258] N=1,186	0.316 [0.270] N=1,100	0.318 [0.287] N=981	0.172 [0.306] N=841	0.154 [0.339] N=696
Stunted: HAZ below -2	0.305 P=359	-0.057 [0.046] N=1,206	-0.054 [0.048] N=1,119	-0.046 [0.051] N=998	-0.068 [0.054] N=859	-0.038 [0.058] N=715
Underweight: WAZ below -2	0.118 P=161	**0.089*** [0.028] N=1,405**	**0.075** [0.029] N=1,302**	**0.058* [0.031] N=1,163**	**0.063** [0.031] N=996**	**0.071** [0.035] N=822**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=143	-0.050 [0.039] N=1,200	-0.039 [0.040] N=1,111	-0.050 [0.042] N=993	-0.034 [0.047] N=854	-0.043 [0.049] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=233	0.028 [0.048] N=1,200	0.035 [0.050] N=1,111	0.030 [0.052] N=993	0.026 [0.058] N=854	0.006 [0.064] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=106	-0.021 [0.036] N=1,200	-0.016 [0.038] N=1,111	-0.002 [0.041] N=993	0.004 [0.045] N=854	0.009 [0.051] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=458	**-0.082* [0.045] N=1,150**	-0.073 [0.047] N=1,065	-0.070 [0.050] N=951	-0.087 [0.055] N=820	-0.049 [0.059] N=680
BMI-for-age z score below -2	0.141 P=163	-0.051 [0.045] N=1,186	-0.043 [0.046] N=1,100	-0.043 [0.051] N=981	-0.006 [0.055] N=841	0.007 [0.059] N=696

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.098 [0.369] N=1,016	-0.085 [0.413] N=929	0.100 [0.490] N=808	0.480 [0.543] N=669	0.369 [0.679] N=525
WHO weight-for-age z score	**-0.399** [0.195] N=1,189**	**-0.390* [0.234] N=1,086**	-0.325 [0.256] N=947	-0.285 [0.287] N=780	**-0.667* [0.379] N=606**
WHO weight-for-length/height z score, 0-60 months	0.221 [0.330] N=1,010	0.246 [0.359] N=921	0.232 [0.379] N=803	0.079 [0.441] N=664	-0.077 [0.544] N=515
WHO BMI-for-age z score	0.311 [0.346] N=996	0.370 [0.370] N=910	0.320 [0.409] N=791	0.016 [0.461] N=651	-0.253 [0.546] N=506
Stunted: HAZ below -2	-0.068 [0.065] N=1,016	-0.064 [0.071] N=929	-0.067 [0.082] N=808	-0.123 [0.093] N=669	-0.118 [0.112] N=525
Underweight: WAZ below -2	**0.122*** [0.042] N=1,189**	**0.102** [0.046] N=1,086**	0.072 [0.053] N=947	0.095 [0.060] N=780	0.125 [0.081] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.073 [0.053] N=1,010	-0.055 [0.057] N=921	-0.082 [0.063] N=803	-0.062 [0.076] N=664	-0.102 [0.092] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.003 [0.061] N=1,010	0.006 [0.068] N=921	-0.018 [0.067] N=803	-0.032 [0.087] N=664	-0.167 [0.101] N=515
Obese: WLZ/WHZ above +3, 0-60 months	-0.060 [0.044] N=1,010	-0.057 [0.050] N=921	-0.042 [0.051] N=803	-0.036 [0.064] N=664	-0.069 [0.087] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.092 [0.066] N=968	-0.079 [0.071] N=883	-0.091 [0.077] N=769	-0.144 [0.093] N=638	-0.121 [0.118] N=498
BMI-for-age z score below -2	**-0.108* [0.056] N=996**	**-0.101* [0.058] N=910**	**-0.113* [0.067] N=791**	-0.061 [0.080] N=651	-0.061 [0.098] N=506

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.765	0.156 [0.254] N=1,206	0.061 [0.267] N=1,119	0.122 [0.292] N=998	0.298 [0.302] N=859	0.157 [0.337] N=715
WHO weight-for-age z score	-0.299	-0.210 [0.132] N=1,405	-0.194 [0.147] N=1,302	-0.150 [0.154] N=1,163	-0.083 [0.155] N=996	-0.225 [0.170] N=822
WHO weight-for-length/height z score, 0-60 months	0.298	0.228 [0.227] N=1,200	0.252 [0.239] N=1,111	0.275 [0.245] N=993	0.187 [0.258] N=854	0.199 [0.293] N=705
WHO BMI-for-age z score	0.282	0.279 [0.260] N=1,186	0.321 [0.274] N=1,100	0.333 [0.291] N=981	0.187 [0.310] N=841	0.168 [0.351] N=696
Stunted: HAZ below -2	0.305 P=359	-0.052 [0.046] N=1,206	-0.045 [0.048] N=1,119	-0.037 [0.051] N=998	-0.064 [0.054] N=859	-0.026 [0.057] N=715
Underweight: WAZ below -2	0.118 P=161	**0.094*** [0.027] N=1,405**	**0.082*** [0.028] N=1,302**	**0.065** [0.030] N=1,163**	**0.068** [0.030] N=996**	**0.080** [0.034] N=822**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=143	-0.048 [0.038] N=1,200	-0.038 [0.040] N=1,111	-0.050 [0.042] N=993	-0.035 [0.046] N=854	-0.042 [0.049] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=233	0.029 [0.048] N=1,200	0.037 [0.050] N=1,111	0.031 [0.052] N=993	0.027 [0.057] N=854	0.008 [0.064] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=106	-0.021 [0.036] N=1,200	-0.015 [0.039] N=1,111	-0.002 [0.041] N=993	0.005 [0.045] N=854	0.009 [0.051] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=458	-0.075 [0.045] N=1,150	-0.061 [0.047] N=1,065	-0.060 [0.050] N=951	-0.081 [0.055] N=820	-0.032 [0.060] N=680
BMI-for-age z score below -2	0.141 P=163	-0.051 [0.045] N=1,186	-0.041 [0.047] N=1,100	-0.043 [0.052] N=981	-0.007 [0.055] N=841	0.007 [0.060] N=696

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.083 [0.368] N=1,016	-0.118 [0.406] N=929	0.041 [0.477] N=808	0.456 [0.525] N=669	0.388 [0.665] N=525
WHO weight-for-age z score	**-0.402** [0.191] N=1,189**	**-0.398* [0.229] N=1,086**	-0.340 [0.250] N=947	-0.259 [0.289] N=780	**-0.645* [0.381] N=606**
WHO weight-for-length/height z score, 0-60 months	0.214 [0.326] N=1,010	0.246 [0.358] N=921	0.238 [0.372] N=803	0.048 [0.430] N=664	-0.136 [0.543] N=515
WHO BMI-for-age z score	0.313 [0.346] N=996	0.367 [0.372] N=910	0.328 [0.404] N=791	-0.008 [0.447] N=651	-0.290 [0.543] N=506
Stunted: HAZ below -2	-0.068 [0.064] N=1,016	-0.060 [0.071] N=929	-0.059 [0.081] N=808	-0.130 [0.090] N=669	-0.127 [0.109] N=525
Underweight: WAZ below -2	**0.121*** [0.041] N=1,189**	**0.102** [0.046] N=1,086**	0.072 [0.052] N=947	0.084 [0.060] N=780	0.120 [0.081] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.073 [0.052] N=1,010	-0.055 [0.057] N=921	-0.087 [0.062] N=803	-0.068 [0.076] N=664	-0.100 [0.093] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.061] N=1,010	0.004 [0.069] N=921	-0.026 [0.067] N=803	-0.049 [0.086] N=664	**-0.175* [0.103] N=515**
Obese: WLZ/WHZ above +3, 0-60 months	-0.063 [0.044] N=1,010	-0.060 [0.051] N=921	-0.048 [0.052] N=803	-0.047 [0.066] N=664	-0.077 [0.089] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.097 [0.066] N=968	-0.080 [0.072] N=883	-0.098 [0.076] N=769	**-0.172* [0.092] N=638**	-0.143 [0.116] N=498
BMI-for-age z score below -2	**-0.110* [0.056] N=996**	**-0.100* [0.059] N=910**	**-0.119* [0.068] N=791**	-0.073 [0.080] N=651	-0.060 [0.099] N=506

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.765	0.144 [0.253] N=1,206	0.068 [0.267] N=1,119	0.131 [0.294] N=998	0.321 [0.304] N=859	0.204 [0.336] N=715
WHO weight-for-age z score	-0.299	**-0.234* [0.131] N=1,405**	-0.216 [0.145] N=1,302	-0.176 [0.154] N=1,163	-0.123 [0.154] N=996	**-0.291* [0.173] N=822**
WHO weight-for-length/height z score, 0-60 months	0.298	0.223 [0.229] N=1,200	0.240 [0.240] N=1,111	0.259 [0.246] N=993	0.157 [0.259] N=854	0.133 [0.293] N=705
WHO BMI-for-age z score	0.282	0.269 [0.260] N=1,186	0.308 [0.272] N=1,100	0.319 [0.289] N=981	0.159 [0.308] N=841	0.100 [0.348] N=696
Stunted: HAZ below -2	0.305 P=359	-0.046 [0.046] N=1,206	-0.044 [0.048] N=1,119	-0.035 [0.051] N=998	-0.062 [0.054] N=859	-0.027 [0.056] N=715
Underweight: WAZ below -2	0.118 P=161	**0.099*** [0.027] N=1,405**	**0.086*** [0.028] N=1,302**	**0.073** [0.030] N=1,163**	**0.078*** [0.029] N=996**	**0.094*** [0.034] N=822**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=143	-0.048 [0.038] N=1,200	-0.036 [0.040] N=1,111	-0.046 [0.042] N=993	-0.031 [0.047] N=854	-0.039 [0.049] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=233	0.030 [0.048] N=1,200	0.039 [0.051] N=1,111	0.036 [0.053] N=993	0.029 [0.058] N=854	0.003 [0.066] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=106	-0.022 [0.036] N=1,200	-0.018 [0.039] N=1,111	-0.004 [0.041] N=993	0.001 [0.044] N=854	-0.001 [0.050] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=458	-0.064 [0.046] N=1,150	-0.052 [0.047] N=1,065	-0.046 [0.051] N=951	-0.067 [0.056] N=820	-0.018 [0.061] N=680
BMI-for-age z score below -2	0.141 P=163	-0.047 [0.045] N=1,186	-0.036 [0.047] N=1,100	-0.035 [0.051] N=981	0.005 [0.055] N=841	0.021 [0.061] N=696

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.045 [0.372] N=1,016	-0.131 [0.412] N=929	0.026 [0.491] N=808	0.474 [0.549] N=669	0.395 [0.680] N=525
WHO weight-for-age z score	**-0.413** [0.190] N=1,189**	**-0.399* [0.226] N=1,086**	-0.345 [0.249] N=947	-0.223 [0.289] N=780	-0.588 [0.377] N=606
WHO weight-for-length/height z score, 0-60 months	0.246 [0.327] N=1,010	0.277 [0.354] N=921	0.264 [0.376] N=803	0.105 [0.434] N=664	-0.074 [0.533] N=515
WHO BMI-for-age z score	0.339 [0.346] N=996	0.396 [0.367] N=910	0.364 [0.410] N=791	0.066 [0.453] N=651	-0.203 [0.538] N=506
Stunted: HAZ below -2	-0.055 [0.065] N=1,016	-0.054 [0.071] N=929	-0.052 [0.081] N=808	-0.124 [0.092] N=669	-0.110 [0.110] N=525
Underweight: WAZ below -2	**0.124*** [0.041] N=1,189**	**0.103** [0.045] N=1,086**	0.075 [0.051] N=947	0.081 [0.060] N=780	0.122 [0.081] N=606
Wasted: WLZ/WHZ below -2, 0-60 months	-0.074 [0.053] N=1,010	-0.059 [0.057] N=921	-0.087 [0.063] N=803	-0.064 [0.076] N=664	-0.098 [0.093] N=515
Overweight: WLZ/WHZ above +2, 0-60 months	-0.005 [0.061] N=1,010	0.006 [0.068] N=921	-0.021 [0.066] N=803	-0.038 [0.086] N=664	-0.158 [0.096] N=515
Obese: WLZ/WHZ above +3, 0-60 months	-0.056 [0.044] N=1,010	-0.053 [0.050] N=921	-0.039 [0.051] N=803	-0.037 [0.063] N=664	-0.071 [0.082] N=515
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.085 [0.067] N=968	-0.075 [0.072] N=883	-0.086 [0.078] N=769	**-0.161* [0.095] N=638**	-0.126 [0.124] N=498
BMI-for-age z score below -2	**-0.114** [0.056] N=996**	**-0.107* [0.058] N=910**	**-0.124* [0.068] N=791**	-0.084 [0.080] N=651	-0.075 [0.100] N=506

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.774	0.145 [0.269] N=1,111	0.062 [0.285] N=1,034	0.127 [0.314] N=920	0.250 [0.320] N=799	0.128 [0.338] N=668
WHO weight-for-age z score	-0.293	-0.083 [0.141] N=1,285	-0.078 [0.153] N=1,194	-0.021 [0.160] N=1,064	0.021 [0.157] N=917	-0.087 [0.171] N=762
WHO weight-for-length/height z score, 0-60 months	0.312	**0.374* [0.224] N=1,105**	**0.390* [0.232] N=1,027**	**0.416* [0.233] N=917**	0.347 [0.246] N=796	0.423 [0.266] N=660
WHO BMI-for-age z score	0.300	0.418 [0.263] N=1,093	0.428 [0.270] N=1,016	0.447 [0.284] N=906	0.330 [0.298] N=784	0.370 [0.329] N=652
Stunted: HAZ below -2	0.306 P=330	-0.053 [0.051] N=1,111	-0.052 [0.054] N=1,034	-0.041 [0.057] N=920	-0.057 [0.061] N=799	-0.021 [0.064] N=668
Underweight: WAZ below -2	0.119 P=149	**0.078*** [0.029] N=1,285**	**0.067** [0.030] N=1,194**	0.050 [0.032] N=1,064	**0.058* [0.033] N=917**	**0.070* [0.036] N=762**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=130	**-0.078** [0.038] N=1,105**	-0.064 [0.039] N=1,027	**-0.070* [0.041] N=917**	-0.056 [0.045] N=796	-0.067 [0.047] N=660
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=216	0.043 [0.048] N=1,105	0.051 [0.051] N=1,027	0.051 [0.053] N=917	0.048 [0.059] N=796	0.044 [0.066] N=660
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=101	-0.022 [0.038] N=1,105	-0.014 [0.041] N=1,027	0.007 [0.043] N=917	0.020 [0.047] N=796	0.033 [0.053] N=660
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=420	**-0.099* [0.052] N=1,059**	-0.087 [0.054] N=984	-0.080 [0.057] N=877	-0.097 [0.063] N=764	-0.056 [0.069] N=636
BMI-for-age z score below -2	0.140 P=147	**-0.077* [0.045] N=1,093**	-0.063 [0.046] N=1,016	-0.064 [0.051] N=906	-0.028 [0.055] N=784	-0.024 [0.060] N=652

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.180 [0.409] N=934	0.031 [0.463] N=857	0.250 [0.550] N=743	0.575 [0.616] N=622	0.560 [0.750] N=491
WHO weight-for-age z score	-0.326 [0.199] N=1,085	-0.343 [0.232] N=994	-0.273 [0.255] N=864	-0.259 [0.272] N=717	**-0.612* [0.351] N=562**
WHO weight-for-length/height z score, 0-60 months	0.256 [0.349] N=928	0.237 [0.375] N=850	0.200 [0.394] N=740	0.098 [0.469] N=619	0.018 [0.566] N=483
WHO BMI-for-age z score	0.308 [0.378] N=916	0.272 [0.397] N=839	0.204 [0.428] N=729	-0.040 [0.486] N=607	-0.303 [0.573] N=475
Stunted: HAZ below -2	-0.099 [0.071] N=934	-0.107 [0.079] N=857	-0.114 [0.089] N=743	-0.171 [0.104] N=622	-0.178 [0.124] N=491
Underweight: WAZ below -2	**0.110** [0.043] N=1,085**	**0.093* [0.048] N=994**	0.065 [0.054] N=864	0.097 [0.062] N=717	**0.142* [0.081] N=562**
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.111** [0.054] N=928**	-0.087 [0.057] N=850	**-0.107* [0.062] N=740**	-0.105 [0.074] N=619	**-0.166* [0.088] N=483**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.009 [0.063] N=928	-0.007 [0.069] N=850	-0.030 [0.068] N=740	-0.046 [0.090] N=619	-0.161 [0.108] N=483
Obese: WLZ/WHZ above +3, 0-60 months	**-0.082* [0.048] N=928**	-0.077 [0.056] N=850	-0.052 [0.057] N=740	-0.034 [0.071] N=619	-0.057 [0.093] N=483
CIAF: any stunting, underweight, or wasting, 0-60 months	**-0.144** [0.073] N=890**	**-0.134* [0.079] N=815**	**-0.152* [0.087] N=708**	**-0.227** [0.102] N=595**	**-0.230* [0.127] N=467**
BMI-for-age z score below -2	**-0.136** [0.059] N=916**	**-0.120** [0.060] N=839**	**-0.130* [0.068] N=729**	-0.093 [0.081] N=607	-0.118 [0.101] N=475

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.774	0.156 [0.267] N=1,111	0.074 [0.283] N=1,034	0.134 [0.313] N=920	0.256 [0.319] N=799	0.130 [0.336] N=668
WHO weight-for-age z score	-0.293	-0.070 [0.140] N=1,285	-0.065 [0.154] N=1,194	-0.014 [0.160] N=1,064	0.012 [0.159] N=917	-0.096 [0.177] N=762
WHO weight-for-length/height z score, 0-60 months	0.312	**0.369* [0.222] N=1,105**	**0.384* [0.230] N=1,027**	**0.403* [0.230] N=917**	0.320 [0.244] N=796	0.381 [0.267] N=660
WHO BMI-for-age z score	0.300	0.410 [0.260] N=1,093	0.416 [0.267] N=1,016	0.437 [0.281] N=906	0.313 [0.296] N=784	0.347 [0.326] N=652
Stunted: HAZ below -2	0.306 P=330	-0.052 [0.050] N=1,111	-0.051 [0.053] N=1,034	-0.040 [0.056] N=920	-0.059 [0.060] N=799	-0.024 [0.062] N=668
Underweight: WAZ below -2	0.119 P=149	**0.077*** [0.029] N=1,285**	**0.066** [0.030] N=1,194**	0.050 [0.032] N=1,064	**0.061* [0.033] N=917**	**0.070* [0.037] N=762**
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=130	**-0.075* [0.038] N=1,105**	-0.061 [0.039] N=1,027	**-0.069* [0.041] N=917**	-0.054 [0.045] N=796	-0.067 [0.048] N=660
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=216	0.044 [0.048] N=1,105	0.051 [0.050] N=1,027	0.048 [0.053] N=917	0.039 [0.059] N=796	0.031 [0.065] N=660
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=101	-0.021 [0.038] N=1,105	-0.012 [0.041] N=1,027	0.007 [0.043] N=917	0.018 [0.047] N=796	0.026 [0.053] N=660
CIAF: any stunting, underweight, or wasting, 0-60 months	0.407 P=420	**-0.095* [0.051] N=1,059**	-0.085 [0.053] N=984	-0.080 [0.057] N=877	-0.100 [0.062] N=764	-0.060 [0.067] N=636
BMI-for-age z score below -2	0.140 P=147	-0.072 [0.045] N=1,093	-0.060 [0.046] N=1,016	-0.062 [0.051] N=906	-0.028 [0.055] N=784	-0.023 [0.060] N=652

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.187 [0.408] N=934	0.039 [0.464] N=857	0.249 [0.552] N=743	0.570 [0.620] N=622	0.576 [0.753] N=491
WHO weight-for-age z score	-0.290 [0.196] N=1,085	-0.304 [0.231] N=994	-0.241 [0.251] N=864	-0.264 [0.276] N=717	-0.607 [0.374] N=562
WHO weight-for-length/height z score, 0-60 months	0.279 [0.345] N=928	0.270 [0.374] N=850	0.224 [0.394] N=740	0.075 [0.468] N=619	-0.045 [0.574] N=483
WHO BMI-for-age z score	0.324 [0.373] N=916	0.286 [0.392] N=839	0.227 [0.427] N=729	-0.047 [0.482] N=607	-0.313 [0.565] N=475
Stunted: HAZ below -2	-0.089 [0.071] N=934	-0.096 [0.078] N=857	-0.105 [0.089] N=743	-0.162 [0.104] N=622	-0.175 [0.123] N=491
Underweight: WAZ below -2	**0.106** [0.044] N=1,085**	**0.088* [0.048] N=994**	0.061 [0.054] N=864	0.098 [0.062] N=717	0.131 [0.083] N=562
Wasted: WLZ/WHZ below -2, 0-60 months	**-0.108** [0.054] N=928**	-0.086 [0.057] N=850	**-0.108* [0.062] N=740**	-0.103 [0.075] N=619	**-0.166* [0.089] N=483**
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.062] N=928	0.004 [0.068] N=850	-0.022 [0.068] N=740	-0.049 [0.090] N=619	-0.170 [0.106] N=483
Obese: WLZ/WHZ above +3, 0-60 months	-0.078 [0.048] N=928	-0.074 [0.055] N=850	-0.049 [0.057] N=740	-0.035 [0.071] N=619	-0.068 [0.093] N=483
CIAF: any stunting, underweight, or wasting, 0-60 months	**-0.130* [0.072] N=890**	-0.122 [0.077] N=815	**-0.145* [0.086] N=708**	**-0.221** [0.101] N=595**	**-0.232* [0.125] N=467**
BMI-for-age z score below -2	**-0.127** [0.059] N=916**	**-0.112* [0.060] N=839**	**-0.126* [0.069] N=729**	-0.090 [0.081] N=607	-0.115 [0.100] N=475

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.674	-0.087 [0.326] N=1,040	-0.163 [0.336] N=933	-0.018 [0.361] N=815	-0.044 [0.392] N=689	-0.133 [0.449] N=560
WHO weight-for-age z score	-0.294	-0.129 [0.161] N=1,203	-0.107 [0.169] N=1,085	-0.171 [0.177] N=947	-0.103 [0.189] N=799	-0.081 [0.215] N=647
WHO weight-for-length/height z score, 0-60 months	0.340	-0.083 [0.302] N=1,040	-0.122 [0.312] N=937	-0.204 [0.338] N=818	-0.015 [0.373] N=691	-0.102 [0.428] N=560
WHO BMI-for-age z score	0.294	0.027 [0.288] N=1,025	0.060 [0.293] N=924	0.024 [0.326] N=805	0.231 [0.359] N=680	0.165 [0.395] N=554
Stunted: HAZ below -2	0.291 P=300	0.050 [0.069] N=1,040	0.053 [0.070] N=933	0.029 [0.075] N=815	0.033 [0.083] N=689	0.008 [0.092] N=560
Underweight: WAZ below -2	0.118 P=137	0.018 [0.037] N=1,203	0.022 [0.039] N=1,085	0.046 [0.041] N=947	0.037 [0.044] N=799	0.023 [0.047] N=647
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=123	0.013 [0.047] N=1,040	0.031 [0.049] N=937	0.052 [0.054] N=818	0.001 [0.054] N=691	0.002 [0.063] N=560
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=207	0.018 [0.055] N=1,040	0.026 [0.057] N=937	-0.000 [0.060] N=818	0.013 [0.066] N=691	0.007 [0.075] N=560
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=99	0.014 [0.042] N=1,040	0.008 [0.042] N=937	0.011 [0.046] N=818	-0.003 [0.051] N=691	-0.045 [0.052] N=560
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=390	0.084 [0.060] N=994	0.096 [0.061] N=892	0.079 [0.065] N=778	0.035 [0.070] N=657	0.009 [0.076] N=532
BMI-for-age z score below -2	0.142 P=143	-0.003 [0.049] N=1,025	0.008 [0.051] N=924	-0.003 [0.057] N=805	-0.050 [0.059] N=680	-0.038 [0.067] N=554

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256 [0.496] N=906	-0.406 [0.523] N=799	-0.168 [0.603] N=681	-0.322 [0.713] N=555	-0.595 [0.901] N=426
WHO weight-for-age z score	-0.108 [0.243] N=1,048	-0.100 [0.261] N=930	-0.236 [0.282] N=792	-0.173 [0.308] N=644	-0.082 [0.356] N=492
WHO weight-for-length/height z score, 0-60 months	-0.205 [0.382] N=905	-0.297 [0.402] N=802	-0.444 [0.456] N=683	-0.113 [0.524] N=556	-0.405 [0.637] N=425
WHO BMI-for-age z score	-0.230 [0.383] N=896	-0.208 [0.397] N=795	-0.267 [0.475] N=676	0.092 [0.549] N=551	-0.146 [0.576] N=425
Stunted: HAZ below -2	0.081 [0.101] N=906	0.095 [0.104] N=799	0.051 [0.116] N=681	0.080 [0.140] N=555	0.033 [0.165] N=426
Underweight: WAZ below -2	0.002 [0.059] N=1,048	0.012 [0.065] N=930	0.057 [0.072] N=792	0.052 [0.081] N=644	0.014 [0.092] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.053 [0.064] N=905	0.086 [0.066] N=802	**0.127* [0.075] N=683**	0.040 [0.078] N=556	0.069 [0.104] N=425
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.073] N=905	0.021 [0.079] N=802	-0.021 [0.084] N=683	0.017 [0.100] N=556	-0.013 [0.125] N=425
Obese: WLZ/WHZ above +3, 0-60 months	0.046 [0.055] N=905	0.032 [0.058] N=802	0.043 [0.066] N=683	0.018 [0.083] N=556	-0.095 [0.094] N=425
CIAF: any stunting, underweight, or wasting, 0-60 months	0.135 [0.098] N=865	0.167 [0.102] N=763	0.139 [0.116] N=649	0.072 [0.137] N=528	0.013 [0.161] N=403
BMI-for-age z score below -2	0.016 [0.064] N=896	0.035 [0.068] N=795	0.003 [0.085] N=676	-0.104 [0.083] N=551	-0.098 [0.104] N=425

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.674	-0.087 [0.325] N=1,040	-0.167 [0.336] N=933	-0.030 [0.360] N=815	-0.051 [0.393] N=689	-0.143 [0.451] N=560
WHO weight-for-age z score	-0.294	-0.123 [0.160] N=1,203	-0.101 [0.167] N=1,085	-0.168 [0.176] N=947	-0.099 [0.189] N=799	-0.078 [0.214] N=647
WHO weight-for-length/height z score, 0-60 months	0.340	-0.078 [0.301] N=1,040	-0.119 [0.311] N=937	-0.201 [0.338] N=818	-0.011 [0.372] N=691	-0.090 [0.425] N=560
WHO BMI-for-age z score	0.294	0.026 [0.288] N=1,025	0.061 [0.294] N=924	0.021 [0.327] N=805	0.229 [0.361] N=680	0.170 [0.397] N=554
Stunted: HAZ below -2	0.291 P=300	0.047 [0.069] N=1,040	0.051 [0.070] N=933	0.028 [0.075] N=815	0.032 [0.083] N=689	0.008 [0.092] N=560
Underweight: WAZ below -2	0.118 P=137	0.017 [0.037] N=1,203	0.021 [0.039] N=1,085	0.045 [0.040] N=947	0.037 [0.044] N=799	0.023 [0.047] N=647
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=123	0.010 [0.047] N=1,040	0.027 [0.049] N=937	0.048 [0.054] N=818	-0.002 [0.053] N=691	-0.003 [0.063] N=560
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=207	0.016 [0.055] N=1,040	0.023 [0.058] N=937	-0.005 [0.060] N=818	0.009 [0.067] N=691	0.004 [0.076] N=560
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=99	0.013 [0.042] N=1,040	0.006 [0.042] N=937	0.009 [0.045] N=818	-0.006 [0.051] N=691	-0.048 [0.052] N=560
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=390	0.078 [0.060] N=994	0.089 [0.062] N=892	0.072 [0.065] N=778	0.031 [0.070] N=657	0.003 [0.078] N=532
BMI-for-age z score below -2	0.142 P=143	-0.004 [0.049] N=1,025	0.006 [0.051] N=924	-0.004 [0.057] N=805	-0.050 [0.059] N=680	-0.039 [0.067] N=554

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.257 [0.495] N=906	-0.422 [0.522] N=799	-0.214 [0.598] N=681	-0.374 [0.710] N=555	-0.659 [0.899] N=426
WHO weight-for-age z score	-0.099 [0.242] N=1,048	-0.089 [0.259] N=930	-0.229 [0.280] N=792	-0.157 [0.307] N=644	-0.059 [0.357] N=492
WHO weight-for-length/height z score, 0-60 months	-0.198 [0.384] N=905	-0.293 [0.405] N=802	-0.442 [0.460] N=683	-0.098 [0.532] N=556	-0.351 [0.655] N=425
WHO BMI-for-age z score	-0.226 [0.386] N=896	-0.197 [0.401] N=795	-0.259 [0.480] N=676	0.113 [0.558] N=551	-0.072 [0.601] N=425
Stunted: HAZ below -2	0.077 [0.101] N=906	0.093 [0.104] N=799	0.053 [0.115] N=681	0.085 [0.140] N=555	0.041 [0.165] N=426
Underweight: WAZ below -2	0.001 [0.059] N=1,048	0.010 [0.066] N=930	0.056 [0.072] N=792	0.052 [0.082] N=644	0.013 [0.093] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.047 [0.064] N=905	0.079 [0.068] N=802	0.118 [0.078] N=683	0.027 [0.079] N=556	0.047 [0.107] N=425
Overweight: WLZ/WHZ above +2, 0-60 months	0.014 [0.073] N=905	0.016 [0.078] N=802	-0.030 [0.083] N=683	0.004 [0.100] N=556	-0.023 [0.125] N=425
Obese: WLZ/WHZ above +3, 0-60 months	0.044 [0.055] N=905	0.028 [0.058] N=802	0.037 [0.066] N=683	0.011 [0.083] N=556	-0.106 [0.094] N=425
CIAF: any stunting, underweight, or wasting, 0-60 months	0.124 [0.097] N=865	0.153 [0.101] N=763	0.123 [0.114] N=649	0.055 [0.136] N=528	-0.011 [0.158] N=403
BMI-for-age z score below -2	0.012 [0.065] N=896	0.029 [0.069] N=795	-0.002 [0.086] N=676	-0.110 [0.085] N=551	-0.110 [0.108] N=425

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.674	-0.082 [0.321] N=1,040	-0.150 [0.332] N=933	0.009 [0.358] N=815	-0.021 [0.391] N=689	-0.119 [0.446] N=560
WHO weight-for-age z score	-0.294	-0.122 [0.154] N=1,203	-0.097 [0.162] N=1,085	-0.148 [0.170] N=947	-0.087 [0.183] N=799	-0.073 [0.206] N=647
WHO weight-for-length/height z score, 0-60 months	0.340	-0.076 [0.302] N=1,040	-0.122 [0.311] N=937	-0.190 [0.336] N=818	-0.006 [0.370] N=691	-0.090 [0.426] N=560
WHO BMI-for-age z score	0.294	0.026 [0.288] N=1,025	0.052 [0.293] N=924	0.029 [0.322] N=805	0.239 [0.353] N=680	0.165 [0.393] N=554
Stunted: HAZ below -2	0.291 P=300	0.045 [0.068] N=1,040	0.045 [0.069] N=933	0.016 [0.073] N=815	0.019 [0.081] N=689	-0.002 [0.089] N=560
Underweight: WAZ below -2	0.118 P=137	0.018 [0.035] N=1,203	0.022 [0.037] N=1,085	0.042 [0.039] N=947	0.035 [0.041] N=799	0.022 [0.042] N=647
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=123	0.010 [0.048] N=1,040	0.027 [0.049] N=937	0.046 [0.054] N=818	-0.004 [0.054] N=691	-0.003 [0.063] N=560
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=207	0.015 [0.055] N=1,040	0.023 [0.058] N=937	-0.009 [0.061] N=818	0.002 [0.068] N=691	0.002 [0.076] N=560
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=99	0.013 [0.043] N=1,040	0.006 [0.043] N=937	0.006 [0.046] N=818	-0.009 [0.052] N=691	-0.051 [0.053] N=560
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=390	0.079 [0.060] N=994	0.086 [0.062] N=892	0.060 [0.066] N=778	0.018 [0.071] N=657	-0.009 [0.075] N=532
BMI-for-age z score below -2	0.142 P=143	-0.005 [0.049] N=1,025	0.007 [0.051] N=924	-0.009 [0.057] N=805	-0.055 [0.059] N=680	-0.042 [0.068] N=554

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.303 [0.494] N=906	-0.436 [0.516] N=799	-0.207 [0.596] N=681	-0.392 [0.704] N=555	-0.727 [0.919] N=426
WHO weight-for-age z score	-0.141 [0.237] N=1,048	-0.129 [0.253] N=930	-0.229 [0.274] N=792	-0.187 [0.303] N=644	-0.153 [0.351] N=492
WHO weight-for-length/height z score, 0-60 months	-0.215 [0.385] N=905	-0.315 [0.406] N=802	-0.439 [0.456] N=683	-0.101 [0.534] N=556	-0.360 [0.682] N=425
WHO BMI-for-age z score	-0.237 [0.387] N=896	-0.225 [0.403] N=795	-0.250 [0.474] N=676	0.115 [0.556] N=551	-0.076 [0.616] N=425
Stunted: HAZ below -2	0.086 [0.099] N=906	0.094 [0.102] N=799	0.052 [0.113] N=681	0.092 [0.138] N=555	0.061 [0.165] N=426
Underweight: WAZ below -2	0.008 [0.057] N=1,048	0.020 [0.063] N=930	0.056 [0.070] N=792	0.057 [0.080] N=644	0.029 [0.088] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.052 [0.066] N=905	0.087 [0.068] N=802	0.118 [0.078] N=683	0.028 [0.080] N=556	0.051 [0.109] N=425
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.074] N=905	0.021 [0.080] N=802	-0.033 [0.085] N=683	0.005 [0.102] N=556	-0.020 [0.130] N=425
Obese: WLZ/WHZ above +3, 0-60 months	0.045 [0.056] N=905	0.031 [0.058] N=802	0.036 [0.067] N=683	0.011 [0.083] N=556	-0.106 [0.095] N=425
CIAF: any stunting, underweight, or wasting, 0-60 months	0.139 [0.097] N=865	0.167 [0.102] N=763	0.123 [0.116] N=649	0.065 [0.138] N=528	0.012 [0.163] N=403
BMI-for-age z score below -2	0.016 [0.065] N=896	0.039 [0.069] N=795	-0.003 [0.084] N=676	-0.109 [0.085] N=551	-0.105 [0.110] N=425

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.674	-0.088 [0.315] N=1,040	-0.148 [0.326] N=933	0.014 [0.348] N=815	-0.024 [0.379] N=689	-0.176 [0.437] N=560
WHO weight-for-age z score	-0.294	-0.122 [0.151] N=1,203	-0.094 [0.159] N=1,085	-0.138 [0.169] N=947	-0.074 [0.178] N=799	-0.066 [0.202] N=647
WHO weight-for-length/height z score, 0-60 months	0.340	-0.064 [0.299] N=1,040	-0.110 [0.309] N=937	-0.187 [0.332] N=818	0.024 [0.365] N=691	-0.018 [0.420] N=560
WHO BMI-for-age z score	0.294	0.039 [0.289] N=1,025	0.071 [0.296] N=924	0.041 [0.323] N=805	0.271 [0.355] N=680	0.245 [0.396] N=554
Stunted: HAZ below -2	0.291 P=300	0.046 [0.068] N=1,040	0.046 [0.069] N=933	0.019 [0.073] N=815	0.026 [0.080] N=689	0.009 [0.089] N=560
Underweight: WAZ below -2	0.118 P=137	0.018 [0.035] N=1,203	0.022 [0.037] N=1,085	0.042 [0.039] N=947	0.036 [0.042] N=799	0.025 [0.044] N=647
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=123	0.008 [0.048] N=1,040	0.027 [0.049] N=937	0.049 [0.054] N=818	-0.004 [0.052] N=691	-0.008 [0.061] N=560
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=207	0.015 [0.054] N=1,040	0.021 [0.057] N=937	-0.006 [0.060] N=818	0.009 [0.067] N=691	0.010 [0.075] N=560
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=99	0.017 [0.041] N=1,040	0.010 [0.042] N=937	0.011 [0.044] N=818	-0.001 [0.050] N=691	-0.040 [0.051] N=560
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=390	0.075 [0.060] N=994	0.084 [0.062] N=892	0.063 [0.065] N=778	0.024 [0.070] N=657	0.001 [0.075] N=532
BMI-for-age z score below -2	0.142 P=143	-0.008 [0.049] N=1,025	0.002 [0.051] N=924	-0.010 [0.057] N=805	-0.056 [0.058] N=680	-0.049 [0.065] N=554

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.283 [0.476] N=906	-0.412 [0.500] N=799	-0.110 [0.571] N=681	-0.272 [0.678] N=555	-0.664 [0.871] N=426
WHO weight-for-age z score	-0.113 [0.231] N=1,048	-0.100 [0.247] N=930	-0.182 [0.276] N=792	-0.128 [0.297] N=644	-0.076 [0.350] N=492
WHO weight-for-length/height z score, 0-60 months	-0.204 [0.384] N=905	-0.313 [0.405] N=802	-0.464 [0.452] N=683	-0.119 [0.523] N=556	-0.358 [0.657] N=425
WHO BMI-for-age z score	-0.221 [0.387] N=896	-0.201 [0.406] N=795	-0.255 [0.480] N=676	0.126 [0.554] N=551	-0.057 [0.603] N=425
Stunted: HAZ below -2	0.086 [0.097] N=906	0.093 [0.100] N=799	0.042 [0.112] N=681	0.082 [0.137] N=555	0.059 [0.163] N=426
Underweight: WAZ below -2	0.003 [0.056] N=1,048	0.014 [0.062] N=930	0.048 [0.071] N=792	0.049 [0.079] N=644	0.016 [0.090] N=492
Wasted: WLZ/WHZ below -2, 0-60 months	0.051 [0.065] N=905	0.084 [0.068] N=802	0.123 [0.076] N=683	0.035 [0.079] N=556	0.054 [0.105] N=425
Overweight: WLZ/WHZ above +2, 0-60 months	0.012 [0.073] N=905	0.013 [0.079] N=802	-0.040 [0.083] N=683	-0.005 [0.099] N=556	-0.022 [0.126] N=425
Obese: WLZ/WHZ above +3, 0-60 months	0.046 [0.055] N=905	0.029 [0.057] N=802	0.035 [0.065] N=683	0.013 [0.082] N=556	-0.099 [0.093] N=425
CIAF: any stunting, underweight, or wasting, 0-60 months	0.134 [0.097] N=865	0.158 [0.102] N=763	0.114 [0.116] N=649	0.056 [0.139] N=528	0.014 [0.162] N=403
BMI-for-age z score below -2	0.014 [0.066] N=896	0.032 [0.070] N=795	-0.004 [0.086] N=676	-0.108 [0.086] N=551	-0.108 [0.107] N=425

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.676	-0.109 [0.348] N=954	-0.193 [0.365] N=856	-0.019 [0.384] N=753	-0.054 [0.406] N=634	-0.162 [0.467] N=518
WHO weight-for-age z score	-0.294	-0.099 [0.171] N=1,097	-0.063 [0.180] N=988	-0.160 [0.188] N=869	-0.127 [0.201] N=734	-0.153 [0.223] N=599
WHO weight-for-length/height z score, 0-60 months	0.347	0.035 [0.318] N=954	0.004 [0.332] N=860	-0.136 [0.354] N=754	0.014 [0.385] N=636	-0.117 [0.432] N=519
WHO BMI-for-age z score	0.303	0.093 [0.306] N=941	0.152 [0.317] N=849	0.042 [0.350] N=744	0.225 [0.389] N=628	0.112 [0.428] N=515
Stunted: HAZ below -2	0.291 P=274	0.045 [0.074] N=954	0.051 [0.077] N=856	0.013 [0.080] N=753	0.010 [0.086] N=634	-0.008 [0.095] N=518
Underweight: WAZ below -2	0.120 P=128	0.016 [0.040] N=1,097	0.020 [0.043] N=988	0.043 [0.044] N=869	0.043 [0.047] N=734	0.035 [0.051] N=599
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=112	0.007 [0.049] N=954	0.025 [0.051] N=860	0.046 [0.056] N=754	-0.003 [0.055] N=636	0.007 [0.064] N=519
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=192	0.026 [0.057] N=954	0.035 [0.060] N=860	-0.005 [0.062] N=754	-0.008 [0.068] N=636	-0.022 [0.075] N=519
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=93	0.016 [0.044] N=954	0.008 [0.044] N=860	0.008 [0.046] N=754	-0.014 [0.050] N=636	-0.054 [0.051] N=519
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=357	0.077 [0.066] N=912	0.086 [0.068] N=819	0.054 [0.071] N=718	0.000 [0.074] N=604	-0.006 [0.082] N=492
BMI-for-age z score below -2	0.140 P=128	0.000 [0.051] N=941	0.011 [0.053] N=849	-0.000 [0.059] N=744	-0.040 [0.062] N=628	-0.016 [0.071] N=515

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.302 [0.540] N=829	-0.479 [0.572] N=731	-0.254 [0.648] N=628	-0.512 [0.742] N=509	-0.770 [0.956] N=393
WHO weight-for-age z score	-0.018 [0.246] N=954	0.025 [0.264] N=845	-0.172 [0.289] N=726	-0.125 [0.324] N=591	-0.035 [0.379] N=456
WHO weight-for-length/height z score, 0-60 months	-0.062 [0.412] N=829	-0.117 [0.443] N=735	-0.353 [0.501] N=629	-0.043 [0.586] N=511	-0.385 [0.739] N=394
WHO BMI-for-age z score	-0.127 [0.389] N=820	-0.035 [0.410] N=728	-0.191 [0.492] N=623	0.193 [0.563] N=507	-0.033 [0.638] N=394
Stunted: HAZ below -2	0.085 [0.111] N=829	0.103 [0.118] N=731	0.042 [0.127] N=628	0.057 [0.151] N=509	-0.001 [0.178] N=393
Underweight: WAZ below -2	-0.012 [0.063] N=954	-0.004 [0.070] N=845	0.037 [0.077] N=726	0.045 [0.090] N=591	-0.004 [0.101] N=456
Wasted: WLZ/WHZ below -2, 0-60 months	0.039 [0.067] N=829	0.067 [0.072] N=735	0.101 [0.084] N=629	0.003 [0.085] N=511	0.036 [0.113] N=394
Overweight: WLZ/WHZ above +2, 0-60 months	0.031 [0.079] N=829	0.040 [0.086] N=735	-0.019 [0.092] N=629	-0.016 [0.111] N=511	-0.068 [0.142] N=394
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.061] N=829	0.025 [0.064] N=735	0.035 [0.073] N=629	-0.007 [0.091] N=511	-0.131 [0.110] N=394
CIAF: any stunting, underweight, or wasting, 0-60 months	0.132 [0.109] N=792	0.154 [0.116] N=699	0.095 [0.128] N=598	-0.019 [0.151] N=484	-0.077 [0.176] N=372
BMI-for-age z score below -2	0.020 [0.066] N=820	0.036 [0.072] N=728	-0.001 [0.090] N=623	-0.103 [0.090] N=507	-0.079 [0.117] N=394

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.676	-0.127 [0.351] N=954	-0.209 [0.368] N=856	-0.035 [0.386] N=753	-0.070 [0.407] N=634	-0.177 [0.470] N=518
WHO weight-for-age z score	-0.294	-0.108 [0.169] N=1,097	-0.077 [0.178] N=988	-0.167 [0.186] N=869	-0.124 [0.198] N=734	-0.143 [0.219] N=599
WHO weight-for-length/height z score, 0-60 months	0.347	0.035 [0.318] N=954	0.004 [0.332] N=860	-0.130 [0.354] N=754	0.019 [0.384] N=636	-0.099 [0.432] N=519
WHO BMI-for-age z score	0.303	0.104 [0.306] N=941	0.166 [0.316] N=849	0.055 [0.349] N=744	0.238 [0.385] N=628	0.127 [0.425] N=515
Stunted: HAZ below -2	0.291 P=274	0.048 [0.074] N=954	0.052 [0.077] N=856	0.014 [0.080] N=753	0.009 [0.086] N=634	-0.012 [0.095] N=518
Underweight: WAZ below -2	0.120 P=128	0.013 [0.039] N=1,097	0.018 [0.042] N=988	0.039 [0.043] N=869	0.038 [0.046] N=734	0.030 [0.049] N=599
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=112	0.004 [0.050] N=954	0.021 [0.052] N=860	0.041 [0.056] N=754	-0.009 [0.055] N=636	0.000 [0.064] N=519
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=192	0.024 [0.056] N=954	0.031 [0.059] N=860	-0.009 [0.061] N=754	-0.012 [0.067] N=636	-0.023 [0.074] N=519
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=93	0.012 [0.043] N=954	0.004 [0.043] N=860	0.005 [0.045] N=754	-0.015 [0.048] N=636	-0.053 [0.049] N=519
CIAF: any stunting, underweight, or wasting, 0-60 months	0.397 P=357	0.077 [0.065] N=912	0.084 [0.068] N=819	0.051 [0.070] N=718	-0.008 [0.074] N=604	-0.019 [0.082] N=492
BMI-for-age z score below -2	0.140 P=128	-0.001 [0.051] N=941	0.008 [0.054] N=849	-0.003 [0.060] N=744	-0.044 [0.062] N=628	-0.023 [0.070] N=515

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.302 [0.536] N=829	-0.468 [0.569] N=731	-0.226 [0.643] N=628	-0.481 [0.731] N=509	-0.773 [0.934] N=393
WHO weight-for-age z score	-0.045 [0.246] N=954	-0.006 [0.267] N=845	-0.189 [0.289] N=726	-0.135 [0.319] N=591	-0.041 [0.370] N=456
WHO weight-for-length/height z score, 0-60 months	-0.105 [0.410] N=829	-0.160 [0.441] N=735	-0.396 [0.498] N=629	-0.091 [0.569] N=511	-0.340 [0.715] N=394
WHO BMI-for-age z score	-0.130 [0.388] N=820	-0.036 [0.410] N=728	-0.196 [0.491] N=623	0.196 [0.560] N=507	0.006 [0.626] N=394
Stunted: HAZ below -2	0.088 [0.111] N=829	0.104 [0.117] N=731	0.042 [0.126] N=628	0.056 [0.149] N=509	-0.001 [0.176] N=393
Underweight: WAZ below -2	-0.013 [0.063] N=954	-0.005 [0.070] N=845	0.034 [0.077] N=726	0.043 [0.089] N=591	0.004 [0.101] N=456
Wasted: WLZ/WHZ below -2, 0-60 months	0.044 [0.067] N=829	0.072 [0.073] N=735	0.107 [0.085] N=629	0.009 [0.084] N=511	0.041 [0.112] N=394
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.078] N=829	0.033 [0.085] N=735	-0.030 [0.090] N=629	-0.025 [0.108] N=511	-0.056 [0.137] N=394
Obese: WLZ/WHZ above +3, 0-60 months	0.032 [0.060] N=829	0.017 [0.063] N=735	0.027 [0.071] N=629	-0.012 [0.088] N=511	-0.120 [0.101] N=394
CIAF: any stunting, underweight, or wasting, 0-60 months	0.140 [0.109] N=792	0.161 [0.116] N=699	0.101 [0.129] N=598	-0.017 [0.153] N=484	-0.077 [0.180] N=372
BMI-for-age z score below -2	0.023 [0.067] N=820	0.038 [0.073] N=728	0.002 [0.092] N=623	-0.099 [0.091] N=507	-0.081 [0.118] N=394

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 35 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.