Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Oldest Child

Age reference: 10 October 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.149 [0.214] N=1,738	0.230 [0.228] N=1,613	0.237 [0.251] N=1,448	0.225 [0.257] N=1,236	-0.048 [0.275] N=1,025
WHO weight-for-age z score	0.306	-0.118 [0.118] N=1,966	-0.054 [0.126] N=1,818	-0.057 [0.132] N=1,633	-0.061 [0.138] N=1,390	-0.165 [0.152] N=1,147
WHO weight-for-length/height z score, 0-60 months	0.329	0.177 [0.191] N=1,715	0.147 [0.202] N=1,596	0.110 [0.226] N=1,434	0.095 [0.244] N=1,224	0.192 [0.270] N=1,011
WHO BMI-for-age z score	0.298	0.147 [0.209] N=1,685	0.162 [0.223] N=1,570	0.120 [0.251] N=1,407	0.019 [0.277] N=1,199	0.174 [0.305] N=992
Stunted: HAZ below -2	0.191 P=319	-0.025 [0.037] N=1,738	-0.038 [0.039] N=1,613	-0.021 [0.041] N=1,448	-0.006 [0.042] N=1,236	0.014 [0.047] N=1,025
Underweight: WAZ below -2	0.052 P=107	0.016 [0.017] N=1,966	0.004 [0.018] N=1,818	0.002 [0.019] N=1,633	-0.002 [0.020] N=1,390	0.004 [0.026] N=1,147
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=208	-0.036 [0.032] N=1,715	-0.030 [0.032] N=1,596	-0.034 [0.035] N=1,434	-0.022 [0.038] N=1,224	-0.042 [0.039] N=1,011
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=327	0.015 [0.038] N=1,715	0.003 [0.041] N=1,596	-0.007 [0.045] N=1,434	0.000 [0.051] N=1,224	0.006 [0.057] N=1,011
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=156	0.025 [0.029] N=1,715	0.024 [0.031] N=1,596	0.025 [0.034] N=1,434	0.031 [0.038] N=1,224	0.041 [0.042] N=1,011
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=441	-0.045 [0.039] N=1,654	-0.050 [0.039] N=1,540	-0.041 [0.042] N=1,383	-0.028 [0.044] N=1,179	-0.023 [0.048] N=976
BMI-for-age z score below -2	0.127 P=218	-0.015 [0.035] N=1,685	-0.011 [0.036] N=1,570	-0.005 [0.040] N=1,407	0.034 [0.043] N=1,199	0.022 [0.045] N=992

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.227 [0.305] N=1,462	0.376 [0.344] N=1,337	0.502 [0.396] N=1,172	0.679 [0.429] N=960	0.143 [0.582] N=749
WHO weight-for-age z score	-0.272 [0.177] N=1,667	-0.181 [0.200] N=1,519	-0.215 [0.216] N=1,334	-0.270 [0.265] N=1,091	**-0.739** [0.360] N=848**
WHO weight-for-length/height z score, 0-60 months	0.092 [0.244] N=1,442	0.041 [0.267] N=1,323	-0.120 [0.315] N=1,161	-0.226 [0.360] N=951	-0.174 [0.444] N=738
WHO BMI-for-age z score	0.119 [0.240] N=1,419	0.156 [0.266] N=1,304	-0.022 [0.322] N=1,141	-0.334 [0.366] N=933	-0.152 [0.429] N=726
Stunted: HAZ below -2	-0.002 [0.053] N=1,462	-0.020 [0.058] N=1,337	0.008 [0.063] N=1,172	0.041 [0.072] N=960	0.122 [0.102] N=749
Underweight: WAZ below -2	0.035 [0.023] N=1,667	0.019 [0.025] N=1,519	0.017 [0.028] N=1,334	0.015 [0.032] N=1,091	0.056 [0.048] N=848
Wasted: WLZ/WHZ below -2, 0-60 months	-0.037 [0.045] N=1,442	-0.029 [0.047] N=1,323	-0.035 [0.054] N=1,161	-0.017 [0.067] N=951	-0.074 [0.079] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.005 [0.045] N=1,442	-0.017 [0.050] N=1,323	-0.049 [0.054] N=1,161	-0.051 [0.070] N=951	-0.079 [0.083] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.018 [0.032] N=1,442	0.017 [0.036] N=1,323	0.018 [0.041] N=1,161	0.031 [0.051] N=951	0.048 [0.062] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.008 [0.058] N=1,390	-0.013 [0.061] N=1,276	0.013 [0.069] N=1,119	0.045 [0.083] N=915	0.097 [0.115] N=712
BMI-for-age z score below -2	-0.044 [0.044] N=1,419	-0.043 [0.046] N=1,304	-0.027 [0.055] N=1,141	0.041 [0.066] N=933	0.014 [0.076] N=726

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.150 [0.215] N=1,738	0.230 [0.229] N=1,613	0.239 [0.252] N=1,448	0.235 [0.258] N=1,236	-0.034 [0.277] N=1,025
WHO weight-for-age z score	0.306	-0.121 [0.117] N=1,966	-0.057 [0.126] N=1,818	-0.060 [0.132] N=1,633	-0.063 [0.138] N=1,390	-0.170 [0.151] N=1,147
WHO weight-for-length/height z score, 0-60 months	0.329	0.173 [0.192] N=1,715	0.141 [0.202] N=1,596	0.103 [0.227] N=1,434	0.083 [0.246] N=1,224	0.172 [0.272] N=1,011
WHO BMI-for-age z score	0.298	0.143 [0.210] N=1,685	0.156 [0.224] N=1,570	0.113 [0.253] N=1,407	0.007 [0.280] N=1,199	0.155 [0.307] N=992
Stunted: HAZ below -2	0.191 P=319	-0.025 [0.037] N=1,738	-0.038 [0.039] N=1,613	-0.021 [0.041] N=1,448	-0.007 [0.042] N=1,236	0.013 [0.048] N=1,025
Underweight: WAZ below -2	0.052 P=107	0.016 [0.017] N=1,966	0.005 [0.018] N=1,818	0.002 [0.019] N=1,633	-0.002 [0.020] N=1,390	0.004 [0.026] N=1,147
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=208	-0.034 [0.032] N=1,715	-0.028 [0.032] N=1,596	-0.031 [0.035] N=1,434	-0.018 [0.038] N=1,224	-0.036 [0.040] N=1,011
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=327	0.016 [0.038] N=1,715	0.003 [0.041] N=1,596	-0.006 [0.045] N=1,434	0.001 [0.051] N=1,224	0.006 [0.057] N=1,011
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=156	0.025 [0.029] N=1,715	0.024 [0.031] N=1,596	0.026 [0.034] N=1,434	0.031 [0.038] N=1,224	0.041 [0.042] N=1,011
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=441	-0.042 [0.039] N=1,654	-0.046 [0.039] N=1,540	-0.036 [0.042] N=1,383	-0.023 [0.044] N=1,179	-0.016 [0.048] N=976
BMI-for-age z score below -2	0.127 P=218	-0.014 [0.035] N=1,685	-0.009 [0.036] N=1,570	-0.002 [0.041] N=1,407	0.038 [0.043] N=1,199	0.028 [0.046] N=992

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.224 [0.303] N=1,462	0.370 [0.342] N=1,337	0.495 [0.393] N=1,172	0.668 [0.420] N=960	0.125 [0.570] N=749
WHO weight-for-age z score	-0.268 [0.177] N=1,667	-0.175 [0.200] N=1,519	-0.211 [0.216] N=1,334	-0.268 [0.266] N=1,091	**-0.729** [0.361] N=848**
WHO weight-for-length/height z score, 0-60 months	0.092 [0.242] N=1,442	0.041 [0.264] N=1,323	-0.120 [0.312] N=1,161	-0.232 [0.357] N=951	-0.184 [0.440] N=738
WHO BMI-for-age z score	0.122 [0.239] N=1,419	0.158 [0.265] N=1,304	-0.021 [0.320] N=1,141	-0.340 [0.366] N=933	-0.164 [0.430] N=726
Stunted: HAZ below -2	-0.001 [0.053] N=1,462	-0.020 [0.058] N=1,337	0.009 [0.063] N=1,172	0.042 [0.071] N=960	0.124 [0.102] N=749
Underweight: WAZ below -2	0.035 [0.023] N=1,667	0.019 [0.025] N=1,519	0.017 [0.028] N=1,334	0.016 [0.032] N=1,091	0.056 [0.048] N=848
Wasted: WLZ/WHZ below -2, 0-60 months	-0.037 [0.045] N=1,442	-0.029 [0.047] N=1,323	-0.034 [0.054] N=1,161	-0.015 [0.067] N=951	-0.072 [0.079] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.005 [0.045] N=1,442	-0.017 [0.050] N=1,323	-0.049 [0.055] N=1,161	-0.051 [0.070] N=951	-0.079 [0.083] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.018 [0.033] N=1,442	0.017 [0.036] N=1,323	0.018 [0.042] N=1,161	0.031 [0.051] N=951	0.048 [0.062] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.009 [0.058] N=1,390	-0.015 [0.060] N=1,276	0.011 [0.068] N=1,119	0.044 [0.082] N=915	0.098 [0.115] N=712
BMI-for-age z score below -2	-0.044 [0.044] N=1,419	-0.043 [0.046] N=1,304	-0.028 [0.056] N=1,141	0.042 [0.067] N=933	0.016 [0.077] N=726

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.118 [0.215] N=1,738	0.185 [0.229] N=1,613	0.199 [0.251] N=1,448	0.212 [0.258] N=1,236	-0.080 [0.273] N=1,025
WHO weight-for-age z score	0.306	-0.160 [0.114] N=1,966	-0.095 [0.123] N=1,818	-0.095 [0.129] N=1,633	-0.086 [0.136] N=1,390	-0.204 [0.150] N=1,147
WHO weight-for-length/height z score, 0-60 months	0.329	0.157 [0.193] N=1,715	0.136 [0.205] N=1,596	0.103 [0.228] N=1,434	0.084 [0.245] N=1,224	0.160 [0.273] N=1,011
WHO BMI-for-age z score	0.298	0.139 [0.212] N=1,685	0.160 [0.226] N=1,570	0.120 [0.254] N=1,407	0.015 [0.279] N=1,199	0.161 [0.309] N=992
Stunted: HAZ below -2	0.191 P=319	-0.019 [0.037] N=1,738	-0.028 [0.039] N=1,613	-0.013 [0.041] N=1,448	-0.002 [0.043] N=1,236	0.022 [0.048] N=1,025
Underweight: WAZ below -2	0.052 P=107	0.020 [0.017] N=1,966	0.010 [0.017] N=1,818	0.007 [0.018] N=1,633	0.001 [0.020] N=1,390	0.010 [0.025] N=1,147
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=208	-0.028 [0.032] N=1,715	-0.020 [0.033] N=1,596	-0.025 [0.035] N=1,434	-0.013 [0.038] N=1,224	-0.031 [0.040] N=1,011
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=327	0.017 [0.039] N=1,715	0.006 [0.042] N=1,596	-0.003 [0.045] N=1,434	0.003 [0.052] N=1,224	0.008 [0.058] N=1,011
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=156	0.027 [0.029] N=1,715	0.028 [0.031] N=1,596	0.028 [0.034] N=1,434	0.032 [0.038] N=1,224	0.041 [0.042] N=1,011
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=441	-0.029 [0.040] N=1,654	-0.028 [0.040] N=1,540	-0.021 [0.042] N=1,383	-0.014 [0.044] N=1,179	0.000 [0.049] N=976
BMI-for-age z score below -2	0.127 P=218	-0.010 [0.035] N=1,685	-0.004 [0.037] N=1,570	0.002 [0.041] N=1,407	0.040 [0.044] N=1,199	0.029 [0.047] N=992

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.201 [0.305] N=1,462	0.321 [0.344] N=1,337	0.454 [0.395] N=1,172	0.675 [0.419] N=960	0.114 [0.564] N=749
WHO weight-for-age z score	**-0.296* [0.171] N=1,667**	-0.204 [0.195] N=1,519	-0.232 [0.210] N=1,334	-0.256 [0.264] N=1,091	**-0.733** [0.360] N=848**
WHO weight-for-length/height z score, 0-60 months	0.087 [0.245] N=1,442	0.053 [0.272] N=1,323	-0.099 [0.316] N=1,161	-0.225 [0.358] N=951	-0.212 [0.447] N=738
WHO BMI-for-age z score	0.123 [0.243] N=1,419	0.170 [0.273] N=1,304	-0.002 [0.324] N=1,141	-0.342 [0.367] N=933	-0.169 [0.438] N=726
Stunted: HAZ below -2	0.002 [0.053] N=1,462	-0.009 [0.058] N=1,337	0.015 [0.062] N=1,172	0.037 [0.070] N=960	0.117 [0.101] N=749
Underweight: WAZ below -2	0.037 [0.023] N=1,667	0.024 [0.025] N=1,519	0.020 [0.027] N=1,334	0.014 [0.031] N=1,091	0.055 [0.049] N=848
Wasted: WLZ/WHZ below -2, 0-60 months	-0.032 [0.045] N=1,442	-0.022 [0.048] N=1,323	-0.031 [0.055] N=1,161	-0.011 [0.067] N=951	-0.062 [0.080] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.008 [0.046] N=1,442	-0.013 [0.052] N=1,323	-0.047 [0.056] N=1,161	-0.050 [0.071] N=951	-0.075 [0.083] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.019 [0.033] N=1,442	0.021 [0.037] N=1,323	0.020 [0.043] N=1,161	0.030 [0.051] N=951	0.043 [0.062] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.002 [0.059] N=1,390	0.002 [0.061] N=1,276	0.019 [0.067] N=1,119	0.037 [0.081] N=915	0.099 [0.114] N=712
BMI-for-age z score below -2	-0.039 [0.044] N=1,419	-0.036 [0.047] N=1,304	-0.023 [0.056] N=1,141	0.048 [0.067] N=933	0.024 [0.079] N=726

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.064 [0.218] N=1,738	0.131 [0.234] N=1,613	0.150 [0.258] N=1,448	0.149 [0.266] N=1,236	-0.153 [0.285] N=1,025
WHO weight-for-age z score	0.306	-0.175 [0.115] N=1,966	-0.113 [0.124] N=1,818	-0.111 [0.132] N=1,633	-0.117 [0.140] N=1,390	-0.244 [0.156] N=1,147
WHO weight-for-length/height z score, 0-60 months	0.329	0.184 [0.194] N=1,715	0.161 [0.206] N=1,596	0.115 [0.230] N=1,434	0.086 [0.251] N=1,224	0.167 [0.278] N=1,011
WHO BMI-for-age z score	0.298	0.167 [0.212] N=1,685	0.190 [0.226] N=1,570	0.136 [0.255] N=1,407	0.023 [0.284] N=1,199	0.168 [0.313] N=992
Stunted: HAZ below -2	0.191 P=319	-0.008 [0.037] N=1,738	-0.016 [0.039] N=1,613	0.001 [0.041] N=1,448	0.014 [0.043] N=1,236	0.040 [0.048] N=1,025
Underweight: WAZ below -2	0.052 P=107	0.023 [0.017] N=1,966	0.013 [0.018] N=1,818	0.011 [0.018] N=1,633	0.006 [0.020] N=1,390	0.016 [0.025] N=1,147
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=208	-0.030 [0.032] N=1,715	-0.023 [0.033] N=1,596	-0.025 [0.035] N=1,434	-0.013 [0.038] N=1,224	-0.033 [0.040] N=1,011
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=327	0.022 [0.039] N=1,715	0.011 [0.042] N=1,596	-0.001 [0.045] N=1,434	0.005 [0.052] N=1,224	0.006 [0.058] N=1,011
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=156	0.029 [0.029] N=1,715	0.028 [0.031] N=1,596	0.029 [0.034] N=1,434	0.033 [0.038] N=1,224	0.042 [0.042] N=1,011
CIAF: any stunting, underweight, or wasting, 0-60 months	0.271 P=441	-0.019 [0.040] N=1,654	-0.018 [0.041] N=1,540	-0.009 [0.043] N=1,383	0.004 [0.046] N=1,179	0.020 [0.050] N=976
BMI-for-age z score below -2	0.127 P=218	-0.012 [0.035] N=1,685	-0.006 [0.037] N=1,570	0.002 [0.041] N=1,407	0.041 [0.044] N=1,199	0.027 [0.046] N=992

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.182 [0.307] N=1,462	0.310 [0.344] N=1,337	0.455 [0.396] N=1,172	0.682 [0.420] N=960	0.083 [0.561] N=749
WHO weight-for-age z score	**-0.310* [0.173] N=1,667**	-0.216 [0.196] N=1,519	-0.247 [0.211] N=1,334	-0.275 [0.264] N=1,091	**-0.728** [0.359] N=848**
WHO weight-for-length/height z score, 0-60 months	0.083 [0.244] N=1,442	0.050 [0.267] N=1,323	-0.116 [0.316] N=1,161	-0.232 [0.364] N=951	-0.156 [0.457] N=738
WHO BMI-for-age z score	0.113 [0.243] N=1,419	0.170 [0.268] N=1,304	-0.016 [0.326] N=1,141	-0.334 [0.376] N=933	-0.119 [0.446] N=726
Stunted: HAZ below -2	0.003 [0.053] N=1,462	-0.009 [0.058] N=1,337	0.012 [0.064] N=1,172	0.028 [0.072] N=960	0.112 [0.102] N=749
Underweight: WAZ below -2	**0.039* [0.023] N=1,667**	0.024 [0.025] N=1,519	0.021 [0.027] N=1,334	0.014 [0.032] N=1,091	0.056 [0.049] N=848
Wasted: WLZ/WHZ below -2, 0-60 months	-0.030 [0.045] N=1,442	-0.023 [0.047] N=1,323	-0.029 [0.054] N=1,161	-0.008 [0.068] N=951	-0.062 [0.081] N=738
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.046] N=1,442	-0.012 [0.051] N=1,323	-0.046 [0.056] N=1,161	-0.047 [0.070] N=951	-0.061 [0.082] N=738
Obese: WLZ/WHZ above +3, 0-60 months	0.022 [0.034] N=1,442	0.022 [0.037] N=1,323	0.021 [0.043] N=1,161	0.031 [0.052] N=951	0.052 [0.062] N=738
CIAF: any stunting, underweight, or wasting, 0-60 months	0.006 [0.060] N=1,390	0.003 [0.063] N=1,276	0.022 [0.070] N=1,119	0.039 [0.085] N=915	0.110 [0.116] N=712
BMI-for-age z score below -2	-0.037 [0.044] N=1,419	-0.038 [0.047] N=1,304	-0.024 [0.056] N=1,141	0.047 [0.068] N=933	0.021 [0.077] N=726

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.011	0.124 [0.213] N=1,611	0.168 [0.229] N=1,500	0.175 [0.250] N=1,342	0.126 [0.258] N=1,153	-0.182 [0.271] N=955
WHO weight-for-age z score	0.307	-0.061 [0.118] N=1,819	0.001 [0.124] N=1,687	0.001 [0.132] N=1,511	0.002 [0.136] N=1,293	-0.075 [0.157] N=1,069
WHO weight-for-length/height z score, 0-60 months	0.343	0.298 [0.192] N=1,591	0.270 [0.201] N=1,487	0.220 [0.226] N=1,333	0.223 [0.240] N=1,146	0.361 [0.268] N=946
WHO BMI-for-age z score	0.313	0.273 [0.210] N=1,561	0.271 [0.221] N=1,460	0.233 [0.249] N=1,306	0.169 [0.271] N=1,121	0.368 [0.300] N=927
Stunted: HAZ below -2	0.193 P=297	-0.015 [0.040] N=1,611	-0.025 [0.042] N=1,500	-0.007 [0.043] N=1,342	0.012 [0.045] N=1,153	0.044 [0.051] N=955
Underweight: WAZ below -2	0.052 P=99	0.010 [0.018] N=1,819	-0.001 [0.019] N=1,687	-0.003 [0.020] N=1,511	-0.006 [0.022] N=1,293	0.001 [0.029] N=1,069
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=192	-0.052 [0.032] N=1,591	-0.045 [0.032] N=1,487	-0.047 [0.035] N=1,333	-0.034 [0.038] N=1,146	-0.055 [0.039] N=946
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=305	0.029 [0.039] N=1,591	0.016 [0.041] N=1,487	0.007 [0.046] N=1,333	0.014 [0.052] N=1,146	0.029 [0.059] N=946
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=149	0.024 [0.031] N=1,591	0.024 [0.032] N=1,487	0.027 [0.035] N=1,333	0.037 [0.039] N=1,146	0.055 [0.043] N=946
CIAF: any stunting, underweight, or wasting, 0-60 months	0.272 P=410	-0.044 [0.043] N=1,534	-0.047 [0.042] N=1,434	-0.039 [0.045] N=1,284	-0.028 [0.048] N=1,103	-0.016 [0.053] N=912
BMI-for-age z score below -2	0.124 P=198	-0.031 [0.033] N=1,561	-0.027 [0.034] N=1,460	-0.021 [0.037] N=1,306	0.012 [0.041] N=1,121	-0.008 [0.043] N=927

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.304 [0.301] N=1,354	0.424 [0.346] N=1,243	0.581 [0.395] N=1,085	**0.721* [0.425] N=896**	0.172 [0.605] N=698
WHO weight-for-age z score	-0.259 [0.176] N=1,540	-0.179 [0.195] N=1,408	-0.223 [0.213] N=1,232	-0.283 [0.254] N=1,014	**-0.703* [0.356] N=790**
WHO weight-for-length/height z score, 0-60 months	0.107 [0.250] N=1,336	0.033 [0.266] N=1,232	-0.165 [0.310] N=1,078	-0.219 [0.365] N=891	-0.103 [0.447] N=691
WHO BMI-for-age z score	0.141 [0.245] N=1,313	0.115 [0.261] N=1,212	-0.069 [0.308] N=1,058	-0.307 [0.360] N=873	-0.086 [0.425] N=679
Stunted: HAZ below -2	-0.006 [0.056] N=1,354	-0.026 [0.061] N=1,243	0.001 [0.066] N=1,085	0.033 [0.075] N=896	0.116 [0.113] N=698
Underweight: WAZ below -2	0.033 [0.024] N=1,540	0.019 [0.026] N=1,408	0.018 [0.029] N=1,232	0.024 [0.032] N=1,014	0.070 [0.049] N=790
Wasted: WLZ/WHZ below -2, 0-60 months	-0.053 [0.046] N=1,336	-0.045 [0.046] N=1,232	-0.047 [0.053] N=1,078	-0.040 [0.067] N=891	-0.115 [0.076] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.047] N=1,336	-0.016 [0.051] N=1,232	-0.048 [0.058] N=1,078	-0.052 [0.075] N=891	-0.064 [0.090] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.002 [0.036] N=1,336	0.001 [0.039] N=1,232	0.001 [0.045] N=1,078	0.019 [0.054] N=891	0.042 [0.066] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.012 [0.062] N=1,288	-0.021 [0.063] N=1,188	0.002 [0.070] N=1,038	0.014 [0.086] N=857	0.047 [0.122] N=666
BMI-for-age z score below -2	-0.038 [0.043] N=1,313	-0.032 [0.043] N=1,212	-0.015 [0.053] N=1,058	0.045 [0.066] N=873	0.007 [0.074] N=679

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.011	0.119 [0.212] N=1,611	0.164 [0.228] N=1,500	0.166 [0.250] N=1,342	0.116 [0.257] N=1,153	-0.191 [0.271] N=955
WHO weight-for-age z score	0.307	-0.064 [0.117] N=1,819	-0.001 [0.123] N=1,687	-0.007 [0.130] N=1,511	-0.012 [0.136] N=1,293	-0.100 [0.158] N=1,069
WHO weight-for-length/height z score, 0-60 months	0.343	0.281 [0.190] N=1,591	0.252 [0.198] N=1,487	0.202 [0.224] N=1,333	0.203 [0.243] N=1,146	0.326 [0.272] N=946
WHO BMI-for-age z score	0.313	0.259 [0.208] N=1,561	0.255 [0.217] N=1,460	0.215 [0.246] N=1,306	0.147 [0.273] N=1,121	0.327 [0.305] N=927
Stunted: HAZ below -2	0.193 P=297	-0.016 [0.040] N=1,611	-0.025 [0.041] N=1,500	-0.006 [0.043] N=1,342	0.013 [0.045] N=1,153	0.042 [0.051] N=955
Underweight: WAZ below -2	0.052 P=99	0.010 [0.018] N=1,819	-0.000 [0.019] N=1,687	-0.003 [0.020] N=1,511	-0.005 [0.022] N=1,293	0.001 [0.029] N=1,069
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=192	-0.049 [0.032] N=1,591	-0.042 [0.033] N=1,487	-0.045 [0.035] N=1,333	-0.033 [0.039] N=1,146	-0.056 [0.040] N=946
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=305	0.028 [0.039] N=1,591	0.015 [0.041] N=1,487	0.006 [0.046] N=1,333	0.012 [0.053] N=1,146	0.025 [0.059] N=946
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=149	0.024 [0.031] N=1,591	0.024 [0.032] N=1,487	0.027 [0.035] N=1,333	0.038 [0.039] N=1,146	0.053 [0.043] N=946
CIAF: any stunting, underweight, or wasting, 0-60 months	0.272 P=410	-0.041 [0.043] N=1,534	-0.045 [0.042] N=1,434	-0.037 [0.045] N=1,284	-0.026 [0.048] N=1,103	-0.018 [0.053] N=912
BMI-for-age z score below -2	0.124 P=198	-0.027 [0.033] N=1,561	-0.024 [0.034] N=1,460	-0.019 [0.038] N=1,306	0.013 [0.041] N=1,121	-0.008 [0.043] N=927

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.328 [0.305] N=1,354	0.441 [0.349] N=1,243	0.586 [0.397] N=1,085	0.703 [0.433] N=896	0.154 [0.620] N=698
WHO weight-for-age z score	-0.232 [0.176] N=1,540	-0.148 [0.195] N=1,408	-0.206 [0.212] N=1,232	-0.280 [0.256] N=1,014	**-0.720* [0.365] N=790**
WHO weight-for-length/height z score, 0-60 months	0.121 [0.248] N=1,336	0.059 [0.265] N=1,232	-0.144 [0.311] N=1,078	-0.222 [0.370] N=891	-0.129 [0.461] N=691
WHO BMI-for-age z score	0.155 [0.244] N=1,313	0.141 [0.260] N=1,212	-0.049 [0.309] N=1,058	-0.301 [0.366] N=873	-0.094 [0.442] N=679
Stunted: HAZ below -2	-0.012 [0.056] N=1,354	-0.031 [0.061] N=1,243	-0.003 [0.065] N=1,085	0.029 [0.074] N=896	0.108 [0.110] N=698
Underweight: WAZ below -2	0.032 [0.024] N=1,540	0.017 [0.026] N=1,408	0.017 [0.029] N=1,232	0.023 [0.032] N=1,014	0.067 [0.051] N=790
Wasted: WLZ/WHZ below -2, 0-60 months	-0.053 [0.046] N=1,336	-0.046 [0.046] N=1,232	-0.047 [0.053] N=1,078	-0.038 [0.067] N=891	-0.116 [0.076] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.047] N=1,336	-0.015 [0.051] N=1,232	-0.046 [0.058] N=1,078	-0.049 [0.076] N=891	-0.064 [0.092] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.000 [0.035] N=1,336	-0.000 [0.039] N=1,232	0.001 [0.045] N=1,078	0.021 [0.054] N=891	0.043 [0.066] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.017 [0.062] N=1,288	-0.027 [0.062] N=1,188	-0.003 [0.069] N=1,038	0.010 [0.085] N=857	0.037 [0.119] N=666
BMI-for-age z score below -2	-0.036 [0.043] N=1,313	-0.032 [0.044] N=1,212	-0.015 [0.053] N=1,058	0.047 [0.066] N=873	0.006 [0.075] N=679

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.068	-0.185 [0.272] N=1,549	-0.266 [0.277] N=1,401	-0.174 [0.290] N=1,220	-0.323 [0.329] N=1,038	-0.370 [0.373] N=834
WHO weight-for-age z score	0.312	0.050 [0.133] N=1,747	0.077 [0.140] N=1,593	-0.005 [0.146] N=1,386	0.051 [0.153] N=1,173	0.087 [0.171] N=938
WHO weight-for-length/height z score, 0-60 months	0.352	0.076 [0.217] N=1,534	0.048 [0.224] N=1,391	-0.005 [0.242] N=1,214	0.114 [0.263] N=1,029	0.016 [0.302] N=827
WHO BMI-for-age z score	0.296	0.087 [0.228] N=1,502	0.078 [0.237] N=1,363	-0.007 [0.258] N=1,189	0.100 [0.288] N=1,009	0.109 [0.327] N=815
Stunted: HAZ below -2	0.182 P=281	0.016 [0.042] N=1,549	0.020 [0.043] N=1,401	-0.002 [0.045] N=1,220	0.014 [0.051] N=1,038	0.008 [0.058] N=834
Underweight: WAZ below -2	0.048 P=89	0.015 [0.021] N=1,747	0.006 [0.021] N=1,593	0.012 [0.022] N=1,386	0.008 [0.022] N=1,173	-0.011 [0.023] N=938
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=180	0.035 [0.036] N=1,534	0.036 [0.037] N=1,391	0.050 [0.040] N=1,214	0.004 [0.040] N=1,029	0.015 [0.044] N=827
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=303	0.066 [0.042] N=1,534	0.056 [0.043] N=1,391	0.040 [0.046] N=1,214	0.047 [0.051] N=1,029	0.040 [0.058] N=827
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=151	0.036 [0.034] N=1,534	0.031 [0.034] N=1,391	0.023 [0.038] N=1,214	0.024 [0.043] N=1,029	-0.008 [0.045] N=827
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=392	0.071 [0.046] N=1,480	0.063 [0.047] N=1,340	0.065 [0.050] N=1,168	0.033 [0.053] N=992	0.048 [0.057] N=796
BMI-for-age z score below -2	0.127 P=191	0.050 [0.042] N=1,502	0.049 [0.044] N=1,363	0.056 [0.049] N=1,189	0.024 [0.049] N=1,009	0.035 [0.055] N=815

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.072 [0.375] N=1,339	-0.208 [0.401] N=1,191	-0.068 [0.446] N=1,010	-0.449 [0.555] N=828	-0.541 [0.690] N=624
WHO weight-for-age z score	-0.009 [0.195] N=1,515	0.030 [0.211] N=1,361	-0.150 [0.229] N=1,154	-0.074 [0.251] N=941	0.002 [0.310] N=706
WHO weight-for-length/height z score, 0-60 months	-0.029 [0.280] N=1,326	-0.090 [0.297] N=1,183	-0.204 [0.350] N=1,006	0.030 [0.382] N=821	-0.384 [0.471] N=619
WHO BMI-for-age z score	-0.218 [0.271] N=1,303	-0.258 [0.290] N=1,164	-0.454 [0.343] N=990	-0.304 [0.394] N=810	-0.490 [0.465] N=616
Stunted: HAZ below -2	0.044 [0.066] N=1,339	0.052 [0.071] N=1,191	0.025 [0.077] N=1,010	0.077 [0.097] N=828	0.054 [0.118] N=624
Underweight: WAZ below -2	0.023 [0.034] N=1,515	0.009 [0.035] N=1,361	0.020 [0.040] N=1,154	0.008 [0.043] N=941	-0.037 [0.045] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	**0.090** [0.045] N=1,326**	**0.097** [0.049] N=1,183**	**0.135** [0.056] N=1,006**	0.039 [0.058] N=821	0.092 [0.066] N=619
Overweight: WLZ/WHZ above +2, 0-60 months	**0.109* [0.055] N=1,326**	0.091 [0.058] N=1,183	0.073 [0.065] N=1,006	0.100 [0.073] N=821	0.064 [0.086] N=619
Obese: WLZ/WHZ above +3, 0-60 months	**0.069* [0.040] N=1,326**	0.056 [0.041] N=1,183	0.046 [0.047] N=1,006	0.049 [0.059] N=821	-0.064 [0.068] N=619
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.134* [0.068] N=1,279**	**0.124* [0.072] N=1,139**	**0.151* [0.079] N=967**	0.093 [0.093] N=791	0.114 [0.108] N=595
BMI-for-age z score below -2	**0.112*** [0.042] N=1,303**	**0.115** [0.046] N=1,164**	**0.144** [0.056] N=990**	0.080 [0.055] N=810	**0.131* [0.067] N=616**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.068	-0.195 [0.270] N=1,549	-0.276 [0.275] N=1,401	-0.182 [0.288] N=1,220	-0.330 [0.327] N=1,038	-0.374 [0.370] N=834
WHO weight-for-age z score	0.312	0.052 [0.133] N=1,747	0.079 [0.140] N=1,593	-0.003 [0.146] N=1,386	0.052 [0.153] N=1,173	0.086 [0.172] N=938
WHO weight-for-length/height z score, 0-60 months	0.352	0.082 [0.217] N=1,534	0.052 [0.224] N=1,391	-0.001 [0.242] N=1,214	0.115 [0.262] N=1,029	0.014 [0.300] N=827
WHO BMI-for-age z score	0.296	0.093 [0.228] N=1,502	0.082 [0.237] N=1,363	-0.005 [0.258] N=1,189	0.100 [0.288] N=1,009	0.107 [0.327] N=815
Stunted: HAZ below -2	0.182 P=281	0.017 [0.042] N=1,549	0.021 [0.043] N=1,401	-0.001 [0.045] N=1,220	0.015 [0.051] N=1,038	0.009 [0.058] N=834
Underweight: WAZ below -2	0.048 P=89	0.015 [0.021] N=1,747	0.006 [0.021] N=1,593	0.011 [0.022] N=1,386	0.008 [0.022] N=1,173	-0.011 [0.023] N=938
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=180	0.033 [0.036] N=1,534	0.034 [0.037] N=1,391	0.048 [0.040] N=1,214	0.003 [0.040] N=1,029	0.015 [0.044] N=827
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=303	0.066 [0.042] N=1,534	0.055 [0.043] N=1,391	0.039 [0.046] N=1,214	0.046 [0.051] N=1,029	0.040 [0.058] N=827
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=151	0.035 [0.034] N=1,534	0.030 [0.034] N=1,391	0.022 [0.038] N=1,214	0.024 [0.044] N=1,029	-0.008 [0.046] N=827
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=392	0.068 [0.047] N=1,480	0.060 [0.048] N=1,340	0.062 [0.050] N=1,168	0.032 [0.053] N=992	0.048 [0.058] N=796
BMI-for-age z score below -2	0.127 P=191	0.047 [0.042] N=1,502	0.047 [0.044] N=1,363	0.054 [0.049] N=1,189	0.024 [0.049] N=1,009	0.036 [0.055] N=815

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.089 [0.372] N=1,339	-0.228 [0.398] N=1,191	-0.081 [0.441] N=1,010	-0.480 [0.549] N=828	-0.578 [0.689] N=624
WHO weight-for-age z score	-0.001 [0.193] N=1,515	0.035 [0.209] N=1,361	-0.139 [0.225] N=1,154	-0.058 [0.245] N=941	0.027 [0.304] N=706
WHO weight-for-length/height z score, 0-60 months	-0.011 [0.283] N=1,326	-0.078 [0.300] N=1,183	-0.189 [0.352] N=1,006	0.051 [0.385] N=821	-0.327 [0.485] N=619
WHO BMI-for-age z score	-0.193 [0.275] N=1,303	-0.239 [0.293] N=1,164	-0.437 [0.345] N=990	-0.279 [0.397] N=810	-0.431 [0.480] N=616
Stunted: HAZ below -2	0.045 [0.066] N=1,339	0.054 [0.071] N=1,191	0.025 [0.077] N=1,010	0.081 [0.098] N=828	0.060 [0.120] N=624
Underweight: WAZ below -2	0.023 [0.034] N=1,515	0.009 [0.035] N=1,361	0.019 [0.040] N=1,154	0.008 [0.044] N=941	-0.039 [0.046] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	**0.083* [0.046] N=1,326**	**0.088* [0.049] N=1,183**	**0.125** [0.057] N=1,006**	0.030 [0.059] N=821	0.074 [0.068] N=619
Overweight: WLZ/WHZ above +2, 0-60 months	**0.107* [0.055] N=1,326**	0.088 [0.058] N=1,183	0.068 [0.065] N=1,006	0.094 [0.072] N=821	0.060 [0.086] N=619
Obese: WLZ/WHZ above +3, 0-60 months	**0.067* [0.040] N=1,326**	0.053 [0.041] N=1,183	0.042 [0.046] N=1,006	0.044 [0.059] N=821	-0.071 [0.068] N=619
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.123* [0.069] N=1,279**	0.111 [0.073] N=1,139	**0.134* [0.080] N=967**	0.081 [0.094] N=791	0.096 [0.111] N=595
BMI-for-age z score below -2	**0.103** [0.042] N=1,303**	**0.104** [0.047] N=1,164**	**0.132** [0.056] N=990**	0.069 [0.055] N=810	**0.115* [0.069] N=616**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.068	-0.197 [0.267] N=1,549	-0.270 [0.272] N=1,401	-0.172 [0.287] N=1,220	-0.320 [0.323] N=1,038	-0.372 [0.360] N=834
WHO weight-for-age z score	0.312	0.056 [0.132] N=1,747	0.086 [0.140] N=1,593	0.015 [0.146] N=1,386	0.061 [0.153] N=1,173	0.093 [0.171] N=938
WHO weight-for-length/height z score, 0-60 months	0.352	0.086 [0.218] N=1,534	0.056 [0.224] N=1,391	0.015 [0.242] N=1,214	0.118 [0.260] N=1,029	0.020 [0.297] N=827
WHO BMI-for-age z score	0.296	0.095 [0.230] N=1,502	0.084 [0.239] N=1,363	0.008 [0.258] N=1,189	0.103 [0.285] N=1,009	0.110 [0.322] N=815
Stunted: HAZ below -2	0.182 P=281	0.018 [0.041] N=1,549	0.021 [0.043] N=1,401	-0.003 [0.044] N=1,220	0.012 [0.050] N=1,038	0.007 [0.056] N=834
Underweight: WAZ below -2	0.048 P=89	0.014 [0.021] N=1,747	0.006 [0.021] N=1,593	0.010 [0.022] N=1,386	0.007 [0.022] N=1,173	-0.011 [0.023] N=938
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=180	0.032 [0.036] N=1,534	0.032 [0.038] N=1,391	0.046 [0.041] N=1,214	0.002 [0.041] N=1,029	0.015 [0.045] N=827
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=303	0.066 [0.042] N=1,534	0.055 [0.044] N=1,391	0.040 [0.046] N=1,214	0.045 [0.051] N=1,029	0.041 [0.058] N=827
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=151	0.036 [0.034] N=1,534	0.030 [0.035] N=1,391	0.022 [0.038] N=1,214	0.024 [0.044] N=1,029	-0.007 [0.046] N=827
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=392	0.069 [0.046] N=1,480	0.060 [0.048] N=1,340	0.058 [0.051] N=1,168	0.028 [0.054] N=992	0.044 [0.058] N=796
BMI-for-age z score below -2	0.127 P=191	0.047 [0.042] N=1,502	0.046 [0.045] N=1,363	0.053 [0.049] N=1,189	0.024 [0.050] N=1,009	0.037 [0.056] N=815

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.076 [0.366] N=1,339	-0.212 [0.392] N=1,191	-0.044 [0.432] N=1,010	-0.434 [0.536] N=828	-0.518 [0.658] N=624
WHO weight-for-age z score	-0.005 [0.195] N=1,515	0.034 [0.212] N=1,361	-0.110 [0.227] N=1,154	-0.041 [0.247] N=941	0.045 [0.301] N=706
WHO weight-for-length/height z score, 0-60 months	-0.031 [0.281] N=1,326	-0.094 [0.298] N=1,183	-0.189 [0.347] N=1,006	0.034 [0.378] N=821	-0.343 [0.478] N=619
WHO BMI-for-age z score	-0.212 [0.273] N=1,303	-0.255 [0.291] N=1,164	-0.438 [0.338] N=990	-0.299 [0.389] N=810	-0.446 [0.476] N=616
Stunted: HAZ below -2	0.041 [0.065] N=1,339	0.049 [0.070] N=1,191	0.019 [0.076] N=1,010	0.070 [0.096] N=828	0.052 [0.118] N=624
Underweight: WAZ below -2	0.021 [0.034] N=1,515	0.007 [0.035] N=1,361	0.014 [0.040] N=1,154	0.003 [0.044] N=941	-0.043 [0.046] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	**0.084* [0.046] N=1,326**	**0.088* [0.050] N=1,183**	**0.124** [0.057] N=1,006**	0.030 [0.060] N=821	0.077 [0.070] N=619
Overweight: WLZ/WHZ above +2, 0-60 months	**0.107* [0.055] N=1,326**	0.087 [0.058] N=1,183	0.066 [0.064] N=1,006	0.089 [0.073] N=821	0.058 [0.086] N=619
Obese: WLZ/WHZ above +3, 0-60 months	0.064 [0.040] N=1,326	0.050 [0.041] N=1,183	0.040 [0.046] N=1,006	0.039 [0.059] N=821	-0.074 [0.068] N=619
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.123* [0.069] N=1,279**	0.110 [0.074] N=1,139	0.128 [0.082] N=967	0.073 [0.096] N=791	0.095 [0.112] N=595
BMI-for-age z score below -2	**0.102** [0.043] N=1,303**	**0.103** [0.047] N=1,164**	**0.131** [0.056] N=990**	0.069 [0.055] N=810	**0.116* [0.069] N=616**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.068	-0.158 [0.268] N=1,549	-0.221 [0.274] N=1,401	-0.120 [0.290] N=1,220	-0.289 [0.325] N=1,038	-0.362 [0.366] N=834
WHO weight-for-age z score	0.312	0.061 [0.133] N=1,747	0.089 [0.141] N=1,593	0.022 [0.147] N=1,386	0.068 [0.153] N=1,173	0.099 [0.172] N=938
WHO weight-for-length/height z score, 0-60 months	0.352	0.064 [0.217] N=1,534	0.032 [0.223] N=1,391	-0.014 [0.241] N=1,214	0.108 [0.262] N=1,029	0.029 [0.301] N=827
WHO BMI-for-age z score	0.296	0.065 [0.228] N=1,502	0.054 [0.237] N=1,363	-0.024 [0.256] N=1,189	0.086 [0.286] N=1,009	0.119 [0.327] N=815
Stunted: HAZ below -2	0.182 P=281	0.011 [0.042] N=1,549	0.012 [0.043] N=1,401	-0.014 [0.045] N=1,220	0.005 [0.051] N=1,038	0.004 [0.057] N=834
Underweight: WAZ below -2	0.048 P=89	0.012 [0.021] N=1,747	0.003 [0.021] N=1,593	0.006 [0.023] N=1,386	0.004 [0.023] N=1,173	-0.016 [0.024] N=938
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=180	0.034 [0.036] N=1,534	0.035 [0.038] N=1,391	0.051 [0.040] N=1,214	0.006 [0.040] N=1,029	0.015 [0.043] N=827
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=303	0.061 [0.041] N=1,534	0.050 [0.043] N=1,391	0.037 [0.046] N=1,214	0.046 [0.051] N=1,029	0.043 [0.058] N=827
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=151	0.037 [0.033] N=1,534	0.032 [0.034] N=1,391	0.023 [0.037] N=1,214	0.026 [0.042] N=1,029	-0.006 [0.045] N=827
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=392	0.066 [0.046] N=1,480	0.058 [0.048] N=1,340	0.057 [0.050] N=1,168	0.029 [0.053] N=992	0.050 [0.056] N=796
BMI-for-age z score below -2	0.127 P=191	0.047 [0.042] N=1,502	0.047 [0.044] N=1,363	0.054 [0.049] N=1,189	0.024 [0.049] N=1,009	0.033 [0.055] N=815

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.043 [0.364] N=1,339	-0.152 [0.392] N=1,191	0.034 [0.439] N=1,010	-0.400 [0.551] N=828	-0.591 [0.682] N=624
WHO weight-for-age z score	0.026 [0.190] N=1,515	0.066 [0.208] N=1,361	-0.072 [0.223] N=1,154	-0.012 [0.240] N=941	0.062 [0.285] N=706
WHO weight-for-length/height z score, 0-60 months	-0.033 [0.280] N=1,326	-0.106 [0.296] N=1,183	-0.220 [0.345] N=1,006	0.030 [0.383] N=821	-0.284 [0.500] N=619
WHO BMI-for-age z score	-0.231 [0.273] N=1,303	-0.281 [0.290] N=1,164	-0.478 [0.337] N=990	-0.313 [0.390] N=810	-0.387 [0.493] N=616
Stunted: HAZ below -2	0.038 [0.064] N=1,339	0.040 [0.069] N=1,191	0.002 [0.077] N=1,010	0.057 [0.098] N=828	0.052 [0.120] N=624
Underweight: WAZ below -2	0.019 [0.034] N=1,515	0.004 [0.035] N=1,361	0.009 [0.040] N=1,154	-0.001 [0.043] N=941	-0.049 [0.045] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	**0.086* [0.045] N=1,326**	**0.092* [0.049] N=1,183**	**0.133** [0.056] N=1,006**	0.035 [0.058] N=821	0.077 [0.067] N=619
Overweight: WLZ/WHZ above +2, 0-60 months	**0.102* [0.055] N=1,326**	0.081 [0.057] N=1,183	0.062 [0.064] N=1,006	0.087 [0.073] N=821	0.065 [0.088] N=619
Obese: WLZ/WHZ above +3, 0-60 months	**0.067* [0.040] N=1,326**	0.052 [0.041] N=1,183	0.039 [0.046] N=1,006	0.040 [0.060] N=821	-0.071 [0.070] N=619
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.121* [0.069] N=1,279**	0.107 [0.074] N=1,139	0.120 [0.082] N=967	0.064 [0.096] N=791	0.099 [0.112] N=595
BMI-for-age z score below -2	**0.105** [0.042] N=1,303**	**0.108** [0.047] N=1,164**	**0.138** [0.056] N=990**	0.074 [0.054] N=810	**0.114* [0.068] N=616**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.060	-0.268 [0.271] N=1,434	-0.368 [0.279] N=1,297	-0.300 [0.290] N=1,133	-0.459 [0.317] N=961	-0.577 [0.354] N=774
WHO weight-for-age z score	0.308	0.069 [0.138] N=1,616	0.109 [0.146] N=1,473	0.006 [0.153] N=1,286	0.044 [0.159] N=1,089	0.033 [0.172] N=874
WHO weight-for-length/height z score, 0-60 months	0.359	0.136 [0.234] N=1,422	0.135 [0.244] N=1,290	0.048 [0.263] N=1,128	0.135 [0.281] N=955	0.045 [0.310] N=771
WHO BMI-for-age z score	0.300	0.105 [0.247] N=1,391	0.128 [0.261] N=1,263	-0.008 [0.284] N=1,105	0.063 [0.310] N=937	0.056 [0.351] N=760
Stunted: HAZ below -2	0.182 P=261	0.025 [0.043] N=1,434	0.035 [0.045] N=1,297	0.011 [0.045] N=1,133	0.025 [0.050] N=961	0.031 [0.055] N=774
Underweight: WAZ below -2	0.048 P=83	0.018 [0.022] N=1,616	0.009 [0.023] N=1,473	0.011 [0.025] N=1,286	0.011 [0.025] N=1,089	-0.001 [0.026] N=874
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=167	0.035 [0.038] N=1,422	0.033 [0.039] N=1,290	0.050 [0.041] N=1,128	0.009 [0.041] N=955	0.022 [0.044] N=771
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=283	0.068 [0.044] N=1,422	0.059 [0.046] N=1,290	0.040 [0.049] N=1,128	0.041 [0.053] N=955	0.036 [0.058] N=771
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=143	0.037 [0.034] N=1,422	0.032 [0.035] N=1,290	0.024 [0.038] N=1,128	0.030 [0.044] N=955	0.002 [0.044] N=771
CIAF: any stunting, underweight, or wasting, 0-60 months	0.265 P=366	0.078 [0.049] N=1,372	0.069 [0.050] N=1,243	0.071 [0.051] N=1,086	0.038 [0.054] N=920	0.068 [0.057] N=741
BMI-for-age z score below -2	0.125 P=174	0.041 [0.043] N=1,391	0.035 [0.045] N=1,263	0.045 [0.050] N=1,105	0.023 [0.051] N=937	0.040 [0.057] N=760

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.039 [0.375] N=1,237	-0.120 [0.404] N=1,100	0.008 [0.450] N=936	-0.344 [0.540] N=764	-0.478 [0.657] N=577
WHO weight-for-age z score	0.053 [0.202] N=1,398	0.103 [0.217] N=1,255	-0.107 [0.237] N=1,068	-0.027 [0.257] N=871	0.030 [0.314] N=656
WHO weight-for-length/height z score, 0-60 months	0.022 [0.312] N=1,225	-0.009 [0.335] N=1,093	-0.194 [0.392] N=931	-0.002 [0.426] N=758	-0.386 [0.542] N=574
WHO BMI-for-age z score	-0.184 [0.300] N=1,202	-0.192 [0.326] N=1,074	-0.476 [0.384] N=916	-0.354 [0.434] N=748	-0.553 [0.529] N=571
Stunted: HAZ below -2	0.029 [0.071] N=1,237	0.047 [0.077] N=1,100	0.015 [0.083] N=936	0.054 [0.103] N=764	0.047 [0.131] N=577
Underweight: WAZ below -2	0.022 [0.037] N=1,398	0.005 [0.038] N=1,255	0.009 [0.044] N=1,068	0.003 [0.051] N=871	-0.042 [0.055] N=656
Wasted: WLZ/WHZ below -2, 0-60 months	**0.085* [0.049] N=1,225**	0.084 [0.052] N=1,093	**0.122** [0.058] N=931**	0.024 [0.062] N=758	0.065 [0.071] N=574
Overweight: WLZ/WHZ above +2, 0-60 months	**0.105* [0.061] N=1,225**	0.086 [0.065] N=1,093	0.058 [0.072] N=931	0.067 [0.082] N=758	0.032 [0.099] N=574
Obese: WLZ/WHZ above +3, 0-60 months	0.055 [0.044] N=1,225	0.040 [0.045] N=1,093	0.027 [0.052] N=931	0.028 [0.066] N=758	-0.088 [0.079] N=574
CIAF: any stunting, underweight, or wasting, 0-60 months	0.121 [0.077] N=1,182	0.108 [0.083] N=1,053	0.129 [0.088] N=896	0.052 [0.106] N=730	0.074 [0.132] N=551
BMI-for-age z score below -2	**0.102** [0.045] N=1,202**	**0.096* [0.050] N=1,074**	**0.124** [0.058] N=916**	0.063 [0.058] N=748	**0.125* [0.072] N=571**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.060	-0.280 [0.274] N=1,434	-0.375 [0.282] N=1,297	-0.302 [0.292] N=1,133	-0.479 [0.318] N=961	**-0.598* [0.357] N=774**
WHO weight-for-age z score	0.308	0.069 [0.138] N=1,616	0.107 [0.146] N=1,473	0.005 [0.153] N=1,286	0.041 [0.159] N=1,089	0.031 [0.174] N=874
WHO weight-for-length/height z score, 0-60 months	0.359	0.145 [0.237] N=1,422	0.137 [0.247] N=1,290	0.053 [0.265] N=1,128	0.145 [0.281] N=955	0.048 [0.318] N=771
WHO BMI-for-age z score	0.300	0.113 [0.251] N=1,391	0.126 [0.264] N=1,263	-0.007 [0.287] N=1,105	0.072 [0.312] N=937	0.061 [0.357] N=760
Stunted: HAZ below -2	0.182 P=261	0.025 [0.043] N=1,434	0.033 [0.045] N=1,297	0.009 [0.045] N=1,133	0.024 [0.049] N=961	0.030 [0.055] N=774
Underweight: WAZ below -2	0.048 P=83	0.017 [0.022] N=1,616	0.008 [0.022] N=1,473	0.010 [0.024] N=1,286	0.010 [0.025] N=1,089	-0.003 [0.025] N=874
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=167	0.031 [0.038] N=1,422	0.029 [0.039] N=1,290	0.045 [0.041] N=1,128	0.003 [0.041] N=955	0.016 [0.044] N=771
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=283	0.067 [0.044] N=1,422	0.058 [0.046] N=1,290	0.038 [0.048] N=1,128	0.040 [0.053] N=955	0.035 [0.059] N=771
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=143	0.036 [0.034] N=1,422	0.030 [0.034] N=1,290	0.022 [0.038] N=1,128	0.028 [0.043] N=955	-0.000 [0.044] N=771
CIAF: any stunting, underweight, or wasting, 0-60 months	0.265 P=366	0.073 [0.048] N=1,372	0.063 [0.050] N=1,243	0.065 [0.050] N=1,086	0.031 [0.053] N=920	0.061 [0.056] N=741
BMI-for-age z score below -2	0.125 P=174	0.038 [0.044] N=1,391	0.033 [0.046] N=1,263	0.043 [0.051] N=1,105	0.019 [0.051] N=937	0.036 [0.056] N=760

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.021 [0.377] N=1,237	-0.125 [0.404] N=1,100	0.007 [0.448] N=936	-0.408 [0.539] N=764	-0.597 [0.658] N=577
WHO weight-for-age z score	0.047 [0.202] N=1,398	0.095 [0.218] N=1,255	-0.114 [0.238] N=1,068	-0.044 [0.255] N=871	-0.014 [0.314] N=656
WHO weight-for-length/height z score, 0-60 months	0.014 [0.309] N=1,225	-0.032 [0.330] N=1,093	-0.214 [0.387] N=931	-0.003 [0.416] N=758	-0.406 [0.521] N=574
WHO BMI-for-age z score	-0.189 [0.298] N=1,202	-0.214 [0.320] N=1,074	-0.492 [0.379] N=916	-0.338 [0.425] N=748	-0.525 [0.520] N=571
Stunted: HAZ below -2	0.030 [0.071] N=1,237	0.044 [0.077] N=1,100	0.011 [0.082] N=936	0.057 [0.102] N=764	0.056 [0.128] N=577
Underweight: WAZ below -2	0.022 [0.036] N=1,398	0.007 [0.038] N=1,255	0.010 [0.044] N=1,068	0.006 [0.050] N=871	-0.033 [0.053] N=656
Wasted: WLZ/WHZ below -2, 0-60 months	**0.086* [0.048] N=1,225**	**0.087* [0.051] N=1,093**	**0.125** [0.058] N=931**	0.024 [0.060] N=758	0.072 [0.070] N=574
Overweight: WLZ/WHZ above +2, 0-60 months	**0.105* [0.060] N=1,225**	0.084 [0.064] N=1,093	0.054 [0.071] N=931	0.068 [0.081] N=758	0.037 [0.096] N=574
Obese: WLZ/WHZ above +3, 0-60 months	0.055 [0.043] N=1,225	0.038 [0.044] N=1,093	0.025 [0.051] N=931	0.029 [0.064] N=758	-0.082 [0.077] N=574
CIAF: any stunting, underweight, or wasting, 0-60 months	0.121 [0.077] N=1,182	0.110 [0.082] N=1,053	0.131 [0.087] N=896	0.055 [0.105] N=730	0.091 [0.127] N=551
BMI-for-age z score below -2	**0.103** [0.044] N=1,202**	**0.101** [0.049] N=1,074**	**0.129** [0.057] N=916**	0.063 [0.055] N=748	**0.125* [0.069] N=571**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.