Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-3: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=1,599	0.001 [0.023] N=1,707	0.014 [0.023] N=1,574	0.004 [0.025] N=1,417	-0.004 [0.027] N=1,208	-0.012 [0.032] N=989
Vegetables consumed at least weekly	0.897 P=1,530	0.022 [0.027] N=1,707	0.022 [0.028] N=1,574	0.003 [0.029] N=1,417	0.005 [0.030] N=1,208	-0.035 [0.035] N=989
Meat fish or poultry consumed at least weekly	0.778 P=1,341	**0.091* [0.047] N=1,707**	**0.111** [0.049] N=1,574**	0.089 [0.054] N=1,417	0.071 [0.060] N=1,208	0.069 [0.069] N=989
Legumes consumed at least weekly	0.777 P=1,336	-0.021 [0.042] N=1,707	-0.032 [0.043] N=1,574	-0.033 [0.043] N=1,417	-0.037 [0.048] N=1,208	-0.087 [0.056] N=989
Animal products consumed at least weekly	0.909 P=1,554	0.024 [0.028] N=1,707	0.024 [0.029] N=1,574	0.011 [0.032] N=1,417	0.007 [0.035] N=1,208	-0.003 [0.039] N=989
Sweets consumed at least weekly	0.646 P=1,105	-0.014 [0.038] N=1,707	0.003 [0.040] N=1,574	0.010 [0.042] N=1,417	-0.004 [0.045] N=1,208	0.012 [0.053] N=989
Snack foods consumed at least weekly	0.528 P=897	-0.036 [0.039] N=1,707	-0.026 [0.041] N=1,574	-0.009 [0.045] N=1,417	-0.011 [0.049] N=1,208	-0.032 [0.055] N=989
Packaged soft drinks consumed at least weekly	0.323 P=543	0.002 [0.047] N=1,707	0.006 [0.051] N=1,574	0.007 [0.054] N=1,417	0.029 [0.058] N=1,208	0.018 [0.067] N=989
Grains consumed at least weekly	0.892 P=1,527	0.018 [0.030] N=1,707	0.028 [0.032] N=1,574	0.011 [0.033] N=1,417	0.020 [0.034] N=1,208	-0.000 [0.040] N=989
Child was ever breastfed	0.968 P=1,836	0.000 [0.014] N=1,902	0.003 [0.015] N=1,756	0.004 [0.016] N=1,580	0.009 [0.018] N=1,345	-0.001 [0.018] N=1,105

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.011 [0.033] N=1,457	0.037 [0.034] N=1,324	0.020 [0.037] N=1,167	0.007 [0.045] N=958	-0.005 [0.062] N=739
Vegetables consumed at least weekly	0.049 [0.041] N=1,457	0.054 [0.044] N=1,324	0.025 [0.047] N=1,167	0.038 [0.056] N=958	-0.064 [0.074] N=739
Meat fish or poultry consumed at least weekly	0.087 [0.055] N=1,457	**0.115* [0.059] N=1,324**	0.076 [0.065] N=1,167	0.012 [0.072] N=958	-0.032 [0.085] N=739
Legumes consumed at least weekly	0.040 [0.059] N=1,457	0.028 [0.063] N=1,324	0.042 [0.065] N=1,167	0.056 [0.080] N=958	-0.027 [0.108] N=739
Animal products consumed at least weekly	0.028 [0.038] N=1,457	0.031 [0.042] N=1,324	0.005 [0.048] N=1,167	0.006 [0.057] N=958	-0.035 [0.073] N=739
Sweets consumed at least weekly	-0.024 [0.051] N=1,457	0.003 [0.055] N=1,324	0.011 [0.061] N=1,167	-0.028 [0.070] N=958	0.024 [0.098] N=739
Snack foods consumed at least weekly	-0.039 [0.049] N=1,457	-0.024 [0.052] N=1,324	0.007 [0.062] N=1,167	0.001 [0.075] N=958	-0.055 [0.092] N=739
Packaged soft drinks consumed at least weekly	-0.024 [0.056] N=1,457	-0.021 [0.062] N=1,324	-0.023 [0.064] N=1,167	0.013 [0.075] N=958	-0.035 [0.090] N=739
Grains consumed at least weekly	0.064 [0.048] N=1,457	0.089 [0.055] N=1,324	0.071 [0.059] N=1,167	0.112 [0.072] N=958	0.098 [0.106] N=739
Child was ever breastfed	0.007 [0.023] N=1,617	0.013 [0.026] N=1,471	0.016 [0.031] N=1,295	0.027 [0.039] N=1,060	0.018 [0.050] N=820

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=1,599	0.001 [0.023] N=1,707	0.014 [0.023] N=1,574	0.004 [0.024] N=1,417	-0.004 [0.027] N=1,208	-0.012 [0.032] N=989
Vegetables consumed at least weekly	0.897 P=1,530	0.022 [0.027] N=1,707	0.022 [0.028] N=1,574	0.004 [0.029] N=1,417	0.005 [0.030] N=1,208	-0.035 [0.034] N=989
Meat fish or poultry consumed at least weekly	0.778 P=1,341	**0.091* [0.047] N=1,707**	**0.111** [0.050] N=1,574**	0.090 [0.054] N=1,417	0.072 [0.061] N=1,208	0.070 [0.071] N=989
Legumes consumed at least weekly	0.777 P=1,336	-0.021 [0.042] N=1,707	-0.032 [0.043] N=1,574	-0.032 [0.044] N=1,417	-0.036 [0.049] N=1,208	-0.086 [0.056] N=989
Animal products consumed at least weekly	0.909 P=1,554	0.024 [0.028] N=1,707	0.025 [0.029] N=1,574	0.012 [0.032] N=1,417	0.008 [0.034] N=1,208	-0.003 [0.038] N=989
Sweets consumed at least weekly	0.646 P=1,105	-0.014 [0.038] N=1,707	0.003 [0.040] N=1,574	0.010 [0.042] N=1,417	-0.004 [0.045] N=1,208	0.012 [0.053] N=989
Snack foods consumed at least weekly	0.528 P=897	-0.036 [0.040] N=1,707	-0.026 [0.042] N=1,574	-0.008 [0.045] N=1,417	-0.010 [0.050] N=1,208	-0.032 [0.056] N=989
Packaged soft drinks consumed at least weekly	0.323 P=543	0.002 [0.047] N=1,707	0.006 [0.051] N=1,574	0.007 [0.054] N=1,417	0.030 [0.058] N=1,208	0.018 [0.068] N=989
Grains consumed at least weekly	0.892 P=1,527	0.018 [0.030] N=1,707	0.028 [0.032] N=1,574	0.012 [0.032] N=1,417	0.019 [0.034] N=1,208	-0.001 [0.040] N=989
Child was ever breastfed	0.968 P=1,836	0.000 [0.014] N=1,902	0.003 [0.015] N=1,756	0.004 [0.016] N=1,580	0.009 [0.018] N=1,345	-0.001 [0.018] N=1,105

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.012 [0.033] N=1,457	0.038 [0.034] N=1,324	0.020 [0.037] N=1,167	0.007 [0.045] N=958	-0.005 [0.063] N=739
Vegetables consumed at least weekly	0.049 [0.041] N=1,457	0.055 [0.044] N=1,324	0.025 [0.047] N=1,167	0.038 [0.056] N=958	-0.064 [0.074] N=739
Meat fish or poultry consumed at least weekly	0.087 [0.055] N=1,457	**0.115* [0.059] N=1,324**	0.077 [0.065] N=1,167	0.014 [0.072] N=958	-0.027 [0.084] N=739
Legumes consumed at least weekly	0.041 [0.059] N=1,457	0.029 [0.063] N=1,324	0.045 [0.065] N=1,167	0.059 [0.080] N=958	-0.025 [0.108] N=739
Animal products consumed at least weekly	0.029 [0.038] N=1,457	0.033 [0.041] N=1,324	0.009 [0.048] N=1,167	0.009 [0.057] N=958	-0.032 [0.072] N=739
Sweets consumed at least weekly	-0.024 [0.051] N=1,457	0.003 [0.055] N=1,324	0.011 [0.062] N=1,167	-0.028 [0.070] N=958	0.023 [0.098] N=739
Snack foods consumed at least weekly	-0.038 [0.050] N=1,457	-0.021 [0.053] N=1,324	0.012 [0.062] N=1,167	0.005 [0.077] N=958	-0.050 [0.093] N=739
Packaged soft drinks consumed at least weekly	-0.024 [0.055] N=1,457	-0.021 [0.062] N=1,324	-0.024 [0.064] N=1,167	0.013 [0.074] N=958	-0.037 [0.090] N=739
Grains consumed at least weekly	0.064 [0.048] N=1,457	0.089 [0.055] N=1,324	0.071 [0.059] N=1,167	0.111 [0.071] N=958	0.097 [0.106] N=739
Child was ever breastfed	0.008 [0.023] N=1,617	0.014 [0.026] N=1,471	0.017 [0.031] N=1,295	0.027 [0.039] N=1,060	0.019 [0.050] N=820

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=1,599	-0.009 [0.023] N=1,707	0.004 [0.023] N=1,574	-0.005 [0.025] N=1,417	-0.011 [0.028] N=1,208	-0.021 [0.033] N=989
Vegetables consumed at least weekly	0.897 P=1,530	0.016 [0.027] N=1,707	0.014 [0.028] N=1,574	-0.003 [0.029] N=1,417	-0.000 [0.030] N=1,208	-0.043 [0.034] N=989
Meat fish or poultry consumed at least weekly	0.778 P=1,341	0.066 [0.047] N=1,707	0.080 [0.049] N=1,574	0.065 [0.054] N=1,417	0.053 [0.059] N=1,208	0.045 [0.068] N=989
Legumes consumed at least weekly	0.777 P=1,336	-0.029 [0.042] N=1,707	-0.041 [0.043] N=1,574	-0.040 [0.044] N=1,417	-0.041 [0.048] N=1,208	-0.090 [0.055] N=989
Animal products consumed at least weekly	0.909 P=1,554	0.018 [0.029] N=1,707	0.017 [0.030] N=1,574	0.005 [0.033] N=1,417	0.003 [0.036] N=1,208	-0.011 [0.040] N=989
Sweets consumed at least weekly	0.646 P=1,105	-0.020 [0.038] N=1,707	-0.003 [0.040] N=1,574	0.003 [0.042] N=1,417	-0.008 [0.045] N=1,208	0.005 [0.052] N=989
Snack foods consumed at least weekly	0.528 P=897	-0.033 [0.040] N=1,707	-0.023 [0.042] N=1,574	-0.007 [0.045] N=1,417	-0.008 [0.048] N=1,208	-0.029 [0.055] N=989
Packaged soft drinks consumed at least weekly	0.323 P=543	0.017 [0.045] N=1,707	0.025 [0.049] N=1,574	0.022 [0.052] N=1,417	0.042 [0.056] N=1,208	0.035 [0.064] N=989
Grains consumed at least weekly	0.892 P=1,527	0.012 [0.030] N=1,707	0.022 [0.032] N=1,574	0.007 [0.033] N=1,417	0.015 [0.034] N=1,208	-0.005 [0.040] N=989
Child was ever breastfed	0.968 P=1,836	-0.003 [0.014] N=1,902	-0.002 [0.015] N=1,756	0.000 [0.016] N=1,580	0.006 [0.018] N=1,345	-0.004 [0.019] N=1,105

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.005 [0.033] N=1,457	0.028 [0.033] N=1,324	0.016 [0.037] N=1,167	0.017 [0.045] N=958	0.007 [0.063] N=739
Vegetables consumed at least weekly	0.045 [0.041] N=1,457	0.047 [0.043] N=1,324	0.021 [0.047] N=1,167	0.044 [0.055] N=958	-0.056 [0.073] N=739
Meat fish or poultry consumed at least weekly	0.066 [0.054] N=1,457	0.086 [0.057] N=1,324	0.059 [0.064] N=1,167	0.015 [0.071] N=958	-0.024 [0.086] N=739
Legumes consumed at least weekly	0.035 [0.060] N=1,457	0.020 [0.063] N=1,324	0.040 [0.065] N=1,167	0.066 [0.080] N=958	-0.014 [0.109] N=739
Animal products consumed at least weekly	0.025 [0.038] N=1,457	0.026 [0.042] N=1,324	0.005 [0.048] N=1,167	0.016 [0.058] N=958	-0.018 [0.073] N=739
Sweets consumed at least weekly	-0.027 [0.052] N=1,457	-0.001 [0.057] N=1,324	0.009 [0.063] N=1,167	-0.021 [0.072] N=958	0.033 [0.102] N=739
Snack foods consumed at least weekly	-0.034 [0.049] N=1,457	-0.018 [0.053] N=1,324	0.015 [0.060] N=1,167	0.014 [0.074] N=958	-0.041 [0.090] N=739
Packaged soft drinks consumed at least weekly	-0.011 [0.054] N=1,457	-0.002 [0.059] N=1,324	-0.009 [0.062] N=1,167	0.023 [0.072] N=958	-0.029 [0.085] N=739
Grains consumed at least weekly	0.058 [0.048] N=1,457	0.084 [0.055] N=1,324	0.068 [0.059] N=1,167	0.114 [0.072] N=958	0.099 [0.106] N=739
Child was ever breastfed	0.005 [0.023] N=1,617	0.010 [0.026] N=1,471	0.015 [0.031] N=1,295	0.030 [0.039] N=1,060	0.018 [0.050] N=820

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.937 P=1,599	-0.010 [0.023] N=1,707	0.003 [0.023] N=1,574	-0.008 [0.025] N=1,417	-0.015 [0.028] N=1,208	-0.023 [0.033] N=989
Vegetables consumed at least weekly	0.897 P=1,530	0.015 [0.027] N=1,707	0.015 [0.028] N=1,574	-0.003 [0.029] N=1,417	-0.002 [0.030] N=1,208	-0.043 [0.035] N=989
Meat fish or poultry consumed at least weekly	0.778 P=1,341	0.060 [0.047] N=1,707	0.075 [0.049] N=1,574	0.057 [0.054] N=1,417	0.043 [0.060] N=1,208	0.033 [0.069] N=989
Legumes consumed at least weekly	0.777 P=1,336	-0.033 [0.043] N=1,707	-0.046 [0.044] N=1,574	-0.049 [0.045] N=1,417	-0.047 [0.050] N=1,208	**-0.096* [0.057] N=989**
Animal products consumed at least weekly	0.909 P=1,554	0.016 [0.029] N=1,707	0.017 [0.030] N=1,574	0.003 [0.033] N=1,417	-0.002 [0.035] N=1,208	-0.015 [0.039] N=989
Sweets consumed at least weekly	0.646 P=1,105	-0.019 [0.038] N=1,707	-0.005 [0.039] N=1,574	-0.002 [0.041] N=1,417	-0.012 [0.045] N=1,208	0.000 [0.053] N=989
Snack foods consumed at least weekly	0.528 P=897	-0.033 [0.039] N=1,707	-0.024 [0.041] N=1,574	-0.012 [0.044] N=1,417	-0.011 [0.048] N=1,208	-0.033 [0.054] N=989
Packaged soft drinks consumed at least weekly	0.323 P=543	0.021 [0.045] N=1,707	0.026 [0.049] N=1,574	0.023 [0.052] N=1,417	0.046 [0.056] N=1,208	0.043 [0.063] N=989
Grains consumed at least weekly	0.892 P=1,527	0.011 [0.030] N=1,707	0.020 [0.032] N=1,574	0.004 [0.033] N=1,417	0.012 [0.035] N=1,208	-0.006 [0.041] N=989
Child was ever breastfed	0.968 P=1,836	-0.004 [0.014] N=1,902	-0.002 [0.016] N=1,756	-0.001 [0.016] N=1,580	0.004 [0.018] N=1,345	-0.005 [0.019] N=1,105

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.033] N=1,457	0.029 [0.034] N=1,324	0.014 [0.037] N=1,167	0.007 [0.045] N=958	-0.009 [0.064] N=739
Vegetables consumed at least weekly	0.043 [0.040] N=1,457	0.047 [0.043] N=1,324	0.019 [0.047] N=1,167	0.036 [0.056] N=958	-0.073 [0.074] N=739
Meat fish or poultry consumed at least weekly	0.061 [0.054] N=1,457	0.085 [0.057] N=1,324	0.056 [0.065] N=1,167	0.008 [0.073] N=958	-0.040 [0.088] N=739
Legumes consumed at least weekly	0.038 [0.059] N=1,457	0.023 [0.063] N=1,324	0.041 [0.064] N=1,167	0.076 [0.079] N=958	-0.008 [0.113] N=739
Animal products consumed at least weekly	0.024 [0.038] N=1,457	0.028 [0.042] N=1,324	0.004 [0.048] N=1,167	0.006 [0.058] N=958	-0.034 [0.076] N=739
Sweets consumed at least weekly	-0.022 [0.051] N=1,457	0.002 [0.055] N=1,324	0.011 [0.061] N=1,167	-0.012 [0.072] N=958	0.041 [0.102] N=739
Snack foods consumed at least weekly	-0.028 [0.049] N=1,457	-0.014 [0.052] N=1,324	0.018 [0.058] N=1,167	0.027 [0.072] N=958	-0.022 [0.089] N=739
Packaged soft drinks consumed at least weekly	-0.004 [0.053] N=1,457	-0.002 [0.059] N=1,324	-0.008 [0.062] N=1,167	0.031 [0.073] N=958	-0.015 [0.085] N=739
Grains consumed at least weekly	0.058 [0.047] N=1,457	0.082 [0.054] N=1,324	0.066 [0.058] N=1,167	0.111 [0.070] N=958	0.088 [0.102] N=739
Child was ever breastfed	0.006 [0.023] N=1,617	0.011 [0.027] N=1,471	0.014 [0.031] N=1,295	0.028 [0.040] N=1,060	0.017 [0.051] N=820

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,486	-0.006 [0.024] N=1,585	0.004 [0.024] N=1,467	-0.007 [0.026] N=1,319	-0.012 [0.029] N=1,128	-0.028 [0.033] N=925
Vegetables consumed at least weekly	0.899 P=1,423	0.011 [0.029] N=1,585	0.008 [0.029] N=1,467	-0.015 [0.030] N=1,319	-0.008 [0.031] N=1,128	-0.050 [0.034] N=925
Meat fish or poultry consumed at least weekly	0.778 P=1,243	**0.078* [0.046] N=1,585**	**0.087* [0.047] N=1,467**	0.058 [0.051] N=1,319	0.039 [0.056] N=1,128	0.022 [0.065] N=925
Legumes consumed at least weekly	0.775 P=1,237	-0.037 [0.042] N=1,585	-0.053 [0.044] N=1,467	-0.059 [0.043] N=1,319	-0.058 [0.049] N=1,128	**-0.112** [0.055] N=925**
Animal products consumed at least weekly	0.911 P=1,446	0.023 [0.029] N=1,585	0.023 [0.030] N=1,467	0.008 [0.033] N=1,319	-0.001 [0.035] N=1,128	-0.016 [0.039] N=925
Sweets consumed at least weekly	0.650 P=1,030	-0.001 [0.039] N=1,585	0.011 [0.042] N=1,467	0.013 [0.043] N=1,319	0.005 [0.048] N=1,128	0.016 [0.057] N=925
Snack foods consumed at least weekly	0.531 P=836	-0.039 [0.043] N=1,585	-0.030 [0.045] N=1,467	-0.015 [0.048] N=1,319	-0.014 [0.054] N=1,128	-0.024 [0.062] N=925
Packaged soft drinks consumed at least weekly	0.329 P=515	0.028 [0.047] N=1,585	0.034 [0.051] N=1,467	0.023 [0.053] N=1,319	0.045 [0.057] N=1,128	0.048 [0.066] N=925
Grains consumed at least weekly	0.894 P=1,421	0.009 [0.031] N=1,585	0.016 [0.033] N=1,467	-0.008 [0.033] N=1,319	0.005 [0.035] N=1,128	-0.015 [0.040] N=925
Child was ever breastfed	0.968 P=1,697	-0.004 [0.015] N=1,758	0.000 [0.016] N=1,628	0.002 [0.017] N=1,461	0.009 [0.019] N=1,251	-0.002 [0.020] N=1,030

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.036] N=1,350	0.027 [0.037] N=1,232	0.007 [0.040] N=1,084	0.000 [0.048] N=893	-0.027 [0.065] N=690
Vegetables consumed at least weekly	0.038 [0.043] N=1,350	0.037 [0.046] N=1,232	-0.004 [0.049] N=1,084	0.022 [0.057] N=893	-0.080 [0.072] N=690
Meat fish or poultry consumed at least weekly	**0.100* [0.056] N=1,350**	**0.111* [0.059] N=1,232**	0.065 [0.064] N=1,084	0.006 [0.073] N=893	-0.048 [0.086] N=690
Legumes consumed at least weekly	0.032 [0.061] N=1,350	0.009 [0.064] N=1,232	0.013 [0.064] N=1,084	0.036 [0.080] N=893	-0.064 [0.109] N=690
Animal products consumed at least weekly	0.032 [0.040] N=1,350	0.034 [0.044] N=1,232	0.005 [0.050] N=1,084	0.004 [0.060] N=893	-0.041 [0.076] N=690
Sweets consumed at least weekly	-0.011 [0.052] N=1,350	0.006 [0.058] N=1,232	-0.001 [0.063] N=1,084	-0.031 [0.072] N=893	-0.003 [0.102] N=690
Snack foods consumed at least weekly	-0.048 [0.050] N=1,350	-0.033 [0.054] N=1,232	-0.010 [0.062] N=1,084	-0.019 [0.076] N=893	-0.075 [0.092] N=690
Packaged soft drinks consumed at least weekly	0.002 [0.056] N=1,350	0.005 [0.060] N=1,232	-0.018 [0.062] N=1,084	0.013 [0.072] N=893	-0.006 [0.092] N=690
Grains consumed at least weekly	0.059 [0.050] N=1,350	0.080 [0.057] N=1,232	0.041 [0.060] N=1,084	0.095 [0.073] N=893	0.079 [0.105] N=690
Child was ever breastfed	-0.000 [0.024] N=1,491	0.008 [0.028] N=1,361	0.011 [0.032] N=1,194	0.024 [0.041] N=984	0.007 [0.052] N=763

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,486	-0.007 [0.024] N=1,585	0.002 [0.023] N=1,467	-0.008 [0.025] N=1,319	-0.013 [0.028] N=1,128	-0.028 [0.032] N=925
Vegetables consumed at least weekly	0.899 P=1,423	0.011 [0.028] N=1,585	0.008 [0.029] N=1,467	-0.016 [0.030] N=1,319	-0.007 [0.031] N=1,128	-0.048 [0.034] N=925
Meat fish or poultry consumed at least weekly	0.778 P=1,243	**0.079* [0.046] N=1,585**	**0.089* [0.048] N=1,467**	0.062 [0.052] N=1,319	0.046 [0.057] N=1,128	0.038 [0.066] N=925
Legumes consumed at least weekly	0.775 P=1,237	-0.039 [0.042] N=1,585	-0.055 [0.044] N=1,467	-0.062 [0.043] N=1,319	-0.062 [0.049] N=1,128	**-0.120** [0.055] N=925**
Animal products consumed at least weekly	0.911 P=1,446	0.021 [0.029] N=1,585	0.021 [0.030] N=1,467	0.007 [0.032] N=1,319	0.001 [0.035] N=1,128	-0.013 [0.039] N=925
Sweets consumed at least weekly	0.650 P=1,030	-0.001 [0.039] N=1,585	0.010 [0.041] N=1,467	0.013 [0.043] N=1,319	0.007 [0.047] N=1,128	0.019 [0.055] N=925
Snack foods consumed at least weekly	0.531 P=836	-0.040 [0.042] N=1,585	-0.032 [0.045] N=1,467	-0.016 [0.048] N=1,319	-0.016 [0.053] N=1,128	-0.030 [0.061] N=925
Packaged soft drinks consumed at least weekly	0.329 P=515	0.026 [0.047] N=1,585	0.031 [0.051] N=1,467	0.022 [0.054] N=1,319	0.042 [0.057] N=1,128	0.039 [0.067] N=925
Grains consumed at least weekly	0.894 P=1,421	0.008 [0.031] N=1,585	0.016 [0.033] N=1,467	-0.009 [0.033] N=1,319	0.004 [0.035] N=1,128	-0.017 [0.039] N=925
Child was ever breastfed	0.968 P=1,697	-0.003 [0.015] N=1,758	0.001 [0.016] N=1,628	0.002 [0.017] N=1,461	0.010 [0.019] N=1,251	-0.002 [0.019] N=1,030

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.007 [0.035] N=1,350	0.026 [0.035] N=1,232	0.008 [0.039] N=1,084	0.001 [0.047] N=893	-0.024 [0.063] N=690
Vegetables consumed at least weekly	0.036 [0.042] N=1,350	0.036 [0.046] N=1,232	-0.004 [0.049] N=1,084	0.021 [0.057] N=893	-0.075 [0.070] N=690
Meat fish or poultry consumed at least weekly	**0.092* [0.056] N=1,350**	**0.104* [0.058] N=1,232**	0.062 [0.064] N=1,084	0.007 [0.073] N=893	-0.034 [0.085] N=690
Legumes consumed at least weekly	0.029 [0.060] N=1,350	0.006 [0.064] N=1,232	0.015 [0.065] N=1,084	0.037 [0.081] N=893	-0.066 [0.109] N=690
Animal products consumed at least weekly	0.029 [0.040] N=1,350	0.031 [0.043] N=1,232	0.005 [0.049] N=1,084	0.003 [0.059] N=893	-0.037 [0.076] N=690
Sweets consumed at least weekly	-0.018 [0.052] N=1,350	-0.004 [0.058] N=1,232	-0.007 [0.063] N=1,084	-0.032 [0.073] N=893	0.003 [0.102] N=690
Snack foods consumed at least weekly	-0.054 [0.050] N=1,350	-0.041 [0.054] N=1,232	-0.014 [0.062] N=1,084	-0.020 [0.077] N=893	-0.074 [0.093] N=690
Packaged soft drinks consumed at least weekly	-0.004 [0.056] N=1,350	-0.001 [0.061] N=1,232	-0.020 [0.063] N=1,084	0.012 [0.073] N=893	-0.011 [0.091] N=690
Grains consumed at least weekly	0.057 [0.050] N=1,350	0.078 [0.057] N=1,232	0.043 [0.061] N=1,084	0.095 [0.074] N=893	0.079 [0.107] N=690
Child was ever breastfed	0.003 [0.024] N=1,491	0.011 [0.028] N=1,361	0.014 [0.033] N=1,194	0.030 [0.041] N=984	0.013 [0.054] N=763

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,412	0.019 [0.028] N=1,503	0.021 [0.029] N=1,364	0.012 [0.032] N=1,191	0.013 [0.033] N=1,003	0.033 [0.040] N=808
Vegetables consumed at least weekly	0.899 P=1,350	0.034 [0.034] N=1,503	0.046 [0.036] N=1,364	0.055 [0.038] N=1,191	0.056 [0.042] N=1,003	0.071 [0.048] N=808
Meat fish or poultry consumed at least weekly	0.768 P=1,169	**0.100** [0.039] N=1,503**	**0.106** [0.041] N=1,364**	**0.094** [0.044] N=1,191**	**0.138*** [0.045] N=1,003**	**0.150*** [0.054] N=808**
Legumes consumed at least weekly	0.778 P=1,180	-0.020 [0.043] N=1,503	-0.000 [0.045] N=1,364	-0.004 [0.049] N=1,191	-0.026 [0.053] N=1,003	0.017 [0.066] N=808
Animal products consumed at least weekly	0.910 P=1,370	0.021 [0.036] N=1,503	0.032 [0.037] N=1,364	0.040 [0.040] N=1,191	0.027 [0.041] N=1,003	0.066 [0.047] N=808
Sweets consumed at least weekly	0.653 P=989	-0.008 [0.049] N=1,503	-0.006 [0.052] N=1,364	-0.011 [0.056] N=1,191	-0.014 [0.059] N=1,003	-0.065 [0.067] N=808
Snack foods consumed at least weekly	0.529 P=793	-0.039 [0.046] N=1,503	-0.031 [0.050] N=1,364	-0.027 [0.053] N=1,191	-0.021 [0.058] N=1,003	-0.060 [0.068] N=808
Packaged soft drinks consumed at least weekly	0.309 P=460	-0.021 [0.044] N=1,503	-0.017 [0.045] N=1,364	0.021 [0.047] N=1,191	0.029 [0.053] N=1,003	0.028 [0.059] N=808
Grains consumed at least weekly	0.892 P=1,345	0.043 [0.038] N=1,503	0.053 [0.041] N=1,364	0.065 [0.044] N=1,191	0.072 [0.048] N=1,003	0.083 [0.055] N=808
Child was ever breastfed	0.966 P=1,619	-0.012 [0.020] N=1,677	-0.011 [0.020] N=1,527	-0.001 [0.021] N=1,331	-0.008 [0.021] N=1,125	-0.025 [0.025] N=903

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.025 [0.037] N=1,300	-0.027 [0.039] N=1,161	-0.049 [0.048] N=988	-0.041 [0.052] N=800	-0.011 [0.079] N=605
Vegetables consumed at least weekly	-0.030 [0.045] N=1,300	-0.021 [0.048] N=1,161	-0.017 [0.054] N=988	-0.014 [0.061] N=800	0.008 [0.081] N=605
Meat fish or poultry consumed at least weekly	0.027 [0.056] N=1,300	0.031 [0.060] N=1,161	-0.020 [0.067] N=988	0.045 [0.068] N=800	0.044 [0.093] N=605
Legumes consumed at least weekly	-0.041 [0.052] N=1,300	-0.012 [0.056] N=1,161	-0.020 [0.062] N=988	-0.044 [0.065] N=800	0.060 [0.086] N=605
Animal products consumed at least weekly	-0.055 [0.045] N=1,300	-0.040 [0.050] N=1,161	-0.037 [0.055] N=988	-0.076 [0.059] N=800	-0.014 [0.073] N=605
Sweets consumed at least weekly	-0.003 [0.062] N=1,300	0.003 [0.067] N=1,161	-0.010 [0.074] N=988	-0.012 [0.090] N=800	-0.154 [0.114] N=605
Snack foods consumed at least weekly	-0.048 [0.065] N=1,300	-0.035 [0.071] N=1,161	-0.031 [0.077] N=988	-0.005 [0.094] N=800	-0.110 [0.126] N=605
Packaged soft drinks consumed at least weekly	-0.064 [0.070] N=1,300	-0.061 [0.075] N=1,161	-0.004 [0.086] N=988	0.005 [0.107] N=800	-0.022 [0.136] N=605
Grains consumed at least weekly	-0.032 [0.040] N=1,300	-0.030 [0.044] N=1,161	-0.023 [0.047] N=988	-0.043 [0.055] N=800	-0.090 [0.074] N=605
Child was ever breastfed	0.007 [0.038] N=1,450	0.012 [0.039] N=1,300	0.039 [0.042] N=1,104	0.031 [0.047] N=898	0.007 [0.064] N=676

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,412	0.019 [0.028] N=1,503	0.021 [0.029] N=1,364	0.012 [0.032] N=1,191	0.013 [0.033] N=1,003	0.033 [0.040] N=808
Vegetables consumed at least weekly	0.899 P=1,350	0.035 [0.034] N=1,503	0.046 [0.036] N=1,364	0.055 [0.038] N=1,191	0.056 [0.041] N=1,003	0.070 [0.047] N=808
Meat fish or poultry consumed at least weekly	0.768 P=1,169	**0.100** [0.039] N=1,503**	**0.107*** [0.041] N=1,364**	**0.095** [0.044] N=1,191**	**0.138*** [0.045] N=1,003**	**0.149*** [0.054] N=808**
Legumes consumed at least weekly	0.778 P=1,180	-0.019 [0.043] N=1,503	-0.000 [0.045] N=1,364	-0.004 [0.049] N=1,191	-0.026 [0.053] N=1,003	0.018 [0.066] N=808
Animal products consumed at least weekly	0.910 P=1,370	0.023 [0.035] N=1,503	0.033 [0.037] N=1,364	0.041 [0.040] N=1,191	0.026 [0.041] N=1,003	0.064 [0.046] N=808
Sweets consumed at least weekly	0.653 P=989	-0.007 [0.050] N=1,503	-0.006 [0.052] N=1,364	-0.010 [0.056] N=1,191	-0.014 [0.059] N=1,003	-0.066 [0.067] N=808
Snack foods consumed at least weekly	0.529 P=793	-0.038 [0.046] N=1,503	-0.030 [0.050] N=1,364	-0.026 [0.052] N=1,191	-0.022 [0.058] N=1,003	-0.062 [0.067] N=808
Packaged soft drinks consumed at least weekly	0.309 P=460	-0.020 [0.043] N=1,503	-0.016 [0.045] N=1,364	0.022 [0.047] N=1,191	0.029 [0.053] N=1,003	0.027 [0.058] N=808
Grains consumed at least weekly	0.892 P=1,345	0.043 [0.038] N=1,503	0.053 [0.041] N=1,364	0.065 [0.044] N=1,191	0.072 [0.048] N=1,003	0.084 [0.055] N=808
Child was ever breastfed	0.966 P=1,619	-0.012 [0.020] N=1,677	-0.011 [0.020] N=1,527	-0.001 [0.021] N=1,331	-0.008 [0.021] N=1,125	-0.025 [0.025] N=903

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.025 [0.037] N=1,300	-0.026 [0.039] N=1,161	-0.049 [0.049] N=988	-0.041 [0.053] N=800	-0.012 [0.079] N=605
Vegetables consumed at least weekly	-0.029 [0.044] N=1,300	-0.020 [0.048] N=1,161	-0.016 [0.054] N=988	-0.012 [0.060] N=800	0.010 [0.080] N=605
Meat fish or poultry consumed at least weekly	0.029 [0.056] N=1,300	0.034 [0.059] N=1,161	-0.017 [0.067] N=988	0.048 [0.068] N=800	0.049 [0.093] N=605
Legumes consumed at least weekly	-0.040 [0.051] N=1,300	-0.012 [0.056] N=1,161	-0.021 [0.062] N=988	-0.045 [0.065] N=800	0.056 [0.086] N=605
Animal products consumed at least weekly	-0.051 [0.043] N=1,300	-0.036 [0.048] N=1,161	-0.033 [0.054] N=988	-0.072 [0.057] N=800	-0.008 [0.070] N=605
Sweets consumed at least weekly	-0.001 [0.063] N=1,300	0.006 [0.067] N=1,161	-0.008 [0.074] N=988	-0.012 [0.090] N=800	-0.153 [0.114] N=605
Snack foods consumed at least weekly	-0.043 [0.065] N=1,300	-0.031 [0.071] N=1,161	-0.027 [0.077] N=988	-0.002 [0.094] N=800	-0.104 [0.127] N=605
Packaged soft drinks consumed at least weekly	-0.062 [0.069] N=1,300	-0.061 [0.075] N=1,161	-0.002 [0.085] N=988	0.006 [0.107] N=800	-0.016 [0.134] N=605
Grains consumed at least weekly	-0.033 [0.040] N=1,300	-0.031 [0.044] N=1,161	-0.026 [0.047] N=988	-0.046 [0.054] N=800	-0.095 [0.075] N=605
Child was ever breastfed	0.009 [0.038] N=1,450	0.014 [0.039] N=1,300	0.041 [0.042] N=1,104	0.033 [0.047] N=898	0.011 [0.065] N=676

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,412	0.018 [0.027] N=1,503	0.021 [0.029] N=1,364	0.014 [0.032] N=1,191	0.016 [0.034] N=1,003	0.035 [0.041] N=808
Vegetables consumed at least weekly	0.899 P=1,350	0.034 [0.034] N=1,503	0.046 [0.035] N=1,364	0.058 [0.038] N=1,191	0.059 [0.041] N=1,003	0.071 [0.048] N=808
Meat fish or poultry consumed at least weekly	0.768 P=1,169	**0.096** [0.038] N=1,503**	**0.106*** [0.040] N=1,364**	**0.103** [0.043] N=1,191**	**0.146*** [0.044] N=1,003**	**0.156*** [0.051] N=808**
Legumes consumed at least weekly	0.778 P=1,180	-0.021 [0.043] N=1,503	-0.001 [0.046] N=1,364	-0.003 [0.049] N=1,191	-0.024 [0.054] N=1,003	0.021 [0.066] N=808
Animal products consumed at least weekly	0.910 P=1,370	0.021 [0.036] N=1,503	0.032 [0.037] N=1,364	0.043 [0.040] N=1,191	0.028 [0.041] N=1,003	0.066 [0.046] N=808
Sweets consumed at least weekly	0.653 P=989	-0.007 [0.049] N=1,503	-0.005 [0.052] N=1,364	-0.008 [0.055] N=1,191	-0.013 [0.058] N=1,003	-0.064 [0.067] N=808
Snack foods consumed at least weekly	0.529 P=793	-0.035 [0.047] N=1,503	-0.029 [0.050] N=1,364	-0.027 [0.053] N=1,191	-0.022 [0.058] N=1,003	-0.062 [0.068] N=808
Packaged soft drinks consumed at least weekly	0.309 P=460	-0.019 [0.045] N=1,503	-0.016 [0.047] N=1,364	0.017 [0.049] N=1,191	0.023 [0.055] N=1,003	0.024 [0.061] N=808
Grains consumed at least weekly	0.892 P=1,345	0.042 [0.038] N=1,503	0.053 [0.041] N=1,364	0.067 [0.044] N=1,191	0.074 [0.049] N=1,003	0.086 [0.056] N=808
Child was ever breastfed	0.966 P=1,619	-0.012 [0.020] N=1,677	-0.010 [0.021] N=1,527	0.001 [0.021] N=1,331	-0.006 [0.021] N=1,125	-0.024 [0.025] N=903

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.025 [0.036] N=1,300	-0.025 [0.038] N=1,161	-0.043 [0.047] N=988	-0.036 [0.053] N=800	-0.008 [0.080] N=605
Vegetables consumed at least weekly	-0.029 [0.044] N=1,300	-0.019 [0.048] N=1,161	-0.011 [0.053] N=988	-0.008 [0.060] N=800	0.014 [0.081] N=605
Meat fish or poultry consumed at least weekly	0.025 [0.054] N=1,300	0.034 [0.056] N=1,161	-0.001 [0.063] N=988	0.062 [0.067] N=800	0.056 [0.087] N=605
Legumes consumed at least weekly	-0.038 [0.051] N=1,300	-0.010 [0.056] N=1,161	-0.019 [0.062] N=988	-0.039 [0.065] N=800	0.057 [0.087] N=605
Animal products consumed at least weekly	-0.054 [0.043] N=1,300	-0.038 [0.047] N=1,161	-0.031 [0.053] N=988	-0.074 [0.059] N=800	-0.009 [0.069] N=605
Sweets consumed at least weekly	-0.001 [0.062] N=1,300	0.004 [0.067] N=1,161	-0.007 [0.073] N=988	-0.015 [0.090] N=800	-0.155 [0.114] N=605
Snack foods consumed at least weekly	-0.036 [0.065] N=1,300	-0.028 [0.071] N=1,161	-0.029 [0.076] N=988	-0.006 [0.093] N=800	-0.104 [0.128] N=605
Packaged soft drinks consumed at least weekly	-0.064 [0.071] N=1,300	-0.065 [0.077] N=1,161	-0.019 [0.089] N=988	-0.014 [0.109] N=800	-0.025 [0.137] N=605
Grains consumed at least weekly	-0.035 [0.040] N=1,300	-0.034 [0.044] N=1,161	-0.023 [0.046] N=988	-0.048 [0.054] N=800	-0.098 [0.074] N=605
Child was ever breastfed	0.008 [0.038] N=1,450	0.014 [0.039] N=1,300	0.044 [0.043] N=1,104	0.036 [0.047] N=898	0.010 [0.063] N=676

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,412	0.017 [0.027] N=1,503	0.020 [0.028] N=1,364	0.013 [0.032] N=1,191	0.010 [0.033] N=1,003	0.026 [0.040] N=808
Vegetables consumed at least weekly	0.899 P=1,350	0.031 [0.035] N=1,503	0.044 [0.036] N=1,364	0.055 [0.039] N=1,191	0.054 [0.042] N=1,003	0.067 [0.047] N=808
Meat fish or poultry consumed at least weekly	0.768 P=1,169	**0.095** [0.040] N=1,503**	**0.104** [0.041] N=1,364**	**0.096** [0.044] N=1,191**	**0.132*** [0.046] N=1,003**	**0.133** [0.053] N=808**
Legumes consumed at least weekly	0.778 P=1,180	-0.016 [0.044] N=1,503	0.005 [0.046] N=1,364	0.003 [0.049] N=1,191	-0.023 [0.053] N=1,003	0.021 [0.065] N=808
Animal products consumed at least weekly	0.910 P=1,370	0.027 [0.035] N=1,503	0.039 [0.036] N=1,364	0.051 [0.039] N=1,191	0.033 [0.041] N=1,003	0.068 [0.047] N=808
Sweets consumed at least weekly	0.653 P=989	0.002 [0.051] N=1,503	0.002 [0.053] N=1,364	-0.003 [0.057] N=1,191	-0.010 [0.061] N=1,003	-0.062 [0.070] N=808
Snack foods consumed at least weekly	0.529 P=793	-0.023 [0.047] N=1,503	-0.018 [0.050] N=1,364	-0.016 [0.053] N=1,191	-0.014 [0.058] N=1,003	-0.050 [0.068] N=808
Packaged soft drinks consumed at least weekly	0.309 P=460	-0.005 [0.045] N=1,503	-0.004 [0.047] N=1,364	0.029 [0.049] N=1,191	0.036 [0.054] N=1,003	0.039 [0.061] N=808
Grains consumed at least weekly	0.892 P=1,345	0.042 [0.038] N=1,503	0.052 [0.040] N=1,364	0.065 [0.043] N=1,191	0.072 [0.048] N=1,003	0.082 [0.055] N=808
Child was ever breastfed	0.966 P=1,619	-0.012 [0.020] N=1,677	-0.010 [0.020] N=1,527	-0.000 [0.021] N=1,331	-0.008 [0.021] N=1,125	-0.027 [0.025] N=903

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.026 [0.036] N=1,300	-0.027 [0.038] N=1,161	-0.044 [0.047] N=988	-0.047 [0.054] N=800	-0.031 [0.078] N=605
Vegetables consumed at least weekly	-0.033 [0.044] N=1,300	-0.021 [0.048] N=1,161	-0.014 [0.054] N=988	-0.015 [0.060] N=800	0.003 [0.078] N=605
Meat fish or poultry consumed at least weekly	0.025 [0.054] N=1,300	0.032 [0.057] N=1,161	-0.009 [0.062] N=988	0.037 [0.069] N=800	0.004 [0.086] N=605
Legumes consumed at least weekly	-0.034 [0.051] N=1,300	-0.004 [0.057] N=1,161	-0.006 [0.063] N=988	-0.035 [0.066] N=800	0.066 [0.085] N=605
Animal products consumed at least weekly	-0.046 [0.041] N=1,300	-0.028 [0.045] N=1,161	-0.019 [0.051] N=988	-0.066 [0.056] N=800	-0.013 [0.065] N=605
Sweets consumed at least weekly	0.012 [0.063] N=1,300	0.017 [0.068] N=1,161	0.003 [0.075] N=988	-0.009 [0.091] N=800	-0.152 [0.112] N=605
Snack foods consumed at least weekly	-0.025 [0.064] N=1,300	-0.017 [0.071] N=1,161	-0.015 [0.076] N=988	0.004 [0.092] N=800	-0.090 [0.123] N=605
Packaged soft drinks consumed at least weekly	-0.044 [0.071] N=1,300	-0.047 [0.076] N=1,161	0.001 [0.087] N=988	0.015 [0.109] N=800	0.019 [0.138] N=605
Grains consumed at least weekly	-0.034 [0.038] N=1,300	-0.033 [0.043] N=1,161	-0.024 [0.046] N=988	-0.050 [0.054] N=800	-0.107 [0.073] N=605
Child was ever breastfed	0.009 [0.037] N=1,450	0.015 [0.039] N=1,300	0.044 [0.042] N=1,104	0.034 [0.047] N=898	0.004 [0.065] N=676

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,311	0.014 [0.029] N=1,393	0.015 [0.030] N=1,262	0.015 [0.034] N=1,107	0.005 [0.035] N=932	0.023 [0.042] N=754
Vegetables consumed at least weekly	0.902 P=1,256	0.032 [0.034] N=1,393	0.041 [0.036] N=1,262	0.061 [0.038] N=1,107	0.053 [0.041] N=932	0.052 [0.048] N=754
Meat fish or poultry consumed at least weekly	0.768 P=1,083	**0.093** [0.042] N=1,393**	**0.096** [0.044] N=1,262**	**0.093* [0.047] N=1,107**	**0.110** [0.050] N=932**	**0.121** [0.058] N=754**
Legumes consumed at least weekly	0.777 P=1,094	-0.019 [0.046] N=1,393	0.002 [0.049] N=1,262	0.001 [0.052] N=1,107	-0.035 [0.058] N=932	0.007 [0.072] N=754
Animal products consumed at least weekly	0.912 P=1,274	0.018 [0.037] N=1,393	0.032 [0.039] N=1,262	0.047 [0.041] N=1,107	0.026 [0.042] N=932	0.063 [0.047] N=754
Sweets consumed at least weekly	0.664 P=930	-0.006 [0.051] N=1,393	-0.003 [0.054] N=1,262	-0.015 [0.058] N=1,107	-0.027 [0.063] N=932	-0.074 [0.074] N=754
Snack foods consumed at least weekly	0.537 P=746	-0.028 [0.047] N=1,393	-0.022 [0.051] N=1,262	-0.030 [0.054] N=1,107	-0.027 [0.059] N=932	-0.050 [0.071] N=754
Packaged soft drinks consumed at least weekly	0.320 P=443	-0.026 [0.044] N=1,393	-0.028 [0.046] N=1,262	0.008 [0.048] N=1,107	0.019 [0.055] N=932	0.016 [0.061] N=754
Grains consumed at least weekly	0.895 P=1,251	0.056 [0.040] N=1,393	0.067 [0.043] N=1,262	**0.080* [0.045] N=1,107**	0.083 [0.051] N=932	0.092 [0.057] N=754
Child was ever breastfed	0.968 P=1,499	-0.017 [0.019] N=1,549	-0.018 [0.020] N=1,410	-0.008 [0.019] N=1,234	-0.019 [0.020] N=1,044	-0.034 [0.025] N=841

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.031 [0.038] N=1,201	-0.033 [0.040] N=1,070	-0.037 [0.050] N=915	-0.055 [0.053] N=740	-0.037 [0.077] N=562
Vegetables consumed at least weekly	-0.033 [0.042] N=1,201	-0.025 [0.045] N=1,070	-0.001 [0.050] N=915	-0.017 [0.056] N=740	-0.038 [0.070] N=562
Meat fish or poultry consumed at least weekly	0.039 [0.064] N=1,201	0.044 [0.068] N=1,070	0.012 [0.076] N=915	0.023 [0.080] N=740	0.007 [0.107] N=562
Legumes consumed at least weekly	-0.025 [0.054] N=1,201	0.009 [0.057] N=1,070	0.014 [0.063] N=915	-0.028 [0.068] N=740	0.067 [0.089] N=562
Animal products consumed at least weekly	-0.065 [0.044] N=1,201	-0.043 [0.049] N=1,070	-0.028 [0.051] N=915	-0.084 [0.055] N=740	-0.028 [0.064] N=562
Sweets consumed at least weekly	0.002 [0.065] N=1,201	0.009 [0.070] N=1,070	-0.024 [0.076] N=915	-0.049 [0.089] N=740	**-0.205* [0.119] N=562**
Snack foods consumed at least weekly	-0.025 [0.067] N=1,201	-0.016 [0.073] N=1,070	-0.036 [0.079] N=915	-0.019 [0.096] N=740	-0.079 [0.136] N=562
Packaged soft drinks consumed at least weekly	-0.075 [0.069] N=1,201	-0.083 [0.074] N=1,070	-0.033 [0.086] N=915	-0.025 [0.109] N=740	-0.070 [0.140] N=562
Grains consumed at least weekly	-0.012 [0.044] N=1,201	-0.007 [0.049] N=1,070	0.004 [0.053] N=915	-0.026 [0.063] N=740	-0.098 [0.085] N=562
Child was ever breastfed	0.005 [0.036] N=1,335	0.006 [0.037] N=1,196	0.034 [0.039] N=1,020	0.019 [0.043] N=830	-0.004 [0.063] N=627

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,311	0.015 [0.029] N=1,393	0.016 [0.030] N=1,262	0.016 [0.034] N=1,107	0.005 [0.035] N=932	0.022 [0.042] N=754
Vegetables consumed at least weekly	0.902 P=1,256	0.031 [0.034] N=1,393	0.041 [0.036] N=1,262	0.061 [0.038] N=1,107	0.050 [0.041] N=932	0.049 [0.048] N=754
Meat fish or poultry consumed at least weekly	0.768 P=1,083	**0.097** [0.042] N=1,393**	**0.101** [0.043] N=1,262**	**0.096** [0.046] N=1,107**	**0.114** [0.048] N=932**	**0.123** [0.056] N=754**
Legumes consumed at least weekly	0.777 P=1,094	-0.021 [0.046] N=1,393	0.001 [0.049] N=1,262	-0.000 [0.052] N=1,107	-0.039 [0.057] N=932	0.003 [0.072] N=754
Animal products consumed at least weekly	0.912 P=1,274	0.017 [0.037] N=1,393	0.031 [0.039] N=1,262	0.046 [0.041] N=1,107	0.024 [0.042] N=932	0.061 [0.046] N=754
Sweets consumed at least weekly	0.664 P=930	-0.007 [0.051] N=1,393	-0.004 [0.054] N=1,262	-0.015 [0.059] N=1,107	-0.028 [0.063] N=932	-0.075 [0.075] N=754
Snack foods consumed at least weekly	0.537 P=746	-0.026 [0.046] N=1,393	-0.022 [0.050] N=1,262	-0.029 [0.053] N=1,107	-0.026 [0.059] N=932	-0.051 [0.071] N=754
Packaged soft drinks consumed at least weekly	0.320 P=443	-0.025 [0.044] N=1,393	-0.027 [0.046] N=1,262	0.010 [0.048] N=1,107	0.018 [0.055] N=932	0.015 [0.062] N=754
Grains consumed at least weekly	0.895 P=1,251	0.055 [0.040] N=1,393	0.066 [0.043] N=1,262	**0.078* [0.045] N=1,107**	0.082 [0.050] N=932	0.090 [0.057] N=754
Child was ever breastfed	0.968 P=1,499	-0.016 [0.019] N=1,549	-0.017 [0.020] N=1,410	-0.006 [0.020] N=1,234	-0.015 [0.021] N=1,044	-0.030 [0.025] N=841

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.030 [0.038] N=1,201	-0.030 [0.041] N=1,070	-0.035 [0.051] N=915	-0.055 [0.054] N=740	-0.038 [0.081] N=562
Vegetables consumed at least weekly	-0.033 [0.042] N=1,201	-0.026 [0.045] N=1,070	-0.002 [0.051] N=915	-0.022 [0.056] N=740	-0.041 [0.072] N=562
Meat fish or poultry consumed at least weekly	0.044 [0.063] N=1,201	0.050 [0.067] N=1,070	0.016 [0.075] N=915	0.029 [0.080] N=740	0.011 [0.108] N=562
Legumes consumed at least weekly	-0.025 [0.054] N=1,201	0.010 [0.058] N=1,070	0.013 [0.064] N=915	-0.038 [0.068] N=740	0.061 [0.087] N=562
Animal products consumed at least weekly	-0.065 [0.044] N=1,201	-0.043 [0.049] N=1,070	-0.029 [0.052] N=915	-0.086 [0.055] N=740	-0.036 [0.065] N=562
Sweets consumed at least weekly	0.005 [0.065] N=1,201	0.012 [0.070] N=1,070	-0.017 [0.077] N=915	-0.039 [0.092] N=740	-0.191 [0.119] N=562
Snack foods consumed at least weekly	-0.024 [0.066] N=1,201	-0.018 [0.073] N=1,070	-0.035 [0.079] N=915	-0.013 [0.096] N=740	-0.084 [0.134] N=562
Packaged soft drinks consumed at least weekly	-0.074 [0.070] N=1,201	-0.082 [0.075] N=1,070	-0.025 [0.086] N=915	-0.017 [0.108] N=740	-0.059 [0.140] N=562
Grains consumed at least weekly	-0.013 [0.043] N=1,201	-0.009 [0.048] N=1,070	0.001 [0.052] N=915	-0.030 [0.062] N=740	-0.103 [0.083] N=562
Child was ever breastfed	0.004 [0.036] N=1,335	0.002 [0.038] N=1,196	0.033 [0.040] N=1,020	0.022 [0.045] N=830	-0.005 [0.065] N=627

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.