Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.613	-0.007 [0.210] N=1,666	-0.026 [0.226] N=1,544	0.004 [0.250] N=1,389	0.012 [0.254] N=1,186	-0.210 [0.270] N=977
WHO weight-for-age z score	-0.147	-0.115 [0.122] N=1,887	-0.086 [0.131] N=1,742	-0.083 [0.140] N=1,567	-0.069 [0.146] N=1,335	-0.168 [0.159] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.319	0.232 [0.196] N=1,642	0.203 [0.208] N=1,526	0.156 [0.232] N=1,373	0.138 [0.254] N=1,171	0.266 [0.284] N=962
WHO BMI-for-age z score	0.287	0.325 [0.218] N=1,609	0.332 [0.234] N=1,496	0.276 [0.260] N=1,344	0.186 [0.294] N=1,148	0.322 [0.329] N=946
Stunted: HAZ below -2	0.274 P=441	-0.009 [0.039] N=1,666	-0.017 [0.042] N=1,544	-0.005 [0.044] N=1,389	-0.010 [0.046] N=1,186	0.029 [0.050] N=977
Underweight: WAZ below -2	0.093 P=171	**0.043* [0.022] N=1,887**	0.038 [0.024] N=1,742	0.027 [0.025] N=1,567	0.027 [0.027] N=1,335	**0.053* [0.031] N=1,097**
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=199	-0.039 [0.033] N=1,642	-0.034 [0.033] N=1,526	-0.036 [0.035] N=1,373	-0.026 [0.039] N=1,171	-0.049 [0.040] N=962
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=310	0.025 [0.040] N=1,642	0.017 [0.043] N=1,526	0.001 [0.046] N=1,373	0.003 [0.052] N=1,171	0.005 [0.060] N=962
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=144	0.029 [0.028] N=1,642	0.030 [0.029] N=1,526	0.031 [0.033] N=1,373	0.042 [0.035] N=1,171	0.062 [0.041] N=962
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=575	-0.039 [0.043] N=1,588	-0.045 [0.044] N=1,476	-0.039 [0.047] N=1,328	-0.048 [0.050] N=1,133	-0.028 [0.057] N=932
BMI-for-age z score below -2	0.132 P=215	-0.033 [0.037] N=1,609	-0.032 [0.038] N=1,496	-0.028 [0.041] N=1,344	-0.000 [0.045] N=1,148	-0.011 [0.050] N=946

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.191 [0.308] N=1,407	0.156 [0.351] N=1,285	0.310 [0.422] N=1,130	0.502 [0.453] N=927	0.173 [0.596] N=718
WHO weight-for-age z score	-0.251 [0.179] N=1,604	-0.228 [0.201] N=1,459	-0.259 [0.225] N=1,284	-0.283 [0.279] N=1,052	**-0.701** [0.344] N=814**
WHO weight-for-length/height z score, 0-60 months	0.076 [0.261] N=1,385	0.011 [0.290] N=1,269	-0.166 [0.343] N=1,116	-0.311 [0.390] N=914	-0.300 [0.478] N=705
WHO BMI-for-age z score	0.274 [0.269] N=1,358	0.275 [0.303] N=1,245	0.105 [0.361] N=1,093	-0.203 [0.417] N=897	-0.117 [0.494] N=695
Stunted: HAZ below -2	-0.063 [0.056] N=1,407	-0.085 [0.062] N=1,285	-0.084 [0.070] N=1,130	-0.133 [0.080] N=927	-0.124 [0.108] N=718
Underweight: WAZ below -2	0.043 [0.033] N=1,604	0.033 [0.037] N=1,459	0.012 [0.043] N=1,284	0.013 [0.053] N=1,052	0.097 [0.070] N=814
Wasted: WLZ/WHZ below -2, 0-60 months	-0.039 [0.047] N=1,385	-0.034 [0.049] N=1,269	-0.038 [0.056] N=1,116	-0.020 [0.070] N=914	-0.079 [0.081] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.008 [0.048] N=1,385	-0.008 [0.054] N=1,269	-0.053 [0.059] N=1,116	-0.058 [0.074] N=914	-0.114 [0.090] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.003 [0.035] N=1,385	0.002 [0.039] N=1,269	-0.006 [0.045] N=1,116	0.014 [0.053] N=914	0.048 [0.069] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.073 [0.061] N=1,341	-0.091 [0.065] N=1,229	-0.096 [0.074] N=1,081	-0.147 [0.089] N=886	-0.173 [0.123] N=685
BMI-for-age z score below -2	-0.051 [0.048] N=1,358	-0.052 [0.049] N=1,245	-0.047 [0.056] N=1,093	0.012 [0.069] N=897	-0.021 [0.083] N=695

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.613	-0.008 [0.210] N=1,666	-0.027 [0.226] N=1,544	0.003 [0.250] N=1,389	0.015 [0.255] N=1,186	-0.202 [0.270] N=977
WHO weight-for-age z score	-0.147	-0.117 [0.121] N=1,887	-0.088 [0.130] N=1,742	-0.084 [0.139] N=1,567	-0.070 [0.146] N=1,335	-0.175 [0.158] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.319	0.226 [0.196] N=1,642	0.196 [0.209] N=1,526	0.149 [0.232] N=1,373	0.130 [0.255] N=1,171	0.247 [0.285] N=962
WHO BMI-for-age z score	0.287	0.323 [0.219] N=1,609	0.327 [0.235] N=1,496	0.272 [0.261] N=1,344	0.178 [0.295] N=1,148	0.306 [0.331] N=946
Stunted: HAZ below -2	0.274 P=441	-0.008 [0.039] N=1,666	-0.017 [0.042] N=1,544	-0.004 [0.044] N=1,389	-0.009 [0.046] N=1,186	0.029 [0.050] N=977
Underweight: WAZ below -2	0.093 P=171	**0.044* [0.022] N=1,887**	0.038 [0.024] N=1,742	0.028 [0.025] N=1,567	0.027 [0.027] N=1,335	**0.054* [0.031] N=1,097**
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=199	-0.037 [0.033] N=1,642	-0.032 [0.034] N=1,526	-0.033 [0.036] N=1,373	-0.023 [0.040] N=1,171	-0.044 [0.041] N=962
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=310	0.025 [0.040] N=1,642	0.017 [0.043] N=1,526	0.001 [0.047] N=1,373	0.003 [0.052] N=1,171	0.005 [0.060] N=962
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=144	0.029 [0.028] N=1,642	0.030 [0.029] N=1,526	0.031 [0.033] N=1,373	0.042 [0.035] N=1,171	0.062 [0.041] N=962
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=575	-0.036 [0.043] N=1,588	-0.042 [0.044] N=1,476	-0.035 [0.047] N=1,328	-0.043 [0.050] N=1,133	-0.019 [0.057] N=932
BMI-for-age z score below -2	0.132 P=215	-0.032 [0.037] N=1,609	-0.030 [0.038] N=1,496	-0.026 [0.041] N=1,344	0.002 [0.046] N=1,148	-0.006 [0.050] N=946

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.187 [0.307] N=1,407	0.151 [0.351] N=1,285	0.306 [0.422] N=1,130	0.490 [0.452] N=927	0.161 [0.589] N=718
WHO weight-for-age z score	-0.245 [0.177] N=1,604	-0.218 [0.200] N=1,459	-0.249 [0.224] N=1,284	-0.275 [0.281] N=1,052	**-0.689* [0.347] N=814**
WHO weight-for-length/height z score, 0-60 months	0.076 [0.258] N=1,385	0.009 [0.286] N=1,269	-0.165 [0.338] N=1,116	-0.311 [0.385] N=914	-0.311 [0.464] N=705
WHO BMI-for-age z score	0.278 [0.268] N=1,358	0.277 [0.301] N=1,245	0.110 [0.357] N=1,093	-0.202 [0.412] N=897	-0.128 [0.481] N=695
Stunted: HAZ below -2	-0.064 [0.055] N=1,407	-0.085 [0.062] N=1,285	-0.084 [0.070] N=1,130	-0.132 [0.080] N=927	-0.122 [0.107] N=718
Underweight: WAZ below -2	0.041 [0.033] N=1,604	0.031 [0.037] N=1,459	0.010 [0.042] N=1,284	0.012 [0.053] N=1,052	0.095 [0.070] N=814
Wasted: WLZ/WHZ below -2, 0-60 months	-0.039 [0.047] N=1,385	-0.033 [0.049] N=1,269	-0.038 [0.056] N=1,116	-0.020 [0.070] N=914	-0.077 [0.081] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.008 [0.048] N=1,385	-0.008 [0.054] N=1,269	-0.053 [0.059] N=1,116	-0.058 [0.074] N=914	-0.114 [0.089] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.003 [0.035] N=1,385	0.002 [0.039] N=1,269	-0.006 [0.045] N=1,116	0.014 [0.053] N=914	0.049 [0.069] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.075 [0.061] N=1,341	-0.093 [0.066] N=1,229	-0.099 [0.074] N=1,081	**-0.150* [0.090] N=886**	-0.173 [0.124] N=685
BMI-for-age z score below -2	-0.052 [0.048] N=1,358	-0.052 [0.049] N=1,245	-0.049 [0.056] N=1,093	0.012 [0.069] N=897	-0.019 [0.083] N=695

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.613	-0.038 [0.212] N=1,666	-0.072 [0.228] N=1,544	-0.032 [0.252] N=1,389	-0.004 [0.255] N=1,186	-0.239 [0.272] N=977
WHO weight-for-age z score	-0.147	-0.155 [0.116] N=1,887	-0.125 [0.126] N=1,742	-0.118 [0.135] N=1,567	-0.093 [0.141] N=1,335	-0.208 [0.151] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.319	0.212 [0.198] N=1,642	0.191 [0.212] N=1,526	0.152 [0.233] N=1,373	0.132 [0.253] N=1,171	0.235 [0.287] N=962
WHO BMI-for-age z score	0.287	0.318 [0.221] N=1,609	0.329 [0.239] N=1,496	0.279 [0.263] N=1,344	0.184 [0.295] N=1,148	0.304 [0.335] N=946
Stunted: HAZ below -2	0.274 P=441	-0.000 [0.039] N=1,666	-0.004 [0.041] N=1,544	0.006 [0.044] N=1,389	-0.004 [0.046] N=1,186	0.038 [0.049] N=977
Underweight: WAZ below -2	0.093 P=171	**0.051** [0.022] N=1,887**	**0.047** [0.023] N=1,742**	0.035 [0.025] N=1,567	0.033 [0.027] N=1,335	**0.062** [0.030] N=1,097**
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=199	-0.031 [0.033] N=1,642	-0.025 [0.034] N=1,526	-0.028 [0.036] N=1,373	-0.019 [0.040] N=1,171	-0.039 [0.041] N=962
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=310	0.026 [0.041] N=1,642	0.021 [0.044] N=1,526	0.004 [0.047] N=1,373	0.006 [0.053] N=1,171	0.006 [0.061] N=962
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=144	0.030 [0.028] N=1,642	0.035 [0.030] N=1,526	0.034 [0.033] N=1,373	0.044 [0.035] N=1,171	0.062 [0.041] N=962
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=575	-0.021 [0.043] N=1,588	-0.020 [0.044] N=1,476	-0.019 [0.047] N=1,328	-0.034 [0.050] N=1,133	-0.003 [0.057] N=932
BMI-for-age z score below -2	0.132 P=215	-0.027 [0.037] N=1,609	-0.023 [0.039] N=1,496	-0.021 [0.042] N=1,344	0.005 [0.047] N=1,148	-0.004 [0.051] N=946

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.157 [0.312] N=1,407	0.092 [0.355] N=1,285	0.257 [0.422] N=1,130	0.476 [0.447] N=927	0.160 [0.592] N=718
WHO weight-for-age z score	-0.274 [0.171] N=1,604	-0.250 [0.195] N=1,459	-0.272 [0.218] N=1,284	-0.269 [0.277] N=1,052	**-0.687** [0.343] N=814**
WHO weight-for-length/height z score, 0-60 months	0.075 [0.261] N=1,385	0.021 [0.295] N=1,269	-0.140 [0.342] N=1,116	-0.303 [0.384] N=914	-0.323 [0.469] N=705
WHO BMI-for-age z score	0.284 [0.273] N=1,358	0.294 [0.312] N=1,245	0.138 [0.363] N=1,093	-0.198 [0.412] N=897	-0.117 [0.490] N=695
Stunted: HAZ below -2	-0.056 [0.056] N=1,407	-0.068 [0.062] N=1,285	-0.072 [0.070] N=1,130	**-0.131* [0.079] N=927**	-0.122 [0.106] N=718
Underweight: WAZ below -2	0.047 [0.032] N=1,604	0.040 [0.036] N=1,459	0.016 [0.041] N=1,284	0.009 [0.053] N=1,052	0.093 [0.071] N=814
Wasted: WLZ/WHZ below -2, 0-60 months	-0.035 [0.047] N=1,385	-0.027 [0.050] N=1,269	-0.035 [0.056] N=1,116	-0.017 [0.070] N=914	-0.071 [0.082] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.049] N=1,385	-0.003 [0.056] N=1,269	-0.049 [0.061] N=1,116	-0.057 [0.075] N=914	-0.112 [0.090] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.005 [0.035] N=1,385	0.008 [0.040] N=1,269	-0.003 [0.047] N=1,116	0.014 [0.054] N=914	0.046 [0.070] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.062 [0.061] N=1,341	-0.070 [0.065] N=1,229	-0.088 [0.073] N=1,081	**-0.153* [0.088] N=886**	-0.174 [0.120] N=685
BMI-for-age z score below -2	-0.048 [0.048] N=1,358	-0.044 [0.050] N=1,245	-0.045 [0.057] N=1,093	0.014 [0.070] N=897	-0.017 [0.085] N=695

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.613	-0.065 [0.212] N=1,666	-0.085 [0.230] N=1,544	-0.047 [0.254] N=1,389	-0.031 [0.259] N=1,186	-0.260 [0.275] N=977
WHO weight-for-age z score	-0.147	-0.169 [0.117] N=1,887	-0.139 [0.126] N=1,742	-0.131 [0.136] N=1,567	-0.119 [0.143] N=1,335	-0.243 [0.155] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.319	0.229 [0.197] N=1,642	0.203 [0.211] N=1,526	0.158 [0.234] N=1,373	0.134 [0.256] N=1,171	0.237 [0.287] N=962
WHO BMI-for-age z score	0.287	0.337 [0.221] N=1,609	0.345 [0.237] N=1,496	0.292 [0.262] N=1,344	0.193 [0.297] N=1,148	0.309 [0.334] N=946
Stunted: HAZ below -2	0.274 P=441	0.006 [0.039] N=1,666	0.001 [0.041] N=1,544	0.012 [0.044] N=1,389	0.006 [0.046] N=1,186	0.046 [0.049] N=977
Underweight: WAZ below -2	0.093 P=171	**0.054** [0.022] N=1,887**	**0.050** [0.023] N=1,742**	0.039 [0.025] N=1,567	0.039 [0.027] N=1,335	**0.068** [0.030] N=1,097**
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=199	-0.032 [0.033] N=1,642	-0.026 [0.034] N=1,526	-0.027 [0.036] N=1,373	-0.019 [0.040] N=1,171	-0.042 [0.041] N=962
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=310	0.028 [0.041] N=1,642	0.022 [0.044] N=1,526	0.006 [0.047] N=1,373	0.007 [0.053] N=1,171	0.005 [0.061] N=962
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=144	0.032 [0.028] N=1,642	0.034 [0.030] N=1,526	0.033 [0.033] N=1,373	0.043 [0.036] N=1,171	0.061 [0.041] N=962
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=575	-0.013 [0.043] N=1,588	-0.014 [0.045] N=1,476	-0.009 [0.048] N=1,328	-0.018 [0.051] N=1,133	0.009 [0.059] N=932
BMI-for-age z score below -2	0.132 P=215	-0.028 [0.037] N=1,609	-0.024 [0.038] N=1,496	-0.019 [0.042] N=1,344	0.008 [0.047] N=1,148	-0.004 [0.051] N=946

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.137 [0.312] N=1,407	0.089 [0.355] N=1,285	0.266 [0.427] N=1,130	0.493 [0.453] N=927	0.169 [0.587] N=718
WHO weight-for-age z score	**-0.294* [0.174] N=1,604**	-0.262 [0.196] N=1,459	-0.287 [0.220] N=1,284	-0.288 [0.279] N=1,052	**-0.690** [0.344] N=814**
WHO weight-for-length/height z score, 0-60 months	0.071 [0.259] N=1,385	0.016 [0.288] N=1,269	-0.154 [0.341] N=1,116	-0.298 [0.390] N=914	-0.289 [0.472] N=705
WHO BMI-for-age z score	0.279 [0.272] N=1,358	0.291 [0.305] N=1,245	0.128 [0.361] N=1,093	-0.184 [0.421] N=897	-0.089 [0.493] N=695
Stunted: HAZ below -2	-0.053 [0.056] N=1,407	-0.070 [0.062] N=1,285	-0.076 [0.071] N=1,130	**-0.136* [0.081] N=927**	-0.123 [0.108] N=718
Underweight: WAZ below -2	0.050 [0.033] N=1,604	0.040 [0.036] N=1,459	0.018 [0.042] N=1,284	0.012 [0.053] N=1,052	0.097 [0.071] N=814
Wasted: WLZ/WHZ below -2, 0-60 months	-0.033 [0.047] N=1,385	-0.028 [0.050] N=1,269	-0.035 [0.056] N=1,116	-0.016 [0.071] N=914	-0.072 [0.081] N=705
Overweight: WLZ/WHZ above +2, 0-60 months	0.012 [0.049] N=1,385	-0.003 [0.055] N=1,269	-0.049 [0.060] N=1,116	-0.053 [0.074] N=914	-0.099 [0.087] N=705
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.036] N=1,385	0.009 [0.040] N=1,269	-0.001 [0.047] N=1,116	0.017 [0.054] N=914	0.051 [0.069] N=705
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.055 [0.063] N=1,341	-0.072 [0.067] N=1,229	-0.086 [0.075] N=1,081	-0.147 [0.091] N=886	-0.162 [0.125] N=685
BMI-for-age z score below -2	-0.048 [0.048] N=1,358	-0.050 [0.049] N=1,245	-0.049 [0.056] N=1,093	0.010 [0.070] N=897	-0.021 [0.083] N=695

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.625	-0.084 [0.212] N=1,547	-0.102 [0.230] N=1,439	-0.058 [0.253] N=1,291	-0.058 [0.252] N=1,111	-0.272 [0.256] N=915
WHO weight-for-age z score	-0.142	-0.070 [0.121] N=1,744	-0.049 [0.128] N=1,615	-0.048 [0.139] N=1,449	-0.031 [0.142] N=1,242	-0.101 [0.161] N=1,023
WHO weight-for-length/height z score, 0-60 months	0.327	**0.354* [0.194] N=1,522**	0.318 [0.205] N=1,421	0.254 [0.230] N=1,276	0.251 [0.248] N=1,097	0.414 [0.279] N=901
WHO BMI-for-age z score	0.298	**0.444** [0.216] N=1,490**	**0.421* [0.228] N=1,391**	0.362 [0.254] N=1,248	0.310 [0.283] N=1,075	0.494 [0.315] N=886
Stunted: HAZ below -2	0.275 P=409	0.002 [0.042] N=1,547	-0.009 [0.044] N=1,439	0.001 [0.046] N=1,291	-0.002 [0.049] N=1,111	0.039 [0.053] N=915
Underweight: WAZ below -2	0.094 P=159	0.037 [0.024] N=1,744	0.033 [0.025] N=1,615	0.023 [0.027] N=1,449	0.022 [0.029] N=1,242	0.048 [0.034] N=1,023
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=183	-0.054 [0.033] N=1,522	-0.048 [0.033] N=1,421	-0.046 [0.035] N=1,276	-0.036 [0.040] N=1,097	-0.057 [0.041] N=901
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=286	0.038 [0.041] N=1,522	0.029 [0.043] N=1,421	0.011 [0.048] N=1,276	0.011 [0.054] N=1,097	0.024 [0.063] N=901
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=137	0.030 [0.030] N=1,522	0.034 [0.032] N=1,421	0.037 [0.035] N=1,276	0.048 [0.037] N=1,097	**0.075* [0.043] N=901**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=531	-0.038 [0.047] N=1,473	-0.045 [0.049] N=1,375	-0.040 [0.052] N=1,234	-0.051 [0.057] N=1,062	-0.024 [0.065] N=873
BMI-for-age z score below -2	0.130 P=195	-0.046 [0.037] N=1,490	-0.041 [0.038] N=1,391	-0.038 [0.041] N=1,248	-0.014 [0.046] N=1,075	-0.028 [0.050] N=886

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.148 [0.320] N=1,303	0.132 [0.367] N=1,195	0.326 [0.441] N=1,047	0.517 [0.477] N=867	0.221 [0.621] N=671
WHO weight-for-age z score	-0.233 [0.177] N=1,479	-0.232 [0.194] N=1,350	-0.282 [0.223] N=1,184	-0.323 [0.268] N=977	**-0.693* [0.359] N=758**
WHO weight-for-length/height z score, 0-60 months	0.115 [0.257] N=1,280	0.016 [0.282] N=1,179	-0.216 [0.334] N=1,034	-0.333 [0.385] N=855	-0.283 [0.473] N=659
WHO BMI-for-age z score	0.291 [0.275] N=1,254	0.220 [0.301] N=1,155	0.022 [0.356] N=1,012	-0.237 [0.413] N=839	-0.112 [0.491] N=650
Stunted: HAZ below -2	-0.061 [0.060] N=1,303	-0.091 [0.064] N=1,195	-0.098 [0.072] N=1,047	**-0.148* [0.085] N=867**	-0.133 [0.113] N=671
Underweight: WAZ below -2	0.036 [0.034] N=1,479	0.030 [0.038] N=1,350	0.010 [0.044] N=1,184	0.018 [0.053] N=977	0.103 [0.075] N=758
Wasted: WLZ/WHZ below -2, 0-60 months	-0.061 [0.047] N=1,280	-0.053 [0.049] N=1,179	-0.052 [0.055] N=1,034	-0.046 [0.070] N=855	-0.116 [0.080] N=659
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.048] N=1,280	-0.006 [0.053] N=1,179	-0.056 [0.060] N=1,034	-0.071 [0.074] N=855	-0.107 [0.093] N=659
Obese: WLZ/WHZ above +3, 0-60 months	-0.009 [0.038] N=1,280	-0.008 [0.043] N=1,179	-0.015 [0.050] N=1,034	0.003 [0.057] N=855	0.044 [0.073] N=659
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.084 [0.065] N=1,241	-0.110 [0.067] N=1,143	-0.124 [0.076] N=1,002	**-0.192** [0.092] N=830**	**-0.221* [0.126] N=641**
BMI-for-age z score below -2	-0.061 [0.047] N=1,254	-0.056 [0.047] N=1,155	-0.050 [0.055] N=1,012	-0.002 [0.069] N=839	-0.047 [0.083] N=650

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.625	-0.084 [0.214] N=1,547	-0.101 [0.231] N=1,439	-0.068 [0.256] N=1,291	-0.076 [0.258] N=1,111	-0.303 [0.269] N=915
WHO weight-for-age z score	-0.142	-0.069 [0.120] N=1,744	-0.047 [0.127] N=1,615	-0.053 [0.137] N=1,449	-0.043 [0.142] N=1,242	-0.123 [0.161] N=1,023
WHO weight-for-length/height z score, 0-60 months	0.327	**0.341* [0.191] N=1,522**	0.305 [0.202] N=1,421	0.239 [0.227] N=1,276	0.236 [0.249] N=1,097	0.389 [0.281] N=901
WHO BMI-for-age z score	0.298	**0.432** [0.213] N=1,490**	**0.408* [0.225] N=1,391**	0.351 [0.253] N=1,248	0.298 [0.285] N=1,075	0.471 [0.320] N=886
Stunted: HAZ below -2	0.275 P=409	0.004 [0.042] N=1,547	-0.007 [0.044] N=1,439	0.004 [0.046] N=1,291	0.001 [0.049] N=1,111	0.042 [0.053] N=915
Underweight: WAZ below -2	0.094 P=159	0.038 [0.023] N=1,744	0.034 [0.025] N=1,615	0.025 [0.026] N=1,449	0.025 [0.028] N=1,242	0.051 [0.033] N=1,023
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=183	-0.051 [0.033] N=1,522	-0.045 [0.033] N=1,421	-0.044 [0.035] N=1,276	-0.035 [0.040] N=1,097	-0.059 [0.042] N=901
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=286	0.038 [0.041] N=1,522	0.029 [0.043] N=1,421	0.011 [0.048] N=1,276	0.011 [0.054] N=1,097	0.022 [0.062] N=901
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=137	0.030 [0.030] N=1,522	0.034 [0.031] N=1,421	0.037 [0.035] N=1,276	0.050 [0.037] N=1,097	**0.074* [0.043] N=901**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.366 P=531	-0.033 [0.047] N=1,473	-0.041 [0.049] N=1,375	-0.037 [0.051] N=1,234	-0.048 [0.057] N=1,062	-0.026 [0.064] N=873
BMI-for-age z score below -2	0.130 P=195	-0.042 [0.037] N=1,490	-0.038 [0.038] N=1,391	-0.036 [0.041] N=1,248	-0.012 [0.047] N=1,075	-0.029 [0.051] N=886

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.178 [0.324] N=1,303	0.159 [0.371] N=1,195	0.347 [0.442] N=1,047	0.515 [0.486] N=867	0.215 [0.644] N=671
WHO weight-for-age z score	-0.202 [0.176] N=1,479	-0.196 [0.194] N=1,350	-0.258 [0.222] N=1,184	-0.305 [0.270] N=977	**-0.697* [0.364] N=758**
WHO weight-for-length/height z score, 0-60 months	0.134 [0.254] N=1,280	0.045 [0.280] N=1,179	-0.189 [0.335] N=1,034	-0.321 [0.390] N=855	-0.289 [0.483] N=659
WHO BMI-for-age z score	0.304 [0.269] N=1,254	0.233 [0.296] N=1,155	0.032 [0.354] N=1,012	-0.230 [0.416] N=839	-0.123 [0.500] N=650
Stunted: HAZ below -2	-0.060 [0.060] N=1,303	-0.088 [0.065] N=1,195	-0.097 [0.072] N=1,047	**-0.145* [0.085] N=867**	-0.140 [0.114] N=671
Underweight: WAZ below -2	0.033 [0.035] N=1,479	0.026 [0.039] N=1,350	0.007 [0.044] N=1,184	0.014 [0.054] N=977	0.098 [0.075] N=758
Wasted: WLZ/WHZ below -2, 0-60 months	-0.059 [0.047] N=1,280	-0.053 [0.049] N=1,179	-0.053 [0.055] N=1,034	-0.045 [0.070] N=855	-0.120 [0.080] N=659
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.048] N=1,280	-0.003 [0.053] N=1,179	-0.054 [0.061] N=1,034	-0.066 [0.075] N=855	-0.106 [0.094] N=659
Obese: WLZ/WHZ above +3, 0-60 months	-0.010 [0.038] N=1,280	-0.008 [0.043] N=1,179	-0.015 [0.050] N=1,034	0.006 [0.057] N=855	0.046 [0.073] N=659
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.079 [0.065] N=1,241	-0.104 [0.068] N=1,143	-0.121 [0.076] N=1,002	**-0.188** [0.092] N=830**	**-0.233* [0.126] N=641**
BMI-for-age z score below -2	-0.057 [0.048] N=1,254	-0.053 [0.047] N=1,155	-0.049 [0.055] N=1,012	-0.001 [0.070] N=839	-0.051 [0.083] N=650

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.546	-0.211 [0.261] N=1,474	-0.273 [0.267] N=1,331	-0.204 [0.284] N=1,160	-0.367 [0.316] N=986	-0.541 [0.371] N=799
WHO weight-for-age z score	-0.157	0.042 [0.141] N=1,664	0.071 [0.150] N=1,515	-0.012 [0.154] N=1,319	0.049 [0.162] N=1,117	0.022 [0.178] N=898
WHO weight-for-length/height z score, 0-60 months	0.336	0.089 [0.223] N=1,460	-0.001 [0.229] N=1,321	-0.104 [0.250] N=1,153	0.045 [0.274] N=979	-0.043 [0.312] N=793
WHO BMI-for-age z score	0.274	0.117 [0.216] N=1,428	0.042 [0.221] N=1,293	-0.059 [0.245] N=1,128	0.064 [0.280] N=957	0.024 [0.303] N=780
Stunted: HAZ below -2	0.266 P=390	0.060 [0.055] N=1,474	0.059 [0.055] N=1,331	0.041 [0.058] N=1,160	0.067 [0.063] N=986	0.091 [0.071] N=799
Underweight: WAZ below -2	0.096 P=156	-0.009 [0.030] N=1,664	-0.015 [0.032] N=1,515	0.000 [0.034] N=1,319	-0.009 [0.037] N=1,117	-0.027 [0.038] N=898
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=177	0.041 [0.037] N=1,460	0.050 [0.039] N=1,321	**0.075* [0.041] N=1,153**	0.031 [0.043] N=979	0.035 [0.050] N=793
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=285	0.068 [0.043] N=1,460	0.054 [0.045] N=1,321	0.033 [0.048] N=1,153	0.046 [0.053] N=979	0.035 [0.058] N=793
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=138	0.043 [0.035] N=1,460	0.028 [0.036] N=1,321	0.020 [0.039] N=1,153	0.015 [0.045] N=979	-0.024 [0.047] N=793
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=515	**0.107* [0.055] N=1,409**	**0.101* [0.056] N=1,271**	**0.107* [0.059] N=1,108**	0.086 [0.061] N=941	**0.120* [0.068] N=762**
BMI-for-age z score below -2	0.134 P=193	0.038 [0.038] N=1,428	0.045 [0.041] N=1,293	0.053 [0.046] N=1,128	0.020 [0.049] N=957	0.025 [0.055] N=780

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.081 [0.380] N=1,271	-0.171 [0.400] N=1,128	-0.022 [0.454] N=957	-0.399 [0.535] N=783	-0.821 [0.723] N=596
WHO weight-for-age z score	0.036 [0.207] N=1,438	0.060 [0.225] N=1,289	-0.107 [0.235] N=1,093	-0.014 [0.257] N=891	-0.004 [0.320] N=672
WHO weight-for-length/height z score, 0-60 months	0.089 [0.298] N=1,255	-0.082 [0.310] N=1,116	-0.267 [0.363] N=948	0.060 [0.402] N=774	-0.182 [0.504] N=588
WHO BMI-for-age z score	-0.096 [0.287] N=1,234	-0.237 [0.293] N=1,099	-0.431 [0.360] N=934	-0.213 [0.435] N=763	-0.400 [0.471] N=586
Stunted: HAZ below -2	0.061 [0.084] N=1,271	0.055 [0.088] N=1,128	0.021 [0.099] N=957	0.087 [0.118] N=783	0.173 [0.150] N=596
Underweight: WAZ below -2	-0.006 [0.046] N=1,438	-0.014 [0.049] N=1,289	0.012 [0.057] N=1,093	-0.007 [0.064] N=891	-0.059 [0.066] N=672
Wasted: WLZ/WHZ below -2, 0-60 months	**0.080* [0.046] N=1,255**	**0.098* [0.050] N=1,116**	**0.150*** [0.054] N=948**	0.059 [0.056] N=774	0.082 [0.073] N=588
Overweight: WLZ/WHZ above +2, 0-60 months	**0.126** [0.059] N=1,255**	**0.103* [0.061] N=1,116**	0.082 [0.067] N=948	**0.138* [0.077] N=774**	0.128 [0.094] N=588
Obese: WLZ/WHZ above +3, 0-60 months	**0.087** [0.043] N=1,255**	0.057 [0.043] N=1,116	0.047 [0.050] N=948	0.040 [0.064] N=774	-0.080 [0.068] N=588
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.141* [0.082] N=1,212**	0.133 [0.085] N=1,074	0.146 [0.096] N=911	0.110 [0.111] N=744	0.211 [0.135] N=565
BMI-for-age z score below -2	**0.082* [0.044] N=1,234**	**0.095* [0.049] N=1,099**	**0.111* [0.061] N=934**	0.035 [0.063] N=763	0.053 [0.076] N=586

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.546	-0.213 [0.260] N=1,474	-0.275 [0.266] N=1,331	-0.204 [0.283] N=1,160	-0.367 [0.316] N=986	-0.542 [0.370] N=799
WHO weight-for-age z score	-0.157	0.048 [0.140] N=1,664	0.075 [0.149] N=1,515	-0.008 [0.152] N=1,319	0.051 [0.161] N=1,117	0.024 [0.178] N=898
WHO weight-for-length/height z score, 0-60 months	0.336	0.095 [0.222] N=1,460	0.004 [0.228] N=1,321	-0.100 [0.249] N=1,153	0.046 [0.272] N=979	-0.039 [0.308] N=793
WHO BMI-for-age z score	0.274	0.120 [0.216] N=1,428	0.045 [0.221] N=1,293	-0.059 [0.244] N=1,128	0.064 [0.280] N=957	0.023 [0.303] N=780
Stunted: HAZ below -2	0.266 P=390	0.060 [0.055] N=1,474	0.057 [0.055] N=1,331	0.040 [0.058] N=1,160	0.067 [0.063] N=986	0.091 [0.071] N=799
Underweight: WAZ below -2	0.096 P=156	-0.010 [0.030] N=1,664	-0.016 [0.032] N=1,515	-0.001 [0.034] N=1,319	-0.010 [0.036] N=1,117	-0.028 [0.038] N=898
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=177	0.038 [0.037] N=1,460	0.047 [0.039] N=1,321	**0.072* [0.041] N=1,153**	0.030 [0.043] N=979	0.034 [0.050] N=793
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=285	0.067 [0.044] N=1,460	0.053 [0.045] N=1,321	0.032 [0.048] N=1,153	0.046 [0.054] N=979	0.035 [0.058] N=793
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=138	0.042 [0.035] N=1,460	0.027 [0.036] N=1,321	0.019 [0.039] N=1,153	0.015 [0.046] N=979	-0.025 [0.047] N=793
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=515	**0.103* [0.056] N=1,409**	**0.096* [0.057] N=1,271**	**0.103* [0.059] N=1,108**	0.086 [0.062] N=941	**0.120* [0.069] N=762**
BMI-for-age z score below -2	0.134 P=193	0.037 [0.038] N=1,428	0.043 [0.041] N=1,293	0.051 [0.046] N=1,128	0.020 [0.050] N=957	0.025 [0.055] N=780

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.086 [0.379] N=1,271	-0.176 [0.399] N=1,128	-0.018 [0.453] N=957	-0.406 [0.533] N=783	-0.830 [0.717] N=596
WHO weight-for-age z score	0.052 [0.202] N=1,438	0.072 [0.221] N=1,289	-0.093 [0.229] N=1,093	0.006 [0.249] N=891	0.035 [0.309] N=672
WHO weight-for-length/height z score, 0-60 months	0.106 [0.300] N=1,255	-0.069 [0.311] N=1,116	-0.255 [0.364] N=948	0.083 [0.406] N=774	-0.101 [0.520] N=588
WHO BMI-for-age z score	-0.077 [0.289] N=1,234	-0.217 [0.295] N=1,099	-0.418 [0.360] N=934	-0.192 [0.438] N=763	-0.328 [0.489] N=586
Stunted: HAZ below -2	0.059 [0.084] N=1,271	0.052 [0.088] N=1,128	0.017 [0.099] N=957	0.087 [0.118] N=783	0.174 [0.150] N=596
Underweight: WAZ below -2	-0.008 [0.045] N=1,438	-0.016 [0.049] N=1,289	0.009 [0.056] N=1,093	-0.009 [0.063] N=891	-0.065 [0.065] N=672
Wasted: WLZ/WHZ below -2, 0-60 months	0.072 [0.046] N=1,255	**0.089* [0.051] N=1,116**	**0.138** [0.054] N=948**	0.048 [0.057] N=774	0.059 [0.076] N=588
Overweight: WLZ/WHZ above +2, 0-60 months	**0.125** [0.059] N=1,255**	0.100 [0.061] N=1,116	0.077 [0.067] N=948	**0.132* [0.076] N=774**	0.125 [0.093] N=588
Obese: WLZ/WHZ above +3, 0-60 months	**0.085** [0.043] N=1,255**	0.054 [0.043] N=1,116	0.043 [0.050] N=948	0.035 [0.064] N=774	-0.086 [0.069] N=588
CIAF: any stunting, underweight, or wasting, 0-60 months	0.128 [0.082] N=1,212	0.117 [0.086] N=1,074	0.124 [0.096] N=911	0.094 [0.111] N=744	0.183 [0.136] N=565
BMI-for-age z score below -2	**0.075* [0.044] N=1,234**	**0.086* [0.049] N=1,099**	**0.102* [0.061] N=934**	0.027 [0.064] N=763	0.040 [0.079] N=586

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.546	-0.216 [0.257] N=1,474	-0.271 [0.264] N=1,331	-0.195 [0.283] N=1,160	-0.357 [0.314] N=986	-0.541 [0.364] N=799
WHO weight-for-age z score	-0.157	0.051 [0.139] N=1,664	0.082 [0.148] N=1,515	0.014 [0.152] N=1,319	0.065 [0.160] N=1,117	0.032 [0.175] N=898
WHO weight-for-length/height z score, 0-60 months	0.336	0.102 [0.223] N=1,460	0.013 [0.229] N=1,321	-0.075 [0.249] N=1,153	0.056 [0.271] N=979	-0.025 [0.306] N=793
WHO BMI-for-age z score	0.274	0.125 [0.219] N=1,428	0.050 [0.223] N=1,293	-0.037 [0.245] N=1,128	0.073 [0.277] N=957	0.033 [0.299] N=780
Stunted: HAZ below -2	0.266 P=390	0.061 [0.054] N=1,474	0.056 [0.055] N=1,331	0.036 [0.057] N=1,160	0.062 [0.062] N=986	0.090 [0.069] N=799
Underweight: WAZ below -2	0.096 P=156	-0.011 [0.030] N=1,664	-0.017 [0.031] N=1,515	-0.004 [0.034] N=1,319	-0.012 [0.035] N=1,117	-0.028 [0.037] N=898
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=177	0.038 [0.037] N=1,460	0.045 [0.039] N=1,321	0.069 [0.042] N=1,153	0.028 [0.044] N=979	0.034 [0.051] N=793
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=285	0.068 [0.043] N=1,460	0.053 [0.045] N=1,321	0.033 [0.048] N=1,153	0.045 [0.054] N=979	0.037 [0.058] N=793
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=138	0.043 [0.035] N=1,460	0.028 [0.036] N=1,321	0.020 [0.039] N=1,153	0.014 [0.046] N=979	-0.023 [0.047] N=793
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=515	**0.106* [0.055] N=1,409**	**0.096* [0.056] N=1,271**	0.098 [0.060] N=1,108	0.080 [0.062] N=941	**0.120* [0.068] N=762**
BMI-for-age z score below -2	0.134 P=193	0.037 [0.039] N=1,428	0.043 [0.041] N=1,293	0.048 [0.047] N=1,128	0.018 [0.050] N=957	0.026 [0.056] N=780

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.084 [0.372] N=1,271	-0.170 [0.391] N=1,128	-0.000 [0.444] N=957	-0.384 [0.520] N=783	-0.809 [0.703] N=596
WHO weight-for-age z score	0.038 [0.203] N=1,438	0.058 [0.223] N=1,289	-0.068 [0.231] N=1,093	0.017 [0.252] N=891	0.032 [0.305] N=672
WHO weight-for-length/height z score, 0-60 months	0.077 [0.299] N=1,255	-0.096 [0.308] N=1,116	-0.253 [0.360] N=948	0.063 [0.401] N=774	-0.119 [0.514] N=588
WHO BMI-for-age z score	-0.106 [0.287] N=1,234	-0.243 [0.292] N=1,099	-0.421 [0.354] N=934	-0.224 [0.429] N=763	-0.349 [0.483] N=586
Stunted: HAZ below -2	0.058 [0.083] N=1,271	0.049 [0.086] N=1,128	0.011 [0.095] N=957	0.078 [0.114] N=783	0.168 [0.143] N=596
Underweight: WAZ below -2	-0.008 [0.045] N=1,438	-0.016 [0.049] N=1,289	0.003 [0.056] N=1,093	-0.015 [0.063] N=891	-0.068 [0.062] N=672
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.047] N=1,255	**0.090* [0.051] N=1,116**	**0.137** [0.055] N=948**	0.049 [0.057] N=774	0.062 [0.077] N=588
Overweight: WLZ/WHZ above +2, 0-60 months	**0.123** [0.059] N=1,255**	0.098 [0.061] N=1,116	0.076 [0.066] N=948	**0.127* [0.076] N=774**	0.123 [0.093] N=588
Obese: WLZ/WHZ above +3, 0-60 months	**0.082* [0.043] N=1,255**	0.051 [0.043] N=1,116	0.041 [0.050] N=948	0.030 [0.065] N=774	-0.090 [0.068] N=588
CIAF: any stunting, underweight, or wasting, 0-60 months	0.130 [0.082] N=1,212	0.116 [0.086] N=1,074	0.118 [0.096] N=911	0.088 [0.110] N=744	0.183 [0.132] N=565
BMI-for-age z score below -2	**0.075* [0.044] N=1,234**	**0.087* [0.049] N=1,099**	0.098 [0.060] N=934	0.024 [0.064] N=763	0.039 [0.080] N=586

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.546	-0.203 [0.254] N=1,474	-0.251 [0.260] N=1,331	-0.175 [0.279] N=1,160	-0.346 [0.307] N=986	-0.555 [0.362] N=799
WHO weight-for-age z score	-0.157	0.054 [0.141] N=1,664	0.080 [0.149] N=1,515	0.012 [0.153] N=1,319	0.064 [0.161] N=1,117	0.022 [0.177] N=898
WHO weight-for-length/height z score, 0-60 months	0.336	0.091 [0.222] N=1,460	-0.003 [0.228] N=1,321	-0.096 [0.248] N=1,153	0.055 [0.271] N=979	-0.018 [0.309] N=793
WHO BMI-for-age z score	0.274	0.107 [0.217] N=1,428	0.032 [0.222] N=1,293	-0.059 [0.243] N=1,128	0.067 [0.278] N=957	0.039 [0.304] N=780
Stunted: HAZ below -2	0.266 P=390	0.057 [0.053] N=1,474	0.052 [0.054] N=1,331	0.033 [0.057] N=1,160	0.062 [0.062] N=986	0.092 [0.070] N=799
Underweight: WAZ below -2	0.096 P=156	-0.012 [0.030] N=1,664	-0.018 [0.031] N=1,515	-0.005 [0.034] N=1,319	-0.013 [0.036] N=1,117	-0.030 [0.038] N=898
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=177	0.039 [0.037] N=1,460	0.048 [0.039] N=1,321	**0.074* [0.041] N=1,153**	0.031 [0.042] N=979	0.034 [0.049] N=793
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=285	0.066 [0.043] N=1,460	0.050 [0.045] N=1,321	0.032 [0.048] N=1,153	0.046 [0.053] N=979	0.038 [0.058] N=793
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=138	0.046 [0.034] N=1,460	0.031 [0.035] N=1,321	0.023 [0.038] N=1,153	0.019 [0.045] N=979	-0.019 [0.046] N=793
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=515	**0.103* [0.055] N=1,409**	**0.094* [0.056] N=1,271**	**0.098* [0.059] N=1,108**	0.083 [0.062] N=941	**0.124* [0.068] N=762**
BMI-for-age z score below -2	0.134 P=193	0.036 [0.038] N=1,428	0.042 [0.041] N=1,293	0.049 [0.046] N=1,128	0.018 [0.049] N=957	0.022 [0.054] N=780

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.073 [0.364] N=1,271	-0.137 [0.382] N=1,128	0.054 [0.435] N=957	-0.348 [0.508] N=783	-0.880 [0.683] N=596
WHO weight-for-age z score	0.070 [0.198] N=1,438	0.089 [0.219] N=1,289	-0.037 [0.226] N=1,093	0.041 [0.241] N=891	0.011 [0.283] N=672
WHO weight-for-length/height z score, 0-60 months	0.092 [0.297] N=1,255	-0.096 [0.307] N=1,116	-0.274 [0.356] N=948	0.063 [0.401] N=774	-0.085 [0.532] N=588
WHO BMI-for-age z score	-0.103 [0.287] N=1,234	-0.251 [0.292] N=1,099	-0.441 [0.354] N=934	-0.213 [0.431] N=763	-0.308 [0.501] N=586
Stunted: HAZ below -2	0.057 [0.081] N=1,271	0.044 [0.084] N=1,128	0.004 [0.095] N=957	0.077 [0.113] N=783	0.190 [0.142] N=596
Underweight: WAZ below -2	-0.011 [0.044] N=1,438	-0.019 [0.048] N=1,289	-0.000 [0.056] N=1,093	-0.017 [0.061] N=891	-0.065 [0.060] N=672
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.047] N=1,255	**0.094* [0.051] N=1,116**	**0.145*** [0.054] N=948**	0.054 [0.056] N=774	0.065 [0.076] N=588
Overweight: WLZ/WHZ above +2, 0-60 months	**0.122** [0.058] N=1,255**	0.095 [0.060] N=1,116	0.073 [0.066] N=948	0.125 [0.077] N=774	0.129 [0.095] N=588
Obese: WLZ/WHZ above +3, 0-60 months	**0.088** [0.043] N=1,255**	0.055 [0.043] N=1,116	0.043 [0.049] N=948	0.034 [0.064] N=774	-0.085 [0.070] N=588
CIAF: any stunting, underweight, or wasting, 0-60 months	0.128 [0.080] N=1,212	0.115 [0.085] N=1,074	0.116 [0.095] N=911	0.089 [0.109] N=744	0.210 [0.130] N=565
BMI-for-age z score below -2	**0.076* [0.045] N=1,234**	**0.088* [0.050] N=1,099**	**0.104* [0.061] N=934**	0.030 [0.063] N=763	0.045 [0.080] N=586

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.552	-0.240 [0.269] N=1,367	-0.305 [0.282] N=1,235	-0.218 [0.298] N=1,081	-0.388 [0.322] N=917	-0.568 [0.364] N=745
WHO weight-for-age z score	-0.160	0.069 [0.148] N=1,537	0.109 [0.159] N=1,399	-0.007 [0.163] N=1,223	0.030 [0.174] N=1,037	-0.035 [0.186] N=837
WHO weight-for-length/height z score, 0-60 months	0.335	0.149 [0.243] N=1,352	0.071 [0.253] N=1,224	-0.080 [0.269] N=1,071	0.023 [0.294] N=909	-0.102 [0.318] N=739
WHO BMI-for-age z score	0.271	0.133 [0.236] N=1,322	0.074 [0.246] N=1,198	-0.091 [0.267] N=1,049	-0.001 [0.303] N=890	-0.105 [0.321] N=728
Stunted: HAZ below -2	0.266 P=360	0.056 [0.057] N=1,367	0.056 [0.059] N=1,235	0.029 [0.061] N=1,081	0.051 [0.064] N=917	0.078 [0.072] N=745
Underweight: WAZ below -2	0.098 P=147	-0.016 [0.031] N=1,537	-0.022 [0.034] N=1,399	-0.007 [0.036] N=1,223	-0.010 [0.039] N=1,037	-0.022 [0.041] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=164	0.040 [0.038] N=1,352	0.046 [0.040] N=1,224	**0.075* [0.042] N=1,071**	0.037 [0.044] N=909	0.048 [0.050] N=739
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=263	0.064 [0.045] N=1,352	0.049 [0.047] N=1,224	0.020 [0.049] N=1,071	0.024 [0.054] N=909	0.007 [0.057] N=739
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=130	0.042 [0.035] N=1,352	0.026 [0.036] N=1,224	0.015 [0.038] N=1,071	0.011 [0.044] N=909	-0.026 [0.044] N=739
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=476	**0.102* [0.060] N=1,306**	0.091 [0.061] N=1,179	0.093 [0.063] N=1,031	0.072 [0.065] N=874	0.118 [0.072] N=710
BMI-for-age z score below -2	0.133 P=176	0.046 [0.041] N=1,322	0.050 [0.044] N=1,198	0.060 [0.048] N=1,049	0.039 [0.051] N=890	0.053 [0.057] N=728

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.072 [0.395] N=1,174	-0.159 [0.425] N=1,042	0.008 [0.486] N=888	-0.418 [0.554] N=724	-0.873 [0.712] N=552
WHO weight-for-age z score	0.116 [0.210] N=1,324	0.166 [0.230] N=1,186	-0.051 [0.239] N=1,010	0.034 [0.262] N=824	0.039 [0.319] N=624
WHO weight-for-length/height z score, 0-60 months	0.185 [0.338] N=1,158	0.039 [0.359] N=1,030	-0.222 [0.418] N=877	0.069 [0.459] N=715	-0.179 [0.592] N=545
WHO BMI-for-age z score	-0.042 [0.322] N=1,137	-0.152 [0.339] N=1,013	-0.445 [0.412] N=864	-0.241 [0.490] N=705	-0.505 [0.542] N=543
Stunted: HAZ below -2	0.053 [0.091] N=1,174	0.047 [0.095] N=1,042	-0.004 [0.105] N=888	0.047 [0.121] N=724	0.130 [0.149] N=552
Underweight: WAZ below -2	-0.021 [0.048] N=1,324	-0.033 [0.052] N=1,186	-0.009 [0.058] N=1,010	-0.020 [0.067] N=824	-0.079 [0.070] N=624
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.049] N=1,158	0.079 [0.054] N=1,030	**0.132** [0.059] N=877**	0.044 [0.061] N=715	0.063 [0.081] N=545
Overweight: WLZ/WHZ above +2, 0-60 months	**0.125* [0.064] N=1,158**	0.101 [0.067] N=1,030	0.067 [0.074] N=877	0.106 [0.085] N=715	0.080 [0.105] N=545
Obese: WLZ/WHZ above +3, 0-60 months	**0.080* [0.048] N=1,158**	0.047 [0.049] N=1,030	0.032 [0.055] N=877	0.025 [0.070] N=715	-0.101 [0.078] N=545
CIAF: any stunting, underweight, or wasting, 0-60 months	0.123 [0.093] N=1,119	0.103 [0.097] N=992	0.100 [0.107] N=844	0.044 [0.125] N=687	0.146 [0.150] N=523
BMI-for-age z score below -2	**0.087* [0.048] N=1,137**	**0.095* [0.054] N=1,013**	**0.112* [0.065] N=864**	0.051 [0.069] N=705	0.087 [0.084] N=543

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.552	-0.257 [0.271] N=1,367	-0.319 [0.285] N=1,235	-0.230 [0.300] N=1,081	-0.411 [0.325] N=917	-0.605 [0.373] N=745
WHO weight-for-age z score	-0.160	0.068 [0.148] N=1,537	0.106 [0.160] N=1,399	-0.006 [0.163] N=1,223	0.029 [0.173] N=1,037	-0.041 [0.187] N=837
WHO weight-for-length/height z score, 0-60 months	0.335	0.165 [0.244] N=1,352	0.080 [0.254] N=1,224	-0.067 [0.271] N=1,071	0.037 [0.291] N=909	-0.089 [0.321] N=739
WHO BMI-for-age z score	0.271	0.148 [0.238] N=1,322	0.083 [0.248] N=1,198	-0.077 [0.269] N=1,049	0.011 [0.302] N=890	-0.085 [0.323] N=728
Stunted: HAZ below -2	0.266 P=360	0.055 [0.057] N=1,367	0.053 [0.058] N=1,235	0.025 [0.061] N=1,081	0.048 [0.064] N=917	0.073 [0.072] N=745
Underweight: WAZ below -2	0.098 P=147	-0.017 [0.031] N=1,537	-0.024 [0.033] N=1,399	-0.009 [0.035] N=1,223	-0.012 [0.038] N=1,037	-0.024 [0.040] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	0.119 P=164	0.035 [0.038] N=1,352	0.041 [0.040] N=1,224	0.068 [0.042] N=1,071	0.030 [0.044] N=909	0.040 [0.050] N=739
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=263	0.065 [0.044] N=1,352	0.048 [0.046] N=1,224	0.019 [0.048] N=1,071	0.023 [0.053] N=909	0.008 [0.057] N=739
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=130	0.042 [0.035] N=1,352	0.025 [0.035] N=1,224	0.015 [0.038] N=1,071	0.010 [0.044] N=909	-0.027 [0.043] N=739
CIAF: any stunting, underweight, or wasting, 0-60 months	0.365 P=476	**0.097* [0.058] N=1,306**	0.084 [0.059] N=1,179	0.083 [0.061] N=1,031	0.060 [0.063] N=874	0.103 [0.070] N=710
BMI-for-age z score below -2	0.133 P=176	0.042 [0.041] N=1,322	0.046 [0.044] N=1,198	0.054 [0.048] N=1,049	0.032 [0.051] N=890	0.044 [0.057] N=728

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.096 [0.395] N=1,174	-0.173 [0.422] N=1,042	-0.003 [0.480] N=888	-0.476 [0.553] N=724	-0.997 [0.712] N=552
WHO weight-for-age z score	0.110 [0.211] N=1,324	0.157 [0.232] N=1,186	-0.056 [0.240] N=1,010	0.022 [0.262] N=824	-0.005 [0.322] N=624
WHO weight-for-length/height z score, 0-60 months	0.178 [0.334] N=1,158	0.018 [0.352] N=1,030	-0.245 [0.409] N=877	0.048 [0.445] N=715	-0.203 [0.564] N=545
WHO BMI-for-age z score	-0.040 [0.320] N=1,137	-0.164 [0.333] N=1,013	-0.453 [0.405] N=864	-0.241 [0.482] N=705	-0.465 [0.540] N=543
Stunted: HAZ below -2	0.055 [0.090] N=1,174	0.045 [0.094] N=1,042	-0.008 [0.104] N=888	0.052 [0.120] N=724	0.139 [0.147] N=552
Underweight: WAZ below -2	-0.021 [0.047] N=1,324	-0.033 [0.052] N=1,186	-0.010 [0.058] N=1,010	-0.020 [0.066] N=824	-0.070 [0.070] N=624
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.049] N=1,158	0.081 [0.053] N=1,030	**0.133** [0.058] N=877**	0.043 [0.059] N=715	0.063 [0.079] N=545
Overweight: WLZ/WHZ above +2, 0-60 months	**0.125* [0.063] N=1,158**	0.099 [0.065] N=1,030	0.062 [0.072] N=877	0.103 [0.083] N=715	0.086 [0.101] N=545
Obese: WLZ/WHZ above +3, 0-60 months	**0.079* [0.047] N=1,158**	0.045 [0.048] N=1,030	0.029 [0.053] N=877	0.023 [0.068] N=715	-0.099 [0.074] N=545
CIAF: any stunting, underweight, or wasting, 0-60 months	0.125 [0.091] N=1,119	0.102 [0.096] N=992	0.095 [0.105] N=844	0.041 [0.123] N=687	0.142 [0.147] N=523
BMI-for-age z score below -2	**0.087* [0.048] N=1,137**	**0.096* [0.054] N=1,013**	**0.113* [0.065] N=864**	0.050 [0.068] N=705	0.081 [0.084] N=543

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.