Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Oldest Child

Age reference: 12 November 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.323	-0.003 [0.207] N=1,705	0.029 [0.219] N=1,581	0.031 [0.245] N=1,420	0.027 [0.247] N=1,218	-0.218 [0.270] N=1,005
WHO weight-for-age z score	0.067	-0.106 [0.119] N=1,935	-0.045 [0.127] N=1,789	-0.054 [0.134] N=1,608	-0.081 [0.141] N=1,373	-0.192 [0.155] N=1,130
WHO weight-for-length/height z score, 0-60 months	0.333	0.197 [0.192] N=1,687	0.163 [0.201] N=1,570	0.126 [0.226] N=1,411	0.117 [0.246] N=1,208	0.195 [0.275] N=995
WHO BMI-for-age z score	0.294	0.239 [0.213] N=1,654	0.245 [0.226] N=1,541	0.188 [0.255] N=1,382	0.109 [0.284] N=1,184	0.240 [0.317] N=977
Stunted: HAZ below -2	0.227 P=367	-0.001 [0.038] N=1,705	-0.015 [0.041] N=1,581	-0.006 [0.043] N=1,420	-0.015 [0.046] N=1,218	0.018 [0.050] N=1,005
Underweight: WAZ below -2	0.076 P=144	0.024 [0.022] N=1,935	0.012 [0.023] N=1,789	0.006 [0.024] N=1,608	0.012 [0.027] N=1,373	0.032 [0.031] N=1,130
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=205	-0.035 [0.032] N=1,687	-0.027 [0.032] N=1,570	-0.031 [0.035] N=1,411	-0.021 [0.038] N=1,208	-0.035 [0.040] N=995
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=324	0.020 [0.040] N=1,687	0.012 [0.042] N=1,570	0.000 [0.047] N=1,411	0.005 [0.053] N=1,208	0.002 [0.061] N=995
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=153	0.021 [0.028] N=1,687	0.023 [0.030] N=1,570	0.023 [0.033] N=1,411	0.030 [0.037] N=1,208	0.046 [0.042] N=995
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=505	-0.037 [0.043] N=1,629	-0.047 [0.044] N=1,515	-0.041 [0.046] N=1,361	-0.047 [0.050] N=1,165	-0.029 [0.055] N=961
BMI-for-age z score below -2	0.129 P=218	-0.031 [0.036] N=1,654	-0.032 [0.036] N=1,541	-0.027 [0.040] N=1,382	-0.001 [0.044] N=1,184	-0.010 [0.047] N=977

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.133 [0.299] N=1,437	0.203 [0.327] N=1,313	0.301 [0.390] N=1,152	0.483 [0.414] N=950	0.000 [0.574] N=737
WHO weight-for-age z score	-0.237 [0.179] N=1,642	-0.152 [0.201] N=1,496	-0.189 [0.219] N=1,315	-0.273 [0.269] N=1,080	**-0.738** [0.355] N=837**
WHO weight-for-length/height z score, 0-60 months	0.103 [0.253] N=1,420	0.033 [0.274] N=1,303	-0.115 [0.323] N=1,144	-0.211 [0.368] N=941	-0.246 [0.448] N=728
WHO BMI-for-age z score	0.207 [0.258] N=1,393	0.208 [0.287] N=1,280	0.030 [0.345] N=1,121	-0.245 [0.389] N=923	-0.131 [0.460] N=716
Stunted: HAZ below -2	0.003 [0.056] N=1,437	-0.024 [0.062] N=1,313	-0.019 [0.069] N=1,152	-0.057 [0.079] N=950	0.003 [0.108] N=737
Underweight: WAZ below -2	0.035 [0.031] N=1,642	0.014 [0.033] N=1,496	0.004 [0.037] N=1,315	0.021 [0.045] N=1,080	0.094 [0.062] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	-0.038 [0.046] N=1,420	-0.025 [0.048] N=1,303	-0.032 [0.054] N=1,144	-0.017 [0.068] N=941	-0.056 [0.079] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.005 [0.047] N=1,420	-0.011 [0.052] N=1,303	-0.044 [0.058] N=1,144	-0.046 [0.072] N=941	-0.105 [0.085] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.003 [0.033] N=1,420	0.003 [0.037] N=1,303	-0.002 [0.042] N=1,144	0.007 [0.052] N=941	0.030 [0.065] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.019 [0.066] N=1,371	-0.042 [0.070] N=1,257	-0.033 [0.078] N=1,103	-0.066 [0.095] N=907	-0.028 [0.130] N=703
BMI-for-age z score below -2	-0.046 [0.047] N=1,393	-0.050 [0.049] N=1,280	-0.043 [0.056] N=1,121	0.009 [0.070] N=923	-0.010 [0.081] N=716

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.323	-0.002 [0.207] N=1,705	0.029 [0.219] N=1,581	0.031 [0.245] N=1,420	0.031 [0.248] N=1,218	-0.210 [0.270] N=1,005
WHO weight-for-age z score	0.067	-0.109 [0.119] N=1,935	-0.047 [0.127] N=1,789	-0.056 [0.134] N=1,608	-0.083 [0.141] N=1,373	-0.198 [0.154] N=1,130
WHO weight-for-length/height z score, 0-60 months	0.333	0.193 [0.192] N=1,687	0.159 [0.201] N=1,570	0.121 [0.226] N=1,411	0.111 [0.247] N=1,208	0.180 [0.277] N=995
WHO BMI-for-age z score	0.294	0.238 [0.213] N=1,654	0.242 [0.227] N=1,541	0.185 [0.256] N=1,382	0.103 [0.286] N=1,184	0.227 [0.319] N=977
Stunted: HAZ below -2	0.227 P=367	-0.001 [0.038] N=1,705	-0.014 [0.041] N=1,581	-0.006 [0.043] N=1,420	-0.014 [0.046] N=1,218	0.019 [0.051] N=1,005
Underweight: WAZ below -2	0.076 P=144	0.024 [0.022] N=1,935	0.012 [0.023] N=1,789	0.006 [0.024] N=1,608	0.012 [0.027] N=1,373	0.032 [0.031] N=1,130
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=205	-0.033 [0.032] N=1,687	-0.025 [0.032] N=1,570	-0.028 [0.035] N=1,411	-0.018 [0.038] N=1,208	-0.031 [0.041] N=995
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=324	0.020 [0.040] N=1,687	0.012 [0.042] N=1,570	0.001 [0.047] N=1,411	0.006 [0.053] N=1,208	0.002 [0.061] N=995
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=153	0.021 [0.028] N=1,687	0.023 [0.030] N=1,570	0.023 [0.033] N=1,411	0.030 [0.037] N=1,208	0.046 [0.042] N=995
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=505	-0.035 [0.043] N=1,629	-0.044 [0.043] N=1,515	-0.038 [0.046] N=1,361	-0.044 [0.050] N=1,165	-0.022 [0.056] N=961
BMI-for-age z score below -2	0.129 P=218	-0.031 [0.036] N=1,654	-0.031 [0.037] N=1,541	-0.026 [0.040] N=1,382	0.001 [0.044] N=1,184	-0.007 [0.048] N=977

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.127 [0.298] N=1,437	0.196 [0.326] N=1,313	0.294 [0.389] N=1,152	0.465 [0.411] N=950	-0.019 [0.567] N=737
WHO weight-for-age z score	-0.232 [0.178] N=1,642	-0.144 [0.201] N=1,496	-0.181 [0.219] N=1,315	-0.266 [0.271] N=1,080	**-0.726** [0.357] N=837**
WHO weight-for-length/height z score, 0-60 months	0.104 [0.250] N=1,420	0.034 [0.271] N=1,303	-0.112 [0.321] N=1,144	-0.211 [0.366] N=941	-0.252 [0.444] N=728
WHO BMI-for-age z score	0.211 [0.257] N=1,393	0.211 [0.285] N=1,280	0.037 [0.342] N=1,121	-0.241 [0.387] N=923	-0.132 [0.455] N=716
Stunted: HAZ below -2	0.004 [0.056] N=1,437	-0.025 [0.062] N=1,313	-0.019 [0.068] N=1,152	-0.057 [0.079] N=950	0.005 [0.107] N=737
Underweight: WAZ below -2	0.034 [0.031] N=1,642	0.014 [0.033] N=1,496	0.003 [0.037] N=1,315	0.021 [0.045] N=1,080	0.093 [0.062] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	-0.038 [0.046] N=1,420	-0.026 [0.048] N=1,303	-0.033 [0.054] N=1,144	-0.017 [0.068] N=941	-0.054 [0.079] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.005 [0.047] N=1,420	-0.011 [0.052] N=1,303	-0.045 [0.058] N=1,144	-0.046 [0.072] N=941	-0.105 [0.085] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.003 [0.033] N=1,420	0.003 [0.037] N=1,303	-0.002 [0.043] N=1,144	0.007 [0.052] N=941	0.030 [0.065] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.022 [0.065] N=1,371	-0.044 [0.070] N=1,257	-0.037 [0.077] N=1,103	-0.069 [0.095] N=907	-0.029 [0.130] N=703
BMI-for-age z score below -2	-0.046 [0.047] N=1,393	-0.051 [0.049] N=1,280	-0.045 [0.056] N=1,121	0.008 [0.070] N=923	-0.010 [0.082] N=716

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.323	-0.026 [0.208] N=1,705	-0.011 [0.219] N=1,581	-0.005 [0.245] N=1,420	0.008 [0.247] N=1,218	-0.253 [0.265] N=1,005
WHO weight-for-age z score	0.067	-0.151 [0.114] N=1,935	-0.090 [0.123] N=1,789	-0.098 [0.130] N=1,608	-0.110 [0.137] N=1,373	-0.235 [0.149] N=1,130
WHO weight-for-length/height z score, 0-60 months	0.333	0.176 [0.193] N=1,687	0.153 [0.205] N=1,570	0.122 [0.228] N=1,411	0.111 [0.247] N=1,208	0.168 [0.278] N=995
WHO BMI-for-age z score	0.294	0.232 [0.215] N=1,654	0.245 [0.230] N=1,541	0.191 [0.258] N=1,382	0.110 [0.286] N=1,184	0.229 [0.323] N=977
Stunted: HAZ below -2	0.227 P=367	0.006 [0.038] N=1,705	-0.002 [0.041] N=1,581	0.004 [0.042] N=1,420	-0.008 [0.045] N=1,218	0.030 [0.049] N=1,005
Underweight: WAZ below -2	0.076 P=144	0.030 [0.022] N=1,935	0.020 [0.022] N=1,789	0.014 [0.024] N=1,608	0.018 [0.026] N=1,373	0.040 [0.031] N=1,130
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=205	-0.027 [0.032] N=1,687	-0.018 [0.033] N=1,570	-0.022 [0.035] N=1,411	-0.013 [0.039] N=1,208	-0.026 [0.041] N=995
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=324	0.020 [0.040] N=1,687	0.015 [0.043] N=1,570	0.004 [0.047] N=1,411	0.008 [0.053] N=1,208	0.003 [0.061] N=995
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=153	0.023 [0.029] N=1,687	0.027 [0.030] N=1,570	0.025 [0.033] N=1,411	0.032 [0.037] N=1,208	0.045 [0.042] N=995
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=505	-0.018 [0.042] N=1,629	-0.021 [0.043] N=1,515	-0.019 [0.046] N=1,361	-0.031 [0.050] N=1,165	-0.004 [0.056] N=961
BMI-for-age z score below -2	0.129 P=218	-0.026 [0.036] N=1,654	-0.025 [0.037] N=1,541	-0.022 [0.040] N=1,382	0.003 [0.044] N=1,184	-0.006 [0.048] N=977

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.111 [0.301] N=1,437	0.150 [0.329] N=1,313	0.253 [0.391] N=1,152	0.458 [0.410] N=950	-0.030 [0.562] N=737
WHO weight-for-age z score	-0.265 [0.171] N=1,642	-0.181 [0.195] N=1,496	-0.212 [0.214] N=1,315	-0.266 [0.268] N=1,080	**-0.733** [0.355] N=837**
WHO weight-for-length/height z score, 0-60 months	0.097 [0.254] N=1,420	0.043 [0.280] N=1,303	-0.093 [0.327] N=1,144	-0.209 [0.367] N=941	-0.269 [0.449] N=728
WHO BMI-for-age z score	0.213 [0.262] N=1,393	0.227 [0.295] N=1,280	0.058 [0.349] N=1,121	-0.241 [0.389] N=923	-0.123 [0.466] N=716
Stunted: HAZ below -2	0.007 [0.056] N=1,437	-0.011 [0.062] N=1,313	-0.010 [0.068] N=1,152	-0.060 [0.077] N=950	0.002 [0.105] N=737
Underweight: WAZ below -2	0.039 [0.030] N=1,642	0.021 [0.033] N=1,496	0.009 [0.036] N=1,315	0.019 [0.044] N=1,080	0.091 [0.063] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	-0.033 [0.045] N=1,420	-0.019 [0.048] N=1,303	-0.029 [0.054] N=1,144	-0.013 [0.068] N=941	-0.047 [0.080] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.048] N=1,420	-0.006 [0.054] N=1,303	-0.041 [0.059] N=1,144	-0.045 [0.072] N=941	-0.100 [0.085] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.004 [0.034] N=1,420	0.007 [0.038] N=1,303	-0.001 [0.043] N=1,144	0.006 [0.053] N=941	0.027 [0.065] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.009 [0.064] N=1,371	-0.021 [0.069] N=1,257	-0.024 [0.076] N=1,103	-0.073 [0.093] N=907	-0.024 [0.126] N=703
BMI-for-age z score below -2	-0.042 [0.047] N=1,393	-0.043 [0.049] N=1,280	-0.040 [0.057] N=1,121	0.012 [0.070] N=923	-0.007 [0.083] N=716

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.323	-0.074 [0.209] N=1,705	-0.051 [0.222] N=1,581	-0.039 [0.250] N=1,420	-0.036 [0.251] N=1,218	-0.305 [0.270] N=1,005
WHO weight-for-age z score	0.067	-0.164 [0.115] N=1,935	-0.104 [0.124] N=1,789	-0.108 [0.132] N=1,608	-0.138 [0.140] N=1,373	**-0.272* [0.154] N=1,130**
WHO weight-for-length/height z score, 0-60 months	0.333	0.199 [0.194] N=1,687	0.172 [0.205] N=1,570	0.130 [0.229] N=1,411	0.113 [0.251] N=1,208	0.172 [0.280] N=995
WHO BMI-for-age z score	0.294	0.259 [0.216] N=1,654	0.271 [0.230] N=1,541	0.207 [0.258] N=1,382	0.119 [0.290] N=1,184	0.234 [0.324] N=977
Stunted: HAZ below -2	0.227 P=367	0.016 [0.039] N=1,705	0.008 [0.041] N=1,581	0.015 [0.043] N=1,420	0.007 [0.046] N=1,218	0.044 [0.050] N=1,005
Underweight: WAZ below -2	0.076 P=144	0.033 [0.022] N=1,935	0.023 [0.023] N=1,789	0.017 [0.024] N=1,608	0.023 [0.027] N=1,373	0.045 [0.032] N=1,130
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=205	-0.029 [0.032] N=1,687	-0.019 [0.033] N=1,570	-0.022 [0.035] N=1,411	-0.013 [0.039] N=1,208	-0.028 [0.041] N=995
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=324	0.025 [0.040] N=1,687	0.020 [0.043] N=1,570	0.006 [0.047] N=1,411	0.011 [0.054] N=1,208	0.002 [0.061] N=995
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=153	0.024 [0.028] N=1,687	0.026 [0.030] N=1,570	0.024 [0.033] N=1,411	0.030 [0.037] N=1,208	0.045 [0.042] N=995
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=505	-0.008 [0.044] N=1,629	-0.012 [0.045] N=1,515	-0.007 [0.048] N=1,361	-0.012 [0.052] N=1,165	0.012 [0.058] N=961
BMI-for-age z score below -2	0.129 P=218	-0.027 [0.036] N=1,654	-0.027 [0.037] N=1,541	-0.021 [0.040] N=1,382	0.005 [0.044] N=1,184	-0.006 [0.048] N=977

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.086 [0.300] N=1,437	0.141 [0.328] N=1,313	0.256 [0.392] N=1,152	0.464 [0.408] N=950	-0.046 [0.548] N=737
WHO weight-for-age z score	-0.279 [0.174] N=1,642	-0.192 [0.196] N=1,496	-0.224 [0.214] N=1,315	-0.284 [0.269] N=1,080	**-0.732** [0.355] N=837**
WHO weight-for-length/height z score, 0-60 months	0.098 [0.252] N=1,420	0.043 [0.274] N=1,303	-0.105 [0.325] N=1,144	-0.204 [0.374] N=941	-0.232 [0.456] N=728
WHO BMI-for-age z score	0.211 [0.260] N=1,393	0.228 [0.288] N=1,280	0.050 [0.347] N=1,121	-0.231 [0.398] N=923	-0.103 [0.470] N=716
Stunted: HAZ below -2	0.010 [0.057] N=1,437	-0.011 [0.063] N=1,313	-0.013 [0.069] N=1,152	-0.066 [0.080] N=950	-0.004 [0.108] N=737
Underweight: WAZ below -2	0.040 [0.031] N=1,642	0.021 [0.033] N=1,496	0.010 [0.037] N=1,315	0.021 [0.045] N=1,080	0.096 [0.062] N=837
Wasted: WLZ/WHZ below -2, 0-60 months	-0.031 [0.046] N=1,420	-0.020 [0.048] N=1,303	-0.028 [0.054] N=1,144	-0.012 [0.069] N=941	-0.048 [0.080] N=728
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.048] N=1,420	-0.005 [0.053] N=1,303	-0.040 [0.059] N=1,144	-0.041 [0.073] N=941	-0.091 [0.084] N=728
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.034] N=1,420	0.009 [0.038] N=1,303	0.001 [0.044] N=1,144	0.009 [0.053] N=941	0.033 [0.064] N=728
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.004 [0.066] N=1,371	-0.022 [0.071] N=1,257	-0.023 [0.079] N=1,103	-0.069 [0.097] N=907	-0.015 [0.130] N=703
BMI-for-age z score below -2	-0.042 [0.047] N=1,393	-0.048 [0.049] N=1,280	-0.043 [0.057] N=1,121	0.010 [0.070] N=923	-0.012 [0.081] N=716

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.333	-0.042 [0.206] N=1,580	-0.033 [0.224] N=1,471	-0.028 [0.250] N=1,317	-0.053 [0.248] N=1,138	-0.323 [0.264] N=938
WHO weight-for-age z score	0.070	-0.052 [0.120] N=1,790	0.004 [0.126] N=1,660	-0.004 [0.133] N=1,488	-0.024 [0.138] N=1,278	-0.099 [0.162] N=1,054
WHO weight-for-length/height z score, 0-60 months	0.344	**0.331* [0.193] N=1,564**	0.295 [0.201] N=1,462	0.246 [0.226] N=1,311	0.254 [0.242] N=1,131	0.365 [0.275] N=931
WHO BMI-for-age z score	0.309	**0.367* [0.214] N=1,532**	0.353 [0.225] N=1,433	0.301 [0.252] N=1,283	0.265 [0.275] N=1,108	0.442 [0.308] N=914
Stunted: HAZ below -2	0.229 P=342	0.013 [0.041] N=1,580	0.000 [0.044] N=1,471	0.007 [0.046] N=1,317	-0.000 [0.049] N=1,138	0.041 [0.053] N=938
Underweight: WAZ below -2	0.076 P=133	0.018 [0.023] N=1,790	0.007 [0.024] N=1,660	-0.000 [0.026] N=1,488	0.007 [0.029] N=1,278	0.024 [0.036] N=1,054
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=189	-0.053 [0.032] N=1,564	-0.043 [0.032] N=1,462	-0.045 [0.034] N=1,311	-0.035 [0.038] N=1,131	-0.049 [0.039] N=931
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=300	0.034 [0.041] N=1,564	0.026 [0.043] N=1,462	0.013 [0.048] N=1,311	0.017 [0.055] N=1,131	0.022 [0.065] N=931
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=146	0.023 [0.030] N=1,564	0.027 [0.032] N=1,462	0.027 [0.035] N=1,311	0.036 [0.039] N=1,131	0.058 [0.044] N=931
CIAF: any stunting, underweight, or wasting, 0-60 months	0.321 P=469	-0.036 [0.046] N=1,509	-0.044 [0.047] N=1,410	-0.042 [0.050] N=1,263	-0.052 [0.055] N=1,090	-0.030 [0.061] N=898
BMI-for-age z score below -2	0.127 P=198	-0.040 [0.036] N=1,532	-0.039 [0.037] N=1,433	-0.036 [0.040] N=1,283	-0.014 [0.044] N=1,108	-0.030 [0.047] N=914

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.168 [0.295] N=1,330	0.224 [0.338] N=1,221	0.358 [0.402] N=1,067	0.531 [0.426] N=888	0.044 [0.609] N=688
WHO weight-for-age z score	-0.223 [0.177] N=1,517	-0.157 [0.194] N=1,387	-0.211 [0.215] N=1,215	-0.294 [0.257] N=1,005	**-0.704* [0.364] N=781**
WHO weight-for-length/height z score, 0-60 months	0.137 [0.253] N=1,315	0.037 [0.269] N=1,213	-0.153 [0.316] N=1,062	-0.208 [0.364] N=882	-0.182 [0.444] N=682
WHO BMI-for-age z score	0.219 [0.264] N=1,289	0.156 [0.285] N=1,190	-0.032 [0.337] N=1,040	-0.240 [0.380] N=865	-0.090 [0.459] N=671
Stunted: HAZ below -2	0.003 [0.060] N=1,330	-0.029 [0.066] N=1,221	-0.031 [0.072] N=1,067	-0.078 [0.085] N=888	-0.010 [0.118] N=688
Underweight: WAZ below -2	0.030 [0.032] N=1,517	0.013 [0.034] N=1,387	0.004 [0.038] N=1,215	0.026 [0.045] N=1,005	0.099 [0.068] N=781
Wasted: WLZ/WHZ below -2, 0-60 months	-0.056 [0.046] N=1,315	-0.041 [0.047] N=1,213	-0.046 [0.052] N=1,062	-0.039 [0.067] N=882	-0.092 [0.076] N=682
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.048] N=1,315	-0.008 [0.052] N=1,213	-0.046 [0.059] N=1,062	-0.052 [0.074] N=882	-0.093 [0.090] N=682
Obese: WLZ/WHZ above +3, 0-60 months	-0.007 [0.037] N=1,315	-0.005 [0.041] N=1,213	-0.011 [0.047] N=1,062	-0.002 [0.055] N=882	0.027 [0.069] N=682
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.033 [0.069] N=1,269	-0.056 [0.071] N=1,170	-0.059 [0.080] N=1,023	-0.117 [0.095] N=850	-0.085 [0.135] N=658
BMI-for-age z score below -2	-0.040 [0.045] N=1,289	-0.039 [0.046] N=1,190	-0.033 [0.053] N=1,040	0.012 [0.069] N=865	-0.021 [0.078] N=671

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.333	-0.044 [0.206] N=1,580	-0.034 [0.223] N=1,471	-0.036 [0.249] N=1,317	-0.065 [0.249] N=1,138	-0.341 [0.269] N=938
WHO weight-for-age z score	0.070	-0.052 [0.119] N=1,790	0.005 [0.125] N=1,660	-0.010 [0.132] N=1,488	-0.037 [0.138] N=1,278	-0.121 [0.162] N=1,054
WHO weight-for-length/height z score, 0-60 months	0.344	**0.319* [0.190] N=1,564**	0.282 [0.198] N=1,462	0.231 [0.224] N=1,311	0.238 [0.244] N=1,131	0.343 [0.279] N=931
WHO BMI-for-age z score	0.309	**0.358* [0.211] N=1,532**	0.341 [0.222] N=1,433	0.288 [0.250] N=1,283	0.249 [0.277] N=1,108	0.416 [0.313] N=914
Stunted: HAZ below -2	0.229 P=342	0.014 [0.041] N=1,580	0.001 [0.044] N=1,471	0.009 [0.046] N=1,317	0.002 [0.049] N=1,138	0.044 [0.054] N=938
Underweight: WAZ below -2	0.076 P=133	0.018 [0.023] N=1,790	0.007 [0.024] N=1,660	0.001 [0.026] N=1,488	0.009 [0.029] N=1,278	0.026 [0.035] N=1,054
Wasted: WLZ/WHZ below -2, 0-60 months	0.118 P=189	-0.050 [0.032] N=1,564	-0.040 [0.032] N=1,462	-0.043 [0.034] N=1,311	-0.034 [0.038] N=1,131	-0.052 [0.041] N=931
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=300	0.034 [0.041] N=1,564	0.025 [0.043] N=1,462	0.013 [0.048] N=1,311	0.016 [0.055] N=1,131	0.021 [0.064] N=931
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=146	0.023 [0.030] N=1,564	0.027 [0.032] N=1,462	0.028 [0.035] N=1,311	0.037 [0.039] N=1,131	0.057 [0.044] N=931
CIAF: any stunting, underweight, or wasting, 0-60 months	0.321 P=469	-0.032 [0.047] N=1,509	-0.040 [0.047] N=1,410	-0.038 [0.050] N=1,263	-0.049 [0.055] N=1,090	-0.026 [0.061] N=898
BMI-for-age z score below -2	0.127 P=198	-0.036 [0.036] N=1,532	-0.035 [0.036] N=1,433	-0.034 [0.040] N=1,283	-0.013 [0.045] N=1,108	-0.030 [0.049] N=914

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.189 [0.299] N=1,330	0.237 [0.341] N=1,221	0.361 [0.403] N=1,067	0.513 [0.436] N=888	0.025 [0.629] N=688
WHO weight-for-age z score	-0.193 [0.177] N=1,517	-0.125 [0.195] N=1,387	-0.193 [0.214] N=1,215	-0.290 [0.259] N=1,005	**-0.719* [0.370] N=781**
WHO weight-for-length/height z score, 0-60 months	0.157 [0.250] N=1,315	0.066 [0.268] N=1,213	-0.132 [0.317] N=1,062	-0.209 [0.370] N=882	-0.207 [0.456] N=682
WHO BMI-for-age z score	0.242 [0.261] N=1,289	0.179 [0.283] N=1,190	-0.018 [0.337] N=1,040	-0.238 [0.385] N=865	-0.109 [0.469] N=671
Stunted: HAZ below -2	0.001 [0.060] N=1,330	-0.030 [0.066] N=1,221	-0.032 [0.071] N=1,067	-0.079 [0.084] N=888	-0.016 [0.116] N=688
Underweight: WAZ below -2	0.028 [0.032] N=1,517	0.009 [0.034] N=1,387	0.001 [0.039] N=1,215	0.024 [0.045] N=1,005	0.094 [0.068] N=781
Wasted: WLZ/WHZ below -2, 0-60 months	-0.055 [0.046] N=1,315	-0.041 [0.047] N=1,213	-0.046 [0.053] N=1,062	-0.038 [0.068] N=882	-0.093 [0.076] N=682
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.048] N=1,315	-0.006 [0.052] N=1,213	-0.044 [0.060] N=1,062	-0.049 [0.075] N=882	-0.092 [0.091] N=682
Obese: WLZ/WHZ above +3, 0-60 months	-0.008 [0.036] N=1,315	-0.006 [0.041] N=1,213	-0.011 [0.047] N=1,062	-0.000 [0.056] N=882	0.027 [0.069] N=682
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.032 [0.068] N=1,269	-0.055 [0.072] N=1,170	-0.059 [0.080] N=1,023	-0.115 [0.096] N=850	-0.094 [0.134] N=658
BMI-for-age z score below -2	-0.038 [0.045] N=1,289	-0.037 [0.046] N=1,190	-0.032 [0.054] N=1,040	0.014 [0.069] N=865	-0.022 [0.078] N=671

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.197 [0.271] N=1,513	-0.249 [0.274] N=1,370	-0.175 [0.287] N=1,196	-0.307 [0.321] N=1,018	-0.431 [0.364] N=826
WHO weight-for-age z score	0.071	0.074 [0.135] N=1,710	0.096 [0.143] N=1,560	0.007 [0.147] N=1,361	0.078 [0.156] N=1,155	0.093 [0.176] N=929
WHO weight-for-length/height z score, 0-60 months	0.349	0.142 [0.217] N=1,501	0.083 [0.224] N=1,361	-0.005 [0.242] N=1,190	0.140 [0.261] N=1,011	0.063 [0.297] N=819
WHO BMI-for-age z score	0.279	0.109 [0.219] N=1,468	0.046 [0.226] N=1,331	-0.055 [0.252] N=1,164	0.099 [0.283] N=990	0.096 [0.313] N=806
Stunted: HAZ below -2	0.220 P=328	0.050 [0.047] N=1,513	0.045 [0.048] N=1,370	0.030 [0.050] N=1,196	0.044 [0.056] N=1,018	0.052 [0.064] N=826
Underweight: WAZ below -2	0.069 P=119	0.006 [0.026] N=1,710	0.002 [0.027] N=1,560	0.003 [0.029] N=1,361	0.004 [0.030] N=1,155	-0.011 [0.030] N=929
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=178	0.029 [0.036] N=1,501	0.033 [0.038] N=1,361	0.055 [0.042] N=1,190	0.007 [0.042] N=1,011	0.012 [0.046] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	**0.071* [0.042] N=1,501**	0.057 [0.043] N=1,361	0.041 [0.046] N=1,190	0.046 [0.050] N=1,011	0.045 [0.055] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=145	0.043 [0.035] N=1,501	0.032 [0.035] N=1,361	0.027 [0.039] N=1,190	0.024 [0.044] N=1,011	-0.009 [0.046] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=448	0.083 [0.051] N=1,448	0.073 [0.052] N=1,311	0.087 [0.055] N=1,144	0.055 [0.056] N=973	0.083 [0.064] N=789
BMI-for-age z score below -2	0.128 P=190	0.056 [0.041] N=1,468	0.062 [0.044] N=1,331	0.069 [0.050] N=1,164	0.031 [0.050] N=990	0.034 [0.055] N=806

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.051 [0.395] N=1,305	-0.114 [0.413] N=1,162	0.029 [0.456] N=988	-0.267 [0.543] N=810	-0.514 [0.706] N=618
WHO weight-for-age z score	0.039 [0.201] N=1,479	0.071 [0.217] N=1,329	-0.110 [0.231] N=1,130	-0.007 [0.251] N=924	0.032 [0.317] N=698
WHO weight-for-length/height z score, 0-60 months	0.075 [0.289] N=1,293	-0.046 [0.302] N=1,153	-0.233 [0.354] N=982	0.041 [0.383] N=803	-0.272 [0.456] N=611
WHO BMI-for-age z score	-0.171 [0.274] N=1,270	-0.302 [0.281] N=1,133	-0.514 [0.342] N=966	-0.269 [0.411] N=792	-0.439 [0.431] N=608
Stunted: HAZ below -2	0.061 [0.073] N=1,305	0.048 [0.075] N=1,162	0.026 [0.082] N=988	0.062 [0.101] N=810	0.087 [0.127] N=618
Underweight: WAZ below -2	0.030 [0.040] N=1,479	0.025 [0.043] N=1,329	0.034 [0.049] N=1,130	0.046 [0.054] N=924	0.010 [0.056] N=698
Wasted: WLZ/WHZ below -2, 0-60 months	**0.081* [0.045] N=1,293**	**0.094* [0.049] N=1,153**	**0.149*** [0.055] N=982**	0.053 [0.055] N=803	0.086 [0.065] N=611
Overweight: WLZ/WHZ above +2, 0-60 months	**0.124** [0.058] N=1,293**	**0.099* [0.059] N=1,153**	0.084 [0.066] N=982	0.113 [0.075] N=803	0.104 [0.088] N=611
Obese: WLZ/WHZ above +3, 0-60 months	**0.083** [0.042] N=1,293**	0.060 [0.042] N=1,153	0.055 [0.048] N=982	0.048 [0.061] N=803	-0.072 [0.069] N=611
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.132* [0.073] N=1,247**	0.114 [0.078] N=1,110	**0.154* [0.086] N=943**	0.092 [0.094] N=772	0.163 [0.120] N=588
BMI-for-age z score below -2	**0.123*** [0.042] N=1,270**	**0.135*** [0.046] N=1,133**	**0.165*** [0.057] N=966**	0.087 [0.055] N=792	0.101 [0.062] N=608

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.206 [0.269] N=1,513	-0.257 [0.273] N=1,370	-0.181 [0.286] N=1,196	-0.312 [0.319] N=1,018	-0.433 [0.362] N=826
WHO weight-for-age z score	0.071	0.078 [0.135] N=1,710	0.099 [0.142] N=1,560	0.010 [0.147] N=1,361	0.079 [0.156] N=1,155	0.091 [0.176] N=929
WHO weight-for-length/height z score, 0-60 months	0.349	0.148 [0.217] N=1,501	0.087 [0.223] N=1,361	-0.002 [0.241] N=1,190	0.140 [0.260] N=1,011	0.061 [0.295] N=819
WHO BMI-for-age z score	0.279	0.113 [0.219] N=1,468	0.048 [0.226] N=1,331	-0.054 [0.251] N=1,164	0.099 [0.283] N=990	0.094 [0.314] N=806
Stunted: HAZ below -2	0.220 P=328	0.050 [0.047] N=1,513	0.044 [0.048] N=1,370	0.030 [0.050] N=1,196	0.044 [0.056] N=1,018	0.052 [0.064] N=826
Underweight: WAZ below -2	0.069 P=119	0.005 [0.026] N=1,710	0.002 [0.027] N=1,560	0.002 [0.029] N=1,361	0.004 [0.030] N=1,155	-0.011 [0.031] N=929
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=178	0.026 [0.036] N=1,501	0.031 [0.039] N=1,361	0.053 [0.042] N=1,190	0.007 [0.042] N=1,011	0.013 [0.046] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	**0.071* [0.042] N=1,501**	0.056 [0.043] N=1,361	0.040 [0.046] N=1,190	0.046 [0.051] N=1,011	0.045 [0.056] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=145	0.042 [0.035] N=1,501	0.031 [0.035] N=1,361	0.025 [0.039] N=1,190	0.023 [0.045] N=1,011	-0.009 [0.046] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=448	0.078 [0.052] N=1,448	0.068 [0.053] N=1,311	0.083 [0.056] N=1,144	0.054 [0.057] N=973	0.084 [0.065] N=789
BMI-for-age z score below -2	0.128 P=190	0.053 [0.041] N=1,468	0.059 [0.044] N=1,331	0.067 [0.050] N=1,164	0.031 [0.050] N=990	0.034 [0.055] N=806

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.065 [0.393] N=1,305	-0.127 [0.411] N=1,162	0.022 [0.453] N=988	-0.289 [0.540] N=810	-0.536 [0.705] N=618
WHO weight-for-age z score	0.050 [0.198] N=1,479	0.079 [0.214] N=1,329	-0.097 [0.227] N=1,130	0.012 [0.245] N=924	0.060 [0.310] N=698
WHO weight-for-length/height z score, 0-60 months	0.092 [0.292] N=1,293	-0.035 [0.305] N=1,153	-0.220 [0.356] N=982	0.061 [0.387] N=803	-0.217 [0.472] N=611
WHO BMI-for-age z score	-0.150 [0.276] N=1,270	-0.284 [0.283] N=1,133	-0.500 [0.342] N=966	-0.245 [0.412] N=792	-0.384 [0.449] N=608
Stunted: HAZ below -2	0.061 [0.073] N=1,305	0.045 [0.075] N=1,162	0.022 [0.082] N=988	0.063 [0.101] N=810	0.091 [0.128] N=618
Underweight: WAZ below -2	0.028 [0.041] N=1,479	0.024 [0.043] N=1,329	0.032 [0.049] N=1,130	0.042 [0.054] N=924	0.006 [0.057] N=698
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.045] N=1,293	**0.085* [0.050] N=1,153**	**0.137** [0.056] N=982**	0.042 [0.056] N=803	0.068 [0.068] N=611
Overweight: WLZ/WHZ above +2, 0-60 months	**0.122** [0.058] N=1,293**	0.096 [0.059] N=1,153	0.078 [0.066] N=982	0.107 [0.075] N=803	0.100 [0.088] N=611
Obese: WLZ/WHZ above +3, 0-60 months	**0.081* [0.042] N=1,293**	0.056 [0.041] N=1,153	0.051 [0.048] N=982	0.043 [0.061] N=803	-0.078 [0.069] N=611
CIAF: any stunting, underweight, or wasting, 0-60 months	0.118 [0.074] N=1,247	0.097 [0.079] N=1,110	0.132 [0.089] N=943	0.076 [0.097] N=772	0.143 [0.123] N=588
BMI-for-age z score below -2	**0.115*** [0.042] N=1,270**	**0.126*** [0.047] N=1,133**	**0.154*** [0.057] N=966**	0.077 [0.056] N=792	0.088 [0.066] N=608

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.208 [0.264] N=1,513	-0.253 [0.268] N=1,370	-0.176 [0.283] N=1,196	-0.303 [0.315] N=1,018	-0.432 [0.354] N=826
WHO weight-for-age z score	0.071	0.081 [0.135] N=1,710	0.105 [0.142] N=1,560	0.029 [0.147] N=1,361	0.092 [0.157] N=1,155	0.099 [0.177] N=929
WHO weight-for-length/height z score, 0-60 months	0.349	0.155 [0.218] N=1,501	0.096 [0.224] N=1,361	0.020 [0.242] N=1,190	0.148 [0.259] N=1,011	0.069 [0.293] N=819
WHO BMI-for-age z score	0.279	0.119 [0.221] N=1,468	0.055 [0.228] N=1,331	-0.032 [0.251] N=1,164	0.106 [0.280] N=990	0.101 [0.310] N=806
Stunted: HAZ below -2	0.220 P=328	0.051 [0.046] N=1,513	0.043 [0.047] N=1,370	0.028 [0.049] N=1,196	0.041 [0.054] N=1,018	0.051 [0.062] N=826
Underweight: WAZ below -2	0.069 P=119	0.005 [0.026] N=1,710	0.001 [0.027] N=1,560	0.000 [0.029] N=1,361	0.003 [0.030] N=1,155	-0.010 [0.030] N=929
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=178	0.025 [0.037] N=1,501	0.028 [0.039] N=1,361	0.050 [0.043] N=1,190	0.005 [0.043] N=1,011	0.013 [0.048] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	**0.072* [0.043] N=1,501**	0.057 [0.044] N=1,361	0.042 [0.046] N=1,190	0.045 [0.051] N=1,011	0.047 [0.056] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=145	0.043 [0.035] N=1,501	0.031 [0.036] N=1,361	0.026 [0.039] N=1,190	0.024 [0.045] N=1,011	-0.008 [0.047] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=448	0.081 [0.051] N=1,448	0.067 [0.053] N=1,311	0.079 [0.057] N=1,144	0.049 [0.057] N=973	0.081 [0.065] N=789
BMI-for-age z score below -2	0.128 P=190	0.053 [0.042] N=1,468	0.058 [0.045] N=1,331	0.065 [0.051] N=1,164	0.029 [0.051] N=990	0.036 [0.056] N=806

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.057 [0.383] N=1,305	-0.118 [0.401] N=1,162	0.035 [0.440] N=988	-0.259 [0.522] N=810	-0.493 [0.673] N=618
WHO weight-for-age z score	0.041 [0.202] N=1,479	0.070 [0.219] N=1,329	-0.072 [0.232] N=1,130	0.032 [0.250] N=924	0.079 [0.310] N=698
WHO weight-for-length/height z score, 0-60 months	0.074 [0.289] N=1,293	-0.048 [0.301] N=1,153	-0.205 [0.351] N=982	0.049 [0.380] N=803	-0.229 [0.464] N=611
WHO BMI-for-age z score	-0.171 [0.273] N=1,270	-0.302 [0.279] N=1,133	-0.497 [0.334] N=966	-0.279 [0.400] N=792	-0.407 [0.440] N=608
Stunted: HAZ below -2	0.058 [0.070] N=1,305	0.041 [0.073] N=1,162	0.019 [0.079] N=988	0.052 [0.097] N=810	0.084 [0.121] N=618
Underweight: WAZ below -2	0.026 [0.040] N=1,479	0.021 [0.044] N=1,329	0.024 [0.049] N=1,130	0.035 [0.054] N=924	-0.001 [0.056] N=698
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.046] N=1,293	**0.084* [0.050] N=1,153**	**0.135** [0.057] N=982**	0.043 [0.057] N=803	0.070 [0.070] N=611
Overweight: WLZ/WHZ above +2, 0-60 months	**0.123** [0.058] N=1,293**	0.096 [0.059] N=1,153	0.080 [0.065] N=982	0.103 [0.075] N=803	0.100 [0.087] N=611
Obese: WLZ/WHZ above +3, 0-60 months	**0.079* [0.042] N=1,293**	0.054 [0.042] N=1,153	0.051 [0.049] N=982	0.039 [0.061] N=803	-0.080 [0.068] N=611
CIAF: any stunting, underweight, or wasting, 0-60 months	0.120 [0.074] N=1,247	0.097 [0.080] N=1,110	0.126 [0.090] N=943	0.065 [0.096] N=772	0.138 [0.119] N=588
BMI-for-age z score below -2	**0.114*** [0.043] N=1,270**	**0.124** [0.048] N=1,133**	**0.151*** [0.057] N=966**	0.074 [0.057] N=792	0.087 [0.069] N=608

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.256	-0.182 [0.265] N=1,513	-0.218 [0.269] N=1,370	-0.135 [0.285] N=1,196	-0.276 [0.316] N=1,018	-0.427 [0.357] N=826
WHO weight-for-age z score	0.071	0.085 [0.136] N=1,710	0.105 [0.144] N=1,560	0.033 [0.148] N=1,361	0.095 [0.157] N=1,155	0.099 [0.176] N=929
WHO weight-for-length/height z score, 0-60 months	0.349	0.138 [0.216] N=1,501	0.074 [0.222] N=1,361	-0.007 [0.238] N=1,190	0.140 [0.258] N=1,011	0.074 [0.293] N=819
WHO BMI-for-age z score	0.279	0.096 [0.219] N=1,468	0.028 [0.225] N=1,331	-0.066 [0.249] N=1,164	0.087 [0.280] N=990	0.101 [0.313] N=806
Stunted: HAZ below -2	0.220 P=328	0.046 [0.046] N=1,513	0.037 [0.047] N=1,370	0.021 [0.050] N=1,196	0.036 [0.056] N=1,018	0.050 [0.063] N=826
Underweight: WAZ below -2	0.069 P=119	0.001 [0.027] N=1,710	-0.002 [0.028] N=1,560	-0.004 [0.029] N=1,361	-0.001 [0.031] N=1,155	-0.017 [0.032] N=929
Wasted: WLZ/WHZ below -2, 0-60 months	0.116 P=178	0.026 [0.036] N=1,501	0.031 [0.039] N=1,361	0.054 [0.042] N=1,190	0.007 [0.041] N=1,011	0.012 [0.046] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	0.067 [0.042] N=1,501	0.052 [0.042] N=1,361	0.038 [0.045] N=1,190	0.044 [0.050] N=1,011	0.046 [0.055] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=145	0.045 [0.034] N=1,501	0.034 [0.035] N=1,361	0.028 [0.038] N=1,190	0.027 [0.043] N=1,011	-0.006 [0.045] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=448	0.076 [0.051] N=1,448	0.065 [0.053] N=1,311	0.076 [0.056] N=1,144	0.049 [0.056] N=973	0.086 [0.064] N=789
BMI-for-age z score below -2	0.128 P=190	0.051 [0.042] N=1,468	0.057 [0.044] N=1,331	0.065 [0.050] N=1,164	0.028 [0.050] N=990	0.030 [0.054] N=806

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.037 [0.381] N=1,305	-0.075 [0.399] N=1,162	0.102 [0.440] N=988	-0.219 [0.528] N=810	-0.548 [0.688] N=618
WHO weight-for-age z score	0.076 [0.196] N=1,479	0.103 [0.214] N=1,329	-0.032 [0.225] N=1,130	0.061 [0.240] N=924	0.081 [0.289] N=698
WHO weight-for-length/height z score, 0-60 months	0.081 [0.288] N=1,293	-0.056 [0.299] N=1,153	-0.234 [0.345] N=982	0.051 [0.379] N=803	-0.184 [0.480] N=611
WHO BMI-for-age z score	-0.173 [0.273] N=1,270	-0.313 [0.278] N=1,133	-0.524 [0.333] N=966	-0.269 [0.404] N=792	-0.359 [0.458] N=608
Stunted: HAZ below -2	0.057 [0.070] N=1,305	0.036 [0.073] N=1,162	0.006 [0.080] N=988	0.044 [0.099] N=810	0.091 [0.125] N=618
Underweight: WAZ below -2	0.022 [0.040] N=1,479	0.018 [0.044] N=1,329	0.019 [0.049] N=1,130	0.031 [0.054] N=924	-0.004 [0.057] N=698
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.045] N=1,293	**0.087* [0.050] N=1,153**	**0.142** [0.056] N=982**	0.045 [0.055] N=803	0.070 [0.067] N=611
Overweight: WLZ/WHZ above +2, 0-60 months	**0.119** [0.057] N=1,293**	0.090 [0.058] N=1,153	0.074 [0.065] N=982	0.101 [0.074] N=803	0.103 [0.088] N=611
Obese: WLZ/WHZ above +3, 0-60 months	**0.083** [0.042] N=1,293**	0.057 [0.041] N=1,153	0.050 [0.047] N=982	0.043 [0.060] N=803	-0.077 [0.070] N=611
CIAF: any stunting, underweight, or wasting, 0-60 months	0.115 [0.073] N=1,247	0.092 [0.079] N=1,110	0.116 [0.089] N=943	0.058 [0.095] N=772	0.149 [0.119] N=588
BMI-for-age z score below -2	**0.114*** [0.043] N=1,270**	**0.125** [0.048] N=1,133**	**0.155*** [0.057] N=966**	0.077 [0.056] N=792	0.088 [0.067] N=608

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.260	-0.245 [0.268] N=1,400	-0.305 [0.275] N=1,268	-0.223 [0.285] N=1,111	-0.359 [0.308] N=944	-0.520 [0.344] N=768
WHO weight-for-age z score	0.067	0.094 [0.140] N=1,581	0.128 [0.148] N=1,442	0.019 [0.153] N=1,263	0.072 [0.161] N=1,073	0.041 [0.176] N=866
WHO weight-for-length/height z score, 0-60 months	0.351	0.187 [0.234] N=1,390	0.144 [0.244] N=1,261	0.012 [0.261] N=1,105	0.116 [0.278] N=938	0.027 [0.303] N=763
WHO BMI-for-age z score	0.281	0.118 [0.236] N=1,359	0.071 [0.246] N=1,233	-0.080 [0.272] N=1,082	0.042 [0.300] N=920	0.001 [0.329] N=752
Stunted: HAZ below -2	0.221 P=304	0.061 [0.049] N=1,400	0.060 [0.050] N=1,268	0.042 [0.052] N=1,111	0.048 [0.055] N=944	0.061 [0.063] N=768
Underweight: WAZ below -2	0.069 P=110	0.005 [0.027] N=1,581	0.002 [0.028] N=1,442	-0.002 [0.029] N=1,263	0.004 [0.031] N=1,073	-0.006 [0.031] N=866
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=165	0.028 [0.038] N=1,390	0.029 [0.040] N=1,261	0.055 [0.042] N=1,105	0.013 [0.042] N=938	0.022 [0.046] N=763
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=275	0.070 [0.044] N=1,390	0.054 [0.044] N=1,261	0.031 [0.047] N=1,105	0.028 [0.050] N=938	0.024 [0.054] N=763
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=137	0.042 [0.035] N=1,390	0.030 [0.035] N=1,261	0.021 [0.038] N=1,105	0.021 [0.043] N=938	-0.011 [0.044] N=763
CIAF: any stunting, underweight, or wasting, 0-60 months	0.314 P=416	**0.095* [0.054] N=1,340**	0.082 [0.056] N=1,214	**0.099* [0.058] N=1,062**	0.062 [0.058] N=902	0.104 [0.066] N=734
BMI-for-age z score below -2	0.127 P=173	0.060 [0.043] N=1,359	0.062 [0.046] N=1,233	0.073 [0.051] N=1,082	0.046 [0.052] N=920	0.056 [0.056] N=752

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.045 [0.399] N=1,203	-0.016 [0.421] N=1,071	0.150 [0.462] N=914	-0.109 [0.524] N=747	-0.349 [0.656] N=571
WHO weight-for-age z score	0.107 [0.207] N=1,363	0.154 [0.222] N=1,224	-0.064 [0.240] N=1,045	0.045 [0.259] N=855	0.050 [0.326] N=648
WHO weight-for-length/height z score, 0-60 months	0.138 [0.323] N=1,193	0.034 [0.342] N=1,064	-0.232 [0.397] N=908	-0.001 [0.432] N=741	-0.346 [0.528] N=566
WHO BMI-for-age z score	-0.142 [0.299] N=1,170	-0.260 [0.313] N=1,044	-0.571 [0.378] N=893	-0.350 [0.450] N=731	-0.607 [0.487] N=563
Stunted: HAZ below -2	0.054 [0.079] N=1,203	0.044 [0.083] N=1,071	0.013 [0.088] N=914	0.026 [0.104] N=747	0.044 [0.137] N=571
Underweight: WAZ below -2	0.026 [0.041] N=1,363	0.020 [0.045] N=1,224	0.021 [0.050] N=1,045	0.039 [0.058] N=855	0.002 [0.062] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.048] N=1,193	0.079 [0.053] N=1,064	**0.135** [0.058] N=908**	0.037 [0.059] N=741	0.067 [0.071] N=566
Overweight: WLZ/WHZ above +2, 0-60 months	**0.124* [0.063] N=1,193**	0.094 [0.065] N=1,064	0.065 [0.072] N=908	0.074 [0.083] N=741	0.052 [0.099] N=566
Obese: WLZ/WHZ above +3, 0-60 months	0.076 [0.047] N=1,193	0.049 [0.047] N=1,064	0.037 [0.054] N=908	0.030 [0.067] N=741	-0.095 [0.080] N=566
CIAF: any stunting, underweight, or wasting, 0-60 months	0.121 [0.084] N=1,150	0.096 [0.090] N=1,024	0.130 [0.099] N=872	0.039 [0.109] N=712	0.110 [0.143] N=544
BMI-for-age z score below -2	**0.124*** [0.045] N=1,170**	**0.132** [0.050] N=1,044**	**0.164*** [0.059] N=893**	0.095 [0.059] N=731	**0.119* [0.068] N=563**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.260	-0.264 [0.272] N=1,400	-0.324 [0.279] N=1,268	-0.240 [0.288] N=1,111	-0.385 [0.311] N=944	-0.554 [0.351] N=768
WHO weight-for-age z score	0.067	0.094 [0.140] N=1,581	0.125 [0.148] N=1,442	0.019 [0.153] N=1,263	0.068 [0.161] N=1,073	0.037 [0.178] N=866
WHO weight-for-length/height z score, 0-60 months	0.351	0.202 [0.236] N=1,390	0.153 [0.246] N=1,261	0.025 [0.263] N=1,105	0.127 [0.278] N=938	0.031 [0.309] N=763
WHO BMI-for-age z score	0.281	0.131 [0.239] N=1,359	0.078 [0.249] N=1,233	-0.069 [0.275] N=1,082	0.050 [0.301] N=920	0.009 [0.336] N=752
Stunted: HAZ below -2	0.221 P=304	0.061 [0.049] N=1,400	0.058 [0.050] N=1,268	0.039 [0.051] N=1,111	0.047 [0.055] N=944	0.062 [0.063] N=768
Underweight: WAZ below -2	0.069 P=110	0.004 [0.026] N=1,581	0.000 [0.028] N=1,442	-0.003 [0.029] N=1,263	0.002 [0.030] N=1,073	-0.007 [0.031] N=866
Wasted: WLZ/WHZ below -2, 0-60 months	0.117 P=165	0.023 [0.038] N=1,390	0.024 [0.040] N=1,261	0.049 [0.043] N=1,105	0.006 [0.042] N=938	0.016 [0.047] N=763
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=275	0.070 [0.043] N=1,390	0.053 [0.044] N=1,261	0.030 [0.046] N=1,105	0.027 [0.050] N=938	0.023 [0.054] N=763
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=137	0.042 [0.035] N=1,390	0.030 [0.035] N=1,261	0.020 [0.038] N=1,105	0.020 [0.043] N=938	-0.012 [0.044] N=763
CIAF: any stunting, underweight, or wasting, 0-60 months	0.314 P=416	0.089 [0.054] N=1,340	0.074 [0.055] N=1,214	0.089 [0.058] N=1,062	0.052 [0.058] N=902	0.095 [0.066] N=734
BMI-for-age z score below -2	0.127 P=173	0.056 [0.043] N=1,359	0.058 [0.046] N=1,233	0.069 [0.052] N=1,082	0.040 [0.051] N=920	0.051 [0.055] N=752

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.016 [0.402] N=1,203	-0.041 [0.423] N=1,071	0.122 [0.461] N=914	-0.186 [0.527] N=747	-0.498 [0.666] N=571
WHO weight-for-age z score	0.101 [0.207] N=1,363	0.145 [0.223] N=1,224	-0.069 [0.241] N=1,045	0.031 [0.258] N=855	0.018 [0.326] N=648
WHO weight-for-length/height z score, 0-60 months	0.135 [0.321] N=1,193	0.023 [0.338] N=1,064	-0.241 [0.392] N=908	-0.008 [0.423] N=741	-0.363 [0.508] N=566
WHO BMI-for-age z score	-0.139 [0.297] N=1,170	-0.266 [0.309] N=1,044	-0.572 [0.373] N=893	-0.337 [0.444] N=731	-0.571 [0.484] N=563
Stunted: HAZ below -2	0.057 [0.079] N=1,203	0.046 [0.082] N=1,071	0.016 [0.087] N=914	0.038 [0.104] N=747	0.075 [0.134] N=571
Underweight: WAZ below -2	0.026 [0.041] N=1,363	0.020 [0.044] N=1,224	0.021 [0.050] N=1,045	0.041 [0.057] N=855	0.014 [0.060] N=648
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,193	0.080 [0.053] N=1,064	**0.134** [0.058] N=908**	0.037 [0.058] N=741	0.073 [0.070] N=566
Overweight: WLZ/WHZ above +2, 0-60 months	**0.124** [0.062] N=1,193**	0.093 [0.063] N=1,064	0.063 [0.071] N=908	0.074 [0.082] N=741	0.058 [0.097] N=566
Obese: WLZ/WHZ above +3, 0-60 months	0.076 [0.046] N=1,193	0.048 [0.046] N=1,064	0.036 [0.053] N=908	0.031 [0.066] N=741	-0.090 [0.077] N=566
CIAF: any stunting, underweight, or wasting, 0-60 months	0.122 [0.083] N=1,150	0.097 [0.089] N=1,024	0.130 [0.098] N=872	0.045 [0.108] N=712	0.136 [0.139] N=544
BMI-for-age z score below -2	**0.124*** [0.044] N=1,170**	**0.133*** [0.050] N=1,044**	**0.165*** [0.059] N=893**	0.095 [0.057] N=731	**0.123* [0.066] N=563**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.