Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-3: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,606	-0.014 [0.022] N=1,721	-0.001 [0.021] N=1,588	-0.005 [0.023] N=1,430	-0.013 [0.025] N=1,226	-0.026 [0.030] N=1,010
Vegetables consumed at least weekly	0.895 P=1,539	0.011 [0.028] N=1,721	0.009 [0.028] N=1,588	-0.010 [0.029] N=1,430	-0.020 [0.029] N=1,226	-0.047 [0.034] N=1,010
Meat fish or poultry consumed at least weekly	0.775 P=1,342	0.070 [0.048] N=1,721	**0.086* [0.050] N=1,588**	0.077 [0.055] N=1,430	0.058 [0.061] N=1,226	0.038 [0.070] N=1,010
Legumes consumed at least weekly	0.773 P=1,336	-0.035 [0.041] N=1,721	-0.048 [0.043] N=1,588	-0.042 [0.044] N=1,430	-0.049 [0.048] N=1,226	-0.085 [0.059] N=1,010
Animal products consumed at least weekly	0.911 P=1,571	0.009 [0.029] N=1,721	0.007 [0.030] N=1,588	-0.008 [0.032] N=1,430	-0.002 [0.036] N=1,226	-0.008 [0.040] N=1,010
Sweets consumed at least weekly	0.622 P=1,073	-0.030 [0.038] N=1,721	-0.019 [0.040] N=1,588	-0.003 [0.043] N=1,430	-0.012 [0.045] N=1,226	0.014 [0.052] N=1,010
Snack foods consumed at least weekly	0.507 P=872	-0.031 [0.040] N=1,721	-0.025 [0.043] N=1,588	-0.025 [0.046] N=1,430	-0.034 [0.049] N=1,226	-0.048 [0.054] N=1,010
Packaged soft drinks consumed at least weekly	0.305 P=521	-0.009 [0.047] N=1,721	-0.000 [0.050] N=1,588	-0.013 [0.054] N=1,430	-0.000 [0.060] N=1,226	-0.009 [0.069] N=1,010
Grains consumed at least weekly	0.886 P=1,530	0.021 [0.030] N=1,721	0.032 [0.031] N=1,588	0.016 [0.032] N=1,430	0.022 [0.035] N=1,226	0.006 [0.042] N=1,010
Child was ever breastfed	0.967 P=1,911	-0.005 [0.014] N=1,980	-0.002 [0.015] N=1,832	-0.004 [0.016] N=1,646	-0.005 [0.017] N=1,400	-0.007 [0.017] N=1,156

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.009 [0.035] N=1,457	0.017 [0.034] N=1,324	0.011 [0.038] N=1,166	-0.005 [0.046] N=962	-0.040 [0.063] N=746
Vegetables consumed at least weekly	0.043 [0.043] N=1,457	0.047 [0.047] N=1,324	0.016 [0.050] N=1,166	-0.006 [0.057] N=962	-0.081 [0.077] N=746
Meat fish or poultry consumed at least weekly	0.062 [0.057] N=1,457	0.087 [0.059] N=1,324	0.068 [0.067] N=1,166	-0.006 [0.071] N=962	-0.080 [0.082] N=746
Legumes consumed at least weekly	0.018 [0.062] N=1,457	0.001 [0.068] N=1,324	0.031 [0.070] N=1,166	0.027 [0.084] N=962	-0.038 [0.120] N=746
Animal products consumed at least weekly	0.007 [0.039] N=1,457	0.003 [0.041] N=1,324	-0.035 [0.046] N=1,166	-0.028 [0.056] N=962	-0.058 [0.075] N=746
Sweets consumed at least weekly	-0.046 [0.053] N=1,457	-0.028 [0.058] N=1,324	0.000 [0.066] N=1,166	-0.031 [0.073] N=962	0.028 [0.101] N=746
Snack foods consumed at least weekly	-0.016 [0.054] N=1,457	-0.002 [0.060] N=1,324	0.004 [0.071] N=1,166	-0.008 [0.082] N=962	-0.057 [0.105] N=746
Packaged soft drinks consumed at least weekly	-0.015 [0.055] N=1,457	0.001 [0.060] N=1,324	-0.021 [0.066] N=1,166	-0.010 [0.078] N=962	-0.041 [0.107] N=746
Grains consumed at least weekly	0.053 [0.047] N=1,457	0.079 [0.052] N=1,324	0.057 [0.057] N=1,166	0.086 [0.072] N=962	0.071 [0.103] N=746
Child was ever breastfed	0.008 [0.022] N=1,678	0.016 [0.026] N=1,530	0.014 [0.030] N=1,344	0.011 [0.037] N=1,098	0.032 [0.044] N=854

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,606	-0.014 [0.022] N=1,721	-0.001 [0.021] N=1,588	-0.006 [0.023] N=1,430	-0.014 [0.025] N=1,226	-0.027 [0.030] N=1,010
Vegetables consumed at least weekly	0.895 P=1,539	0.011 [0.028] N=1,721	0.008 [0.028] N=1,588	-0.010 [0.028] N=1,430	-0.021 [0.029] N=1,226	-0.048 [0.034] N=1,010
Meat fish or poultry consumed at least weekly	0.775 P=1,342	0.070 [0.048] N=1,721	**0.086* [0.050] N=1,588**	0.077 [0.055] N=1,430	0.056 [0.062] N=1,226	0.036 [0.071] N=1,010
Legumes consumed at least weekly	0.773 P=1,336	-0.035 [0.041] N=1,721	-0.049 [0.043] N=1,588	-0.043 [0.044] N=1,430	-0.050 [0.049] N=1,226	-0.086 [0.059] N=1,010
Animal products consumed at least weekly	0.911 P=1,571	0.009 [0.029] N=1,721	0.006 [0.029] N=1,588	-0.009 [0.032] N=1,430	-0.003 [0.035] N=1,226	-0.009 [0.040] N=1,010
Sweets consumed at least weekly	0.622 P=1,073	-0.030 [0.038] N=1,721	-0.019 [0.040] N=1,588	-0.003 [0.043] N=1,430	-0.012 [0.045] N=1,226	0.014 [0.052] N=1,010
Snack foods consumed at least weekly	0.507 P=872	-0.032 [0.040] N=1,721	-0.026 [0.043] N=1,588	-0.026 [0.047] N=1,430	-0.035 [0.049] N=1,226	-0.048 [0.054] N=1,010
Packaged soft drinks consumed at least weekly	0.305 P=521	-0.009 [0.047] N=1,721	-0.001 [0.050] N=1,588	-0.013 [0.054] N=1,430	-0.001 [0.060] N=1,226	-0.009 [0.069] N=1,010
Grains consumed at least weekly	0.886 P=1,530	0.021 [0.030] N=1,721	0.031 [0.031] N=1,588	0.016 [0.032] N=1,430	0.021 [0.036] N=1,226	0.005 [0.042] N=1,010
Child was ever breastfed	0.967 P=1,911	-0.005 [0.014] N=1,980	-0.002 [0.015] N=1,832	-0.004 [0.016] N=1,646	-0.004 [0.017] N=1,400	-0.007 [0.017] N=1,156

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.035] N=1,457	0.018 [0.034] N=1,324	0.012 [0.038] N=1,166	-0.005 [0.045] N=962	-0.037 [0.063] N=746
Vegetables consumed at least weekly	0.043 [0.044] N=1,457	0.048 [0.047] N=1,324	0.017 [0.050] N=1,166	-0.005 [0.057] N=962	-0.079 [0.076] N=746
Meat fish or poultry consumed at least weekly	0.062 [0.057] N=1,457	0.087 [0.059] N=1,324	0.069 [0.067] N=1,166	-0.005 [0.072] N=962	-0.075 [0.081] N=746
Legumes consumed at least weekly	0.018 [0.062] N=1,457	0.001 [0.068] N=1,324	0.033 [0.070] N=1,166	0.029 [0.084] N=962	-0.035 [0.120] N=746
Animal products consumed at least weekly	0.009 [0.039] N=1,457	0.004 [0.041] N=1,324	-0.031 [0.046] N=1,166	-0.026 [0.056] N=962	-0.054 [0.074] N=746
Sweets consumed at least weekly	-0.046 [0.053] N=1,457	-0.028 [0.058] N=1,324	0.000 [0.066] N=1,166	-0.031 [0.073] N=962	0.028 [0.101] N=746
Snack foods consumed at least weekly	-0.015 [0.054] N=1,457	-0.000 [0.060] N=1,324	0.007 [0.070] N=1,166	-0.006 [0.082] N=962	-0.053 [0.105] N=746
Packaged soft drinks consumed at least weekly	-0.015 [0.055] N=1,457	0.001 [0.060] N=1,324	-0.021 [0.066] N=1,166	-0.010 [0.078] N=962	-0.041 [0.106] N=746
Grains consumed at least weekly	0.053 [0.047] N=1,457	0.079 [0.052] N=1,324	0.058 [0.057] N=1,166	0.086 [0.071] N=962	0.072 [0.103] N=746
Child was ever breastfed	0.008 [0.023] N=1,678	0.016 [0.026] N=1,530	0.014 [0.030] N=1,344	0.010 [0.037] N=1,098	0.031 [0.044] N=854

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,606	-0.024 [0.022] N=1,721	-0.011 [0.021] N=1,588	-0.015 [0.023] N=1,430	-0.021 [0.026] N=1,226	-0.035 [0.030] N=1,010
Vegetables consumed at least weekly	0.895 P=1,539	0.003 [0.028] N=1,721	-0.001 [0.028] N=1,588	-0.019 [0.029] N=1,430	-0.029 [0.029] N=1,226	**-0.059* [0.034] N=1,010**
Meat fish or poultry consumed at least weekly	0.775 P=1,342	0.045 [0.046] N=1,721	0.058 [0.049] N=1,588	0.053 [0.054] N=1,430	0.036 [0.060] N=1,226	0.010 [0.068] N=1,010
Legumes consumed at least weekly	0.773 P=1,336	-0.042 [0.041] N=1,721	-0.056 [0.043] N=1,588	-0.048 [0.044] N=1,430	-0.053 [0.048] N=1,226	-0.088 [0.058] N=1,010
Animal products consumed at least weekly	0.911 P=1,571	0.002 [0.029] N=1,721	-0.001 [0.030] N=1,588	-0.015 [0.033] N=1,430	-0.008 [0.036] N=1,226	-0.018 [0.040] N=1,010
Sweets consumed at least weekly	0.622 P=1,073	-0.032 [0.039] N=1,721	-0.022 [0.040] N=1,588	-0.008 [0.044] N=1,430	-0.014 [0.046] N=1,226	0.011 [0.052] N=1,010
Snack foods consumed at least weekly	0.507 P=872	-0.026 [0.040] N=1,721	-0.021 [0.042] N=1,588	-0.023 [0.046] N=1,430	-0.030 [0.049] N=1,226	-0.040 [0.053] N=1,010
Packaged soft drinks consumed at least weekly	0.305 P=521	0.005 [0.046] N=1,721	0.016 [0.049] N=1,588	0.002 [0.053] N=1,430	0.012 [0.058] N=1,226	0.008 [0.067] N=1,010
Grains consumed at least weekly	0.886 P=1,530	0.016 [0.030] N=1,721	0.027 [0.031] N=1,588	0.013 [0.032] N=1,430	0.018 [0.035] N=1,226	0.003 [0.042] N=1,010
Child was ever breastfed	0.967 P=1,911	-0.009 [0.014] N=1,980	-0.007 [0.015] N=1,832	-0.009 [0.016] N=1,646	-0.008 [0.017] N=1,400	-0.011 [0.018] N=1,156

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.034] N=1,457	0.009 [0.033] N=1,324	0.007 [0.037] N=1,166	0.003 [0.044] N=962	-0.028 [0.061] N=746
Vegetables consumed at least weekly	0.038 [0.043] N=1,457	0.041 [0.046] N=1,324	0.014 [0.050] N=1,166	0.003 [0.057] N=962	-0.071 [0.075] N=746
Meat fish or poultry consumed at least weekly	0.042 [0.055] N=1,457	0.062 [0.056] N=1,324	0.052 [0.065] N=1,166	-0.010 [0.071] N=962	-0.085 [0.083] N=746
Legumes consumed at least weekly	0.011 [0.062] N=1,457	-0.007 [0.067] N=1,324	0.030 [0.069] N=1,166	0.034 [0.084] N=962	-0.025 [0.121] N=746
Animal products consumed at least weekly	0.005 [0.038] N=1,457	-0.000 [0.041] N=1,324	-0.033 [0.047] N=1,166	-0.020 [0.057] N=962	-0.045 [0.074] N=746
Sweets consumed at least weekly	-0.047 [0.054] N=1,457	-0.031 [0.058] N=1,324	-0.002 [0.066] N=1,166	-0.029 [0.072] N=962	0.036 [0.102] N=746
Snack foods consumed at least weekly	-0.010 [0.053] N=1,457	0.004 [0.059] N=1,324	0.009 [0.068] N=1,166	-0.002 [0.078] N=962	-0.043 [0.100] N=746
Packaged soft drinks consumed at least weekly	-0.002 [0.054] N=1,457	0.020 [0.058] N=1,324	-0.005 [0.065] N=1,166	0.001 [0.077] N=962	-0.027 [0.102] N=746
Grains consumed at least weekly	0.048 [0.047] N=1,457	0.076 [0.052] N=1,324	0.057 [0.057] N=1,166	0.088 [0.071] N=962	0.072 [0.102] N=746
Child was ever breastfed	0.005 [0.023] N=1,678	0.012 [0.026] N=1,530	0.011 [0.030] N=1,344	0.012 [0.037] N=1,098	0.028 [0.044] N=854

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,606	-0.026 [0.022] N=1,721	-0.013 [0.022] N=1,588	-0.019 [0.023] N=1,430	-0.026 [0.026] N=1,226	-0.041 [0.031] N=1,010
Vegetables consumed at least weekly	0.895 P=1,539	0.003 [0.028] N=1,721	0.000 [0.028] N=1,588	-0.020 [0.029] N=1,430	-0.031 [0.029] N=1,226	**-0.059* [0.034] N=1,010**
Meat fish or poultry consumed at least weekly	0.775 P=1,342	0.043 [0.046] N=1,721	0.056 [0.049] N=1,588	0.044 [0.053] N=1,430	0.026 [0.059] N=1,226	-0.005 [0.068] N=1,010
Legumes consumed at least weekly	0.773 P=1,336	-0.043 [0.042] N=1,721	-0.059 [0.044] N=1,588	-0.056 [0.045] N=1,430	-0.060 [0.050] N=1,226	-0.096 [0.060] N=1,010
Animal products consumed at least weekly	0.911 P=1,571	0.001 [0.029] N=1,721	-0.001 [0.030] N=1,588	-0.016 [0.032] N=1,430	-0.010 [0.036] N=1,226	-0.019 [0.040] N=1,010
Sweets consumed at least weekly	0.622 P=1,073	-0.031 [0.038] N=1,721	-0.023 [0.040] N=1,588	-0.011 [0.043] N=1,430	-0.015 [0.045] N=1,226	0.010 [0.053] N=1,010
Snack foods consumed at least weekly	0.507 P=872	-0.027 [0.040] N=1,721	-0.023 [0.043] N=1,588	-0.026 [0.047] N=1,430	-0.032 [0.050] N=1,226	-0.043 [0.054] N=1,010
Packaged soft drinks consumed at least weekly	0.305 P=521	0.006 [0.046] N=1,721	0.013 [0.049] N=1,588	-0.001 [0.053] N=1,430	0.014 [0.058] N=1,226	0.012 [0.067] N=1,010
Grains consumed at least weekly	0.886 P=1,530	0.019 [0.030] N=1,721	0.030 [0.031] N=1,588	0.015 [0.032] N=1,430	0.020 [0.036] N=1,226	0.008 [0.043] N=1,010
Child was ever breastfed	0.967 P=1,911	-0.010 [0.014] N=1,980	-0.007 [0.015] N=1,832	-0.010 [0.017] N=1,646	-0.009 [0.018] N=1,400	-0.012 [0.018] N=1,156

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.034] N=1,457	0.010 [0.034] N=1,324	0.004 [0.037] N=1,166	-0.003 [0.045] N=962	-0.041 [0.063] N=746
Vegetables consumed at least weekly	0.039 [0.043] N=1,457	0.042 [0.046] N=1,324	0.011 [0.050] N=1,166	-0.003 [0.057] N=962	-0.087 [0.076] N=746
Meat fish or poultry consumed at least weekly	0.043 [0.055] N=1,457	0.066 [0.056] N=1,324	0.051 [0.065] N=1,166	-0.003 [0.070] N=962	-0.090 [0.085] N=746
Legumes consumed at least weekly	0.016 [0.062] N=1,457	-0.003 [0.067] N=1,324	0.030 [0.069] N=1,166	0.045 [0.083] N=962	-0.021 [0.124] N=746
Animal products consumed at least weekly	0.003 [0.039] N=1,457	-0.001 [0.041] N=1,324	-0.036 [0.046] N=1,166	-0.032 [0.057] N=962	-0.066 [0.078] N=746
Sweets consumed at least weekly	-0.040 [0.053] N=1,457	-0.026 [0.056] N=1,324	0.001 [0.064] N=1,166	-0.017 [0.073] N=962	0.040 [0.102] N=746
Snack foods consumed at least weekly	-0.004 [0.052] N=1,457	0.005 [0.058] N=1,324	0.011 [0.067] N=1,166	0.009 [0.077] N=962	-0.034 [0.098] N=746
Packaged soft drinks consumed at least weekly	0.002 [0.053] N=1,457	0.014 [0.057] N=1,324	-0.010 [0.064] N=1,166	0.005 [0.076] N=962	-0.014 [0.100] N=746
Grains consumed at least weekly	0.049 [0.047] N=1,457	0.076 [0.051] N=1,324	0.055 [0.056] N=1,166	0.083 [0.070] N=962	0.060 [0.099] N=746
Child was ever breastfed	0.006 [0.023] N=1,678	0.012 [0.026] N=1,530	0.010 [0.030] N=1,344	0.012 [0.038] N=1,098	0.030 [0.046] N=854

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,490	-0.024 [0.023] N=1,596	-0.014 [0.022] N=1,478	-0.017 [0.023] N=1,329	-0.025 [0.026] N=1,143	-0.045 [0.029] N=943
Vegetables consumed at least weekly	0.897 P=1,429	-0.004 [0.029] N=1,596	-0.008 [0.029] N=1,478	-0.030 [0.030] N=1,329	-0.035 [0.030] N=1,143	**-0.064* [0.035] N=943**
Meat fish or poultry consumed at least weekly	0.776 P=1,244	0.055 [0.046] N=1,596	0.062 [0.048] N=1,478	0.048 [0.053] N=1,329	0.021 [0.058] N=1,143	-0.012 [0.065] N=943
Legumes consumed at least weekly	0.771 P=1,234	-0.052 [0.042] N=1,596	-0.071 [0.044] N=1,478	-0.067 [0.044] N=1,329	-0.075 [0.050] N=1,143	**-0.115* [0.060] N=943**
Animal products consumed at least weekly	0.914 P=1,461	0.008 [0.030] N=1,596	0.005 [0.031] N=1,478	-0.012 [0.033] N=1,329	-0.012 [0.037] N=1,143	-0.022 [0.041] N=943
Sweets consumed at least weekly	0.624 P=997	-0.015 [0.038] N=1,596	-0.010 [0.040] N=1,478	0.004 [0.043] N=1,329	-0.002 [0.045] N=1,143	0.027 [0.052] N=943
Snack foods consumed at least weekly	0.510 P=811	-0.033 [0.042] N=1,596	-0.027 [0.045] N=1,478	-0.029 [0.049] N=1,329	-0.038 [0.053] N=1,143	-0.034 [0.060] N=943
Packaged soft drinks consumed at least weekly	0.312 P=494	0.021 [0.047] N=1,596	0.030 [0.049] N=1,478	0.009 [0.053] N=1,329	0.016 [0.058] N=1,143	0.023 [0.067] N=943
Grains consumed at least weekly	0.888 P=1,421	0.012 [0.031] N=1,596	0.020 [0.032] N=1,478	-0.002 [0.033] N=1,329	0.006 [0.037] N=1,143	-0.011 [0.044] N=943
Child was ever breastfed	0.967 P=1,768	-0.010 [0.015] N=1,832	-0.005 [0.016] N=1,700	-0.006 [0.017] N=1,523	-0.005 [0.018] N=1,302	-0.010 [0.019] N=1,077

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.014 [0.036] N=1,349	0.005 [0.037] N=1,231	-0.001 [0.042] N=1,082	-0.015 [0.049] N=896	-0.062 [0.067] N=696
Vegetables consumed at least weekly	0.029 [0.045] N=1,349	0.027 [0.049] N=1,231	-0.010 [0.052] N=1,082	-0.020 [0.059] N=896	-0.103 [0.073] N=696
Meat fish or poultry consumed at least weekly	0.070 [0.058] N=1,349	0.080 [0.059] N=1,231	0.058 [0.067] N=1,082	-0.023 [0.073] N=896	-0.101 [0.080] N=696
Legumes consumed at least weekly	0.005 [0.063] N=1,349	-0.025 [0.068] N=1,231	-0.000 [0.070] N=1,082	-0.008 [0.085] N=896	-0.092 [0.119] N=696
Animal products consumed at least weekly	0.012 [0.041] N=1,349	0.005 [0.044] N=1,231	-0.032 [0.049] N=1,082	-0.034 [0.060] N=896	-0.074 [0.076] N=696
Sweets consumed at least weekly	-0.033 [0.055] N=1,349	-0.027 [0.061] N=1,231	-0.006 [0.069] N=1,082	-0.036 [0.076] N=896	0.005 [0.106] N=696
Snack foods consumed at least weekly	-0.024 [0.055] N=1,349	-0.011 [0.061] N=1,231	-0.014 [0.071] N=1,082	-0.037 [0.082] N=896	-0.077 [0.104] N=696
Packaged soft drinks consumed at least weekly	0.012 [0.057] N=1,349	0.027 [0.060] N=1,231	-0.014 [0.067] N=1,082	-0.017 [0.079] N=896	-0.028 [0.111] N=696
Grains consumed at least weekly	0.048 [0.049] N=1,349	0.070 [0.054] N=1,231	0.032 [0.058] N=1,082	0.065 [0.073] N=896	0.047 [0.103] N=696
Child was ever breastfed	0.001 [0.024] N=1,550	0.011 [0.027] N=1,418	0.009 [0.031] N=1,241	0.009 [0.038] N=1,020	0.023 [0.046] N=795

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,490	-0.025 [0.022] N=1,596	-0.015 [0.022] N=1,478	-0.017 [0.023] N=1,329	-0.025 [0.025] N=1,143	-0.044 [0.028] N=943
Vegetables consumed at least weekly	0.897 P=1,429	-0.004 [0.029] N=1,596	-0.009 [0.029] N=1,478	-0.031 [0.030] N=1,329	-0.035 [0.030] N=1,143	**-0.063* [0.035] N=943**
Meat fish or poultry consumed at least weekly	0.776 P=1,244	0.060 [0.046] N=1,596	0.066 [0.048] N=1,478	0.052 [0.053] N=1,329	0.027 [0.059] N=1,143	-0.000 [0.066] N=943
Legumes consumed at least weekly	0.771 P=1,234	-0.054 [0.042] N=1,596	**-0.074* [0.044] N=1,478**	-0.071 [0.045] N=1,329	-0.079 [0.050] N=1,143	**-0.122** [0.060] N=943**
Animal products consumed at least weekly	0.914 P=1,461	0.007 [0.030] N=1,596	0.004 [0.031] N=1,478	-0.012 [0.033] N=1,329	-0.011 [0.037] N=1,143	-0.020 [0.041] N=943
Sweets consumed at least weekly	0.624 P=997	-0.012 [0.038] N=1,596	-0.008 [0.040] N=1,478	0.008 [0.042] N=1,329	0.002 [0.044] N=1,143	0.032 [0.050] N=943
Snack foods consumed at least weekly	0.510 P=811	-0.033 [0.042] N=1,596	-0.028 [0.045] N=1,478	-0.029 [0.049] N=1,329	-0.038 [0.053] N=1,143	-0.039 [0.058] N=943
Packaged soft drinks consumed at least weekly	0.312 P=494	0.018 [0.047] N=1,596	0.027 [0.049] N=1,478	0.007 [0.053] N=1,329	0.014 [0.059] N=1,143	0.016 [0.068] N=943
Grains consumed at least weekly	0.888 P=1,421	0.011 [0.031] N=1,596	0.020 [0.032] N=1,478	-0.003 [0.033] N=1,329	0.005 [0.037] N=1,143	-0.012 [0.043] N=943
Child was ever breastfed	0.967 P=1,768	-0.009 [0.014] N=1,832	-0.004 [0.016] N=1,700	-0.006 [0.017] N=1,523	-0.005 [0.018] N=1,302	-0.009 [0.018] N=1,077

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.036] N=1,349	0.004 [0.036] N=1,231	0.000 [0.041] N=1,082	-0.013 [0.048] N=896	-0.059 [0.063] N=696
Vegetables consumed at least weekly	0.026 [0.045] N=1,349	0.025 [0.048] N=1,231	-0.010 [0.052] N=1,082	-0.022 [0.058] N=896	-0.101 [0.071] N=696
Meat fish or poultry consumed at least weekly	0.071 [0.058] N=1,349	0.081 [0.059] N=1,231	0.059 [0.066] N=1,082	-0.021 [0.073] N=896	-0.097 [0.080] N=696
Legumes consumed at least weekly	0.004 [0.063] N=1,349	-0.025 [0.069] N=1,231	0.003 [0.070] N=1,082	-0.006 [0.086] N=896	-0.090 [0.119] N=696
Animal products consumed at least weekly	0.009 [0.040] N=1,349	0.003 [0.043] N=1,231	-0.032 [0.048] N=1,082	-0.035 [0.059] N=896	-0.072 [0.075] N=696
Sweets consumed at least weekly	-0.039 [0.055] N=1,349	-0.037 [0.060] N=1,231	-0.014 [0.069] N=1,082	-0.040 [0.076] N=896	0.006 [0.105] N=696
Snack foods consumed at least weekly	-0.031 [0.055] N=1,349	-0.021 [0.061] N=1,231	-0.021 [0.071] N=1,082	-0.040 [0.083] N=896	-0.078 [0.106] N=696
Packaged soft drinks consumed at least weekly	0.006 [0.057] N=1,349	0.021 [0.061] N=1,231	-0.017 [0.068] N=1,082	-0.017 [0.080] N=896	-0.027 [0.112] N=696
Grains consumed at least weekly	0.047 [0.049] N=1,349	0.069 [0.054] N=1,231	0.035 [0.059] N=1,082	0.067 [0.075] N=896	0.050 [0.105] N=696
Child was ever breastfed	0.004 [0.024] N=1,550	0.014 [0.027] N=1,418	0.011 [0.032] N=1,241	0.013 [0.039] N=1,020	0.028 [0.048] N=795

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,447	0.019 [0.028] N=1,548	0.020 [0.029] N=1,410	0.011 [0.032] N=1,229	-0.004 [0.032] N=1,034	0.002 [0.037] N=829
Vegetables consumed at least weekly	0.897 P=1,390	0.038 [0.032] N=1,548	0.044 [0.034] N=1,410	0.040 [0.037] N=1,229	0.029 [0.040] N=1,034	0.028 [0.044] N=829
Meat fish or poultry consumed at least weekly	0.767 P=1,201	**0.079* [0.043] N=1,548**	**0.092** [0.045] N=1,410**	**0.092* [0.048] N=1,229**	**0.128** [0.050] N=1,034**	**0.123** [0.056] N=829**
Legumes consumed at least weekly	0.770 P=1,206	-0.021 [0.041] N=1,548	0.002 [0.044] N=1,410	-0.009 [0.048] N=1,229	-0.021 [0.052] N=1,034	0.000 [0.059] N=829
Animal products consumed at least weekly	0.909 P=1,412	0.036 [0.031] N=1,548	0.041 [0.032] N=1,410	0.035 [0.035] N=1,229	0.027 [0.036] N=1,034	0.046 [0.041] N=829
Sweets consumed at least weekly	0.628 P=984	-0.007 [0.050] N=1,548	-0.011 [0.052] N=1,410	-0.024 [0.056] N=1,229	-0.028 [0.059] N=1,034	-0.067 [0.067] N=829
Snack foods consumed at least weekly	0.509 P=792	-0.021 [0.049] N=1,548	-0.017 [0.052] N=1,410	-0.008 [0.056] N=1,229	0.005 [0.059] N=1,034	-0.035 [0.066] N=829
Packaged soft drinks consumed at least weekly	0.297 P=461	-0.002 [0.043] N=1,548	-0.000 [0.045] N=1,410	0.027 [0.048] N=1,229	0.026 [0.052] N=1,034	0.002 [0.060] N=829
Grains consumed at least weekly	0.885 P=1,379	0.055 [0.038] N=1,548	0.066 [0.040] N=1,410	0.068 [0.043] N=1,229	0.075 [0.047] N=1,034	0.069 [0.056] N=829
Child was ever breastfed	0.965 P=1,698	-0.020 [0.018] N=1,761	-0.020 [0.018] N=1,605	-0.011 [0.018] N=1,398	-0.007 [0.019] N=1,180	-0.002 [0.021] N=944

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.018 [0.039] N=1,344	-0.022 [0.041] N=1,206	-0.046 [0.047] N=1,025	-0.061 [0.050] N=830	-0.069 [0.070] N=625
Vegetables consumed at least weekly	-0.010 [0.045] N=1,344	-0.004 [0.049] N=1,206	-0.025 [0.056] N=1,025	-0.045 [0.064] N=830	-0.066 [0.086] N=625
Meat fish or poultry consumed at least weekly	-0.001 [0.057] N=1,344	0.010 [0.059] N=1,206	-0.018 [0.065] N=1,025	0.030 [0.069] N=830	-0.031 [0.084] N=625
Legumes consumed at least weekly	-0.061 [0.052] N=1,344	-0.027 [0.058] N=1,206	-0.052 [0.068] N=1,025	-0.068 [0.074] N=830	-0.026 [0.085] N=625
Animal products consumed at least weekly	-0.014 [0.044] N=1,344	-0.006 [0.047] N=1,206	-0.030 [0.053] N=1,025	-0.059 [0.055] N=830	-0.049 [0.070] N=625
Sweets consumed at least weekly	-0.002 [0.063] N=1,344	-0.009 [0.068] N=1,206	-0.031 [0.078] N=1,025	-0.048 [0.090] N=830	-0.173 [0.108] N=625
Snack foods consumed at least weekly	-0.009 [0.068] N=1,344	-0.006 [0.074] N=1,206	0.014 [0.083] N=1,025	0.048 [0.096] N=830	-0.043 [0.120] N=625
Packaged soft drinks consumed at least weekly	-0.018 [0.065] N=1,344	-0.011 [0.071] N=1,206	0.040 [0.081] N=1,025	0.043 [0.095] N=830	-0.015 [0.120] N=625
Grains consumed at least weekly	-0.015 [0.040] N=1,344	-0.011 [0.045] N=1,206	-0.025 [0.049] N=1,025	-0.049 [0.056] N=830	**-0.133* [0.077] N=625**
Child was ever breastfed	-0.007 [0.033] N=1,529	-0.005 [0.034] N=1,373	0.014 [0.037] N=1,166	0.029 [0.042] N=948	0.053 [0.056] N=712

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,447	0.019 [0.028] N=1,548	0.020 [0.029] N=1,410	0.011 [0.032] N=1,229	-0.004 [0.032] N=1,034	0.002 [0.038] N=829
Vegetables consumed at least weekly	0.897 P=1,390	0.038 [0.032] N=1,548	0.045 [0.034] N=1,410	0.040 [0.037] N=1,229	0.029 [0.040] N=1,034	0.027 [0.044] N=829
Meat fish or poultry consumed at least weekly	0.767 P=1,201	**0.080* [0.043] N=1,548**	**0.093** [0.045] N=1,410**	**0.093* [0.048] N=1,229**	**0.128** [0.050] N=1,034**	**0.122** [0.057] N=829**
Legumes consumed at least weekly	0.770 P=1,206	-0.021 [0.041] N=1,548	0.002 [0.044] N=1,410	-0.009 [0.048] N=1,229	-0.021 [0.052] N=1,034	0.001 [0.059] N=829
Animal products consumed at least weekly	0.909 P=1,412	0.037 [0.031] N=1,548	0.042 [0.032] N=1,410	0.036 [0.034] N=1,229	0.027 [0.036] N=1,034	0.044 [0.041] N=829
Sweets consumed at least weekly	0.628 P=984	-0.006 [0.050] N=1,548	-0.010 [0.052] N=1,410	-0.023 [0.056] N=1,229	-0.028 [0.059] N=1,034	-0.068 [0.067] N=829
Snack foods consumed at least weekly	0.509 P=792	-0.019 [0.049] N=1,548	-0.016 [0.052] N=1,410	-0.007 [0.056] N=1,229	0.005 [0.059] N=1,034	-0.036 [0.066] N=829
Packaged soft drinks consumed at least weekly	0.297 P=461	-0.002 [0.043] N=1,548	0.000 [0.045] N=1,410	0.028 [0.048] N=1,229	0.026 [0.052] N=1,034	-0.000 [0.059] N=829
Grains consumed at least weekly	0.885 P=1,379	0.055 [0.038] N=1,548	**0.067* [0.040] N=1,410**	0.068 [0.043] N=1,229	0.075 [0.048] N=1,034	0.069 [0.056] N=829
Child was ever breastfed	0.965 P=1,698	-0.020 [0.018] N=1,761	-0.020 [0.019] N=1,605	-0.011 [0.018] N=1,398	-0.007 [0.019] N=1,180	-0.003 [0.021] N=944

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.017 [0.039] N=1,344	-0.020 [0.041] N=1,206	-0.045 [0.048] N=1,025	-0.061 [0.051] N=830	-0.069 [0.071] N=625
Vegetables consumed at least weekly	-0.009 [0.045] N=1,344	-0.004 [0.049] N=1,206	-0.025 [0.056] N=1,025	-0.043 [0.064] N=830	-0.066 [0.086] N=625
Meat fish or poultry consumed at least weekly	0.002 [0.057] N=1,344	0.015 [0.059] N=1,206	-0.013 [0.065] N=1,025	0.036 [0.069] N=830	-0.023 [0.085] N=625
Legumes consumed at least weekly	-0.061 [0.051] N=1,344	-0.028 [0.057] N=1,206	-0.055 [0.067] N=1,025	-0.071 [0.074] N=830	-0.036 [0.085] N=625
Animal products consumed at least weekly	-0.009 [0.042] N=1,344	0.000 [0.046] N=1,206	-0.024 [0.052] N=1,025	-0.052 [0.054] N=830	-0.041 [0.069] N=625
Sweets consumed at least weekly	0.002 [0.063] N=1,344	-0.003 [0.069] N=1,206	-0.025 [0.078] N=1,025	-0.047 [0.090] N=830	-0.168 [0.108] N=625
Snack foods consumed at least weekly	-0.002 [0.069] N=1,344	0.000 [0.075] N=1,206	0.020 [0.083] N=1,025	0.051 [0.096] N=830	-0.035 [0.121] N=625
Packaged soft drinks consumed at least weekly	-0.017 [0.065] N=1,344	-0.009 [0.070] N=1,206	0.043 [0.081] N=1,025	0.043 [0.095] N=830	-0.008 [0.119] N=625
Grains consumed at least weekly	-0.015 [0.041] N=1,344	-0.011 [0.046] N=1,206	-0.026 [0.050] N=1,025	-0.053 [0.057] N=830	**-0.137* [0.079] N=625**
Child was ever breastfed	-0.006 [0.033] N=1,529	-0.004 [0.035] N=1,373	0.014 [0.037] N=1,166	0.031 [0.043] N=948	0.055 [0.057] N=712

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,447	0.018 [0.027] N=1,548	0.019 [0.029] N=1,410	0.012 [0.032] N=1,229	-0.002 [0.033] N=1,034	0.003 [0.038] N=829
Vegetables consumed at least weekly	0.897 P=1,390	0.038 [0.032] N=1,548	0.044 [0.033] N=1,410	0.042 [0.036] N=1,229	0.030 [0.040] N=1,034	0.029 [0.045] N=829
Meat fish or poultry consumed at least weekly	0.767 P=1,201	**0.078* [0.042] N=1,548**	**0.091** [0.045] N=1,410**	**0.100** [0.047] N=1,229**	**0.134*** [0.049] N=1,034**	**0.127** [0.055] N=829**
Legumes consumed at least weekly	0.770 P=1,206	-0.022 [0.042] N=1,548	0.001 [0.045] N=1,410	-0.008 [0.049] N=1,229	-0.019 [0.052] N=1,034	0.005 [0.061] N=829
Animal products consumed at least weekly	0.909 P=1,412	0.036 [0.031] N=1,548	0.040 [0.032] N=1,410	0.037 [0.035] N=1,229	0.029 [0.036] N=1,034	0.046 [0.041] N=829
Sweets consumed at least weekly	0.628 P=984	-0.005 [0.050] N=1,548	-0.010 [0.052] N=1,410	-0.023 [0.056] N=1,229	-0.028 [0.059] N=1,034	-0.068 [0.067] N=829
Snack foods consumed at least weekly	0.509 P=792	-0.018 [0.049] N=1,548	-0.016 [0.053] N=1,410	-0.009 [0.056] N=1,229	0.002 [0.059] N=1,034	-0.038 [0.067] N=829
Packaged soft drinks consumed at least weekly	0.297 P=461	-0.001 [0.044] N=1,548	-0.001 [0.046] N=1,410	0.024 [0.049] N=1,229	0.021 [0.054] N=1,034	-0.002 [0.062] N=829
Grains consumed at least weekly	0.885 P=1,379	0.054 [0.038] N=1,548	0.066 [0.040] N=1,410	0.069 [0.043] N=1,229	0.076 [0.048] N=1,034	0.071 [0.057] N=829
Child was ever breastfed	0.965 P=1,698	-0.020 [0.018] N=1,761	-0.019 [0.019] N=1,605	-0.009 [0.018] N=1,398	-0.004 [0.019] N=1,180	-0.000 [0.021] N=944

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.038] N=1,344	-0.020 [0.040] N=1,206	-0.040 [0.046] N=1,025	-0.057 [0.051] N=830	-0.067 [0.071] N=625
Vegetables consumed at least weekly	-0.007 [0.045] N=1,344	-0.003 [0.049] N=1,206	-0.019 [0.055] N=1,025	-0.038 [0.064] N=830	-0.063 [0.088] N=625
Meat fish or poultry consumed at least weekly	0.006 [0.054] N=1,344	0.020 [0.056] N=1,206	0.007 [0.062] N=1,025	0.055 [0.069] N=830	-0.016 [0.080] N=625
Legumes consumed at least weekly	-0.059 [0.051] N=1,344	-0.027 [0.058] N=1,206	-0.052 [0.068] N=1,025	-0.066 [0.074] N=830	-0.036 [0.085] N=625
Animal products consumed at least weekly	-0.010 [0.042] N=1,344	-0.003 [0.045] N=1,206	-0.021 [0.052] N=1,025	-0.051 [0.055] N=830	-0.041 [0.068] N=625
Sweets consumed at least weekly	0.002 [0.062] N=1,344	-0.006 [0.068] N=1,206	-0.027 [0.077] N=1,025	-0.053 [0.088] N=830	**-0.174* [0.103] N=625**
Snack foods consumed at least weekly	-0.000 [0.068] N=1,344	0.000 [0.074] N=1,206	0.012 [0.082] N=1,025	0.041 [0.093] N=830	-0.040 [0.119] N=625
Packaged soft drinks consumed at least weekly	-0.023 [0.066] N=1,344	-0.020 [0.071] N=1,206	0.026 [0.082] N=1,025	0.021 [0.095] N=830	-0.019 [0.122] N=625
Grains consumed at least weekly	-0.016 [0.041] N=1,344	-0.013 [0.046] N=1,206	-0.025 [0.050] N=1,025	-0.054 [0.057] N=830	**-0.140* [0.078] N=625**
Child was ever breastfed	-0.006 [0.033] N=1,529	-0.005 [0.035] N=1,373	0.017 [0.037] N=1,166	0.034 [0.042] N=948	0.055 [0.056] N=712

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.935 P=1,447	0.017 [0.027] N=1,548	0.019 [0.028] N=1,410	0.013 [0.031] N=1,229	-0.008 [0.032] N=1,034	-0.007 [0.037] N=829
Vegetables consumed at least weekly	0.897 P=1,390	0.038 [0.033] N=1,548	0.046 [0.034] N=1,410	0.044 [0.037] N=1,229	0.027 [0.040] N=1,034	0.026 [0.044] N=829
Meat fish or poultry consumed at least weekly	0.767 P=1,201	**0.078* [0.043] N=1,548**	**0.091** [0.045] N=1,410**	**0.096** [0.048] N=1,229**	**0.122** [0.051] N=1,034**	**0.105* [0.056] N=829**
Legumes consumed at least weekly	0.770 P=1,206	-0.017 [0.042] N=1,548	0.005 [0.045] N=1,410	-0.003 [0.048] N=1,229	-0.017 [0.052] N=1,034	0.004 [0.059] N=829
Animal products consumed at least weekly	0.909 P=1,412	0.039 [0.030] N=1,548	0.046 [0.031] N=1,410	0.045 [0.034] N=1,229	0.032 [0.036] N=1,034	0.046 [0.041] N=829
Sweets consumed at least weekly	0.628 P=984	0.001 [0.050] N=1,548	-0.002 [0.053] N=1,410	-0.018 [0.057] N=1,229	-0.024 [0.060] N=1,034	-0.064 [0.069] N=829
Snack foods consumed at least weekly	0.509 P=792	-0.010 [0.049] N=1,548	-0.007 [0.052] N=1,410	-0.002 [0.055] N=1,229	0.009 [0.058] N=1,034	-0.028 [0.067] N=829
Packaged soft drinks consumed at least weekly	0.297 P=461	0.009 [0.043] N=1,548	0.008 [0.045] N=1,410	0.029 [0.048] N=1,229	0.028 [0.053] N=1,034	0.007 [0.061] N=829
Grains consumed at least weekly	0.885 P=1,379	0.051 [0.037] N=1,548	0.062 [0.040] N=1,410	0.065 [0.043] N=1,229	0.069 [0.048] N=1,034	0.062 [0.056] N=829
Child was ever breastfed	0.965 P=1,698	-0.019 [0.018] N=1,761	-0.019 [0.018] N=1,605	-0.010 [0.018] N=1,398	-0.006 [0.019] N=1,180	-0.003 [0.020] N=944

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.037] N=1,344	-0.019 [0.040] N=1,206	-0.037 [0.046] N=1,025	-0.066 [0.052] N=830	-0.088 [0.071] N=625
Vegetables consumed at least weekly	-0.007 [0.045] N=1,344	-0.001 [0.050] N=1,206	-0.017 [0.056] N=1,025	-0.043 [0.065] N=830	-0.072 [0.086] N=625
Meat fish or poultry consumed at least weekly	0.008 [0.055] N=1,344	0.022 [0.059] N=1,206	0.005 [0.065] N=1,025	0.036 [0.075] N=830	-0.057 [0.085] N=625
Legumes consumed at least weekly	-0.055 [0.051] N=1,344	-0.022 [0.058] N=1,206	-0.041 [0.069] N=1,025	-0.060 [0.075] N=830	-0.029 [0.085] N=625
Animal products consumed at least weekly	-0.004 [0.040] N=1,344	0.007 [0.043] N=1,206	-0.011 [0.050] N=1,025	-0.049 [0.053] N=830	-0.048 [0.064] N=625
Sweets consumed at least weekly	0.007 [0.063] N=1,344	0.002 [0.069] N=1,206	-0.024 [0.078] N=1,025	-0.053 [0.089] N=830	-0.175 [0.107] N=625
Snack foods consumed at least weekly	0.003 [0.067] N=1,344	0.003 [0.073] N=1,206	0.014 [0.081] N=1,025	0.038 [0.092] N=830	-0.042 [0.117] N=625
Packaged soft drinks consumed at least weekly	-0.011 [0.064] N=1,344	-0.010 [0.069] N=1,206	0.029 [0.079] N=1,025	0.031 [0.093] N=830	0.000 [0.120] N=625
Grains consumed at least weekly	-0.017 [0.040] N=1,344	-0.016 [0.045] N=1,206	-0.031 [0.050] N=1,025	-0.063 [0.058] N=830	**-0.155* [0.079] N=625**
Child was ever breastfed	-0.004 [0.032] N=1,529	-0.003 [0.034] N=1,373	0.019 [0.037] N=1,166	0.034 [0.042] N=948	0.054 [0.056] N=712

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.936 P=1,342	0.015 [0.029] N=1,434	0.015 [0.031] N=1,304	0.018 [0.034] N=1,141	-0.004 [0.034] N=959	0.001 [0.040] N=772
Vegetables consumed at least weekly	0.900 P=1,292	0.034 [0.032] N=1,434	0.040 [0.034] N=1,304	0.052 [0.037] N=1,141	0.034 [0.041] N=959	0.022 [0.046] N=772
Meat fish or poultry consumed at least weekly	0.768 P=1,113	**0.081* [0.044] N=1,434**	**0.091** [0.046] N=1,304**	**0.099** [0.048] N=1,141**	**0.111** [0.050] N=959**	**0.103* [0.056] N=772**
Legumes consumed at least weekly	0.770 P=1,117	-0.018 [0.044] N=1,434	0.005 [0.047] N=1,304	-0.001 [0.051] N=1,141	-0.021 [0.056] N=959	-0.006 [0.065] N=772
Animal products consumed at least weekly	0.912 P=1,313	0.035 [0.032] N=1,434	0.043 [0.034] N=1,304	0.044 [0.036] N=1,141	0.035 [0.038] N=959	0.052 [0.043] N=772
Sweets consumed at least weekly	0.638 P=924	-0.012 [0.049] N=1,434	-0.015 [0.052] N=1,304	-0.034 [0.056] N=1,141	-0.042 [0.060] N=959	-0.076 [0.071] N=772
Snack foods consumed at least weekly	0.517 P=744	-0.014 [0.050] N=1,434	-0.014 [0.054] N=1,304	-0.012 [0.057] N=1,141	0.000 [0.059] N=959	-0.029 [0.068] N=772
Packaged soft drinks consumed at least weekly	0.307 P=443	-0.011 [0.045] N=1,434	-0.011 [0.047] N=1,304	0.015 [0.050] N=1,141	0.015 [0.055] N=959	-0.012 [0.064] N=772
Grains consumed at least weekly	0.888 P=1,281	**0.067* [0.039] N=1,434**	**0.078* [0.041] N=1,304**	**0.087** [0.044] N=1,141**	**0.091* [0.049] N=959**	0.081 [0.058] N=772
Child was ever breastfed	0.966 P=1,574	-0.023 [0.017] N=1,629	-0.025 [0.018] N=1,484	-0.017 [0.018] N=1,297	-0.017 [0.018] N=1,095	-0.012 [0.020] N=879

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.028 [0.039] N=1,242	-0.033 [0.042] N=1,112	-0.035 [0.049] N=949	-0.073 [0.051] N=767	-0.095 [0.068] N=580
Vegetables consumed at least weekly	-0.023 [0.043] N=1,242	-0.022 [0.047] N=1,112	-0.016 [0.053] N=949	-0.053 [0.061] N=767	-0.121 [0.075] N=580
Meat fish or poultry consumed at least weekly	0.020 [0.064] N=1,242	0.033 [0.067] N=1,112	0.025 [0.073] N=949	0.018 [0.078] N=767	-0.052 [0.097] N=580
Legumes consumed at least weekly	-0.048 [0.054] N=1,242	-0.014 [0.060] N=1,112	-0.019 [0.069] N=949	-0.042 [0.080] N=767	-0.021 [0.089] N=580
Animal products consumed at least weekly	-0.023 [0.043] N=1,242	-0.010 [0.046] N=1,112	-0.017 [0.051] N=949	-0.052 [0.054] N=767	-0.047 [0.066] N=580
Sweets consumed at least weekly	-0.008 [0.064] N=1,242	-0.015 [0.069] N=1,112	-0.055 [0.078] N=949	-0.094 [0.088] N=767	**-0.221* [0.112] N=580**
Snack foods consumed at least weekly	0.009 [0.071] N=1,242	0.006 [0.079] N=1,112	0.010 [0.089] N=949	0.033 [0.098] N=767	-0.018 [0.129] N=580
Packaged soft drinks consumed at least weekly	-0.033 [0.067] N=1,242	-0.028 [0.072] N=1,112	0.016 [0.083] N=949	0.013 [0.097] N=767	-0.058 [0.125] N=580
Grains consumed at least weekly	-0.003 [0.043] N=1,242	-0.000 [0.048] N=1,112	0.002 [0.053] N=949	-0.041 [0.060] N=767	**-0.151* [0.080] N=580**
Child was ever breastfed	-0.007 [0.031] N=1,411	-0.009 [0.034] N=1,266	0.010 [0.036] N=1,079	0.018 [0.039] N=877	0.037 [0.052] N=661

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.936 P=1,342	0.016 [0.029] N=1,434	0.017 [0.031] N=1,304	0.018 [0.034] N=1,141	-0.005 [0.034] N=959	-0.001 [0.039] N=772
Vegetables consumed at least weekly	0.900 P=1,292	0.035 [0.032] N=1,434	0.040 [0.035] N=1,304	0.052 [0.037] N=1,141	0.033 [0.041] N=959	0.020 [0.046] N=772
Meat fish or poultry consumed at least weekly	0.768 P=1,113	**0.081* [0.043] N=1,434**	**0.092** [0.045] N=1,304**	**0.099** [0.048] N=1,141**	**0.111** [0.049] N=959**	**0.104* [0.055] N=772**
Legumes consumed at least weekly	0.770 P=1,117	-0.017 [0.044] N=1,434	0.006 [0.047] N=1,304	0.001 [0.050] N=1,141	-0.024 [0.056] N=959	-0.011 [0.065] N=772
Animal products consumed at least weekly	0.912 P=1,313	0.035 [0.032] N=1,434	0.043 [0.034] N=1,304	0.044 [0.036] N=1,141	0.034 [0.038] N=959	0.052 [0.042] N=772
Sweets consumed at least weekly	0.638 P=924	-0.012 [0.049] N=1,434	-0.016 [0.052] N=1,304	-0.034 [0.057] N=1,141	-0.043 [0.061] N=959	-0.078 [0.072] N=772
Snack foods consumed at least weekly	0.517 P=744	-0.012 [0.049] N=1,434	-0.013 [0.053] N=1,304	-0.011 [0.056] N=1,141	0.001 [0.059] N=959	-0.028 [0.068] N=772
Packaged soft drinks consumed at least weekly	0.307 P=443	-0.008 [0.045] N=1,434	-0.009 [0.047] N=1,304	0.017 [0.050] N=1,141	0.015 [0.055] N=959	-0.013 [0.065] N=772
Grains consumed at least weekly	0.888 P=1,281	**0.066* [0.039] N=1,434**	**0.077* [0.041] N=1,304**	**0.086* [0.044] N=1,141**	**0.089* [0.049] N=959**	0.079 [0.058] N=772
Child was ever breastfed	0.966 P=1,574	-0.023 [0.018] N=1,629	-0.025 [0.018] N=1,484	-0.017 [0.018] N=1,297	-0.014 [0.019] N=1,095	-0.009 [0.020] N=879

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.027 [0.040] N=1,242	-0.030 [0.043] N=1,112	-0.034 [0.049] N=949	-0.076 [0.053] N=767	-0.096 [0.071] N=580
Vegetables consumed at least weekly	-0.023 [0.043] N=1,242	-0.022 [0.047] N=1,112	-0.017 [0.053] N=949	-0.057 [0.062] N=767	-0.122 [0.077] N=580
Meat fish or poultry consumed at least weekly	0.017 [0.063] N=1,242	0.029 [0.067] N=1,112	0.020 [0.073] N=949	0.014 [0.079] N=767	-0.049 [0.095] N=580
Legumes consumed at least weekly	-0.047 [0.054] N=1,242	-0.011 [0.060] N=1,112	-0.018 [0.069] N=949	-0.048 [0.080] N=767	-0.022 [0.088] N=580
Animal products consumed at least weekly	-0.023 [0.043] N=1,242	-0.010 [0.047] N=1,112	-0.018 [0.051] N=949	-0.054 [0.054] N=767	-0.051 [0.066] N=580
Sweets consumed at least weekly	-0.006 [0.064] N=1,242	-0.012 [0.069] N=1,112	-0.050 [0.078] N=949	-0.085 [0.089] N=767	**-0.210* [0.111] N=580**
Snack foods consumed at least weekly	0.010 [0.071] N=1,242	0.005 [0.079] N=1,112	0.011 [0.088] N=949	0.040 [0.098] N=767	-0.019 [0.128] N=580
Packaged soft drinks consumed at least weekly	-0.030 [0.067] N=1,242	-0.027 [0.073] N=1,112	0.021 [0.083] N=949	0.020 [0.096] N=767	-0.051 [0.124] N=580
Grains consumed at least weekly	-0.004 [0.043] N=1,242	-0.002 [0.047] N=1,112	-0.000 [0.053] N=949	-0.045 [0.060] N=767	**-0.155* [0.080] N=580**
Child was ever breastfed	-0.008 [0.032] N=1,411	-0.012 [0.034] N=1,266	0.009 [0.036] N=1,079	0.022 [0.041] N=877	0.040 [0.054] N=661

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.