Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.010	0.241 [0.212] N=1,714	0.295 [0.222] N=1,588	0.304 [0.245] N=1,426	0.372 [0.252] N=1,219	0.147 [0.277] N=1,009
WHO weight-for-age z score	0.296	-0.174 [0.114] N=1,960	-0.166 [0.124] N=1,814	-0.158 [0.130] N=1,630	-0.158 [0.135] N=1,387	**-0.256* [0.147] N=1,144**
WHO weight-for-length/height z score, 0-60 months	0.330	0.151 [0.180] N=1,692	0.119 [0.188] N=1,571	0.133 [0.202] N=1,412	0.035 [0.218] N=1,208	0.111 [0.229] N=998
WHO BMI-for-age z score	0.300	0.134 [0.197] N=1,670	0.155 [0.211] N=1,553	0.159 [0.231] N=1,394	-0.028 [0.244] N=1,188	0.068 [0.261] N=983
Stunted: HAZ below -2	0.192 P=318	-0.041 [0.039] N=1,714	-0.047 [0.040] N=1,588	-0.035 [0.042] N=1,426	-0.040 [0.044] N=1,219	-0.028 [0.050] N=1,009
Underweight: WAZ below -2	0.054 P=110	0.027 [0.019] N=1,960	0.023 [0.020] N=1,814	0.024 [0.021] N=1,630	0.011 [0.022] N=1,387	0.015 [0.029] N=1,144
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=197	-0.044 [0.032] N=1,692	-0.034 [0.033] N=1,571	-0.042 [0.035] N=1,412	-0.029 [0.038] N=1,208	-0.055 [0.039] N=998
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=328	0.015 [0.039] N=1,692	0.005 [0.042] N=1,571	0.008 [0.045] N=1,412	0.001 [0.051] N=1,208	-0.004 [0.057] N=998
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=155	0.005 [0.032] N=1,692	0.006 [0.034] N=1,571	0.018 [0.036] N=1,412	0.012 [0.041] N=1,208	0.011 [0.046] N=998
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=428	-0.053 [0.040] N=1,627	-0.044 [0.041] N=1,511	-0.031 [0.043] N=1,357	-0.042 [0.047] N=1,161	-0.041 [0.054] N=960
BMI-for-age z score below -2	0.126 P=211	-0.027 [0.034] N=1,670	-0.025 [0.036] N=1,553	-0.020 [0.041] N=1,394	0.016 [0.043] N=1,188	0.008 [0.046] N=983

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.195 [0.303] N=1,442	0.288 [0.340] N=1,316	0.377 [0.394] N=1,154	0.664 [0.427] N=947	0.236 [0.588] N=737
WHO weight-for-age z score	**-0.324* [0.169] N=1,661**	**-0.331* [0.194] N=1,515**	**-0.345* [0.208] N=1,331**	-0.406 [0.250] N=1,088	**-0.800** [0.342] N=845**
WHO weight-for-length/height z score, 0-60 months	0.089 [0.241] N=1,423	0.028 [0.261] N=1,302	-0.048 [0.298] N=1,143	-0.276 [0.344] N=939	-0.324 [0.408] N=729
WHO BMI-for-age z score	0.179 [0.245] N=1,406	0.222 [0.275] N=1,289	0.130 [0.325] N=1,130	-0.261 [0.359] N=924	-0.230 [0.444] N=719
Stunted: HAZ below -2	0.002 [0.053] N=1,442	-0.002 [0.057] N=1,316	0.021 [0.064] N=1,154	0.022 [0.074] N=947	0.088 [0.106] N=737
Underweight: WAZ below -2	**0.048* [0.026] N=1,661**	0.045 [0.029] N=1,515	0.051 [0.031] N=1,331	0.037 [0.035] N=1,088	0.068 [0.058] N=845
Wasted: WLZ/WHZ below -2, 0-60 months	-0.043 [0.044] N=1,423	-0.027 [0.047] N=1,302	-0.034 [0.053] N=1,143	-0.012 [0.065] N=939	-0.077 [0.076] N=729
Overweight: WLZ/WHZ above +2, 0-60 months	-0.002 [0.045] N=1,423	-0.023 [0.049] N=1,302	-0.035 [0.053] N=1,143	-0.060 [0.066] N=939	**-0.143* [0.081] N=729**
Obese: WLZ/WHZ above +3, 0-60 months	-0.016 [0.034] N=1,423	-0.016 [0.038] N=1,302	0.003 [0.040] N=1,143	-0.012 [0.050] N=939	-0.052 [0.059] N=729
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.004 [0.056] N=1,366	0.019 [0.060] N=1,250	0.057 [0.065] N=1,096	0.048 [0.076] N=900	0.097 [0.109] N=699
BMI-for-age z score below -2	-0.061 [0.040] N=1,406	-0.066 [0.043] N=1,289	-0.057 [0.052] N=1,130	-0.003 [0.060] N=924	-0.026 [0.069] N=719

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.010	0.238 [0.212] N=1,714	0.291 [0.222] N=1,588	0.303 [0.245] N=1,426	0.376 [0.253] N=1,219	0.155 [0.278] N=1,009
WHO weight-for-age z score	0.296	-0.181 [0.113] N=1,960	-0.174 [0.124] N=1,814	-0.164 [0.129] N=1,630	-0.165 [0.135] N=1,387	**-0.265* [0.147] N=1,144**
WHO weight-for-length/height z score, 0-60 months	0.330	0.150 [0.180] N=1,692	0.119 [0.189] N=1,571	0.131 [0.203] N=1,412	0.031 [0.218] N=1,208	0.102 [0.230] N=998
WHO BMI-for-age z score	0.300	0.134 [0.197] N=1,670	0.153 [0.211] N=1,553	0.156 [0.231] N=1,394	-0.029 [0.244] N=1,188	0.063 [0.260] N=983
Stunted: HAZ below -2	0.192 P=318	-0.040 [0.038] N=1,714	-0.046 [0.039] N=1,588	-0.034 [0.042] N=1,426	-0.039 [0.044] N=1,219	-0.028 [0.050] N=1,009
Underweight: WAZ below -2	0.054 P=110	0.028 [0.019] N=1,960	0.024 [0.020] N=1,814	0.024 [0.021] N=1,630	0.011 [0.022] N=1,387	0.016 [0.030] N=1,144
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=197	-0.042 [0.032] N=1,692	-0.032 [0.033] N=1,571	-0.039 [0.036] N=1,412	-0.025 [0.039] N=1,208	-0.050 [0.040] N=998
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=328	0.016 [0.040] N=1,692	0.006 [0.042] N=1,571	0.009 [0.045] N=1,412	0.003 [0.051] N=1,208	-0.002 [0.057] N=998
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=155	0.007 [0.032] N=1,692	0.007 [0.034] N=1,571	0.020 [0.036] N=1,412	0.014 [0.041] N=1,208	0.013 [0.045] N=998
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=428	-0.050 [0.040] N=1,627	-0.040 [0.041] N=1,511	-0.026 [0.043] N=1,357	-0.037 [0.046] N=1,161	-0.035 [0.053] N=960
BMI-for-age z score below -2	0.126 P=211	-0.026 [0.034] N=1,670	-0.023 [0.036] N=1,553	-0.018 [0.041] N=1,394	0.019 [0.043] N=1,188	0.011 [0.046] N=983

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.196 [0.303] N=1,442	0.290 [0.341] N=1,316	0.374 [0.392] N=1,154	0.650 [0.424] N=947	0.195 [0.579] N=737
WHO weight-for-age z score	**-0.318* [0.169] N=1,661**	**-0.326* [0.195] N=1,515**	-0.339 [0.209] N=1,331	-0.402 [0.251] N=1,088	**-0.789** [0.342] N=845**
WHO weight-for-length/height z score, 0-60 months	0.089 [0.241] N=1,423	0.028 [0.261] N=1,302	-0.046 [0.297] N=1,143	-0.276 [0.343] N=939	-0.316 [0.408] N=729
WHO BMI-for-age z score	0.182 [0.245] N=1,406	0.224 [0.274] N=1,289	0.134 [0.323] N=1,130	-0.258 [0.359] N=924	-0.219 [0.445] N=719
Stunted: HAZ below -2	0.002 [0.053] N=1,442	-0.003 [0.057] N=1,316	0.021 [0.063] N=1,154	0.023 [0.073] N=947	0.092 [0.105] N=737
Underweight: WAZ below -2	**0.047* [0.026] N=1,661**	0.045 [0.029] N=1,515	0.051 [0.031] N=1,331	0.037 [0.035] N=1,088	0.067 [0.058] N=845
Wasted: WLZ/WHZ below -2, 0-60 months	-0.043 [0.044] N=1,423	-0.027 [0.047] N=1,302	-0.035 [0.053] N=1,143	-0.012 [0.065] N=939	-0.079 [0.076] N=729
Overweight: WLZ/WHZ above +2, 0-60 months	-0.002 [0.045] N=1,423	-0.023 [0.049] N=1,302	-0.035 [0.053] N=1,143	-0.060 [0.067] N=939	**-0.143* [0.081] N=729**
Obese: WLZ/WHZ above +3, 0-60 months	-0.016 [0.034] N=1,423	-0.017 [0.038] N=1,302	0.002 [0.041] N=1,143	-0.012 [0.050] N=939	-0.053 [0.060] N=729
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.006 [0.055] N=1,366	0.016 [0.059] N=1,250	0.053 [0.064] N=1,096	0.045 [0.075] N=900	0.092 [0.108] N=699
BMI-for-age z score below -2	-0.062 [0.040] N=1,406	-0.067 [0.043] N=1,289	-0.058 [0.052] N=1,130	-0.004 [0.061] N=924	-0.029 [0.070] N=719

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.010	0.204 [0.212] N=1,714	0.245 [0.222] N=1,588	0.258 [0.245] N=1,426	0.346 [0.249] N=1,219	0.100 [0.272] N=1,009
WHO weight-for-age z score	0.296	**-0.224** [0.109] N=1,960**	**-0.214* [0.121] N=1,814**	-0.201 [0.126] N=1,630	-0.187 [0.133] N=1,387	**-0.297** [0.144] N=1,144**
WHO weight-for-length/height z score, 0-60 months	0.330	0.139 [0.182] N=1,692	0.119 [0.192] N=1,571	0.139 [0.203] N=1,412	0.036 [0.218] N=1,208	0.100 [0.231] N=998
WHO BMI-for-age z score	0.300	0.128 [0.199] N=1,670	0.155 [0.214] N=1,553	0.164 [0.232] N=1,394	-0.020 [0.244] N=1,188	0.069 [0.262] N=983
Stunted: HAZ below -2	0.192 P=318	-0.034 [0.039] N=1,714	-0.037 [0.040] N=1,588	-0.027 [0.042] N=1,426	-0.035 [0.044] N=1,219	-0.020 [0.050] N=1,009
Underweight: WAZ below -2	0.054 P=110	**0.031* [0.018] N=1,960**	0.028 [0.020] N=1,814	0.029 [0.020] N=1,630	0.015 [0.022] N=1,387	0.021 [0.029] N=1,144
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=197	-0.037 [0.032] N=1,692	-0.026 [0.033] N=1,571	-0.035 [0.036] N=1,412	-0.022 [0.039] N=1,208	-0.045 [0.040] N=998
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=328	0.018 [0.040] N=1,692	0.010 [0.042] N=1,571	0.012 [0.045] N=1,412	0.006 [0.050] N=1,208	0.001 [0.057] N=998
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=155	0.007 [0.032] N=1,692	0.009 [0.034] N=1,571	0.021 [0.036] N=1,412	0.015 [0.041] N=1,208	0.013 [0.046] N=998
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=428	-0.038 [0.040] N=1,627	-0.024 [0.041] N=1,511	-0.014 [0.042] N=1,357	-0.030 [0.046] N=1,161	-0.021 [0.053] N=960
BMI-for-age z score below -2	0.126 P=211	-0.022 [0.034] N=1,670	-0.018 [0.037] N=1,553	-0.015 [0.041] N=1,394	0.021 [0.043] N=1,188	0.013 [0.046] N=983

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.164 [0.304] N=1,442	0.234 [0.342] N=1,316	0.314 [0.393] N=1,154	0.632 [0.421] N=947	0.150 [0.572] N=737
WHO weight-for-age z score	**-0.356** [0.163] N=1,661**	**-0.361* [0.190] N=1,515**	**-0.368* [0.203] N=1,331**	-0.397 [0.251] N=1,088	**-0.799** [0.341] N=845**
WHO weight-for-length/height z score, 0-60 months	0.080 [0.244] N=1,423	0.039 [0.268] N=1,302	-0.026 [0.298] N=1,143	-0.275 [0.342] N=939	-0.344 [0.414] N=729
WHO BMI-for-age z score	0.180 [0.249] N=1,406	0.236 [0.282] N=1,289	0.158 [0.325] N=1,130	-0.248 [0.361] N=924	-0.218 [0.452] N=719
Stunted: HAZ below -2	0.007 [0.053] N=1,442	0.008 [0.058] N=1,316	0.029 [0.063] N=1,154	0.020 [0.072] N=947	0.093 [0.104] N=737
Underweight: WAZ below -2	**0.050* [0.026] N=1,661**	**0.049* [0.029] N=1,515**	**0.054* [0.031] N=1,331**	0.036 [0.035] N=1,088	0.067 [0.058] N=845
Wasted: WLZ/WHZ below -2, 0-60 months	-0.038 [0.044] N=1,423	-0.021 [0.048] N=1,302	-0.033 [0.054] N=1,143	-0.012 [0.066] N=939	-0.072 [0.078] N=729
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.047] N=1,423	-0.018 [0.051] N=1,302	-0.033 [0.054] N=1,143	-0.062 [0.067] N=939	**-0.139* [0.080] N=729**
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.034] N=1,423	-0.015 [0.039] N=1,302	0.003 [0.041] N=1,143	-0.014 [0.051] N=939	-0.058 [0.059] N=729
CIAF: any stunting, underweight, or wasting, 0-60 months	0.006 [0.056] N=1,366	0.033 [0.060] N=1,250	0.061 [0.063] N=1,096	0.034 [0.076] N=900	0.094 [0.107] N=699
BMI-for-age z score below -2	-0.057 [0.041] N=1,406	-0.060 [0.044] N=1,289	-0.055 [0.053] N=1,130	-0.002 [0.062] N=924	-0.022 [0.072] N=719

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.010	0.165 [0.213] N=1,714	0.210 [0.224] N=1,588	0.227 [0.247] N=1,426	0.304 [0.254] N=1,219	0.042 [0.277] N=1,009
WHO weight-for-age z score	0.296	**-0.239** [0.108] N=1,960**	**-0.239** [0.119] N=1,814**	**-0.228* [0.126] N=1,630**	-0.220 [0.133] N=1,387	**-0.347** [0.147] N=1,144**
WHO weight-for-length/height z score, 0-60 months	0.330	0.152 [0.183] N=1,692	0.124 [0.193] N=1,571	0.139 [0.206] N=1,412	0.038 [0.223] N=1,208	0.097 [0.236] N=998
WHO BMI-for-age z score	0.300	0.141 [0.199] N=1,670	0.161 [0.214] N=1,553	0.165 [0.234] N=1,394	-0.022 [0.249] N=1,188	0.059 [0.269] N=983
Stunted: HAZ below -2	0.192 P=318	-0.026 [0.039] N=1,714	-0.027 [0.040] N=1,588	-0.016 [0.043] N=1,426	-0.023 [0.044] N=1,219	-0.005 [0.050] N=1,009
Underweight: WAZ below -2	0.054 P=110	**0.034* [0.018] N=1,960**	**0.032* [0.019] N=1,814**	**0.034* [0.020] N=1,630**	0.019 [0.022] N=1,387	0.028 [0.029] N=1,144
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=197	-0.038 [0.032] N=1,692	-0.028 [0.033] N=1,571	-0.036 [0.036] N=1,412	-0.022 [0.039] N=1,208	-0.048 [0.040] N=998
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=328	0.021 [0.040] N=1,692	0.013 [0.042] N=1,571	0.014 [0.045] N=1,412	0.007 [0.051] N=1,208	-0.001 [0.057] N=998
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=155	0.007 [0.032] N=1,692	0.007 [0.034] N=1,571	0.020 [0.036] N=1,412	0.014 [0.040] N=1,208	0.011 [0.045] N=998
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=428	-0.029 [0.040] N=1,627	-0.013 [0.041] N=1,511	-0.000 [0.043] N=1,357	-0.013 [0.046] N=1,161	-0.001 [0.054] N=960
BMI-for-age z score below -2	0.126 P=211	-0.023 [0.034] N=1,670	-0.019 [0.037] N=1,553	-0.014 [0.041] N=1,394	0.023 [0.043] N=1,188	0.014 [0.047] N=983

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.152 [0.306] N=1,442	0.238 [0.341] N=1,316	0.326 [0.391] N=1,154	0.653 [0.425] N=947	0.106 [0.568] N=737
WHO weight-for-age z score	**-0.355** [0.163] N=1,661**	**-0.365* [0.188] N=1,515**	**-0.377* [0.201] N=1,331**	-0.402 [0.250] N=1,088	**-0.793** [0.339] N=845**
WHO weight-for-length/height z score, 0-60 months	0.077 [0.243] N=1,423	0.028 [0.265] N=1,302	-0.044 [0.302] N=1,143	-0.279 [0.350] N=939	-0.296 [0.421] N=729
WHO BMI-for-age z score	0.179 [0.248] N=1,406	0.228 [0.277] N=1,289	0.143 [0.328] N=1,130	-0.235 [0.366] N=924	-0.165 [0.454] N=719
Stunted: HAZ below -2	0.007 [0.054] N=1,442	0.006 [0.058] N=1,316	0.025 [0.064] N=1,154	0.012 [0.074] N=947	0.090 [0.103] N=737
Underweight: WAZ below -2	**0.049* [0.026] N=1,661**	**0.049* [0.029] N=1,515**	**0.054* [0.030] N=1,331**	0.033 [0.035] N=1,088	0.066 [0.058] N=845
Wasted: WLZ/WHZ below -2, 0-60 months	-0.036 [0.044] N=1,423	-0.022 [0.047] N=1,302	-0.031 [0.054] N=1,143	-0.007 [0.067] N=939	-0.069 [0.079] N=729
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.046] N=1,423	-0.018 [0.050] N=1,302	-0.033 [0.054] N=1,143	-0.060 [0.067] N=939	-0.128 [0.081] N=729
Obese: WLZ/WHZ above +3, 0-60 months	-0.014 [0.035] N=1,423	-0.013 [0.039] N=1,302	0.005 [0.042] N=1,143	-0.012 [0.051] N=939	-0.052 [0.058] N=729
CIAF: any stunting, underweight, or wasting, 0-60 months	0.007 [0.057] N=1,366	0.032 [0.061] N=1,250	0.061 [0.064] N=1,096	0.033 [0.078] N=900	0.097 [0.110] N=699
BMI-for-age z score below -2	-0.056 [0.041] N=1,406	-0.063 [0.043] N=1,289	-0.056 [0.053] N=1,130	-0.004 [0.062] N=924	-0.028 [0.071] N=719

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.215 [0.213] N=1,589	0.222 [0.224] N=1,477	0.208 [0.247] N=1,322	0.242 [0.258] N=1,138	-0.046 [0.282] N=941
WHO weight-for-age z score	0.298	-0.134 [0.118] N=1,813	-0.134 [0.127] N=1,683	-0.121 [0.134] N=1,508	-0.117 [0.140] N=1,290	-0.211 [0.158] N=1,066
WHO weight-for-length/height z score, 0-60 months	0.341	0.274 [0.182] N=1,571	0.249 [0.188] N=1,465	0.264 [0.199] N=1,314	0.176 [0.212] N=1,133	0.290 [0.223] N=936
WHO BMI-for-age z score	0.311	0.262 [0.202] N=1,549	0.271 [0.212] N=1,446	0.289 [0.228] N=1,296	0.127 [0.236] N=1,113	0.256 [0.252] N=921
Stunted: HAZ below -2	0.192 P=294	-0.034 [0.041] N=1,589	-0.036 [0.042] N=1,477	-0.020 [0.045] N=1,322	-0.020 [0.047] N=1,138	0.003 [0.053] N=941
Underweight: WAZ below -2	0.053 P=101	0.024 [0.019] N=1,813	0.021 [0.020] N=1,683	0.022 [0.021] N=1,508	0.012 [0.023] N=1,290	0.021 [0.030] N=1,066
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=181	**-0.059* [0.032] N=1,571**	-0.049 [0.033] N=1,465	-0.053 [0.035] N=1,314	-0.039 [0.037] N=1,133	**-0.064* [0.038] N=936**
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=305	0.033 [0.041] N=1,571	0.023 [0.042] N=1,465	0.029 [0.045] N=1,314	0.020 [0.051] N=1,133	0.025 [0.058] N=936
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=148	0.004 [0.034] N=1,571	0.007 [0.036] N=1,465	0.024 [0.038] N=1,314	0.022 [0.042] N=1,133	0.028 [0.046] N=936
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=396	-0.054 [0.043] N=1,510	-0.041 [0.044] N=1,408	-0.027 [0.046] N=1,261	-0.041 [0.050] N=1,088	-0.034 [0.058] N=899
BMI-for-age z score below -2	0.122 P=191	-0.043 [0.033] N=1,549	-0.041 [0.035] N=1,446	-0.037 [0.039] N=1,296	-0.005 [0.041] N=1,113	-0.019 [0.044] N=921

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.339 [0.309] N=1,335	0.394 [0.351] N=1,223	0.460 [0.403] N=1,068	0.720 [0.442] N=884	0.188 [0.619] N=687
WHO weight-for-age z score	**-0.313* [0.171] N=1,534**	**-0.338* [0.194] N=1,404**	**-0.359* [0.208] N=1,229**	**-0.420* [0.246] N=1,011**	**-0.811** [0.351] N=787**
WHO weight-for-length/height z score, 0-60 months	0.075 [0.255] N=1,319	-0.003 [0.268] N=1,213	-0.094 [0.302] N=1,062	-0.257 [0.362] N=881	-0.199 [0.419] N=684
WHO BMI-for-age z score	0.178 [0.264] N=1,302	0.169 [0.285] N=1,199	0.095 [0.329] N=1,049	-0.217 [0.371] N=866	-0.128 [0.456] N=674
Stunted: HAZ below -2	-0.012 [0.056] N=1,335	-0.017 [0.060] N=1,223	0.012 [0.066] N=1,068	0.013 [0.077] N=884	0.090 [0.116] N=687
Underweight: WAZ below -2	**0.046* [0.026] N=1,534**	0.046 [0.030] N=1,404	**0.055* [0.032] N=1,229**	0.050 [0.035] N=1,011	**0.095* [0.056] N=787**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.061 [0.044] N=1,319	-0.045 [0.045] N=1,213	-0.050 [0.051] N=1,062	-0.040 [0.064] N=881	**-0.122* [0.070] N=684**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.049] N=1,319	-0.029 [0.051] N=1,213	-0.038 [0.057] N=1,062	-0.064 [0.071] N=881	-0.116 [0.090] N=684
Obese: WLZ/WHZ above +3, 0-60 months	-0.040 [0.038] N=1,319	-0.039 [0.043] N=1,213	-0.018 [0.045] N=1,062	-0.023 [0.054] N=881	-0.054 [0.064] N=684
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.016 [0.059] N=1,266	0.004 [0.062] N=1,164	0.043 [0.066] N=1,017	0.011 [0.079] N=844	0.040 [0.119] N=655
BMI-for-age z score below -2	-0.057 [0.040] N=1,302	-0.060 [0.041] N=1,199	-0.051 [0.051] N=1,049	-0.009 [0.061] N=866	-0.040 [0.066] N=674

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003	0.209 [0.214] N=1,589	0.216 [0.225] N=1,477	0.201 [0.248] N=1,322	0.238 [0.257] N=1,138	-0.042 [0.279] N=941
WHO weight-for-age z score	0.298	-0.132 [0.117] N=1,813	-0.133 [0.126] N=1,683	-0.125 [0.132] N=1,508	-0.125 [0.139] N=1,290	-0.225 [0.157] N=1,066
WHO weight-for-length/height z score, 0-60 months	0.341	0.254 [0.181] N=1,571	0.230 [0.186] N=1,465	0.242 [0.199] N=1,314	0.154 [0.215] N=1,133	0.256 [0.230] N=936
WHO BMI-for-age z score	0.311	0.250 [0.200] N=1,549	0.258 [0.210] N=1,446	0.274 [0.227] N=1,296	0.111 [0.239] N=1,113	0.228 [0.260] N=921
Stunted: HAZ below -2	0.192 P=294	-0.034 [0.040] N=1,589	-0.036 [0.041] N=1,477	-0.019 [0.044] N=1,322	-0.018 [0.046] N=1,138	0.002 [0.052] N=941
Underweight: WAZ below -2	0.053 P=101	0.023 [0.019] N=1,813	0.020 [0.020] N=1,683	0.022 [0.021] N=1,508	0.013 [0.023] N=1,290	0.021 [0.030] N=1,066
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=181	**-0.057* [0.032] N=1,571**	-0.047 [0.033] N=1,465	-0.053 [0.035] N=1,314	-0.039 [0.038] N=1,133	**-0.066* [0.039] N=936**
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=305	0.030 [0.041] N=1,571	0.020 [0.042] N=1,465	0.025 [0.045] N=1,314	0.014 [0.051] N=1,133	0.017 [0.058] N=936
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=148	0.003 [0.034] N=1,571	0.006 [0.036] N=1,465	0.022 [0.038] N=1,314	0.020 [0.042] N=1,133	0.024 [0.047] N=936
CIAF: any stunting, underweight, or wasting, 0-60 months	0.267 P=396	-0.052 [0.043] N=1,510	-0.039 [0.043] N=1,408	-0.024 [0.045] N=1,261	-0.038 [0.049] N=1,088	-0.033 [0.056] N=899
BMI-for-age z score below -2	0.122 P=191	-0.040 [0.033] N=1,549	-0.039 [0.035] N=1,446	-0.036 [0.039] N=1,296	-0.004 [0.042] N=1,113	-0.019 [0.045] N=921

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.338 [0.314] N=1,335	0.385 [0.357] N=1,223	0.448 [0.408] N=1,068	0.691 [0.453] N=884	0.190 [0.635] N=687
WHO weight-for-age z score	**-0.288* [0.169] N=1,534**	-0.309 [0.192] N=1,404	**-0.340* [0.205] N=1,229**	-0.408 [0.247] N=1,011	**-0.799** [0.352] N=787**
WHO weight-for-length/height z score, 0-60 months	0.095 [0.254] N=1,319	0.028 [0.267] N=1,213	-0.068 [0.303] N=1,062	-0.265 [0.366] N=881	-0.247 [0.430] N=684
WHO BMI-for-age z score	0.196 [0.263] N=1,302	0.198 [0.285] N=1,199	0.119 [0.329] N=1,049	-0.211 [0.374] N=866	-0.142 [0.458] N=674
Stunted: HAZ below -2	-0.018 [0.056] N=1,335	-0.021 [0.060] N=1,223	0.008 [0.065] N=1,068	0.008 [0.075] N=884	0.081 [0.112] N=687
Underweight: WAZ below -2	0.042 [0.026] N=1,534	0.041 [0.029] N=1,404	0.051 [0.031] N=1,229	0.045 [0.035] N=1,011	0.088 [0.057] N=787
Wasted: WLZ/WHZ below -2, 0-60 months	-0.061 [0.044] N=1,319	-0.046 [0.046] N=1,213	-0.051 [0.051] N=1,062	-0.040 [0.065] N=881	**-0.125* [0.072] N=684**
Overweight: WLZ/WHZ above +2, 0-60 months	0.004 [0.049] N=1,319	-0.022 [0.051] N=1,213	-0.032 [0.057] N=1,062	-0.063 [0.072] N=881	-0.125 [0.090] N=684
Obese: WLZ/WHZ above +3, 0-60 months	-0.039 [0.037] N=1,319	-0.039 [0.042] N=1,213	-0.017 [0.044] N=1,062	-0.024 [0.054] N=881	-0.062 [0.063] N=684
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.022 [0.058] N=1,266	-0.002 [0.061] N=1,164	0.039 [0.064] N=1,017	0.004 [0.076] N=844	0.027 [0.112] N=655
BMI-for-age z score below -2	-0.056 [0.041] N=1,302	-0.061 [0.043] N=1,199	-0.051 [0.052] N=1,049	-0.007 [0.062] N=866	-0.041 [0.069] N=674

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062	-0.198 [0.279] N=1,534	-0.257 [0.290] N=1,390	-0.198 [0.300] N=1,212	-0.285 [0.337] N=1,028	-0.317 [0.380] N=826
WHO weight-for-age z score	0.301	0.118 [0.125] N=1,745	0.147 [0.134] N=1,590	0.052 [0.136] N=1,383	0.131 [0.145] N=1,169	0.139 [0.168] N=935
WHO weight-for-length/height z score, 0-60 months	0.365	0.091 [0.213] N=1,520	0.093 [0.221] N=1,380	0.058 [0.245] N=1,204	0.176 [0.264] N=1,019	0.120 [0.307] N=819
WHO BMI-for-age z score	0.318	0.113 [0.229] N=1,494	0.123 [0.236] N=1,356	0.065 [0.264] N=1,184	0.170 [0.288] N=1,002	0.177 [0.327] N=808
Stunted: HAZ below -2	0.184 P=283	-0.004 [0.043] N=1,534	0.002 [0.045] N=1,390	-0.024 [0.047] N=1,212	-0.008 [0.053] N=1,028	-0.018 [0.059] N=826
Underweight: WAZ below -2	0.052 P=95	0.002 [0.022] N=1,745	-0.006 [0.022] N=1,590	0.001 [0.023] N=1,383	-0.001 [0.022] N=1,169	-0.014 [0.025] N=935
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=173	0.028 [0.036] N=1,520	0.029 [0.038] N=1,380	0.042 [0.042] N=1,204	-0.004 [0.041] N=1,019	0.001 [0.045] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=304	0.062 [0.041] N=1,520	0.063 [0.044] N=1,380	0.052 [0.046] N=1,204	0.061 [0.052] N=1,019	0.055 [0.060] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=148	0.033 [0.035] N=1,520	0.026 [0.036] N=1,380	0.019 [0.039] N=1,204	0.018 [0.045] N=1,019	-0.008 [0.048] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=388	0.039 [0.044] N=1,462	0.036 [0.045] N=1,325	0.034 [0.047] N=1,156	0.003 [0.049] N=979	0.015 [0.052] N=785
BMI-for-age z score below -2	0.127 P=189	0.050 [0.042] N=1,494	0.051 [0.044] N=1,356	0.052 [0.051] N=1,184	0.026 [0.052] N=1,002	0.039 [0.058] N=808

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.113 [0.386] N=1,328	-0.207 [0.423] N=1,184	-0.112 [0.463] N=1,006	-0.385 [0.579] N=822	-0.401 [0.722] N=620
WHO weight-for-age z score	0.091 [0.177] N=1,516	0.147 [0.194] N=1,361	-0.059 [0.204] N=1,154	0.075 [0.218] N=940	0.094 [0.269] N=706
WHO weight-for-length/height z score, 0-60 months	-0.069 [0.286] N=1,316	-0.075 [0.306] N=1,176	-0.182 [0.360] N=1,000	0.076 [0.397] N=815	-0.239 [0.464] N=615
WHO BMI-for-age z score	-0.143 [0.275] N=1,297	-0.141 [0.293] N=1,159	-0.291 [0.351] N=987	-0.080 [0.391] N=805	-0.194 [0.426] N=611
Stunted: HAZ below -2	0.023 [0.068] N=1,328	0.032 [0.073] N=1,184	-0.004 [0.079] N=1,006	0.046 [0.098] N=822	0.001 [0.118] N=620
Underweight: WAZ below -2	0.006 [0.036] N=1,516	-0.008 [0.038] N=1,361	0.007 [0.042] N=1,154	0.001 [0.044] N=940	-0.029 [0.052] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	**0.080* [0.047] N=1,316**	0.084 [0.051] N=1,176	**0.122** [0.059] N=1,000**	0.033 [0.060] N=815	0.069 [0.067] N=615
Overweight: WLZ/WHZ above +2, 0-60 months	0.088 [0.056] N=1,316	0.087 [0.060] N=1,176	0.074 [0.065] N=1,000	0.112 [0.073] N=815	0.081 [0.088] N=615
Obese: WLZ/WHZ above +3, 0-60 months	0.065 [0.041] N=1,316	0.053 [0.042] N=1,176	0.046 [0.047] N=1,000	0.048 [0.061] N=815	-0.044 [0.073] N=615
CIAF: any stunting, underweight, or wasting, 0-60 months	0.097 [0.071] N=1,266	0.091 [0.075] N=1,129	0.108 [0.083] N=960	0.060 [0.095] N=783	0.072 [0.109] N=589
BMI-for-age z score below -2	**0.093** [0.042] N=1,297**	**0.095** [0.047] N=1,159**	**0.110* [0.057] N=987**	0.050 [0.056] N=805	0.094 [0.066] N=611

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062	-0.203 [0.279] N=1,534	-0.265 [0.290] N=1,390	-0.208 [0.299] N=1,212	-0.296 [0.335] N=1,028	-0.327 [0.378] N=826
WHO weight-for-age z score	0.301	0.124 [0.125] N=1,745	0.153 [0.134] N=1,590	0.056 [0.136] N=1,383	0.134 [0.145] N=1,169	0.141 [0.168] N=935
WHO weight-for-length/height z score, 0-60 months	0.365	0.095 [0.213] N=1,520	0.093 [0.222] N=1,380	0.057 [0.246] N=1,204	0.176 [0.264] N=1,019	0.121 [0.306] N=819
WHO BMI-for-age z score	0.318	0.116 [0.230] N=1,494	0.122 [0.237] N=1,356	0.063 [0.264] N=1,184	0.169 [0.288] N=1,002	0.178 [0.327] N=808
Stunted: HAZ below -2	0.184 P=283	-0.004 [0.044] N=1,534	0.003 [0.046] N=1,390	-0.023 [0.047] N=1,212	-0.007 [0.053] N=1,028	-0.017 [0.058] N=826
Underweight: WAZ below -2	0.052 P=95	0.001 [0.022] N=1,745	-0.007 [0.022] N=1,590	0.000 [0.023] N=1,383	-0.001 [0.022] N=1,169	-0.014 [0.025] N=935
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=173	0.026 [0.036] N=1,520	0.026 [0.038] N=1,380	0.039 [0.042] N=1,204	-0.004 [0.041] N=1,019	0.001 [0.045] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=304	0.061 [0.041] N=1,520	0.061 [0.044] N=1,380	0.049 [0.047] N=1,204	0.060 [0.052] N=1,019	0.055 [0.060] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=148	0.031 [0.035] N=1,520	0.024 [0.036] N=1,380	0.017 [0.039] N=1,204	0.017 [0.045] N=1,019	-0.008 [0.049] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=388	0.036 [0.045] N=1,462	0.033 [0.045] N=1,325	0.031 [0.048] N=1,156	0.002 [0.049] N=979	0.015 [0.052] N=785
BMI-for-age z score below -2	0.127 P=189	0.048 [0.042] N=1,494	0.048 [0.044] N=1,356	0.051 [0.051] N=1,184	0.025 [0.052] N=1,002	0.039 [0.058] N=808

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.122 [0.386] N=1,328	-0.224 [0.423] N=1,184	-0.142 [0.460] N=1,006	-0.438 [0.576] N=822	-0.485 [0.730] N=620
WHO weight-for-age z score	0.107 [0.175] N=1,516	0.164 [0.192] N=1,361	-0.045 [0.200] N=1,154	0.095 [0.213] N=940	0.116 [0.266] N=706
WHO weight-for-length/height z score, 0-60 months	-0.058 [0.289] N=1,316	-0.073 [0.309] N=1,176	-0.185 [0.363] N=1,000	0.078 [0.401] N=815	-0.207 [0.473] N=615
WHO BMI-for-age z score	-0.127 [0.279] N=1,297	-0.134 [0.296] N=1,159	-0.289 [0.354] N=987	-0.073 [0.394] N=805	-0.153 [0.440] N=611
Stunted: HAZ below -2	0.021 [0.068] N=1,328	0.031 [0.073] N=1,184	-0.006 [0.080] N=1,006	0.048 [0.099] N=822	0.008 [0.120] N=620
Underweight: WAZ below -2	0.005 [0.036] N=1,516	-0.010 [0.038] N=1,361	0.006 [0.042] N=1,154	0.001 [0.045] N=940	-0.029 [0.054] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,316	0.076 [0.052] N=1,176	**0.113* [0.060] N=1,000**	0.025 [0.062] N=815	0.054 [0.070] N=615
Overweight: WLZ/WHZ above +2, 0-60 months	0.085 [0.056] N=1,316	0.081 [0.060] N=1,176	0.065 [0.065] N=1,000	0.101 [0.074] N=815	0.072 [0.088] N=615
Obese: WLZ/WHZ above +3, 0-60 months	0.061 [0.041] N=1,316	0.047 [0.042] N=1,176	0.038 [0.047] N=1,000	0.037 [0.060] N=815	-0.055 [0.073] N=615
CIAF: any stunting, underweight, or wasting, 0-60 months	0.087 [0.072] N=1,266	0.079 [0.075] N=1,129	0.092 [0.084] N=960	0.048 [0.096] N=783	0.058 [0.112] N=589
BMI-for-age z score below -2	**0.087** [0.042] N=1,297**	**0.087* [0.047] N=1,159**	**0.102* [0.057] N=987**	0.042 [0.057] N=805	0.083 [0.069] N=611

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062	-0.204 [0.275] N=1,534	-0.260 [0.286] N=1,390	-0.195 [0.298] N=1,212	-0.284 [0.330] N=1,028	-0.330 [0.368] N=826
WHO weight-for-age z score	0.301	0.130 [0.124] N=1,745	0.161 [0.133] N=1,590	0.075 [0.135] N=1,383	0.146 [0.145] N=1,169	0.148 [0.168] N=935
WHO weight-for-length/height z score, 0-60 months	0.365	0.102 [0.213] N=1,520	0.100 [0.222] N=1,380	0.068 [0.247] N=1,204	0.172 [0.263] N=1,019	0.120 [0.304] N=819
WHO BMI-for-age z score	0.318	0.120 [0.231] N=1,494	0.125 [0.238] N=1,356	0.073 [0.264] N=1,184	0.167 [0.285] N=1,002	0.175 [0.322] N=808
Stunted: HAZ below -2	0.184 P=283	-0.003 [0.042] N=1,534	0.002 [0.045] N=1,390	-0.026 [0.046] N=1,212	-0.010 [0.051] N=1,028	-0.020 [0.057] N=826
Underweight: WAZ below -2	0.052 P=95	0.001 [0.022] N=1,745	-0.007 [0.022] N=1,590	-0.002 [0.023] N=1,383	-0.003 [0.022] N=1,169	-0.016 [0.025] N=935
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=173	0.025 [0.037] N=1,520	0.025 [0.038] N=1,380	0.038 [0.042] N=1,204	-0.005 [0.042] N=1,019	0.002 [0.045] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=304	0.062 [0.042] N=1,520	0.061 [0.044] N=1,380	0.048 [0.047] N=1,204	0.056 [0.052] N=1,019	0.054 [0.061] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=148	0.032 [0.035] N=1,520	0.024 [0.036] N=1,380	0.017 [0.039] N=1,204	0.015 [0.045] N=1,019	-0.008 [0.049] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=388	0.037 [0.045] N=1,462	0.033 [0.045] N=1,325	0.026 [0.048] N=1,156	-0.004 [0.050] N=979	0.009 [0.052] N=785
BMI-for-age z score below -2	0.127 P=189	0.048 [0.042] N=1,494	0.048 [0.045] N=1,356	0.049 [0.051] N=1,184	0.024 [0.053] N=1,002	0.040 [0.059] N=808

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.104 [0.380] N=1,328	-0.208 [0.416] N=1,184	-0.095 [0.448] N=1,006	-0.385 [0.559] N=822	-0.424 [0.694] N=620
WHO weight-for-age z score	0.105 [0.175] N=1,516	0.166 [0.192] N=1,361	-0.012 [0.198] N=1,154	0.115 [0.213] N=940	0.132 [0.259] N=706
WHO weight-for-length/height z score, 0-60 months	-0.070 [0.287] N=1,316	-0.081 [0.308] N=1,176	-0.185 [0.358] N=1,000	0.060 [0.393] N=815	-0.220 [0.464] N=615
WHO BMI-for-age z score	-0.140 [0.278] N=1,297	-0.141 [0.296] N=1,159	-0.286 [0.348] N=987	-0.085 [0.387] N=805	-0.152 [0.435] N=611
Stunted: HAZ below -2	0.018 [0.067] N=1,328	0.028 [0.072] N=1,184	-0.013 [0.077] N=1,006	0.037 [0.096] N=822	0.001 [0.116] N=620
Underweight: WAZ below -2	0.004 [0.036] N=1,516	-0.011 [0.038] N=1,361	0.001 [0.042] N=1,154	-0.004 [0.046] N=940	-0.031 [0.054] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,316	0.075 [0.052] N=1,176	**0.110* [0.061] N=1,000**	0.023 [0.062] N=815	0.055 [0.071] N=615
Overweight: WLZ/WHZ above +2, 0-60 months	0.085 [0.056] N=1,316	0.081 [0.060] N=1,176	0.060 [0.066] N=1,000	0.093 [0.074] N=815	0.068 [0.089] N=615
Obese: WLZ/WHZ above +3, 0-60 months	0.060 [0.041] N=1,316	0.045 [0.043] N=1,176	0.035 [0.047] N=1,000	0.032 [0.060] N=815	-0.059 [0.073] N=615
CIAF: any stunting, underweight, or wasting, 0-60 months	0.087 [0.072] N=1,266	0.078 [0.076] N=1,129	0.084 [0.085] N=960	0.037 [0.097] N=783	0.058 [0.112] N=589
BMI-for-age z score below -2	**0.087** [0.043] N=1,297**	**0.086* [0.047] N=1,159**	**0.100* [0.056] N=987**	0.043 [0.056] N=805	0.084 [0.068] N=611

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062	-0.157 [0.277] N=1,534	-0.202 [0.288] N=1,390	-0.133 [0.302] N=1,212	-0.243 [0.333] N=1,028	-0.301 [0.376] N=826
WHO weight-for-age z score	0.301	0.137 [0.126] N=1,745	0.174 [0.136] N=1,590	0.095 [0.139] N=1,383	0.164 [0.149] N=1,169	0.168 [0.173] N=935
WHO weight-for-length/height z score, 0-60 months	0.365	0.086 [0.212] N=1,520	0.087 [0.221] N=1,380	0.056 [0.244] N=1,204	0.185 [0.261] N=1,019	0.156 [0.304] N=819
WHO BMI-for-age z score	0.318	0.103 [0.229] N=1,494	0.111 [0.237] N=1,356	0.057 [0.262] N=1,184	0.169 [0.285] N=1,002	0.201 [0.326] N=808
Stunted: HAZ below -2	0.184 P=283	-0.013 [0.042] N=1,534	-0.009 [0.045] N=1,390	-0.038 [0.047] N=1,212	-0.020 [0.052] N=1,028	-0.026 [0.057] N=826
Underweight: WAZ below -2	0.052 P=95	-0.002 [0.022] N=1,745	-0.011 [0.022] N=1,590	-0.005 [0.023] N=1,383	-0.005 [0.023] N=1,169	-0.019 [0.027] N=935
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=173	0.027 [0.036] N=1,520	0.027 [0.038] N=1,380	0.041 [0.042] N=1,204	-0.005 [0.041] N=1,019	-0.002 [0.044] N=819
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=304	0.057 [0.041] N=1,520	0.057 [0.043] N=1,380	0.047 [0.046] N=1,204	0.058 [0.051] N=1,019	0.058 [0.060] N=819
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=148	0.034 [0.034] N=1,520	0.028 [0.035] N=1,380	0.019 [0.038] N=1,204	0.020 [0.044] N=1,019	-0.003 [0.048] N=819
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=388	0.031 [0.044] N=1,462	0.027 [0.045] N=1,325	0.020 [0.047] N=1,156	-0.008 [0.048] N=979	0.010 [0.051] N=785
BMI-for-age z score below -2	0.127 P=189	0.047 [0.042] N=1,494	0.048 [0.044] N=1,356	0.050 [0.050] N=1,184	0.025 [0.052] N=1,002	0.037 [0.058] N=808

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.043 [0.375] N=1,328	-0.112 [0.413] N=1,184	0.027 [0.451] N=1,006	-0.305 [0.571] N=822	-0.431 [0.721] N=620
WHO weight-for-age z score	0.142 [0.170] N=1,516	0.212 [0.188] N=1,361	0.047 [0.196] N=1,154	0.166 [0.211] N=940	0.186 [0.255] N=706
WHO weight-for-length/height z score, 0-60 months	-0.079 [0.286] N=1,316	-0.098 [0.305] N=1,176	-0.215 [0.355] N=1,000	0.056 [0.395] N=815	-0.150 [0.482] N=615
WHO BMI-for-age z score	-0.153 [0.276] N=1,297	-0.162 [0.294] N=1,159	-0.321 [0.346] N=987	-0.096 [0.386] N=805	-0.094 [0.448] N=611
Stunted: HAZ below -2	0.007 [0.066] N=1,328	0.012 [0.071] N=1,184	-0.037 [0.077] N=1,006	0.016 [0.097] N=822	-0.010 [0.117] N=620
Underweight: WAZ below -2	0.000 [0.036] N=1,516	-0.017 [0.038] N=1,361	-0.006 [0.043] N=1,154	-0.008 [0.046] N=940	-0.038 [0.056] N=706
Wasted: WLZ/WHZ below -2, 0-60 months	0.076 [0.048] N=1,316	0.081 [0.052] N=1,176	**0.121** [0.059] N=1,000**	0.030 [0.060] N=815	0.059 [0.068] N=615
Overweight: WLZ/WHZ above +2, 0-60 months	0.079 [0.056] N=1,316	0.074 [0.059] N=1,176	0.055 [0.065] N=1,000	0.089 [0.075] N=815	0.075 [0.090] N=615
Obese: WLZ/WHZ above +3, 0-60 months	0.063 [0.041] N=1,316	0.049 [0.042] N=1,176	0.037 [0.047] N=1,000	0.036 [0.061] N=815	-0.051 [0.076] N=615
CIAF: any stunting, underweight, or wasting, 0-60 months	0.079 [0.071] N=1,266	0.068 [0.075] N=1,129	0.069 [0.084] N=960	0.021 [0.096] N=783	0.052 [0.112] N=589
BMI-for-age z score below -2	**0.088** [0.043] N=1,297**	**0.089* [0.048] N=1,159**	**0.106* [0.056] N=987**	0.047 [0.055] N=805	0.086 [0.068] N=611

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.058	-0.293 [0.282] N=1,421	-0.376 [0.294] N=1,288	-0.319 [0.302] N=1,126	-0.419 [0.328] N=952	-0.520 [0.365] N=766
WHO weight-for-age z score	0.298	0.144 [0.129] N=1,614	0.184 [0.137] N=1,470	0.072 [0.139] N=1,283	0.134 [0.146] N=1,085	0.102 [0.162] N=871
WHO weight-for-length/height z score, 0-60 months	0.368	0.154 [0.223] N=1,410	0.181 [0.235] N=1,281	0.107 [0.259] N=1,119	0.203 [0.273] N=946	0.145 [0.310] N=763
WHO BMI-for-age z score	0.318	0.137 [0.247] N=1,385	0.174 [0.259] N=1,258	0.058 [0.289] N=1,101	0.137 [0.310] N=931	0.113 [0.356] N=753
Stunted: HAZ below -2	0.184 P=263	0.002 [0.045] N=1,421	0.015 [0.048] N=1,288	-0.013 [0.048] N=1,126	0.002 [0.052] N=952	0.003 [0.056] N=766
Underweight: WAZ below -2	0.052 P=88	0.003 [0.023] N=1,614	-0.006 [0.024] N=1,470	-0.001 [0.025] N=1,283	0.002 [0.025] N=1,085	-0.007 [0.028] N=871
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=160	0.028 [0.037] N=1,410	0.026 [0.039] N=1,281	0.041 [0.042] N=1,119	0.002 [0.042] N=946	0.008 [0.044] N=763
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=283	0.068 [0.043] N=1,410	0.071 [0.045] N=1,281	0.051 [0.048] N=1,119	0.056 [0.052] N=946	0.049 [0.058] N=763
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=140	0.033 [0.034] N=1,410	0.026 [0.035] N=1,281	0.019 [0.038] N=1,119	0.022 [0.044] N=946	0.001 [0.045] N=763
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=362	0.045 [0.046] N=1,356	0.043 [0.048] N=1,230	0.038 [0.048] N=1,075	0.007 [0.050] N=908	0.032 [0.052] N=730
BMI-for-age z score below -2	0.125 P=172	0.040 [0.044] N=1,385	0.036 [0.046] N=1,258	0.041 [0.053] N=1,101	0.024 [0.055] N=931	0.043 [0.062] N=753

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.022 [0.393] N=1,228	-0.148 [0.433] N=1,095	-0.033 [0.471] N=933	-0.270 [0.577] N=759	-0.318 [0.709] N=573
WHO weight-for-age z score	0.162 [0.183] N=1,399	0.227 [0.199] N=1,255	-0.008 [0.210] N=1,068	0.137 [0.224] N=870	0.115 [0.272] N=656
WHO weight-for-length/height z score, 0-60 months	-0.022 [0.310] N=1,217	-0.004 [0.338] N=1,088	-0.199 [0.393] N=926	0.018 [0.422] N=753	-0.317 [0.497] N=570
WHO BMI-for-age z score	-0.111 [0.297] N=1,198	-0.079 [0.323] N=1,071	-0.339 [0.384] N=914	-0.155 [0.410] N=744	-0.321 [0.450] N=566
Stunted: HAZ below -2	0.002 [0.074] N=1,228	0.021 [0.081] N=1,095	-0.022 [0.086] N=933	0.012 [0.105] N=759	-0.024 [0.131] N=573
Underweight: WAZ below -2	0.001 [0.039] N=1,399	-0.014 [0.042] N=1,255	-0.004 [0.047] N=1,068	-0.004 [0.052] N=870	-0.033 [0.062] N=656
Wasted: WLZ/WHZ below -2, 0-60 months	0.076 [0.050] N=1,217	0.074 [0.054] N=1,088	**0.113* [0.063] N=926**	0.024 [0.065] N=753	0.049 [0.071] N=570
Overweight: WLZ/WHZ above +2, 0-60 months	0.091 [0.061] N=1,217	0.088 [0.066] N=1,088	0.060 [0.071] N=926	0.078 [0.080] N=753	0.041 [0.095] N=570
Obese: WLZ/WHZ above +3, 0-60 months	0.052 [0.044] N=1,217	0.037 [0.046] N=1,088	0.026 [0.051] N=926	0.024 [0.065] N=753	-0.070 [0.080] N=570
CIAF: any stunting, underweight, or wasting, 0-60 months	0.084 [0.079] N=1,171	0.076 [0.084] N=1,045	0.087 [0.090] N=890	0.016 [0.107] N=723	0.025 [0.132] N=545
BMI-for-age z score below -2	**0.084* [0.045] N=1,198**	0.078 [0.051] N=1,071	0.096 [0.061] N=914	0.040 [0.061] N=744	0.095 [0.072] N=566

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.058	-0.302 [0.285] N=1,421	-0.381 [0.297] N=1,288	-0.320 [0.304] N=1,126	-0.438 [0.329] N=952	-0.539 [0.368] N=766
WHO weight-for-age z score	0.298	0.144 [0.128] N=1,614	0.182 [0.137] N=1,470	0.072 [0.138] N=1,283	0.134 [0.145] N=1,085	0.104 [0.163] N=871
WHO weight-for-length/height z score, 0-60 months	0.368	0.164 [0.225] N=1,410	0.186 [0.237] N=1,281	0.113 [0.260] N=1,119	0.215 [0.272] N=946	0.158 [0.316] N=763
WHO BMI-for-age z score	0.318	0.145 [0.250] N=1,385	0.177 [0.260] N=1,258	0.061 [0.291] N=1,101	0.148 [0.310] N=931	0.128 [0.359] N=753
Stunted: HAZ below -2	0.184 P=263	0.002 [0.045] N=1,421	0.013 [0.048] N=1,288	-0.015 [0.048] N=1,126	0.001 [0.052] N=952	0.002 [0.056] N=766
Underweight: WAZ below -2	0.052 P=88	0.002 [0.023] N=1,614	-0.006 [0.024] N=1,470	-0.002 [0.025] N=1,283	-0.000 [0.024] N=1,085	-0.010 [0.027] N=871
Wasted: WLZ/WHZ below -2, 0-60 months	0.113 P=160	0.025 [0.037] N=1,410	0.023 [0.039] N=1,281	0.037 [0.042] N=1,119	-0.004 [0.042] N=946	0.001 [0.045] N=763
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=283	0.067 [0.043] N=1,410	0.068 [0.045] N=1,281	0.048 [0.047] N=1,119	0.054 [0.052] N=946	0.048 [0.059] N=763
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=140	0.032 [0.034] N=1,410	0.025 [0.035] N=1,281	0.018 [0.038] N=1,119	0.021 [0.043] N=946	-0.001 [0.044] N=763
CIAF: any stunting, underweight, or wasting, 0-60 months	0.266 P=362	0.039 [0.046] N=1,356	0.036 [0.047] N=1,230	0.031 [0.047] N=1,075	-0.002 [0.049] N=908	0.022 [0.050] N=730
BMI-for-age z score below -2	0.125 P=172	0.037 [0.044] N=1,385	0.034 [0.046] N=1,258	0.038 [0.053] N=1,101	0.019 [0.054] N=931	0.037 [0.061] N=753

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.032 [0.394] N=1,228	-0.147 [0.432] N=1,095	-0.031 [0.467] N=933	-0.336 [0.573] N=759	-0.438 [0.707] N=573
WHO weight-for-age z score	0.156 [0.183] N=1,399	0.221 [0.201] N=1,255	-0.012 [0.210] N=1,068	0.129 [0.222] N=870	0.096 [0.270] N=656
WHO weight-for-length/height z score, 0-60 months	-0.033 [0.310] N=1,217	-0.024 [0.335] N=1,088	-0.218 [0.389] N=926	0.022 [0.417] N=753	-0.298 [0.488] N=570
WHO BMI-for-age z score	-0.109 [0.296] N=1,198	-0.087 [0.319] N=1,071	-0.344 [0.379] N=914	-0.125 [0.406] N=744	-0.245 [0.450] N=566
Stunted: HAZ below -2	0.003 [0.073] N=1,228	0.019 [0.080] N=1,095	-0.023 [0.085] N=933	0.019 [0.105] N=759	-0.009 [0.129] N=573
Underweight: WAZ below -2	0.003 [0.040] N=1,399	-0.012 [0.042] N=1,255	-0.002 [0.047] N=1,068	-0.001 [0.052] N=870	-0.025 [0.060] N=656
Wasted: WLZ/WHZ below -2, 0-60 months	0.078 [0.050] N=1,217	0.078 [0.054] N=1,088	**0.116* [0.062] N=926**	0.024 [0.064] N=753	0.051 [0.070] N=570
Overweight: WLZ/WHZ above +2, 0-60 months	0.089 [0.060] N=1,217	0.085 [0.065] N=1,088	0.054 [0.070] N=926	0.077 [0.079] N=753	0.044 [0.093] N=570
Obese: WLZ/WHZ above +3, 0-60 months	0.050 [0.043] N=1,217	0.034 [0.045] N=1,088	0.023 [0.051] N=926	0.023 [0.064] N=753	-0.066 [0.077] N=570
CIAF: any stunting, underweight, or wasting, 0-60 months	0.083 [0.079] N=1,171	0.076 [0.084] N=1,045	0.088 [0.090] N=890	0.018 [0.106] N=723	0.041 [0.128] N=545
BMI-for-age z score below -2	**0.085* [0.044] N=1,198**	0.081 [0.050] N=1,071	0.098 [0.060] N=914	0.038 [0.059] N=744	0.089 [0.070] N=566

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.