Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Youngest Child

Age reference: 12 November 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.098 [0.203] N=1,681	0.110 [0.212] N=1,556	0.097 [0.234] N=1,398	0.195 [0.238] N=1,201	-0.003 [0.263] N=992
WHO weight-for-age z score	0.017	-0.160 [0.117] N=1,930	-0.140 [0.128] N=1,786	-0.147 [0.134] N=1,606	-0.160 [0.142] N=1,371	**-0.272* [0.155] N=1,128**
WHO weight-for-length/height z score, 0-60 months	0.326	0.185 [0.181] N=1,664	0.163 [0.189] N=1,545	0.176 [0.202] N=1,389	0.075 [0.217] N=1,192	0.146 [0.232] N=982
WHO BMI-for-age z score	0.290	0.183 [0.203] N=1,643	0.182 [0.216] N=1,527	0.199 [0.235] N=1,371	0.024 [0.250] N=1,175	0.104 [0.273] N=970
Stunted: HAZ below -2	0.236 P=381	-0.026 [0.040] N=1,681	-0.037 [0.042] N=1,556	-0.029 [0.044] N=1,398	-0.057 [0.047] N=1,201	-0.039 [0.050] N=992
Underweight: WAZ below -2	0.082 P=156	**0.038* [0.023] N=1,930**	0.030 [0.024] N=1,786	0.027 [0.025] N=1,606	0.028 [0.028] N=1,371	0.044 [0.032] N=1,128
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=194	-0.043 [0.032] N=1,664	-0.034 [0.033] N=1,545	-0.042 [0.035] N=1,389	-0.031 [0.039] N=1,192	-0.047 [0.040] N=982
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=321	0.028 [0.041] N=1,664	0.020 [0.044] N=1,545	0.021 [0.046] N=1,389	0.010 [0.052] N=1,192	0.001 [0.059] N=982
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=150	0.007 [0.030] N=1,664	0.008 [0.032] N=1,545	0.021 [0.034] N=1,389	0.015 [0.039] N=1,192	0.024 [0.044] N=982
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=511	-0.051 [0.043] N=1,606	-0.056 [0.045] N=1,490	-0.050 [0.047] N=1,339	-0.073 [0.052] N=1,150	-0.062 [0.059] N=949
BMI-for-age z score below -2	0.128 P=213	-0.039 [0.035] N=1,643	-0.041 [0.037] N=1,527	-0.039 [0.040] N=1,371	-0.017 [0.044] N=1,175	-0.016 [0.047] N=970

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.076 [0.299] N=1,417	0.103 [0.327] N=1,292	0.146 [0.388] N=1,134	0.458 [0.416] N=937	0.121 [0.571] N=728
WHO weight-for-age z score	**-0.292* [0.170] N=1,637**	-0.273 [0.195] N=1,493	-0.307 [0.210] N=1,313	-0.388 [0.256] N=1,078	**-0.788** [0.347] N=835**
WHO weight-for-length/height z score, 0-60 months	0.127 [0.256] N=1,401	0.072 [0.276] N=1,282	0.027 [0.313] N=1,126	-0.191 [0.368] N=929	-0.215 [0.421] N=719
WHO BMI-for-age z score	0.226 [0.268] N=1,384	0.213 [0.300] N=1,268	0.179 [0.352] N=1,112	-0.192 [0.396] N=916	-0.140 [0.477] N=711
Stunted: HAZ below -2	0.003 [0.059] N=1,417	-0.016 [0.065] N=1,292	-0.008 [0.072] N=1,134	-0.070 [0.086] N=937	-0.041 [0.116] N=728
Underweight: WAZ below -2	0.046 [0.033] N=1,637	0.029 [0.036] N=1,493	0.025 [0.038] N=1,313	0.035 [0.044] N=1,078	0.083 [0.066] N=835
Wasted: WLZ/WHZ below -2, 0-60 months	-0.044 [0.044] N=1,401	-0.026 [0.047] N=1,282	-0.037 [0.053] N=1,126	-0.020 [0.066] N=929	-0.064 [0.076] N=719
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.048] N=1,401	-0.010 [0.052] N=1,282	-0.023 [0.056] N=1,126	-0.052 [0.069] N=929	**-0.141* [0.082] N=719**
Obese: WLZ/WHZ above +3, 0-60 months	-0.023 [0.034] N=1,401	-0.024 [0.039] N=1,282	-0.008 [0.040] N=1,126	-0.025 [0.049] N=929	-0.040 [0.061] N=719
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.019 [0.061] N=1,351	-0.027 [0.067] N=1,235	-0.014 [0.071] N=1,084	-0.067 [0.085] N=895	-0.053 [0.116] N=694
BMI-for-age z score below -2	-0.062 [0.044] N=1,384	-0.070 [0.046] N=1,268	-0.071 [0.054] N=1,112	-0.043 [0.064] N=916	-0.051 [0.074] N=711

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.096 [0.203] N=1,681	0.107 [0.211] N=1,556	0.095 [0.234] N=1,398	0.196 [0.238] N=1,201	0.000 [0.263] N=992
WHO weight-for-age z score	0.017	-0.165 [0.117] N=1,930	-0.147 [0.128] N=1,786	-0.153 [0.134] N=1,606	-0.166 [0.142] N=1,371	**-0.280* [0.154] N=1,128**
WHO weight-for-length/height z score, 0-60 months	0.326	0.184 [0.182] N=1,664	0.161 [0.190] N=1,545	0.173 [0.202] N=1,389	0.071 [0.218] N=1,192	0.137 [0.233] N=982
WHO BMI-for-age z score	0.290	0.183 [0.203] N=1,643	0.181 [0.216] N=1,527	0.196 [0.235] N=1,371	0.022 [0.251] N=1,175	0.098 [0.273] N=970
Stunted: HAZ below -2	0.236 P=381	-0.025 [0.040] N=1,681	-0.035 [0.041] N=1,556	-0.028 [0.043] N=1,398	-0.056 [0.047] N=1,201	-0.038 [0.050] N=992
Underweight: WAZ below -2	0.082 P=156	**0.039* [0.023] N=1,930**	0.031 [0.024] N=1,786	0.028 [0.025] N=1,606	0.028 [0.027] N=1,371	0.045 [0.033] N=1,128
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=194	-0.041 [0.032] N=1,664	-0.032 [0.033] N=1,545	-0.039 [0.035] N=1,389	-0.027 [0.039] N=1,192	-0.043 [0.041] N=982
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=321	0.029 [0.041] N=1,664	0.021 [0.043] N=1,545	0.022 [0.046] N=1,389	0.011 [0.052] N=1,192	0.003 [0.059] N=982
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=150	0.007 [0.030] N=1,664	0.010 [0.032] N=1,545	0.022 [0.034] N=1,389	0.016 [0.039] N=1,192	0.025 [0.044] N=982
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=511	-0.049 [0.043] N=1,606	-0.052 [0.044] N=1,490	-0.046 [0.047] N=1,339	-0.069 [0.052] N=1,150	-0.056 [0.059] N=949
BMI-for-age z score below -2	0.128 P=213	-0.038 [0.035] N=1,643	-0.040 [0.037] N=1,527	-0.038 [0.040] N=1,371	-0.015 [0.044] N=1,175	-0.014 [0.047] N=970

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.075 [0.299] N=1,417	0.104 [0.327] N=1,292	0.142 [0.386] N=1,134	0.447 [0.415] N=937	0.101 [0.569] N=728
WHO weight-for-age z score	**-0.287* [0.169] N=1,637**	-0.269 [0.196] N=1,493	-0.300 [0.211] N=1,313	-0.385 [0.258] N=1,078	**-0.780** [0.346] N=835**
WHO weight-for-length/height z score, 0-60 months	0.128 [0.255] N=1,401	0.073 [0.275] N=1,282	0.029 [0.312] N=1,126	-0.191 [0.367] N=929	-0.212 [0.419] N=719
WHO BMI-for-age z score	0.228 [0.267] N=1,384	0.214 [0.299] N=1,268	0.184 [0.349] N=1,112	-0.190 [0.395] N=916	-0.136 [0.475] N=711
Stunted: HAZ below -2	0.001 [0.059] N=1,417	-0.019 [0.065] N=1,292	-0.011 [0.071] N=1,134	-0.072 [0.086] N=937	-0.041 [0.116] N=728
Underweight: WAZ below -2	0.045 [0.032] N=1,637	0.029 [0.035] N=1,493	0.025 [0.038] N=1,313	0.035 [0.044] N=1,078	0.083 [0.066] N=835
Wasted: WLZ/WHZ below -2, 0-60 months	-0.044 [0.044] N=1,401	-0.026 [0.047] N=1,282	-0.038 [0.053] N=1,126	-0.020 [0.066] N=929	-0.065 [0.076] N=719
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.048] N=1,401	-0.011 [0.052] N=1,282	-0.023 [0.056] N=1,126	-0.052 [0.069] N=929	**-0.141* [0.082] N=719**
Obese: WLZ/WHZ above +3, 0-60 months	-0.024 [0.034] N=1,401	-0.025 [0.039] N=1,282	-0.008 [0.040] N=1,126	-0.025 [0.050] N=929	-0.040 [0.061] N=719
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.022 [0.060] N=1,351	-0.031 [0.066] N=1,235	-0.020 [0.070] N=1,084	-0.072 [0.084] N=895	-0.058 [0.116] N=694
BMI-for-age z score below -2	-0.063 [0.044] N=1,384	-0.071 [0.045] N=1,268	-0.072 [0.053] N=1,112	-0.043 [0.064] N=916	-0.052 [0.074] N=711

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.071 [0.202] N=1,681	0.068 [0.211] N=1,556	0.054 [0.233] N=1,398	0.164 [0.234] N=1,201	-0.055 [0.253] N=992
WHO weight-for-age z score	0.017	**-0.207* [0.112] N=1,930**	-0.188 [0.124] N=1,786	-0.193 [0.130] N=1,606	-0.193 [0.138] N=1,371	**-0.316** [0.150] N=1,128**
WHO weight-for-length/height z score, 0-60 months	0.326	0.173 [0.184] N=1,664	0.163 [0.193] N=1,545	0.182 [0.204] N=1,389	0.076 [0.218] N=1,192	0.133 [0.235] N=982
WHO BMI-for-age z score	0.290	0.180 [0.206] N=1,643	0.187 [0.220] N=1,527	0.206 [0.237] N=1,371	0.030 [0.252] N=1,175	0.099 [0.275] N=970
Stunted: HAZ below -2	0.236 P=381	-0.017 [0.039] N=1,681	-0.023 [0.041] N=1,556	-0.017 [0.043] N=1,398	-0.049 [0.045] N=1,201	-0.025 [0.048] N=992
Underweight: WAZ below -2	0.082 P=156	**0.045** [0.022] N=1,930**	0.038 [0.023] N=1,786	0.035 [0.024] N=1,606	0.034 [0.027] N=1,371	0.052 [0.032] N=1,128
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=194	-0.037 [0.032] N=1,664	-0.026 [0.033] N=1,545	-0.035 [0.035] N=1,389	-0.024 [0.039] N=1,192	-0.039 [0.041] N=982
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=321	0.031 [0.041] N=1,664	0.026 [0.044] N=1,545	0.026 [0.046] N=1,389	0.014 [0.052] N=1,192	0.005 [0.059] N=982
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=150	0.007 [0.031] N=1,664	0.011 [0.033] N=1,545	0.023 [0.034] N=1,389	0.017 [0.039] N=1,192	0.024 [0.044] N=982
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=511	-0.032 [0.042] N=1,606	-0.032 [0.044] N=1,490	-0.030 [0.046] N=1,339	-0.058 [0.051] N=1,150	-0.038 [0.058] N=949
BMI-for-age z score below -2	0.128 P=213	-0.034 [0.035] N=1,643	-0.035 [0.037] N=1,527	-0.034 [0.041] N=1,371	-0.013 [0.043] N=1,175	-0.012 [0.047] N=970

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.053 [0.299] N=1,417	0.055 [0.327] N=1,292	0.084 [0.384] N=1,134	0.416 [0.408] N=937	0.041 [0.554] N=728
WHO weight-for-age z score	**-0.323** [0.163] N=1,637**	-0.308 [0.191] N=1,493	-0.334 [0.206] N=1,313	-0.390 [0.258] N=1,078	**-0.794** [0.346] N=835**
WHO weight-for-length/height z score, 0-60 months	0.121 [0.258] N=1,401	0.083 [0.283] N=1,282	0.051 [0.313] N=1,126	-0.196 [0.365] N=929	-0.236 [0.423] N=719
WHO BMI-for-age z score	0.230 [0.273] N=1,384	0.231 [0.309] N=1,268	0.209 [0.353] N=1,112	-0.182 [0.396] N=916	-0.122 [0.482] N=711
Stunted: HAZ below -2	0.006 [0.058] N=1,417	-0.005 [0.064] N=1,292	0.000 [0.071] N=1,134	-0.075 [0.083] N=937	-0.036 [0.111] N=728
Underweight: WAZ below -2	0.050 [0.032] N=1,637	0.035 [0.035] N=1,493	0.030 [0.037] N=1,313	0.033 [0.043] N=1,078	0.083 [0.066] N=835
Wasted: WLZ/WHZ below -2, 0-60 months	-0.039 [0.045] N=1,401	-0.020 [0.048] N=1,282	-0.036 [0.053] N=1,126	-0.018 [0.068] N=929	-0.057 [0.078] N=719
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.049] N=1,401	-0.005 [0.054] N=1,282	-0.020 [0.057] N=1,126	-0.053 [0.069] N=929	**-0.135* [0.081] N=719**
Obese: WLZ/WHZ above +3, 0-60 months	-0.024 [0.035] N=1,401	-0.023 [0.040] N=1,282	-0.007 [0.041] N=1,126	-0.026 [0.050] N=929	-0.044 [0.061] N=719
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.008 [0.059] N=1,351	-0.008 [0.065] N=1,235	-0.006 [0.068] N=1,084	-0.078 [0.083] N=895	-0.049 [0.110] N=694
BMI-for-age z score below -2	-0.058 [0.044] N=1,384	-0.064 [0.046] N=1,268	-0.069 [0.054] N=1,112	-0.042 [0.064] N=916	-0.049 [0.076] N=711

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.366	0.035 [0.202] N=1,681	0.044 [0.212] N=1,556	0.037 [0.235] N=1,398	0.141 [0.239] N=1,201	-0.080 [0.258] N=992
WHO weight-for-age z score	0.017	**-0.223** [0.111] N=1,930**	**-0.214* [0.122] N=1,786**	**-0.218* [0.130] N=1,606**	-0.225 [0.139] N=1,371	**-0.365** [0.153] N=1,128**
WHO weight-for-length/height z score, 0-60 months	0.326	0.183 [0.184] N=1,664	0.161 [0.193] N=1,545	0.177 [0.205] N=1,389	0.078 [0.221] N=1,192	0.129 [0.238] N=982
WHO BMI-for-age z score	0.290	0.187 [0.206] N=1,643	0.185 [0.219] N=1,527	0.203 [0.238] N=1,371	0.033 [0.256] N=1,175	0.096 [0.281] N=970
Stunted: HAZ below -2	0.236 P=381	-0.010 [0.040] N=1,681	-0.016 [0.041] N=1,556	-0.010 [0.044] N=1,398	-0.039 [0.047] N=1,201	-0.015 [0.049] N=992
Underweight: WAZ below -2	0.082 P=156	**0.048** [0.022] N=1,930**	**0.041* [0.024] N=1,786**	0.039 [0.025] N=1,606	0.040 [0.028] N=1,371	**0.058* [0.033] N=1,128**
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=194	-0.038 [0.032] N=1,664	-0.027 [0.033] N=1,545	-0.035 [0.035] N=1,389	-0.025 [0.039] N=1,192	-0.041 [0.041] N=982
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=321	0.033 [0.041] N=1,664	0.028 [0.044] N=1,545	0.027 [0.047] N=1,389	0.016 [0.052] N=1,192	0.003 [0.059] N=982
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=150	0.007 [0.030] N=1,664	0.008 [0.032] N=1,545	0.020 [0.034] N=1,389	0.015 [0.039] N=1,192	0.021 [0.044] N=982
CIAF: any stunting, underweight, or wasting, 0-60 months	0.327 P=511	-0.024 [0.043] N=1,606	-0.022 [0.044] N=1,490	-0.018 [0.047] N=1,339	-0.043 [0.052] N=1,150	-0.023 [0.059] N=949
BMI-for-age z score below -2	0.128 P=213	-0.034 [0.035] N=1,643	-0.034 [0.037] N=1,527	-0.032 [0.041] N=1,371	-0.010 [0.044] N=1,175	-0.010 [0.048] N=970

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.030 [0.298] N=1,417	0.057 [0.325] N=1,292	0.098 [0.382] N=1,134	0.440 [0.410] N=937	0.013 [0.544] N=728
WHO weight-for-age z score	**-0.324** [0.164] N=1,637**	**-0.315* [0.188] N=1,493**	**-0.343* [0.204] N=1,313**	-0.388 [0.257] N=1,078	**-0.775** [0.343] N=835**
WHO weight-for-length/height z score, 0-60 months	0.122 [0.256] N=1,401	0.073 [0.279] N=1,282	0.034 [0.317] N=1,126	-0.189 [0.375] N=929	-0.192 [0.428] N=719
WHO BMI-for-age z score	0.238 [0.270] N=1,384	0.227 [0.302] N=1,268	0.202 [0.354] N=1,112	-0.161 [0.403] N=916	-0.080 [0.483] N=711
Stunted: HAZ below -2	0.010 [0.059] N=1,417	-0.006 [0.065] N=1,292	-0.002 [0.072] N=1,134	-0.078 [0.085] N=937	-0.035 [0.112] N=728
Underweight: WAZ below -2	0.050 [0.032] N=1,637	0.036 [0.034] N=1,493	0.032 [0.036] N=1,313	0.034 [0.043] N=1,078	0.085 [0.064] N=835
Wasted: WLZ/WHZ below -2, 0-60 months	-0.038 [0.044] N=1,401	-0.021 [0.047] N=1,282	-0.035 [0.053] N=1,126	-0.014 [0.068] N=929	-0.056 [0.079] N=719
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.049] N=1,401	-0.005 [0.053] N=1,282	-0.020 [0.057] N=1,126	-0.050 [0.070] N=929	-0.127 [0.082] N=719
Obese: WLZ/WHZ above +3, 0-60 months	-0.020 [0.035] N=1,401	-0.021 [0.039] N=1,282	-0.006 [0.041] N=1,126	-0.024 [0.050] N=929	-0.039 [0.059] N=719
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.004 [0.060] N=1,351	-0.008 [0.066] N=1,235	-0.005 [0.069] N=1,084	-0.076 [0.085] N=895	-0.045 [0.114] N=694
BMI-for-age z score below -2	-0.060 [0.044] N=1,384	-0.068 [0.046] N=1,268	-0.072 [0.055] N=1,112	-0.045 [0.064] N=916	-0.058 [0.074] N=711

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.058 [0.205] N=1,558	0.035 [0.218] N=1,448	-0.003 [0.240] N=1,297	0.080 [0.242] N=1,123	-0.165 [0.261] N=927
WHO weight-for-age z score	0.019	-0.107 [0.120] N=1,785	-0.100 [0.129] N=1,657	-0.101 [0.137] N=1,486	-0.104 [0.143] N=1,276	-0.197 [0.160] N=1,052
WHO weight-for-length/height z score, 0-60 months	0.334	**0.325* [0.184] N=1,544**	0.305 [0.190] N=1,440	0.323 [0.201] N=1,292	0.233 [0.215] N=1,118	0.334 [0.229] N=921
WHO BMI-for-age z score	0.301	0.312 [0.210] N=1,524	0.298 [0.219] N=1,422	0.334 [0.235] N=1,275	0.188 [0.247] N=1,102	0.307 [0.269] N=910
Stunted: HAZ below -2	0.238 P=354	-0.015 [0.042] N=1,558	-0.024 [0.044] N=1,448	-0.014 [0.046] N=1,297	-0.039 [0.049] N=1,123	-0.011 [0.053] N=927
Underweight: WAZ below -2	0.082 P=144	0.032 [0.024] N=1,785	0.025 [0.025] N=1,657	0.022 [0.027] N=1,486	0.025 [0.029] N=1,276	0.042 [0.035] N=1,052
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=178	**-0.061* [0.032] N=1,544**	-0.050 [0.032] N=1,440	-0.055 [0.034] N=1,292	-0.043 [0.037] N=1,118	-0.057 [0.038] N=921
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	0.046 [0.042] N=1,544	0.039 [0.044] N=1,440	0.043 [0.048] N=1,292	0.030 [0.054] N=1,118	0.030 [0.062] N=921
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=143	0.009 [0.032] N=1,544	0.013 [0.035] N=1,440	0.030 [0.037] N=1,292	0.025 [0.040] N=1,118	0.038 [0.045] N=921
CIAF: any stunting, underweight, or wasting, 0-60 months	0.329 P=474	-0.052 [0.046] N=1,489	-0.053 [0.048] N=1,388	-0.048 [0.051] N=1,244	-0.075 [0.056] N=1,078	-0.059 [0.063] N=889
BMI-for-age z score below -2	0.126 P=193	-0.047 [0.036] N=1,524	-0.047 [0.037] N=1,422	-0.047 [0.040] N=1,275	-0.027 [0.043] N=1,102	-0.031 [0.046] N=910

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.172 [0.303] N=1,311	0.170 [0.344] N=1,201	0.191 [0.409] N=1,050	0.515 [0.439] N=876	0.121 [0.619] N=680
WHO weight-for-age z score	**-0.283* [0.168] N=1,512**	-0.295 [0.191] N=1,384	-0.335 [0.206] N=1,213	-0.402 [0.247] N=1,003	**-0.757** [0.342] N=779**
WHO weight-for-length/height z score, 0-60 months	0.143 [0.264] N=1,298	0.067 [0.280] N=1,194	0.010 [0.316] N=1,046	-0.154 [0.374] N=872	-0.058 [0.424] N=675
WHO BMI-for-age z score	0.224 [0.285] N=1,282	0.164 [0.310] N=1,180	0.147 [0.357] N=1,033	-0.165 [0.400] N=860	-0.054 [0.483] N=668
Stunted: HAZ below -2	-0.013 [0.063] N=1,311	-0.037 [0.069] N=1,201	-0.028 [0.076] N=1,050	-0.097 [0.092] N=876	-0.052 [0.127] N=680
Underweight: WAZ below -2	0.040 [0.033] N=1,512	0.027 [0.037] N=1,384	0.026 [0.039] N=1,213	0.044 [0.044] N=1,003	0.099 [0.068] N=779
Wasted: WLZ/WHZ below -2, 0-60 months	-0.065 [0.045] N=1,298	-0.045 [0.046] N=1,194	-0.055 [0.051] N=1,046	-0.046 [0.065] N=872	-0.107 [0.071] N=675
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.050] N=1,298	-0.013 [0.053] N=1,194	-0.023 [0.059] N=1,046	-0.051 [0.071] N=872	-0.109 [0.089] N=675
Obese: WLZ/WHZ above +3, 0-60 months	-0.040 [0.038] N=1,298	-0.039 [0.044] N=1,194	-0.018 [0.046] N=1,046	-0.029 [0.054] N=872	-0.033 [0.066] N=675
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.044 [0.063] N=1,251	-0.053 [0.068] N=1,150	-0.050 [0.074] N=1,006	-0.128 [0.087] N=840	-0.121 [0.123] N=651
BMI-for-age z score below -2	-0.059 [0.043] N=1,282	-0.064 [0.044] N=1,180	-0.067 [0.052] N=1,033	-0.047 [0.063] N=860	-0.073 [0.069] N=668

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.057 [0.205] N=1,558	0.035 [0.217] N=1,448	-0.008 [0.239] N=1,297	0.073 [0.243] N=1,123	-0.175 [0.262] N=927
WHO weight-for-age z score	0.019	-0.110 [0.119] N=1,785	-0.103 [0.128] N=1,657	-0.112 [0.136] N=1,486	-0.122 [0.143] N=1,276	-0.223 [0.162] N=1,052
WHO weight-for-length/height z score, 0-60 months	0.334	**0.307* [0.183] N=1,544**	0.288 [0.188] N=1,440	0.303 [0.199] N=1,292	0.213 [0.217] N=1,118	0.312 [0.235] N=921
WHO BMI-for-age z score	0.301	0.305 [0.207] N=1,524	0.289 [0.217] N=1,422	0.323 [0.234] N=1,275	0.177 [0.249] N=1,102	0.294 [0.275] N=910
Stunted: HAZ below -2	0.238 P=354	-0.014 [0.042] N=1,558	-0.022 [0.044] N=1,448	-0.012 [0.046] N=1,297	-0.038 [0.050] N=1,123	-0.009 [0.052] N=927
Underweight: WAZ below -2	0.082 P=144	0.032 [0.024] N=1,785	0.025 [0.025] N=1,657	0.024 [0.026] N=1,486	0.027 [0.029] N=1,276	0.045 [0.035] N=1,052
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=178	**-0.059* [0.032] N=1,544**	-0.048 [0.032] N=1,440	-0.055 [0.034] N=1,292	-0.044 [0.038] N=1,118	-0.062 [0.039] N=921
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=297	0.045 [0.042] N=1,544	0.037 [0.044] N=1,440	0.040 [0.047] N=1,292	0.025 [0.054] N=1,118	0.025 [0.062] N=921
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=143	0.007 [0.032] N=1,544	0.012 [0.035] N=1,440	0.028 [0.037] N=1,292	0.024 [0.040] N=1,118	0.036 [0.045] N=921
CIAF: any stunting, underweight, or wasting, 0-60 months	0.329 P=474	-0.048 [0.046] N=1,489	-0.049 [0.048] N=1,388	-0.045 [0.051] N=1,244	-0.072 [0.056] N=1,078	-0.054 [0.062] N=889
BMI-for-age z score below -2	0.126 P=193	-0.044 [0.035] N=1,524	-0.045 [0.037] N=1,422	-0.046 [0.041] N=1,275	-0.027 [0.044] N=1,102	-0.032 [0.047] N=910

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.176 [0.308] N=1,311	0.167 [0.349] N=1,201	0.183 [0.413] N=1,050	0.487 [0.454] N=876	0.118 [0.637] N=680
WHO weight-for-age z score	-0.248 [0.168] N=1,512	-0.258 [0.190] N=1,384	-0.313 [0.205] N=1,213	-0.402 [0.250] N=1,003	**-0.785** [0.356] N=779**
WHO weight-for-length/height z score, 0-60 months	0.169 [0.262] N=1,298	0.101 [0.279] N=1,194	0.038 [0.317] N=1,046	-0.161 [0.380] N=872	-0.114 [0.435] N=675
WHO BMI-for-age z score	0.246 [0.284] N=1,282	0.186 [0.310] N=1,180	0.165 [0.357] N=1,033	-0.161 [0.404] N=860	-0.070 [0.486] N=668
Stunted: HAZ below -2	-0.012 [0.062] N=1,311	-0.034 [0.068] N=1,201	-0.027 [0.075] N=1,050	-0.096 [0.090] N=876	-0.060 [0.124] N=680
Underweight: WAZ below -2	0.036 [0.033] N=1,512	0.022 [0.036] N=1,384	0.022 [0.038] N=1,213	0.040 [0.044] N=1,003	0.092 [0.068] N=779
Wasted: WLZ/WHZ below -2, 0-60 months	-0.064 [0.045] N=1,298	-0.045 [0.047] N=1,194	-0.055 [0.051] N=1,046	-0.046 [0.066] N=872	-0.111 [0.071] N=675
Overweight: WLZ/WHZ above +2, 0-60 months	0.017 [0.050] N=1,298	-0.005 [0.053] N=1,194	-0.017 [0.059] N=1,046	-0.050 [0.073] N=872	-0.120 [0.088] N=675
Obese: WLZ/WHZ above +3, 0-60 months	-0.039 [0.038] N=1,298	-0.037 [0.043] N=1,194	-0.016 [0.045] N=1,046	-0.030 [0.054] N=872	-0.042 [0.065] N=675
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.041 [0.063] N=1,251	-0.050 [0.068] N=1,150	-0.047 [0.073] N=1,006	-0.127 [0.086] N=840	-0.132 [0.119] N=651
BMI-for-age z score below -2	-0.056 [0.044] N=1,282	-0.061 [0.045] N=1,180	-0.065 [0.053] N=1,033	-0.045 [0.064] N=860	-0.073 [0.070] N=668

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.307	-0.187 [0.276] N=1,499	-0.206 [0.286] N=1,360	-0.138 [0.296] N=1,188	-0.191 [0.328] N=1,009	-0.326 [0.370] N=818
WHO weight-for-age z score	0.030	0.137 [0.128] N=1,709	0.175 [0.137] N=1,558	0.073 [0.139] N=1,359	0.169 [0.148] N=1,152	0.160 [0.171] N=927
WHO weight-for-length/height z score, 0-60 months	0.362	0.162 [0.211] N=1,487	0.132 [0.220] N=1,351	0.068 [0.243] N=1,181	0.214 [0.261] N=1,002	0.186 [0.303] N=811
WHO BMI-for-age z score	0.307	0.152 [0.222] N=1,464	0.143 [0.228] N=1,329	0.065 [0.257] N=1,163	0.214 [0.284] N=987	0.213 [0.317] N=801
Stunted: HAZ below -2	0.228 P=338	0.034 [0.049] N=1,499	0.026 [0.050] N=1,360	0.015 [0.052] N=1,188	0.024 [0.058] N=1,009	0.029 [0.065] N=818
Underweight: WAZ below -2	0.079 P=132	-0.004 [0.027] N=1,709	-0.008 [0.028] N=1,558	-0.002 [0.029] N=1,359	0.001 [0.031] N=1,152	-0.006 [0.034] N=927
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=170	0.022 [0.037] N=1,487	0.026 [0.039] N=1,351	0.046 [0.043] N=1,181	-0.005 [0.042] N=1,002	-0.006 [0.047] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=296	0.067 [0.041] N=1,487	0.064 [0.043] N=1,351	0.053 [0.045] N=1,181	0.060 [0.050] N=1,002	0.058 [0.057] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=142	0.039 [0.036] N=1,487	0.028 [0.036] N=1,351	0.025 [0.039] N=1,181	0.020 [0.045] N=1,002	-0.006 [0.047] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=452	0.056 [0.050] N=1,434	0.047 [0.051] N=1,301	0.063 [0.054] N=1,136	0.024 [0.053] N=964	0.046 [0.061] N=780
BMI-for-age z score below -2	0.128 P=188	0.050 [0.042] N=1,464	0.054 [0.045] N=1,329	0.057 [0.052] N=1,163	0.020 [0.052] N=987	0.025 [0.057] N=801

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.061 [0.407] N=1,295	-0.067 [0.436] N=1,156	0.089 [0.476] N=984	-0.034 [0.569] N=805	-0.286 [0.747] N=614
WHO weight-for-age z score	0.115 [0.186] N=1,481	0.177 [0.205] N=1,330	-0.020 [0.212] N=1,131	0.147 [0.217] N=924	0.118 [0.281] N=699
WHO weight-for-length/height z score, 0-60 months	0.037 [0.291] N=1,283	-0.034 [0.306] N=1,147	-0.212 [0.356] N=977	0.065 [0.386] N=798	-0.120 [0.450] N=607
WHO BMI-for-age z score	-0.061 [0.279] N=1,268	-0.096 [0.285] N=1,133	-0.270 [0.346] N=967	0.031 [0.399] N=791	-0.024 [0.395] N=605
Stunted: HAZ below -2	0.036 [0.077] N=1,295	0.018 [0.080] N=1,156	-0.001 [0.086] N=984	0.023 [0.104] N=805	0.035 [0.130] N=614
Underweight: WAZ below -2	0.013 [0.042] N=1,481	0.007 [0.046] N=1,330	0.024 [0.050] N=1,131	0.037 [0.056] N=924	0.032 [0.067] N=699
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,283	0.085 [0.052] N=1,147	**0.138** [0.059]** **N=977**	0.045 [0.057] N=798	0.061 [0.067] N=607
Overweight: WLZ/WHZ above +2, 0-60 months	**0.105** [0.057]* **N=1,283**	0.096 [0.060] N=1,147	0.084 [0.066] N=977	0.121 [0.075] N=798	0.115 [0.089] N=607
Obese: WLZ/WHZ above +3, 0-60 months	**0.077** [0.043]* **N=1,283**	0.055 [0.043] N=1,147	0.053 [0.049] N=977	0.046 [0.063] N=798	-0.049 [0.074] N=607
CIAF: any stunting, underweight, or wasting, 0-60 months	0.098 [0.078] N=1,238	0.081 [0.084] N=1,105	0.124 [0.093] N=940	0.058 [0.100] N=768	0.112 [0.127] N=584
BMI-for-age z score below -2	**0.107** [0.044]** **N=1,268**	**0.116** [0.048]** **N=1,133**	**0.136** [0.058]** **N=967**	0.058 [0.056] N=791	0.065 [0.064] N=605

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.307	-0.190 [0.276] N=1,499	-0.211 [0.285] N=1,360	-0.146 [0.295] N=1,188	-0.197 [0.326] N=1,009	-0.332 [0.368] N=818
WHO weight-for-age z score	0.030	0.142 [0.129] N=1,709	0.179 [0.137] N=1,558	0.076 [0.138] N=1,359	0.171 [0.148] N=1,152	0.161 [0.171] N=927
WHO weight-for-length/height z score, 0-60 months	0.362	0.167 [0.211] N=1,487	0.134 [0.220] N=1,351	0.068 [0.243] N=1,181	0.214 [0.261] N=1,002	0.186 [0.302] N=811
WHO BMI-for-age z score	0.307	0.153 [0.222] N=1,464	0.142 [0.228] N=1,329	0.062 [0.257] N=1,163	0.212 [0.284] N=987	0.213 [0.317] N=801
Stunted: HAZ below -2	0.228 P=338	0.032 [0.049] N=1,499	0.025 [0.050] N=1,360	0.014 [0.052] N=1,188	0.024 [0.058] N=1,009	0.030 [0.065] N=818
Underweight: WAZ below -2	0.079 P=132	-0.005 [0.027] N=1,709	-0.009 [0.029] N=1,558	-0.003 [0.029] N=1,359	-0.000 [0.031] N=1,152	-0.006 [0.034] N=927
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=170	0.019 [0.037] N=1,487	0.023 [0.039] N=1,351	0.043 [0.043] N=1,181	-0.006 [0.042] N=1,002	-0.006 [0.047] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=296	0.066 [0.041] N=1,487	0.062 [0.043] N=1,351	0.051 [0.045] N=1,181	0.059 [0.051] N=1,002	0.058 [0.057] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=142	0.038 [0.036] N=1,487	0.026 [0.036] N=1,351	0.023 [0.039] N=1,181	0.019 [0.045] N=1,002	-0.006 [0.048] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=452	0.052 [0.051] N=1,434	0.042 [0.052] N=1,301	0.059 [0.054] N=1,136	0.023 [0.054] N=964	0.046 [0.061] N=780
BMI-for-age z score below -2	0.128 P=188	0.049 [0.042] N=1,464	0.053 [0.045] N=1,329	0.056 [0.052] N=1,163	0.020 [0.052] N=987	0.025 [0.057] N=801

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.064 [0.407] N=1,295	-0.076 [0.436] N=1,156	0.066 [0.473] N=984	-0.066 [0.568] N=805	-0.347 [0.755] N=614
WHO weight-for-age z score	0.128 [0.185] N=1,481	0.189 [0.203] N=1,330	-0.011 [0.209] N=1,131	0.162 [0.214] N=924	0.138 [0.277] N=699
WHO weight-for-length/height z score, 0-60 months	0.052 [0.293] N=1,283	-0.027 [0.309] N=1,147	-0.207 [0.358] N=977	0.073 [0.390] N=798	-0.087 [0.461] N=607
WHO BMI-for-age z score	-0.049 [0.281] N=1,268	-0.086 [0.288] N=1,133	-0.264 [0.346] N=967	0.040 [0.400] N=791	0.013 [0.409] N=605
Stunted: HAZ below -2	0.033 [0.077] N=1,295	0.013 [0.080] N=1,156	-0.004 [0.086] N=984	0.024 [0.105] N=805	0.042 [0.132] N=614
Underweight: WAZ below -2	0.010 [0.042] N=1,481	0.005 [0.046] N=1,330	0.022 [0.050] N=1,131	0.033 [0.056] N=924	0.029 [0.067] N=699
Wasted: WLZ/WHZ below -2, 0-60 months	0.067 [0.049] N=1,283	0.076 [0.053] N=1,147	**0.127** [0.060] N=977**	0.036 [0.058] N=798	0.046 [0.070] N=607
Overweight: WLZ/WHZ above +2, 0-60 months	**0.103* [0.057] N=1,283**	0.091 [0.060] N=1,147	0.076 [0.065] N=977	0.111 [0.075] N=798	0.107 [0.089] N=607
Obese: WLZ/WHZ above +3, 0-60 months	**0.073* [0.043] N=1,283**	0.049 [0.043] N=1,147	0.045 [0.048] N=977	0.037 [0.062] N=798	-0.060 [0.074] N=607
CIAF: any stunting, underweight, or wasting, 0-60 months	0.085 [0.080] N=1,238	0.065 [0.085] N=1,105	0.107 [0.094] N=940	0.045 [0.102] N=768	0.097 [0.129] N=584
BMI-for-age z score below -2	**0.102** [0.044] N=1,268**	**0.109** [0.050] N=1,133**	**0.129** [0.059] N=967**	0.052 [0.057] N=791	0.057 [0.067] N=605

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.307	-0.189 [0.269] N=1,499	-0.207 [0.279] N=1,360	-0.138 [0.292] N=1,188	-0.185 [0.320] N=1,009	-0.338 [0.358] N=818
WHO weight-for-age z score	0.030	0.149 [0.128] N=1,709	0.188 [0.136] N=1,558	0.096 [0.138] N=1,359	0.187 [0.149] N=1,152	0.170 [0.171] N=927
WHO weight-for-length/height z score, 0-60 months	0.362	0.178 [0.211] N=1,487	0.144 [0.220] N=1,351	0.088 [0.244] N=1,181	0.216 [0.259] N=1,002	0.187 [0.301] N=811
WHO BMI-for-age z score	0.307	0.159 [0.223] N=1,464	0.148 [0.229] N=1,329	0.082 [0.257] N=1,163	0.218 [0.282] N=987	0.217 [0.315] N=801
Stunted: HAZ below -2	0.228 P=338	0.033 [0.047] N=1,499	0.024 [0.049] N=1,360	0.012 [0.051] N=1,188	0.019 [0.055] N=1,009	0.028 [0.062] N=818
Underweight: WAZ below -2	0.079 P=132	-0.005 [0.027] N=1,709	-0.010 [0.029] N=1,558	-0.005 [0.029] N=1,359	-0.002 [0.031] N=1,152	-0.007 [0.034] N=927
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=170	0.018 [0.037] N=1,487	0.021 [0.039] N=1,351	0.041 [0.044] N=1,181	-0.007 [0.042] N=1,002	-0.005 [0.047] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=296	0.068 [0.041] N=1,487	0.063 [0.043] N=1,351	0.051 [0.046] N=1,181	0.055 [0.051] N=1,002	0.058 [0.057] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=142	0.039 [0.036] N=1,487	0.027 [0.036] N=1,351	0.024 [0.039] N=1,181	0.018 [0.045] N=1,002	-0.007 [0.048] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=452	0.054 [0.050] N=1,434	0.042 [0.051] N=1,301	0.054 [0.054] N=1,136	0.016 [0.054] N=964	0.041 [0.059] N=780
BMI-for-age z score below -2	0.128 P=188	0.048 [0.042] N=1,464	0.052 [0.045] N=1,329	0.053 [0.052] N=1,163	0.018 [0.053] N=987	0.026 [0.058] N=801

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.047 [0.396] N=1,295	-0.064 [0.424] N=1,156	0.091 [0.455] N=984	-0.020 [0.543] N=805	-0.297 [0.716] N=614
WHO weight-for-age z score	0.122 [0.186] N=1,481	0.185 [0.204] N=1,330	0.021 [0.209] N=1,131	0.185 [0.217] N=924	0.153 [0.271] N=699
WHO weight-for-length/height z score, 0-60 months	0.040 [0.290] N=1,283	-0.033 [0.306] N=1,147	-0.191 [0.353] N=977	0.058 [0.381] N=798	-0.103 [0.451] N=607
WHO BMI-for-age z score	-0.065 [0.278] N=1,268	-0.098 [0.285] N=1,133	-0.256 [0.339] N=967	0.013 [0.387] N=791	-0.008 [0.398] N=605
Stunted: HAZ below -2	0.030 [0.075] N=1,295	0.011 [0.078] N=1,156	-0.009 [0.082] N=984	0.011 [0.099] N=805	0.035 [0.124] N=614
Underweight: WAZ below -2	0.010 [0.042] N=1,481	0.004 [0.046] N=1,330	0.015 [0.050] N=1,131	0.027 [0.056] N=924	0.025 [0.067] N=699
Wasted: WLZ/WHZ below -2, 0-60 months	0.067 [0.049] N=1,283	0.076 [0.053] N=1,147	**0.124** [0.060]** **N=977**	0.034 [0.058] N=798	0.047 [0.071] N=607
Overweight: WLZ/WHZ above +2, 0-60 months	**0.103* [0.058]** **N=1,283**	0.092 [0.060] N=1,147	0.075 [0.066] N=977	0.102 [0.076] N=798	0.103 [0.090] N=607
Obese: WLZ/WHZ above +3, 0-60 months	**0.072* [0.043]** **N=1,283**	0.047 [0.044] N=1,147	0.045 [0.049] N=977	0.032 [0.063] N=798	-0.063 [0.074] N=607
CIAF: any stunting, underweight, or wasting, 0-60 months	0.089 [0.079] N=1,238	0.068 [0.085] N=1,105	0.099 [0.095] N=940	0.033 [0.101] N=768	0.092 [0.123] N=584
BMI-for-age z score below -2	**0.100** [0.045]** **N=1,268**	**0.107** [0.050]** **N=1,133**	**0.124** [0.058]** **N=967**	0.049 [0.057] N=791	0.055 [0.068] N=605

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.307	-0.164 [0.272] N=1,499	-0.170 [0.282] N=1,360	-0.088 [0.295] N=1,188	-0.151 [0.323] N=1,009	-0.314 [0.362] N=818
WHO weight-for-age z score	0.030	0.157 [0.130] N=1,709	0.198 [0.139] N=1,558	0.113 [0.142] N=1,359	0.202 [0.153] N=1,152	0.185 [0.176] N=927
WHO weight-for-length/height z score, 0-60 months	0.362	0.169 [0.209] N=1,487	0.134 [0.219] N=1,351	0.074 [0.241] N=1,181	0.225 [0.258] N=1,002	0.218 [0.301] N=811
WHO BMI-for-age z score	0.307	0.150 [0.222] N=1,464	0.137 [0.228] N=1,329	0.063 [0.255] N=1,163	0.214 [0.282] N=987	0.231 [0.317] N=801
Stunted: HAZ below -2	0.228 P=338	0.027 [0.048] N=1,499	0.017 [0.050] N=1,360	0.002 [0.052] N=1,188	0.013 [0.057] N=1,009	0.025 [0.064] N=818
Underweight: WAZ below -2	0.079 P=132	-0.009 [0.027] N=1,709	-0.014 [0.029] N=1,558	-0.009 [0.030] N=1,359	-0.005 [0.032] N=1,152	-0.012 [0.036] N=927
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=170	0.018 [0.037] N=1,487	0.023 [0.039] N=1,351	0.044 [0.043] N=1,181	-0.008 [0.042] N=1,002	-0.009 [0.046] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=296	0.063 [0.041] N=1,487	0.058 [0.042] N=1,351	0.049 [0.045] N=1,181	0.056 [0.049] N=1,002	0.060 [0.056] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=142	0.042 [0.035] N=1,487	0.030 [0.035] N=1,351	0.026 [0.038] N=1,181	0.023 [0.044] N=1,002	-0.001 [0.046] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=452	0.048 [0.050] N=1,434	0.037 [0.051] N=1,301	0.048 [0.054] N=1,136	0.012 [0.053] N=964	0.042 [0.059] N=780
BMI-for-age z score below -2	0.128 P=188	0.045 [0.042] N=1,464	0.049 [0.045] N=1,329	0.052 [0.051] N=1,163	0.016 [0.052] N=987	0.020 [0.056] N=801

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.014 [0.393] N=1,295	-0.002 [0.422] N=1,156	0.195 [0.455] N=984	0.048 [0.552] N=805	-0.296 [0.735] N=614
WHO weight-for-age z score	0.165 [0.180] N=1,481	0.234 [0.199] N=1,330	0.081 [0.205] N=1,131	0.237 [0.211] N=924	0.201 [0.260] N=699
WHO weight-for-length/height z score, 0-60 months	0.042 [0.288] N=1,283	-0.045 [0.303] N=1,147	-0.223 [0.348] N=977	0.061 [0.382] N=798	-0.039 [0.466] N=607
WHO BMI-for-age z score	-0.059 [0.277] N=1,268	-0.103 [0.284] N=1,133	-0.280 [0.338] N=967	0.027 [0.393] N=791	0.055 [0.418] N=605
Stunted: HAZ below -2	0.023 [0.074] N=1,295	-0.001 [0.078] N=1,156	-0.030 [0.083] N=984	-0.004 [0.101] N=805	0.033 [0.127] N=614
Underweight: WAZ below -2	0.003 [0.042] N=1,481	-0.002 [0.046] N=1,330	0.007 [0.051] N=1,131	0.023 [0.057] N=924	0.018 [0.069] N=699
Wasted: WLZ/WHZ below -2, 0-60 months	0.068 [0.048] N=1,283	0.079 [0.053] N=1,147	**0.132** [0.059] N=977**	0.038 [0.057] N=798	0.050 [0.068] N=607
Overweight: WLZ/WHZ above +2, 0-60 months	**0.098* [0.057] N=1,283**	0.085 [0.059] N=1,147	0.069 [0.064] N=977	0.101 [0.076] N=798	0.110 [0.091] N=607
Obese: WLZ/WHZ above +3, 0-60 months	**0.076* [0.042] N=1,283**	0.052 [0.043] N=1,147	0.047 [0.048] N=977	0.038 [0.063] N=798	-0.053 [0.076] N=607
CIAF: any stunting, underweight, or wasting, 0-60 months	0.078 [0.078] N=1,238	0.055 [0.084] N=1,105	0.081 [0.094] N=940	0.017 [0.099] N=768	0.095 [0.123] N=584
BMI-for-age z score below -2	**0.100** [0.045] N=1,268**	**0.107** [0.050] N=1,133**	**0.128** [0.058] N=967**	0.052 [0.057] N=791	0.058 [0.068] N=605

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.310	-0.235 [0.279] N=1,388	-0.270 [0.291] N=1,260	-0.175 [0.298] N=1,104	-0.229 [0.321] N=936	-0.409 [0.358] N=760
WHO weight-for-age z score	0.027	0.151 [0.132] N=1,580	0.198 [0.141] N=1,440	0.087 [0.141] N=1,261	0.167 [0.148] N=1,070	0.111 [0.166] N=864
WHO weight-for-length/height z score, 0-60 months	0.361	0.201 [0.221] N=1,378	0.187 [0.232] N=1,253	0.075 [0.254] N=1,097	0.192 [0.269] N=930	0.151 [0.303] N=755
WHO BMI-for-age z score	0.307	0.159 [0.237] N=1,357	0.172 [0.246] N=1,233	0.032 [0.275] N=1,082	0.160 [0.302] N=918	0.121 [0.339] N=747
Stunted: HAZ below -2	0.229 P=314	0.043 [0.052] N=1,388	0.039 [0.054] N=1,260	0.024 [0.055] N=1,104	0.025 [0.058] N=936	0.035 [0.065] N=760
Underweight: WAZ below -2	0.078 P=122	-0.002 [0.028] N=1,580	-0.006 [0.030] N=1,440	-0.006 [0.030] N=1,261	0.001 [0.031] N=1,070	0.003 [0.035] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=157	0.022 [0.038] N=1,378	0.023 [0.040] N=1,253	0.047 [0.043] N=1,097	0.002 [0.042] N=930	0.005 [0.046] N=755
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=274	0.066 [0.042] N=1,378	0.062 [0.043] N=1,253	0.043 [0.045] N=1,097	0.044 [0.049] N=930	0.039 [0.054] N=755
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=134	0.037 [0.035] N=1,378	0.024 [0.035] N=1,253	0.018 [0.038] N=1,097	0.016 [0.043] N=930	-0.008 [0.044] N=755
CIAF: any stunting, underweight, or wasting, 0-60 months	0.321 P=420	0.068 [0.053] N=1,328	0.057 [0.055] N=1,206	0.074 [0.057] N=1,055	0.030 [0.056] N=894	0.064 [0.062] N=725
BMI-for-age z score below -2	0.127 P=171	0.054 [0.044] N=1,357	0.055 [0.047] N=1,233	0.062 [0.053] N=1,082	0.035 [0.054] N=918	0.047 [0.060] N=747

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.038 [0.416] N=1,195	0.014 [0.450] N=1,067	0.217 [0.483] N=911	0.166 [0.559] N=743	-0.106 [0.732] N=567
WHO weight-for-age z score	0.188 [0.192] N=1,365	0.264 [0.210] N=1,225	0.036 [0.220] N=1,046	0.222 [0.224] N=855	0.141 [0.294] N=649
WHO weight-for-length/height z score, 0-60 months	0.093 [0.317] N=1,185	0.042 [0.340] N=1,060	-0.227 [0.389] N=904	0.004 [0.414] N=737	-0.213 [0.485] N=562
WHO BMI-for-age z score	-0.034 [0.298] N=1,170	-0.041 [0.315] N=1,046	-0.328 [0.376] N=895	-0.055 [0.423] N=731	-0.177 [0.427] N=560
Stunted: HAZ below -2	0.023 [0.084] N=1,195	0.009 [0.088] N=1,067	-0.018 [0.093] N=911	-0.019 [0.109] N=743	-0.016 [0.143] N=567
Underweight: WAZ below -2	0.007 [0.044] N=1,365	0.002 [0.048] N=1,225	0.009 [0.053] N=1,046	0.027 [0.060] N=855	0.023 [0.073] N=649
Wasted: WLZ/WHZ below -2, 0-60 months	0.068 [0.051] N=1,185	0.073 [0.056] N=1,060	**0.127** [0.062] N=904**	0.036 [0.061] N=737	0.045 [0.071] N=562
Overweight: WLZ/WHZ above +2, 0-60 months	**0.107* [0.062] N=1,185**	0.094 [0.065] N=1,060	0.066 [0.070] N=904	0.083 [0.080] N=737	0.067 [0.095] N=562
Obese: WLZ/WHZ above +3, 0-60 months	0.068 [0.047] N=1,185	0.043 [0.048] N=1,060	0.034 [0.053] N=904	0.026 [0.067] N=737	-0.073 [0.083] N=562
CIAF: any stunting, underweight, or wasting, 0-60 months	0.085 [0.087] N=1,143	0.063 [0.094] N=1,021	0.102 [0.103] N=870	0.005 [0.114] N=709	0.049 [0.147] N=540
BMI-for-age z score below -2	**0.109** [0.047] N=1,170**	**0.113** [0.053] N=1,046**	**0.138** [0.063] N=895**	0.073 [0.061] N=731	0.083 [0.069] N=560

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.310	-0.254 [0.282] N=1,388	-0.287 [0.295] N=1,260	-0.187 [0.300] N=1,104	-0.252 [0.323] N=936	-0.439 [0.362] N=760
WHO weight-for-age z score	0.027	0.152 [0.132] N=1,580	0.195 [0.142] N=1,440	0.087 [0.141] N=1,261	0.166 [0.148] N=1,070	0.111 [0.166] N=864
WHO weight-for-length/height z score, 0-60 months	0.361	0.215 [0.222] N=1,378	0.198 [0.234] N=1,253	0.088 [0.255] N=1,097	0.205 [0.266] N=930	0.165 [0.308] N=755
WHO BMI-for-age z score	0.307	0.171 [0.239] N=1,357	0.182 [0.248] N=1,233	0.042 [0.277] N=1,082	0.169 [0.302] N=918	0.139 [0.344] N=747
Stunted: HAZ below -2	0.229 P=314	0.041 [0.051] N=1,388	0.037 [0.053] N=1,260	0.021 [0.054] N=1,104	0.024 [0.058] N=936	0.035 [0.066] N=760
Underweight: WAZ below -2	0.078 P=122	-0.004 [0.027] N=1,580	-0.008 [0.029] N=1,440	-0.007 [0.030] N=1,261	-0.000 [0.031] N=1,070	0.000 [0.034] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=157	0.017 [0.038] N=1,378	0.019 [0.040] N=1,253	0.041 [0.043] N=1,097	-0.005 [0.042] N=930	-0.004 [0.046] N=755
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=274	0.065 [0.041] N=1,378	0.061 [0.042] N=1,253	0.041 [0.044] N=1,097	0.041 [0.048] N=930	0.037 [0.054] N=755
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=134	0.036 [0.035] N=1,378	0.024 [0.035] N=1,253	0.017 [0.037] N=1,097	0.015 [0.042] N=930	-0.010 [0.044] N=755
CIAF: any stunting, underweight, or wasting, 0-60 months	0.321 P=420	0.062 [0.053] N=1,328	0.049 [0.054] N=1,206	0.065 [0.056] N=1,055	0.019 [0.055] N=894	0.053 [0.062] N=725
BMI-for-age z score below -2	0.127 P=171	0.050 [0.044] N=1,357	0.051 [0.047] N=1,233	0.057 [0.054] N=1,082	0.029 [0.053] N=918	0.040 [0.058] N=747

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.014 [0.418] N=1,195	-0.005 [0.450] N=1,067	0.197 [0.480] N=911	0.091 [0.557] N=743	-0.239 [0.727] N=567
WHO weight-for-age z score	0.178 [0.193] N=1,365	0.252 [0.213] N=1,225	0.028 [0.222] N=1,046	0.206 [0.224] N=855	0.119 [0.291] N=649
WHO weight-for-length/height z score, 0-60 months	0.084 [0.316] N=1,185	0.029 [0.338] N=1,060	-0.239 [0.385] N=904	0.003 [0.407] N=737	-0.200 [0.475] N=562
WHO BMI-for-age z score	-0.028 [0.297] N=1,170	-0.039 [0.311] N=1,046	-0.324 [0.371] N=895	-0.029 [0.419] N=731	-0.110 [0.424] N=560
Stunted: HAZ below -2	0.025 [0.083] N=1,195	0.010 [0.087] N=1,067	-0.015 [0.092] N=911	-0.007 [0.109] N=743	0.012 [0.140] N=567
Underweight: WAZ below -2	0.007 [0.044] N=1,365	0.002 [0.048] N=1,225	0.010 [0.052] N=1,046	0.031 [0.059] N=855	0.037 [0.070] N=649
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.051] N=1,185	0.074 [0.056] N=1,060	**0.128** [0.062] N=904**	0.035 [0.060] N=737	0.048 [0.070] N=562
Overweight: WLZ/WHZ above +2, 0-60 months	**0.105* [0.061] N=1,185**	0.091 [0.064] N=1,060	0.062 [0.069] N=904	0.081 [0.079] N=737	0.071 [0.094] N=562
Obese: WLZ/WHZ above +3, 0-60 months	0.066 [0.046] N=1,185	0.041 [0.047] N=1,060	0.032 [0.052] N=904	0.025 [0.066] N=737	-0.070 [0.079] N=562
CIAF: any stunting, underweight, or wasting, 0-60 months	0.087 [0.087] N=1,143	0.065 [0.094] N=1,021	0.105 [0.102] N=870	0.012 [0.113] N=709	0.076 [0.143] N=540
BMI-for-age z score below -2	**0.109** [0.046] N=1,170**	**0.113** [0.053] N=1,046**	**0.138** [0.062] N=895**	0.070 [0.060] N=731	0.082 [0.067] N=560

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.