Table 9 Reduced Form Effects on Weekly Nutrition for Ages 0-3: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,527	-0.012 [0.023] N=1,639	0.002 [0.023] N=1,509	-0.002 [0.024] N=1,361	-0.007 [0.027] N=1,168	-0.020 [0.032] N=957
Vegetables consumed at least weekly	0.895 P=1,466	0.010 [0.028] N=1,639	0.010 [0.028] N=1,509	-0.007 [0.029] N=1,361	-0.010 [0.030] N=1,168	-0.042 [0.035] N=957
Meat fish or poultry consumed at least weekly	0.770 P=1,272	0.069 [0.050] N=1,639	**0.087* [0.053] N=1,509**	0.078 [0.057] N=1,361	0.053 [0.064] N=1,168	0.046 [0.075] N=957
Legumes consumed at least weekly	0.770 P=1,269	-0.027 [0.042] N=1,639	-0.039 [0.044] N=1,509	-0.032 [0.045] N=1,361	-0.042 [0.049] N=1,168	-0.078 [0.057] N=957
Animal products consumed at least weekly	0.908 P=1,492	0.002 [0.030] N=1,639	0.002 [0.031] N=1,509	-0.013 [0.034] N=1,361	-0.011 [0.037] N=1,168	-0.016 [0.041] N=957
Sweets consumed at least weekly	0.608 P=1,001	-0.032 [0.040] N=1,639	-0.022 [0.042] N=1,509	-0.004 [0.044] N=1,361	-0.023 [0.047] N=1,168	0.008 [0.056] N=957
Snack foods consumed at least weekly	0.494 P=810	-0.038 [0.041] N=1,639	-0.032 [0.043] N=1,509	-0.017 [0.047] N=1,361	-0.021 [0.049] N=1,168	-0.021 [0.053] N=957
Packaged soft drinks consumed at least weekly	0.291 P=473	-0.012 [0.049] N=1,639	-0.008 [0.052] N=1,509	-0.009 [0.056] N=1,361	-0.006 [0.061] N=1,168	-0.009 [0.072] N=957
Grains consumed at least weekly	0.880 P=1,448	0.023 [0.032] N=1,639	0.037 [0.033] N=1,509	0.023 [0.034] N=1,361	0.031 [0.037] N=1,168	0.010 [0.044] N=957
Child was ever breastfed	0.967 P=1,834	-0.004 [0.014] N=1,901	0.000 [0.015] N=1,756	0.000 [0.015] N=1,580	0.003 [0.017] N=1,344	0.001 [0.016] N=1,104

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.037] N=1,391	0.019 [0.037] N=1,261	0.015 [0.041] N=1,113	0.008 [0.050] N=920	-0.018 [0.068] N=709
Vegetables consumed at least weekly	0.032 [0.045] N=1,391	0.038 [0.047] N=1,261	0.014 [0.051] N=1,113	0.011 [0.059] N=920	-0.065 [0.077] N=709
Meat fish or poultry consumed at least weekly	0.063 [0.057] N=1,391	0.091 [0.059] N=1,261	0.077 [0.068] N=1,113	-0.006 [0.071] N=920	-0.061 [0.086] N=709
Legumes consumed at least weekly	0.034 [0.063] N=1,391	0.020 [0.068] N=1,261	0.053 [0.070] N=1,113	0.054 [0.084] N=920	-0.003 [0.120] N=709
Animal products consumed at least weekly	0.001 [0.041] N=1,391	0.003 [0.044] N=1,261	-0.030 [0.050] N=1,113	-0.025 [0.059] N=920	-0.049 [0.074] N=709
Sweets consumed at least weekly	-0.036 [0.057] N=1,391	-0.023 [0.062] N=1,261	0.011 [0.069] N=1,113	-0.045 [0.076] N=920	0.042 [0.107] N=709
Snack foods consumed at least weekly	-0.026 [0.056] N=1,391	-0.019 [0.061] N=1,261	0.013 [0.071] N=1,113	0.003 [0.083] N=920	0.014 [0.103] N=709
Packaged soft drinks consumed at least weekly	-0.020 [0.056] N=1,391	-0.018 [0.061] N=1,261	-0.021 [0.066] N=1,113	-0.025 [0.076] N=920	-0.071 [0.098] N=709
Grains consumed at least weekly	0.061 [0.051] N=1,391	**0.093* [0.056] N=1,261**	0.080 [0.063] N=1,113	0.113 [0.077] N=920	0.100 [0.114] N=709
Child was ever breastfed	0.005 [0.023] N=1,615	0.014 [0.027] N=1,470	0.016 [0.031] N=1,294	0.020 [0.037] N=1,058	0.035 [0.046] N=818

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,527	-0.013 [0.023] N=1,639	0.001 [0.023] N=1,509	-0.002 [0.024] N=1,361	-0.008 [0.027] N=1,168	-0.021 [0.032] N=957
Vegetables consumed at least weekly	0.895 P=1,466	0.010 [0.028] N=1,639	0.010 [0.028] N=1,509	-0.007 [0.029] N=1,361	-0.011 [0.030] N=1,168	-0.043 [0.035] N=957
Meat fish or poultry consumed at least weekly	0.770 P=1,272	0.069 [0.050] N=1,639	0.087 [0.053] N=1,509	0.078 [0.057] N=1,361	0.051 [0.065] N=1,168	0.043 [0.076] N=957
Legumes consumed at least weekly	0.770 P=1,269	-0.027 [0.042] N=1,639	-0.040 [0.044] N=1,509	-0.032 [0.045] N=1,361	-0.043 [0.049] N=1,168	-0.079 [0.057] N=957
Animal products consumed at least weekly	0.908 P=1,492	0.001 [0.030] N=1,639	0.001 [0.031] N=1,509	-0.014 [0.033] N=1,361	-0.012 [0.036] N=1,168	-0.018 [0.041] N=957
Sweets consumed at least weekly	0.608 P=1,001	-0.032 [0.040] N=1,639	-0.023 [0.042] N=1,509	-0.004 [0.044] N=1,361	-0.023 [0.047] N=1,168	0.007 [0.056] N=957
Snack foods consumed at least weekly	0.494 P=810	-0.038 [0.041] N=1,639	-0.033 [0.043] N=1,509	-0.017 [0.047] N=1,361	-0.022 [0.049] N=1,168	-0.022 [0.053] N=957
Packaged soft drinks consumed at least weekly	0.291 P=473	-0.012 [0.049] N=1,639	-0.008 [0.052] N=1,509	-0.010 [0.056] N=1,361	-0.007 [0.061] N=1,168	-0.010 [0.072] N=957
Grains consumed at least weekly	0.880 P=1,448	0.023 [0.032] N=1,639	0.036 [0.033] N=1,509	0.023 [0.035] N=1,361	0.030 [0.038] N=1,168	0.010 [0.044] N=957
Child was ever breastfed	0.967 P=1,834	-0.004 [0.014] N=1,901	0.000 [0.015] N=1,756	0.000 [0.015] N=1,580	0.004 [0.017] N=1,344	0.001 [0.016] N=1,104

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.037] N=1,391	0.020 [0.037] N=1,261	0.016 [0.041] N=1,113	0.008 [0.050] N=920	-0.016 [0.068] N=709
Vegetables consumed at least weekly	0.032 [0.045] N=1,391	0.038 [0.047] N=1,261	0.014 [0.051] N=1,113	0.011 [0.059] N=920	-0.064 [0.077] N=709
Meat fish or poultry consumed at least weekly	0.063 [0.057] N=1,391	0.091 [0.059] N=1,261	0.078 [0.067] N=1,113	-0.005 [0.072] N=920	-0.056 [0.085] N=709
Legumes consumed at least weekly	0.034 [0.063] N=1,391	0.021 [0.068] N=1,261	0.055 [0.070] N=1,113	0.055 [0.084] N=920	0.001 [0.120] N=709
Animal products consumed at least weekly	0.003 [0.041] N=1,391	0.004 [0.044] N=1,261	-0.027 [0.050] N=1,113	-0.024 [0.059] N=920	-0.046 [0.073] N=709
Sweets consumed at least weekly	-0.037 [0.057] N=1,391	-0.023 [0.062] N=1,261	0.011 [0.069] N=1,113	-0.045 [0.076] N=920	0.042 [0.107] N=709
Snack foods consumed at least weekly	-0.025 [0.056] N=1,391	-0.018 [0.061] N=1,261	0.017 [0.071] N=1,113	0.004 [0.083] N=920	0.017 [0.103] N=709
Packaged soft drinks consumed at least weekly	-0.020 [0.056] N=1,391	-0.018 [0.061] N=1,261	-0.022 [0.065] N=1,113	-0.025 [0.076] N=920	-0.072 [0.098] N=709
Grains consumed at least weekly	0.061 [0.051] N=1,391	**0.094* [0.056] N=1,261**	0.081 [0.062] N=1,113	0.113 [0.077] N=920	0.100 [0.114] N=709
Child was ever breastfed	0.005 [0.023] N=1,615	0.014 [0.027] N=1,470	0.016 [0.031] N=1,294	0.019 [0.037] N=1,058	0.034 [0.046] N=818

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,527	-0.022 [0.023] N=1,639	-0.010 [0.023] N=1,509	-0.012 [0.025] N=1,361	-0.015 [0.028] N=1,168	-0.030 [0.033] N=957
Vegetables consumed at least weekly	0.895 P=1,466	0.003 [0.028] N=1,639	0.001 [0.028] N=1,509	-0.015 [0.028] N=1,361	-0.017 [0.030] N=1,168	-0.052 [0.035] N=957
Meat fish or poultry consumed at least weekly	0.770 P=1,272	0.045 [0.047] N=1,639	0.059 [0.050] N=1,509	0.055 [0.055] N=1,361	0.034 [0.062] N=1,168	0.021 [0.072] N=957
Legumes consumed at least weekly	0.770 P=1,269	-0.033 [0.042] N=1,639	-0.048 [0.044] N=1,509	-0.038 [0.045] N=1,361	-0.046 [0.049] N=1,168	-0.082 [0.056] N=957
Animal products consumed at least weekly	0.908 P=1,492	-0.004 [0.030] N=1,639	-0.005 [0.032] N=1,509	-0.019 [0.034] N=1,361	-0.016 [0.037] N=1,168	-0.024 [0.042] N=957
Sweets consumed at least weekly	0.608 P=1,001	-0.035 [0.040] N=1,639	-0.027 [0.042] N=1,509	-0.009 [0.045] N=1,361	-0.025 [0.048] N=1,168	0.003 [0.055] N=957
Snack foods consumed at least weekly	0.494 P=810	-0.034 [0.040] N=1,639	-0.030 [0.042] N=1,509	-0.016 [0.046] N=1,361	-0.018 [0.048] N=1,168	-0.018 [0.053] N=957
Packaged soft drinks consumed at least weekly	0.291 P=473	0.002 [0.048] N=1,639	0.009 [0.051] N=1,509	0.005 [0.054] N=1,361	0.006 [0.059] N=1,168	0.005 [0.068] N=957
Grains consumed at least weekly	0.880 P=1,448	0.018 [0.032] N=1,639	0.032 [0.033] N=1,509	0.021 [0.035] N=1,361	0.028 [0.037] N=1,168	0.009 [0.044] N=957
Child was ever breastfed	0.967 P=1,834	-0.007 [0.014] N=1,901	-0.004 [0.015] N=1,756	-0.004 [0.016] N=1,580	0.001 [0.017] N=1,344	-0.002 [0.017] N=1,104

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.016 [0.036] N=1,391	0.010 [0.035] N=1,261	0.010 [0.041] N=1,113	0.017 [0.049] N=920	-0.003 [0.068] N=709
Vegetables consumed at least weekly	0.028 [0.045] N=1,391	0.030 [0.046] N=1,261	0.010 [0.050] N=1,113	0.019 [0.058] N=920	-0.055 [0.076] N=709
Meat fish or poultry consumed at least weekly	0.043 [0.054] N=1,391	0.063 [0.056] N=1,261	0.059 [0.066] N=1,113	-0.008 [0.070] N=920	-0.059 [0.086] N=709
Legumes consumed at least weekly	0.028 [0.063] N=1,391	0.011 [0.068] N=1,261	0.051 [0.070] N=1,113	0.061 [0.085] N=920	0.013 [0.122] N=709
Animal products consumed at least weekly	-0.002 [0.041] N=1,391	-0.001 [0.045] N=1,261	-0.030 [0.051] N=1,113	-0.017 [0.060] N=920	-0.030 [0.074] N=709
Sweets consumed at least weekly	-0.038 [0.057] N=1,391	-0.026 [0.063] N=1,261	0.010 [0.070] N=1,113	-0.039 [0.077] N=920	0.051 [0.110] N=709
Snack foods consumed at least weekly	-0.019 [0.055] N=1,391	-0.013 [0.060] N=1,261	0.022 [0.069] N=1,113	0.014 [0.080] N=920	0.026 [0.101] N=709
Packaged soft drinks consumed at least weekly	-0.006 [0.054] N=1,391	0.002 [0.059] N=1,261	-0.003 [0.064] N=1,113	-0.012 [0.075] N=920	-0.060 [0.094] N=709
Grains consumed at least weekly	0.056 [0.051] N=1,391	0.089 [0.056] N=1,261	0.078 [0.062] N=1,113	0.115 [0.076] N=920	0.103 [0.113] N=709
Child was ever breastfed	0.003 [0.023] N=1,615	0.010 [0.027] N=1,470	0.012 [0.031] N=1,294	0.022 [0.037] N=1,058	0.031 [0.047] N=818

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.932 P=1,527	-0.023 [0.024] N=1,639	-0.010 [0.023] N=1,509	-0.015 [0.025] N=1,361	-0.018 [0.028] N=1,168	-0.033 [0.032] N=957
Vegetables consumed at least weekly	0.895 P=1,466	0.003 [0.028] N=1,639	0.002 [0.028] N=1,509	-0.014 [0.029] N=1,361	-0.018 [0.030] N=1,168	-0.049 [0.035] N=957
Meat fish or poultry consumed at least weekly	0.770 P=1,272	0.042 [0.048] N=1,639	0.056 [0.051] N=1,509	0.046 [0.055] N=1,361	0.024 [0.062] N=1,168	0.007 [0.072] N=957
Legumes consumed at least weekly	0.770 P=1,269	-0.036 [0.043] N=1,639	-0.052 [0.045] N=1,509	-0.047 [0.046] N=1,361	-0.054 [0.050] N=1,168	-0.091 [0.059] N=957
Animal products consumed at least weekly	0.908 P=1,492	-0.006 [0.030] N=1,639	-0.005 [0.032] N=1,509	-0.020 [0.034] N=1,361	-0.020 [0.037] N=1,168	-0.026 [0.041] N=957
Sweets consumed at least weekly	0.608 P=1,001	-0.034 [0.040] N=1,639	-0.027 [0.042] N=1,509	-0.011 [0.045] N=1,361	-0.026 [0.047] N=1,168	0.002 [0.056] N=957
Snack foods consumed at least weekly	0.494 P=810	-0.034 [0.041] N=1,639	-0.029 [0.043] N=1,509	-0.019 [0.047] N=1,361	-0.020 [0.049] N=1,168	-0.022 [0.053] N=957
Packaged soft drinks consumed at least weekly	0.291 P=473	0.003 [0.047] N=1,639	0.005 [0.051] N=1,509	0.000 [0.054] N=1,361	0.005 [0.058] N=1,168	0.007 [0.067] N=957
Grains consumed at least weekly	0.880 P=1,448	0.021 [0.032] N=1,639	0.035 [0.034] N=1,509	0.023 [0.035] N=1,361	0.030 [0.038] N=1,168	0.013 [0.046] N=957
Child was ever breastfed	0.967 P=1,834	-0.008 [0.014] N=1,901	-0.005 [0.016] N=1,756	-0.005 [0.016] N=1,580	-0.000 [0.017] N=1,344	-0.004 [0.018] N=1,104

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.015 [0.037] N=1,391	0.012 [0.036] N=1,261	0.007 [0.041] N=1,113	0.010 [0.050] N=920	-0.017 [0.069] N=709
Vegetables consumed at least weekly	0.028 [0.044] N=1,391	0.032 [0.046] N=1,261	0.008 [0.050] N=1,113	0.013 [0.058] N=920	-0.071 [0.076] N=709
Meat fish or poultry consumed at least weekly	0.044 [0.055] N=1,391	0.067 [0.056] N=1,261	0.058 [0.065] N=1,113	-0.003 [0.070] N=920	-0.060 [0.087] N=709
Legumes consumed at least weekly	0.033 [0.063] N=1,391	0.015 [0.068] N=1,261	0.051 [0.070] N=1,113	0.073 [0.083] N=920	0.019 [0.124] N=709
Animal products consumed at least weekly	-0.003 [0.041] N=1,391	-0.001 [0.045] N=1,261	-0.033 [0.050] N=1,113	-0.029 [0.059] N=920	-0.054 [0.076] N=709
Sweets consumed at least weekly	-0.030 [0.057] N=1,391	-0.020 [0.062] N=1,261	0.012 [0.069] N=1,113	-0.029 [0.078] N=920	0.057 [0.112] N=709
Snack foods consumed at least weekly	-0.013 [0.054] N=1,391	-0.009 [0.059] N=1,261	0.023 [0.068] N=1,113	0.024 [0.078] N=920	0.039 [0.099] N=709
Packaged soft drinks consumed at least weekly	-0.000 [0.053] N=1,391	-0.001 [0.058] N=1,261	-0.007 [0.063] N=1,113	-0.009 [0.074] N=920	-0.050 [0.092] N=709
Grains consumed at least weekly	0.057 [0.051] N=1,391	0.089 [0.056] N=1,261	0.077 [0.062] N=1,113	0.111 [0.075] N=920	0.090 [0.109] N=709
Child was ever breastfed	0.003 [0.024] N=1,615	0.011 [0.027] N=1,470	0.011 [0.031] N=1,294	0.021 [0.038] N=1,058	0.034 [0.048] N=818

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,414	-0.021 [0.024] N=1,517	-0.012 [0.023] N=1,402	-0.015 [0.025] N=1,263	-0.019 [0.028] N=1,088	-0.041 [0.032] N=893
Vegetables consumed at least weekly	0.896 P=1,358	-0.005 [0.030] N=1,517	-0.008 [0.029] N=1,402	-0.028 [0.030] N=1,263	-0.026 [0.031] N=1,088	**-0.060* [0.035] N=893**
Meat fish or poultry consumed at least weekly	0.771 P=1,177	0.057 [0.047] N=1,517	0.064 [0.050] N=1,402	0.048 [0.054] N=1,263	0.017 [0.060] N=1,088	-0.007 [0.071] N=893
Legumes consumed at least weekly	0.768 P=1,171	-0.043 [0.044] N=1,517	-0.062 [0.045] N=1,402	-0.059 [0.045] N=1,263	-0.069 [0.051] N=1,088	**-0.111* [0.059] N=893**
Animal products consumed at least weekly	0.911 P=1,385	0.000 [0.030] N=1,517	-0.001 [0.032] N=1,402	-0.020 [0.034] N=1,263	-0.024 [0.037] N=1,088	-0.037 [0.042] N=893
Sweets consumed at least weekly	0.610 P=928	-0.022 [0.041] N=1,517	-0.020 [0.043] N=1,402	-0.003 [0.045] N=1,263	-0.019 [0.048] N=1,088	0.009 [0.057] N=893
Snack foods consumed at least weekly	0.496 P=751	-0.042 [0.043] N=1,517	-0.039 [0.045] N=1,402	-0.026 [0.049] N=1,263	-0.030 [0.053] N=1,088	-0.020 [0.058] N=893
Packaged soft drinks consumed at least weekly	0.296 P=447	0.015 [0.049] N=1,517	0.019 [0.052] N=1,402	0.007 [0.055] N=1,263	0.005 [0.060] N=1,088	0.011 [0.070] N=893
Grains consumed at least weekly	0.883 P=1,342	0.013 [0.033] N=1,517	0.025 [0.035] N=1,402	0.005 [0.036] N=1,263	0.015 [0.039] N=1,088	-0.008 [0.046] N=893
Child was ever breastfed	0.967 P=1,695	-0.009 [0.015] N=1,757	-0.004 [0.016] N=1,628	-0.003 [0.016] N=1,461	0.003 [0.018] N=1,250	-0.001 [0.018] N=1,029

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.011 [0.039] N=1,284	0.007 [0.040] N=1,169	0.002 [0.044] N=1,030	-0.001 [0.053] N=855	-0.048 [0.071] N=660
Vegetables consumed at least weekly	0.017 [0.047] N=1,284	0.016 [0.050] N=1,169	-0.017 [0.053] N=1,030	-0.006 [0.061] N=855	-0.090 [0.075] N=660
Meat fish or poultry consumed at least weekly	0.077 [0.057] N=1,284	0.088 [0.060] N=1,169	0.067 [0.067] N=1,030	-0.019 [0.072] N=855	-0.091 [0.087] N=660
Legumes consumed at least weekly	0.025 [0.064] N=1,284	-0.002 [0.068] N=1,169	0.023 [0.069] N=1,030	0.028 [0.084] N=855	-0.054 [0.120] N=660
Animal products consumed at least weekly	0.008 [0.043] N=1,284	0.005 [0.047] N=1,169	-0.031 [0.052] N=1,030	-0.032 [0.062] N=855	-0.070 [0.077] N=660
Sweets consumed at least weekly	-0.024 [0.059] N=1,284	-0.024 [0.065] N=1,169	0.001 [0.072] N=1,030	-0.054 [0.078] N=855	0.009 [0.109] N=660
Snack foods consumed at least weekly	-0.031 [0.056] N=1,284	-0.028 [0.062] N=1,169	-0.002 [0.071] N=1,030	-0.025 [0.082] N=855	-0.021 [0.101] N=660
Packaged soft drinks consumed at least weekly	0.011 [0.057] N=1,284	0.012 [0.060] N=1,169	-0.013 [0.064] N=1,030	-0.030 [0.076] N=855	-0.062 [0.102] N=660
Grains consumed at least weekly	0.055 [0.053] N=1,284	0.083 [0.058] N=1,169	0.053 [0.064] N=1,030	0.095 [0.079] N=855	0.069 [0.115] N=660
Child was ever breastfed	-0.003 [0.025] N=1,489	0.008 [0.028] N=1,360	0.009 [0.032] N=1,193	0.017 [0.038] N=982	0.026 [0.047] N=761

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,414	-0.022 [0.024] N=1,517	-0.013 [0.023] N=1,402	-0.015 [0.025] N=1,263	-0.019 [0.028] N=1,088	-0.040 [0.031] N=893
Vegetables consumed at least weekly	0.896 P=1,358	-0.006 [0.029] N=1,517	-0.009 [0.029] N=1,402	-0.028 [0.030] N=1,263	-0.024 [0.031] N=1,088	**-0.058* [0.035] N=893**
Meat fish or poultry consumed at least weekly	0.771 P=1,177	0.060 [0.048] N=1,517	0.068 [0.050] N=1,402	0.052 [0.054] N=1,263	0.023 [0.061] N=1,088	0.004 [0.071] N=893
Legumes consumed at least weekly	0.768 P=1,171	-0.045 [0.043] N=1,517	-0.065 [0.045] N=1,402	-0.063 [0.046] N=1,263	-0.073 [0.051] N=1,088	**-0.118** [0.059] N=893**
Animal products consumed at least weekly	0.911 P=1,385	-0.001 [0.030] N=1,517	-0.002 [0.032] N=1,402	-0.020 [0.034] N=1,263	-0.022 [0.037] N=1,088	-0.034 [0.041] N=893
Sweets consumed at least weekly	0.610 P=928	-0.021 [0.040] N=1,517	-0.019 [0.043] N=1,402	-0.000 [0.045] N=1,263	-0.016 [0.047] N=1,088	0.013 [0.056] N=893
Snack foods consumed at least weekly	0.496 P=751	-0.042 [0.043] N=1,517	-0.040 [0.045] N=1,402	-0.026 [0.049] N=1,263	-0.030 [0.052] N=1,088	-0.024 [0.057] N=893
Packaged soft drinks consumed at least weekly	0.296 P=447	0.012 [0.049] N=1,517	0.016 [0.052] N=1,402	0.006 [0.056] N=1,263	0.002 [0.060] N=1,088	0.005 [0.071] N=893
Grains consumed at least weekly	0.883 P=1,342	0.013 [0.033] N=1,517	0.024 [0.034] N=1,402	0.004 [0.036] N=1,263	0.015 [0.039] N=1,088	-0.010 [0.045] N=893
Child was ever breastfed	0.967 P=1,695	-0.008 [0.015] N=1,757	-0.003 [0.016] N=1,628	-0.003 [0.016] N=1,461	0.004 [0.018] N=1,250	0.000 [0.017] N=1,029

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.013 [0.038] N=1,284	0.006 [0.038] N=1,169	0.003 [0.044] N=1,030	-0.000 [0.052] N=855	-0.045 [0.068] N=660
Vegetables consumed at least weekly	0.014 [0.047] N=1,284	0.014 [0.049] N=1,169	-0.018 [0.052] N=1,030	-0.008 [0.061] N=855	-0.087 [0.073] N=660
Meat fish or poultry consumed at least weekly	0.076 [0.057] N=1,284	0.088 [0.059] N=1,169	0.068 [0.066] N=1,030	-0.016 [0.072] N=855	-0.084 [0.086] N=660
Legumes consumed at least weekly	0.024 [0.064] N=1,284	-0.001 [0.069] N=1,169	0.026 [0.070] N=1,030	0.030 [0.085] N=855	-0.052 [0.120] N=660
Animal products consumed at least weekly	0.005 [0.043] N=1,284	0.003 [0.046] N=1,169	-0.030 [0.051] N=1,030	-0.034 [0.061] N=855	-0.068 [0.075] N=660
Sweets consumed at least weekly	-0.031 [0.058] N=1,284	-0.034 [0.064] N=1,169	-0.005 [0.072] N=1,030	-0.055 [0.078] N=855	0.015 [0.109] N=660
Snack foods consumed at least weekly	-0.038 [0.057] N=1,284	-0.037 [0.062] N=1,169	-0.007 [0.072] N=1,030	-0.027 [0.083] N=855	-0.019 [0.102] N=660
Packaged soft drinks consumed at least weekly	0.003 [0.057] N=1,284	0.003 [0.061] N=1,169	-0.016 [0.066] N=1,030	-0.030 [0.077] N=855	-0.062 [0.102] N=660
Grains consumed at least weekly	0.053 [0.053] N=1,284	0.083 [0.058] N=1,169	0.056 [0.065] N=1,030	0.095 [0.080] N=855	0.071 [0.118] N=660
Child was ever breastfed	0.000 [0.025] N=1,489	0.010 [0.028] N=1,360	0.011 [0.033] N=1,193	0.022 [0.039] N=982	0.031 [0.050] N=761

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,363	0.032 [0.029] N=1,461	0.034 [0.030] N=1,329	0.028 [0.034] N=1,160	0.020 [0.034] N=976	0.029 [0.043] N=787
Vegetables consumed at least weekly	0.897 P=1,311	0.052 [0.032] N=1,461	**0.062* [0.033] N=1,329**	**0.064* [0.037] N=1,160**	0.054 [0.040] N=976	0.052 [0.046] N=787
Meat fish or poultry consumed at least weekly	0.762 P=1,128	**0.094** [0.041] N=1,461**	**0.100** [0.042] N=1,329**	**0.088* [0.046] N=1,160**	**0.124** [0.049] N=976**	**0.117** [0.056] N=787**
Legumes consumed at least weekly	0.769 P=1,137	-0.000 [0.040] N=1,461	0.021 [0.042] N=1,329	0.011 [0.046] N=1,160	-0.020 [0.048] N=976	-0.002 [0.058] N=787
Animal products consumed at least weekly	0.907 P=1,330	0.045 [0.034] N=1,461	0.053 [0.036] N=1,329	0.052 [0.039] N=1,160	0.037 [0.040] N=976	0.058 [0.047] N=787
Sweets consumed at least weekly	0.615 P=910	-0.019 [0.050] N=1,461	-0.022 [0.053] N=1,329	-0.026 [0.057] N=1,160	-0.036 [0.060] N=976	-0.087 [0.069] N=787
Snack foods consumed at least weekly	0.497 P=729	-0.040 [0.046] N=1,461	-0.036 [0.050] N=1,329	-0.029 [0.053] N=1,160	-0.029 [0.057] N=976	-0.075 [0.066] N=787
Packaged soft drinks consumed at least weekly	0.283 P=412	-0.021 [0.044] N=1,461	-0.023 [0.046] N=1,329	0.011 [0.049] N=1,160	0.003 [0.054] N=976	-0.014 [0.061] N=787
Grains consumed at least weekly	0.878 P=1,291	0.058 [0.039] N=1,461	**0.070* [0.041] N=1,329**	**0.073* [0.044] N=1,160**	0.077 [0.049] N=976	0.070 [0.058] N=787
Child was ever breastfed	0.965 P=1,618	-0.020 [0.018] N=1,678	-0.019 [0.019] N=1,528	-0.010 [0.018] N=1,332	-0.014 [0.019] N=1,125	-0.014 [0.021] N=904

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.012 [0.040] N=1,264	-0.014 [0.042] N=1,132	-0.029 [0.050] N=963	-0.032 [0.054] N=779	-0.036 [0.083] N=590
Vegetables consumed at least weekly	0.003 [0.045] N=1,264	0.011 [0.049] N=1,132	0.001 [0.058] N=963	-0.013 [0.065] N=779	-0.037 [0.086] N=590
Meat fish or poultry consumed at least weekly	0.026 [0.057] N=1,264	0.031 [0.058] N=1,132	-0.017 [0.065] N=963	0.025 [0.067] N=779	-0.038 [0.087] N=590
Legumes consumed at least weekly	-0.005 [0.052] N=1,264	0.027 [0.057] N=1,132	0.015 [0.065] N=963	-0.024 [0.067] N=779	0.009 [0.084] N=590
Animal products consumed at least weekly	-0.016 [0.048] N=1,264	-0.003 [0.052] N=1,132	-0.011 [0.058] N=963	-0.046 [0.060] N=779	-0.041 [0.076] N=590
Sweets consumed at least weekly	-0.021 [0.064] N=1,264	-0.025 [0.068] N=1,132	-0.031 [0.077] N=963	-0.059 [0.091] N=779	**-0.215* [0.112] N=590**
Snack foods consumed at least weekly	-0.045 [0.061] N=1,264	-0.039 [0.067] N=1,132	-0.023 [0.075] N=963	-0.015 [0.088] N=779	-0.149 [0.108] N=590
Packaged soft drinks consumed at least weekly	-0.054 [0.068] N=1,264	-0.057 [0.073] N=1,132	-0.001 [0.084] N=963	-0.022 [0.102] N=779	-0.096 [0.130] N=590
Grains consumed at least weekly	-0.020 [0.040] N=1,264	-0.014 [0.044] N=1,132	-0.019 [0.048] N=963	-0.054 [0.053] N=779	**-0.168** [0.071] N=590**
Child was ever breastfed	-0.009 [0.033] N=1,452	-0.005 [0.034] N=1,302	0.015 [0.035] N=1,106	0.010 [0.040] N=899	0.022 [0.053] N=678

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,363	0.032 [0.029] N=1,461	0.034 [0.030] N=1,329	0.028 [0.034] N=1,160	0.020 [0.034] N=976	0.029 [0.043] N=787
Vegetables consumed at least weekly	0.897 P=1,311	0.052 [0.032] N=1,461	**0.062* [0.033] N=1,329**	**0.064* [0.037] N=1,160**	0.054 [0.040] N=976	0.052 [0.045] N=787
Meat fish or poultry consumed at least weekly	0.762 P=1,128	**0.095** [0.041] N=1,461**	**0.101** [0.042] N=1,329**	**0.089* [0.046] N=1,160**	**0.123** [0.048] N=976**	**0.116** [0.056] N=787**
Legumes consumed at least weekly	0.769 P=1,137	-0.000 [0.040] N=1,461	0.021 [0.042] N=1,329	0.011 [0.046] N=1,160	-0.020 [0.048] N=976	-0.001 [0.058] N=787
Animal products consumed at least weekly	0.907 P=1,330	0.046 [0.034] N=1,461	0.054 [0.035] N=1,329	0.053 [0.038] N=1,160	0.036 [0.040] N=976	0.056 [0.046] N=787
Sweets consumed at least weekly	0.615 P=910	-0.018 [0.050] N=1,461	-0.021 [0.053] N=1,329	-0.025 [0.057] N=1,160	-0.036 [0.060] N=976	-0.088 [0.069] N=787
Snack foods consumed at least weekly	0.497 P=729	-0.039 [0.046] N=1,461	-0.035 [0.050] N=1,329	-0.028 [0.053] N=1,160	-0.029 [0.057] N=976	-0.076 [0.066] N=787
Packaged soft drinks consumed at least weekly	0.283 P=412	-0.020 [0.044] N=1,461	-0.023 [0.046] N=1,329	0.012 [0.048] N=1,160	0.003 [0.054] N=976	-0.016 [0.060] N=787
Grains consumed at least weekly	0.878 P=1,291	0.058 [0.039] N=1,461	**0.070* [0.041] N=1,329**	**0.073* [0.044] N=1,160**	0.077 [0.049] N=976	0.071 [0.058] N=787
Child was ever breastfed	0.965 P=1,618	-0.019 [0.018] N=1,678	-0.018 [0.019] N=1,528	-0.010 [0.018] N=1,332	-0.013 [0.019] N=1,125	-0.013 [0.021] N=904

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.011 [0.040] N=1,264	-0.012 [0.042] N=1,132	-0.028 [0.051] N=963	-0.032 [0.055] N=779	-0.036 [0.084] N=590
Vegetables consumed at least weekly	0.004 [0.045] N=1,264	0.012 [0.049] N=1,132	0.002 [0.058] N=963	-0.010 [0.064] N=779	-0.035 [0.086] N=590
Meat fish or poultry consumed at least weekly	0.029 [0.057] N=1,264	0.037 [0.058] N=1,132	-0.010 [0.064] N=963	0.032 [0.067] N=779	-0.026 [0.086] N=590
Legumes consumed at least weekly	-0.003 [0.051] N=1,264	0.029 [0.056] N=1,132	0.016 [0.065] N=963	-0.021 [0.067] N=779	0.004 [0.084] N=590
Animal products consumed at least weekly	-0.011 [0.046] N=1,264	0.004 [0.050] N=1,132	-0.005 [0.057] N=963	-0.039 [0.059] N=779	-0.030 [0.074] N=590
Sweets consumed at least weekly	-0.017 [0.064] N=1,264	-0.019 [0.068] N=1,132	-0.026 [0.077] N=963	-0.057 [0.091] N=779	**-0.212* [0.112] N=590**
Snack foods consumed at least weekly	-0.039 [0.061] N=1,264	-0.033 [0.067] N=1,132	-0.017 [0.075] N=963	-0.012 [0.088] N=779	-0.141 [0.109] N=590
Packaged soft drinks consumed at least weekly	-0.052 [0.067] N=1,264	-0.056 [0.072] N=1,132	0.003 [0.083] N=963	-0.020 [0.102] N=779	-0.085 [0.127] N=590
Grains consumed at least weekly	-0.020 [0.040] N=1,264	-0.014 [0.045] N=1,132	-0.020 [0.049] N=963	-0.057 [0.054] N=779	**-0.174** [0.074] N=590**
Child was ever breastfed	-0.008 [0.033] N=1,452	-0.003 [0.034] N=1,302	0.017 [0.035] N=1,106	0.013 [0.041] N=899	0.025 [0.054] N=678

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,363	0.031 [0.029] N=1,461	0.034 [0.030] N=1,329	0.031 [0.034] N=1,160	0.024 [0.035] N=976	0.031 [0.043] N=787
Vegetables consumed at least weekly	0.897 P=1,311	**0.052* [0.031] N=1,461**	**0.062* [0.033] N=1,329**	**0.067* [0.036] N=1,160**	0.057 [0.040] N=976	0.053 [0.046] N=787
Meat fish or poultry consumed at least weekly	0.762 P=1,128	**0.092** [0.040] N=1,461**	**0.099** [0.042] N=1,329**	**0.097** [0.045] N=1,160**	**0.133*** [0.048] N=976**	**0.123** [0.054] N=787**
Legumes consumed at least weekly	0.769 P=1,137	-0.001 [0.040] N=1,461	0.020 [0.043] N=1,329	0.012 [0.047] N=1,160	-0.017 [0.049] N=976	0.002 [0.059] N=787
Animal products consumed at least weekly	0.907 P=1,330	0.045 [0.034] N=1,461	0.053 [0.035] N=1,329	0.054 [0.038] N=1,160	0.039 [0.040] N=976	0.058 [0.046] N=787
Sweets consumed at least weekly	0.615 P=910	-0.017 [0.050] N=1,461	-0.021 [0.053] N=1,329	-0.023 [0.057] N=1,160	-0.036 [0.059] N=976	-0.088 [0.068] N=787
Snack foods consumed at least weekly	0.497 P=729	-0.036 [0.046] N=1,461	-0.033 [0.050] N=1,329	-0.029 [0.053] N=1,160	-0.031 [0.057] N=976	-0.078 [0.066] N=787
Packaged soft drinks consumed at least weekly	0.283 P=412	-0.019 [0.045] N=1,461	-0.023 [0.047] N=1,329	0.008 [0.050] N=1,160	-0.003 [0.056] N=976	-0.019 [0.063] N=787
Grains consumed at least weekly	0.878 P=1,291	0.057 [0.039] N=1,461	**0.069* [0.041] N=1,329**	**0.075* [0.045] N=1,160**	0.079 [0.050] N=976	0.073 [0.059] N=787
Child was ever breastfed	0.965 P=1,618	-0.019 [0.019] N=1,678	-0.018 [0.019] N=1,528	-0.008 [0.019] N=1,332	-0.011 [0.019] N=1,125	-0.013 [0.021] N=904

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.012 [0.040] N=1,264	-0.013 [0.041] N=1,132	-0.024 [0.049] N=963	-0.029 [0.055] N=779	-0.037 [0.084] N=590
Vegetables consumed at least weekly	0.004 [0.045] N=1,264	0.012 [0.048] N=1,132	0.008 [0.056] N=963	-0.008 [0.064] N=779	-0.034 [0.087] N=590
Meat fish or poultry consumed at least weekly	0.027 [0.053] N=1,264	0.034 [0.055] N=1,132	0.005 [0.060] N=963	0.045 [0.067] N=779	-0.026 [0.081] N=590
Legumes consumed at least weekly	-0.002 [0.051] N=1,264	0.029 [0.057] N=1,132	0.016 [0.066] N=963	-0.017 [0.068] N=779	0.001 [0.087] N=590
Animal products consumed at least weekly	-0.014 [0.046] N=1,264	-0.001 [0.049] N=1,132	-0.004 [0.056] N=963	-0.041 [0.059] N=779	-0.032 [0.073] N=590
Sweets consumed at least weekly	-0.015 [0.063] N=1,264	-0.020 [0.068] N=1,132	-0.026 [0.076] N=963	-0.061 [0.091] N=779	**-0.212* [0.110] N=590**
Snack foods consumed at least weekly	-0.031 [0.060] N=1,264	-0.029 [0.067] N=1,132	-0.020 [0.074] N=963	-0.017 [0.085] N=779	-0.138 [0.107] N=590
Packaged soft drinks consumed at least weekly	-0.051 [0.068] N=1,264	-0.059 [0.073] N=1,132	-0.012 [0.085] N=963	-0.039 [0.104] N=779	-0.088 [0.129] N=590
Grains consumed at least weekly	-0.022 [0.041] N=1,264	-0.017 [0.045] N=1,132	-0.020 [0.049] N=963	-0.060 [0.054] N=779	**-0.178** [0.073] N=590**
Child was ever breastfed	-0.010 [0.034] N=1,452	-0.006 [0.035] N=1,302	0.018 [0.036] N=1,106	0.014 [0.041] N=899	0.023 [0.053] N=678

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.933 P=1,363	0.030 [0.029] N=1,461	0.033 [0.030] N=1,329	0.031 [0.034] N=1,160	0.016 [0.035] N=976	0.019 [0.042] N=787
Vegetables consumed at least weekly	0.897 P=1,311	0.051 [0.032] N=1,461	**0.062* [0.034] N=1,329**	**0.067* [0.038] N=1,160**	0.052 [0.041] N=976	0.047 [0.046] N=787
Meat fish or poultry consumed at least weekly	0.762 P=1,128	**0.094** [0.041] N=1,461**	**0.101** [0.042] N=1,329**	**0.094** [0.046] N=1,160**	**0.120** [0.050] N=976**	**0.098* [0.055] N=787**
Legumes consumed at least weekly	0.769 P=1,137	0.005 [0.040] N=1,461	0.027 [0.042] N=1,329	0.019 [0.046] N=1,160	-0.016 [0.048] N=976	0.001 [0.058] N=787
Animal products consumed at least weekly	0.907 P=1,330	0.050 [0.033] N=1,461	**0.060* [0.034] N=1,329**	**0.064* [0.037] N=1,160**	0.043 [0.040] N=976	0.059 [0.046] N=787
Sweets consumed at least weekly	0.615 P=910	-0.009 [0.051] N=1,461	-0.012 [0.053] N=1,329	-0.017 [0.058] N=1,160	-0.032 [0.061] N=976	-0.082 [0.070] N=787
Snack foods consumed at least weekly	0.497 P=729	-0.026 [0.046] N=1,461	-0.023 [0.050] N=1,329	-0.019 [0.053] N=1,160	-0.023 [0.057] N=976	-0.066 [0.066] N=787
Packaged soft drinks consumed at least weekly	0.283 P=412	-0.007 [0.044] N=1,461	-0.012 [0.046] N=1,329	0.017 [0.049] N=1,160	0.008 [0.055] N=976	-0.003 [0.062] N=787
Grains consumed at least weekly	0.878 P=1,291	0.055 [0.039] N=1,461	0.068 [0.041] N=1,329	0.072 [0.044] N=1,160	0.074 [0.049] N=976	0.066 [0.058] N=787
Child was ever breastfed	0.965 P=1,618	-0.019 [0.018] N=1,678	-0.018 [0.018] N=1,528	-0.009 [0.018] N=1,332	-0.013 [0.018] N=1,125	-0.016 [0.021] N=904

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.014 [0.039] N=1,264	-0.014 [0.041] N=1,132	-0.023 [0.050] N=963	-0.042 [0.056] N=779	-0.066 [0.083] N=590
Vegetables consumed at least weekly	0.003 [0.045] N=1,264	0.012 [0.048] N=1,132	0.007 [0.057] N=963	-0.015 [0.064] N=779	-0.051 [0.084] N=590
Meat fish or poultry consumed at least weekly	0.030 [0.054] N=1,264	0.037 [0.057] N=1,132	0.003 [0.062] N=963	0.023 [0.072] N=779	-0.079 [0.082] N=590
Legumes consumed at least weekly	0.005 [0.052] N=1,264	0.035 [0.057] N=1,132	0.030 [0.066] N=963	-0.015 [0.068] N=779	0.006 [0.085] N=590
Animal products consumed at least weekly	-0.006 [0.043] N=1,264	0.011 [0.047] N=1,132	0.011 [0.053] N=963	-0.035 [0.056] N=779	-0.038 [0.069] N=590
Sweets consumed at least weekly	-0.007 [0.064] N=1,264	-0.009 [0.069] N=1,132	-0.018 [0.077] N=963	-0.057 [0.091] N=779	**-0.207* [0.110] N=590**
Snack foods consumed at least weekly	-0.026 [0.060] N=1,264	-0.022 [0.067] N=1,132	-0.011 [0.074] N=963	-0.011 [0.085] N=779	-0.130 [0.105] N=590
Packaged soft drinks consumed at least weekly	-0.037 [0.067] N=1,264	-0.044 [0.072] N=1,132	0.003 [0.084] N=963	-0.014 [0.102] N=779	-0.046 [0.129] N=590
Grains consumed at least weekly	-0.023 [0.040] N=1,264	-0.019 [0.044] N=1,132	-0.023 [0.049] N=963	-0.067 [0.055] N=779	**-0.195** [0.075] N=590**
Child was ever breastfed	-0.008 [0.033] N=1,452	-0.003 [0.033] N=1,302	0.020 [0.034] N=1,106	0.014 [0.039] N=899	0.019 [0.052] N=678

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,262	0.027 [0.031] N=1,351	0.029 [0.032] N=1,227	0.032 [0.036] N=1,076	0.012 [0.037] N=905	0.018 [0.046] N=733
Vegetables consumed at least weekly	0.900 P=1,216	0.045 [0.032] N=1,351	0.054 [0.034] N=1,227	**0.070* [0.037] N=1,076**	0.048 [0.040] N=905	0.028 [0.046] N=733
Meat fish or poultry consumed at least weekly	0.763 P=1,043	**0.088** [0.043] N=1,351**	**0.090** [0.045] N=1,227**	**0.087* [0.048] N=1,076**	**0.095* [0.051] N=905**	0.084 [0.058] N=733
Legumes consumed at least weekly	0.769 P=1,052	0.001 [0.043] N=1,351	0.024 [0.045] N=1,227	0.017 [0.049] N=1,076	-0.030 [0.053] N=905	-0.014 [0.064] N=733
Animal products consumed at least weekly	0.910 P=1,235	0.041 [0.035] N=1,351	0.052 [0.037] N=1,227	0.060 [0.040] N=1,076	0.035 [0.041] N=905	0.052 [0.048] N=733
Sweets consumed at least weekly	0.625 P=853	-0.019 [0.051] N=1,351	-0.021 [0.053] N=1,227	-0.031 [0.058] N=1,076	-0.050 [0.063] N=905	-0.097 [0.075] N=733
Snack foods consumed at least weekly	0.504 P=683	-0.033 [0.046] N=1,351	-0.032 [0.051] N=1,227	-0.035 [0.054] N=1,076	-0.035 [0.058] N=905	-0.064 [0.068] N=733
Packaged soft drinks consumed at least weekly	0.292 P=395	-0.029 [0.045] N=1,351	-0.035 [0.047] N=1,227	-0.002 [0.049] N=1,076	-0.009 [0.056] N=905	-0.029 [0.064] N=733
Grains consumed at least weekly	0.880 P=1,197	**0.068* [0.040] N=1,351**	**0.080* [0.043] N=1,227**	**0.091** [0.046] N=1,076**	**0.089* [0.051] N=905**	0.078 [0.060] N=733
Child was ever breastfed	0.967 P=1,498	-0.024 [0.018] N=1,550	-0.026 [0.018] N=1,411	-0.017 [0.017] N=1,235	-0.025 [0.018] N=1,044	-0.023 [0.020] N=842

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.019 [0.042] N=1,165	-0.018 [0.044] N=1,041	-0.016 [0.052] N=890	-0.047 [0.055] N=719	-0.070 [0.083] N=547
Vegetables consumed at least weekly	-0.005 [0.043] N=1,165	0.002 [0.047] N=1,041	0.019 [0.054] N=890	-0.018 [0.059] N=719	-0.095 [0.075] N=547
Meat fish or poultry consumed at least weekly	0.040 [0.064] N=1,165	0.045 [0.067] N=1,041	0.017 [0.074] N=890	-0.001 [0.080] N=719	-0.089 [0.104] N=547
Legumes consumed at least weekly	0.012 [0.054] N=1,165	0.048 [0.058] N=1,041	0.050 [0.066] N=890	-0.007 [0.070] N=719	0.008 [0.087] N=547
Animal products consumed at least weekly	-0.022 [0.048] N=1,165	-0.001 [0.053] N=1,041	0.006 [0.056] N=890	-0.045 [0.058] N=719	-0.051 [0.071] N=547
Sweets consumed at least weekly	-0.020 [0.065] N=1,165	-0.022 [0.071] N=1,041	-0.049 [0.078] N=890	-0.105 [0.089] N=719	**-0.282** [0.116] N=547**
Snack foods consumed at least weekly	-0.028 [0.063] N=1,165	-0.027 [0.070] N=1,041	-0.031 [0.077] N=890	-0.033 [0.086] N=719	-0.121 [0.112] N=547
Packaged soft drinks consumed at least weekly	-0.068 [0.068] N=1,165	-0.080 [0.073] N=1,041	-0.030 [0.084] N=890	-0.057 [0.104] N=719	-0.137 [0.136] N=547
Grains consumed at least weekly	-0.007 [0.043] N=1,165	0.002 [0.049] N=1,041	0.010 [0.053] N=890	-0.042 [0.059] N=719	**-0.189** [0.078] N=547**
Child was ever breastfed	-0.013 [0.032] N=1,337	-0.013 [0.034] N=1,198	0.008 [0.033] N=1,022	-0.004 [0.037] N=831	0.010 [0.049] N=629

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.934 P=1,262	0.028 [0.031] N=1,351	0.030 [0.032] N=1,227	0.033 [0.036] N=1,076	0.012 [0.036] N=905	0.017 [0.045] N=733
Vegetables consumed at least weekly	0.900 P=1,216	0.046 [0.032] N=1,351	0.054 [0.034] N=1,227	**0.070* [0.036] N=1,076**	0.045 [0.040] N=905	0.025 [0.046] N=733
Meat fish or poultry consumed at least weekly	0.763 P=1,043	**0.090** [0.043] N=1,351**	**0.092** [0.044] N=1,227**	**0.089* [0.047] N=1,076**	**0.096* [0.050] N=905**	0.085 [0.057] N=733
Legumes consumed at least weekly	0.769 P=1,052	0.000 [0.042] N=1,351	0.023 [0.045] N=1,227	0.016 [0.049] N=1,076	-0.033 [0.052] N=905	-0.019 [0.064] N=733
Animal products consumed at least weekly	0.910 P=1,235	0.041 [0.035] N=1,351	0.052 [0.037] N=1,227	0.059 [0.039] N=1,076	0.034 [0.041] N=905	0.051 [0.047] N=733
Sweets consumed at least weekly	0.625 P=853	-0.018 [0.051] N=1,351	-0.020 [0.053] N=1,227	-0.030 [0.058] N=1,076	-0.050 [0.063] N=905	-0.097 [0.075] N=733
Snack foods consumed at least weekly	0.504 P=683	-0.030 [0.046] N=1,351	-0.029 [0.050] N=1,227	-0.030 [0.053] N=1,076	-0.032 [0.057] N=905	-0.062 [0.068] N=733
Packaged soft drinks consumed at least weekly	0.292 P=395	-0.025 [0.045] N=1,351	-0.032 [0.047] N=1,227	0.001 [0.049] N=1,076	-0.008 [0.056] N=905	-0.029 [0.064] N=733
Grains consumed at least weekly	0.880 P=1,197	**0.067* [0.040] N=1,351**	**0.079* [0.043] N=1,227**	**0.088* [0.046] N=1,076**	**0.087* [0.051] N=905**	0.076 [0.060] N=733
Child was ever breastfed	0.967 P=1,498	-0.023 [0.018] N=1,550	-0.025 [0.019] N=1,411	-0.014 [0.018] N=1,235	-0.020 [0.018] N=1,044	-0.018 [0.021] N=842

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.017 [0.042] N=1,165	-0.016 [0.044] N=1,041	-0.014 [0.053] N=890	-0.046 [0.057] N=719	-0.067 [0.087] N=547
Vegetables consumed at least weekly	-0.004 [0.043] N=1,165	0.002 [0.047] N=1,041	0.020 [0.054] N=890	-0.021 [0.060] N=719	-0.096 [0.078] N=547
Meat fish or poultry consumed at least weekly	0.040 [0.064] N=1,165	0.045 [0.067] N=1,041	0.016 [0.074] N=890	0.002 [0.080] N=719	-0.082 [0.105] N=547
Legumes consumed at least weekly	0.013 [0.054] N=1,165	0.050 [0.059] N=1,041	0.051 [0.066] N=890	-0.014 [0.070] N=719	0.007 [0.087] N=547
Animal products consumed at least weekly	-0.021 [0.048] N=1,165	-0.002 [0.053] N=1,041	0.003 [0.057] N=890	-0.049 [0.058] N=719	-0.060 [0.071] N=547
Sweets consumed at least weekly	-0.015 [0.066] N=1,165	-0.017 [0.071] N=1,041	-0.039 [0.080] N=890	-0.090 [0.092] N=719	**-0.260** [0.119] N=547**
Snack foods consumed at least weekly	-0.025 [0.063] N=1,165	-0.025 [0.070] N=1,041	-0.025 [0.077] N=890	-0.022 [0.087] N=719	-0.118 [0.113] N=547
Packaged soft drinks consumed at least weekly	-0.063 [0.069] N=1,165	-0.075 [0.074] N=1,041	-0.020 [0.084] N=890	-0.045 [0.103] N=719	-0.127 [0.135] N=547
Grains consumed at least weekly	-0.008 [0.043] N=1,165	0.000 [0.048] N=1,041	0.007 [0.052] N=890	-0.046 [0.058] N=719	**-0.193** [0.077] N=547**
Child was ever breastfed	-0.013 [0.032] N=1,337	-0.015 [0.034] N=1,198	0.009 [0.034] N=1,022	0.001 [0.039] N=831	0.013 [0.052] N=629

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.