Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-3: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.689	0.102 [0.214] N=1,642	0.053 [0.230] N=1,519	0.054 [0.253] N=1,367	0.181 [0.263] N=1,168	0.014 [0.296] N=964
WHO weight-for-age z score	-0.218	-0.117 [0.122] N=1,883	-0.112 [0.134] N=1,740	-0.107 [0.141] N=1,566	-0.111 [0.150] N=1,334	-0.233 [0.163] N=1,096
WHO weight-for-length/height z score, 0-60 months	0.308	0.226 [0.180] N=1,619	0.214 [0.191] N=1,501	0.218 [0.202] N=1,351	0.103 [0.219] N=1,155	0.208 [0.239] N=949
WHO BMI-for-age z score	0.281	0.275 [0.208] N=1,601	0.284 [0.222] N=1,486	0.286 [0.242] N=1,336	0.115 [0.258] N=1,141	0.195 [0.282] N=940
Stunted: HAZ below -2	0.286 P=458	-0.039 [0.038] N=1,642	-0.045 [0.040] N=1,519	-0.037 [0.042] N=1,367	-0.055 [0.045] N=1,168	-0.024 [0.049] N=964
Underweight: WAZ below -2	0.105 P=191	**0.049** [0.023] N=1,883**	**0.044* [0.025] N=1,740**	0.034 [0.026] N=1,566	0.040 [0.027] N=1,334	**0.058* [0.032] N=1,096**
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=189	-0.050 [0.033] N=1,619	-0.043 [0.034] N=1,501	-0.049 [0.035] N=1,351	-0.039 [0.040] N=1,155	-0.059 [0.041] N=949
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=306	0.033 [0.042] N=1,619	0.027 [0.044] N=1,501	0.024 [0.047] N=1,351	0.006 [0.052] N=1,155	-0.000 [0.059] N=949
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=141	0.007 [0.030] N=1,619	0.010 [0.032] N=1,501	0.026 [0.034] N=1,351	0.021 [0.037] N=1,155	0.031 [0.043] N=949
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=588	-0.063 [0.041] N=1,564	-0.067 [0.043] N=1,450	-0.060 [0.045] N=1,305	-0.078 [0.049] N=1,117	-0.056 [0.054] N=920
BMI-for-age z score below -2	0.135 P=216	-0.050 [0.038] N=1,601	-0.049 [0.040] N=1,486	-0.048 [0.043] N=1,336	-0.021 [0.048] N=1,141	-0.018 [0.052] N=940

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.168 [0.306] N=1,387	0.067 [0.348] N=1,264	0.146 [0.416] N=1,112	0.487 [0.457] N=913	0.268 [0.634] N=709
WHO weight-for-age z score	-0.260 [0.173] N=1,600	-0.278 [0.200] N=1,457	-0.289 [0.221] N=1,283	-0.349 [0.267] N=1,051	**-0.766** [0.330] N=813**
WHO weight-for-length/height z score, 0-60 months	0.099 [0.252] N=1,366	0.064 [0.279] N=1,248	0.003 [0.312] N=1,098	-0.274 [0.374] N=902	-0.287 [0.451] N=696
WHO BMI-for-age z score	0.264 [0.275] N=1,351	0.269 [0.307] N=1,236	0.207 [0.358] N=1,086	-0.171 [0.407] N=891	-0.174 [0.489] N=690
Stunted: HAZ below -2	-0.075 [0.054] N=1,387	-0.090 [0.059] N=1,264	-0.095 [0.068] N=1,112	**-0.154* [0.078] N=913**	-0.146 [0.106] N=709
Underweight: WAZ below -2	0.055 [0.034] N=1,600	0.046 [0.038] N=1,457	0.027 [0.042] N=1,283	0.045 [0.050] N=1,051	0.098 [0.070] N=813
Wasted: WLZ/WHZ below -2, 0-60 months	-0.048 [0.045] N=1,366	-0.037 [0.047] N=1,248	-0.046 [0.052] N=1,098	-0.030 [0.066] N=902	-0.082 [0.080] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.048] N=1,366	-0.002 [0.054] N=1,248	-0.021 [0.056] N=1,098	-0.058 [0.071] N=902	**-0.148* [0.082] N=696**
Obese: WLZ/WHZ above +3, 0-60 months	-0.030 [0.035] N=1,366	-0.030 [0.040] N=1,248	-0.008 [0.042] N=1,098	-0.021 [0.052] N=902	-0.033 [0.069] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.080 [0.056] N=1,320	-0.091 [0.061] N=1,206	-0.094 [0.067] N=1,061	**-0.149* [0.078] N=873**	-0.154 [0.102] N=676
BMI-for-age z score below -2	**-0.082* [0.048] N=1,351**	**-0.089* [0.049] N=1,236**	-0.093 [0.056] N=1,086	-0.059 [0.068] N=891	-0.083 [0.085] N=690

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.689	0.101 [0.214] N=1,642	0.050 [0.229] N=1,519	0.053 [0.252] N=1,367	0.184 [0.263] N=1,168	0.019 [0.293] N=964
WHO weight-for-age z score	-0.218	-0.122 [0.121] N=1,883	-0.119 [0.133] N=1,740	-0.113 [0.141] N=1,566	-0.118 [0.149] N=1,334	-0.246 [0.162] N=1,096
WHO weight-for-length/height z score, 0-60 months	0.308	0.223 [0.180] N=1,619	0.211 [0.191] N=1,501	0.213 [0.203] N=1,351	0.096 [0.219] N=1,155	0.192 [0.240] N=949
WHO BMI-for-age z score	0.281	0.276 [0.208] N=1,601	0.283 [0.222] N=1,486	0.281 [0.241] N=1,336	0.111 [0.258] N=1,141	0.184 [0.282] N=940
Stunted: HAZ below -2	0.286 P=458	-0.038 [0.038] N=1,642	-0.043 [0.040] N=1,519	-0.035 [0.042] N=1,367	-0.053 [0.045] N=1,168	-0.023 [0.049] N=964
Underweight: WAZ below -2	0.105 P=191	**0.050** [0.023] N=1,883**	**0.045* [0.024] N=1,740**	0.035 [0.026] N=1,566	0.041 [0.027] N=1,334	**0.059* [0.032] N=1,096**
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=189	-0.048 [0.033] N=1,619	-0.041 [0.034] N=1,501	-0.045 [0.036] N=1,351	-0.035 [0.040] N=1,155	-0.054 [0.042] N=949
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=306	0.034 [0.042] N=1,619	0.028 [0.044] N=1,501	0.025 [0.047] N=1,351	0.008 [0.052] N=1,155	0.001 [0.059] N=949
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=141	0.008 [0.030] N=1,619	0.011 [0.032] N=1,501	0.027 [0.034] N=1,351	0.023 [0.037] N=1,155	0.033 [0.043] N=949
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=588	-0.060 [0.041] N=1,564	-0.064 [0.043] N=1,450	-0.055 [0.045] N=1,305	-0.072 [0.048] N=1,117	-0.048 [0.054] N=920
BMI-for-age z score below -2	0.135 P=216	-0.049 [0.038] N=1,601	-0.048 [0.040] N=1,486	-0.045 [0.043] N=1,336	-0.018 [0.048] N=1,141	-0.014 [0.053] N=940

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.168 [0.306] N=1,387	0.068 [0.347] N=1,264	0.143 [0.415] N=1,112	0.481 [0.457] N=913	0.253 [0.633] N=709
WHO weight-for-age z score	-0.255 [0.173] N=1,600	-0.274 [0.200] N=1,457	-0.283 [0.222] N=1,283	-0.347 [0.268] N=1,051	**-0.757** [0.329] N=813**
WHO weight-for-length/height z score, 0-60 months	0.101 [0.251] N=1,366	0.065 [0.278] N=1,248	0.004 [0.309] N=1,098	-0.275 [0.373] N=902	-0.285 [0.446] N=696
WHO BMI-for-age z score	0.265 [0.275] N=1,351	0.270 [0.306] N=1,236	0.209 [0.355] N=1,086	-0.172 [0.406] N=891	-0.175 [0.483] N=690
Stunted: HAZ below -2	-0.077 [0.054] N=1,387	-0.092 [0.059] N=1,264	-0.097 [0.067] N=1,112	**-0.155** [0.078] N=913**	-0.145 [0.106] N=709
Underweight: WAZ below -2	0.054 [0.034] N=1,600	0.046 [0.038] N=1,457	0.026 [0.042] N=1,283	0.045 [0.050] N=1,051	0.098 [0.070] N=813
Wasted: WLZ/WHZ below -2, 0-60 months	-0.049 [0.045] N=1,366	-0.037 [0.047] N=1,248	-0.047 [0.052] N=1,098	-0.029 [0.067] N=902	-0.083 [0.079] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.049] N=1,366	-0.002 [0.054] N=1,248	-0.021 [0.057] N=1,098	-0.057 [0.071] N=902	**-0.148* [0.082] N=696**
Obese: WLZ/WHZ above +3, 0-60 months	-0.030 [0.036] N=1,366	-0.030 [0.041] N=1,248	-0.008 [0.042] N=1,098	-0.021 [0.052] N=902	-0.033 [0.070] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.083 [0.056] N=1,320	-0.094 [0.061] N=1,206	-0.098 [0.067] N=1,061	**-0.152* [0.079] N=873**	-0.158 [0.103] N=676
BMI-for-age z score below -2	**-0.082* [0.048] N=1,351**	**-0.089* [0.049] N=1,236**	**-0.093* [0.056] N=1,086**	-0.059 [0.069] N=891	-0.083 [0.086] N=690

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.689	0.071 [0.214] N=1,642	0.010 [0.230] N=1,519	0.016 [0.253] N=1,367	0.159 [0.259] N=1,168	-0.018 [0.292] N=964
WHO weight-for-age z score	-0.218	-0.159 [0.117] N=1,883	-0.156 [0.130] N=1,740	-0.147 [0.137] N=1,566	-0.140 [0.146] N=1,334	**-0.277* [0.158] N=1,096**
WHO weight-for-length/height z score, 0-60 months	0.308	0.214 [0.182] N=1,619	0.211 [0.194] N=1,501	0.221 [0.203] N=1,351	0.101 [0.219] N=1,155	0.185 [0.241] N=949
WHO BMI-for-age z score	0.281	0.271 [0.210] N=1,601	0.285 [0.225] N=1,486	0.289 [0.242] N=1,336	0.117 [0.258] N=1,141	0.182 [0.284] N=940
Stunted: HAZ below -2	0.286 P=458	-0.030 [0.038] N=1,642	-0.031 [0.040] N=1,519	-0.025 [0.042] N=1,367	-0.047 [0.044] N=1,168	-0.013 [0.048] N=964
Underweight: WAZ below -2	0.105 P=191	**0.058** [0.022] N=1,883**	**0.053** [0.024] N=1,740**	**0.043* [0.025] N=1,566**	**0.047* [0.026] N=1,334**	**0.068** [0.031] N=1,096**
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=189	-0.044 [0.033] N=1,619	-0.036 [0.034] N=1,501	-0.042 [0.036] N=1,351	-0.032 [0.040] N=1,155	-0.050 [0.042] N=949
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=306	0.036 [0.042] N=1,619	0.032 [0.045] N=1,501	0.028 [0.047] N=1,351	0.011 [0.052] N=1,155	0.003 [0.059] N=949
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=141	0.009 [0.030] N=1,619	0.014 [0.032] N=1,501	0.029 [0.034] N=1,351	0.024 [0.037] N=1,155	0.032 [0.043] N=949
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=588	-0.046 [0.041] N=1,564	-0.045 [0.043] N=1,450	-0.042 [0.045] N=1,305	-0.064 [0.049] N=1,117	-0.033 [0.054] N=920
BMI-for-age z score below -2	0.135 P=216	-0.044 [0.038] N=1,601	-0.042 [0.040] N=1,486	-0.041 [0.044] N=1,336	-0.016 [0.048] N=1,141	-0.012 [0.052] N=940

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.136 [0.308] N=1,387	0.009 [0.350] N=1,264	0.079 [0.412] N=1,112	0.442 [0.448] N=913	0.218 [0.633] N=709
WHO weight-for-age z score	**-0.286* [0.167] N=1,600**	-0.308 [0.195] N=1,457	-0.312 [0.217] N=1,283	-0.348 [0.268] N=1,051	**-0.761** [0.329] N=813**
WHO weight-for-length/height z score, 0-60 months	0.098 [0.254] N=1,366	0.075 [0.285] N=1,248	0.028 [0.310] N=1,098	-0.274 [0.369] N=902	-0.309 [0.447] N=696
WHO BMI-for-age z score	0.267 [0.280] N=1,351	0.283 [0.316] N=1,236	0.236 [0.358] N=1,086	-0.160 [0.405] N=891	-0.165 [0.487] N=690
Stunted: HAZ below -2	-0.068 [0.054] N=1,387	-0.075 [0.059] N=1,264	-0.082 [0.067] N=1,112	**-0.150** [0.075] N=913**	-0.138 [0.103] N=709
Underweight: WAZ below -2	**0.059* [0.033] N=1,600**	0.054 [0.036] N=1,457	0.032 [0.041] N=1,283	0.042 [0.049] N=1,051	0.097 [0.071] N=813
Wasted: WLZ/WHZ below -2, 0-60 months	-0.045 [0.045] N=1,366	-0.032 [0.048] N=1,248	-0.046 [0.053] N=1,098	-0.029 [0.068] N=902	-0.078 [0.081] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.050] N=1,366	0.003 [0.056] N=1,248	-0.019 [0.058] N=1,098	-0.059 [0.072] N=902	**-0.145* [0.082] N=696**
Obese: WLZ/WHZ above +3, 0-60 months	-0.029 [0.036] N=1,366	-0.027 [0.042] N=1,248	-0.007 [0.043] N=1,098	-0.022 [0.053] N=902	-0.036 [0.069] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.070 [0.055] N=1,320	-0.072 [0.060] N=1,206	-0.085 [0.065] N=1,061	**-0.156** [0.077] N=873**	-0.157 [0.098] N=676
BMI-for-age z score below -2	-0.078 [0.048] N=1,351	-0.081 [0.050] N=1,236	-0.090 [0.057] N=1,086	-0.061 [0.070] N=891	-0.081 [0.087] N=690

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.689	0.050 [0.214] N=1,642	0.001 [0.232] N=1,519	0.007 [0.256] N=1,367	0.145 [0.264] N=1,168	-0.029 [0.296] N=964
WHO weight-for-age z score	-0.218	-0.174 [0.117] N=1,883	-0.179 [0.128] N=1,740	-0.172 [0.137] N=1,566	-0.171 [0.147] N=1,334	**-0.325** [0.162] N=1,096**
WHO weight-for-length/height z score, 0-60 months	0.308	0.218 [0.182] N=1,619	0.204 [0.194] N=1,501	0.213 [0.205] N=1,351	0.101 [0.221] N=1,155	0.177 [0.243] N=949
WHO BMI-for-age z score	0.281	0.274 [0.210] N=1,601	0.277 [0.225] N=1,486	0.283 [0.244] N=1,336	0.119 [0.261] N=1,141	0.177 [0.288] N=940
Stunted: HAZ below -2	0.286 P=458	-0.024 [0.038] N=1,642	-0.028 [0.040] N=1,519	-0.021 [0.042] N=1,367	-0.041 [0.044] N=1,168	-0.008 [0.048] N=964
Underweight: WAZ below -2	0.105 P=191	**0.060*** [0.022] N=1,883**	**0.057** [0.023] N=1,740**	**0.047* [0.025] N=1,566**	**0.053* [0.027] N=1,334**	**0.075** [0.031] N=1,096**
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=189	-0.045 [0.033] N=1,619	-0.036 [0.034] N=1,501	-0.042 [0.036] N=1,351	-0.033 [0.040] N=1,155	-0.053 [0.042] N=949
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=306	0.037 [0.042] N=1,619	0.032 [0.045] N=1,501	0.029 [0.047] N=1,351	0.012 [0.053] N=1,155	0.000 [0.059] N=949
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=141	0.009 [0.030] N=1,619	0.011 [0.032] N=1,501	0.025 [0.034] N=1,351	0.022 [0.037] N=1,155	0.029 [0.043] N=949
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=588	-0.039 [0.041] N=1,564	-0.038 [0.043] N=1,450	-0.031 [0.045] N=1,305	-0.053 [0.049] N=1,117	-0.023 [0.056] N=920
BMI-for-age z score below -2	0.135 P=216	-0.044 [0.039] N=1,601	-0.041 [0.040] N=1,486	-0.038 [0.044] N=1,336	-0.012 [0.048] N=1,141	-0.009 [0.054] N=940

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.115 [0.308] N=1,387	0.008 [0.349] N=1,264	0.089 [0.416] N=1,112	0.473 [0.455] N=913	0.220 [0.632] N=709
WHO weight-for-age z score	**-0.289* [0.168] N=1,600**	-0.315 [0.192] N=1,457	-0.321 [0.215] N=1,283	-0.342 [0.267] N=1,051	**-0.740** [0.327] N=813**
WHO weight-for-length/height z score, 0-60 months	0.097 [0.252] N=1,366	0.065 [0.280] N=1,248	0.014 [0.312] N=1,098	-0.266 [0.377] N=902	-0.264 [0.451] N=696
WHO BMI-for-age z score	0.279 [0.276] N=1,351	0.282 [0.308] N=1,236	0.233 [0.358] N=1,086	-0.135 [0.410] N=891	-0.112 [0.488] N=690
Stunted: HAZ below -2	-0.064 [0.055] N=1,387	-0.077 [0.059] N=1,264	-0.085 [0.067] N=1,112	**-0.152** [0.077] N=913**	-0.131 [0.103] N=709
Underweight: WAZ below -2	**0.061* [0.033] N=1,600**	0.055 [0.036] N=1,457	0.035 [0.041] N=1,283	0.043 [0.050] N=1,051	0.100 [0.071] N=813
Wasted: WLZ/WHZ below -2, 0-60 months	-0.044 [0.045] N=1,366	-0.033 [0.048] N=1,248	-0.045 [0.053] N=1,098	-0.026 [0.068] N=902	-0.078 [0.081] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.049] N=1,366	0.002 [0.055] N=1,248	-0.019 [0.058] N=1,098	-0.056 [0.072] N=902	-0.135 [0.081] N=696
Obese: WLZ/WHZ above +3, 0-60 months	-0.026 [0.036] N=1,366	-0.025 [0.041] N=1,248	-0.004 [0.043] N=1,098	-0.019 [0.053] N=902	-0.032 [0.068] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.064 [0.056] N=1,320	-0.073 [0.061] N=1,206	-0.084 [0.066] N=1,061	**-0.152* [0.079] N=873**	-0.146 [0.103] N=676
BMI-for-age z score below -2	**-0.081* [0.048] N=1,351**	**-0.087* [0.049] N=1,236**	**-0.095* [0.057] N=1,086**	-0.066 [0.069] N=891	-0.090 [0.086] N=690

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.704	0.041 [0.225] N=1,525	-0.016 [0.239] N=1,416	-0.019 [0.263] N=1,271	0.116 [0.270] N=1,095	-0.044 [0.291] N=904
WHO weight-for-age z score	-0.216	-0.068 [0.124] N=1,740	-0.080 [0.135] N=1,613	-0.071 [0.144] N=1,448	-0.075 [0.150] N=1,241	-0.180 [0.167] N=1,022
WHO weight-for-length/height z score, 0-60 months	0.313	**0.344* [0.182] N=1,501**	**0.327* [0.191] N=1,398**	0.328 [0.200] N=1,256	0.222 [0.214] N=1,083	0.364 [0.231] N=890
WHO BMI-for-age z score	0.286	**0.393* [0.213] N=1,485**	0.369 [0.223] N=1,383	0.376 [0.240] N=1,242	0.234 [0.252] N=1,070	0.353 [0.272] N=882
Stunted: HAZ below -2	0.287 P=424	-0.030 [0.042] N=1,525	-0.038 [0.044] N=1,416	-0.028 [0.046] N=1,271	-0.047 [0.049] N=1,095	-0.014 [0.052] N=904
Underweight: WAZ below -2	0.106 P=178	**0.042* [0.024] N=1,740**	0.039 [0.026] N=1,613	0.029 [0.028] N=1,448	0.039 [0.029] N=1,241	**0.059* [0.034] N=1,022**
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=173	**-0.067** [0.033] N=1,501**	**-0.058* [0.033] N=1,398**	**-0.059* [0.035] N=1,256**	-0.049 [0.039] N=1,083	**-0.067* [0.040] N=890**
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=282	0.048 [0.043] N=1,501	0.041 [0.046] N=1,398	0.041 [0.049] N=1,256	0.021 [0.055] N=1,083	0.024 [0.062] N=890
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=134	0.009 [0.032] N=1,501	0.014 [0.034] N=1,398	0.034 [0.036] N=1,256	0.032 [0.039] N=1,083	0.048 [0.045] N=890
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=542	-0.064 [0.046] N=1,451	-0.067 [0.048] N=1,351	-0.058 [0.050] N=1,213	-0.079 [0.056] N=1,048	-0.052 [0.061] N=863
BMI-for-age z score below -2	0.134 P=197	**-0.067* [0.039] N=1,485**	-0.062 [0.040] N=1,383	-0.059 [0.043] N=1,242	-0.035 [0.048] N=1,070	-0.034 [0.051] N=882

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.185 [0.328] N=1,285	0.092 [0.372] N=1,176	0.164 [0.444] N=1,031	0.534 [0.491] N=855	0.351 [0.672] N=664
WHO weight-for-age z score	-0.245 [0.173] N=1,475	-0.295 [0.196] N=1,348	-0.310 [0.220] N=1,183	-0.389 [0.254] N=976	**-0.771** [0.325] N=757**
WHO weight-for-length/height z score, 0-60 months	0.112 [0.261] N=1,263	0.043 [0.284] N=1,160	-0.039 [0.315] N=1,018	-0.273 [0.385] N=845	-0.177 [0.452] N=652
WHO BMI-for-age z score	0.271 [0.296] N=1,250	0.204 [0.323] N=1,148	0.139 [0.369] N=1,007	-0.185 [0.423] N=835	-0.131 [0.497] N=647
Stunted: HAZ below -2	-0.082 [0.060] N=1,285	**-0.107* [0.063] N=1,176**	-0.111 [0.073] N=1,031	**-0.178** [0.084] N=855**	-0.167 [0.113] N=664
Underweight: WAZ below -2	0.045 [0.035] N=1,475	0.042 [0.038] N=1,348	0.025 [0.044] N=1,183	0.055 [0.050] N=976	**0.118* [0.070] N=757**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.072 [0.045] N=1,263	-0.058 [0.047] N=1,160	-0.061 [0.051] N=1,018	-0.057 [0.066] N=845	-0.124 [0.077] N=652
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.050] N=1,263	-0.009 [0.055] N=1,160	-0.025 [0.058] N=1,018	-0.066 [0.073] N=845	-0.125 [0.088] N=652
Obese: WLZ/WHZ above +3, 0-60 months	-0.049 [0.040] N=1,263	-0.048 [0.046] N=1,160	-0.020 [0.048] N=1,018	-0.029 [0.058] N=845	-0.030 [0.073] N=652
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.098 [0.061] N=1,222	**-0.115* [0.065] N=1,122**	**-0.121* [0.072] N=984**	**-0.199** [0.084] N=819**	**-0.212* [0.110] N=634**
BMI-for-age z score below -2	**-0.099** [0.049] N=1,250**	**-0.101** [0.048] N=1,148**	**-0.103* [0.057] N=1,007**	-0.082 [0.070] N=835	-0.122 [0.086] N=647

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.704	0.043 [0.225] N=1,525	-0.013 [0.239] N=1,416	-0.023 [0.263] N=1,271	0.105 [0.273] N=1,095	-0.065 [0.297] N=904
WHO weight-for-age z score	-0.216	-0.071 [0.123] N=1,740	-0.082 [0.134] N=1,613	-0.080 [0.142] N=1,448	-0.090 [0.149] N=1,241	-0.205 [0.168] N=1,022
WHO weight-for-length/height z score, 0-60 months	0.313	**0.325* [0.180] N=1,501**	0.309 [0.189] N=1,398	0.307 [0.199] N=1,256	0.202 [0.217] N=1,083	0.334 [0.238] N=890
WHO BMI-for-age z score	0.286	**0.380* [0.211] N=1,485**	0.357 [0.221] N=1,383	0.365 [0.239] N=1,242	0.225 [0.255] N=1,070	0.340 [0.279] N=882
Stunted: HAZ below -2	0.287 P=424	-0.028 [0.042] N=1,525	-0.035 [0.044] N=1,416	-0.025 [0.046] N=1,271	-0.045 [0.049] N=1,095	-0.013 [0.051] N=904
Underweight: WAZ below -2	0.106 P=178	**0.042* [0.024] N=1,740**	0.040 [0.025] N=1,613	0.031 [0.027] N=1,448	0.041 [0.028] N=1,241	**0.061* [0.034] N=1,022**
Wasted: WLZ/WHZ below -2, 0-60 months	0.114 P=173	**-0.065** [0.033] N=1,501**	**-0.056* [0.033] N=1,398**	-0.058 [0.035] N=1,256	-0.049 [0.040] N=1,083	**-0.071* [0.042] N=890**
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=282	0.046 [0.043] N=1,501	0.039 [0.045] N=1,398	0.037 [0.048] N=1,256	0.016 [0.054] N=1,083	0.018 [0.062] N=890
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=134	0.007 [0.032] N=1,501	0.013 [0.034] N=1,398	0.033 [0.036] N=1,256	0.031 [0.040] N=1,083	0.044 [0.045] N=890
CIAF: any stunting, underweight, or wasting, 0-60 months	0.382 P=542	-0.061 [0.046] N=1,451	-0.064 [0.048] N=1,351	-0.056 [0.050] N=1,213	-0.079 [0.055] N=1,048	-0.055 [0.060] N=863
BMI-for-age z score below -2	0.134 P=197	-0.063 [0.039] N=1,485	-0.059 [0.040] N=1,383	-0.058 [0.044] N=1,242	-0.035 [0.049] N=1,070	-0.037 [0.054] N=882

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.191 [0.331] N=1,285	0.095 [0.376] N=1,176	0.166 [0.447] N=1,031	0.519 [0.504] N=855	0.343 [0.691] N=664
WHO weight-for-age z score	-0.209 [0.171] N=1,475	-0.256 [0.195] N=1,348	-0.284 [0.217] N=1,183	-0.378 [0.256] N=976	**-0.789** [0.337] N=757**
WHO weight-for-length/height z score, 0-60 months	0.138 [0.258] N=1,263	0.080 [0.282] N=1,160	-0.011 [0.315] N=1,018	-0.272 [0.390] N=845	-0.223 [0.467] N=652
WHO BMI-for-age z score	0.284 [0.291] N=1,250	0.218 [0.318] N=1,148	0.151 [0.367] N=1,007	-0.184 [0.425] N=835	-0.146 [0.503] N=647
Stunted: HAZ below -2	-0.077 [0.060] N=1,285	-0.099 [0.064] N=1,176	-0.106 [0.073] N=1,031	**-0.172** [0.085] N=855**	-0.172 [0.114] N=664
Underweight: WAZ below -2	0.042 [0.035] N=1,475	0.038 [0.039] N=1,348	0.021 [0.043] N=1,183	0.050 [0.050] N=976	0.109 [0.071] N=757
Wasted: WLZ/WHZ below -2, 0-60 months	-0.070 [0.045] N=1,263	-0.058 [0.047] N=1,160	-0.062 [0.051] N=1,018	-0.058 [0.067] N=845	-0.130 [0.078] N=652
Overweight: WLZ/WHZ above +2, 0-60 months	0.020 [0.050] N=1,263	-0.000 [0.054] N=1,160	-0.019 [0.058] N=1,018	-0.064 [0.074] N=845	-0.135 [0.087] N=652
Obese: WLZ/WHZ above +3, 0-60 months	-0.047 [0.039] N=1,263	-0.046 [0.045] N=1,160	-0.018 [0.047] N=1,018	-0.028 [0.058] N=845	-0.038 [0.073] N=652
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.091 [0.060] N=1,222	**-0.107* [0.064] N=1,122**	-0.116 [0.071] N=984	**-0.194** [0.083] N=819**	**-0.223** [0.107] N=634**
BMI-for-age z score below -2	**-0.092* [0.049] N=1,250**	**-0.095* [0.049] N=1,148**	**-0.100* [0.057] N=1,007**	-0.079 [0.071] N=835	-0.124 [0.087] N=647

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.619	-0.209 [0.263] N=1,459	-0.229 [0.274] N=1,320	-0.192 [0.290] N=1,151	-0.315 [0.322] N=976	-0.515 [0.373] N=790
WHO weight-for-age z score	-0.209	0.055 [0.134] N=1,664	0.091 [0.141] N=1,514	-0.020 [0.145] N=1,318	0.062 [0.153] N=1,115	0.013 [0.174] N=897
WHO weight-for-length/height z score, 0-60 months	0.349	0.086 [0.212] N=1,446	0.032 [0.222] N=1,311	-0.052 [0.248] N=1,144	0.119 [0.271] N=970	0.103 [0.313] N=784
WHO BMI-for-age z score	0.298	0.127 [0.217] N=1,426	0.112 [0.221] N=1,293	0.035 [0.248] N=1,129	0.167 [0.281] N=956	0.170 [0.304] N=776
Stunted: HAZ below -2	0.276 P=401	0.051 [0.057] N=1,459	0.041 [0.058] N=1,320	0.033 [0.061] N=1,151	0.059 [0.066] N=976	0.080 [0.074] N=790
Underweight: WAZ below -2	0.106 P=169	-0.015 [0.029] N=1,664	-0.018 [0.030] N=1,514	0.001 [0.032] N=1,318	-0.007 [0.034] N=1,115	-0.012 [0.037] N=897
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=168	0.034 [0.037] N=1,446	0.042 [0.039] N=1,311	0.064 [0.043] N=1,144	0.016 [0.044] N=970	0.009 [0.051] N=784
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=283	0.062 [0.042] N=1,446	0.059 [0.044] N=1,311	0.043 [0.047] N=1,144	0.059 [0.053] N=970	0.052 [0.058] N=784
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=135	0.036 [0.035] N=1,446	0.023 [0.036] N=1,311	0.017 [0.039] N=1,144	0.009 [0.045] N=970	-0.021 [0.047] N=784
CIAF: any stunting, underweight, or wasting, 0-60 months	0.376 P=521	**0.090* [0.054] N=1,393**	0.081 [0.055] N=1,259	0.094 [0.058] N=1,098	0.070 [0.060] N=930	0.092 [0.066] N=751
BMI-for-age z score below -2	0.135 P=193	0.033 [0.040] N=1,426	0.036 [0.042] N=1,293	0.040 [0.048] N=1,129	0.006 [0.051] N=956	0.004 [0.056] N=776

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.105 [0.385] N=1,260	-0.122 [0.411] N=1,121	0.001 [0.465] N=952	-0.282 [0.558] N=777	-0.729 [0.752] N=591
WHO weight-for-age z score	0.082 [0.193] N=1,440	0.126 [0.208] N=1,290	-0.083 [0.217] N=1,094	0.050 [0.222] N=891	0.007 [0.285] N=673
WHO weight-for-length/height z score, 0-60 months	-0.019 [0.295] N=1,245	-0.131 [0.311] N=1,110	-0.324 [0.361] N=943	0.004 [0.401] N=769	-0.081 [0.505] N=583
WHO BMI-for-age z score	-0.066 [0.288] N=1,234	-0.115 [0.295] N=1,101	-0.271 [0.359] N=937	0.001 [0.423] N=764	-0.011 [0.429] N=584
Stunted: HAZ below -2	0.044 [0.089] N=1,260	0.026 [0.093] N=1,121	0.002 [0.104] N=952	0.064 [0.124] N=777	0.132 [0.160] N=591
Underweight: WAZ below -2	-0.023 [0.044] N=1,440	-0.026 [0.047] N=1,290	0.007 [0.053] N=1,094	-0.006 [0.059] N=891	-0.023 [0.068] N=673
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.049] N=1,245	**0.091* [0.053] N=1,110**	**0.144** [0.058] N=943**	0.056 [0.059] N=769	0.050 [0.076] N=583
Overweight: WLZ/WHZ above +2, 0-60 months	**0.098* [0.058] N=1,245**	0.092 [0.062] N=1,110	0.073 [0.067] N=943	**0.132* [0.078] N=769**	0.131 [0.099] N=583
Obese: WLZ/WHZ above +3, 0-60 months	**0.074* [0.043] N=1,245**	0.048 [0.044] N=1,110	0.041 [0.049] N=943	0.028 [0.063] N=769	-0.065 [0.071] N=583
CIAF: any stunting, underweight, or wasting, 0-60 months	0.122 [0.086] N=1,201	0.110 [0.090] N=1,067	0.135 [0.103] N=906	0.101 [0.118] N=738	0.170 [0.146] N=559
BMI-for-age z score below -2	0.070 [0.048] N=1,234	0.078 [0.053] N=1,101	0.085 [0.065] N=937	0.009 [0.066] N=764	-0.003 [0.077] N=584

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.619	-0.210 [0.263] N=1,459	-0.232 [0.273] N=1,320	-0.197 [0.288] N=1,151	-0.318 [0.322] N=976	-0.519 [0.373] N=790
WHO weight-for-age z score	-0.209	0.062 [0.134] N=1,664	0.096 [0.141] N=1,514	-0.016 [0.145] N=1,318	0.064 [0.152] N=1,115	0.015 [0.174] N=897
WHO weight-for-length/height z score, 0-60 months	0.349	0.091 [0.212] N=1,446	0.035 [0.222] N=1,311	-0.050 [0.248] N=1,144	0.120 [0.271] N=970	0.107 [0.312] N=784
WHO BMI-for-age z score	0.298	0.129 [0.217] N=1,426	0.112 [0.222] N=1,293	0.032 [0.248] N=1,129	0.166 [0.281] N=956	0.170 [0.305] N=776
Stunted: HAZ below -2	0.276 P=401	0.050 [0.057] N=1,459	0.040 [0.058] N=1,320	0.032 [0.061] N=1,151	0.059 [0.066] N=976	0.080 [0.074] N=790
Underweight: WAZ below -2	0.106 P=169	-0.016 [0.029] N=1,664	-0.019 [0.030] N=1,514	0.000 [0.032] N=1,318	-0.007 [0.034] N=1,115	-0.013 [0.037] N=897
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=168	0.032 [0.037] N=1,446	0.039 [0.039] N=1,311	0.062 [0.043] N=1,144	0.015 [0.044] N=970	0.008 [0.050] N=784
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=283	0.061 [0.042] N=1,446	0.057 [0.044] N=1,311	0.040 [0.047] N=1,144	0.057 [0.053] N=970	0.051 [0.058] N=784
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=135	0.035 [0.035] N=1,446	0.022 [0.036] N=1,311	0.015 [0.039] N=1,144	0.008 [0.045] N=970	-0.022 [0.048] N=784
CIAF: any stunting, underweight, or wasting, 0-60 months	0.376 P=521	0.087 [0.055] N=1,393	0.076 [0.056] N=1,259	0.089 [0.058] N=1,098	0.069 [0.060] N=930	0.091 [0.067] N=751
BMI-for-age z score below -2	0.135 P=193	0.031 [0.040] N=1,426	0.035 [0.042] N=1,293	0.039 [0.048] N=1,129	0.006 [0.051] N=956	0.004 [0.056] N=776

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.108 [0.384] N=1,260	-0.136 [0.409] N=1,121	-0.027 [0.458] N=952	-0.314 [0.552] N=777	-0.793 [0.744] N=591
WHO weight-for-age z score	0.098 [0.190] N=1,440	0.142 [0.205] N=1,290	-0.074 [0.213] N=1,094	0.064 [0.217] N=891	0.033 [0.279] N=673
WHO weight-for-length/height z score, 0-60 months	-0.005 [0.297] N=1,245	-0.121 [0.314] N=1,110	-0.315 [0.363] N=943	0.019 [0.405] N=769	-0.024 [0.518] N=583
WHO BMI-for-age z score	-0.055 [0.291] N=1,234	-0.100 [0.298] N=1,101	-0.263 [0.361] N=937	0.010 [0.426] N=764	0.042 [0.447] N=584
Stunted: HAZ below -2	0.041 [0.089] N=1,260	0.024 [0.092] N=1,121	-0.001 [0.103] N=952	0.064 [0.123] N=777	0.139 [0.158] N=591
Underweight: WAZ below -2	-0.025 [0.044] N=1,440	-0.028 [0.047] N=1,290	0.006 [0.053] N=1,094	-0.007 [0.059] N=891	-0.024 [0.069] N=673
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.050] N=1,245	0.082 [0.054] N=1,110	**0.133** [0.059] N=943**	0.046 [0.060] N=769	0.030 [0.080] N=583
Overweight: WLZ/WHZ above +2, 0-60 months	**0.097* [0.058] N=1,245**	0.088 [0.062] N=1,110	0.066 [0.066] N=943	0.124 [0.077] N=769	0.124 [0.098] N=583
Obese: WLZ/WHZ above +3, 0-60 months	0.071 [0.043] N=1,245	0.044 [0.044] N=1,110	0.034 [0.049] N=943	0.021 [0.063] N=769	-0.075 [0.071] N=583
CIAF: any stunting, underweight, or wasting, 0-60 months	0.112 [0.087] N=1,201	0.096 [0.090] N=1,067	0.118 [0.102] N=906	0.088 [0.119] N=738	0.151 [0.146] N=559
BMI-for-age z score below -2	0.065 [0.049] N=1,234	0.071 [0.053] N=1,101	0.080 [0.065] N=937	0.005 [0.067] N=764	-0.010 [0.080] N=584

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.619	-0.211 [0.258] N=1,459	-0.230 [0.269] N=1,320	-0.182 [0.287] N=1,151	-0.302 [0.318] N=976	-0.522 [0.366] N=790
WHO weight-for-age z score	-0.209	0.066 [0.132] N=1,664	0.103 [0.140] N=1,514	0.005 [0.144] N=1,318	0.080 [0.153] N=1,115	0.022 [0.174] N=897
WHO weight-for-length/height z score, 0-60 months	0.349	0.100 [0.212] N=1,446	0.045 [0.222] N=1,311	-0.031 [0.249] N=1,144	0.123 [0.270] N=970	0.113 [0.313] N=784
WHO BMI-for-age z score	0.298	0.134 [0.218] N=1,426	0.119 [0.223] N=1,293	0.054 [0.248] N=1,129	0.178 [0.278] N=956	0.183 [0.304] N=776
Stunted: HAZ below -2	0.276 P=401	0.051 [0.056] N=1,459	0.040 [0.057] N=1,320	0.027 [0.060] N=1,151	0.052 [0.064] N=976	0.079 [0.071] N=790
Underweight: WAZ below -2	0.106 P=169	-0.017 [0.028] N=1,664	-0.020 [0.030] N=1,514	-0.003 [0.032] N=1,318	-0.011 [0.034] N=1,115	-0.014 [0.036] N=897
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=168	0.031 [0.038] N=1,446	0.037 [0.040] N=1,311	0.059 [0.043] N=1,144	0.014 [0.044] N=970	0.009 [0.051] N=784
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=283	0.062 [0.042] N=1,446	0.058 [0.044] N=1,311	0.040 [0.047] N=1,144	0.054 [0.053] N=970	0.052 [0.059] N=784
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=135	0.036 [0.035] N=1,446	0.022 [0.036] N=1,311	0.015 [0.039] N=1,144	0.006 [0.045] N=970	-0.022 [0.048] N=784
CIAF: any stunting, underweight, or wasting, 0-60 months	0.376 P=521	**0.090* [0.054] N=1,393**	0.077 [0.055] N=1,259	0.083 [0.058] N=1,098	0.062 [0.060] N=930	0.089 [0.065] N=751
BMI-for-age z score below -2	0.135 P=193	0.031 [0.040] N=1,426	0.034 [0.042] N=1,293	0.035 [0.048] N=1,129	0.002 [0.051] N=956	0.003 [0.057] N=776

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.107 [0.378] N=1,260	-0.134 [0.400] N=1,121	-0.001 [0.449] N=952	-0.284 [0.537] N=777	-0.768 [0.733] N=591
WHO weight-for-age z score	0.086 [0.188] N=1,440	0.132 [0.203] N=1,290	-0.045 [0.211] N=1,094	0.082 [0.218] N=891	0.033 [0.269] N=673
WHO weight-for-length/height z score, 0-60 months	-0.027 [0.295] N=1,245	-0.140 [0.312] N=1,110	-0.312 [0.361] N=943	-0.000 [0.400] N=769	-0.041 [0.513] N=583
WHO BMI-for-age z score	-0.076 [0.288] N=1,234	-0.121 [0.296] N=1,101	-0.258 [0.355] N=937	-0.013 [0.417] N=764	0.029 [0.438] N=584
Stunted: HAZ below -2	0.042 [0.087] N=1,260	0.024 [0.090] N=1,121	-0.006 [0.099] N=952	0.056 [0.119] N=777	0.135 [0.151] N=591
Underweight: WAZ below -2	-0.024 [0.043] N=1,440	-0.027 [0.047] N=1,290	-0.001 [0.053] N=1,094	-0.012 [0.059] N=891	-0.027 [0.067] N=673
Wasted: WLZ/WHZ below -2, 0-60 months	0.070 [0.050] N=1,245	0.083 [0.054] N=1,110	**0.131** [0.060] N=943**	0.046 [0.060] N=769	0.033 [0.080] N=583
Overweight: WLZ/WHZ above +2, 0-60 months	0.096 [0.058] N=1,245	0.087 [0.062] N=1,110	0.062 [0.067] N=943	0.116 [0.078] N=769	0.120 [0.098] N=583
Obese: WLZ/WHZ above +3, 0-60 months	0.069 [0.043] N=1,245	0.041 [0.044] N=1,110	0.031 [0.049] N=943	0.016 [0.063] N=769	-0.079 [0.070] N=583
CIAF: any stunting, underweight, or wasting, 0-60 months	0.116 [0.086] N=1,201	0.099 [0.090] N=1,067	0.111 [0.101] N=906	0.082 [0.117] N=738	0.152 [0.142] N=559
BMI-for-age z score below -2	0.065 [0.049] N=1,234	0.072 [0.053] N=1,101	0.074 [0.063] N=937	0.002 [0.067] N=764	-0.011 [0.081] N=584

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.619	-0.198 [0.256] N=1,459	-0.211 [0.266] N=1,320	-0.159 [0.283] N=1,151	-0.284 [0.313] N=976	-0.517 [0.365] N=790
WHO weight-for-age z score	-0.209	0.077 [0.134] N=1,664	0.116 [0.142] N=1,514	0.022 [0.147] N=1,318	0.099 [0.156] N=1,115	0.036 [0.177] N=897
WHO weight-for-length/height z score, 0-60 months	0.349	0.099 [0.212] N=1,446	0.044 [0.223] N=1,311	-0.031 [0.248] N=1,144	0.145 [0.270] N=970	0.148 [0.312] N=784
WHO BMI-for-age z score	0.298	0.131 [0.218] N=1,426	0.117 [0.223] N=1,293	0.048 [0.247] N=1,129	0.186 [0.278] N=956	0.199 [0.305] N=776
Stunted: HAZ below -2	0.276 P=401	0.047 [0.055] N=1,459	0.036 [0.057] N=1,320	0.024 [0.059] N=1,151	0.051 [0.064] N=976	0.076 [0.071] N=790
Underweight: WAZ below -2	0.106 P=169	-0.019 [0.029] N=1,664	-0.022 [0.030] N=1,514	-0.005 [0.032] N=1,318	-0.012 [0.034] N=1,115	-0.017 [0.038] N=897
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=168	0.031 [0.038] N=1,446	0.039 [0.039] N=1,311	0.061 [0.043] N=1,144	0.013 [0.043] N=970	0.005 [0.050] N=784
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=283	0.060 [0.041] N=1,446	0.056 [0.044] N=1,311	0.041 [0.047] N=1,144	0.057 [0.052] N=970	0.055 [0.057] N=784
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=135	0.041 [0.034] N=1,446	0.027 [0.035] N=1,311	0.020 [0.038] N=1,144	0.013 [0.044] N=970	-0.016 [0.047] N=784
CIAF: any stunting, underweight, or wasting, 0-60 months	0.376 P=521	0.084 [0.054] N=1,393	0.073 [0.055] N=1,259	0.081 [0.058] N=1,098	0.060 [0.059] N=930	0.086 [0.065] N=751
BMI-for-age z score below -2	0.135 P=193	0.029 [0.040] N=1,426	0.032 [0.042] N=1,293	0.035 [0.047] N=1,129	0.002 [0.050] N=956	-0.001 [0.055] N=776

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.081 [0.366] N=1,260	-0.087 [0.388] N=1,121	0.082 [0.435] N=952	-0.210 [0.525] N=777	-0.773 [0.720] N=591
WHO weight-for-age z score	0.127 [0.183] N=1,440	0.177 [0.199] N=1,290	0.008 [0.206] N=1,094	0.131 [0.211] N=891	0.061 [0.256] N=673
WHO weight-for-length/height z score, 0-60 months	-0.014 [0.294] N=1,245	-0.137 [0.311] N=1,110	-0.322 [0.358] N=943	0.012 [0.402] N=769	0.012 [0.528] N=583
WHO BMI-for-age z score	-0.065 [0.288] N=1,234	-0.117 [0.297] N=1,101	-0.267 [0.357] N=937	0.010 [0.420] N=764	0.082 [0.454] N=584
Stunted: HAZ below -2	0.037 [0.085] N=1,260	0.017 [0.088] N=1,121	-0.016 [0.098] N=952	0.050 [0.119] N=777	0.146 [0.152] N=591
Underweight: WAZ below -2	-0.029 [0.042] N=1,440	-0.031 [0.046] N=1,290	-0.006 [0.052] N=1,094	-0.015 [0.058] N=891	-0.029 [0.066] N=673
Wasted: WLZ/WHZ below -2, 0-60 months	0.072 [0.050] N=1,245	0.087 [0.054] N=1,110	**0.139** [0.058] N=943**	0.051 [0.059] N=769	0.037 [0.079] N=583
Overweight: WLZ/WHZ above +2, 0-60 months	0.094 [0.057] N=1,245	0.083 [0.061] N=1,110	0.061 [0.066] N=943	0.115 [0.078] N=769	0.129 [0.099] N=583
Obese: WLZ/WHZ above +3, 0-60 months	**0.075* [0.043] N=1,245**	0.048 [0.044] N=1,110	0.037 [0.049] N=943	0.022 [0.064] N=769	-0.069 [0.072] N=583
CIAF: any stunting, underweight, or wasting, 0-60 months	0.110 [0.085] N=1,201	0.094 [0.089] N=1,067	0.105 [0.102] N=906	0.076 [0.117] N=738	0.166 [0.141] N=559
BMI-for-age z score below -2	0.065 [0.049] N=1,234	0.072 [0.054] N=1,101	0.078 [0.064] N=937	0.006 [0.067] N=764	-0.005 [0.080] N=584

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.627	-0.245 [0.276] N=1,353	-0.275 [0.293] N=1,225	-0.205 [0.307] N=1,072	-0.333 [0.336] N=907	-0.553 [0.379] N=736
WHO weight-for-age z score	-0.213	0.072 [0.141] N=1,537	0.117 [0.151] N=1,398	-0.016 [0.153] N=1,222	0.041 [0.161] N=1,035	-0.054 [0.177] N=836
WHO weight-for-length/height z score, 0-60 months	0.346	0.138 [0.224] N=1,339	0.094 [0.238] N=1,215	-0.042 [0.259] N=1,062	0.092 [0.281] N=900	0.046 [0.314] N=730
WHO BMI-for-age z score	0.294	0.138 [0.234] N=1,321	0.145 [0.243] N=1,199	-0.007 [0.269] N=1,050	0.099 [0.304] N=889	0.051 [0.328] N=724
Stunted: HAZ below -2	0.276 P=370	0.048 [0.060] N=1,353	0.039 [0.062] N=1,225	0.018 [0.064] N=1,072	0.041 [0.067] N=907	0.065 [0.074] N=736
Underweight: WAZ below -2	0.107 P=159	-0.020 [0.031] N=1,537	-0.023 [0.033] N=1,398	-0.006 [0.034] N=1,222	-0.007 [0.036] N=1,035	-0.004 [0.040] N=836
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=155	0.034 [0.038] N=1,339	0.040 [0.040] N=1,215	0.066 [0.043] N=1,062	0.024 [0.044] N=900	0.024 [0.050] N=730
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=261	0.060 [0.043] N=1,339	0.056 [0.045] N=1,215	0.030 [0.048] N=1,062	0.040 [0.053] N=900	0.027 [0.057] N=730
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=127	0.035 [0.035] N=1,339	0.020 [0.035] N=1,215	0.011 [0.037] N=1,062	0.005 [0.043] N=900	-0.023 [0.045] N=730
CIAF: any stunting, underweight, or wasting, 0-60 months	0.375 P=481	0.088 [0.058] N=1,291	0.076 [0.060] N=1,168	0.078 [0.062] N=1,021	0.055 [0.063] N=863	0.091 [0.069] N=699
BMI-for-age z score below -2	0.134 P=176	0.041 [0.042] N=1,321	0.043 [0.045] N=1,199	0.049 [0.050] N=1,050	0.025 [0.053] N=889	0.031 [0.059] N=724

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.116 [0.409] N=1,164	-0.154 [0.444] N=1,036	0.001 [0.502] N=883	-0.309 [0.589] N=718	-0.854 [0.775] N=547
WHO weight-for-age z score	0.159 [0.198] N=1,326	0.226 [0.214] N=1,187	-0.027 [0.222] N=1,011	0.097 [0.230] N=824	0.019 [0.292] N=625
WHO weight-for-length/height z score, 0-60 months	0.073 [0.327] N=1,149	-0.010 [0.353] N=1,025	-0.300 [0.408] N=872	-0.009 [0.444] N=710	-0.065 [0.578] N=540
WHO BMI-for-age z score	-0.007 [0.312] N=1,138	-0.006 [0.331] N=1,016	-0.282 [0.401] N=867	-0.033 [0.457] N=706	-0.053 [0.480] N=541
Stunted: HAZ below -2	0.038 [0.096] N=1,164	0.023 [0.101] N=1,036	-0.026 [0.111] N=883	0.020 [0.128] N=718	0.089 [0.160] N=547
Underweight: WAZ below -2	-0.038 [0.047] N=1,326	-0.043 [0.051] N=1,187	-0.015 [0.057] N=1,011	-0.018 [0.064] N=824	-0.034 [0.076] N=625
Wasted: WLZ/WHZ below -2, 0-60 months	0.066 [0.052] N=1,149	0.075 [0.057] N=1,025	**0.130** [0.064]** **N=872**	0.046 [0.064] N=710	0.031 [0.082] N=540
Overweight: WLZ/WHZ above +2, 0-60 months	0.102 [0.063] N=1,149	0.094 [0.067] N=1,025	0.059 [0.073] N=872	0.104 [0.084] N=710	0.092 [0.108] N=540
Obese: WLZ/WHZ above +3, 0-60 months	0.068 [0.047] N=1,149	0.039 [0.049] N=1,025	0.026 [0.054] N=872	0.014 [0.069] N=710	-0.084 [0.080] N=540
CIAF: any stunting, underweight, or wasting, 0-60 months	0.110 [0.095] N=1,109	0.089 [0.101] N=986	0.092 [0.113] N=839	0.037 [0.131] N=681	0.112 [0.156] N=517
BMI-for-age z score below -2	0.076 [0.051] N=1,138	0.078 [0.058] N=1,016	0.089 [0.070] N=867	0.028 [0.072] N=706	0.028 [0.086] N=541

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.627	-0.262 [0.279] N=1,353	-0.287 [0.295] N=1,225	-0.213 [0.310] N=1,072	-0.349 [0.338] N=907	-0.583 [0.384] N=736
WHO weight-for-age z score	-0.213	0.073 [0.141] N=1,537	0.115 [0.151] N=1,398	-0.014 [0.153] N=1,222	0.041 [0.161] N=1,035	-0.054 [0.178] N=836
WHO weight-for-length/height z score, 0-60 months	0.346	0.156 [0.225] N=1,339	0.110 [0.238] N=1,215	-0.025 [0.260] N=1,062	0.111 [0.276] N=900	0.072 [0.315] N=730
WHO BMI-for-age z score	0.294	0.157 [0.235] N=1,321	0.159 [0.244] N=1,199	0.009 [0.270] N=1,050	0.113 [0.301] N=889	0.081 [0.328] N=724
Stunted: HAZ below -2	0.276 P=370	0.047 [0.059] N=1,353	0.037 [0.062] N=1,225	0.016 [0.064] N=1,072	0.037 [0.067] N=907	0.061 [0.074] N=736
Underweight: WAZ below -2	0.107 P=159	-0.022 [0.031] N=1,537	-0.025 [0.033] N=1,398	-0.008 [0.034] N=1,222	-0.010 [0.036] N=1,035	-0.007 [0.039] N=836
Wasted: WLZ/WHZ below -2, 0-60 months	0.115 P=155	0.029 [0.038] N=1,339	0.035 [0.040] N=1,215	0.060 [0.044] N=1,062	0.017 [0.044] N=900	0.014 [0.050] N=730
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=261	0.060 [0.042] N=1,339	0.055 [0.044] N=1,215	0.027 [0.047] N=1,062	0.037 [0.052] N=900	0.027 [0.056] N=730
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=127	0.035 [0.034] N=1,339	0.020 [0.035] N=1,215	0.011 [0.037] N=1,062	0.004 [0.043] N=900	-0.023 [0.044] N=730
CIAF: any stunting, underweight, or wasting, 0-60 months	0.375 P=481	0.083 [0.057] N=1,291	0.069 [0.059] N=1,168	0.069 [0.060] N=1,021	0.042 [0.062] N=863	0.074 [0.067] N=699
BMI-for-age z score below -2	0.134 P=176	0.036 [0.042] N=1,321	0.037 [0.045] N=1,199	0.043 [0.050] N=1,050	0.018 [0.053] N=889	0.021 [0.058] N=724

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.127 [0.409] N=1,164	-0.154 [0.442] N=1,036	0.009 [0.496] N=883	-0.345 [0.586] N=718	-0.932 [0.763] N=547
WHO weight-for-age z score	0.152 [0.198] N=1,326	0.216 [0.217] N=1,187	-0.034 [0.224] N=1,011	0.082 [0.230] N=824	-0.015 [0.293] N=625
WHO weight-for-length/height z score, 0-60 months	0.064 [0.324] N=1,149	-0.026 [0.349] N=1,025	-0.320 [0.401] N=872	-0.024 [0.434] N=710	-0.056 [0.559] N=540
WHO BMI-for-age z score	0.003 [0.311] N=1,138	-0.005 [0.327] N=1,016	-0.282 [0.396] N=867	-0.020 [0.451] N=706	0.013 [0.478] N=541
Stunted: HAZ below -2	0.038 [0.095] N=1,164	0.021 [0.100] N=1,036	-0.029 [0.110] N=883	0.023 [0.128] N=718	0.095 [0.157] N=547
Underweight: WAZ below -2	-0.038 [0.047] N=1,326	-0.044 [0.051] N=1,187	-0.016 [0.057] N=1,011	-0.018 [0.064] N=824	-0.026 [0.075] N=625
Wasted: WLZ/WHZ below -2, 0-60 months	0.067 [0.051] N=1,149	0.077 [0.057] N=1,025	**0.132** [0.063] N=872**	0.046 [0.063] N=710	0.031 [0.081] N=540
Overweight: WLZ/WHZ above +2, 0-60 months	0.099 [0.062] N=1,149	0.091 [0.066] N=1,025	0.052 [0.071] N=872	0.098 [0.082] N=710	0.096 [0.105] N=540
Obese: WLZ/WHZ above +3, 0-60 months	0.065 [0.047] N=1,149	0.037 [0.048] N=1,025	0.022 [0.052] N=872	0.010 [0.066] N=710	-0.082 [0.075] N=540
CIAF: any stunting, underweight, or wasting, 0-60 months	0.111 [0.095] N=1,109	0.089 [0.100] N=986	0.089 [0.111] N=839	0.033 [0.130] N=681	0.104 [0.154] N=517
BMI-for-age z score below -2	0.074 [0.051] N=1,138	0.077 [0.057] N=1,016	0.089 [0.070] N=867	0.026 [0.071] N=706	0.018 [0.086] N=541

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.