Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-6

WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height and related classifications are available through month 60 only. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.594	0.227 [0.164] N=2,676	0.202 [0.176] N=2,470	0.227 [0.191] N=2,212	**0.396** [0.192] N=1,901**	0.349 [0.221] N=1,571
WHO weight-for-age z score	-0.082	-0.021 [0.096] N=2,964	-0.013 [0.106] N=2,736	0.031 [0.106] N=2,452	0.001 [0.118] N=2,104	-0.017 [0.123] N=1,734
WHO weight-for-length/height z score, 0-60 months	0.329	0.086 [0.155] N=1,994	0.044 [0.159] N=1,853	-0.006 [0.170] N=1,668	-0.153 [0.185] N=1,430	-0.057 [0.200] N=1,171
WHO BMI-for-age z score	0.346	0.030 [0.154] N=2,592	0.006 [0.158] N=2,394	-0.007 [0.173] N=2,139	-0.245 [0.177] N=1,839	-0.189 [0.193] N=1,518
Stunted: HAZ below -2	0.257 P=671	-0.043 [0.031] N=2,676	-0.047 [0.033] N=2,470	-0.046 [0.033] N=2,212	**-0.082** [0.035] N=1,901**	-0.050 [0.040] N=1,571
Underweight: WAZ below -2	0.088 P=250	0.030 [0.020] N=2,964	0.021 [0.022] N=2,736	0.003 [0.023] N=2,452	0.005 [0.024] N=2,104	0.011 [0.027] N=1,734
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=227	-0.034 [0.028] N=1,994	-0.025 [0.028] N=1,853	-0.018 [0.031] N=1,668	-0.011 [0.034] N=1,430	-0.034 [0.035] N=1,171
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=382	0.016 [0.038] N=1,994	0.008 [0.040] N=1,853	0.003 [0.041] N=1,668	-0.020 [0.046] N=1,430	-0.031 [0.051] N=1,171
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=179	-0.014 [0.027] N=1,994	-0.017 [0.029] N=1,853	-0.005 [0.031] N=1,668	-0.015 [0.034] N=1,430	-0.018 [0.040] N=1,171
CIAF: any stunting, underweight, or wasting, 0-60 months	0.363 P=694	**-0.063* [0.035] N=1,935**	**-0.067* [0.037] N=1,798**	-0.055 [0.038] N=1,618	**-0.085** [0.041] N=1,388**	-0.064 [0.043] N=1,139
BMI-for-age z score below -2	0.121 P=315	-0.009 [0.028] N=2,592	-0.010 [0.028] N=2,394	-0.008 [0.032] N=2,139	0.014 [0.034] N=1,839	0.013 [0.037] N=1,518
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=228	-0.040 [0.074] N=599	-0.041 [0.078] N=543	-0.042 [0.082] N=477	-0.105 [0.088] N=416	-0.128 [0.104] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=137	-0.032 [0.064] N=599	-0.048 [0.067] N=543	-0.021 [0.073] N=477	-0.072 [0.080] N=416	-0.122 [0.093] N=348
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=60	0.060 [0.069] N=599	0.041 [0.073] N=543	0.019 [0.078] N=477	0.050 [0.082] N=416	0.027 [0.102] N=348

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.060 [0.218] N=2,249	-0.171 [0.241] N=2,043	-0.179 [0.284] N=1,785	0.073 [0.293] N=1,474	-0.185 [0.392] N=1,144
WHO weight-for-age z score	-0.191 [0.134] N=2,497	-0.204 [0.155] N=2,269	-0.144 [0.161] N=1,985	-0.266 [0.197] N=1,637	**-0.400* [0.231] N=1,267**
WHO weight-for-length/height z score, 0-60 months	0.106 [0.230] N=1,680	0.042 [0.245] N=1,539	-0.073 [0.276] N=1,354	-0.382 [0.339] N=1,116	-0.283 [0.425] N=857
WHO BMI-for-age z score	0.179 [0.223] N=2,176	0.179 [0.234] N=1,978	0.203 [0.280] N=1,723	-0.219 [0.313] N=1,423	-0.091 [0.399] N=1,102
Stunted: HAZ below -2	-0.006 [0.041] N=2,249	-0.006 [0.046] N=2,043	-0.003 [0.050] N=1,785	-0.068 [0.055] N=1,474	0.022 [0.072] N=1,144
Underweight: WAZ below -2	**0.046* [0.028] N=2,497**	0.033 [0.031] N=2,269	-0.003 [0.035] N=1,985	0.005 [0.040] N=1,637	0.026 [0.054] N=1,267

Outcome	8 years	7 years	6 years	5 years	4 years
Wasted: WLZ/WHZ below -2, 0-60 months	-0.032 [0.038] N=1,680	-0.017 [0.039] N=1,539	-0.003 [0.045] N=1,354	0.012 [0.055] N=1,116	-0.040 [0.064] N=857
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.045] N=1,680	0.014 [0.050] N=1,539	0.001 [0.052] N=1,354	-0.036 [0.066] N=1,116	-0.097 [0.084] N=857
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.031] N=1,680	-0.029 [0.034] N=1,539	-0.007 [0.036] N=1,354	-0.026 [0.044] N=1,116	-0.047 [0.062] N=857
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.039 [0.048] N=1,632	-0.043 [0.055] N=1,495	-0.027 [0.059] N=1,315	-0.084 [0.069] N=1,085	-0.044 [0.083] N=836
BMI-for-age z score below -2	-0.033 [0.036] N=2,176	-0.042 [0.035] N=1,978	-0.052 [0.041] N=1,723	-0.025 [0.045] N=1,423	-0.055 [0.058] N=1,102
WHO 2007 BMI overweight above +1, 61+ months	0.008 [0.119] N=496	0.036 [0.130] N=440	0.054 [0.140] N=374	-0.049 [0.158] N=313	-0.125 [0.217] N=245
WHO 2007 BMI obesity above +2, 61+ months	0.058 [0.090] N=496	0.055 [0.097] N=440	0.127 [0.109] N=374	0.056 [0.125] N=313	-0.035 [0.161] N=245
WHO 2007 BMI thinness below -2, 61+ months	0.023 [0.067] N=496	-0.026 [0.073] N=440	-0.081 [0.081] N=374	-0.038 [0.088] N=313	-0.165 [0.132] N=245

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.594	0.228 [0.164] N=2,676	0.203 [0.176] N=2,470	0.230 [0.190] N=2,212	**0.399** [0.190] N=1,901**	0.350 [0.219] N=1,571
WHO weight-for-age z score	-0.082	-0.021 [0.096] N=2,964	-0.013 [0.106] N=2,736	0.031 [0.106] N=2,452	0.002 [0.118] N=2,104	-0.017 [0.123] N=1,734
WHO weight-for-length/height z score, 0-60 months	0.329	0.085 [0.155] N=1,994	0.042 [0.160] N=1,853	-0.010 [0.171] N=1,668	-0.157 [0.185] N=1,430	-0.068 [0.200] N=1,171
WHO BMI-for-age z score	0.346	0.032 [0.154] N=2,592	0.008 [0.158] N=2,394	-0.006 [0.173] N=2,139	-0.244 [0.178] N=1,839	-0.189 [0.193] N=1,518
Stunted: HAZ below -2	0.257 P=671	-0.042 [0.031] N=2,676	-0.046 [0.033] N=2,470	-0.045 [0.034] N=2,212	**-0.081** [0.035] N=1,901**	-0.050 [0.040] N=1,571
Underweight: WAZ below -2	0.088 P=250	0.030 [0.020] N=2,964	0.021 [0.022] N=2,736	0.003 [0.023] N=2,452	0.005 [0.024] N=2,104	0.011 [0.027] N=1,734
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=227	-0.032 [0.027] N=1,994	-0.022 [0.028] N=1,853	-0.014 [0.030] N=1,668	-0.007 [0.033] N=1,430	-0.029 [0.035] N=1,171
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=382	0.018 [0.037] N=1,994	0.010 [0.039] N=1,853	0.005 [0.041] N=1,668	-0.017 [0.046] N=1,430	-0.028 [0.051] N=1,171
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=179	-0.013 [0.027] N=1,994	-0.015 [0.029] N=1,853	-0.003 [0.031] N=1,668	-0.012 [0.034] N=1,430	-0.015 [0.040] N=1,171
CIAF: any stunting, underweight, or wasting, 0-60 months	0.363 P=694	**-0.060* [0.034] N=1,935**	**-0.063* [0.037] N=1,798**	-0.050 [0.037] N=1,618	**-0.079** [0.040] N=1,388**	-0.056 [0.042] N=1,139
BMI-for-age z score below -2	0.121 P=315	-0.008 [0.028] N=2,592	-0.009 [0.028] N=2,394	-0.007 [0.031] N=2,139	0.015 [0.033] N=1,839	0.013 [0.037] N=1,518
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=228	-0.043 [0.075] N=599	-0.045 [0.078] N=543	-0.049 [0.082] N=477	-0.126 [0.089] N=416	-0.151 [0.105] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=137	-0.035 [0.064] N=599	-0.053 [0.067] N=543	-0.028 [0.074] N=477	-0.096 [0.081] N=416	-0.149 [0.095] N=348
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=60	0.063 [0.069] N=599	0.043 [0.073] N=543	0.023 [0.078] N=477	0.057 [0.080] N=416	0.036 [0.100] N=348

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.062 [0.218] N=2,249	-0.171 [0.241] N=2,043	-0.181 [0.282] N=1,785	0.061 [0.292] N=1,474	-0.225 [0.394] N=1,144
WHO weight-for-age z score	-0.191 [0.134] N=2,497	-0.204 [0.155] N=2,269	-0.144 [0.161] N=1,985	-0.269 [0.196] N=1,637	**-0.406* [0.230] N=1,267**
WHO weight-for-length/height z score, 0-60 months	0.106 [0.230] N=1,680	0.040 [0.245] N=1,539	-0.077 [0.276] N=1,354	-0.387 [0.339] N=1,116	-0.288 [0.424] N=857
WHO BMI-for-age z score	0.179 [0.223] N=2,176	0.180 [0.234] N=1,978	0.203 [0.280] N=1,723	-0.221 [0.312] N=1,423	-0.084 [0.399] N=1,102
Stunted: HAZ below -2	-0.006 [0.041] N=2,249	-0.006 [0.046] N=2,043	-0.003 [0.050] N=1,785	-0.068 [0.055] N=1,474	0.027 [0.073] N=1,144
Underweight: WAZ below -2	**0.046* [0.028] N=2,497**	0.033 [0.031] N=2,269	-0.003 [0.035] N=1,985	0.006 [0.040] N=1,637	0.027 [0.053] N=1,267
Wasted: WLZ/WHZ below -2, 0-60 months	-0.032 [0.038] N=1,680	-0.016 [0.039] N=1,539	-0.002 [0.046] N=1,354	0.014 [0.056] N=1,116	-0.039 [0.064] N=857
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.045] N=1,680	0.015 [0.050] N=1,539	0.002 [0.052] N=1,354	-0.034 [0.066] N=1,116	-0.096 [0.084] N=857
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.031] N=1,680	-0.029 [0.035] N=1,539	-0.006 [0.036] N=1,354	-0.024 [0.044] N=1,116	-0.047 [0.063] N=857
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.040 [0.048] N=1,632	-0.043 [0.055] N=1,495	-0.027 [0.058] N=1,315	-0.083 [0.069] N=1,085	-0.044 [0.083] N=836
BMI-for-age z score below -2	-0.033 [0.036] N=2,176	-0.042 [0.035] N=1,978	-0.052 [0.041] N=1,723	-0.026 [0.045] N=1,423	-0.057 [0.059] N=1,102
WHO 2007 BMI overweight above +1, 61+ months	0.008 [0.119] N=496	0.033 [0.131] N=440	0.047 [0.142] N=374	-0.078 [0.160] N=313	-0.154 [0.217] N=245
WHO 2007 BMI obesity above +2, 61+ months	0.058 [0.090] N=496	0.049 [0.096] N=440	0.115 [0.108] N=374	0.014 [0.120] N=313	-0.094 [0.155] N=245
WHO 2007 BMI thinness below -2, 61+ months	0.024 [0.066] N=496	-0.021 [0.072] N=440	-0.070 [0.080] N=374	-0.014 [0.084] N=313	-0.125 [0.126] N=245

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.594	0.207 [0.166] N=2,676	0.181 [0.178] N=2,470	0.208 [0.194] N=2,212	**0.377* [0.194] N=1,901**	0.310 [0.223] N=1,571
WHO weight-for-age z score	-0.082	-0.051 [0.094] N=2,964	-0.040 [0.104] N=2,736	-0.000 [0.105] N=2,452	-0.028 [0.116] N=2,104	-0.053 [0.121] N=1,734
WHO weight-for-length/height z score, 0-60 months	0.329	0.077 [0.158] N=1,994	0.039 [0.164] N=1,853	-0.008 [0.174] N=1,668	-0.155 [0.187] N=1,430	-0.069 [0.203] N=1,171
WHO BMI-for-age z score	0.346	0.025 [0.155] N=2,592	0.005 [0.159] N=2,394	-0.008 [0.174] N=2,139	-0.243 [0.178] N=1,839	-0.188 [0.195] N=1,518
Stunted: HAZ below -2	0.257 P=671	-0.037 [0.030] N=2,676	-0.040 [0.032] N=2,470	-0.039 [0.033] N=2,212	**-0.074** [0.035] N=1,901**	-0.038 [0.040] N=1,571
Underweight: WAZ below -2	0.088 P=250	**0.037* [0.019] N=2,964**	0.028 [0.021] N=2,736	0.011 [0.022] N=2,452	0.013 [0.023] N=2,104	0.023 [0.026] N=1,734
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=227	-0.028 [0.027] N=1,994	-0.017 [0.028] N=1,853	-0.011 [0.031] N=1,668	-0.005 [0.033] N=1,430	-0.026 [0.035] N=1,171
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=382	0.021 [0.038] N=1,994	0.014 [0.040] N=1,853	0.009 [0.041] N=1,668	-0.014 [0.046] N=1,430	-0.024 [0.051] N=1,171
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=179	-0.012 [0.027] N=1,994	-0.013 [0.029] N=1,853	-0.001 [0.031] N=1,668	-0.011 [0.034] N=1,430	-0.015 [0.040] N=1,171
CIAF: any stunting, underweight, or wasting, 0-60 months	0.363 P=694	-0.046 [0.033] N=1,935	-0.047 [0.035] N=1,798	-0.039 [0.036] N=1,618	**-0.071* [0.038] N=1,388**	-0.041 [0.041] N=1,139

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
BMI-for-age z score below -2	0.121 P=315	-0.006 [0.028] N=2,592	-0.006 [0.028] N=2,394	-0.004 [0.032] N=2,139	0.016 [0.033] N=1,839	0.016 [0.037] N=1,518
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=228	-0.045 [0.075] N=599	-0.044 [0.079] N=543	-0.046 [0.083] N=477	-0.122 [0.090] N=416	-0.146 [0.106] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=137	-0.040 [0.064] N=599	-0.058 [0.068] N=543	-0.033 [0.075] N=477	-0.096 [0.081] N=416	-0.146 [0.094] N=348
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=60	0.062 [0.070] N=599	0.041 [0.073] N=543	0.024 [0.078] N=477	0.060 [0.080] N=416	0.038 [0.101] N=348

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.082 [0.219] N=2,249	-0.199 [0.240] N=2,043	-0.215 [0.280] N=1,785	0.013 [0.292] N=1,474	-0.311 [0.389] N=1,144
WHO weight-for-age z score	-0.216 [0.131] N=2,497	-0.227 [0.153] N=2,269	-0.173 [0.158] N=1,985	-0.296 [0.191] N=1,637	**-0.442* [0.226] N=1,267**
WHO weight-for-length/height z score, 0-60 months	0.100 [0.232] N=1,680	0.041 [0.251] N=1,539	-0.068 [0.277] N=1,354	-0.393 [0.338] N=1,116	-0.314 [0.425] N=857
WHO BMI-for-age z score	0.176 [0.224] N=2,176	0.182 [0.236] N=1,978	0.213 [0.280] N=1,723	-0.205 [0.311] N=1,423	-0.061 [0.400] N=1,102
Stunted: HAZ below -2	-0.002 [0.041] N=2,249	0.001 [0.046] N=2,043	0.004 [0.049] N=1,785	-0.057 [0.055] N=1,474	0.049 [0.071] N=1,144
Underweight: WAZ below -2	**0.051* [0.027] N=2,497**	0.038 [0.031] N=2,269	0.004 [0.034] N=1,985	0.010 [0.040] N=1,637	0.037 [0.052] N=1,267
Wasted: WLZ/WHZ below -2, 0-60 months	-0.028 [0.038] N=1,680	-0.010 [0.040] N=1,539	0.001 [0.046] N=1,354	0.017 [0.057] N=1,116	-0.033 [0.066] N=857
Overweight: WLZ/WHZ above +2, 0-60 months	0.028 [0.046] N=1,680	0.020 [0.051] N=1,539	0.004 [0.053] N=1,354	-0.035 [0.066] N=1,116	-0.091 [0.084] N=857
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.031] N=1,680	-0.026 [0.035] N=1,539	-0.006 [0.036] N=1,354	-0.026 [0.044] N=1,116	-0.052 [0.062] N=857
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.027 [0.046] N=1,632	-0.026 [0.052] N=1,495	-0.018 [0.055] N=1,315	-0.081 [0.065] N=1,085	-0.034 [0.078] N=836
BMI-for-age z score below -2	-0.030 [0.036] N=2,176	-0.039 [0.035] N=1,978	-0.051 [0.042] N=1,723	-0.027 [0.046] N=1,423	-0.054 [0.060] N=1,102
WHO 2007 BMI overweight above +1, 61+ months	0.008 [0.118] N=496	0.042 [0.131] N=440	0.051 [0.143] N=374	-0.082 [0.160] N=313	-0.151 [0.217] N=245
WHO 2007 BMI obesity above +2, 61+ months	0.056 [0.088] N=496	0.054 [0.094] N=440	0.116 [0.106] N=374	0.012 [0.115] N=313	-0.093 [0.152] N=245
WHO 2007 BMI thinness below -2, 61+ months	0.027 [0.065] N=496	-0.023 [0.071] N=440	-0.071 [0.079] N=374	-0.003 [0.085] N=313	-0.121 [0.130] N=245

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.594	0.195 [0.166] N=2,676	0.173 [0.178] N=2,470	0.202 [0.194] N=2,212	**0.361* [0.196] N=1,901**	0.298 [0.225] N=1,571
WHO weight-for-age z score	-0.082	-0.053 [0.094] N=2,964	-0.051 [0.104] N=2,736	-0.011 [0.105] N=2,452	-0.042 [0.117] N=2,104	-0.077 [0.123] N=1,734
WHO weight-for-length/height z score, 0-60 months	0.329	0.090 [0.157] N=1,994	0.044 [0.162] N=1,853	-0.006 [0.173] N=1,668	-0.152 [0.187] N=1,430	-0.072 [0.203] N=1,171
WHO BMI-for-age z score	0.346	0.037 [0.156] N=2,592	0.010 [0.160] N=2,394	-0.003 [0.175] N=2,139	-0.235 [0.181] N=1,839	-0.182 [0.197] N=1,518
Stunted: HAZ below -2	0.257 P=671	-0.035 [0.030] N=2,676	-0.038 [0.032] N=2,470	-0.037 [0.033] N=2,212	**-0.070** [0.034] N=1,901**	-0.037 [0.039] N=1,571

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Underweight: WAZ below -2	0.088 P=250	**0.037* [0.019] N=2,964**	0.030 [0.021] N=2,736	0.014 [0.022] N=2,452	0.017 [0.023] N=2,104	0.026 [0.026] N=1,734
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=227	-0.029 [0.027] N=1,994	-0.018 [0.028] N=1,853	-0.010 [0.030] N=1,668	-0.005 [0.033] N=1,430	-0.027 [0.035] N=1,171
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=382	0.024 [0.038] N=1,994	0.017 [0.040] N=1,853	0.011 [0.041] N=1,668	-0.011 [0.046] N=1,430	-0.025 [0.051] N=1,171
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=179	-0.011 [0.027] N=1,994	-0.015 [0.029] N=1,853	-0.003 [0.030] N=1,668	-0.011 [0.034] N=1,430	-0.017 [0.040] N=1,171
CIAF: any stunting, underweight, or wasting, 0-60 months	0.363 P=694	-0.040 [0.033] N=1,935	-0.041 [0.035] N=1,798	-0.031 [0.036] N=1,618	-0.061 [0.038] N=1,388	-0.033 [0.041] N=1,139
BMI-for-age z score below -2	0.121 P=315	-0.006 [0.028] N=2,592	-0.006 [0.028] N=2,394	-0.004 [0.032] N=2,139	0.018 [0.034] N=1,839	0.016 [0.037] N=1,518
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=228	-0.046 [0.074] N=599	-0.049 [0.079] N=543	-0.052 [0.083] N=477	-0.125 [0.089] N=416	-0.150 [0.106] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=137	-0.041 [0.065] N=599	-0.060 [0.067] N=543	-0.037 [0.074] N=477	-0.100 [0.081] N=416	-0.144 [0.096] N=348
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=60	0.060 [0.069] N=599	0.045 [0.073] N=543	0.024 [0.077] N=477	0.057 [0.080] N=416	0.045 [0.094] N=348

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.088 [0.217] N=2,249	-0.203 [0.238] N=2,043	-0.216 [0.279] N=1,785	0.026 [0.290] N=1,474	-0.301 [0.386] N=1,144
WHO weight-for-age z score	-0.208 [0.132] N=2,497	-0.230 [0.153] N=2,269	-0.181 [0.160] N=1,985	-0.295 [0.194] N=1,637	**-0.424* [0.225] N=1,267**
WHO weight-for-length/height z score, 0-60 months	0.107 [0.232] N=1,680	0.044 [0.249] N=1,539	-0.073 [0.280] N=1,354	-0.392 [0.342] N=1,116	-0.293 [0.426] N=857
WHO BMI-for-age z score	0.187 [0.224] N=2,176	0.187 [0.234] N=1,978	0.216 [0.281] N=1,723	-0.204 [0.314] N=1,423	-0.047 [0.399] N=1,102
Stunted: HAZ below -2	-0.001 [0.041] N=2,249	0.002 [0.046] N=2,043	0.004 [0.050] N=1,785	-0.060 [0.055] N=1,474	0.048 [0.071] N=1,144
Underweight: WAZ below -2	**0.050* [0.027] N=2,497**	0.039 [0.031] N=2,269	0.007 [0.034] N=1,985	0.012 [0.039] N=1,637	0.038 [0.052] N=1,267
Wasted: WLZ/WHZ below -2, 0-60 months	-0.027 [0.038] N=1,680	-0.013 [0.040] N=1,539	0.000 [0.046] N=1,354	0.018 [0.057] N=1,116	-0.033 [0.066] N=857
Overweight: WLZ/WHZ above +2, 0-60 months	0.029 [0.046] N=1,680	0.021 [0.050] N=1,539	0.004 [0.053] N=1,354	-0.035 [0.066] N=1,116	-0.090 [0.084] N=857
Obese: WLZ/WHZ above +3, 0-60 months	-0.019 [0.031] N=1,680	-0.025 [0.035] N=1,539	-0.004 [0.037] N=1,354	-0.025 [0.044] N=1,116	-0.052 [0.060] N=857
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.023 [0.047] N=1,632	-0.027 [0.052] N=1,495	-0.017 [0.056] N=1,315	-0.082 [0.067] N=1,085	-0.030 [0.081] N=836
BMI-for-age z score below -2	-0.030 [0.036] N=2,176	-0.041 [0.035] N=1,978	-0.052 [0.041] N=1,723	-0.025 [0.046] N=1,423	-0.054 [0.059] N=1,102
WHO 2007 BMI overweight above +1, 61+ months	0.011 [0.117] N=496	0.035 [0.130] N=440	0.045 [0.139] N=374	-0.087 [0.155] N=313	-0.152 [0.218] N=245
WHO 2007 BMI obesity above +2, 61+ months	0.052 [0.088] N=496	0.044 [0.094] N=440	0.103 [0.105] N=374	0.000 [0.118] N=313	-0.094 [0.159] N=245
WHO 2007 BMI thinness below -2, 61+ months	0.025 [0.066] N=496	-0.008 [0.073] N=440	-0.065 [0.079] N=374	-0.009 [0.084] N=313	-0.080 [0.121] N=245

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.605	0.193 [0.175] N=2,503	0.159 [0.188] N=2,316	0.177 [0.204] N=2,068	0.333 [0.203] N=1,785	0.275 [0.230] N=1,471
WHO weight-for-age z score	-0.082	0.008 [0.100] N=2,763	0.004 [0.108] N=2,556	0.048 [0.111] N=2,284	0.004 [0.122] N=1,967	-0.006 [0.130] N=1,619
WHO weight-for-length/height z score, 0-60 months	0.338	0.193 [0.160] N=1,859	0.141 [0.163] N=1,735	0.099 [0.172] N=1,559	-0.039 [0.190] N=1,346	0.096 [0.212] N=1,100
WHO BMI-for-age z score	0.353	0.116 [0.158] N=2,423	0.072 [0.161] N=2,243	0.074 [0.177] N=2,000	-0.151 [0.185] N=1,728	-0.068 [0.203] N=1,423
Stunted: HAZ below -2	0.258 P=627	-0.037 [0.032] N=2,503	-0.040 [0.035] N=2,316	-0.036 [0.035] N=2,068	**-0.070* [0.037] N=1,785**	-0.033 [0.042] N=1,471
Underweight: WAZ below -2	0.089 P=236	0.026 [0.021] N=2,763	0.019 [0.022] N=2,556	0.001 [0.024] N=2,284	0.007 [0.026] N=1,967	0.016 [0.030] N=1,619
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=211	-0.043 [0.028] N=1,859	-0.032 [0.028] N=1,735	-0.020 [0.031] N=1,559	-0.014 [0.034] N=1,346	-0.035 [0.036] N=1,100
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=357	0.028 [0.039] N=1,859	0.019 [0.041] N=1,735	0.018 [0.043] N=1,559	-0.006 [0.048] N=1,346	-0.006 [0.056] N=1,100
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=172	-0.013 [0.029] N=1,859	-0.014 [0.031] N=1,735	0.004 [0.033] N=1,559	-0.003 [0.036] N=1,346	-0.001 [0.042] N=1,100
CIAF: any stunting, underweight, or wasting, 0-60 months	0.364 P=647	-0.060 [0.039] N=1,805	-0.062 [0.041] N=1,684	-0.046 [0.042] N=1,512	**-0.078* [0.046] N=1,307**	-0.048 [0.049] N=1,070
BMI-for-age z score below -2	0.120 P=291	-0.017 [0.029] N=2,423	-0.015 [0.029] N=2,243	-0.014 [0.033] N=2,000	0.006 [0.036] N=1,728	0.001 [0.039] N=1,423
WHO 2007 BMI overweight above +1, 61+ months	0.363 P=211	-0.048 [0.079] N=565	-0.040 [0.084] N=512	-0.049 [0.088] N=448	-0.142 [0.093] N=390	-0.166 [0.114] N=325
WHO 2007 BMI obesity above +2, 61+ months	0.219 P=127	-0.051 [0.068] N=565	-0.060 [0.071] N=512	-0.037 [0.079] N=448	-0.124 [0.085] N=390	**-0.201* [0.103] N=325**
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=56	0.061 [0.072] N=565	0.040 [0.078] N=512	0.016 [0.082] N=448	0.052 [0.086] N=390	0.013 [0.102] N=325

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.028 [0.242] N=2,105	-0.144 [0.266] N=1,918	-0.158 [0.310] N=1,670	0.078 [0.331] N=1,387	-0.160 [0.439] N=1,073
WHO weight-for-age z score	-0.171 [0.137] N=2,328	-0.206 [0.154] N=2,121	-0.162 [0.164] N=1,849	**-0.333* [0.195] N=1,532**	**-0.465** [0.233] N=1,184**
WHO weight-for-length/height z score, 0-60 months	0.125 [0.239] N=1,565	0.018 [0.250] N=1,441	-0.088 [0.280] N=1,265	-0.383 [0.351] N=1,052	-0.194 [0.441] N=806
WHO BMI-for-age z score	0.179 [0.231] N=2,035	0.136 [0.241] N=1,855	0.171 [0.289] N=1,612	-0.232 [0.328] N=1,340	-0.059 [0.410] N=1,035
Stunted: HAZ below -2	-0.017 [0.044] N=2,105	-0.017 [0.049] N=1,918	-0.013 [0.054] N=1,670	-0.080 [0.061] N=1,387	0.012 [0.081] N=1,073
Underweight: WAZ below -2	0.041 [0.029] N=2,328	0.031 [0.032] N=2,121	-0.002 [0.036] N=1,849	0.016 [0.043] N=1,532	0.046 [0.057] N=1,184
Wasted: WLZ/WHZ below -2, 0-60 months	-0.048 [0.038] N=1,565	-0.029 [0.039] N=1,441	-0.010 [0.045] N=1,265	-0.003 [0.056] N=1,052	-0.066 [0.061] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.047] N=1,565	0.004 [0.050] N=1,441	-0.002 [0.052] N=1,265	-0.045 [0.068] N=1,052	-0.079 [0.089] N=806
Obese: WLZ/WHZ above +3, 0-60 months	-0.036 [0.033] N=1,565	-0.043 [0.038] N=1,441	-0.013 [0.039] N=1,265	-0.028 [0.047] N=1,052	-0.042 [0.066] N=806

Outcome	8 years	7 years	6 years	5 years	4 years
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.053 [0.051] N=1,522	-0.061 [0.057] N=1,401	-0.040 [0.064] N=1,229	-0.117 [0.075] N=1,024	-0.077 [0.092] N=787
BMI-for-age z score below -2	-0.037 [0.037] N=2,035	-0.042 [0.035] N=1,855	-0.054 [0.041] N=1,612	-0.035 [0.047] N=1,340	-0.082 [0.059] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	-0.006 [0.130] N=469	0.035 [0.142] N=416	0.040 [0.156] N=352	-0.108 [0.170] N=294	-0.200 [0.238] N=229
WHO 2007 BMI obesity above +2, 61+ months	0.023 [0.095] N=469	0.031 [0.100] N=416	0.088 [0.116] N=352	-0.040 [0.122] N=294	-0.215 [0.169] N=229
WHO 2007 BMI thinness below -2, 61+ months	0.040 [0.071] N=469	-0.013 [0.079] N=416	-0.070 [0.086] N=352	-0.021 [0.092] N=294	-0.170 [0.130] N=229

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.605	0.200 [0.173] N=2,503	0.166 [0.185] N=2,316	0.180 [0.203] N=2,068	**0.341* [0.203] N=1,785**	0.296 [0.228] N=1,471
WHO weight-for-age z score	-0.082	0.014 [0.099] N=2,763	0.009 [0.108] N=2,556	0.051 [0.110] N=2,284	0.010 [0.122] N=1,967	0.005 [0.131] N=1,619
WHO weight-for-length/height z score, 0-60 months	0.338	0.193 [0.159] N=1,859	0.143 [0.162] N=1,735	0.091 [0.171] N=1,559	-0.046 [0.190] N=1,346	0.072 [0.212] N=1,100
WHO BMI-for-age z score	0.353	0.117 [0.158] N=2,423	0.073 [0.161] N=2,243	0.074 [0.176] N=2,000	-0.151 [0.185] N=1,728	-0.075 [0.205] N=1,423
Stunted: HAZ below -2	0.258 P=627	-0.039 [0.032] N=2,503	-0.042 [0.034] N=2,316	-0.038 [0.035] N=2,068	**-0.074** [0.036] N=1,785**	-0.041 [0.041] N=1,471
Underweight: WAZ below -2	0.089 P=236	0.024 [0.021] N=2,763	0.017 [0.022] N=2,556	-0.000 [0.024] N=2,284	0.005 [0.025] N=1,967	0.012 [0.029] N=1,619
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=211	-0.043 [0.028] N=1,859	-0.032 [0.028] N=1,735	-0.021 [0.031] N=1,559	-0.015 [0.034] N=1,346	-0.037 [0.036] N=1,100
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=357	0.028 [0.039] N=1,859	0.019 [0.041] N=1,735	0.016 [0.043] N=1,559	-0.010 [0.048] N=1,346	-0.013 [0.055] N=1,100
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=172	-0.013 [0.029] N=1,859	-0.013 [0.031] N=1,735	0.003 [0.033] N=1,559	-0.004 [0.036] N=1,346	-0.004 [0.042] N=1,100
CIAF: any stunting, underweight, or wasting, 0-60 months	0.364 P=647	-0.060 [0.038] N=1,805	-0.062 [0.041] N=1,684	-0.047 [0.041] N=1,512	**-0.082* [0.045] N=1,307**	-0.057 [0.046] N=1,070
BMI-for-age z score below -2	0.120 P=291	-0.017 [0.029] N=2,423	-0.015 [0.029] N=2,243	-0.014 [0.033] N=2,000	0.006 [0.036] N=1,728	0.001 [0.040] N=1,423
WHO 2007 BMI overweight above +1, 61+ months	0.363 P=211	-0.043 [0.078] N=565	-0.038 [0.083] N=512	-0.046 [0.087] N=448	-0.136 [0.091] N=390	-0.169 [0.112] N=325
WHO 2007 BMI obesity above +2, 61+ months	0.219 P=127	-0.053 [0.067] N=565	-0.064 [0.070] N=512	-0.037 [0.079] N=448	-0.125 [0.084] N=390	**-0.198* [0.102] N=325**
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=56	0.058 [0.072] N=565	0.038 [0.077] N=512	0.016 [0.083] N=448	0.052 [0.086] N=390	0.025 [0.108] N=325

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.020 [0.240] N=2,105	-0.140 [0.265] N=1,918	-0.162 [0.311] N=1,670	0.081 [0.334] N=1,387	-0.117 [0.441] N=1,073
WHO weight-for-age z score	-0.159 [0.136] N=2,328	-0.196 [0.153] N=2,121	-0.154 [0.161] N=1,849	-0.320 [0.193] N=1,532	**-0.427* [0.229] N=1,184**

Outcome	8 years	7 years	6 years	5 years	4 years
WHO weight-for-length/height z score, 0-60 months	0.155 [0.238] N=1,565	0.061 [0.251] N=1,441	-0.075 [0.281] N=1,265	-0.372 [0.355] N=1,052	-0.200 [0.450] N=806
WHO BMI-for-age z score	0.189 [0.229] N=2,035	0.146 [0.239] N=1,855	0.176 [0.287] N=1,612	-0.230 [0.325] N=1,340	-0.064 [0.412] N=1,035
Stunted: HAZ below -2	-0.018 [0.044] N=2,105	-0.017 [0.050] N=1,918	-0.014 [0.055] N=1,670	-0.083 [0.061] N=1,387	-0.000 [0.080] N=1,073
Underweight: WAZ below -2	0.038 [0.029] N=2,328	0.029 [0.033] N=2,121	-0.005 [0.036] N=1,849	0.012 [0.042] N=1,532	0.036 [0.056] N=1,184
Wasted: WLZ/WHZ below -2, 0-60 months	-0.048 [0.038] N=1,565	-0.031 [0.039] N=1,441	-0.010 [0.046] N=1,265	-0.005 [0.056] N=1,052	-0.073 [0.062] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.031 [0.046] N=1,565	0.016 [0.050] N=1,441	0.003 [0.053] N=1,265	-0.041 [0.069] N=1,052	-0.079 [0.090] N=806
Obese: WLZ/WHZ above +3, 0-60 months	-0.034 [0.033] N=1,565	-0.040 [0.037] N=1,441	-0.012 [0.039] N=1,265	-0.028 [0.047] N=1,052	-0.043 [0.066] N=806
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.049 [0.051] N=1,522	-0.056 [0.058] N=1,401	-0.039 [0.063] N=1,229	-0.119 [0.075] N=1,024	-0.095 [0.088] N=787
BMI-for-age z score below -2	-0.037 [0.037] N=2,035	-0.043 [0.035] N=1,855	-0.053 [0.041] N=1,612	-0.035 [0.047] N=1,340	-0.082 [0.059] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	0.008 [0.126] N=469	0.046 [0.139] N=416	0.053 [0.151] N=352	-0.089 [0.167] N=294	-0.192 [0.238] N=229
WHO 2007 BMI obesity above +2, 61+ months	0.021 [0.092] N=469	0.027 [0.098] N=416	0.093 [0.113] N=352	-0.044 [0.121] N=294	-0.212 [0.169] N=229
WHO 2007 BMI thinness below -2, 61+ months	0.030 [0.069] N=469	-0.018 [0.078] N=416	-0.073 [0.087] N=352	-0.016 [0.090] N=294	-0.164 [0.136] N=229

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.498	-0.221 [0.178] N=2,665	-0.250 [0.189] N=2,419	-0.303 [0.198] N=2,132	**-0.443** [0.213] N=1,819**	**-0.520** [0.250] N=1,480**
WHO weight-for-age z score	-0.020	-0.096 [0.099] N=2,926	-0.071 [0.104] N=2,668	-0.169 [0.110] N=2,345	-0.148 [0.116] N=1,993	-0.144 [0.131] N=1,614
WHO weight-for-length/height z score, 0-60 months	0.366	0.065 [0.182] N=1,842	0.082 [0.187] N=1,665	0.026 [0.211] N=1,454	0.142 [0.232] N=1,236	0.119 [0.273] N=992
WHO BMI-for-age z score	0.371	0.084 [0.171] N=2,583	0.086 [0.178] N=2,348	0.079 [0.196] N=2,070	0.197 [0.210] N=1,764	0.179 [0.232] N=1,434
Stunted: HAZ below -2	0.245 P=650	0.049 [0.036] N=2,665	0.051 [0.037] N=2,419	0.053 [0.039] N=2,132	**0.087** [0.041] N=1,819**	**0.093* [0.050] N=1,480**
Underweight: WAZ below -2	0.082 P=231	-0.008 [0.017] N=2,926	-0.009 [0.018] N=2,668	0.004 [0.019] N=2,345	0.004 [0.020] N=1,993	0.005 [0.020] N=1,614
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=206	0.041 [0.029] N=1,842	0.037 [0.030] N=1,665	**0.054* [0.033] N=1,454**	0.012 [0.034] N=1,236	0.003 [0.041] N=992
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=364	**0.060* [0.036] N=1,842**	**0.063* [0.037] N=1,665**	0.054 [0.041] N=1,454	0.068 [0.044] N=1,236	0.060 [0.050] N=992
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=180	0.026 [0.030] N=1,842	0.023 [0.031] N=1,665	0.016 [0.034] N=1,454	0.019 [0.038] N=1,236	-0.007 [0.042] N=992
CIAF: any stunting, underweight, or wasting, 0-60 months	0.358 P=638	**0.111** [0.044] N=1,785**	**0.103** [0.045] N=1,609**	**0.115** [0.048] N=1,404**	**0.093* [0.050] N=1,192**	**0.096* [0.055] N=957**
BMI-for-age z score below -2	0.118 P=305	0.023 [0.028] N=2,583	0.024 [0.030] N=2,348	0.021 [0.033] N=2,070	0.000 [0.035] N=1,764	0.002 [0.040] N=1,434
WHO 2007 BMI overweight above +1, 61+ months	0.377 P=281	-0.033 [0.071] N=746	-0.075 [0.072] N=686	-0.079 [0.077] N=617	-0.045 [0.079] N=530	-0.039 [0.087] N=439
WHO 2007 BMI obesity above +2, 61+ months	0.230 P=168	-0.022 [0.062] N=746	-0.064 [0.063] N=686	-0.062 [0.068] N=617	-0.017 [0.070] N=530	-0.016 [0.082] N=439
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=72	0.019 [0.046] N=746	0.033 [0.048] N=686	0.018 [0.050] N=617	0.009 [0.053] N=530	0.009 [0.060] N=439

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.020 [0.225] N=2,284	-0.003 [0.242] N=2,038	-0.042 [0.258] N=1,751	-0.319 [0.305] N=1,438	-0.369 [0.434] N=1,099
WHO weight-for-age z score	-0.150 [0.138] N=2,513	-0.112 [0.150] N=2,255	**-0.324** [0.161] N=1,932**	**-0.357** [0.177] N=1,580**	**-0.391* [0.229] N=1,201**
WHO weight-for-length/height z score, 0-60 months	-0.099 [0.253] N=1,595	-0.111 [0.259] N=1,418	-0.249 [0.307] N=1,207	-0.074 [0.337] N=989	-0.272 [0.427] N=745
WHO BMI-for-age z score	-0.283 [0.205] N=2,217	-0.353 [0.213] N=1,982	**-0.441* [0.252] N=1,704**	-0.303 [0.294] N=1,398	**-0.599* [0.319] N=1,068**
Stunted: HAZ below -2	0.018 [0.049] N=2,284	0.015 [0.053] N=2,038	0.004 [0.056] N=1,751	0.071 [0.062] N=1,438	0.080 [0.097] N=1,099
Underweight: WAZ below -2	-0.017 [0.025] N=2,513	-0.020 [0.028] N=2,255	0.002 [0.031] N=1,932	0.002 [0.035] N=1,580	0.002 [0.039] N=1,201
Wasted: WLZ/WHZ below -2, 0-60 months	**0.098** [0.039] N=1,595**	**0.098** [0.041] N=1,418**	**0.142*** [0.045] N=1,207**	0.077 [0.047] N=989	0.086 [0.058] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.068 [0.050] N=1,595	0.067 [0.053] N=1,418	0.058 [0.060] N=1,207	0.094 [0.069] N=989	0.067 [0.092] N=745

Outcome	8 years	7 years	6 years	5 years	4 years
Obese: WLZ/WHZ above +3, 0-60 months	0.033 [0.039] N=1,595	0.023 [0.040] N=1,418	0.007 [0.045] N=1,207	0.016 [0.057] N=989	-0.078 [0.071] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.151** [0.069] N=1,547**	**0.146* [0.074] N=1,371**	**0.173** [0.084] N=1,166**	0.146 [0.101] N=954	0.173 [0.128] N=719
BMI-for-age z score below -2	**0.074** [0.030] N=2,217**	**0.083** [0.034] N=1,982**	**0.086** [0.039] N=1,704**	0.057 [0.039] N=1,398	**0.087* [0.044] N=1,068**
WHO 2007 BMI overweight above +1, 61+ months	-0.126 [0.104] N=623	**-0.225** [0.109] N=563**	**-0.277** [0.121] N=494**	**-0.261* [0.148] N=407**	**-0.370* [0.199] N=316**
WHO 2007 BMI obesity above +2, 61+ months	-0.102 [0.086] N=623	**-0.198** [0.088] N=563**	**-0.231** [0.101] N=494**	-0.164 [0.112] N=407	-0.250 [0.176] N=316
WHO 2007 BMI thinness below -2, 61+ months	0.067 [0.055] N=623	**0.104* [0.057] N=563**	0.090 [0.063] N=494	0.093 [0.066] N=407	**0.134* [0.073] N=316**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.498	-0.220 [0.178] N=2,665	-0.249 [0.189] N=2,419	-0.303 [0.198] N=2,132	**-0.442** [0.214] N=1,819**	**-0.511** [0.251] N=1,480**
WHO weight-for-age z score	-0.020	-0.096 [0.099] N=2,926	-0.071 [0.104] N=2,668	-0.167 [0.110] N=2,345	-0.147 [0.117] N=1,993	-0.140 [0.132] N=1,614
WHO weight-for-length/height z score, 0-60 months	0.366	0.070 [0.181] N=1,842	0.084 [0.187] N=1,665	0.027 [0.211] N=1,454	0.142 [0.232] N=1,236	0.118 [0.271] N=992
WHO BMI-for-age z score	0.371	0.085 [0.171] N=2,583	0.087 [0.178] N=2,348	0.081 [0.195] N=2,070	0.198 [0.210] N=1,764	0.180 [0.232] N=1,434
Stunted: HAZ below -2	0.245 P=650	0.049 [0.036] N=2,665	0.051 [0.037] N=2,419	0.054 [0.039] N=2,132	**0.087** [0.042] N=1,819**	**0.094* [0.050] N=1,480**
Underweight: WAZ below -2	0.082 P=231	-0.008 [0.017] N=2,926	-0.009 [0.018] N=2,668	0.004 [0.019] N=2,345	0.004 [0.020] N=1,993	0.005 [0.020] N=1,614
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=206	0.039 [0.029] N=1,842	0.034 [0.030] N=1,665	0.052 [0.033] N=1,454	0.011 [0.034] N=1,236	0.004 [0.040] N=992
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=364	0.059 [0.036] N=1,842	0.061 [0.038] N=1,665	0.052 [0.041] N=1,454	0.066 [0.045] N=1,236	0.060 [0.051] N=992
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=180	0.024 [0.030] N=1,842	0.021 [0.031] N=1,665	0.014 [0.034] N=1,454	0.017 [0.038] N=1,236	-0.006 [0.043] N=992
CIAF: any stunting, underweight, or wasting, 0-60 months	0.358 P=638	**0.109** [0.045] N=1,785**	**0.100** [0.046] N=1,609**	**0.111** [0.048] N=1,404**	**0.092* [0.051] N=1,192**	**0.097* [0.056] N=957**
BMI-for-age z score below -2	0.118 P=305	0.024 [0.028] N=2,583	0.024 [0.030] N=2,348	0.022 [0.033] N=2,070	0.001 [0.035] N=1,764	0.002 [0.040] N=1,434
WHO 2007 BMI overweight above +1, 61+ months	0.377 P=281	-0.020 [0.071] N=746	-0.062 [0.073] N=686	-0.069 [0.077] N=617	-0.040 [0.080] N=530	-0.025 [0.088] N=439
WHO 2007 BMI obesity above +2, 61+ months	0.230 P=168	-0.012 [0.063] N=746	-0.053 [0.064] N=686	-0.054 [0.069] N=617	-0.012 [0.071] N=530	-0.009 [0.084] N=439
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=72	0.019 [0.047] N=746	0.033 [0.048] N=686	0.016 [0.050] N=617	0.008 [0.053] N=530	0.004 [0.061] N=439

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.021 [0.224] N=2,284	-0.006 [0.241] N=2,038	-0.046 [0.256] N=1,751	-0.329 [0.304] N=1,438	-0.368 [0.431] N=1,099
WHO weight-for-age z score	-0.150 [0.138] N=2,513	-0.112 [0.150] N=2,255	**-0.325** [0.161] N=1,932**	**-0.359** [0.177] N=1,580**	**-0.386* [0.229] N=1,201**

Outcome	8 years	7 years	6 years	5 years	4 years
WHO weight-for-length/height z score, 0-60 months	-0.087 [0.254] N=1,595	-0.103 [0.260] N=1,418	-0.246 [0.308] N=1,207	-0.072 [0.337] N=989	-0.244 [0.431] N=745
WHO BMI-for-age z score	-0.283 [0.205] N=2,217	-0.354 [0.214] N=1,982	**-0.441* [0.252] N=1,704**	-0.305 [0.294] N=1,398	**-0.601* [0.318] N=1,068**
Stunted: HAZ below -2	0.018 [0.049] N=2,284	0.014 [0.053] N=2,038	0.003 [0.056] N=1,751	0.071 [0.062] N=1,438	0.080 [0.097] N=1,099
Underweight: WAZ below -2	-0.017 [0.025] N=2,513	-0.020 [0.028] N=2,255	0.002 [0.031] N=1,932	0.002 [0.035] N=1,580	0.002 [0.039] N=1,201
Wasted: WLZ/WHZ below -2, 0-60 months	**0.093** [0.039] N=1,595**	**0.090** [0.042] N=1,418**	**0.134*** [0.045] N=1,207**	0.067 [0.047] N=989	0.073 [0.059] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.066 [0.050] N=1,595	0.062 [0.053] N=1,418	0.050 [0.059] N=1,207	0.084 [0.068] N=989	0.059 [0.091] N=745
Obese: WLZ/WHZ above +3, 0-60 months	0.029 [0.039] N=1,595	0.017 [0.040] N=1,418	-0.002 [0.044] N=1,207	0.006 [0.056] N=989	-0.087 [0.070] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.144** [0.069] N=1,547**	**0.135* [0.074] N=1,371**	**0.159* [0.084] N=1,166**	0.132 [0.100] N=954	0.161 [0.127] N=719
BMI-for-age z score below -2	**0.074** [0.030] N=2,217**	**0.083** [0.033] N=1,982**	**0.086** [0.039] N=1,704**	0.055 [0.039] N=1,398	**0.088** [0.043] N=1,068**
WHO 2007 BMI overweight above +1, 61+ months	-0.107 [0.104] N=623	**-0.206* [0.109] N=563**	**-0.263** [0.122] N=494**	**-0.254* [0.149] N=407**	**-0.339* [0.202] N=316**
WHO 2007 BMI obesity above +2, 61+ months	-0.086 [0.088] N=623	**-0.180** [0.090] N=563**	**-0.217** [0.102] N=494**	-0.156 [0.114] N=407	-0.232 [0.179] N=316
WHO 2007 BMI thinness below -2, 61+ months	0.071 [0.057] N=623	**0.110* [0.059] N=563**	0.090 [0.064] N=494	0.096 [0.067] N=407	**0.134* [0.076] N=316**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.498	-0.218 [0.174] N=2,665	-0.241 [0.184] N=2,419	-0.290 [0.195] N=2,132	**-0.426** [0.207] N=1,819**	**-0.497** [0.241] N=1,480**
WHO weight-for-age z score	-0.020	-0.096 [0.098] N=2,926	-0.069 [0.104] N=2,668	-0.161 [0.110] N=2,345	-0.145 [0.116] N=1,993	-0.139 [0.132] N=1,614
WHO weight-for-length/height z score, 0-60 months	0.366	0.079 [0.182] N=1,842	0.090 [0.188] N=1,665	0.040 [0.213] N=1,454	0.142 [0.233] N=1,236	0.118 [0.271] N=992
WHO BMI-for-age z score	0.371	0.080 [0.171] N=2,583	0.082 [0.179] N=2,348	0.079 [0.195] N=2,070	0.192 [0.209] N=1,764	0.175 [0.232] N=1,434
Stunted: HAZ below -2	0.245 P=650	0.048 [0.035] N=2,665	0.049 [0.037] N=2,419	0.049 [0.039] N=2,132	**0.081** [0.040] N=1,819**	**0.089* [0.048] N=1,480**
Underweight: WAZ below -2	0.082 P=231	-0.008 [0.017] N=2,926	-0.009 [0.018] N=2,668	0.002 [0.019] N=2,345	0.003 [0.020] N=1,993	0.005 [0.019] N=1,614
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=206	0.038 [0.030] N=1,842	0.032 [0.031] N=1,665	0.049 [0.033] N=1,454	0.010 [0.035] N=1,236	0.003 [0.040] N=992
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=364	**0.062* [0.036] N=1,842**	0.062 [0.038] N=1,665	0.052 [0.042] N=1,454	0.064 [0.045] N=1,236	0.060 [0.051] N=992
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=180	0.026 [0.031] N=1,842	0.022 [0.031] N=1,665	0.014 [0.034] N=1,454	0.016 [0.039] N=1,236	-0.006 [0.044] N=992
CIAF: any stunting, underweight, or wasting, 0-60 months	0.358 P=638	**0.111** [0.044] N=1,785**	**0.098** [0.046] N=1,609**	**0.106** [0.048] N=1,404**	**0.085* [0.050] N=1,192**	**0.093* [0.054] N=957**
BMI-for-age z score below -2	0.118 P=305	0.024 [0.028] N=2,583	0.024 [0.030] N=2,348	0.021 [0.033] N=2,070	-0.000 [0.035] N=1,764	0.002 [0.040] N=1,434
WHO 2007 BMI overweight above +1, 61+ months	0.377 P=281	-0.025 [0.072] N=746	-0.065 [0.074] N=686	-0.072 [0.078] N=617	-0.044 [0.082] N=530	-0.027 [0.089] N=439

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO 2007 BMI obesity above +2, 61+ months	0.230 P=168	-0.019 [0.063] N=746	-0.057 [0.065] N=686	-0.056 [0.069] N=617	-0.018 [0.071] N=530	-0.017 [0.083] N=439
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=72	0.019 [0.046] N=746	0.031 [0.047] N=686	0.017 [0.049] N=617	0.007 [0.053] N=530	-0.003 [0.058] N=439

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.013 [0.223] N=2,284	0.009 [0.239] N=2,038	-0.019 [0.255] N=1,751	-0.294 [0.297] N=1,438	-0.327 [0.420] N=1,099
WHO weight-for-age z score	-0.152 [0.139] N=2,513	-0.113 [0.151] N=2,255	**-0.314* [0.163] N=1,932**	**-0.360** [0.175] N=1,580**	**-0.390* [0.226] N=1,201**
WHO weight-for-length/height z score, 0-60 months	-0.096 [0.254] N=1,595	-0.111 [0.262] N=1,418	-0.246 [0.309] N=1,207	-0.089 [0.336] N=989	-0.253 [0.426] N=745
WHO BMI-for-age z score	-0.301 [0.204] N=2,217	**-0.370* [0.214] N=1,982**	**-0.455* [0.250] N=1,704**	-0.331 [0.292] N=1,398	**-0.626* [0.317] N=1,068**
Stunted: HAZ below -2	0.016 [0.049] N=2,284	0.011 [0.052] N=2,038	-0.002 [0.056] N=1,751	0.063 [0.061] N=1,438	0.070 [0.095] N=1,099
Underweight: WAZ below -2	-0.018 [0.025] N=2,513	-0.020 [0.027] N=2,255	-0.002 [0.031] N=1,932	0.000 [0.035] N=1,580	0.001 [0.038] N=1,201
Wasted: WLZ/WHZ below -2, 0-60 months	**0.093** [0.040] N=1,595**	**0.089** [0.042] N=1,418**	**0.130*** [0.046] N=1,207**	0.064 [0.048] N=989	0.073 [0.060] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.067 [0.051] N=1,595	0.063 [0.054] N=1,418	0.047 [0.061] N=1,207	0.078 [0.070] N=989	0.058 [0.093] N=745
Obese: WLZ/WHZ above +3, 0-60 months	0.029 [0.039] N=1,595	0.015 [0.040] N=1,418	-0.003 [0.045] N=1,207	0.003 [0.057] N=989	-0.089 [0.071] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.145** [0.069] N=1,547**	**0.132* [0.075] N=1,371**	**0.148* [0.085] N=1,166**	0.122 [0.100] N=954	0.158 [0.125] N=719
BMI-for-age z score below -2	**0.075** [0.030] N=2,217**	**0.083** [0.033] N=1,982**	**0.084** [0.039] N=1,704**	0.054 [0.040] N=1,398	**0.083* [0.044] N=1,068**
WHO 2007 BMI overweight above +1, 61+ months	-0.116 [0.105] N=623	**-0.212* [0.110] N=563**	**-0.265** [0.122] N=494**	**-0.260* [0.153] N=407**	**-0.360* [0.203] N=316**
WHO 2007 BMI obesity above +2, 61+ months	-0.093 [0.088] N=623	**-0.185** [0.090] N=563**	**-0.217** [0.101] N=494**	-0.158 [0.114] N=407	-0.258 [0.174] N=316
WHO 2007 BMI thinness below -2, 61+ months	0.074 [0.056] N=623	**0.108* [0.059] N=563**	0.094 [0.064] N=494	0.099 [0.067] N=407	0.117 [0.079] N=316

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.498	-0.212 [0.173] N=2,665	-0.233 [0.183] N=2,419	-0.283 [0.195] N=2,132	**-0.414** [0.207] N=1,819**	**-0.495** [0.241] N=1,480**
WHO weight-for-age z score	-0.020	-0.094 [0.096] N=2,926	-0.066 [0.102] N=2,668	-0.156 [0.108] N=2,345	-0.132 [0.113] N=1,993	-0.137 [0.128] N=1,614
WHO weight-for-length/height z score, 0-60 months	0.366	0.083 [0.181] N=1,842	0.096 [0.186] N=1,665	0.042 [0.211] N=1,454	0.157 [0.230] N=1,236	0.140 [0.270] N=992
WHO BMI-for-age z score	0.371	0.079 [0.172] N=2,583	0.082 [0.179] N=2,348	0.078 [0.196] N=2,070	0.198 [0.210] N=1,764	0.176 [0.233] N=1,434
Stunted: HAZ below -2	0.245 P=650	0.048 [0.035] N=2,665	0.048 [0.037] N=2,419	0.050 [0.039] N=2,132	**0.082** [0.040] N=1,819**	**0.088* [0.048] N=1,480**
Underweight: WAZ below -2	0.082 P=231	-0.008 [0.016] N=2,926	-0.010 [0.017] N=2,668	0.002 [0.018] N=2,345	0.002 [0.020] N=1,993	0.004 [0.019] N=1,614
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=206	0.037 [0.029] N=1,842	0.033 [0.030] N=1,665	0.051 [0.033] N=1,454	0.009 [0.034] N=1,236	0.000 [0.040] N=992

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=364	**0.060* [0.036] N=1,842**	0.062 [0.037] N=1,665	0.054 [0.041] N=1,454	0.067 [0.044] N=1,236	0.062 [0.050] N=992
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=180	0.028 [0.030] N=1,842	0.025 [0.030] N=1,665	0.017 [0.033] N=1,454	0.021 [0.038] N=1,236	-0.002 [0.043] N=992
CIAF: any stunting, underweight, or wasting, 0-60 months	0.358 P=638	**0.108** [0.044] N=1,785**	**0.096** [0.046] N=1,609**	**0.106** [0.048] N=1,404**	**0.085* [0.050] N=1,192**	**0.092* [0.054] N=957**
BMI-for-age z score below -2	0.118 P=305	0.023 [0.028] N=2,583	0.023 [0.030] N=2,348	0.020 [0.033] N=2,070	-0.002 [0.035] N=1,764	-0.002 [0.039] N=1,434
WHO 2007 BMI overweight above +1, 61+ months	0.377 P=281	-0.028 [0.073] N=746	-0.068 [0.075] N=686	-0.073 [0.080] N=617	-0.043 [0.083] N=530	-0.032 [0.092] N=439
WHO 2007 BMI obesity above +2, 61+ months	0.230 P=168	-0.024 [0.063] N=746	-0.067 [0.065] N=686	-0.065 [0.070] N=617	-0.030 [0.072] N=530	-0.034 [0.083] N=439
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=72	0.017 [0.046] N=746	0.032 [0.047] N=686	0.017 [0.049] N=617	0.006 [0.052] N=530	-0.008 [0.058] N=439

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.015 [0.222] N=2,284	0.012 [0.239] N=2,038	-0.006 [0.256] N=1,751	-0.271 [0.297] N=1,438	-0.334 [0.412] N=1,099
WHO weight-for-age z score	-0.136 [0.134] N=2,513	-0.093 [0.146] N=2,255	**-0.289* [0.159] N=1,932**	**-0.307* [0.170] N=1,580**	-0.358 [0.216] N=1,201
WHO weight-for-length/height z score, 0-60 months	-0.084 [0.253] N=1,595	-0.107 [0.258] N=1,418	-0.252 [0.301] N=1,207	-0.085 [0.329] N=989	-0.230 [0.425] N=745
WHO BMI-for-age z score	-0.290 [0.204] N=2,217	**-0.360* [0.213] N=1,982**	**-0.453* [0.251] N=1,704**	-0.314 [0.292] N=1,398	**-0.613* [0.319] N=1,068**
Stunted: HAZ below -2	0.018 [0.049] N=2,284	0.011 [0.052] N=2,038	-0.003 [0.056] N=1,751	0.061 [0.061] N=1,438	0.073 [0.093] N=1,099
Underweight: WAZ below -2	-0.019 [0.024] N=2,513	-0.023 [0.027] N=2,255	-0.004 [0.030] N=1,932	-0.006 [0.034] N=1,580	-0.004 [0.038] N=1,201
Wasted: WLZ/WHZ below -2, 0-60 months	**0.092** [0.040] N=1,595**	**0.089** [0.042] N=1,418**	**0.135*** [0.045] N=1,207**	0.068 [0.047] N=989	0.075 [0.058] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.065 [0.050] N=1,595	0.060 [0.052] N=1,418	0.046 [0.059] N=1,207	0.075 [0.069] N=989	0.060 [0.093] N=745
Obese: WLZ/WHZ above +3, 0-60 months	0.031 [0.039] N=1,595	0.018 [0.040] N=1,418	-0.001 [0.045] N=1,207	0.006 [0.057] N=989	-0.085 [0.070] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.142** [0.068] N=1,547**	**0.128* [0.074] N=1,371**	**0.146* [0.084] N=1,166**	0.116 [0.098] N=954	0.162 [0.121] N=719
BMI-for-age z score below -2	**0.076** [0.030] N=2,217**	**0.082** [0.034] N=1,982**	**0.085** [0.039] N=1,704**	0.055 [0.040] N=1,398	**0.089* [0.044] N=1,068**
WHO 2007 BMI overweight above +1, 61+ months	-0.123 [0.104] N=623	**-0.215* [0.110] N=563**	**-0.275** [0.122] N=494**	**-0.264* [0.150] N=407**	**-0.383* [0.201] N=316**
WHO 2007 BMI obesity above +2, 61+ months	-0.089 [0.086] N=623	**-0.186** [0.089] N=563**	**-0.220** [0.099] N=494**	-0.159 [0.109] N=407	-0.260 [0.172] N=316
WHO 2007 BMI thinness below -2, 61+ months	0.070 [0.056] N=623	**0.108* [0.059] N=563**	0.095 [0.063] N=494	0.100 [0.069] N=407	0.107 [0.080] N=316

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.511	-0.211 [0.177] N=2,499	-0.208 [0.189] N=2,270	-0.249 [0.199] N=2,003	**-0.384* [0.216] N=1,703**	**-0.490* [0.249] N=1,388**

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO weight-for-age z score	-0.015	-0.079 [0.102] N=2,736	-0.042 [0.108] N=2,495	-0.140 [0.114] N=2,196	-0.122 [0.120] N=1,863	-0.149 [0.134] N=1,513
WHO weight-for-length/height z score, 0-60 months	0.368	0.116 [0.192] N=1,719	0.144 [0.199] N=1,554	0.055 [0.222] N=1,358	0.154 [0.243] N=1,153	0.094 [0.279] N=928
WHO BMI-for-age z score	0.386	0.069 [0.176] N=2,421	0.071 [0.184] N=2,202	0.039 [0.202] N=1,943	0.144 [0.219] N=1,652	0.097 [0.242] N=1,346
Stunted: HAZ below -2	0.245 P=609	0.048 [0.035] N=2,499	0.048 [0.037] N=2,270	0.046 [0.038] N=2,003	**0.078** [0.039] N=1,703**	**0.090** [0.045] N=1,388**
Underweight: WAZ below -2	0.083 P=219	-0.014 [0.018] N=2,736	-0.016 [0.019] N=2,495	-0.006 [0.019] N=2,196	-0.001 [0.020] N=1,863	0.008 [0.021] N=1,513
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=193	0.042 [0.030] N=1,719	0.035 [0.031] N=1,554	0.054 [0.033] N=1,358	0.017 [0.035] N=1,153	0.016 [0.041] N=928
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=341	0.059 [0.037] N=1,719	0.062 [0.038] N=1,554	0.048 [0.041] N=1,358	0.059 [0.044] N=1,153	0.048 [0.050] N=928
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=172	0.024 [0.031] N=1,719	0.021 [0.031] N=1,554	0.012 [0.033] N=1,358	0.019 [0.038] N=1,153	-0.001 [0.042] N=928
CIAF: any stunting, underweight, or wasting, 0-60 months	0.359 P=597	**0.116** [0.047] N=1,667**	**0.104** [0.049] N=1,503**	**0.109** [0.050] N=1,313**	0.087 [0.053] N=1,112	**0.103* [0.057] N=895**
BMI-for-age z score below -2	0.116 P=278	0.031 [0.028] N=2,421	0.032 [0.030] N=2,202	0.030 [0.032] N=1,943	0.012 [0.034] N=1,652	0.017 [0.037] N=1,346
WHO 2007 BMI overweight above +1, 61+ months	0.380 P=269	-0.037 [0.072] N=708	-0.080 [0.073] N=652	-0.075 [0.078] N=586	-0.045 [0.082] N=501	-0.022 [0.089] N=415
WHO 2007 BMI obesity above +2, 61+ months	0.237 P=164	-0.031 [0.065] N=708	-0.076 [0.067] N=652	-0.080 [0.072] N=586	-0.048 [0.075] N=501	-0.029 [0.088] N=415
WHO 2007 BMI thinness below -2, 61+ months	0.094 P=64	0.025 [0.046] N=708	0.042 [0.047] N=652	0.025 [0.050] N=586	0.015 [0.053] N=501	0.021 [0.059] N=415

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.008 [0.228] N=2,141	0.055 [0.247] N=1,912	0.027 [0.266] N=1,645	-0.255 [0.316] N=1,345	-0.330 [0.428] N=1,030
WHO weight-for-age z score	-0.140 [0.141] N=2,350	-0.082 [0.154] N=2,109	**-0.308* [0.164] N=1,810**	**-0.338* [0.181] N=1,477**	**-0.396* [0.230] N=1,127**
WHO weight-for-length/height z score, 0-60 months	-0.034 [0.267] N=1,487	-0.027 [0.280] N=1,322	-0.241 [0.331] N=1,126	-0.093 [0.354] N=921	-0.244 [0.466] N=696
WHO BMI-for-age z score	-0.280 [0.206] N=2,076	-0.338 [0.216] N=1,857	**-0.454* [0.256] N=1,598**	-0.325 [0.290] N=1,307	**-0.642** [0.305] N=1,001**
Stunted: HAZ below -2	0.013 [0.050] N=2,141	0.007 [0.054] N=1,912	-0.014 [0.057] N=1,645	0.045 [0.060] N=1,345	0.051 [0.086] N=1,030
Underweight: WAZ below -2	-0.028 [0.026] N=2,350	-0.033 [0.029] N=2,109	-0.016 [0.032] N=1,810	-0.010 [0.037] N=1,477	-0.004 [0.041] N=1,127
Wasted: WLZ/WHZ below -2, 0-60 months	**0.095** [0.040] N=1,487**	**0.086** [0.043] N=1,322**	**0.132*** [0.047] N=1,126**	0.070 [0.051] N=921	0.074 [0.061] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.066 [0.052] N=1,487	0.063 [0.055] N=1,322	0.041 [0.061] N=1,126	0.066 [0.071] N=921	0.046 [0.098] N=696
Obese: WLZ/WHZ above +3, 0-60 months	0.023 [0.041] N=1,487	0.010 [0.042] N=1,322	-0.014 [0.047] N=1,126	-0.001 [0.059] N=921	-0.084 [0.077] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.150** [0.074] N=1,443**	**0.135* [0.081] N=1,279**	0.143 [0.090] N=1,089	0.093 [0.110] N=888	0.130 [0.131] N=671
BMI-for-age z score below -2	**0.080*** [0.031] N=2,076**	**0.088** [0.035] N=1,857**	**0.091** [0.039] N=1,598**	**0.067* [0.040] N=1,307**	**0.100** [0.044] N=1,001**

Outcome	8 years	7 years	6 years	5 years	4 years
WHO 2007 BMI overweight above +1, 61+ months	-0.133 [0.104] N=592	**-0.233** [0.108] N=536**	**-0.263** [0.120] N=470**	-0.244 [0.147] N=385	-0.310 [0.199] N=299
WHO 2007 BMI obesity above +2, 61+ months	-0.104 [0.090] N=592	**-0.200** [0.094] N=536**	**-0.237** [0.104] N=470**	-0.177 [0.118] N=385	-0.239 [0.181] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.071 [0.057] N=592	**0.114* [0.059] N=536**	0.093 [0.063] N=470	0.088 [0.063] N=385	**0.129* [0.075] N=299**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.511	-0.216 [0.177] N=2,499	-0.212 [0.190] N=2,270	-0.252 [0.199] N=2,003	**-0.381* [0.217] N=1,703**	**-0.479* [0.250] N=1,388**
WHO weight-for-age z score	-0.015	-0.080 [0.101] N=2,736	-0.044 [0.108] N=2,495	-0.145 [0.115] N=2,196	-0.122 [0.121] N=1,863	-0.143 [0.136] N=1,513
WHO weight-for-length/height z score, 0-60 months	0.368	0.124 [0.192] N=1,719	0.148 [0.200] N=1,554	0.060 [0.222] N=1,358	0.169 [0.240] N=1,153	0.112 [0.280] N=928
WHO BMI-for-age z score	0.386	0.070 [0.176] N=2,421	0.074 [0.184] N=2,202	0.041 [0.201] N=1,943	0.146 [0.219] N=1,652	0.097 [0.241] N=1,346
Stunted: HAZ below -2	0.245 P=609	0.047 [0.035] N=2,499	0.047 [0.037] N=2,270	0.045 [0.038] N=2,003	**0.076* [0.039] N=1,703**	**0.088* [0.045] N=1,388**
Underweight: WAZ below -2	0.083 P=219	-0.013 [0.017] N=2,736	-0.015 [0.019] N=2,495	-0.005 [0.019] N=2,196	-0.000 [0.021] N=1,863	0.007 [0.021] N=1,513
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=193	0.040 [0.030] N=1,719	0.033 [0.031] N=1,554	0.053 [0.033] N=1,358	0.012 [0.035] N=1,153	0.010 [0.040] N=928
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=341	0.059 [0.036] N=1,719	0.061 [0.038] N=1,554	0.047 [0.041] N=1,358	0.058 [0.044] N=1,153	0.048 [0.050] N=928
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=172	0.024 [0.030] N=1,719	0.021 [0.030] N=1,554	0.011 [0.033] N=1,358	0.018 [0.037] N=1,153	-0.002 [0.042] N=928
CIAF: any stunting, underweight, or wasting, 0-60 months	0.359 P=597	**0.112** [0.046] N=1,667**	**0.100** [0.048] N=1,503**	**0.104** [0.050] N=1,313**	0.080 [0.052] N=1,112	**0.093* [0.055] N=895**
BMI-for-age z score below -2	0.116 P=278	0.030 [0.028] N=2,421	0.031 [0.030] N=2,202	0.029 [0.032] N=1,943	0.011 [0.033] N=1,652	0.015 [0.037] N=1,346
WHO 2007 BMI overweight above +1, 61+ months	0.380 P=269	-0.041 [0.072] N=708	-0.085 [0.074] N=652	-0.084 [0.078] N=586	-0.053 [0.082] N=501	-0.029 [0.089] N=415
WHO 2007 BMI obesity above +2, 61+ months	0.237 P=164	-0.031 [0.065] N=708	-0.077 [0.067] N=652	-0.081 [0.072] N=586	-0.049 [0.075] N=501	-0.029 [0.088] N=415
WHO 2007 BMI thinness below -2, 61+ months	0.094 P=64	0.021 [0.046] N=708	0.039 [0.047] N=652	0.017 [0.049] N=586	0.007 [0.051] N=501	0.009 [0.057] N=415

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.031 [0.225] N=2,141	0.033 [0.244] N=1,912	0.005 [0.261] N=1,645	-0.293 [0.316] N=1,345	-0.367 [0.426] N=1,030
WHO weight-for-age z score	-0.143 [0.140] N=2,350	-0.085 [0.154] N=2,109	**-0.313* [0.164] N=1,810**	**-0.340* [0.180] N=1,477**	**-0.410* [0.230] N=1,127**
WHO weight-for-length/height z score, 0-60 months	-0.035 [0.266] N=1,487	-0.030 [0.276] N=1,322	-0.240 [0.327] N=1,126	-0.063 [0.354] N=921	-0.223 [0.456] N=696
WHO BMI-for-age z score	-0.275 [0.206] N=2,076	-0.332 [0.215] N=1,857	**-0.452* [0.254] N=1,598**	-0.306 [0.289] N=1,307	**-0.626** [0.308] N=1,001**
Stunted: HAZ below -2	0.016 [0.049] N=2,141	0.010 [0.054] N=1,912	-0.011 [0.057] N=1,645	0.052 [0.060] N=1,345	0.064 [0.086] N=1,030

Outcome	8 years	7 years	6 years	5 years	4 years
Underweight: WAZ below -2	-0.027 [0.026] N=2,350	-0.032 [0.029] N=2,109	-0.014 [0.031] N=1,810	-0.008 [0.036] N=1,477	0.002 [0.041] N=1,127
Wasted: WLZ/WHZ below -2, 0-60 months	**0.095** [0.040] N=1,487**	**0.087** [0.043] N=1,322**	**0.134*** [0.046] N=1,126**	0.066 [0.050] N=921	0.073 [0.059] N=696
Overweight: WLZ/WHZ above +2, 0-60 months	0.066 [0.051] N=1,487	0.063 [0.053] N=1,322	0.040 [0.060] N=1,126	0.070 [0.070] N=921	0.048 [0.096] N=696
Obese: WLZ/WHZ above +3, 0-60 months	0.023 [0.041] N=1,487	0.010 [0.042] N=1,322	-0.013 [0.046] N=1,126	-0.001 [0.059] N=921	-0.084 [0.075] N=696
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.149** [0.073] N=1,443**	**0.134* [0.080] N=1,279**	0.144 [0.090] N=1,089	0.098 [0.109] N=888	0.136 [0.130] N=671
BMI-for-age z score below -2	**0.079** [0.030] N=2,076**	**0.088** [0.034] N=1,857**	**0.090** [0.039] N=1,598**	0.064 [0.039] N=1,307	**0.099** [0.043] N=1,001**
WHO 2007 BMI overweight above +1, 61+ months	-0.129 [0.103] N=592	**-0.228** [0.107] N=536**	**-0.264** [0.120] N=470**	-0.243 [0.147] N=385	-0.307 [0.199] N=299
WHO 2007 BMI obesity above +2, 61+ months	-0.098 [0.090] N=592	**-0.196** [0.093] N=536**	**-0.232** [0.104] N=470**	-0.176 [0.118] N=385	-0.237 [0.183] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.069 [0.057] N=592	**0.113* [0.059] N=536**	0.082 [0.063] N=470	0.078 [0.063] N=385	0.106 [0.076] N=299

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 83 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.