Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-5 Using 10 October 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 71 months on 10 October 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	0.160 [0.155] N=2,523	0.165 [0.160] N=2,342	0.080 [0.167] N=2,111	0.153 [0.171] N=1,812	0.077 [0.186] N=1,493
WHO weight-for-age z score	0.329	0.050 [0.098] N=2,726	0.112 [0.105] N=2,526	0.139 [0.107] N=2,276	0.101 [0.110] N=1,953	0.083 [0.117] N=1,608
WHO weight-for-length/height z score, 0-60 months	0.342	0.028 [0.169] N=1,880	0.038 [0.174] N=1,741	0.005 [0.187] N=1,566	-0.161 [0.211] N=1,342	-0.116 [0.225] N=1,097
WHO BMI-for-age z score	0.355	-0.028 [0.172] N=2,419	-0.003 [0.178] N=2,252	0.002 [0.193] N=2,021	-0.172 [0.214] N=1,731	-0.109 [0.234] N=1,426
Stunted: HAZ below -2	0.194 P=475	-0.025 [0.032] N=2,523	-0.020 [0.033] N=2,342	-0.005 [0.035] N=2,111	-0.014 [0.036] N=1,812	-0.004 [0.041] N=1,493
Underweight: WAZ below -2	0.048 P=128	-0.017 [0.016] N=2,726	-0.023 [0.016] N=2,526	**-0.030* [0.017] N=2,276**	-0.028 [0.018] N=1,953	-0.029 [0.021] N=1,608
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=210	-0.013 [0.026] N=1,880	-0.008 [0.027] N=1,741	0.000 [0.030] N=1,566	0.011 [0.034] N=1,342	0.004 [0.035] N=1,097
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=361	0.030 [0.038] N=1,880	0.030 [0.041] N=1,741	0.029 [0.043] N=1,566	-0.008 [0.048] N=1,342	-0.005 [0.052] N=1,097
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=160	0.008 [0.028] N=1,880	0.009 [0.030] N=1,741	0.019 [0.033] N=1,566	-0.003 [0.036] N=1,342	-0.008 [0.041] N=1,097
CIAF: any stunting, underweight, or wasting, 0-60 months	0.254 P=458	-0.030 [0.039] N=1,838	-0.030 [0.040] N=1,703	-0.005 [0.043] N=1,533	-0.013 [0.043] N=1,311	0.002 [0.046] N=1,073
BMI-for-age z score below -2	0.118 P=290	0.012 [0.028] N=2,419	0.014 [0.029] N=2,252	0.019 [0.031] N=2,021	0.043 [0.034] N=1,731	0.048 [0.034] N=1,426
WHO 2007 BMI overweight above +1, 61+ months	0.383 P=201	0.029 [0.078] N=527	0.026 [0.079] N=497	0.064 [0.086] N=448	0.059 [0.094] N=388	0.079 [0.108] N=324
WHO 2007 BMI obesity above +2, 61+ months	0.208 P=110	-0.018 [0.060] N=527	-0.029 [0.061] N=497	-0.040 [0.064] N=448	-0.043 [0.069] N=388	-0.080 [0.084] N=324
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=66	0.013 [0.059] N=527	0.007 [0.060] N=497	-0.040 [0.066] N=448	-0.046 [0.073] N=388	-0.059 [0.085] N=324

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.100 [0.239] N=2,121	0.099 [0.257] N=1,940	-0.078 [0.280] N=1,709	0.094 [0.314] N=1,410	-0.190 [0.426] N=1,091
WHO weight-for-age z score	-0.044 [0.138] N=2,298	0.061 [0.153] N=2,098	0.093 [0.162] N=1,848	0.042 [0.188] N=1,525	-0.090 [0.238] N=1,180
WHO weight-for-length/height z score, 0-60 months	0.029 [0.225] N=1,594	0.063 [0.235] N=1,455	-0.030 [0.269] N=1,280	-0.369 [0.326] N=1,056	-0.362 [0.414] N=811
WHO BMI-for-age z score	0.107 [0.214] N=2,038	0.186 [0.224] N=1,871	0.222 [0.251] N=1,640	-0.061 [0.294] N=1,350	0.169 [0.380] N=1,045
Stunted: HAZ below -2	0.020 [0.045] N=2,121	0.037 [0.048] N=1,940	0.077 [0.054] N=1,709	0.075 [0.058] N=1,410	**0.143* [0.083] N=1,091**
Underweight: WAZ below -2	-0.006 [0.023] N=2,298	-0.014 [0.024] N=2,098	-0.023 [0.027] N=1,848	-0.020 [0.032] N=1,525	-0.007 [0.043] N=1,180
Wasted: WLZ/WHZ below -2, 0-60 months	0.001 [0.034] N=1,594	0.010 [0.035] N=1,455	0.032 [0.040] N=1,280	0.053 [0.052] N=1,056	0.053 [0.056] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.044 [0.049] N=1,594	0.049 [0.053] N=1,455	0.049 [0.059] N=1,280	-0.022 [0.072] N=1,056	-0.012 [0.094] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.032] N=1,594	0.037 [0.035] N=1,455	0.063 [0.040] N=1,280	0.024 [0.048] N=1,056	0.035 [0.067] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.026 [0.052] N=1,557	0.032 [0.056] N=1,422	0.094 [0.063] N=1,252	0.087 [0.070] N=1,030	**0.183** [0.081] N=792**
BMI-for-age z score below -2	-0.018 [0.033] N=2,038	-0.021 [0.035] N=1,871	-0.015 [0.039] N=1,640	0.008 [0.045] N=1,350	0.003 [0.051] N=1,045
WHO 2007 BMI overweight above +1, 61+ months	0.033 [0.117] N=431	0.022 [0.121] N=401	0.094 [0.132] N=352	0.086 [0.148] N=292	0.177 [0.203] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.057 [0.095] N=431	0.040 [0.098] N=401	0.041 [0.101] N=352	0.061 [0.117] N=292	0.037 [0.166] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.002 [0.067] N=431	-0.016 [0.069] N=401	-0.117 [0.075] N=352	**-0.171** [0.081] N=292**	**-0.256** [0.101] N=228**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	0.166 [0.155] N=2,523	0.171 [0.160] N=2,342	0.090 [0.168] N=2,111	0.166 [0.171] N=1,812	0.086 [0.187] N=1,493
WHO weight-for-age z score	0.329	0.052 [0.099] N=2,726	0.113 [0.106] N=2,526	0.143 [0.108] N=2,276	0.107 [0.112] N=1,953	0.085 [0.118] N=1,608
WHO weight-for-length/height z score, 0-60 months	0.342	0.026 [0.170] N=1,880	0.036 [0.175] N=1,741	0.001 [0.188] N=1,566	-0.162 [0.213] N=1,342	-0.120 [0.227] N=1,097
WHO BMI-for-age z score	0.355	-0.029 [0.172] N=2,419	-0.006 [0.178] N=2,252	-0.001 [0.194] N=2,021	-0.173 [0.215] N=1,731	-0.112 [0.235] N=1,426
Stunted: HAZ below -2	0.194 P=475	-0.025 [0.032] N=2,523	-0.020 [0.033] N=2,342	-0.005 [0.035] N=2,111	-0.015 [0.036] N=1,812	-0.005 [0.041] N=1,493
Underweight: WAZ below -2	0.048 P=128	-0.018 [0.016] N=2,726	-0.023 [0.016] N=2,526	**-0.030* [0.017] N=2,276**	-0.028 [0.018] N=1,953	-0.029 [0.021] N=1,608
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=210	-0.012 [0.026] N=1,880	-0.007 [0.027] N=1,741	0.002 [0.030] N=1,566	0.012 [0.034] N=1,342	0.006 [0.036] N=1,097
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=361	0.030 [0.038] N=1,880	0.030 [0.040] N=1,741	0.029 [0.043] N=1,566	-0.006 [0.047] N=1,342	-0.003 [0.052] N=1,097
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=160	0.009 [0.028] N=1,880	0.010 [0.030] N=1,741	0.020 [0.033] N=1,566	-0.002 [0.036] N=1,342	-0.007 [0.041] N=1,097
CIAF: any stunting, underweight, or wasting, 0-60 months	0.254 P=458	-0.030 [0.038] N=1,838	-0.029 [0.040] N=1,703	-0.003 [0.043] N=1,533	-0.012 [0.043] N=1,311	0.005 [0.045] N=1,073
BMI-for-age z score below -2	0.118 P=290	0.012 [0.028] N=2,419	0.015 [0.029] N=2,252	0.020 [0.031] N=2,021	0.044 [0.033] N=1,731	0.050 [0.034] N=1,426
WHO 2007 BMI overweight above +1, 61+ months	0.383 P=201	0.029 [0.078] N=527	0.026 [0.079] N=497	0.065 [0.086] N=448	0.060 [0.095] N=388	0.080 [0.109] N=324
WHO 2007 BMI obesity above +2, 61+ months	0.208 P=110	-0.018 [0.060] N=527	-0.029 [0.061] N=497	-0.038 [0.064] N=448	-0.041 [0.070] N=388	-0.082 [0.084] N=324
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=66	0.013 [0.059] N=527	0.007 [0.060] N=497	-0.041 [0.066] N=448	-0.048 [0.073] N=388	-0.058 [0.085] N=324

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.094 [0.238] N=2,121	0.090 [0.255] N=1,940	-0.079 [0.276] N=1,709	0.090 [0.311] N=1,410	-0.224 [0.427] N=1,091
WHO weight-for-age z score	-0.046 [0.137] N=2,298	0.057 [0.152] N=2,098	0.091 [0.161] N=1,848	0.039 [0.186] N=1,525	-0.105 [0.236] N=1,180
WHO weight-for-length/height z score, 0-60 months	0.028 [0.226] N=1,594	0.060 [0.237] N=1,455	-0.037 [0.270] N=1,280	-0.374 [0.328] N=1,056	-0.378 [0.418] N=811
WHO BMI-for-age z score	0.109 [0.214] N=2,038	0.189 [0.224] N=1,871	0.222 [0.250] N=1,640	-0.060 [0.294] N=1,350	0.181 [0.381] N=1,045
Stunted: HAZ below -2	0.021 [0.045] N=2,121	0.037 [0.048] N=1,940	0.077 [0.054] N=1,709	0.075 [0.058] N=1,410	**0.146* [0.083] N=1,091**
Underweight: WAZ below -2	-0.006 [0.023] N=2,298	-0.014 [0.024] N=2,098	-0.023 [0.027] N=1,848	-0.020 [0.032] N=1,525	-0.006 [0.043] N=1,180
Wasted: WLZ/WHZ below -2, 0-60 months	0.002 [0.034] N=1,594	0.011 [0.036] N=1,455	0.034 [0.041] N=1,280	0.055 [0.052] N=1,056	0.057 [0.057] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.044 [0.049] N=1,594	0.049 [0.053] N=1,455	0.049 [0.059] N=1,280	-0.021 [0.071] N=1,056	-0.009 [0.093] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.032] N=1,594	0.037 [0.035] N=1,455	0.064 [0.040] N=1,280	0.025 [0.048] N=1,056	0.036 [0.067] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.026 [0.053] N=1,557	0.032 [0.056] N=1,422	0.094 [0.063] N=1,252	0.088 [0.070] N=1,030	**0.185** [0.081] N=792**
BMI-for-age z score below -2	-0.018 [0.033] N=2,038	-0.022 [0.035] N=1,871	-0.016 [0.039] N=1,640	0.008 [0.045] N=1,350	0.001 [0.051] N=1,045
WHO 2007 BMI overweight above +1, 61+ months	0.033 [0.118] N=431	0.023 [0.122] N=401	0.091 [0.133] N=352	0.088 [0.150] N=292	0.212 [0.209] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.049 [0.094] N=431	0.031 [0.098] N=401	0.027 [0.101] N=352	0.049 [0.116] N=292	0.013 [0.161] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.008 [0.067] N=431	-0.013 [0.069] N=401	-0.112 [0.075] N=352	**-0.159* [0.082] N=292**	**-0.229** [0.103] N=228**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	0.129 [0.157] N=2,523	0.134 [0.162] N=2,342	0.061 [0.170] N=2,111	0.135 [0.175] N=1,812	0.044 [0.193] N=1,493
WHO weight-for-age z score	0.329	0.006 [0.096] N=2,726	0.069 [0.104] N=2,526	0.097 [0.106] N=2,276	0.066 [0.109] N=1,953	0.038 [0.116] N=1,608
WHO weight-for-length/height z score, 0-60 months	0.342	0.004 [0.170] N=1,880	0.019 [0.176] N=1,741	-0.010 [0.188] N=1,566	-0.169 [0.213] N=1,342	-0.130 [0.228] N=1,097
WHO BMI-for-age z score	0.355	-0.035 [0.172] N=2,419	-0.009 [0.179] N=2,252	-0.006 [0.193] N=2,021	-0.174 [0.213] N=1,731	-0.106 [0.236] N=1,426
Stunted: HAZ below -2	0.194 P=475	-0.018 [0.032] N=2,523	-0.011 [0.032] N=2,342	0.002 [0.035] N=2,111	-0.008 [0.036] N=1,812	0.005 [0.041] N=1,493
Underweight: WAZ below -2	0.048 P=128	-0.012 [0.015] N=2,726	-0.016 [0.016] N=2,526	-0.023 [0.017] N=2,276	-0.021 [0.018] N=1,953	-0.019 [0.021] N=1,608
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=210	-0.005 [0.026] N=1,880	0.002 [0.027] N=1,741	0.007 [0.030] N=1,566	0.018 [0.033] N=1,342	0.012 [0.035] N=1,097
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=361	0.031 [0.038] N=1,880	0.032 [0.041] N=1,741	0.030 [0.043] N=1,566	-0.006 [0.048] N=1,342	-0.002 [0.052] N=1,097
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=160	0.010 [0.028] N=1,880	0.012 [0.030] N=1,741	0.021 [0.033] N=1,566	-0.002 [0.036] N=1,342	-0.007 [0.041] N=1,097
CIAF: any stunting, underweight, or wasting, 0-60 months	0.254 P=458	-0.016 [0.038] N=1,838	-0.013 [0.039] N=1,703	0.007 [0.042] N=1,533	-0.004 [0.041] N=1,311	0.018 [0.044] N=1,073
BMI-for-age z score below -2	0.118 P=290	0.019 [0.028] N=2,419	0.023 [0.029] N=2,252	0.028 [0.031] N=2,021	0.051 [0.033] N=1,731	**0.058* [0.034] N=1,426**
WHO 2007 BMI overweight above +1, 61+ months	0.383 P=201	0.042 [0.077] N=527	0.042 [0.079] N=497	0.082 [0.083] N=448	0.077 [0.092] N=388	0.097 [0.107] N=324
WHO 2007 BMI obesity above +2, 61+ months	0.208 P=110	-0.006 [0.060] N=527	-0.016 [0.061] N=497	-0.027 [0.063] N=448	-0.037 [0.067] N=388	-0.074 [0.084] N=324
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=66	0.022 [0.058] N=527	0.016 [0.059] N=497	-0.028 [0.064] N=448	-0.031 [0.071] N=388	-0.041 [0.082] N=324

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.064 [0.237] N=2,121	0.056 [0.252] N=1,940	-0.106 [0.275] N=1,709	0.051 [0.313] N=1,410	-0.282 [0.430] N=1,091
WHO weight-for-age z score	-0.082 [0.133] N=2,298	0.023 [0.150] N=2,098	0.056 [0.157] N=1,848	0.006 [0.180] N=1,525	-0.149 [0.234] N=1,180
WHO weight-for-length/height z score, 0-60 months	0.004 [0.226] N=1,594	0.047 [0.239] N=1,455	-0.043 [0.269] N=1,280	-0.392 [0.329] N=1,056	-0.410 [0.417] N=811
WHO BMI-for-age z score	0.103 [0.216] N=2,038	0.186 [0.226] N=1,871	0.219 [0.251] N=1,640	-0.056 [0.296] N=1,350	0.194 [0.385] N=1,045
Stunted: HAZ below -2	0.026 [0.045] N=2,121	0.044 [0.047] N=1,940	0.082 [0.053] N=1,709	0.079 [0.057] N=1,410	**0.153* [0.083] N=1,091**
Underweight: WAZ below -2	-0.002 [0.023] N=2,298	-0.008 [0.024] N=2,098	-0.018 [0.027] N=1,848	-0.016 [0.032] N=1,525	0.002 [0.043] N=1,180
Wasted: WLZ/WHZ below -2, 0-60 months	0.011 [0.034] N=1,594	0.020 [0.036] N=1,455	0.039 [0.041] N=1,280	0.063 [0.052] N=1,056	0.067 [0.059] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.047 [0.049] N=1,594	0.053 [0.054] N=1,455	0.051 [0.059] N=1,280	-0.021 [0.072] N=1,056	-0.001 [0.091] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.031 [0.032] N=1,594	0.040 [0.035] N=1,455	0.064 [0.040] N=1,280	0.022 [0.048] N=1,056	0.029 [0.066] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.041 [0.052] N=1,557	0.049 [0.056] N=1,422	0.101 [0.062] N=1,252	0.090 [0.069] N=1,030	**0.193** [0.083] N=792**
BMI-for-age z score below -2	-0.012 [0.033] N=2,038	-0.014 [0.035] N=1,871	-0.010 [0.040] N=1,640	0.014 [0.045] N=1,350	0.008 [0.053] N=1,045
WHO 2007 BMI overweight above +1, 61+ months	0.037 [0.117] N=431	0.028 [0.122] N=401	0.095 [0.132] N=352	0.084 [0.147] N=292	0.212 [0.209] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.058 [0.094] N=431	0.041 [0.099] N=401	0.040 [0.099] N=352	0.049 [0.114] N=292	0.038 [0.166] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.015 [0.067] N=431	-0.006 [0.069] N=401	-0.104 [0.075] N=352	**-0.134* [0.080] N=292**	**-0.217** [0.101] N=228**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	0.107 [0.156] N=2,523	0.113 [0.162] N=2,342	0.033 [0.172] N=2,111	0.093 [0.176] N=1,812	-0.010 [0.191] N=1,493
WHO weight-for-age z score	0.329	0.017 [0.096] N=2,726	0.076 [0.104] N=2,526	0.101 [0.107] N=2,276	0.061 [0.111] N=1,953	0.021 [0.119] N=1,608
WHO weight-for-length/height z score, 0-60 months	0.342	0.034 [0.171] N=1,880	0.048 [0.176] N=1,741	0.010 [0.189] N=1,566	-0.154 [0.215] N=1,342	-0.107 [0.228] N=1,097
WHO BMI-for-age z score	0.355	-0.017 [0.174] N=2,419	0.009 [0.180] N=2,252	0.010 [0.195] N=2,021	-0.156 [0.217] N=1,731	-0.082 [0.237] N=1,426
Stunted: HAZ below -2	0.194 P=475	-0.014 [0.032] N=2,523	-0.008 [0.033] N=2,342	0.007 [0.036] N=2,111	0.002 [0.036] N=1,812	0.014 [0.041] N=1,493
Underweight: WAZ below -2	0.048 P=128	-0.012 [0.015] N=2,726	-0.017 [0.016] N=2,526	-0.023 [0.017] N=2,276	-0.020 [0.018] N=1,953	-0.018 [0.021] N=1,608
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=210	-0.006 [0.026] N=1,880	-0.000 [0.027] N=1,741	0.007 [0.029] N=1,566	0.019 [0.033] N=1,342	0.011 [0.035] N=1,097
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=361	0.035 [0.038] N=1,880	0.035 [0.041] N=1,741	0.033 [0.043] N=1,566	-0.002 [0.048] N=1,342	-0.003 [0.052] N=1,097
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=160	0.015 [0.028] N=1,880	0.017 [0.030] N=1,741	0.024 [0.033] N=1,566	0.003 [0.037] N=1,342	0.001 [0.042] N=1,097
CIAF: any stunting, underweight, or wasting, 0-60 months	0.254 P=458	-0.014 [0.037] N=1,838	-0.012 [0.039] N=1,703	0.011 [0.042] N=1,533	0.006 [0.042] N=1,311	0.025 [0.045] N=1,073
BMI-for-age z score below -2	0.118 P=290	0.015 [0.027] N=2,419	0.018 [0.028] N=2,252	0.023 [0.031] N=2,021	0.048 [0.033] N=1,731	0.052 [0.034] N=1,426
WHO 2007 BMI overweight above +1, 61+ months	0.383 P=201	0.035 [0.077] N=527	0.032 [0.079] N=497	0.067 [0.086] N=448	0.065 [0.094] N=388	0.091 [0.108] N=324
WHO 2007 BMI obesity above +2, 61+ months	0.208 P=110	-0.011 [0.060] N=527	-0.022 [0.061] N=497	-0.034 [0.064] N=448	-0.040 [0.070] N=388	-0.076 [0.084] N=324
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=66	0.014 [0.058] N=527	0.009 [0.059] N=497	-0.043 [0.064] N=448	-0.046 [0.071] N=388	-0.053 [0.082] N=324

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.062 [0.235] N=2,121	0.059 [0.251] N=1,940	-0.102 [0.273] N=1,709	0.064 [0.309] N=1,410	-0.301 [0.419] N=1,091
WHO weight-for-age z score	-0.066 [0.135] N=2,298	0.034 [0.151] N=2,098	0.064 [0.160] N=1,848	0.009 [0.184] N=1,525	-0.143 [0.234] N=1,180
WHO weight-for-length/height z score, 0-60 months	0.022 [0.228] N=1,594	0.071 [0.237] N=1,455	-0.037 [0.271] N=1,280	-0.387 [0.331] N=1,056	-0.379 [0.421] N=811
WHO BMI-for-age z score	0.108 [0.217] N=2,038	0.189 [0.226] N=1,871	0.219 [0.254] N=1,640	-0.069 [0.296] N=1,350	0.180 [0.382] N=1,045
Stunted: HAZ below -2	0.027 [0.045] N=2,121	0.044 [0.047] N=1,940	0.082 [0.053] N=1,709	0.079 [0.057] N=1,410	**0.157* [0.081] N=1,091**
Underweight: WAZ below -2	-0.003 [0.023] N=2,298	-0.010 [0.024] N=2,098	-0.018 [0.027] N=1,848	-0.016 [0.032] N=1,525	0.002 [0.043] N=1,180
Wasted: WLZ/WHZ below -2, 0-60 months	0.010 [0.034] N=1,594	0.016 [0.037] N=1,455	0.038 [0.040] N=1,280	0.062 [0.052] N=1,056	0.065 [0.059] N=811
Overweight: WLZ/WHZ above +2, 0-60 months	0.050 [0.050] N=1,594	0.055 [0.053] N=1,455	0.052 [0.060] N=1,280	-0.018 [0.072] N=1,056	0.002 [0.092] N=811
Obese: WLZ/WHZ above +3, 0-60 months	0.034 [0.032] N=1,594	0.043 [0.035] N=1,455	0.065 [0.040] N=1,280	0.022 [0.048] N=1,056	0.033 [0.065] N=811
CIAF: any stunting, underweight, or wasting, 0-60 months	0.043 [0.052] N=1,557	0.047 [0.056] N=1,422	0.102 [0.063] N=1,252	0.093 [0.069] N=1,030	**0.197** [0.083] N=792**
BMI-for-age z score below -2	-0.011 [0.033] N=2,038	-0.016 [0.035] N=1,871	-0.010 [0.040] N=1,640	0.018 [0.046] N=1,350	0.014 [0.053] N=1,045
WHO 2007 BMI overweight above +1, 61+ months	0.039 [0.118] N=431	0.027 [0.123] N=401	0.093 [0.133] N=352	0.096 [0.149] N=292	0.257 [0.210] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.065 [0.092] N=431	0.045 [0.097] N=401	0.030 [0.099] N=352	0.034 [0.116] N=292	0.021 [0.168] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.020 [0.067] N=431	0.003 [0.070] N=401	-0.106 [0.075] N=352	**-0.147* [0.082] N=292**	**-0.199* [0.105] N=228**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.104	0.133 [0.168] N=2,356	0.110 [0.177] N=2,191	0.017 [0.183] N=1,969	0.064 [0.188] N=1,698	-0.052 [0.207] N=1,395
WHO weight-for-age z score	0.323	0.075 [0.096] N=2,542	0.137 [0.100] N=2,360	0.169 [0.106] N=2,121	0.130 [0.110] N=1,828	0.116 [0.122] N=1,504
WHO weight-for-length/height z score, 0-60 months	0.360	0.122 [0.172] N=1,748	0.145 [0.176] N=1,624	0.099 [0.188] N=1,457	-0.049 [0.215] N=1,258	0.063 [0.229] N=1,026
WHO BMI-for-age z score	0.371	0.033 [0.169] N=2,261	0.052 [0.173] N=2,109	0.070 [0.191] N=1,889	-0.077 [0.214] N=1,626	0.016 [0.237] N=1,336
Stunted: HAZ below -2	0.196 P=447	-0.019 [0.034] N=2,356	-0.010 [0.035] N=2,191	0.006 [0.038] N=1,969	0.002 [0.039] N=1,698	0.017 [0.046] N=1,395
Underweight: WAZ below -2	0.049 P=122	-0.021 [0.017] N=2,542	-0.027 [0.017] N=2,360	**-0.033* [0.019] N=2,121**	-0.030 [0.020] N=1,828	-0.030 [0.024] N=1,504
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=194	-0.016 [0.027] N=1,748	-0.012 [0.028] N=1,624	0.000 [0.031] N=1,457	0.010 [0.036] N=1,258	-0.002 [0.037] N=1,026
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=339	0.040 [0.040] N=1,748	0.040 [0.041] N=1,624	0.038 [0.044] N=1,457	0.001 [0.049] N=1,258	0.012 [0.054] N=1,026
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=153	0.012 [0.029] N=1,748	0.017 [0.031] N=1,624	0.027 [0.033] N=1,457	0.008 [0.037] N=1,258	0.008 [0.043] N=1,026
CIAF: any stunting, underweight, or wasting, 0-60 months	0.255 P=428	-0.022 [0.043] N=1,708	-0.022 [0.045] N=1,588	0.003 [0.049] N=1,425	-0.008 [0.049] N=1,228	0.010 [0.053] N=1,002
BMI-for-age z score below -2	0.115 P=264	0.007 [0.026] N=2,261	0.008 [0.026] N=2,109	0.012 [0.029] N=1,889	0.031 [0.032] N=1,626	0.026 [0.032] N=1,336
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=191	0.025 [0.081] N=505	0.022 [0.083] N=476	0.073 [0.089] N=429	0.061 [0.096] N=371	0.089 [0.112] N=309
WHO 2007 BMI obesity above +2, 61+ months	0.201 P=103	-0.025 [0.061] N=505	-0.031 [0.062] N=476	-0.039 [0.065] N=429	-0.051 [0.068] N=371	-0.099 [0.085] N=309
WHO 2007 BMI thinness below -2, 61+ months	0.116 P=62	0.019 [0.058] N=505	0.016 [0.059] N=476	-0.034 [0.063] N=429	-0.046 [0.069] N=371	-0.060 [0.081] N=309

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.166 [0.249] N=1,982	0.134 [0.270] N=1,817	-0.035 [0.293] N=1,595	0.122 [0.333] N=1,324	-0.242 [0.475] N=1,021
WHO weight-for-age z score	-0.045 [0.133] N=2,143	0.053 [0.144] N=1,961	0.081 [0.160] N=1,722	0.015 [0.187] N=1,429	-0.129 [0.242] N=1,105
WHO weight-for-length/height z score, 0-60 months	0.025 [0.234] N=1,482	0.063 [0.241] N=1,358	-0.061 [0.268] N=1,191	-0.361 [0.339] N=992	-0.245 [0.431] N=760
WHO BMI-for-age z score	0.085 [0.219] N=1,906	0.145 [0.222] N=1,754	0.195 [0.245] N=1,534	-0.037 [0.291] N=1,271	0.252 [0.393] N=981
Stunted: HAZ below -2	0.016 [0.047] N=1,982	0.035 [0.050] N=1,817	0.076 [0.056] N=1,595	0.075 [0.062] N=1,324	0.149 [0.090] N=1,021
Underweight: WAZ below -2	-0.007 [0.024] N=2,143	-0.014 [0.025] N=1,961	-0.020 [0.029] N=1,722	-0.014 [0.034] N=1,429	0.003 [0.043] N=1,105
Wasted: WLZ/WHZ below -2, 0-60 months	-0.000 [0.034] N=1,482	0.008 [0.035] N=1,358	0.038 [0.039] N=1,191	0.049 [0.051] N=992	0.027 [0.053] N=760
Overweight: WLZ/WHZ above +2, 0-60 months	0.048 [0.051] N=1,482	0.054 [0.054] N=1,358	0.056 [0.060] N=1,191	-0.017 [0.074] N=992	0.013 [0.099] N=760
Obese: WLZ/WHZ above +3, 0-60 months	0.025 [0.034] N=1,482	0.035 [0.037] N=1,358	0.060 [0.041] N=1,191	0.025 [0.051] N=992	0.046 [0.072] N=760
CIAF: any stunting, underweight, or wasting, 0-60 months	0.033 [0.056] N=1,447	0.038 [0.059] N=1,327	0.098 [0.067] N=1,164	0.071 [0.075] N=967	**0.148* [0.087] N=741**
BMI-for-age z score below -2	-0.002 [0.033] N=1,906	-0.005 [0.033] N=1,754	0.002 [0.037] N=1,534	0.018 [0.043] N=1,271	-0.012 [0.048] N=981
WHO 2007 BMI overweight above +1, 61+ months	0.017 [0.125] N=413	0.006 [0.129] N=384	0.096 [0.141] N=337	0.068 [0.154] N=279	0.187 [0.216] N=217
WHO 2007 BMI obesity above +2, 61+ months	0.034 [0.097] N=413	0.022 [0.102] N=384	0.014 [0.105] N=337	0.013 [0.116] N=279	-0.062 [0.163] N=217
WHO 2007 BMI thinness below -2, 61+ months	0.033 [0.071] N=413	0.012 [0.075] N=384	-0.091 [0.080] N=337	**-0.156* [0.083] N=279**	**-0.237** [0.108] N=217**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.104	0.145 [0.166] N=2,356	0.120 [0.174] N=2,191	0.016 [0.182] N=1,969	0.063 [0.186] N=1,698	-0.049 [0.205] N=1,395
WHO weight-for-age z score	0.323	0.087 [0.097] N=2,542	0.146 [0.102] N=2,360	**0.177* [0.106] N=2,121**	0.139 [0.111] N=1,828	0.121 [0.123] N=1,504
WHO weight-for-length/height z score, 0-60 months	0.360	0.122 [0.173] N=1,748	0.146 [0.176] N=1,624	0.099 [0.189] N=1,457	-0.053 [0.215] N=1,258	0.049 [0.232] N=1,026
WHO BMI-for-age z score	0.371	0.043 [0.169] N=2,261	0.060 [0.172] N=2,109	0.075 [0.191] N=1,889	-0.074 [0.214] N=1,626	0.011 [0.238] N=1,336
Stunted: HAZ below -2	0.196 P=447	-0.021 [0.034] N=2,356	-0.011 [0.035] N=2,191	0.006 [0.038] N=1,969	0.001 [0.039] N=1,698	0.015 [0.045] N=1,395
Underweight: WAZ below -2	0.049 P=122	-0.024 [0.017] N=2,542	**-0.029* [0.017] N=2,360**	**-0.036* [0.019] N=2,121**	-0.033 [0.020] N=1,828	-0.033 [0.024] N=1,504
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=194	-0.016 [0.027] N=1,748	-0.012 [0.028] N=1,624	0.000 [0.031] N=1,457	0.010 [0.036] N=1,258	-0.003 [0.037] N=1,026
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=339	0.039 [0.040] N=1,748	0.040 [0.042] N=1,624	0.038 [0.045] N=1,457	0.001 [0.050] N=1,258	0.012 [0.055] N=1,026
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=153	0.012 [0.029] N=1,748	0.016 [0.031] N=1,624	0.027 [0.033] N=1,457	0.008 [0.037] N=1,258	0.009 [0.043] N=1,026
CIAF: any stunting, underweight, or wasting, 0-60 months	0.255 P=428	-0.023 [0.043] N=1,708	-0.022 [0.044] N=1,588	0.004 [0.049] N=1,425	-0.007 [0.049] N=1,228	0.008 [0.052] N=1,002
BMI-for-age z score below -2	0.115 P=264	0.006 [0.026] N=2,261	0.007 [0.026] N=2,109	0.012 [0.029] N=1,889	0.031 [0.032] N=1,626	0.026 [0.033] N=1,336
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=191	0.021 [0.078] N=505	0.017 [0.080] N=476	0.059 [0.085] N=429	0.051 [0.093] N=371	0.064 [0.111] N=309
WHO 2007 BMI obesity above +2, 61+ months	0.201 P=103	-0.026 [0.060] N=505	-0.037 [0.062] N=476	-0.043 [0.064] N=429	-0.052 [0.068] N=371	-0.105 [0.083] N=309
WHO 2007 BMI thinness below -2, 61+ months	0.116 P=62	0.016 [0.057] N=505	0.012 [0.058] N=476	-0.035 [0.062] N=429	-0.048 [0.069] N=371	-0.055 [0.079] N=309

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.183 [0.245] N=1,982	0.151 [0.267] N=1,817	-0.031 [0.289] N=1,595	0.118 [0.329] N=1,324	-0.221 [0.468] N=1,021
WHO weight-for-age z score	-0.026 [0.134] N=2,143	0.070 [0.145] N=1,961	0.099 [0.159] N=1,722	0.040 [0.185] N=1,429	-0.101 [0.241] N=1,105
WHO weight-for-length/height z score, 0-60 months	0.045 [0.235] N=1,482	0.089 [0.242] N=1,358	-0.052 [0.268] N=1,191	-0.352 [0.339] N=992	-0.246 [0.434] N=760
WHO BMI-for-age z score	0.111 [0.218] N=1,906	0.173 [0.223] N=1,754	0.210 [0.245] N=1,534	-0.025 [0.291] N=1,271	0.255 [0.396] N=981
Stunted: HAZ below -2	0.013 [0.047] N=1,982	0.033 [0.050] N=1,817	0.073 [0.056] N=1,595	0.071 [0.061] N=1,324	0.140 [0.090] N=1,021
Underweight: WAZ below -2	-0.009 [0.024] N=2,143	-0.016 [0.025] N=1,961	-0.024 [0.029] N=1,722	-0.020 [0.035] N=1,429	-0.004 [0.044] N=1,105
Wasted: WLZ/WHZ below -2, 0-60 months	-0.002 [0.034] N=1,482	0.006 [0.035] N=1,358	0.038 [0.039] N=1,191	0.048 [0.051] N=992	0.021 [0.052] N=760
Overweight: WLZ/WHZ above +2, 0-60 months	0.051 [0.050] N=1,482	0.057 [0.054] N=1,358	0.057 [0.060] N=1,191	-0.014 [0.075] N=992	0.016 [0.101] N=760
Obese: WLZ/WHZ above +3, 0-60 months	0.024 [0.034] N=1,482	0.034 [0.037] N=1,358	0.060 [0.041] N=1,191	0.027 [0.051] N=992	0.050 [0.072] N=760
CIAF: any stunting, underweight, or wasting, 0-60 months	0.031 [0.056] N=1,447	0.037 [0.059] N=1,327	0.097 [0.067] N=1,164	0.068 [0.074] N=967	0.132 [0.085] N=741
BMI-for-age z score below -2	-0.006 [0.033] N=1,906	-0.010 [0.033] N=1,754	0.000 [0.037] N=1,534	0.017 [0.043] N=1,271	-0.013 [0.049] N=981
WHO 2007 BMI overweight above +1, 61+ months	0.026 [0.120] N=413	0.019 [0.126] N=384	0.095 [0.136] N=337	0.073 [0.148] N=279	0.182 [0.213] N=217
WHO 2007 BMI obesity above +2, 61+ months	0.035 [0.096] N=413	0.016 [0.101] N=384	0.016 [0.104] N=337	0.013 [0.114] N=279	-0.064 [0.158] N=217
WHO 2007 BMI thinness below -2, 61+ months	0.025 [0.070] N=413	0.006 [0.074] N=384	-0.096 [0.080] N=337	**-0.160* [0.083] N=279**	**-0.239** [0.107] N=217**

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.075	-0.211 [0.202] N=2,423	-0.261 [0.216] N=2,189	-0.287 [0.229] N=1,918	**-0.457* [0.247] N=1,632**	**-0.471* [0.275] N=1,328**
WHO weight-for-age z score	0.345	-0.081 [0.099] N=2,615	-0.082 [0.103] N=2,375	-0.157 [0.110] N=2,078	-0.200 [0.121] N=1,761	-0.188 [0.129] N=1,426
WHO weight-for-length/height z score, 0-60 months	0.380	-0.099 [0.186] N=1,765	-0.077 [0.195] N=1,605	-0.122 [0.216] N=1,395	-0.035 [0.233] N=1,193	-0.121 [0.279] N=961
WHO BMI-for-age z score	0.384	-0.056 [0.189] N=2,323	-0.062 [0.199] N=2,100	-0.052 [0.219] N=1,841	0.073 [0.238] N=1,565	0.026 [0.275] N=1,275
Stunted: HAZ below -2	0.196 P=468	0.031 [0.035] N=2,423	0.032 [0.037] N=2,189	0.023 [0.040] N=1,918	0.031 [0.045] N=1,632	0.032 [0.051] N=1,328
Underweight: WAZ below -2	0.042 P=113	0.011 [0.014] N=2,615	0.006 [0.015] N=2,375	0.013 [0.016] N=2,078	0.006 [0.016] N=1,761	0.006 [0.017] N=1,426
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=190	0.028 [0.032] N=1,765	0.015 [0.035] N=1,605	0.026 [0.037] N=1,395	-0.004 [0.038] N=1,193	0.009 [0.047] N=961
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.006 [0.036] N=1,765	0.004 [0.037] N=1,605	-0.006 [0.040] N=1,395	0.002 [0.043] N=1,193	-0.001 [0.049] N=961
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=167	0.000 [0.027] N=1,765	-0.005 [0.028] N=1,605	-0.016 [0.030] N=1,395	-0.009 [0.034] N=1,193	-0.036 [0.036] N=961
CIAF: any stunting, underweight, or wasting, 0-60 months	0.261 P=445	0.060 [0.041] N=1,727	0.047 [0.045] N=1,567	0.045 [0.046] N=1,360	0.005 [0.050] N=1,163	0.036 [0.055] N=935
BMI-for-age z score below -2	0.118 P=272	0.041 [0.031] N=2,323	0.033 [0.034] N=2,100	0.033 [0.036] N=1,841	0.014 [0.037] N=1,565	0.026 [0.044] N=1,275
WHO 2007 BMI overweight above +1, 61+ months	0.406 P=219	0.030 [0.079] N=552	-0.021 [0.081] N=489	0.016 [0.082] N=441	0.066 [0.089] N=369	-0.011 [0.100] N=304
WHO 2007 BMI obesity above +2, 61+ months	0.221 P=118	0.052 [0.078] N=552	-0.016 [0.081] N=489	-0.005 [0.088] N=441	0.036 [0.096] N=369	-0.031 [0.113] N=304
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=67	0.040 [0.055] N=552	0.057 [0.057] N=489	0.035 [0.060] N=441	0.031 [0.065] N=369	0.049 [0.074] N=304

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.007 [0.256] N=2,080	-0.060 [0.286] N=1,846	-0.072 [0.317] N=1,575	-0.361 [0.355] N=1,289	-0.283 [0.431] N=985
WHO weight-for-age z score	-0.058 [0.149] N=2,248	-0.050 [0.160] N=2,008	-0.219 [0.178] N=1,711	-0.311 [0.213] N=1,394	-0.319 [0.245] N=1,059
WHO weight-for-length/height z score, 0-60 months	-0.232 [0.244] N=1,531	-0.224 [0.256] N=1,371	-0.386 [0.305] N=1,161	-0.302 [0.337] N=959	**-0.848** [0.415] N=727**
WHO BMI-for-age z score	**-0.426** [0.212] N=1,998**	**-0.510** [0.219] N=1,775**	**-0.620** [0.260] N=1,516**	**-0.544* [0.299] N=1,240**	**-1.093*** [0.369] N=950**
Stunted: HAZ below -2	0.027 [0.046] N=2,080	0.031 [0.050] N=1,846	0.015 [0.054] N=1,575	0.024 [0.067] N=1,289	0.013 [0.090] N=985
Underweight: WAZ below -2	0.004 [0.022] N=2,248	-0.007 [0.024] N=2,008	0.007 [0.028] N=1,711	-0.018 [0.030] N=1,394	-0.033 [0.033] N=1,059
Wasted: WLZ/WHZ below -2, 0-60 months	**0.082** [0.037] N=1,531**	0.067 [0.042] N=1,371	**0.110** [0.047] N=1,161**	0.058 [0.048] N=959	**0.144*** [0.053] N=727**
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.053] N=1,531	-0.001 [0.056] N=1,371	-0.021 [0.062] N=1,161	-0.025 [0.074] N=959	-0.083 [0.091] N=727
Obese: WLZ/WHZ above +3, 0-60 months	0.014 [0.041] N=1,531	0.001 [0.042] N=1,371	-0.015 [0.049] N=1,161	-0.007 [0.062] N=959	-0.109 [0.076] N=727
CIAF: any stunting, underweight, or wasting, 0-60 months	0.093 [0.062] N=1,500	0.075 [0.069] N=1,340	0.087 [0.078] N=1,133	0.003 [0.090] N=936	0.078 [0.109] N=708
BMI-for-age z score below -2	**0.096*** [0.030] N=1,998**	**0.093*** [0.035] N=1,775**	**0.115*** [0.040] N=1,516**	**0.099*** [0.037] N=1,240**	**0.186*** [0.047] N=950**
WHO 2007 BMI overweight above +1, 61+ months	-0.047 [0.131] N=464	-0.155 [0.140] N=401	-0.110 [0.154] N=353	-0.026 [0.200] N=281	-0.358 [0.299] N=216
WHO 2007 BMI obesity above +2, 61+ months	0.080 [0.115] N=464	-0.045 [0.121] N=401	-0.038 [0.141] N=353	0.028 [0.162] N=281	-0.144 [0.245] N=216
WHO 2007 BMI thinness below -2, 61+ months	0.097 [0.070] N=464	**0.149* [0.077] N=401**	0.114 [0.083] N=353	0.136 [0.092] N=281	**0.264** [0.115] N=216**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.075	-0.208 [0.202] N=2,423	-0.259 [0.216] N=2,189	-0.285 [0.229] N=1,918	**-0.454* [0.247] N=1,632**	**-0.459* [0.275] N=1,328**
WHO weight-for-age z score	0.345	-0.078 [0.099] N=2,615	-0.079 [0.102] N=2,375	-0.155 [0.110] N=2,078	-0.198 [0.121] N=1,761	-0.186 [0.130] N=1,426
WHO weight-for-length/height z score, 0-60 months	0.380	-0.096 [0.186] N=1,765	-0.076 [0.194] N=1,605	-0.122 [0.215] N=1,395	-0.037 [0.233] N=1,193	-0.133 [0.277] N=961
WHO BMI-for-age z score	0.384	-0.058 [0.189] N=2,323	-0.065 [0.199] N=2,100	-0.053 [0.219] N=1,841	0.071 [0.238] N=1,565	0.018 [0.276] N=1,275
Stunted: HAZ below -2	0.196 P=468	0.031 [0.035] N=2,423	0.032 [0.037] N=2,189	0.024 [0.040] N=1,918	0.032 [0.045] N=1,632	0.033 [0.051] N=1,328
Underweight: WAZ below -2	0.042 P=113	0.011 [0.014] N=2,615	0.005 [0.015] N=2,375	0.013 [0.016] N=2,078	0.007 [0.016] N=1,761	0.006 [0.017] N=1,426
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=190	0.027 [0.032] N=1,765	0.015 [0.035] N=1,605	0.026 [0.037] N=1,395	-0.003 [0.038] N=1,193	0.013 [0.046] N=961
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.006 [0.036] N=1,765	0.004 [0.037] N=1,605	-0.006 [0.040] N=1,395	0.002 [0.043] N=1,193	-0.000 [0.049] N=961
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=167	-0.000 [0.027] N=1,765	-0.005 [0.028] N=1,605	-0.016 [0.031] N=1,395	-0.009 [0.034] N=1,193	-0.033 [0.036] N=961
CIAF: any stunting, underweight, or wasting, 0-60 months	0.261 P=445	0.060 [0.041] N=1,727	0.046 [0.045] N=1,567	0.044 [0.046] N=1,360	0.005 [0.050] N=1,163	0.038 [0.055] N=935
BMI-for-age z score below -2	0.118 P=272	0.042 [0.031] N=2,323	0.034 [0.034] N=2,100	0.035 [0.036] N=1,841	0.015 [0.037] N=1,565	0.028 [0.045] N=1,275
WHO 2007 BMI overweight above +1, 61+ months	0.406 P=219	0.038 [0.079] N=552	-0.014 [0.081] N=489	0.020 [0.082] N=441	0.070 [0.089] N=369	-0.005 [0.099] N=304
WHO 2007 BMI obesity above +2, 61+ months	0.221 P=118	0.064 [0.079] N=552	-0.007 [0.082] N=489	0.002 [0.089] N=441	0.040 [0.097] N=369	-0.025 [0.113] N=304
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=67	0.035 [0.055] N=552	0.053 [0.057] N=489	0.029 [0.059] N=441	0.028 [0.065] N=369	0.042 [0.073] N=304

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.003 [0.255] N=2,080	-0.067 [0.285] N=1,846	-0.076 [0.316] N=1,575	-0.377 [0.350] N=1,289	-0.290 [0.431] N=985
WHO weight-for-age z score	-0.058 [0.149] N=2,248	-0.051 [0.159] N=2,008	-0.220 [0.177] N=1,711	-0.313 [0.213] N=1,394	-0.319 [0.245] N=1,059
WHO weight-for-length/height z score, 0-60 months	-0.218 [0.246] N=1,531	-0.212 [0.258] N=1,371	-0.377 [0.307] N=1,161	-0.290 [0.338] N=959	**-0.838** [0.420] N=727**
WHO BMI-for-age z score	**-0.423** [0.213] N=1,998**	**-0.507** [0.220] N=1,775**	**-0.619** [0.261] N=1,516**	**-0.541* [0.299] N=1,240**	**-1.092*** [0.372] N=950**
Stunted: HAZ below -2	0.027 [0.046] N=2,080	0.032 [0.050] N=1,846	0.015 [0.054] N=1,575	0.025 [0.067] N=1,289	0.013 [0.090] N=985
Underweight: WAZ below -2	0.004 [0.022] N=2,248	-0.007 [0.024] N=2,008	0.007 [0.028] N=1,711	-0.018 [0.030] N=1,394	-0.033 [0.033] N=1,059
Wasted: WLZ/WHZ below -2, 0-60 months	**0.078** [0.037] N=1,531**	0.064 [0.042] N=1,371	**0.106** [0.047] N=1,161**	0.054 [0.049] N=959	**0.141** [0.054] N=727**
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.053] N=1,531	-0.001 [0.057] N=1,371	-0.023 [0.062] N=1,161	-0.026 [0.074] N=959	-0.083 [0.091] N=727
Obese: WLZ/WHZ above +3, 0-60 months	0.013 [0.041] N=1,531	-0.000 [0.042] N=1,371	-0.018 [0.049] N=1,161	-0.010 [0.062] N=959	-0.110 [0.076] N=727
CIAF: any stunting, underweight, or wasting, 0-60 months	0.089 [0.062] N=1,500	0.071 [0.069] N=1,340	0.081 [0.078] N=1,133	-0.002 [0.092] N=936	0.076 [0.110] N=708
BMI-for-age z score below -2	**0.095*** [0.030] N=1,998**	**0.092*** [0.035] N=1,775**	**0.113*** [0.040] N=1,516**	**0.097*** [0.037] N=1,240**	**0.186*** [0.047] N=950**
WHO 2007 BMI overweight above +1, 61+ months	-0.041 [0.132] N=464	-0.151 [0.141] N=401	-0.107 [0.155] N=353	-0.026 [0.201] N=281	-0.354 [0.301] N=216
WHO 2007 BMI obesity above +2, 61+ months	0.091 [0.118] N=464	-0.038 [0.125] N=401	-0.032 [0.145] N=353	0.027 [0.165] N=281	-0.140 [0.249] N=216
WHO 2007 BMI thinness below -2, 61+ months	0.092 [0.071] N=464	**0.146* [0.077] N=401**	0.108 [0.082] N=353	0.137 [0.091] N=281	**0.254** [0.116] N=216**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.075	-0.208 [0.200] N=2,423	-0.248 [0.213] N=2,189	-0.270 [0.228] N=1,918	**-0.445* [0.242] N=1,632**	-0.446 [0.269] N=1,328
WHO weight-for-age z score	0.345	-0.076 [0.097] N=2,615	-0.072 [0.101] N=2,375	-0.135 [0.109] N=2,078	-0.191 [0.119] N=1,761	-0.180 [0.130] N=1,426
WHO weight-for-length/height z score, 0-60 months	0.380	-0.099 [0.187] N=1,765	-0.077 [0.195] N=1,605	-0.104 [0.219] N=1,395	-0.021 [0.236] N=1,193	-0.105 [0.282] N=961
WHO BMI-for-age z score	0.384	-0.055 [0.188] N=2,323	-0.060 [0.199] N=2,100	-0.038 [0.218] N=1,841	0.078 [0.237] N=1,565	0.017 [0.276] N=1,275
Stunted: HAZ below -2	0.196 P=468	0.031 [0.035] N=2,423	0.031 [0.037] N=2,189	0.021 [0.040] N=1,918	0.030 [0.044] N=1,632	0.030 [0.050] N=1,328
Underweight: WAZ below -2	0.042 P=113	0.011 [0.014] N=2,615	0.005 [0.014] N=2,375	0.011 [0.015] N=2,078	0.006 [0.015] N=1,761	0.006 [0.016] N=1,426
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=190	0.029 [0.033] N=1,765	0.015 [0.035] N=1,605	0.025 [0.038] N=1,395	-0.003 [0.039] N=1,193	0.011 [0.046] N=961
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.008 [0.036] N=1,765	0.006 [0.038] N=1,605	-0.001 [0.040] N=1,395	0.007 [0.044] N=1,193	0.006 [0.050] N=961
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=167	0.002 [0.028] N=1,765	-0.004 [0.028] N=1,605	-0.013 [0.031] N=1,395	-0.006 [0.035] N=1,193	-0.029 [0.038] N=961
CIAF: any stunting, underweight, or wasting, 0-60 months	0.261 P=445	**0.069* [0.041] N=1,727**	0.052 [0.044] N=1,567	0.048 [0.046] N=1,360	0.010 [0.050] N=1,163	0.039 [0.055] N=935
BMI-for-age z score below -2	0.118 P=272	0.041 [0.031] N=2,323	0.032 [0.034] N=2,100	0.032 [0.036] N=1,841	0.013 [0.037] N=1,565	0.027 [0.045] N=1,275
WHO 2007 BMI overweight above +1, 61+ months	0.406 P=219	0.010 [0.082] N=552	-0.043 [0.084] N=489	-0.007 [0.085] N=441	0.042 [0.095] N=369	-0.045 [0.105] N=304
WHO 2007 BMI obesity above +2, 61+ months	0.221 P=118	0.042 [0.079] N=552	-0.025 [0.083] N=489	-0.016 [0.089] N=441	0.021 [0.097] N=369	-0.034 [0.114] N=304
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=67	0.024 [0.054] N=552	0.035 [0.056] N=489	0.013 [0.057] N=441	0.018 [0.062] N=369	0.027 [0.070] N=304

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.022 [0.256] N=2,080	-0.035 [0.285] N=1,846	-0.017 [0.316] N=1,575	-0.322 [0.347] N=1,289	-0.202 [0.425] N=985
WHO weight-for-age z score	-0.056 [0.147] N=2,248	-0.046 [0.158] N=2,008	-0.188 [0.176] N=1,711	-0.305 [0.211] N=1,394	-0.307 [0.243] N=1,059
WHO weight-for-length/height z score, 0-60 months	-0.235 [0.249] N=1,531	-0.226 [0.263] N=1,371	-0.369 [0.312] N=1,161	-0.286 [0.342] N=959	**-0.810* [0.424] N=727**
WHO BMI-for-age z score	**-0.433** [0.213] N=1,998**	**-0.515** [0.221] N=1,775**	**-0.626** [0.260] N=1,516**	**-0.563* [0.299] N=1,240**	**-1.132*** [0.379] N=950**
Stunted: HAZ below -2	0.023 [0.046] N=2,080	0.025 [0.051] N=1,846	0.003 [0.055] N=1,575	0.012 [0.067] N=1,289	-0.005 [0.091] N=985
Underweight: WAZ below -2	0.002 [0.021] N=2,248	-0.010 [0.023] N=2,008	0.001 [0.027] N=1,711	-0.023 [0.029] N=1,394	-0.037 [0.032] N=1,059
Wasted: WLZ/WHZ below -2, 0-60 months	**0.082** [0.038] N=1,531**	0.065 [0.043] N=1,371	**0.103** [0.049] N=1,161**	0.051 [0.051] N=959	**0.136** [0.054] N=727**
Overweight: WLZ/WHZ above +2, 0-60 months	0.007 [0.054] N=1,531	-0.002 [0.058] N=1,371	-0.021 [0.064] N=1,161	-0.026 [0.076] N=959	-0.080 [0.094] N=727
Obese: WLZ/WHZ above +3, 0-60 months	0.014 [0.041] N=1,531	-0.000 [0.043] N=1,371	-0.017 [0.050] N=1,161	-0.010 [0.063] N=959	-0.107 [0.077] N=727
CIAF: any stunting, underweight, or wasting, 0-60 months	0.098 [0.063] N=1,500	0.074 [0.071] N=1,340	0.078 [0.080] N=1,133	-0.004 [0.094] N=936	0.065 [0.113] N=708
BMI-for-age z score below -2	**0.095*** [0.030] N=1,998**	**0.089** [0.035] N=1,775**	**0.109*** [0.040] N=1,516**	**0.094** [0.038] N=1,240**	**0.183*** [0.048] N=950**
WHO 2007 BMI overweight above +1, 61+ months	-0.088 [0.130] N=464	-0.200 [0.138] N=401	-0.164 [0.152] N=353	-0.103 [0.201] N=281	**-0.504* [0.296] N=216**
WHO 2007 BMI obesity above +2, 61+ months	0.052 [0.118] N=464	-0.075 [0.124] N=401	-0.062 [0.143] N=353	-0.014 [0.164] N=281	-0.191 [0.246] N=216
WHO 2007 BMI thinness below -2, 61+ months	0.081 [0.071] N=464	0.127 [0.078] N=401	0.085 [0.084] N=353	0.122 [0.092] N=281	**0.237** [0.112] N=216**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.075	-0.212 [0.200] N=2,423	-0.250 [0.213] N=2,189	-0.276 [0.229] N=1,918	**-0.441* [0.243] N=1,632**	**-0.457* [0.268] N=1,328**
WHO weight-for-age z score	0.345	-0.081 [0.095] N=2,615	-0.075 [0.099] N=2,375	-0.145 [0.106] N=2,078	**-0.191* [0.115] N=1,761**	-0.192 [0.121] N=1,426
WHO weight-for-length/height z score, 0-60 months	0.380	-0.093 [0.187] N=1,765	-0.072 [0.194] N=1,605	-0.110 [0.216] N=1,395	-0.020 [0.233] N=1,193	-0.110 [0.280] N=961
WHO BMI-for-age z score	0.384	-0.058 [0.189] N=2,323	-0.065 [0.199] N=2,100	-0.046 [0.218] N=1,841	0.077 [0.235] N=1,565	0.021 [0.275] N=1,275
Stunted: HAZ below -2	0.196 P=468	0.033 [0.035] N=2,423	0.032 [0.037] N=2,189	0.023 [0.040] N=1,918	0.032 [0.044] N=1,632	0.033 [0.049] N=1,328
Underweight: WAZ below -2	0.042 P=113	0.011 [0.014] N=2,615	0.005 [0.014] N=2,375	0.012 [0.015] N=2,078	0.006 [0.016] N=1,761	0.006 [0.017] N=1,426
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=190	0.029 [0.032] N=1,765	0.017 [0.035] N=1,605	0.029 [0.037] N=1,395	-0.002 [0.038] N=1,193	0.012 [0.046] N=961
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.008 [0.037] N=1,765	0.006 [0.038] N=1,605	-0.000 [0.040] N=1,395	0.008 [0.043] N=1,193	0.004 [0.050] N=961
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=167	0.003 [0.028] N=1,765	-0.003 [0.028] N=1,605	-0.013 [0.031] N=1,395	-0.005 [0.034] N=1,193	-0.030 [0.037] N=961
CIAF: any stunting, underweight, or wasting, 0-60 months	0.261 P=445	**0.071* [0.040] N=1,727**	0.055 [0.044] N=1,567	0.052 [0.045] N=1,360	0.015 [0.049] N=1,163	0.046 [0.053] N=935
BMI-for-age z score below -2	0.118 P=272	0.041 [0.031] N=2,323	0.032 [0.034] N=2,100	0.033 [0.036] N=1,841	0.013 [0.037] N=1,565	0.023 [0.044] N=1,275
WHO 2007 BMI overweight above +1, 61+ months	0.406 P=219	0.021 [0.081] N=552	-0.030 [0.083] N=489	0.013 [0.082] N=441	0.065 [0.089] N=369	-0.003 [0.098] N=304
WHO 2007 BMI obesity above +2, 61+ months	0.221 P=118	0.043 [0.080] N=552	-0.024 [0.083] N=489	-0.014 [0.089] N=441	0.022 [0.097] N=369	-0.039 [0.114] N=304
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=67	0.024 [0.055] N=552	0.040 [0.057] N=489	0.007 [0.058] N=441	0.011 [0.063] N=369	0.017 [0.069] N=304

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.010 [0.252] N=2,080	-0.041 [0.281] N=1,846	-0.023 [0.316] N=1,575	-0.298 [0.345] N=1,289	-0.248 [0.420] N=985
WHO weight-for-age z score	-0.047 [0.144] N=2,248	-0.031 [0.155] N=2,008	-0.178 [0.171] N=1,711	-0.261 [0.203] N=1,394	-0.277 [0.224] N=1,059
WHO weight-for-length/height z score, 0-60 months	-0.210 [0.245] N=1,531	-0.207 [0.256] N=1,371	-0.376 [0.302] N=1,161	-0.287 [0.330] N=959	**-0.818* [0.423] N=727**
WHO BMI-for-age z score	**-0.432** [0.212] N=1,998**	**-0.517** [0.219] N=1,775**	**-0.632** [0.258] N=1,516**	**-0.558* [0.289] N=1,240**	**-1.095*** [0.374] N=950**
Stunted: HAZ below -2	0.027 [0.045] N=2,080	0.028 [0.050] N=1,846	0.004 [0.055] N=1,575	0.009 [0.067] N=1,289	0.005 [0.090] N=985
Underweight: WAZ below -2	0.003 [0.021] N=2,248	-0.009 [0.024] N=2,008	0.002 [0.028] N=1,711	-0.024 [0.029] N=1,394	-0.038 [0.032] N=1,059
Wasted: WLZ/WHZ below -2, 0-60 months	**0.081** [0.037] N=1,531**	0.066 [0.042] N=1,371	**0.110** [0.047] N=1,161**	0.057 [0.048] N=959	**0.145*** [0.052] N=727**
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.054] N=1,531	0.001 [0.057] N=1,371	-0.020 [0.062] N=1,161	-0.024 [0.074] N=959	-0.078 [0.093] N=727
Obese: WLZ/WHZ above +3, 0-60 months	0.016 [0.040] N=1,531	0.001 [0.042] N=1,371	-0.019 [0.049] N=1,161	-0.011 [0.061] N=959	-0.109 [0.076] N=727
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.103* [0.061] N=1,500**	0.077 [0.069] N=1,340	0.080 [0.079] N=1,133	-0.002 [0.092] N=936	0.082 [0.108] N=708
BMI-for-age z score below -2	**0.098*** [0.030] N=1,998**	**0.092*** [0.035] N=1,775**	**0.114*** [0.040] N=1,516**	**0.100*** [0.037] N=1,240**	**0.188*** [0.046] N=950**
WHO 2007 BMI overweight above +1, 61+ months	-0.083 [0.126] N=464	-0.188 [0.135] N=401	-0.149 [0.143] N=353	-0.079 [0.183] N=281	-0.394 [0.261] N=216
WHO 2007 BMI obesity above +2, 61+ months	0.052 [0.119] N=464	-0.070 [0.125] N=401	-0.070 [0.144] N=353	-0.019 [0.162] N=281	-0.164 [0.250] N=216
WHO 2007 BMI thinness below -2, 61+ months	0.092 [0.069] N=464	**0.140* [0.076] N=401**	0.092 [0.080] N=353	0.130 [0.089] N=281	**0.245** [0.112] N=216**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	-0.247 [0.199] N=2,271	-0.286 [0.214] N=2,051	-0.314 [0.227] N=1,801	**-0.499** [0.240] N=1,528**	**-0.585** [0.265] N=1,246**
WHO weight-for-age z score	0.343	-0.072 [0.100] N=2,446	-0.060 [0.103] N=2,220	-0.144 [0.112] N=1,947	-0.197 [0.122] N=1,648	**-0.245* [0.127] N=1,339**
WHO weight-for-length/height z score, 0-60 months	0.389	-0.023 [0.196] N=1,645	0.013 [0.206] N=1,496	-0.066 [0.228] N=1,301	0.012 [0.243] N=1,112	-0.096 [0.282] N=900
WHO BMI-for-age z score	0.401	-0.068 [0.196] N=2,178	-0.066 [0.208] N=1,969	-0.098 [0.229] N=1,730	0.002 [0.244] N=1,468	-0.083 [0.280] N=1,200
Stunted: HAZ below -2	0.198 P=443	0.040 [0.035] N=2,271	0.043 [0.037] N=2,051	0.034 [0.039] N=1,801	0.045 [0.043] N=1,528	0.060 [0.049] N=1,246
Underweight: WAZ below -2	0.042 P=107	0.009 [0.014] N=2,446	0.003 [0.015] N=2,220	0.007 [0.016] N=1,947	0.004 [0.017] N=1,648	0.009 [0.018] N=1,339
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=178	0.027 [0.033] N=1,645	0.011 [0.036] N=1,496	0.025 [0.037] N=1,301	0.000 [0.038] N=1,112	0.018 [0.046] N=900
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=333	0.008 [0.038] N=1,645	0.008 [0.039] N=1,496	-0.003 [0.041] N=1,301	0.004 [0.044] N=1,112	0.002 [0.050] N=900
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=159	0.005 [0.027] N=1,645	0.001 [0.028] N=1,496	-0.007 [0.030] N=1,301	0.006 [0.034] N=1,112	-0.010 [0.036] N=900
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=420	**0.076* [0.044] N=1,609**	0.061 [0.048] N=1,460	0.059 [0.049] N=1,268	0.024 [0.052] N=1,083	0.076 [0.057] N=874
BMI-for-age z score below -2	0.116 P=249	0.039 [0.031] N=2,178	0.028 [0.033] N=1,969	0.033 [0.035] N=1,730	0.020 [0.034] N=1,468	0.039 [0.040] N=1,200
WHO 2007 BMI overweight above +1, 61+ months	0.411 P=213	-0.006 [0.085] N=531	-0.068 [0.085] N=471	-0.026 [0.087] N=427	0.020 [0.096] N=356	-0.067 [0.105] N=293
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=115	0.026 [0.084] N=531	-0.053 [0.087] N=471	-0.047 [0.094] N=427	-0.020 [0.102] N=356	-0.084 [0.118] N=293
WHO 2007 BMI thinness below -2, 61+ months	0.114 P=62	0.045 [0.058] N=531	0.062 [0.061] N=471	0.035 [0.064] N=427	0.033 [0.069] N=356	0.054 [0.077] N=293

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.058 [0.260] N=1,950	0.025 [0.293] N=1,730	0.017 [0.331] N=1,480	-0.239 [0.360] N=1,207	-0.288 [0.447] N=925
WHO weight-for-age z score	-0.051 [0.153] N=2,103	-0.023 [0.163] N=1,877	-0.217 [0.183] N=1,604	-0.291 [0.224] N=1,305	-0.352 [0.267] N=996
WHO weight-for-length/height z score, 0-60 months	-0.166 [0.266] N=1,423	-0.151 [0.281] N=1,274	-0.392 [0.336] N=1,079	-0.337 [0.366] N=890	**-0.933** [0.462] N=678**
WHO BMI-for-age z score	**-0.406* [0.228] N=1,871**	**-0.474** [0.238] N=1,662**	**-0.655** [0.289] N=1,423**	**-0.588* [0.331] N=1,161**	**-1.172*** [0.411] N=893**
Stunted: HAZ below -2	0.016 [0.049] N=1,950	0.020 [0.055] N=1,730	-0.001 [0.060] N=1,480	-0.001 [0.071] N=1,207	0.009 [0.097] N=925
Underweight: WAZ below -2	0.001 [0.023] N=2,103	-0.012 [0.025] N=1,877	-0.005 [0.028] N=1,604	-0.029 [0.033] N=1,305	-0.043 [0.037] N=996
Wasted: WLZ/WHZ below -2, 0-60 months	**0.076* [0.039] N=1,423**	0.057 [0.044] N=1,274	**0.102** [0.049] N=1,079**	0.050 [0.053] N=890	**0.138** [0.060] N=678**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.004 [0.056] N=1,423	-0.013 [0.059] N=1,274	-0.040 [0.065] N=1,079	-0.055 [0.079] N=890	-0.112 [0.101] N=678
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.043] N=1,423	-0.017 [0.044] N=1,274	-0.036 [0.051] N=1,079	-0.028 [0.064] N=890	-0.120 [0.080] N=678
CIAF: any stunting, underweight, or wasting, 0-60 months	0.089 [0.068] N=1,394	0.067 [0.077] N=1,245	0.071 [0.085] N=1,053	-0.033 [0.101] N=868	0.069 [0.127] N=659
BMI-for-age z score below -2	**0.090*** [0.033] N=1,871**	**0.082** [0.037] N=1,662**	**0.106** [0.043] N=1,423**	**0.094** [0.040] N=1,161**	**0.190*** [0.051] N=893**
WHO 2007 BMI overweight above +1, 61+ months	-0.076 [0.136] N=447	-0.193 [0.144] N=387	-0.138 [0.160] N=343	-0.038 [0.208] N=272	-0.377 [0.303] N=209
WHO 2007 BMI obesity above +2, 61+ months	0.067 [0.124] N=447	-0.064 [0.133] N=387	-0.057 [0.154] N=343	0.019 [0.173] N=272	-0.121 [0.262] N=209
WHO 2007 BMI thinness below -2, 61+ months	0.083 [0.073] N=447	**0.135* [0.079] N=387**	0.094 [0.084] N=343	0.115 [0.093] N=272	**0.214* [0.117] N=209**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.091	-0.249 [0.199] N=2,271	-0.289 [0.214] N=2,051	-0.315 [0.227] N=1,801	**-0.495** [0.241] N=1,528**	**-0.574** [0.264] N=1,246**
WHO weight-for-age z score	0.343	-0.072 [0.102] N=2,446	-0.061 [0.105] N=2,220	-0.149 [0.113] N=1,947	-0.198 [0.124] N=1,648	**-0.239* [0.132] N=1,339**
WHO weight-for-length/height z score, 0-60 months	0.389	-0.018 [0.197] N=1,645	0.016 [0.206] N=1,496	-0.058 [0.229] N=1,301	0.024 [0.245] N=1,112	-0.093 [0.292] N=900
WHO BMI-for-age z score	0.401	-0.064 [0.197] N=2,178	-0.063 [0.208] N=1,969	-0.095 [0.230] N=1,730	0.005 [0.246] N=1,468	-0.085 [0.282] N=1,200
Stunted: HAZ below -2	0.198 P=443	0.039 [0.035] N=2,271	0.042 [0.037] N=2,051	0.033 [0.039] N=1,801	0.044 [0.043] N=1,528	0.058 [0.049] N=1,246
Underweight: WAZ below -2	0.042 P=107	0.009 [0.015] N=2,446	0.003 [0.015] N=2,220	0.009 [0.016] N=1,947	0.004 [0.017] N=1,648	0.009 [0.018] N=1,339
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=178	0.024 [0.033] N=1,645	0.008 [0.035] N=1,496	0.021 [0.037] N=1,301	-0.005 [0.038] N=1,112	0.013 [0.046] N=900
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=333	0.007 [0.037] N=1,645	0.007 [0.039] N=1,496	-0.004 [0.041] N=1,301	0.003 [0.045] N=1,112	-0.001 [0.051] N=900
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=159	0.004 [0.027] N=1,645	-0.000 [0.027] N=1,496	-0.009 [0.030] N=1,301	0.004 [0.034] N=1,112	-0.013 [0.036] N=900
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=420	0.070 [0.044] N=1,609	0.054 [0.047] N=1,460	0.053 [0.049] N=1,268	0.017 [0.052] N=1,083	0.068 [0.057] N=874
BMI-for-age z score below -2	0.116 P=249	0.038 [0.031] N=2,178	0.027 [0.033] N=1,969	0.031 [0.035] N=1,730	0.018 [0.035] N=1,468	0.038 [0.040] N=1,200
WHO 2007 BMI overweight above +1, 61+ months	0.411 P=213	-0.006 [0.085] N=531	-0.068 [0.086] N=471	-0.028 [0.088] N=427	0.009 [0.096] N=356	-0.069 [0.105] N=293
WHO 2007 BMI obesity above +2, 61+ months	0.225 P=115	0.028 [0.083] N=531	-0.054 [0.087] N=471	-0.049 [0.093] N=427	-0.026 [0.101] N=356	-0.089 [0.117] N=293
WHO 2007 BMI thinness below -2, 61+ months	0.114 P=62	0.044 [0.058] N=531	0.062 [0.060] N=471	0.034 [0.062] N=427	0.032 [0.068] N=356	0.055 [0.074] N=293

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.041 [0.258] N=1,950	0.006 [0.288] N=1,730	0.003 [0.325] N=1,480	-0.285 [0.362] N=1,207	-0.339 [0.441] N=925
WHO weight-for-age z score	-0.050 [0.153] N=2,103	-0.024 [0.163] N=1,877	-0.226 [0.183] N=1,604	-0.299 [0.223] N=1,305	-0.386 [0.267] N=996
WHO weight-for-length/height z score, 0-60 months	-0.161 [0.265] N=1,423	-0.146 [0.280] N=1,274	-0.386 [0.335] N=1,079	-0.295 [0.373] N=890	**-0.932** [0.461] N=678**
WHO BMI-for-age z score	**-0.396* [0.228] N=1,871**	**-0.469* [0.238] N=1,662**	**-0.652** [0.288] N=1,423**	**-0.567* [0.332] N=1,161**	**-1.175*** [0.411] N=893**
Stunted: HAZ below -2	0.017 [0.048] N=1,950	0.022 [0.054] N=1,730	0.001 [0.059] N=1,480	0.006 [0.072] N=1,207	0.017 [0.096] N=925
Underweight: WAZ below -2	0.000 [0.022] N=2,103	-0.012 [0.024] N=1,877	-0.003 [0.028] N=1,604	-0.027 [0.032] N=1,305	-0.035 [0.037] N=996
Wasted: WLZ/WHZ below -2, 0-60 months	**0.074* [0.039] N=1,423**	0.055 [0.044] N=1,274	**0.100** [0.049] N=1,079**	0.045 [0.051] N=890	**0.139** [0.058] N=678**
Overweight: WLZ/WHZ above +2, 0-60 months	-0.002 [0.056] N=1,423	-0.011 [0.059] N=1,274	-0.039 [0.065] N=1,079	-0.048 [0.080] N=890	-0.112 [0.101] N=678
Obese: WLZ/WHZ above +3, 0-60 months	0.000 [0.043] N=1,423	-0.015 [0.044] N=1,274	-0.035 [0.051] N=1,079	-0.023 [0.065] N=890	-0.119 [0.080] N=678
CIAF: any stunting, underweight, or wasting, 0-60 months	0.087 [0.067] N=1,394	0.066 [0.076] N=1,245	0.072 [0.086] N=1,053	-0.025 [0.101] N=868	0.083 [0.128] N=659
BMI-for-age z score below -2	**0.090*** [0.033] N=1,871**	**0.083** [0.037] N=1,662**	**0.107** [0.043] N=1,423**	**0.094** [0.040] N=1,161**	**0.195*** [0.049] N=893**
WHO 2007 BMI overweight above +1, 61+ months	-0.069 [0.137] N=447	-0.181 [0.146] N=387	-0.133 [0.162] N=343	-0.042 [0.211] N=272	-0.363 [0.313] N=209
WHO 2007 BMI obesity above +2, 61+ months	0.074 [0.122] N=447	-0.060 [0.131] N=387	-0.056 [0.152] N=343	0.012 [0.172] N=272	-0.140 [0.261] N=209
WHO 2007 BMI thinness below -2, 61+ months	0.087 [0.072] N=447	**0.138* [0.077] N=387**	0.092 [0.082] N=343	0.113 [0.090] N=272	**0.215* [0.110] N=209**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 71 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.