Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-5 Using 16 December 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 71 months on 16 December 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.518	0.106 [0.163] N=2,458	0.099 [0.169] N=2,285	0.047 [0.185] N=2,059	0.101 [0.194] N=1,765	0.035 [0.220] N=1,457
WHO weight-for-age z score	0.005	-0.013 [0.103] N=2,660	0.025 [0.109] N=2,466	0.046 [0.115] N=2,223	0.019 [0.126] N=1,906	-0.023 [0.133] N=1,573
WHO weight-for-length/height z score, 0-60 months	0.338	0.099 [0.173] N=1,839	0.069 [0.177] N=1,707	-0.036 [0.192] N=1,539	-0.123 [0.217] N=1,319	-0.042 [0.234] N=1,078
WHO BMI-for-age z score	0.346	0.010 [0.175] N=2,355	0.020 [0.181] N=2,192	-0.000 [0.199] N=1,967	-0.135 [0.227] N=1,684	-0.042 [0.255] N=1,390
Stunted: HAZ below -2	0.248 P=600	-0.032 [0.033] N=2,458	-0.035 [0.035] N=2,285	-0.022 [0.037] N=2,059	-0.044 [0.038] N=1,765	-0.015 [0.042] N=1,457
Underweight: WAZ below -2	0.064 P=166	0.006 [0.017] N=2,660	0.003 [0.018] N=2,466	-0.007 [0.020] N=2,223	-0.001 [0.021] N=1,906	0.004 [0.024] N=1,573
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=208	-0.023 [0.028] N=1,839	-0.013 [0.029] N=1,707	0.004 [0.031] N=1,539	0.008 [0.036] N=1,319	-0.011 [0.037] N=1,078
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=345	0.038 [0.039] N=1,839	0.034 [0.041] N=1,707	0.019 [0.044] N=1,539	-0.002 [0.048] N=1,319	-0.010 [0.053] N=1,078
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=153	0.011 [0.028] N=1,839	0.009 [0.029] N=1,707	0.017 [0.032] N=1,539	0.013 [0.035] N=1,319	0.006 [0.041] N=1,078
CIAF: any stunting, underweight, or wasting, 0-60 months	0.330 P=588	-0.047 [0.041] N=1,800	-0.053 [0.044] N=1,672	-0.027 [0.046] N=1,508	-0.047 [0.048] N=1,291	-0.026 [0.051] N=1,057
BMI-for-age z score below -2	0.121 P=289	-0.005 [0.029] N=2,355	-0.003 [0.029] N=2,192	0.003 [0.032] N=1,967	0.016 [0.035] N=1,684	0.005 [0.038] N=1,390
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=191	-0.049 [0.071] N=509	-0.060 [0.071] N=479	0.002 [0.076] N=427	-0.021 [0.082] N=368	-0.030 [0.097] N=310
WHO 2007 BMI obesity above +2, 61+ months	0.215 P=111	-0.038 [0.060] N=509	-0.048 [0.061] N=479	-0.047 [0.067] N=427	-0.086 [0.074] N=368	-0.103 [0.088] N=310
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=63	0.020 [0.058] N=509	0.002 [0.057] N=479	-0.068 [0.061] N=427	-0.081 [0.069] N=368	-0.110 [0.081] N=310

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.059 [0.238] N=2,073	0.017 [0.257] N=1,900	-0.112 [0.287] N=1,674	0.026 [0.309] N=1,380	-0.181 [0.409] N=1,072
WHO weight-for-age z score	-0.091 [0.142] N=2,247	-0.032 [0.156] N=2,053	-0.009 [0.172] N=1,810	-0.076 [0.210] N=1,493	-0.224 [0.257] N=1,160
WHO weight-for-length/height z score, 0-60 months	0.144 [0.239] N=1,556	0.109 [0.251] N=1,424	-0.108 [0.284] N=1,256	-0.353 [0.352] N=1,036	-0.310 [0.436] N=795
WHO BMI-for-age z score	0.163 [0.224] N=1,989	0.224 [0.231] N=1,826	0.232 [0.264] N=1,601	-0.043 [0.331] N=1,318	0.264 [0.408] N=1,024
Stunted: HAZ below -2	-0.015 [0.048] N=2,073	-0.016 [0.052] N=1,900	0.011 [0.059] N=1,674	-0.042 [0.065] N=1,380	0.036 [0.085] N=1,072
Underweight: WAZ below -2	0.020 [0.023] N=2,247	0.014 [0.025] N=2,053	-0.000 [0.029] N=1,810	0.017 [0.034] N=1,493	0.060 [0.047] N=1,160
Wasted: WLZ/WHZ below -2, 0-60 months	-0.016 [0.036] N=1,556	-0.000 [0.036] N=1,424	0.036 [0.041] N=1,256	0.052 [0.053] N=1,036	0.020 [0.053] N=795
Overweight: WLZ/WHZ above +2, 0-60 months	0.062 [0.050] N=1,556	0.061 [0.054] N=1,424	0.034 [0.059] N=1,256	-0.002 [0.072] N=1,036	-0.036 [0.089] N=795
Obese: WLZ/WHZ above +3, 0-60 months	0.031 [0.033] N=1,556	0.027 [0.035] N=1,424	0.044 [0.040] N=1,256	0.046 [0.050] N=1,036	0.042 [0.073] N=795
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.011 [0.055] N=1,524	-0.019 [0.061] N=1,396	0.028 [0.069] N=1,232	-0.024 [0.081] N=1,015	0.035 [0.097] N=781
BMI-for-age z score below -2	-0.019 [0.035] N=1,989	-0.022 [0.034] N=1,826	-0.014 [0.039] N=1,601	0.011 [0.046] N=1,318	-0.025 [0.058] N=1,024
WHO 2007 BMI overweight above +1, 61+ months	-0.039 [0.114] N=425	-0.047 [0.118] N=395	0.070 [0.133] N=343	0.059 [0.157] N=284	0.101 [0.216] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.013 [0.084] N=425	0.004 [0.087] N=395	0.024 [0.098] N=343	-0.031 [0.117] N=284	-0.029 [0.153] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.062 [0.075] N=425	0.028 [0.071] N=395	-0.091 [0.076] N=343	-0.140 [0.092] N=284	**-0.243** [0.116] N=226**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.518	0.110 [0.163] N=2,458	0.103 [0.169] N=2,285	0.054 [0.185] N=2,059	0.109 [0.194] N=1,765	0.044 [0.219] N=1,457
WHO weight-for-age z score	0.005	-0.010 [0.103] N=2,660	0.027 [0.110] N=2,466	0.050 [0.116] N=2,223	0.025 [0.127] N=1,906	-0.021 [0.134] N=1,573
WHO weight-for-length/height z score, 0-60 months	0.338	0.094 [0.174] N=1,839	0.062 [0.179] N=1,707	-0.045 [0.194] N=1,539	-0.130 [0.219] N=1,319	-0.054 [0.237] N=1,078
WHO BMI-for-age z score	0.346	0.009 [0.175] N=2,355	0.018 [0.181] N=2,192	-0.004 [0.200] N=1,967	-0.137 [0.228] N=1,684	-0.045 [0.254] N=1,390
Stunted: HAZ below -2	0.248 P=600	-0.033 [0.033] N=2,458	-0.036 [0.035] N=2,285	-0.023 [0.037] N=2,059	-0.045 [0.038] N=1,765	-0.016 [0.042] N=1,457
Underweight: WAZ below -2	0.064 P=166	0.007 [0.017] N=2,660	0.003 [0.018] N=2,466	-0.006 [0.020] N=2,223	-0.001 [0.021] N=1,906	0.004 [0.024] N=1,573
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=208	-0.022 [0.029] N=1,839	-0.012 [0.029] N=1,707	0.007 [0.031] N=1,539	0.010 [0.036] N=1,319	-0.008 [0.037] N=1,078
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=345	0.038 [0.039] N=1,839	0.035 [0.041] N=1,707	0.020 [0.043] N=1,539	-0.001 [0.048] N=1,319	-0.009 [0.052] N=1,078
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=153	0.011 [0.028] N=1,839	0.009 [0.029] N=1,707	0.017 [0.031] N=1,539	0.014 [0.035] N=1,319	0.007 [0.040] N=1,078
CIAF: any stunting, underweight, or wasting, 0-60 months	0.330 P=588	-0.046 [0.041] N=1,800	-0.051 [0.043] N=1,672	-0.025 [0.045] N=1,508	-0.046 [0.047] N=1,291	-0.025 [0.050] N=1,057
BMI-for-age z score below -2	0.121 P=289	-0.005 [0.029] N=2,355	-0.003 [0.029] N=2,192	0.004 [0.032] N=1,967	0.017 [0.035] N=1,684	0.006 [0.038] N=1,390
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=191	-0.049 [0.071] N=509	-0.061 [0.071] N=479	0.002 [0.076] N=427	-0.025 [0.082] N=368	-0.035 [0.097] N=310
WHO 2007 BMI obesity above +2, 61+ months	0.215 P=111	-0.038 [0.060] N=509	-0.050 [0.061] N=479	-0.047 [0.068] N=427	-0.090 [0.075] N=368	-0.109 [0.089] N=310
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=63	0.021 [0.058] N=509	0.003 [0.057] N=479	-0.067 [0.061] N=427	-0.077 [0.069] N=368	-0.107 [0.081] N=310

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.058 [0.238] N=2,073	0.016 [0.256] N=1,900	-0.104 [0.286] N=1,674	0.032 [0.306] N=1,380	-0.189 [0.408] N=1,072
WHO weight-for-age z score	-0.091 [0.142] N=2,247	-0.033 [0.155] N=2,053	-0.007 [0.171] N=1,810	-0.074 [0.207] N=1,493	-0.228 [0.255] N=1,160
WHO weight-for-length/height z score, 0-60 months	0.141 [0.240] N=1,556	0.099 [0.253] N=1,424	-0.124 [0.285] N=1,256	-0.375 [0.354] N=1,036	-0.346 [0.439] N=795
WHO BMI-for-age z score	0.164 [0.224] N=1,989	0.223 [0.232] N=1,826	0.228 [0.264] N=1,601	-0.044 [0.331] N=1,318	0.269 [0.407] N=1,024
Stunted: HAZ below -2	-0.015 [0.048] N=2,073	-0.016 [0.052] N=1,900	0.010 [0.059] N=1,674	-0.043 [0.065] N=1,380	0.037 [0.085] N=1,072
Underweight: WAZ below -2	0.020 [0.023] N=2,247	0.014 [0.025] N=2,053	-0.000 [0.029] N=1,810	0.017 [0.034] N=1,493	0.060 [0.047] N=1,160
Wasted: WLZ/WHZ below -2, 0-60 months	-0.015 [0.036] N=1,556	0.001 [0.037] N=1,424	0.039 [0.041] N=1,256	0.055 [0.053] N=1,036	0.026 [0.054] N=795
Overweight: WLZ/WHZ above +2, 0-60 months	0.062 [0.050] N=1,556	0.061 [0.054] N=1,424	0.034 [0.059] N=1,256	-0.001 [0.072] N=1,036	-0.034 [0.089] N=795
Obese: WLZ/WHZ above +3, 0-60 months	0.031 [0.033] N=1,556	0.027 [0.035] N=1,424	0.044 [0.040] N=1,256	0.046 [0.050] N=1,036	0.042 [0.073] N=795
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.011 [0.055] N=1,524	-0.018 [0.061] N=1,396	0.030 [0.069] N=1,232	-0.023 [0.081] N=1,015	0.035 [0.097] N=781
BMI-for-age z score below -2	-0.019 [0.035] N=1,989	-0.021 [0.034] N=1,826	-0.014 [0.039] N=1,601	0.011 [0.046] N=1,318	-0.026 [0.058] N=1,024
WHO 2007 BMI overweight above +1, 61+ months	-0.039 [0.114] N=425	-0.047 [0.119] N=395	0.069 [0.134] N=343	0.052 [0.161] N=284	0.096 [0.222] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.010 [0.084] N=425	-0.001 [0.087] N=395	0.018 [0.098] N=343	-0.048 [0.117] N=284	-0.054 [0.153] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.066 [0.076] N=425	0.033 [0.071] N=395	-0.087 [0.076] N=343	-0.122 [0.090] N=284	**-0.231** [0.114] N=226**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.518	0.073 [0.163] N=2,458	0.065 [0.170] N=2,285	0.020 [0.187] N=2,059	0.070 [0.195] N=1,765	-0.006 [0.222] N=1,457
WHO weight-for-age z score	0.005	-0.050 [0.101] N=2,660	-0.011 [0.108] N=2,466	0.012 [0.114] N=2,223	-0.012 [0.125] N=1,906	-0.063 [0.132] N=1,573
WHO weight-for-length/height z score, 0-60 months	0.338	0.068 [0.174] N=1,839	0.041 [0.181] N=1,707	-0.061 [0.194] N=1,539	-0.143 [0.219] N=1,319	-0.066 [0.240] N=1,078
WHO BMI-for-age z score	0.346	0.007 [0.175] N=2,355	0.017 [0.182] N=2,192	-0.002 [0.200] N=1,967	-0.132 [0.226] N=1,684	-0.033 [0.256] N=1,390
Stunted: HAZ below -2	0.248 P=600	-0.023 [0.032] N=2,458	-0.024 [0.033] N=2,285	-0.012 [0.036] N=2,059	-0.034 [0.037] N=1,765	-0.001 [0.041] N=1,457
Underweight: WAZ below -2	0.064 P=166	0.013 [0.017] N=2,660	0.010 [0.018] N=2,466	0.001 [0.019] N=2,223	0.006 [0.020] N=1,906	0.014 [0.023] N=1,573
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=208	-0.014 [0.028] N=1,839	-0.003 [0.029] N=1,707	0.013 [0.031] N=1,539	0.015 [0.035] N=1,319	-0.002 [0.037] N=1,078
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=345	0.038 [0.039] N=1,839	0.036 [0.041] N=1,707	0.021 [0.044] N=1,539	-0.001 [0.048] N=1,319	-0.009 [0.054] N=1,078
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=153	0.012 [0.028] N=1,839	0.011 [0.029] N=1,707	0.018 [0.031] N=1,539	0.013 [0.034] N=1,319	0.005 [0.040] N=1,078
CIAF: any stunting, underweight, or wasting, 0-60 months	0.330 P=588	-0.025 [0.039] N=1,800	-0.029 [0.041] N=1,672	-0.010 [0.043] N=1,508	-0.034 [0.044] N=1,291	-0.005 [0.048] N=1,057
BMI-for-age z score below -2	0.121 P=289	0.001 [0.029] N=2,355	0.004 [0.029] N=2,192	0.010 [0.032] N=1,967	0.023 [0.035] N=1,684	0.013 [0.038] N=1,390
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=191	-0.046 [0.070] N=509	-0.057 [0.070] N=479	0.012 [0.073] N=427	-0.004 [0.080] N=368	-0.029 [0.096] N=310
WHO 2007 BMI obesity above +2, 61+ months	0.215 P=111	-0.032 [0.059] N=509	-0.043 [0.060] N=479	-0.034 [0.065] N=427	-0.069 [0.070] N=368	-0.096 [0.085] N=310
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=63	0.023 [0.058] N=509	0.005 [0.057] N=479	-0.064 [0.061] N=427	-0.068 [0.070] N=368	-0.091 [0.081] N=310

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.036 [0.235] N=2,073	-0.010 [0.253] N=1,900	-0.126 [0.284] N=1,674	-0.013 [0.309] N=1,380	-0.228 [0.409] N=1,072
WHO weight-for-age z score	-0.119 [0.138] N=2,247	-0.057 [0.152] N=2,053	-0.029 [0.167] N=1,810	-0.089 [0.201] N=1,493	-0.249 [0.249] N=1,160
WHO weight-for-length/height z score, 0-60 months	0.123 [0.240] N=1,556	0.089 [0.256] N=1,424	-0.128 [0.283] N=1,256	-0.387 [0.354] N=1,036	-0.373 [0.442] N=795
WHO BMI-for-age z score	0.165 [0.226] N=1,989	0.225 [0.233] N=1,826	0.234 [0.264] N=1,601	-0.028 [0.333] N=1,318	0.296 [0.410] N=1,024
Stunted: HAZ below -2	-0.009 [0.047] N=2,073	-0.008 [0.051] N=1,900	0.016 [0.058] N=1,674	-0.034 [0.065] N=1,380	0.047 [0.084] N=1,072
Underweight: WAZ below -2	0.024 [0.023] N=2,247	0.019 [0.025] N=2,053	0.003 [0.028] N=1,810	0.018 [0.033] N=1,493	0.063 [0.046] N=1,160
Wasted: WLZ/WHZ below -2, 0-60 months	-0.007 [0.036] N=1,556	0.010 [0.037] N=1,424	0.043 [0.041] N=1,256	0.061 [0.054] N=1,036	0.032 [0.056] N=795
Overweight: WLZ/WHZ above +2, 0-60 months	0.065 [0.051] N=1,556	0.066 [0.055] N=1,424	0.037 [0.060] N=1,256	-0.002 [0.073] N=1,036	-0.031 [0.089] N=795
Obese: WLZ/WHZ above +3, 0-60 months	0.032 [0.033] N=1,556	0.030 [0.036] N=1,424	0.045 [0.040] N=1,256	0.043 [0.050] N=1,036	0.032 [0.073] N=795
CIAF: any stunting, underweight, or wasting, 0-60 months	0.010 [0.054] N=1,524	0.005 [0.060] N=1,396	0.039 [0.067] N=1,232	-0.020 [0.079] N=1,015	0.041 [0.095] N=781
BMI-for-age z score below -2	-0.014 [0.035] N=1,989	-0.015 [0.034] N=1,826	-0.011 [0.040] N=1,601	0.014 [0.046] N=1,318	-0.023 [0.058] N=1,024
WHO 2007 BMI overweight above +1, 61+ months	-0.043 [0.113] N=425	-0.057 [0.118] N=395	0.073 [0.131] N=343	0.093 [0.153] N=284	0.096 [0.215] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.006 [0.085] N=425	-0.008 [0.089] N=395	0.023 [0.097] N=343	-0.022 [0.109] N=284	-0.052 [0.149] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.064 [0.075] N=425	0.027 [0.069] N=395	-0.097 [0.076] N=343	-0.121 [0.087] N=284	**-0.204* [0.110] N=226**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.518	0.060 [0.163] N=2,458	0.059 [0.171] N=2,285	0.010 [0.188] N=2,059	0.054 [0.197] N=1,765	-0.028 [0.225] N=1,457
WHO weight-for-age z score	0.005	-0.043 [0.101] N=2,660	-0.008 [0.108] N=2,466	0.013 [0.116] N=2,223	-0.014 [0.127] N=1,906	-0.082 [0.134] N=1,573
WHO weight-for-length/height z score, 0-60 months	0.338	0.104 [0.175] N=1,839	0.075 [0.181] N=1,707	-0.033 [0.196] N=1,539	-0.122 [0.223] N=1,319	-0.047 [0.241] N=1,078
WHO BMI-for-age z score	0.346	0.026 [0.178] N=2,355	0.034 [0.183] N=2,192	0.011 [0.201] N=1,967	-0.118 [0.230] N=1,684	-0.026 [0.259] N=1,390
Stunted: HAZ below -2	0.248 P=600	-0.020 [0.033] N=2,458	-0.023 [0.034] N=2,285	-0.010 [0.037] N=2,059	-0.028 [0.037] N=1,765	0.001 [0.041] N=1,457
Underweight: WAZ below -2	0.064 P=166	0.014 [0.017] N=2,660	0.011 [0.018] N=2,466	0.003 [0.019] N=2,223	0.008 [0.021] N=1,906	0.017 [0.024] N=1,573
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=208	-0.015 [0.028] N=1,839	-0.004 [0.029] N=1,707	0.013 [0.031] N=1,539	0.017 [0.035] N=1,319	-0.001 [0.037] N=1,078
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=345	0.042 [0.039] N=1,839	0.039 [0.042] N=1,707	0.024 [0.044] N=1,539	0.004 [0.049] N=1,319	-0.009 [0.053] N=1,078
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=153	0.019 [0.028] N=1,839	0.017 [0.030] N=1,707	0.023 [0.032] N=1,539	0.020 [0.036] N=1,319	0.014 [0.042] N=1,078
CIAF: any stunting, underweight, or wasting, 0-60 months	0.330 P=588	-0.022 [0.039] N=1,800	-0.026 [0.042] N=1,672	-0.003 [0.044] N=1,508	-0.020 [0.045] N=1,291	0.004 [0.049] N=1,057
BMI-for-age z score below -2	0.121 P=289	-0.001 [0.029] N=2,355	0.001 [0.029] N=2,192	0.008 [0.032] N=1,967	0.022 [0.035] N=1,684	0.011 [0.039] N=1,390
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=191	-0.048 [0.071] N=509	-0.061 [0.071] N=479	0.006 [0.076] N=427	-0.012 [0.080] N=368	-0.023 [0.096] N=310
WHO 2007 BMI obesity above +2, 61+ months	0.215 P=111	-0.035 [0.060] N=509	-0.046 [0.061] N=479	-0.040 [0.067] N=427	-0.079 [0.075] N=368	-0.100 [0.091] N=310
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=63	0.023 [0.058] N=509	0.010 [0.057] N=479	-0.065 [0.060] N=427	-0.070 [0.069] N=368	-0.101 [0.080] N=310

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.030 [0.235] N=2,073	-0.007 [0.253] N=1,900	-0.120 [0.282] N=1,674	0.005 [0.306] N=1,380	-0.252 [0.406] N=1,072
WHO weight-for-age z score	-0.109 [0.141] N=2,247	-0.053 [0.154] N=2,053	-0.029 [0.171] N=1,810	-0.089 [0.206] N=1,493	-0.241 [0.252] N=1,160
WHO weight-for-length/height z score, 0-60 months	0.146 [0.241] N=1,556	0.118 [0.253] N=1,424	-0.115 [0.285] N=1,256	-0.386 [0.355] N=1,036	-0.372 [0.444] N=795
WHO BMI-for-age z score	0.167 [0.226] N=1,989	0.227 [0.232] N=1,826	0.232 [0.265] N=1,601	-0.043 [0.333] N=1,318	0.279 [0.406] N=1,024
Stunted: HAZ below -2	-0.008 [0.048] N=2,073	-0.010 [0.052] N=1,900	0.014 [0.058] N=1,674	-0.037 [0.065] N=1,380	0.049 [0.085] N=1,072
Underweight: WAZ below -2	0.023 [0.023] N=2,247	0.019 [0.025] N=2,053	0.005 [0.028] N=1,810	0.019 [0.033] N=1,493	0.063 [0.047] N=1,160
Wasted: WLZ/WHZ below -2, 0-60 months	-0.008 [0.036] N=1,556	0.007 [0.038] N=1,424	0.042 [0.042] N=1,256	0.060 [0.054] N=1,036	0.031 [0.055] N=795
Overweight: WLZ/WHZ above +2, 0-60 months	0.068 [0.051] N=1,556	0.068 [0.055] N=1,424	0.038 [0.060] N=1,256	-0.001 [0.073] N=1,036	-0.033 [0.089] N=795
Obese: WLZ/WHZ above +3, 0-60 months	0.036 [0.033] N=1,556	0.034 [0.035] N=1,424	0.047 [0.040] N=1,256	0.044 [0.049] N=1,036	0.032 [0.071] N=795
CIAF: any stunting, underweight, or wasting, 0-60 months	0.010 [0.054] N=1,524	0.000 [0.060] N=1,396	0.038 [0.067] N=1,232	-0.016 [0.078] N=1,015	0.039 [0.096] N=781
BMI-for-age z score below -2	-0.013 [0.035] N=1,989	-0.017 [0.034] N=1,826	-0.011 [0.040] N=1,601	0.018 [0.046] N=1,318	-0.020 [0.058] N=1,024
WHO 2007 BMI overweight above +1, 61+ months	-0.036 [0.114] N=425	-0.048 [0.118] N=395	0.074 [0.132] N=343	0.075 [0.155] N=284	0.132 [0.212] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.016 [0.082] N=425	0.003 [0.085] N=395	0.023 [0.093] N=343	-0.035 [0.109] N=284	-0.033 [0.148] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.078 [0.075] N=425	0.052 [0.072] N=395	-0.079 [0.076] N=343	-0.104 [0.089] N=284	**-0.215* [0.116] N=226**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.524	0.073 [0.172] N=2,296	0.058 [0.180] N=2,138	0.009 [0.193] N=1,921	0.056 [0.200] N=1,655	-0.034 [0.225] N=1,363
WHO weight-for-age z score	0.005	0.004 [0.100] N=2,478	0.035 [0.104] N=2,302	0.059 [0.114] N=2,070	0.029 [0.126] N=1,783	0.002 [0.135] N=1,471
WHO weight-for-length/height z score, 0-60 months	0.354	0.195 [0.172] N=1,709	0.164 [0.177] N=1,592	0.037 [0.191] N=1,432	-0.034 [0.218] N=1,236	0.100 [0.237] N=1,008
WHO BMI-for-age z score	0.365	0.064 [0.173] N=2,198	0.066 [0.176] N=2,050	0.055 [0.196] N=1,836	-0.044 [0.225] N=1,580	0.100 [0.254] N=1,302
Stunted: HAZ below -2	0.250 P=563	-0.022 [0.036] N=2,296	-0.024 [0.038] N=2,138	-0.012 [0.041] N=1,921	-0.034 [0.042] N=1,655	-0.002 [0.048] N=1,363
Underweight: WAZ below -2	0.066 P=159	0.004 [0.018] N=2,478	0.001 [0.019] N=2,302	-0.008 [0.021] N=2,070	-0.002 [0.023] N=1,783	0.005 [0.026] N=1,471
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	-0.029 [0.030] N=1,709	-0.019 [0.030] N=1,592	0.006 [0.033] N=1,432	0.008 [0.038] N=1,236	-0.013 [0.040] N=1,008
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=323	0.045 [0.040] N=1,709	0.042 [0.042] N=1,592	0.025 [0.045] N=1,432	0.002 [0.050] N=1,236	-0.000 [0.056] N=1,008
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=146	0.018 [0.028] N=1,709	0.018 [0.030] N=1,592	0.026 [0.032] N=1,432	0.023 [0.036] N=1,236	0.018 [0.042] N=1,008
CIAF: any stunting, underweight, or wasting, 0-60 months	0.331 P=547	-0.039 [0.047] N=1,673	-0.046 [0.050] N=1,559	-0.019 [0.052] N=1,402	-0.042 [0.056] N=1,209	-0.019 [0.062] N=987
BMI-for-age z score below -2	0.118 P=263	-0.009 [0.030] N=2,198	-0.007 [0.029] N=2,050	-0.000 [0.032] N=1,836	0.007 [0.037] N=1,580	-0.012 [0.040] N=1,302
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=181	-0.057 [0.074] N=484	-0.064 [0.075] N=455	0.003 [0.079] N=405	-0.028 [0.081] N=349	-0.016 [0.101] N=294
WHO 2007 BMI obesity above +2, 61+ months	0.212 P=104	-0.051 [0.064] N=484	-0.058 [0.065] N=455	-0.053 [0.072] N=405	-0.101 [0.077] N=349	-0.122 [0.093] N=294
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=59	0.029 [0.056] N=484	0.012 [0.055] N=455	-0.063 [0.056] N=405	-0.081 [0.063] N=349	-0.118 [0.074] N=294

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.099 [0.254] N=1,937	0.058 [0.276] N=1,779	-0.057 [0.305] N=1,562	0.102 [0.334] N=1,296	-0.137 [0.453] N=1,004
WHO weight-for-age z score	-0.091 [0.138] N=2,093	-0.047 [0.145] N=1,917	-0.033 [0.168] N=1,685	-0.129 [0.206] N=1,398	-0.262 [0.255] N=1,086
WHO weight-for-length/height z score, 0-60 months	0.172 [0.241] N=1,446	0.124 [0.251] N=1,329	-0.135 [0.280] N=1,169	-0.370 [0.357] N=973	-0.250 [0.453] N=745
WHO BMI-for-age z score	0.121 [0.226] N=1,857	0.158 [0.229] N=1,709	0.166 [0.263] N=1,495	-0.078 [0.335] N=1,239	0.329 [0.426] N=961
Stunted: HAZ below -2	-0.012 [0.050] N=1,937	-0.015 [0.054] N=1,779	0.009 [0.062] N=1,562	-0.049 [0.069] N=1,296	0.031 [0.091] N=1,004
Underweight: WAZ below -2	0.018 [0.025] N=2,093	0.015 [0.027] N=1,917	0.002 [0.031] N=1,685	0.023 [0.037] N=1,398	0.069 [0.051] N=1,086
Wasted: WLZ/WHZ below -2, 0-60 months	-0.024 [0.036] N=1,446	-0.009 [0.036] N=1,329	0.039 [0.041] N=1,169	0.048 [0.053] N=973	0.006 [0.054] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.065 [0.050] N=1,446	0.064 [0.054] N=1,329	0.037 [0.059] N=1,169	-0.005 [0.072] N=973	-0.019 [0.092] N=745
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.034] N=1,446	0.029 [0.037] N=1,329	0.046 [0.042] N=1,169	0.047 [0.052] N=973	0.048 [0.079] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.008 [0.058] N=1,417	-0.021 [0.064] N=1,303	0.030 [0.073] N=1,146	-0.034 [0.086] N=953	0.019 [0.102] N=731
BMI-for-age z score below -2	-0.012 [0.035] N=1,857	-0.015 [0.033] N=1,709	-0.005 [0.039] N=1,495	0.010 [0.046] N=1,239	-0.048 [0.060] N=961
WHO 2007 BMI overweight above +1, 61+ months	-0.072 [0.120] N=403	-0.079 [0.126] N=374	0.045 [0.141] N=324	0.021 [0.165] N=268	0.109 [0.237] N=213
WHO 2007 BMI obesity above +2, 61+ months	-0.022 [0.089] N=403	-0.028 [0.093] N=374	-0.018 [0.103] N=324	-0.098 [0.121] N=268	-0.112 [0.163] N=213
WHO 2007 BMI thinness below -2, 61+ months	0.097 [0.080] N=403	0.064 [0.075] N=374	-0.065 [0.080] N=324	-0.124 [0.098] N=268	**-0.245* [0.127] N=213**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.524	0.084 [0.170] N=2,296	0.068 [0.178] N=2,138	0.006 [0.192] N=1,921	0.053 [0.199] N=1,655	-0.031 [0.224] N=1,363
WHO weight-for-age z score	0.005	0.017 [0.101] N=2,478	0.046 [0.106] N=2,302	0.066 [0.114] N=2,070	0.038 [0.127] N=1,783	0.008 [0.136] N=1,471
WHO weight-for-length/height z score, 0-60 months	0.354	0.197 [0.173] N=1,709	0.168 [0.178] N=1,592	0.039 [0.191] N=1,432	-0.032 [0.217] N=1,236	0.091 [0.237] N=1,008
WHO BMI-for-age z score	0.365	0.072 [0.172] N=2,198	0.072 [0.176] N=2,050	0.058 [0.196] N=1,836	-0.042 [0.225] N=1,580	0.097 [0.254] N=1,302
Stunted: HAZ below -2	0.250 P=563	-0.025 [0.036] N=2,296	-0.027 [0.038] N=2,138	-0.014 [0.040] N=1,921	-0.036 [0.041] N=1,655	-0.006 [0.047] N=1,363
Underweight: WAZ below -2	0.066 P=159	0.001 [0.018] N=2,478	-0.002 [0.019] N=2,302	-0.011 [0.021] N=2,070	-0.005 [0.022] N=1,783	0.002 [0.026] N=1,471
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	-0.028 [0.030] N=1,709	-0.018 [0.030] N=1,592	0.006 [0.033] N=1,432	0.008 [0.038] N=1,236	-0.016 [0.039] N=1,008
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=323	0.045 [0.040] N=1,709	0.042 [0.043] N=1,592	0.026 [0.046] N=1,432	0.002 [0.051] N=1,236	-0.000 [0.056] N=1,008
Obese: WLZ/WHZ above +3, 0-60 months	0.086 P=146	0.018 [0.028] N=1,709	0.017 [0.030] N=1,592	0.027 [0.032] N=1,432	0.023 [0.036] N=1,236	0.019 [0.041] N=1,008
CIAF: any stunting, underweight, or wasting, 0-60 months	0.331 P=547	-0.038 [0.047] N=1,673	-0.045 [0.050] N=1,559	-0.018 [0.052] N=1,402	-0.042 [0.055] N=1,209	-0.026 [0.060] N=987
BMI-for-age z score below -2	0.118 P=263	-0.009 [0.029] N=2,198	-0.008 [0.029] N=2,050	-0.000 [0.032] N=1,836	0.007 [0.037] N=1,580	-0.013 [0.040] N=1,302
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=181	-0.059 [0.071] N=484	-0.067 [0.072] N=455	-0.002 [0.077] N=405	-0.026 [0.081] N=349	-0.027 [0.099] N=294
WHO 2007 BMI obesity above +2, 61+ months	0.212 P=104	-0.052 [0.063] N=484	-0.062 [0.064] N=455	-0.052 [0.072] N=405	-0.097 [0.077] N=349	-0.121 [0.093] N=294
WHO 2007 BMI thinness below -2, 61+ months	0.121 P=59	0.025 [0.056] N=484	0.008 [0.054] N=455	-0.066 [0.056] N=405	-0.085 [0.064] N=349	-0.120 [0.073] N=294

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.115 [0.250] N=1,937	0.075 [0.272] N=1,779	-0.058 [0.303] N=1,562	0.100 [0.331] N=1,296	-0.111 [0.451] N=1,004
WHO weight-for-age z score	-0.071 [0.137] N=2,093	-0.028 [0.145] N=1,917	-0.016 [0.166] N=1,685	-0.104 [0.204] N=1,398	-0.239 [0.254] N=1,086
WHO weight-for-length/height z score, 0-60 months	0.191 [0.242] N=1,446	0.151 [0.254] N=1,329	-0.128 [0.283] N=1,169	-0.358 [0.359] N=973	-0.250 [0.461] N=745
WHO BMI-for-age z score	0.143 [0.226] N=1,857	0.179 [0.230] N=1,709	0.171 [0.263] N=1,495	-0.074 [0.335] N=1,239	0.327 [0.431] N=961
Stunted: HAZ below -2	-0.015 [0.050] N=1,937	-0.017 [0.054] N=1,779	0.006 [0.061] N=1,562	-0.053 [0.068] N=1,296	0.022 [0.091] N=1,004
Underweight: WAZ below -2	0.015 [0.025] N=2,093	0.011 [0.027] N=1,917	-0.003 [0.031] N=1,685	0.017 [0.037] N=1,398	0.063 [0.051] N=1,086
Wasted: WLZ/WHZ below -2, 0-60 months	-0.025 [0.036] N=1,446	-0.010 [0.036] N=1,329	0.039 [0.041] N=1,169	0.046 [0.053] N=973	-0.003 [0.054] N=745
Overweight: WLZ/WHZ above +2, 0-60 months	0.069 [0.050] N=1,446	0.070 [0.054] N=1,329	0.040 [0.059] N=1,169	-0.000 [0.073] N=973	-0.016 [0.094] N=745
Obese: WLZ/WHZ above +3, 0-60 months	0.029 [0.034] N=1,446	0.029 [0.037] N=1,329	0.048 [0.042] N=1,169	0.050 [0.052] N=973	0.054 [0.078] N=745
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.004 [0.058] N=1,417	-0.016 [0.064] N=1,303	0.030 [0.073] N=1,146	-0.035 [0.086] N=953	-0.004 [0.100] N=731
BMI-for-age z score below -2	-0.015 [0.035] N=1,857	-0.018 [0.033] N=1,709	-0.005 [0.039] N=1,495	0.010 [0.046] N=1,239	-0.050 [0.060] N=961
WHO 2007 BMI overweight above +1, 61+ months	-0.060 [0.117] N=403	-0.065 [0.123] N=374	0.055 [0.138] N=324	0.031 [0.162] N=268	0.107 [0.230] N=213
WHO 2007 BMI obesity above +2, 61+ months	-0.021 [0.088] N=403	-0.033 [0.091] N=374	-0.009 [0.102] N=324	-0.097 [0.116] N=268	-0.119 [0.156] N=213
WHO 2007 BMI thinness below -2, 61+ months	0.091 [0.079] N=403	0.059 [0.075] N=374	-0.065 [0.080] N=324	-0.115 [0.097] N=268	**-0.233* [0.124] N=213**

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.475	-0.199 [0.196] N=2,342	-0.241 [0.207] N=2,116	-0.230 [0.221] N=1,859	**-0.385* [0.231] N=1,583**	-0.449 [0.273] N=1,288
WHO weight-for-age z score	0.021	-0.006 [0.110] N=2,537	-0.005 [0.114] N=2,304	-0.068 [0.121] N=2,018	-0.081 [0.129] N=1,711	-0.050 [0.138] N=1,384
WHO weight-for-length/height z score, 0-60 months	0.364	-0.094 [0.193] N=1,712	-0.095 [0.202] N=1,551	-0.152 [0.227] N=1,351	-0.062 [0.249] N=1,154	-0.060 [0.291] N=927
WHO BMI-for-age z score	0.361	-0.079 [0.187] N=2,246	-0.100 [0.199] N=2,029	-0.111 [0.217] N=1,784	0.003 [0.232] N=1,518	-0.012 [0.265] N=1,235
Stunted: HAZ below -2	0.248 P=580	0.052 [0.036] N=2,342	0.055 [0.038] N=2,116	0.047 [0.040] N=1,859	0.069 [0.042] N=1,583	**0.081* [0.048] N=1,288**
Underweight: WAZ below -2	0.065 P=161	-0.015 [0.019] N=2,537	-0.017 [0.020] N=2,304	-0.007 [0.020] N=2,018	-0.014 [0.022] N=1,711	-0.014 [0.022] N=1,384
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=190	0.047 [0.033] N=1,712	0.036 [0.036] N=1,551	0.052 [0.038] N=1,351	0.019 [0.041] N=1,154	0.015 [0.049] N=927
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=334	0.021 [0.039] N=1,712	0.017 [0.041] N=1,551	0.006 [0.044] N=1,351	0.010 [0.047] N=1,154	0.008 [0.053] N=927
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=160	0.005 [0.029] N=1,712	-0.004 [0.030] N=1,551	-0.012 [0.033] N=1,351	-0.016 [0.037] N=1,154	-0.040 [0.038] N=927
CIAF: any stunting, underweight, or wasting, 0-60 months	0.336 P=561	**0.086* [0.045] N=1,674**	0.064 [0.048] N=1,513	0.067 [0.048] N=1,316	0.038 [0.053] N=1,123	0.064 [0.056] N=901
BMI-for-age z score below -2	0.120 P=270	0.043 [0.030] N=2,246	0.038 [0.033] N=2,029	0.042 [0.036] N=1,784	0.025 [0.036] N=1,518	0.026 [0.042] N=1,235
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=205	-0.043 [0.083] N=532	-0.077 [0.086] N=477	-0.019 [0.087] N=430	0.033 [0.090] N=363	-0.016 [0.100] N=301
WHO 2007 BMI obesity above +2, 61+ months	0.233 P=121	0.024 [0.078] N=532	-0.040 [0.081] N=477	-0.016 [0.086] N=430	0.040 [0.091] N=363	0.011 [0.104] N=301
WHO 2007 BMI thinness below -2, 61+ months	0.117 P=62	0.049 [0.053] N=532	0.050 [0.055] N=477	0.043 [0.057] N=430	0.073 [0.059] N=363	0.110 [0.067] N=301

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.040 [0.257] N=2,011	-0.075 [0.280] N=1,785	-0.016 [0.317] N=1,528	-0.378 [0.334] N=1,252	-0.419 [0.456] N=957
WHO weight-for-age z score	-0.044 [0.157] N=2,182	-0.040 [0.168] N=1,949	-0.185 [0.188] N=1,663	-0.289 [0.216] N=1,356	-0.264 [0.255] N=1,029
WHO weight-for-length/height z score, 0-60 months	-0.185 [0.242] N=1,484	-0.204 [0.248] N=1,323	-0.344 [0.304] N=1,123	-0.199 [0.329] N=926	-0.446 [0.430] N=699
WHO BMI-for-age z score	**-0.424** [0.214] N=1,930**	**-0.534** [0.225] N=1,713**	**-0.632** [0.268] N=1,468**	-0.484 [0.301] N=1,202	**-0.930** [0.397] N=919**
Stunted: HAZ below -2	0.025 [0.053] N=2,011	0.023 [0.057] N=1,785	-0.004 [0.062] N=1,528	0.035 [0.068] N=1,252	0.036 [0.096] N=957
Underweight: WAZ below -2	-0.032 [0.028] N=2,182	-0.037 [0.032] N=1,949	-0.020 [0.035] N=1,663	-0.037 [0.041] N=1,356	-0.051 [0.041] N=1,029
Wasted: WLZ/WHZ below -2, 0-60 months	**0.104*** [0.037] N=1,484**	**0.092** [0.041] N=1,323**	**0.131*** [0.046] N=1,123**	0.071 [0.047] N=926	0.088 [0.053] N=699
Overweight: WLZ/WHZ above +2, 0-60 months	0.039 [0.054] N=1,484	0.030 [0.058] N=1,323	0.019 [0.064] N=1,123	0.030 [0.076] N=926	-0.022 [0.096] N=699
Obese: WLZ/WHZ above +3, 0-60 months	0.025 [0.041] N=1,484	0.011 [0.043] N=1,323	-0.004 [0.050] N=1,123	-0.009 [0.064] N=926	-0.125 [0.076] N=699
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.117* [0.070] N=1,454**	0.082 [0.077] N=1,293	0.081 [0.083] N=1,096	0.018 [0.098] N=903	0.044 [0.119] N=681
BMI-for-age z score below -2	**0.101*** [0.031] N=1,930**	**0.102*** [0.037] N=1,713**	**0.119*** [0.043] N=1,468**	**0.095** [0.037] N=1,202**	**0.148*** [0.049] N=919**
WHO 2007 BMI overweight above +1, 61+ months	-0.154 [0.135] N=443	**-0.258* [0.147] N=388**	-0.175 [0.162] N=341	-0.100 [0.206] N=274	-0.346 [0.309] N=212
WHO 2007 BMI obesity above +2, 61+ months	0.026 [0.119] N=443	-0.108 [0.131] N=388	-0.080 [0.149] N=341	0.049 [0.170] N=274	-0.025 [0.259] N=212
WHO 2007 BMI thinness below -2, 61+ months	0.106 [0.068] N=443	0.121 [0.073] N=388	0.115 [0.079] N=341	**0.213** [0.084] N=274**	**0.413*** [0.113] N=212**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.475	-0.198 [0.196] N=2,342	-0.240 [0.207] N=2,116	-0.230 [0.222] N=1,859	-0.382 [0.232] N=1,583	-0.441 [0.274] N=1,288
WHO weight-for-age z score	0.021	-0.004 [0.110] N=2,537	-0.003 [0.114] N=2,304	-0.066 [0.121] N=2,018	-0.079 [0.129] N=1,711	-0.049 [0.139] N=1,384
WHO weight-for-length/height z score, 0-60 months	0.364	-0.092 [0.192] N=1,712	-0.094 [0.201] N=1,551	-0.152 [0.226] N=1,351	-0.063 [0.249] N=1,154	-0.072 [0.289] N=927
WHO BMI-for-age z score	0.361	-0.083 [0.187] N=2,246	-0.105 [0.199] N=2,029	-0.115 [0.217] N=1,784	-0.002 [0.233] N=1,518	-0.029 [0.269] N=1,235
Stunted: HAZ below -2	0.248 P=580	0.053 [0.037] N=2,342	0.055 [0.038] N=2,116	0.048 [0.040] N=1,859	0.069 [0.042] N=1,583	**0.082* [0.048] N=1,288**
Underweight: WAZ below -2	0.065 P=161	-0.015 [0.019] N=2,537	-0.017 [0.020] N=2,304	-0.007 [0.020] N=2,018	-0.015 [0.022] N=1,711	-0.014 [0.022] N=1,384
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=190	0.046 [0.033] N=1,712	0.036 [0.035] N=1,551	0.052 [0.038] N=1,351	0.020 [0.040] N=1,154	0.018 [0.049] N=927
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=334	0.021 [0.039] N=1,712	0.016 [0.041] N=1,551	0.006 [0.044] N=1,351	0.010 [0.048] N=1,154	0.009 [0.053] N=927
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=160	0.004 [0.029] N=1,712	-0.004 [0.030] N=1,551	-0.012 [0.033] N=1,351	-0.016 [0.038] N=1,154	-0.038 [0.039] N=927
CIAF: any stunting, underweight, or wasting, 0-60 months	0.336 P=561	**0.085* [0.045] N=1,674**	0.064 [0.048] N=1,513	0.067 [0.049] N=1,316	0.039 [0.053] N=1,123	0.067 [0.056] N=901
BMI-for-age z score below -2	0.120 P=270	0.044 [0.030] N=2,246	0.039 [0.033] N=2,029	0.043 [0.036] N=1,784	0.026 [0.036] N=1,518	0.029 [0.043] N=1,235
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=205	-0.035 [0.083] N=532	-0.071 [0.087] N=477	-0.014 [0.088] N=430	0.034 [0.091] N=363	-0.012 [0.101] N=301
WHO 2007 BMI obesity above +2, 61+ months	0.233 P=121	0.033 [0.079] N=532	-0.032 [0.082] N=477	-0.011 [0.087] N=430	0.040 [0.092] N=363	0.012 [0.105] N=301
WHO 2007 BMI thinness below -2, 61+ months	0.117 P=62	0.045 [0.053] N=532	0.048 [0.055] N=477	0.040 [0.056] N=430	0.071 [0.059] N=363	0.108 [0.068] N=301

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.041 [0.256] N=2,011	-0.077 [0.279] N=1,785	-0.016 [0.316] N=1,528	-0.389 [0.328] N=1,252	-0.426 [0.453] N=957
WHO weight-for-age z score	-0.044 [0.157] N=2,182	-0.040 [0.168] N=1,949	-0.186 [0.188] N=1,663	-0.291 [0.215] N=1,356	-0.264 [0.255] N=1,029
WHO weight-for-length/height z score, 0-60 months	-0.172 [0.242] N=1,484	-0.191 [0.249] N=1,323	-0.336 [0.304] N=1,123	-0.186 [0.327] N=926	-0.422 [0.434] N=699
WHO BMI-for-age z score	**-0.422* [0.214] N=1,930**	**-0.530** [0.225] N=1,713**	**-0.630** [0.268] N=1,468**	-0.473 [0.301] N=1,202	**-0.922** [0.402] N=919**
Stunted: HAZ below -2	0.025 [0.053] N=2,011	0.023 [0.057] N=1,785	-0.004 [0.062] N=1,528	0.035 [0.067] N=1,252	0.036 [0.096] N=957
Underweight: WAZ below -2	-0.032 [0.028] N=2,182	-0.037 [0.032] N=1,949	-0.020 [0.035] N=1,663	-0.037 [0.041] N=1,356	-0.051 [0.041] N=1,029
Wasted: WLZ/WHZ below -2, 0-60 months	**0.101*** [0.038] N=1,484**	**0.088** [0.041] N=1,323**	**0.127*** [0.046] N=1,123**	0.065 [0.047] N=926	0.081 [0.053] N=699
Overweight: WLZ/WHZ above +2, 0-60 months	0.039 [0.054] N=1,484	0.029 [0.058] N=1,323	0.017 [0.064] N=1,123	0.027 [0.075] N=926	-0.023 [0.096] N=699
Obese: WLZ/WHZ above +3, 0-60 months	0.024 [0.041] N=1,484	0.009 [0.043] N=1,323	-0.008 [0.049] N=1,123	-0.013 [0.064] N=926	**-0.129* [0.076] N=699**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.114 [0.071] N=1,454	0.078 [0.078] N=1,293	0.076 [0.083] N=1,096	0.012 [0.099] N=903	0.039 [0.120] N=681
BMI-for-age z score below -2	**0.101*** [0.031] N=1,930**	**0.101*** [0.037] N=1,713**	**0.118*** [0.043] N=1,468**	**0.093** [0.037] N=1,202**	**0.147*** [0.049] N=919**
WHO 2007 BMI overweight above +1, 61+ months	-0.148 [0.135] N=443	**-0.255* [0.148] N=388**	-0.173 [0.162] N=341	-0.100 [0.206] N=274	-0.347 [0.311] N=212
WHO 2007 BMI obesity above +2, 61+ months	0.035 [0.121] N=443	-0.101 [0.133] N=388	-0.076 [0.151] N=341	0.049 [0.169] N=274	-0.027 [0.261] N=212
WHO 2007 BMI thinness below -2, 61+ months	0.102 [0.068] N=443	0.121 [0.073] N=388	0.112 [0.078] N=341	**0.213** [0.083] N=274**	**0.412*** [0.115] N=212**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.475	-0.197 [0.192] N=2,342	-0.229 [0.202] N=2,116	-0.213 [0.217] N=1,859	-0.366 [0.225] N=1,583	-0.422 [0.263] N=1,288
WHO weight-for-age z score	0.021	-0.003 [0.109] N=2,537	0.003 [0.113] N=2,304	-0.047 [0.121] N=2,018	-0.076 [0.129] N=1,711	-0.043 [0.140] N=1,384
WHO weight-for-length/height z score, 0-60 months	0.364	-0.093 [0.192] N=1,712	-0.092 [0.202] N=1,551	-0.134 [0.228] N=1,351	-0.053 [0.250] N=1,154	-0.052 [0.290] N=927
WHO BMI-for-age z score	0.361	-0.079 [0.188] N=2,246	-0.101 [0.199] N=2,029	-0.104 [0.216] N=1,784	-0.001 [0.232] N=1,518	-0.030 [0.269] N=1,235
Stunted: HAZ below -2	0.248 P=580	0.053 [0.036] N=2,342	0.053 [0.038] N=2,116	0.044 [0.039] N=1,859	0.065 [0.041] N=1,583	**0.078* [0.046] N=1,288**
Underweight: WAZ below -2	0.065 P=161	-0.014 [0.018] N=2,537	-0.017 [0.020] N=2,304	-0.010 [0.020] N=2,018	-0.014 [0.021] N=1,711	-0.013 [0.022] N=1,384
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=190	0.047 [0.033] N=1,712	0.034 [0.036] N=1,551	0.049 [0.039] N=1,351	0.018 [0.041] N=1,154	0.016 [0.049] N=927
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=334	0.023 [0.039] N=1,712	0.018 [0.041] N=1,551	0.010 [0.044] N=1,351	0.013 [0.048] N=1,154	0.013 [0.053] N=927
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=160	0.007 [0.030] N=1,712	-0.002 [0.030] N=1,551	-0.009 [0.033] N=1,351	-0.014 [0.038] N=1,154	-0.034 [0.040] N=927
CIAF: any stunting, underweight, or wasting, 0-60 months	0.336 P=561	**0.093** [0.045] N=1,674**	0.067 [0.048] N=1,513	0.068 [0.049] N=1,316	0.041 [0.053] N=1,123	0.072 [0.056] N=901
BMI-for-age z score below -2	0.120 P=270	0.044 [0.030] N=2,246	0.038 [0.033] N=2,029	0.040 [0.036] N=1,784	0.024 [0.037] N=1,518	0.027 [0.043] N=1,235
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=205	-0.061 [0.082] N=532	-0.096 [0.087] N=477	-0.041 [0.088] N=430	-0.000 [0.093] N=363	-0.059 [0.106] N=301
WHO 2007 BMI obesity above +2, 61+ months	0.233 P=121	0.012 [0.078] N=532	-0.051 [0.082] N=477	-0.027 [0.086] N=430	0.011 [0.091] N=363	-0.019 [0.106] N=301
WHO 2007 BMI thinness below -2, 61+ months	0.117 P=62	0.038 [0.052] N=532	0.039 [0.054] N=477	0.030 [0.055] N=430	0.063 [0.057] N=363	0.097 [0.066] N=301

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.029 [0.254] N=2,011	-0.053 [0.277] N=1,785	0.026 [0.314] N=1,528	-0.347 [0.323] N=1,252	-0.342 [0.439] N=957
WHO weight-for-age z score	-0.046 [0.154] N=2,182	-0.040 [0.166] N=1,949	-0.162 [0.188] N=1,663	-0.301 [0.214] N=1,356	-0.270 [0.250] N=1,029
WHO weight-for-length/height z score, 0-60 months	-0.193 [0.244] N=1,484	-0.209 [0.253] N=1,323	-0.340 [0.310] N=1,123	-0.203 [0.335] N=926	-0.406 [0.441] N=699
WHO BMI-for-age z score	**-0.437** [0.214] N=1,930**	**-0.543** [0.226] N=1,713**	**-0.646** [0.267] N=1,468**	**-0.511* [0.303] N=1,202**	**-0.969** [0.405] N=919**
Stunted: HAZ below -2	0.022 [0.053] N=2,011	0.017 [0.057] N=1,785	-0.015 [0.062] N=1,528	0.023 [0.067] N=1,252	0.016 [0.092] N=957
Underweight: WAZ below -2	-0.033 [0.027] N=2,182	-0.039 [0.031] N=1,949	-0.026 [0.034] N=1,663	-0.039 [0.039] N=1,356	-0.053 [0.039] N=1,029
Wasted: WLZ/WHZ below -2, 0-60 months	**0.104*** [0.038] N=1,484**	**0.088** [0.042] N=1,323**	**0.124*** [0.047] N=1,123**	0.064 [0.048] N=926	0.079 [0.053] N=699
Overweight: WLZ/WHZ above +2, 0-60 months	0.037 [0.055] N=1,484	0.028 [0.059] N=1,323	0.016 [0.066] N=1,123	0.023 [0.078] N=926	-0.022 [0.100] N=699
Obese: WLZ/WHZ above +3, 0-60 months	0.023 [0.042] N=1,484	0.007 [0.044] N=1,323	-0.009 [0.051] N=1,123	-0.016 [0.066] N=926	-0.127 [0.078] N=699
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.119* [0.071] N=1,454**	0.077 [0.079] N=1,293	0.068 [0.085] N=1,096	0.009 [0.100] N=903	0.041 [0.118] N=681
BMI-for-age z score below -2	**0.101*** [0.031] N=1,930**	**0.099*** [0.037] N=1,713**	**0.114*** [0.043] N=1,468**	**0.089** [0.038] N=1,202**	**0.143*** [0.051] N=919**
WHO 2007 BMI overweight above +1, 61+ months	-0.180 [0.132] N=443	**-0.288** [0.144] N=388**	-0.213 [0.159] N=341	-0.170 [0.208] N=274	**-0.527* [0.300] N=212**
WHO 2007 BMI obesity above +2, 61+ months	0.011 [0.121] N=443	-0.123 [0.132] N=388	-0.094 [0.150] N=341	-0.001 [0.170] N=274	-0.140 [0.263] N=212
WHO 2007 BMI thinness below -2, 61+ months	0.094 [0.068] N=443	0.111 [0.073] N=388	0.098 [0.078] N=341	**0.201** [0.084] N=274**	**0.396*** [0.124] N=212**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.475	-0.206 [0.191] N=2,342	-0.236 [0.201] N=2,116	-0.230 [0.218] N=1,859	**-0.387* [0.225] N=1,583**	**-0.452* [0.264] N=1,288**
WHO weight-for-age z score	0.021	-0.007 [0.107] N=2,537	0.001 [0.110] N=2,304	-0.053 [0.117] N=2,018	-0.067 [0.123] N=1,711	-0.052 [0.131] N=1,384
WHO weight-for-length/height z score, 0-60 months	0.364	-0.082 [0.191] N=1,712	-0.085 [0.200] N=1,551	-0.136 [0.226] N=1,351	-0.042 [0.245] N=1,154	-0.049 [0.289] N=927
WHO BMI-for-age z score	0.361	-0.080 [0.187] N=2,246	-0.102 [0.199] N=2,029	-0.103 [0.216] N=1,784	0.018 [0.230] N=1,518	-0.014 [0.267] N=1,235
Stunted: HAZ below -2	0.248 P=580	0.054 [0.036] N=2,342	0.054 [0.037] N=2,116	0.047 [0.040] N=1,859	**0.069* [0.041] N=1,583**	**0.081* [0.046] N=1,288**
Underweight: WAZ below -2	0.065 P=161	-0.014 [0.018] N=2,537	-0.017 [0.019] N=2,304	-0.008 [0.020] N=2,018	-0.015 [0.021] N=1,711	-0.012 [0.021] N=1,384
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=190	0.046 [0.033] N=1,712	0.035 [0.036] N=1,551	0.052 [0.038] N=1,351	0.019 [0.040] N=1,154	0.014 [0.049] N=927
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=334	0.023 [0.039] N=1,712	0.018 [0.041] N=1,551	0.009 [0.044] N=1,351	0.014 [0.047] N=1,154	0.009 [0.053] N=927
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=160	0.009 [0.029] N=1,712	-0.000 [0.030] N=1,551	-0.009 [0.033] N=1,351	-0.010 [0.037] N=1,154	-0.033 [0.039] N=927
CIAF: any stunting, underweight, or wasting, 0-60 months	0.336 P=561	**0.091** [0.044] N=1,674**	0.066 [0.047] N=1,513	0.070 [0.048] N=1,316	0.043 [0.052] N=1,123	0.069 [0.055] N=901
BMI-for-age z score below -2	0.120 P=270	0.044 [0.030] N=2,246	0.037 [0.033] N=2,029	0.041 [0.036] N=1,784	0.024 [0.036] N=1,518	0.024 [0.043] N=1,235
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=205	-0.047 [0.082] N=532	-0.079 [0.086] N=477	-0.017 [0.087] N=430	0.029 [0.091] N=363	-0.029 [0.102] N=301
WHO 2007 BMI obesity above +2, 61+ months	0.233 P=121	0.020 [0.078] N=532	-0.041 [0.082] N=477	-0.017 [0.087] N=430	0.030 [0.093] N=363	-0.005 [0.107] N=301
WHO 2007 BMI thinness below -2, 61+ months	0.117 P=62	0.044 [0.053] N=532	0.044 [0.054] N=477	0.033 [0.055] N=430	0.064 [0.056] N=363	0.094 [0.065] N=301

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.037 [0.251] N=2,011	-0.052 [0.272] N=1,785	0.021 [0.314] N=1,528	-0.348 [0.320] N=1,252	-0.408 [0.423] N=957
WHO weight-for-age z score	-0.036 [0.151] N=2,182	-0.022 [0.163] N=1,949	-0.146 [0.182] N=1,663	-0.242 [0.204] N=1,356	-0.240 [0.233] N=1,029
WHO weight-for-length/height z score, 0-60 months	-0.165 [0.242] N=1,484	-0.192 [0.247] N=1,323	-0.335 [0.300] N=1,123	-0.179 [0.319] N=926	-0.375 [0.429] N=699
WHO BMI-for-age z score	**-0.433** [0.214] N=1,930**	**-0.545** [0.225] N=1,713**	**-0.646** [0.267] N=1,468**	-0.477 [0.293] N=1,202	**-0.913** [0.400] N=919**
Stunted: HAZ below -2	0.026 [0.052] N=2,011	0.020 [0.056] N=1,785	-0.011 [0.062] N=1,528	0.028 [0.066] N=1,252	0.034 [0.094] N=957
Underweight: WAZ below -2	-0.033 [0.026] N=2,182	-0.040 [0.030] N=1,949	-0.027 [0.033] N=1,663	-0.045 [0.038] N=1,356	-0.055 [0.037] N=1,029
Wasted: WLZ/WHZ below -2, 0-60 months	**0.102*** [0.037] N=1,484**	**0.089** [0.042] N=1,323**	**0.129*** [0.045] N=1,123**	0.066 [0.046] N=926	0.083 [0.053] N=699
Overweight: WLZ/WHZ above +2, 0-60 months	0.041 [0.054] N=1,484	0.030 [0.058] N=1,323	0.018 [0.064] N=1,123	0.028 [0.075] N=926	-0.014 [0.096] N=699
Obese: WLZ/WHZ above +3, 0-60 months	0.027 [0.041] N=1,484	0.009 [0.042] N=1,323	-0.009 [0.050] N=1,123	-0.014 [0.063] N=926	-0.124 [0.076] N=699
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.121* [0.069] N=1,454**	0.079 [0.077] N=1,293	0.071 [0.083] N=1,096	0.012 [0.095] N=903	0.062 [0.113] N=681
BMI-for-age z score below -2	**0.103*** [0.031] N=1,930**	**0.102*** [0.038] N=1,713**	**0.118*** [0.043] N=1,468**	**0.093** [0.037] N=1,202**	**0.147*** [0.049] N=919**
WHO 2007 BMI overweight above +1, 61+ months	-0.175 [0.127] N=443	**-0.274* [0.140] N=388**	-0.181 [0.150] N=341	-0.128 [0.193] N=274	-0.447 [0.281] N=212
WHO 2007 BMI obesity above +2, 61+ months	0.012 [0.119] N=443	-0.118 [0.129] N=388	-0.086 [0.148] N=341	0.025 [0.169] N=274	-0.089 [0.264] N=212
WHO 2007 BMI thinness below -2, 61+ months	**0.111* [0.067] N=443**	**0.125* [0.072] N=388**	0.112 [0.075] N=341	**0.203** [0.081] N=274**	**0.378*** [0.121] N=212**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.482	-0.198 [0.198] N=2,195	-0.214 [0.212] N=1,983	-0.201 [0.227] N=1,746	-0.348 [0.239] N=1,483	**-0.467* [0.278] N=1,210**
WHO weight-for-age z score	0.024	0.018 [0.115] N=2,370	0.037 [0.119] N=2,151	-0.034 [0.126] N=1,888	-0.059 [0.135] N=1,599	-0.074 [0.145] N=1,298
WHO weight-for-length/height z score, 0-60 months	0.369	-0.013 [0.207] N=1,596	0.003 [0.216] N=1,446	-0.085 [0.239] N=1,260	-0.022 [0.262] N=1,075	-0.051 [0.300] N=866
WHO BMI-for-age z score	0.377	-0.079 [0.196] N=2,103	-0.097 [0.209] N=1,900	-0.145 [0.227] N=1,674	-0.058 [0.241] N=1,422	-0.121 [0.275] N=1,161
Stunted: HAZ below -2	0.249 P=546	0.053 [0.037] N=2,195	0.055 [0.039] N=1,983	0.044 [0.040] N=1,746	0.067 [0.042] N=1,483	**0.090* [0.048] N=1,210**
Underweight: WAZ below -2	0.067 P=154	-0.019 [0.019] N=2,370	-0.022 [0.021] N=2,151	-0.016 [0.021] N=1,888	-0.017 [0.023] N=1,599	-0.010 [0.025] N=1,298
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=178	0.045 [0.034] N=1,596	0.031 [0.036] N=1,446	0.048 [0.038] N=1,260	0.021 [0.041] N=1,075	0.023 [0.050] N=866
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=313	0.018 [0.040] N=1,596	0.016 [0.042] N=1,446	0.005 [0.044] N=1,260	0.006 [0.047] N=1,075	0.004 [0.052] N=866
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=152	0.006 [0.029] N=1,596	-0.001 [0.030] N=1,446	-0.009 [0.033] N=1,260	-0.008 [0.037] N=1,075	-0.022 [0.039] N=866
CIAF: any stunting, underweight, or wasting, 0-60 months	0.338 P=526	**0.091* [0.047] N=1,561**	0.066 [0.050] N=1,411	0.067 [0.050] N=1,228	0.042 [0.055] N=1,046	0.085 [0.059] N=841
BMI-for-age z score below -2	0.118 P=247	**0.051* [0.030] N=2,103**	0.044 [0.033] N=1,900	0.050 [0.035] N=1,674	0.041 [0.035] N=1,422	0.047 [0.041] N=1,161
WHO 2007 BMI overweight above +1, 61+ months	0.395 P=198	-0.072 [0.089] N=507	-0.125 [0.093] N=455	-0.056 [0.096] N=412	-0.018 [0.102] N=347	-0.077 [0.111] N=289
WHO 2007 BMI obesity above +2, 61+ months	0.238 P=117	-0.000 [0.086] N=507	-0.080 [0.089] N=455	-0.059 [0.095] N=412	-0.026 [0.102] N=347	-0.045 [0.115] N=289
WHO 2007 BMI thinness below -2, 61+ months	0.114 P=57	0.067 [0.056] N=507	0.073 [0.058] N=455	0.066 [0.060] N=412	0.096 [0.064] N=347	**0.145** [0.072] N=289**

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.013 [0.269] N=1,883	0.000 [0.299] N=1,671	0.054 [0.347] N=1,434	-0.305 [0.373] N=1,171	-0.462 [0.514] N=898
WHO weight-for-age z score	-0.024 [0.164] N=2,038	0.014 [0.175] N=1,819	-0.150 [0.195] N=1,556	-0.254 [0.230] N=1,267	-0.277 [0.284] N=966
WHO weight-for-length/height z score, 0-60 months	-0.098 [0.261] N=1,380	-0.098 [0.273] N=1,230	-0.302 [0.332] N=1,044	-0.179 [0.358] N=859	-0.456 [0.480] N=650
WHO BMI-for-age z score	**-0.395* [0.229] N=1,804**	**-0.488** [0.244] N=1,601**	**-0.629** [0.293] N=1,375**	-0.479 [0.333] N=1,123	**-0.977** [0.446] N=862**
Stunted: HAZ below -2	0.014 [0.055] N=1,883	0.009 [0.061] N=1,671	-0.026 [0.066] N=1,434	0.008 [0.071] N=1,171	0.022 [0.096] N=898
Underweight: WAZ below -2	-0.038 [0.029] N=2,038	-0.046 [0.033] N=1,819	-0.032 [0.035] N=1,556	-0.041 [0.043] N=1,267	-0.050 [0.045] N=966
Wasted: WLZ/WHZ below -2, 0-60 months	**0.098** [0.039] N=1,380**	**0.079* [0.043] N=1,230**	**0.118** [0.048] N=1,044**	0.058 [0.052] N=859	0.073 [0.060] N=650
Overweight: WLZ/WHZ above +2, 0-60 months	0.028 [0.057] N=1,380	0.019 [0.061] N=1,230	0.005 [0.067] N=1,044	0.006 [0.080] N=859	-0.046 [0.105] N=650
Obese: WLZ/WHZ above +3, 0-60 months	0.013 [0.044] N=1,380	-0.003 [0.045] N=1,230	-0.022 [0.052] N=1,044	-0.026 [0.067] N=859	**-0.139* [0.081] N=650**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.109 [0.077] N=1,352	0.065 [0.084] N=1,202	0.053 [0.091] N=1,019	-0.022 [0.109] N=837	0.024 [0.132] N=632
BMI-for-age z score below -2	**0.105*** [0.033] N=1,804**	**0.103** [0.040] N=1,601**	**0.122*** [0.046] N=1,375**	**0.106** [0.042] N=1,123**	**0.167*** [0.056] N=862**
WHO 2007 BMI overweight above +1, 61+ months	-0.181 [0.140] N=423	**-0.305** [0.151] N=371**	-0.196 [0.167] N=328	-0.126 [0.216] N=263	-0.399 [0.325] N=205
WHO 2007 BMI obesity above +2, 61+ months	0.021 [0.129] N=423	-0.123 [0.143] N=371	-0.092 [0.162] N=328	0.020 [0.184] N=263	-0.028 [0.281] N=205
WHO 2007 BMI thinness below -2, 61+ months	0.104 [0.071] N=423	**0.127* [0.076] N=371**	0.118 [0.081] N=328	**0.208** [0.088] N=263**	**0.406*** [0.118] N=205**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.482	-0.199 [0.198] N=2,195	-0.219 [0.211] N=1,983	-0.202 [0.228] N=1,746	-0.343 [0.239] N=1,483	-0.459 [0.277] N=1,210
WHO weight-for-age z score	0.024	0.017 [0.117] N=2,370	0.034 [0.121] N=2,151	-0.043 [0.128] N=1,888	-0.063 [0.138] N=1,599	-0.075 [0.149] N=1,298
WHO weight-for-length/height z score, 0-60 months	0.369	-0.008 [0.209] N=1,596	0.006 [0.220] N=1,446	-0.081 [0.243] N=1,260	-0.008 [0.265] N=1,075	-0.044 [0.310] N=866
WHO BMI-for-age z score	0.377	-0.078 [0.197] N=2,103	-0.097 [0.210] N=1,900	-0.143 [0.228] N=1,674	-0.053 [0.243] N=1,422	-0.120 [0.277] N=1,161
Stunted: HAZ below -2	0.249 P=546	0.051 [0.037] N=2,195	0.054 [0.039] N=1,983	0.043 [0.041] N=1,746	0.064 [0.042] N=1,483	**0.087* [0.048] N=1,210**
Underweight: WAZ below -2	0.067 P=154	-0.019 [0.019] N=2,370	-0.022 [0.021] N=2,151	-0.014 [0.021] N=1,888	-0.017 [0.023] N=1,599	-0.010 [0.025] N=1,298
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=178	0.043 [0.034] N=1,596	0.029 [0.036] N=1,446	0.046 [0.038] N=1,260	0.017 [0.041] N=1,075	0.019 [0.050] N=866
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=313	0.019 [0.040] N=1,596	0.016 [0.042] N=1,446	0.005 [0.044] N=1,260	0.007 [0.048] N=1,075	0.005 [0.054] N=866
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=152	0.006 [0.029] N=1,596	-0.001 [0.030] N=1,446	-0.009 [0.033] N=1,260	-0.008 [0.037] N=1,075	-0.023 [0.039] N=866
CIAF: any stunting, underweight, or wasting, 0-60 months	0.338 P=526	**0.087* [0.046] N=1,561**	0.062 [0.049] N=1,411	0.063 [0.049] N=1,228	0.037 [0.053] N=1,046	0.077 [0.056] N=841
BMI-for-age z score below -2	0.118 P=247	**0.050* [0.030] N=2,103**	0.043 [0.033] N=1,900	0.049 [0.035] N=1,674	0.039 [0.035] N=1,422	0.045 [0.040] N=1,161
WHO 2007 BMI overweight above +1, 61+ months	0.395 P=198	-0.082 [0.089] N=507	-0.133 [0.093] N=455	-0.066 [0.095] N=412	-0.028 [0.101] N=347	-0.074 [0.109] N=289
WHO 2007 BMI obesity above +2, 61+ months	0.238 P=117	-0.005 [0.086] N=507	-0.087 [0.089] N=455	-0.065 [0.094] N=412	-0.030 [0.101] N=347	-0.047 [0.114] N=289
WHO 2007 BMI thinness below -2, 61+ months	0.114 P=57	0.067 [0.054] N=507	0.071 [0.056] N=455	0.061 [0.057] N=412	0.093 [0.060] N=347	**0.141** [0.066] N=289**

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.034 [0.266] N=1,883	-0.028 [0.293] N=1,671	0.032 [0.341] N=1,434	-0.357 [0.367] N=1,171	-0.522 [0.503] N=898
WHO weight-for-age z score	-0.024 [0.163] N=2,038	0.009 [0.175] N=1,819	-0.164 [0.196] N=1,556	-0.262 [0.230] N=1,267	-0.312 [0.283] N=966
WHO weight-for-length/height z score, 0-60 months	-0.096 [0.260] N=1,380	-0.099 [0.270] N=1,230	-0.298 [0.329] N=1,044	-0.139 [0.356] N=859	-0.449 [0.475] N=650
WHO BMI-for-age z score	**-0.391* [0.229] N=1,804**	**-0.491** [0.244] N=1,601**	**-0.633** [0.291] N=1,375**	-0.462 [0.329] N=1,123	**-0.980** [0.441] N=862**
Stunted: HAZ below -2	0.016 [0.055] N=1,883	0.011 [0.060] N=1,671	-0.024 [0.065] N=1,434	0.016 [0.070] N=1,171	0.031 [0.095] N=898
Underweight: WAZ below -2	-0.038 [0.028] N=2,038	-0.046 [0.032] N=1,819	-0.030 [0.035] N=1,556	-0.038 [0.042] N=1,267	-0.040 [0.044] N=966
Wasted: WLZ/WHZ below -2, 0-60 months	**0.097** [0.039] N=1,380**	**0.078* [0.043] N=1,230**	**0.117** [0.047] N=1,044**	0.053 [0.050] N=859	0.073 [0.059] N=650
Overweight: WLZ/WHZ above +2, 0-60 months	0.029 [0.056] N=1,380	0.019 [0.060] N=1,230	0.005 [0.067] N=1,044	0.013 [0.080] N=859	-0.045 [0.104] N=650
Obese: WLZ/WHZ above +3, 0-60 months	0.014 [0.044] N=1,380	-0.001 [0.045] N=1,230	-0.021 [0.051] N=1,044	-0.021 [0.067] N=859	**-0.136* [0.081] N=650**
CIAF: any stunting, underweight, or wasting, 0-60 months	0.108 [0.076] N=1,352	0.063 [0.084] N=1,202	0.053 [0.090] N=1,019	-0.019 [0.108] N=837	0.027 [0.132] N=632
BMI-for-age z score below -2	**0.105*** [0.033] N=1,804**	**0.103** [0.040] N=1,601**	**0.122*** [0.046] N=1,375**	**0.105** [0.042] N=1,123**	**0.167*** [0.056] N=862**
WHO 2007 BMI overweight above +1, 61+ months	-0.179 [0.139] N=423	**-0.298* [0.151] N=371**	-0.197 [0.168] N=328	-0.125 [0.217] N=263	-0.363 [0.325] N=205
WHO 2007 BMI obesity above +2, 61+ months	0.022 [0.127] N=423	-0.126 [0.141] N=371	-0.093 [0.160] N=328	0.021 [0.182] N=263	-0.024 [0.276] N=205
WHO 2007 BMI thinness below -2, 61+ months	0.107 [0.069] N=423	**0.125* [0.074] N=371**	0.109 [0.078] N=328	**0.198** [0.083] N=263**	**0.387*** [0.111] N=205**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 71 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.