Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-5 Using 12 November 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 71 months on 12 November 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.306	0.149 [0.164] N=2,496	0.156 [0.171] N=2,320	0.093 [0.183] N=2,090	0.153 [0.199] N=1,792	0.049 [0.221] N=1,475
WHO weight-for-age z score	0.162	0.043 [0.098] N=2,699	0.102 [0.105] N=2,503	0.128 [0.108] N=2,255	0.080 [0.113] N=1,933	0.056 [0.123] N=1,591
WHO weight-for-length/height z score, 0-60 months	0.341	0.042 [0.170] N=1,864	0.025 [0.174] N=1,727	-0.068 [0.186] N=1,555	-0.180 [0.213] N=1,334	-0.094 [0.228] N=1,092
WHO BMI-for-age z score	0.336	-0.001 [0.173] N=2,389	0.011 [0.178] N=2,225	-0.014 [0.195] N=1,996	-0.153 [0.223] N=1,709	-0.069 [0.249] N=1,406
Stunted: HAZ below -2	0.215 P=523	-0.028 [0.034] N=2,496	-0.033 [0.036] N=2,320	-0.024 [0.038] N=2,090	-0.046 [0.040] N=1,792	-0.017 [0.044] N=1,475
Underweight: WAZ below -2	0.057 P=148	-0.014 [0.017] N=2,699	-0.022 [0.018] N=2,503	-0.031 [0.019] N=2,255	-0.024 [0.020] N=1,933	-0.022 [0.023] N=1,591
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=209	-0.013 [0.028] N=1,864	-0.003 [0.028] N=1,727	0.013 [0.031] N=1,555	0.019 [0.035] N=1,334	0.007 [0.038] N=1,092
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=353	0.036 [0.038] N=1,864	0.033 [0.040] N=1,727	0.024 [0.043] N=1,555	-0.002 [0.048] N=1,334	-0.006 [0.054] N=1,092
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=157	0.010 [0.028] N=1,864	0.010 [0.030] N=1,727	0.014 [0.032] N=1,555	0.005 [0.036] N=1,334	0.003 [0.041] N=1,092
CIAF: any stunting, underweight, or wasting, 0-60 months	0.287 P=516	-0.042 [0.043] N=1,827	-0.053 [0.044] N=1,693	-0.025 [0.048] N=1,525	-0.048 [0.049] N=1,306	-0.022 [0.053] N=1,071
BMI-for-age z score below -2	0.120 P=292	0.002 [0.029] N=2,389	0.002 [0.029] N=2,225	0.010 [0.032] N=1,996	0.025 [0.035] N=1,709	0.027 [0.038] N=1,406
WHO 2007 BMI overweight above +1, 61+ months	0.375 P=194	-0.005 [0.082] N=516	0.000 [0.084] N=489	0.037 [0.090] N=438	0.021 [0.102] N=376	0.036 [0.115] N=311
WHO 2007 BMI obesity above +2, 61+ months	0.205 P=108	0.008 [0.067] N=516	-0.002 [0.069] N=489	0.001 [0.072] N=438	-0.007 [0.081] N=376	-0.048 [0.097] N=311
WHO 2007 BMI thinness below -2, 61+ months	0.127 P=67	0.012 [0.060] N=516	-0.006 [0.060] N=489	-0.062 [0.066] N=438	-0.069 [0.075] N=376	-0.085 [0.090] N=311

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.134 [0.243] N=2,105	0.140 [0.261] N=1,929	0.000 [0.288] N=1,699	0.158 [0.326] N=1,401	-0.129 [0.434] N=1,084
WHO weight-for-age z score	-0.028 [0.138] N=2,280	0.072 [0.152] N=2,084	0.110 [0.166] N=1,836	0.035 [0.193] N=1,514	-0.065 [0.245] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.063 [0.232] N=1,576	0.045 [0.244] N=1,439	-0.153 [0.274] N=1,267	-0.410 [0.346] N=1,046	-0.353 [0.430] N=804
WHO BMI-for-age z score	0.132 [0.218] N=2,018	0.191 [0.225] N=1,854	0.191 [0.257] N=1,625	-0.048 [0.317] N=1,338	0.222 [0.396] N=1,035
Stunted: HAZ below -2	0.005 [0.049] N=2,105	-0.003 [0.053] N=1,929	0.022 [0.061] N=1,699	-0.022 [0.068] N=1,401	0.061 [0.090] N=1,084
Underweight: WAZ below -2	-0.008 [0.025] N=2,280	-0.023 [0.026] N=2,084	-0.041 [0.030] N=1,836	-0.031 [0.034] N=1,514	-0.011 [0.047] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.005 [0.035] N=1,576	0.012 [0.036] N=1,439	0.048 [0.041] N=1,267	0.058 [0.053] N=1,046	0.044 [0.057] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.050 [0.047] N=1,576	0.049 [0.052] N=1,439	0.034 [0.058] N=1,267	-0.022 [0.071] N=1,046	-0.045 [0.093] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.032 [0.033] N=1,576	0.032 [0.036] N=1,439	0.045 [0.041] N=1,267	0.030 [0.051] N=1,046	0.038 [0.073] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.009 [0.060] N=1,544	-0.012 [0.065] N=1,410	0.051 [0.075] N=1,242	-0.007 [0.083] N=1,023	0.086 [0.097] N=788
BMI-for-age z score below -2	-0.025 [0.035] N=2,018	-0.033 [0.035] N=1,854	-0.024 [0.040] N=1,625	-0.015 [0.048] N=1,338	-0.032 [0.058] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	0.017 [0.117] N=433	0.033 [0.124] N=406	0.104 [0.139] N=355	0.069 [0.166] N=293	0.154 [0.219] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.082 [0.093] N=433	0.075 [0.099] N=406	0.106 [0.104] N=355	0.112 [0.123] N=293	0.085 [0.165] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.004 [0.069] N=433	-0.032 [0.066] N=406	**-0.133* [0.074] N=355**	**-0.196** [0.086] N=293**	**-0.318*** [0.113] N=228**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.306	0.153 [0.165] N=2,496	0.160 [0.171] N=2,320	0.101 [0.183] N=2,090	0.162 [0.199] N=1,792	0.054 [0.222] N=1,475
WHO weight-for-age z score	0.162	0.046 [0.099] N=2,699	0.104 [0.105] N=2,503	0.132 [0.109] N=2,255	0.086 [0.113] N=1,933	0.058 [0.124] N=1,591
WHO weight-for-length/height z score, 0-60 months	0.341	0.039 [0.171] N=1,864	0.020 [0.176] N=1,727	-0.074 [0.188] N=1,555	-0.183 [0.215] N=1,334	-0.100 [0.230] N=1,092
WHO BMI-for-age z score	0.336	-0.002 [0.173] N=2,389	0.010 [0.179] N=2,225	-0.016 [0.196] N=1,996	-0.153 [0.224] N=1,709	-0.071 [0.249] N=1,406
Stunted: HAZ below -2	0.215 P=523	-0.028 [0.034] N=2,496	-0.034 [0.036] N=2,320	-0.024 [0.038] N=2,090	-0.046 [0.040] N=1,792	-0.017 [0.044] N=1,475
Underweight: WAZ below -2	0.057 P=148	-0.014 [0.017] N=2,699	-0.022 [0.018] N=2,503	-0.031 [0.019] N=2,255	-0.024 [0.020] N=1,933	-0.022 [0.023] N=1,591
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=209	-0.012 [0.028] N=1,864	-0.002 [0.028] N=1,727	0.015 [0.031] N=1,555	0.021 [0.035] N=1,334	0.009 [0.038] N=1,092
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=353	0.036 [0.038] N=1,864	0.033 [0.040] N=1,727	0.025 [0.043] N=1,555	-0.001 [0.048] N=1,334	-0.005 [0.053] N=1,092
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=157	0.010 [0.028] N=1,864	0.010 [0.029] N=1,727	0.014 [0.032] N=1,555	0.006 [0.035] N=1,334	0.003 [0.041] N=1,092
CIAF: any stunting, underweight, or wasting, 0-60 months	0.287 P=516	-0.041 [0.042] N=1,827	-0.052 [0.044] N=1,693	-0.023 [0.048] N=1,525	-0.047 [0.048] N=1,306	-0.020 [0.052] N=1,071
BMI-for-age z score below -2	0.120 P=292	0.002 [0.029] N=2,389	0.002 [0.029] N=2,225	0.011 [0.032] N=1,996	0.026 [0.035] N=1,709	0.027 [0.038] N=1,406
WHO 2007 BMI overweight above +1, 61+ months	0.375 P=194	-0.006 [0.082] N=516	-0.001 [0.084] N=489	0.037 [0.090] N=438	0.021 [0.101] N=376	0.035 [0.115] N=311
WHO 2007 BMI obesity above +2, 61+ months	0.205 P=108	0.006 [0.066] N=516	-0.005 [0.068] N=489	-0.000 [0.073] N=438	-0.007 [0.081] N=376	-0.051 [0.097] N=311
WHO 2007 BMI thinness below -2, 61+ months	0.127 P=67	0.014 [0.060] N=516	-0.004 [0.060] N=489	-0.061 [0.066] N=438	-0.068 [0.075] N=376	-0.083 [0.091] N=311

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.130 [0.242] N=2,105	0.134 [0.259] N=1,929	0.003 [0.285] N=1,699	0.159 [0.323] N=1,401	-0.153 [0.435] N=1,084
WHO weight-for-age z score	-0.030 [0.137] N=2,280	0.069 [0.151] N=2,084	0.110 [0.165] N=1,836	0.034 [0.191] N=1,514	-0.073 [0.243] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.060 [0.233] N=1,576	0.038 [0.246] N=1,439	-0.167 [0.275] N=1,267	-0.422 [0.348] N=1,046	-0.379 [0.434] N=804
WHO BMI-for-age z score	0.133 [0.218] N=2,018	0.193 [0.226] N=1,854	0.191 [0.257] N=1,625	-0.049 [0.316] N=1,338	0.232 [0.396] N=1,035
Stunted: HAZ below -2	0.006 [0.049] N=2,105	-0.003 [0.053] N=1,929	0.022 [0.061] N=1,699	-0.022 [0.068] N=1,401	0.062 [0.090] N=1,084
Underweight: WAZ below -2	-0.008 [0.025] N=2,280	-0.023 [0.026] N=2,084	-0.041 [0.030] N=1,836	-0.031 [0.034] N=1,514	-0.011 [0.047] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.004 [0.035] N=1,576	0.013 [0.036] N=1,439	0.051 [0.041] N=1,267	0.061 [0.054] N=1,046	0.049 [0.059] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.049 [0.048] N=1,576	0.049 [0.052] N=1,439	0.034 [0.058] N=1,267	-0.021 [0.070] N=1,046	-0.044 [0.092] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.032 [0.032] N=1,576	0.032 [0.035] N=1,439	0.045 [0.040] N=1,267	0.030 [0.051] N=1,046	0.038 [0.073] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.009 [0.060] N=1,544	-0.011 [0.065] N=1,410	0.052 [0.075] N=1,242	-0.005 [0.083] N=1,023	0.088 [0.097] N=788
BMI-for-age z score below -2	-0.025 [0.035] N=2,018	-0.033 [0.035] N=1,854	-0.024 [0.040] N=1,625	-0.015 [0.048] N=1,338	-0.034 [0.057] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	0.015 [0.117] N=433	0.033 [0.125] N=406	0.099 [0.140] N=355	0.067 [0.170] N=293	0.167 [0.224] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.074 [0.092] N=433	0.063 [0.099] N=406	0.089 [0.104] N=355	0.093 [0.123] N=293	0.049 [0.167] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.011 [0.069] N=433	-0.025 [0.067] N=406	**-0.125* [0.073] N=355**	**-0.179** [0.085] N=293**	**-0.297*** [0.111] N=228**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.306	0.116 [0.166] N=2,496	0.122 [0.173] N=2,320	0.067 [0.186] N=2,090	0.125 [0.203] N=1,792	0.004 [0.226] N=1,475
WHO weight-for-age z score	0.162	0.003 [0.097] N=2,699	0.062 [0.104] N=2,503	0.089 [0.108] N=2,255	0.047 [0.111] N=1,933	0.016 [0.123] N=1,591
WHO weight-for-length/height z score, 0-60 months	0.341	0.016 [0.172] N=1,864	0.005 [0.177] N=1,727	-0.084 [0.188] N=1,555	-0.190 [0.215] N=1,334	-0.108 [0.233] N=1,092
WHO BMI-for-age z score	0.336	-0.007 [0.173] N=2,389	0.008 [0.179] N=2,225	-0.018 [0.196] N=1,996	-0.153 [0.222] N=1,709	-0.059 [0.252] N=1,406
Stunted: HAZ below -2	0.215 P=523	-0.019 [0.033] N=2,496	-0.022 [0.035] N=2,320	-0.014 [0.038] N=2,090	-0.036 [0.040] N=1,792	-0.003 [0.043] N=1,475
Underweight: WAZ below -2	0.057 P=148	-0.007 [0.017] N=2,699	-0.015 [0.018] N=2,503	-0.023 [0.019] N=2,255	-0.017 [0.020] N=1,933	-0.011 [0.023] N=1,591
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=209	-0.005 [0.028] N=1,864	0.006 [0.028] N=1,727	0.021 [0.031] N=1,555	0.026 [0.035] N=1,334	0.015 [0.037] N=1,092
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=353	0.036 [0.038] N=1,864	0.035 [0.041] N=1,727	0.025 [0.043] N=1,555	-0.002 [0.048] N=1,334	-0.005 [0.054] N=1,092
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=157	0.011 [0.028] N=1,864	0.012 [0.029] N=1,727	0.015 [0.031] N=1,555	0.006 [0.035] N=1,334	0.003 [0.041] N=1,092
CIAF: any stunting, underweight, or wasting, 0-60 months	0.287 P=516	-0.022 [0.041] N=1,827	-0.030 [0.042] N=1,693	-0.008 [0.046] N=1,525	-0.035 [0.046] N=1,306	-0.002 [0.050] N=1,071
BMI-for-age z score below -2	0.120 P=292	0.008 [0.029] N=2,389	0.010 [0.029] N=2,225	0.018 [0.032] N=1,996	0.032 [0.035] N=1,709	0.035 [0.038] N=1,406
WHO 2007 BMI overweight above +1, 61+ months	0.375 P=194	0.001 [0.081] N=516	0.007 [0.084] N=489	0.046 [0.089] N=438	0.034 [0.100] N=376	0.041 [0.114] N=311
WHO 2007 BMI obesity above +2, 61+ months	0.205 P=108	0.018 [0.066] N=516	0.005 [0.068] N=489	0.010 [0.071] N=438	-0.002 [0.077] N=376	-0.042 [0.096] N=311
WHO 2007 BMI thinness below -2, 61+ months	0.127 P=67	0.020 [0.060] N=516	0.002 [0.060] N=489	-0.054 [0.066] N=438	-0.059 [0.075] N=376	-0.067 [0.091] N=311

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.106 [0.240] N=2,105	0.105 [0.256] N=1,929	-0.022 [0.283] N=1,699	0.118 [0.326] N=1,401	-0.198 [0.434] N=1,084
WHO weight-for-age z score	-0.063 [0.133] N=2,280	0.037 [0.149] N=2,084	0.076 [0.162] N=1,836	0.006 [0.185] N=1,514	-0.110 [0.239] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.042 [0.234] N=1,576	0.030 [0.249] N=1,439	-0.167 [0.276] N=1,267	-0.431 [0.350] N=1,046	-0.399 [0.436] N=804
WHO BMI-for-age z score	0.130 [0.219] N=2,018	0.193 [0.228] N=1,854	0.190 [0.258] N=1,625	-0.044 [0.318] N=1,338	0.251 [0.401] N=1,035
Stunted: HAZ below -2	0.011 [0.047] N=2,105	0.006 [0.051] N=1,929	0.028 [0.059] N=1,699	-0.016 [0.066] N=1,401	0.070 [0.088] N=1,084
Underweight: WAZ below -2	-0.003 [0.024] N=2,280	-0.017 [0.026] N=2,084	-0.034 [0.029] N=1,836	-0.028 [0.033] N=1,514	-0.004 [0.046] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	0.004 [0.035] N=1,576	0.022 [0.037] N=1,439	0.055 [0.041] N=1,267	0.067 [0.054] N=1,046	0.056 [0.059] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.051 [0.048] N=1,576	0.053 [0.053] N=1,439	0.036 [0.058] N=1,267	-0.021 [0.071] N=1,046	-0.037 [0.091] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.033 [0.033] N=1,576	0.035 [0.036] N=1,439	0.045 [0.040] N=1,267	0.027 [0.051] N=1,046	0.030 [0.073] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.028 [0.058] N=1,544	0.012 [0.062] N=1,410	0.062 [0.072] N=1,242	-0.003 [0.080] N=1,023	0.095 [0.094] N=788
BMI-for-age z score below -2	-0.019 [0.035] N=2,018	-0.026 [0.035] N=1,854	-0.019 [0.040] N=1,625	-0.011 [0.048] N=1,338	-0.028 [0.058] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	0.018 [0.117] N=433	0.034 [0.124] N=406	0.103 [0.139] N=355	0.073 [0.168] N=293	0.159 [0.226] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.079 [0.093] N=433	0.065 [0.101] N=406	0.095 [0.104] N=355	0.089 [0.119] N=293	0.063 [0.169] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.016 [0.068] N=433	-0.023 [0.066] N=406	**-0.124* [0.074] N=355**	**-0.164* [0.084] N=293**	**-0.269** [0.106] N=228**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.306	0.098 [0.166] N=2,496	0.110 [0.173] N=2,320	0.053 [0.186] N=2,090	0.100 [0.202] N=1,792	-0.032 [0.225] N=1,475
WHO weight-for-age z score	0.162	0.014 [0.097] N=2,699	0.070 [0.104] N=2,503	0.095 [0.109] N=2,255	0.043 [0.113] N=1,933	-0.003 [0.124] N=1,591
WHO weight-for-length/height z score, 0-60 months	0.341	0.048 [0.172] N=1,864	0.036 [0.177] N=1,727	-0.060 [0.189] N=1,555	-0.170 [0.217] N=1,334	-0.088 [0.232] N=1,092
WHO BMI-for-age z score	0.336	0.014 [0.175] N=2,389	0.027 [0.180] N=2,225	-0.000 [0.197] N=1,996	-0.133 [0.225] N=1,709	-0.043 [0.252] N=1,406
Stunted: HAZ below -2	0.215 P=523	-0.015 [0.034] N=2,496	-0.019 [0.035] N=2,320	-0.010 [0.038] N=2,090	-0.028 [0.039] N=1,792	0.004 [0.043] N=1,475
Underweight: WAZ below -2	0.057 P=148	-0.008 [0.017] N=2,699	-0.015 [0.018] N=2,503	-0.023 [0.019] N=2,255	-0.015 [0.020] N=1,933	-0.010 [0.023] N=1,591
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=209	-0.007 [0.027] N=1,864	0.005 [0.028] N=1,727	0.021 [0.031] N=1,555	0.027 [0.034] N=1,334	0.015 [0.037] N=1,092
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=353	0.040 [0.038] N=1,864	0.038 [0.041] N=1,727	0.029 [0.043] N=1,555	0.004 [0.048] N=1,334	-0.005 [0.054] N=1,092
Obese: WLZ/WHZ above +3, 0-60 months	0.085 P=157	0.017 [0.028] N=1,864	0.017 [0.030] N=1,727	0.020 [0.032] N=1,555	0.012 [0.036] N=1,334	0.011 [0.042] N=1,092
CIAF: any stunting, underweight, or wasting, 0-60 months	0.287 P=516	-0.018 [0.041] N=1,827	-0.027 [0.043] N=1,693	-0.001 [0.047] N=1,525	-0.021 [0.047] N=1,306	0.007 [0.051] N=1,071
BMI-for-age z score below -2	0.120 P=292	0.005 [0.029] N=2,389	0.005 [0.029] N=2,225	0.014 [0.032] N=1,996	0.030 [0.035] N=1,709	0.031 [0.038] N=1,406
WHO 2007 BMI overweight above +1, 61+ months	0.375 P=194	-0.002 [0.082] N=516	0.001 [0.084] N=489	0.038 [0.091] N=438	0.025 [0.101] N=376	0.042 [0.115] N=311
WHO 2007 BMI obesity above +2, 61+ months	0.205 P=108	0.014 [0.066] N=516	-0.000 [0.068] N=489	0.004 [0.072] N=438	-0.010 [0.081] N=376	-0.049 [0.098] N=311
WHO 2007 BMI thinness below -2, 61+ months	0.127 P=67	0.015 [0.059] N=516	-0.000 [0.060] N=489	-0.060 [0.066] N=438	-0.061 [0.075] N=376	-0.073 [0.091] N=311

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.098 [0.238] N=2,105	0.108 [0.254] N=1,929	-0.014 [0.280] N=1,699	0.132 [0.322] N=1,401	-0.236 [0.428] N=1,084
WHO weight-for-age z score	-0.048 [0.136] N=2,280	0.049 [0.151] N=2,084	0.085 [0.165] N=1,836	0.008 [0.190] N=1,514	-0.105 [0.241] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.060 [0.234] N=1,576	0.053 [0.245] N=1,439	-0.158 [0.275] N=1,267	-0.426 [0.348] N=1,046	-0.371 [0.435] N=804
WHO BMI-for-age z score	0.136 [0.219] N=2,018	0.198 [0.227] N=1,854	0.191 [0.258] N=1,625	-0.061 [0.319] N=1,338	0.231 [0.397] N=1,035
Stunted: HAZ below -2	0.012 [0.048] N=2,105	0.005 [0.052] N=1,929	0.027 [0.059] N=1,699	-0.018 [0.066] N=1,401	0.073 [0.088] N=1,084
Underweight: WAZ below -2	-0.005 [0.024] N=2,280	-0.019 [0.026] N=2,084	-0.035 [0.029] N=1,836	-0.027 [0.034] N=1,514	-0.003 [0.046] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	0.003 [0.035] N=1,576	0.018 [0.037] N=1,439	0.053 [0.041] N=1,267	0.067 [0.054] N=1,046	0.054 [0.059] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.055 [0.048] N=1,576	0.055 [0.052] N=1,439	0.037 [0.058] N=1,267	-0.019 [0.071] N=1,046	-0.036 [0.091] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.036 [0.033] N=1,576	0.038 [0.035] N=1,439	0.047 [0.040] N=1,267	0.030 [0.050] N=1,046	0.036 [0.071] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.028 [0.058] N=1,544	0.008 [0.063] N=1,410	0.062 [0.072] N=1,242	0.003 [0.079] N=1,023	0.096 [0.093] N=788
BMI-for-age z score below -2	-0.019 [0.035] N=2,018	-0.029 [0.035] N=1,854	-0.020 [0.040] N=1,625	-0.007 [0.048] N=1,338	-0.025 [0.058] N=1,035
WHO 2007 BMI overweight above +1, 61+ months	0.016 [0.117] N=433	0.030 [0.124] N=406	0.101 [0.140] N=355	0.076 [0.166] N=293	0.200 [0.220] N=228
WHO 2007 BMI obesity above +2, 61+ months	0.082 [0.092] N=433	0.066 [0.099] N=406	0.086 [0.103] N=355	0.075 [0.122] N=293	0.045 [0.171] N=228
WHO 2007 BMI thinness below -2, 61+ months	0.020 [0.068] N=433	-0.013 [0.066] N=406	-0.117 [0.073] N=355	**-0.158* [0.086] N=293**	**-0.261** [0.112] N=228**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.312	0.125 [0.174] N=2,331	0.110 [0.184] N=2,171	0.045 [0.193] N=1,950	0.088 [0.208] N=1,680	-0.042 [0.231] N=1,379
WHO weight-for-age z score	0.160	0.063 [0.096] N=2,516	0.119 [0.100] N=2,338	0.148 [0.108] N=2,101	0.099 [0.111] N=1,809	0.096 [0.126] N=1,488
WHO weight-for-length/height z score, 0-60 months	0.358	0.154 [0.171] N=1,731	0.143 [0.176] N=1,609	0.031 [0.187] N=1,445	-0.069 [0.216] N=1,249	0.066 [0.236] N=1,020
WHO BMI-for-age z score	0.354	0.059 [0.172] N=2,232	0.066 [0.175] N=2,083	0.055 [0.193] N=1,865	-0.054 [0.223] N=1,605	0.072 [0.254] N=1,317
Stunted: HAZ below -2	0.217 P=493	-0.018 [0.036] N=2,331	-0.022 [0.038] N=2,171	-0.014 [0.041] N=1,950	-0.033 [0.043] N=1,680	-0.002 [0.048] N=1,379
Underweight: WAZ below -2	0.058 P=141	-0.017 [0.018] N=2,516	-0.025 [0.019] N=2,338	-0.034 [0.021] N=2,101	-0.026 [0.022] N=1,809	-0.023 [0.027] N=1,488
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=193	-0.019 [0.028] N=1,731	-0.008 [0.029] N=1,609	0.014 [0.032] N=1,445	0.019 [0.036] N=1,249	0.001 [0.039] N=1,020
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=330	0.046 [0.040] N=1,731	0.045 [0.042] N=1,609	0.033 [0.045] N=1,445	0.007 [0.051] N=1,249	0.008 [0.058] N=1,020
Obese: WLZ/WHZ above +3, 0-60 months	0.087 P=150	0.018 [0.029] N=1,731	0.021 [0.031] N=1,609	0.025 [0.033] N=1,445	0.017 [0.037] N=1,249	0.016 [0.044] N=1,020
CIAF: any stunting, underweight, or wasting, 0-60 months	0.290 P=483	-0.032 [0.047] N=1,696	-0.043 [0.048] N=1,577	-0.017 [0.052] N=1,416	-0.044 [0.054] N=1,222	-0.019 [0.059] N=999
BMI-for-age z score below -2	0.117 P=266	0.003 [0.029] N=2,232	0.003 [0.028] N=2,083	0.009 [0.032] N=1,865	0.020 [0.036] N=1,605	0.012 [0.039] N=1,317
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=185	-0.021 [0.084] N=494	-0.013 [0.087] N=468	0.037 [0.093] N=419	0.016 [0.103] N=359	0.056 [0.118] N=296
WHO 2007 BMI obesity above +2, 61+ months	0.200 P=101	-0.017 [0.067] N=494	-0.025 [0.069] N=468	-0.014 [0.074] N=419	-0.025 [0.080] N=359	-0.057 [0.100] N=296
WHO 2007 BMI thinness below -2, 61+ months	0.125 P=63	0.025 [0.058] N=494	0.009 [0.058] N=468	-0.051 [0.062] N=419	-0.066 [0.071] N=359	-0.090 [0.088] N=296

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.194 [0.254] N=1,967	0.168 [0.276] N=1,807	0.042 [0.302] N=1,586	0.200 [0.347] N=1,316	-0.142 [0.487] N=1,015
WHO weight-for-age z score	-0.026 [0.132] N=2,126	0.062 [0.141] N=1,948	0.092 [0.161] N=1,711	0.010 [0.188] N=1,419	-0.069 [0.240] N=1,098
WHO weight-for-length/height z score, 0-60 months	0.096 [0.234] N=1,463	0.075 [0.242] N=1,341	-0.166 [0.267] N=1,177	-0.400 [0.349] N=981	-0.243 [0.450] N=752
WHO BMI-for-age z score	0.109 [0.218] N=1,887	0.150 [0.221] N=1,738	0.158 [0.251] N=1,520	-0.040 [0.314] N=1,260	0.328 [0.413] N=972
Stunted: HAZ below -2	0.005 [0.050] N=1,967	-0.002 [0.054] N=1,807	0.021 [0.062] N=1,586	-0.026 [0.070] N=1,316	0.057 [0.097] N=1,015
Underweight: WAZ below -2	-0.010 [0.026] N=2,126	-0.024 [0.027] N=1,948	-0.039 [0.032] N=1,711	-0.025 [0.037] N=1,419	-0.008 [0.050] N=1,098
Wasted: WLZ/WHZ below -2, 0-60 months	-0.009 [0.035] N=1,463	0.009 [0.035] N=1,341	0.054 [0.040] N=1,177	0.059 [0.053] N=981	0.023 [0.056] N=752
Overweight: WLZ/WHZ above +2, 0-60 months	0.057 [0.047] N=1,463	0.058 [0.051] N=1,341	0.042 [0.056] N=1,177	-0.015 [0.070] N=981	-0.022 [0.096] N=752
Obese: WLZ/WHZ above +3, 0-60 months	0.033 [0.034] N=1,463	0.038 [0.037] N=1,341	0.050 [0.042] N=1,177	0.036 [0.053] N=981	0.049 [0.079] N=752
CIAF: any stunting, underweight, or wasting, 0-60 months	0.012 [0.062] N=1,433	-0.009 [0.066] N=1,314	0.050 [0.077] N=1,153	-0.025 [0.084] N=959	0.052 [0.096] N=736
BMI-for-age z score below -2	-0.009 [0.034] N=1,887	-0.017 [0.032] N=1,738	-0.007 [0.038] N=1,520	-0.005 [0.046] N=1,260	-0.053 [0.057] N=972
WHO 2007 BMI overweight above +1, 61+ months	-0.018 [0.123] N=415	-0.001 [0.130] N=389	0.086 [0.147] N=340	0.037 [0.177] N=280	0.184 [0.239] N=217
WHO 2007 BMI obesity above +2, 61+ months	0.037 [0.093] N=415	0.029 [0.100] N=389	0.058 [0.105] N=340	0.052 [0.122] N=280	0.029 [0.173] N=217
WHO 2007 BMI thinness below -2, 61+ months	0.040 [0.071] N=415	0.004 [0.069] N=389	-0.100 [0.074] N=340	**-0.177** [0.087] N=280**	**-0.318*** [0.119] N=217**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.312	0.134 [0.173] N=2,331	0.118 [0.182] N=2,171	0.041 [0.193] N=1,950	0.085 [0.208] N=1,680	-0.041 [0.231] N=1,379
WHO weight-for-age z score	0.160	0.075 [0.097] N=2,516	0.127 [0.101] N=2,338	0.155 [0.107] N=2,101	0.107 [0.111] N=1,809	0.097 [0.126] N=1,488
WHO weight-for-length/height z score, 0-60 months	0.358	0.155 [0.172] N=1,731	0.145 [0.177] N=1,609	0.031 [0.188] N=1,445	-0.070 [0.215] N=1,249	0.055 [0.237] N=1,020
WHO BMI-for-age z score	0.354	0.068 [0.171] N=2,232	0.073 [0.174] N=2,083	0.059 [0.193] N=1,865	-0.052 [0.223] N=1,605	0.067 [0.255] N=1,317
Stunted: HAZ below -2	0.217 P=493	-0.020 [0.036] N=2,331	-0.024 [0.038] N=2,171	-0.015 [0.041] N=1,950	-0.034 [0.042] N=1,680	-0.004 [0.047] N=1,379
Underweight: WAZ below -2	0.058 P=141	-0.020 [0.019] N=2,516	-0.027 [0.019] N=2,338	**-0.036* [0.021] N=2,101**	-0.028 [0.022] N=1,809	-0.026 [0.026] N=1,488
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=193	-0.019 [0.028] N=1,731	-0.008 [0.029] N=1,609	0.013 [0.032] N=1,445	0.018 [0.036] N=1,249	-0.001 [0.039] N=1,020
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=330	0.045 [0.040] N=1,731	0.044 [0.042] N=1,609	0.034 [0.045] N=1,445	0.006 [0.051] N=1,249	0.008 [0.058] N=1,020
Obese: WLZ/WHZ above +3, 0-60 months	0.087 P=150	0.017 [0.029] N=1,731	0.020 [0.031] N=1,609	0.025 [0.033] N=1,445	0.017 [0.037] N=1,249	0.017 [0.043] N=1,020
CIAF: any stunting, underweight, or wasting, 0-60 months	0.290 P=483	-0.032 [0.046] N=1,696	-0.042 [0.048] N=1,577	-0.015 [0.052] N=1,416	-0.042 [0.054] N=1,222	-0.020 [0.058] N=999
BMI-for-age z score below -2	0.117 P=266	0.002 [0.029] N=2,232	0.001 [0.028] N=2,083	0.009 [0.032] N=1,865	0.020 [0.036] N=1,605	0.011 [0.039] N=1,317
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=185	-0.024 [0.082] N=494	-0.018 [0.084] N=468	0.026 [0.090] N=419	0.012 [0.101] N=359	0.038 [0.117] N=296
WHO 2007 BMI obesity above +2, 61+ months	0.200 P=101	-0.015 [0.066] N=494	-0.026 [0.068] N=468	-0.012 [0.073] N=419	-0.019 [0.079] N=359	-0.054 [0.099] N=296
WHO 2007 BMI thinness below -2, 61+ months	0.125 P=63	0.019 [0.057] N=494	0.002 [0.057] N=468	-0.056 [0.062] N=419	-0.072 [0.071] N=359	-0.090 [0.086] N=296

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.207 [0.251] N=1,967	0.180 [0.273] N=1,807	0.041 [0.300] N=1,586	0.198 [0.345] N=1,316	-0.123 [0.483] N=1,015
WHO weight-for-age z score	-0.009 [0.132] N=2,126	0.077 [0.142] N=1,948	0.108 [0.160] N=1,711	0.033 [0.186] N=1,419	-0.047 [0.239] N=1,098
WHO weight-for-length/height z score, 0-60 months	0.115 [0.235] N=1,463	0.100 [0.245] N=1,341	-0.163 [0.269] N=1,177	-0.394 [0.349] N=981	-0.254 [0.455] N=752
WHO BMI-for-age z score	0.134 [0.218] N=1,887	0.173 [0.221] N=1,738	0.167 [0.251] N=1,520	-0.034 [0.315] N=1,260	0.320 [0.418] N=972
Stunted: HAZ below -2	0.002 [0.050] N=1,967	-0.004 [0.054] N=1,807	0.018 [0.061] N=1,586	-0.030 [0.069] N=1,316	0.048 [0.096] N=1,015
Underweight: WAZ below -2	-0.013 [0.026] N=2,126	-0.027 [0.028] N=1,948	-0.043 [0.032] N=1,711	-0.031 [0.037] N=1,419	-0.014 [0.050] N=1,098
Wasted: WLZ/WHZ below -2, 0-60 months	-0.009 [0.035] N=1,463	0.008 [0.035] N=1,341	0.055 [0.040] N=1,177	0.058 [0.053] N=981	0.017 [0.055] N=752
Overweight: WLZ/WHZ above +2, 0-60 months	0.060 [0.047] N=1,463	0.062 [0.051] N=1,341	0.043 [0.057] N=1,177	-0.012 [0.071] N=981	-0.020 [0.098] N=752
Obese: WLZ/WHZ above +3, 0-60 months	0.032 [0.034] N=1,463	0.036 [0.037] N=1,341	0.051 [0.042] N=1,177	0.038 [0.053] N=981	0.053 [0.078] N=752
CIAF: any stunting, underweight, or wasting, 0-60 months	0.013 [0.062] N=1,433	-0.008 [0.066] N=1,314	0.051 [0.077] N=1,153	-0.027 [0.084] N=959	0.037 [0.093] N=736
BMI-for-age z score below -2	-0.013 [0.034] N=1,887	-0.021 [0.032] N=1,738	-0.008 [0.038] N=1,520	-0.005 [0.046] N=1,260	-0.053 [0.057] N=972
WHO 2007 BMI overweight above +1, 61+ months	-0.010 [0.119] N=415	0.011 [0.127] N=389	0.089 [0.143] N=340	0.045 [0.173] N=280	0.180 [0.236] N=217
WHO 2007 BMI obesity above +2, 61+ months	0.036 [0.092] N=415	0.023 [0.098] N=389	0.060 [0.104] N=340	0.046 [0.118] N=280	0.016 [0.167] N=217
WHO 2007 BMI thinness below -2, 61+ months	0.030 [0.071] N=415	-0.004 [0.070] N=389	-0.106 [0.076] N=340	**-0.174** [0.087] N=280**	**-0.306** [0.118] N=217**

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.272	-0.212 [0.195] N=2,387	-0.262 [0.207] N=2,157	-0.277 [0.220] N=1,893	**-0.395*** **[0.234]** **N=1,610**	**-0.461*** **[0.268]** **N=1,310**
WHO weight-for-age z score	0.183	-0.090 [0.101] N=2,580	-0.088 [0.106] N=2,343	-0.162 [0.112] N=2,051	-0.167 [0.121] N=1,738	-0.167 [0.132] N=1,406
WHO weight-for-length/height z score, 0-60 months	0.366	-0.026 [0.189] N=1,746	-0.004 [0.200] N=1,585	-0.052 [0.223] N=1,378	0.081 [0.236] N=1,181	0.044 [0.284] N=950
WHO BMI-for-age z score	0.356	-0.082 [0.193] N=2,288	-0.108 [0.207] N=2,066	-0.107 [0.227] N=1,815	0.045 [0.245] N=1,543	-0.007 [0.282] N=1,255
Stunted: HAZ below -2	0.215 P=510	0.034 [0.035] N=2,387	0.035 [0.036] N=2,157	0.028 [0.038] N=1,893	0.037 [0.042] N=1,610	0.051 [0.048] N=1,310
Underweight: WAZ below -2	0.053 P=134	0.004 [0.017] N=2,580	0.003 [0.018] N=2,343	0.012 [0.018] N=2,051	0.004 [0.018] N=1,738	0.009 [0.020] N=1,406
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	0.030 [0.032] N=1,746	0.020 [0.035] N=1,585	0.035 [0.037] N=1,378	0.001 [0.039] N=1,181	0.012 [0.047] N=950
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=344	0.020 [0.037] N=1,746	0.018 [0.039] N=1,585	0.010 [0.042] N=1,378	0.023 [0.043] N=1,181	0.030 [0.049] N=950
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=164	0.010 [0.028] N=1,746	0.003 [0.029] N=1,585	-0.001 [0.033] N=1,378	0.010 [0.036] N=1,181	-0.009 [0.038] N=950
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=498	0.055 [0.042] N=1,710	0.039 [0.045] N=1,549	0.049 [0.045] N=1,344	0.015 [0.048] N=1,152	0.052 [0.053] N=926
BMI-for-age z score below -2	0.119 P=274	0.050 [0.032] N=2,288	0.048 [0.035] N=2,066	0.052 [0.038] N=1,815	0.032 [0.039] N=1,543	0.041 [0.045] N=1,255
WHO 2007 BMI overweight above +1, 61+ months	0.402 P=212	-0.037 [0.083] N=538	-0.096 [0.085] N=479	-0.044 [0.087] N=435	0.003 [0.095] N=362	-0.085 [0.105] N=299
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=118	0.018 [0.079] N=538	-0.056 [0.082] N=479	-0.041 [0.089] N=435	-0.001 [0.097] N=362	-0.074 [0.110] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.122 P=67	0.062 [0.057] N=538	0.074 [0.061] N=479	0.053 [0.062] N=435	0.064 [0.069] N=362	0.086 [0.078] N=299

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.039 [0.246] N=2,047	-0.104 [0.275] N=1,817	-0.059 [0.304] N=1,553	-0.269 [0.325] N=1,270	-0.299 [0.423] N=970
WHO weight-for-age z score	-0.090 [0.155] N=2,216	-0.081 [0.168] N=1,979	-0.258 [0.183] N=1,687	-0.316 [0.218] N=1,374	-0.354 [0.252] N=1,042
WHO weight-for-length/height z score, 0-60 months	-0.245 [0.244] N=1,513	-0.246 [0.256] N=1,352	-0.469 [0.304] N=1,145	-0.340 [0.332] N=948	**-0.776* [0.428] N=717**
WHO BMI-for-age z score	**-0.472** [0.215] N=1,966**	**-0.598*** [0.225] N=1,744**	**-0.720*** [0.267] N=1,493**	**-0.570* [0.310] N=1,221**	**-1.105*** [0.370] N=933**
Stunted: HAZ below -2	0.021 [0.046] N=2,047	0.019 [0.050] N=1,817	-0.009 [0.054] N=1,553	-0.005 [0.063] N=1,270	0.008 [0.087] N=970
Underweight: WAZ below -2	-0.006 [0.025] N=2,216	-0.008 [0.028] N=1,979	0.014 [0.030] N=1,687	-0.006 [0.034] N=1,374	0.001 [0.039] N=1,042
Wasted: WLZ/WHZ below -2, 0-60 months	**0.085** [0.038] N=1,513**	**0.075* [0.042] N=1,352**	**0.130*** [0.047] N=1,145**	0.070 [0.048] N=948	**0.136** [0.051] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.054] N=1,513	0.001 [0.057] N=1,352	-0.021 [0.064] N=1,145	-0.021 [0.076] N=948	-0.060 [0.094] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.009 [0.040] N=1,513	-0.008 [0.042] N=1,352	-0.022 [0.049] N=1,145	-0.013 [0.062] N=948	-0.118 [0.078] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.083 [0.063] N=1,483	0.059 [0.071] N=1,322	0.083 [0.079] N=1,117	0.006 [0.092] N=925	0.083 [0.115] N=699
BMI-for-age z score below -2	**0.107*** [0.031] N=1,966**	**0.115*** [0.037] N=1,744**	**0.137*** [0.042] N=1,493**	**0.114*** [0.038] N=1,221**	**0.179*** [0.048] N=933**
WHO 2007 BMI overweight above +1, 61+ months	-0.134 [0.139] N=449	**-0.273* [0.145] N=390**	-0.198 [0.159] N=346	-0.127 [0.205] N=273	-0.460 [0.292] N=210
WHO 2007 BMI obesity above +2, 61+ months	0.086 [0.116] N=449	-0.042 [0.124] N=390	-0.029 [0.146] N=346	0.067 [0.172] N=273	-0.066 [0.255] N=210
WHO 2007 BMI thinness below -2, 61+ months	0.111 [0.071] N=449	**0.152* [0.079] N=390**	0.107 [0.084] N=346	0.157 [0.097] N=273	**0.274** [0.116] N=210**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.272	-0.208 [0.195] N=2,387	-0.258 [0.207] N=2,157	-0.274 [0.220] N=1,893	**-0.390* [0.235] N=1,610**	-0.447 [0.269] N=1,310
WHO weight-for-age z score	0.183	-0.086 [0.101] N=2,580	-0.083 [0.106] N=2,343	-0.158 [0.112] N=2,051	-0.164 [0.121] N=1,738	-0.164 [0.134] N=1,406
WHO weight-for-length/height z score, 0-60 months	0.366	-0.024 [0.188] N=1,746	-0.003 [0.199] N=1,585	-0.052 [0.222] N=1,378	0.079 [0.236] N=1,181	0.031 [0.283] N=950
WHO BMI-for-age z score	0.356	-0.086 [0.193] N=2,288	-0.111 [0.207] N=2,066	-0.110 [0.228] N=1,815	0.042 [0.246] N=1,543	-0.019 [0.285] N=1,255
Stunted: HAZ below -2	0.215 P=510	0.034 [0.035] N=2,387	0.035 [0.036] N=2,157	0.029 [0.038] N=1,893	0.037 [0.042] N=1,610	0.052 [0.048] N=1,310
Underweight: WAZ below -2	0.053 P=134	0.004 [0.017] N=2,580	0.003 [0.018] N=2,343	0.013 [0.018] N=2,051	0.004 [0.018] N=1,738	0.010 [0.020] N=1,406
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	0.030 [0.032] N=1,746	0.019 [0.035] N=1,585	0.035 [0.037] N=1,378	0.002 [0.038] N=1,181	0.016 [0.047] N=950
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=344	0.020 [0.037] N=1,746	0.018 [0.039] N=1,585	0.010 [0.042] N=1,378	0.023 [0.044] N=1,181	0.032 [0.050] N=950
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=164	0.010 [0.028] N=1,746	0.003 [0.029] N=1,585	-0.001 [0.033] N=1,378	0.011 [0.036] N=1,181	-0.006 [0.039] N=950
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=498	0.055 [0.042] N=1,710	0.038 [0.045] N=1,549	0.049 [0.045] N=1,344	0.016 [0.049] N=1,152	0.055 [0.053] N=926
BMI-for-age z score below -2	0.119 P=274	0.051 [0.032] N=2,288	0.049 [0.035] N=2,066	0.053 [0.038] N=1,815	0.033 [0.039] N=1,543	0.043 [0.045] N=1,255
WHO 2007 BMI overweight above +1, 61+ months	0.402 P=212	-0.025 [0.083] N=538	-0.086 [0.085] N=479	-0.038 [0.087] N=435	0.007 [0.095] N=362	-0.079 [0.106] N=299
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=118	0.036 [0.081] N=538	-0.042 [0.083] N=479	-0.030 [0.090] N=435	0.005 [0.099] N=362	-0.067 [0.112] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.122 P=67	0.054 [0.057] N=538	0.067 [0.061] N=479	0.042 [0.062] N=435	0.057 [0.069] N=362	0.075 [0.079] N=299

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.041 [0.245] N=2,047	-0.109 [0.274] N=1,817	-0.061 [0.302] N=1,553	-0.284 [0.320] N=1,270	-0.301 [0.424] N=970
WHO weight-for-age z score	-0.089 [0.154] N=2,216	-0.081 [0.167] N=1,979	-0.258 [0.182] N=1,687	-0.318 [0.218] N=1,374	-0.353 [0.253] N=1,042
WHO weight-for-length/height z score, 0-60 months	-0.228 [0.246] N=1,513	-0.229 [0.259] N=1,352	-0.454 [0.307] N=1,145	-0.323 [0.333] N=948	**-0.763* [0.435] N=717**
WHO BMI-for-age z score	**-0.471** [0.215] N=1,966**	**-0.595*** [0.225] N=1,744**	**-0.718*** [0.268] N=1,493**	**-0.563* [0.310] N=1,221**	**-1.106*** [0.377] N=933**
Stunted: HAZ below -2	0.021 [0.046] N=2,047	0.018 [0.050] N=1,817	-0.010 [0.054] N=1,553	-0.005 [0.063] N=1,270	0.008 [0.087] N=970
Underweight: WAZ below -2	-0.006 [0.025] N=2,216	-0.008 [0.028] N=1,979	0.014 [0.030] N=1,687	-0.006 [0.034] N=1,374	0.002 [0.039] N=1,042
Wasted: WLZ/WHZ below -2, 0-60 months	**0.082** [0.038] N=1,513**	0.070 [0.043] N=1,352	**0.125** [0.048] N=1,145**	0.063 [0.048] N=948	**0.132** [0.052] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.054] N=1,513	0.002 [0.057] N=1,352	-0.022 [0.064] N=1,145	-0.022 [0.075] N=948	-0.061 [0.094] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.040] N=1,513	-0.010 [0.042] N=1,352	-0.026 [0.050] N=1,145	-0.016 [0.062] N=948	-0.120 [0.078] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.079 [0.064] N=1,483	0.053 [0.072] N=1,322	0.075 [0.080] N=1,117	-0.000 [0.094] N=925	0.080 [0.117] N=699
BMI-for-age z score below -2	**0.107*** [0.031] N=1,966**	**0.114*** [0.037] N=1,744**	**0.136*** [0.042] N=1,493**	**0.113*** [0.038] N=1,221**	**0.179*** [0.048] N=933**
WHO 2007 BMI overweight above +1, 61+ months	-0.123 [0.139] N=449	**-0.263* [0.145] N=390**	-0.194 [0.159] N=346	-0.126 [0.206] N=273	-0.460 [0.296] N=210
WHO 2007 BMI obesity above +2, 61+ months	0.106 [0.120] N=449	-0.029 [0.128] N=390	-0.018 [0.150] N=346	0.070 [0.175] N=273	-0.062 [0.260] N=210
WHO 2007 BMI thinness below -2, 61+ months	0.104 [0.073] N=449	**0.147* [0.081] N=390**	0.096 [0.084] N=346	0.153 [0.096] N=273	**0.261** [0.117] N=210**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.272	-0.211 [0.192] N=2,387	-0.250 [0.202] N=2,157	-0.259 [0.216] N=1,893	**-0.381* [0.228] N=1,610**	-0.429 [0.260] N=1,310
WHO weight-for-age z score	0.183	-0.088 [0.100] N=2,580	-0.081 [0.104] N=2,343	-0.142 [0.111] N=2,051	-0.163 [0.120] N=1,738	-0.158 [0.134] N=1,406
WHO weight-for-length/height z score, 0-60 months	0.366	-0.031 [0.189] N=1,746	-0.008 [0.200] N=1,585	-0.039 [0.224] N=1,378	0.087 [0.237] N=1,181	0.050 [0.284] N=950
WHO BMI-for-age z score	0.356	-0.082 [0.192] N=2,288	-0.107 [0.207] N=2,066	-0.097 [0.226] N=1,815	0.046 [0.245] N=1,543	-0.023 [0.285] N=1,255
Stunted: HAZ below -2	0.215 P=510	0.036 [0.034] N=2,387	0.035 [0.036] N=2,157	0.027 [0.038] N=1,893	0.036 [0.041] N=1,610	0.049 [0.046] N=1,310
Underweight: WAZ below -2	0.053 P=134	0.005 [0.016] N=2,580	0.004 [0.017] N=2,343	0.011 [0.017] N=2,051	0.005 [0.017] N=1,738	0.011 [0.018] N=1,406
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	0.031 [0.033] N=1,746	0.019 [0.035] N=1,585	0.033 [0.038] N=1,378	-0.000 [0.039] N=1,181	0.012 [0.047] N=950
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=344	0.020 [0.037] N=1,746	0.018 [0.039] N=1,585	0.013 [0.042] N=1,378	0.025 [0.044] N=1,181	0.035 [0.050] N=950
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=164	0.012 [0.029] N=1,746	0.005 [0.030] N=1,585	0.001 [0.033] N=1,378	0.012 [0.037] N=1,181	-0.004 [0.040] N=950
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=498	0.066 [0.042] N=1,710	0.045 [0.045] N=1,549	0.054 [0.046] N=1,344	0.024 [0.050] N=1,152	0.059 [0.054] N=926
BMI-for-age z score below -2	0.119 P=274	0.051 [0.032] N=2,288	0.049 [0.035] N=2,066	0.050 [0.038] N=1,815	0.032 [0.039] N=1,543	0.042 [0.046] N=1,255
WHO 2007 BMI overweight above +1, 61+ months	0.402 P=212	-0.048 [0.084] N=538	-0.106 [0.086] N=479	-0.060 [0.088] N=435	-0.019 [0.097] N=362	-0.121 [0.110] N=299
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=118	0.017 [0.080] N=538	-0.055 [0.083] N=479	-0.044 [0.090] N=435	-0.014 [0.096] N=362	-0.086 [0.110] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.122 P=67	0.046 [0.056] N=538	0.053 [0.059] N=479	0.030 [0.060] N=435	0.051 [0.066] N=362	0.060 [0.077] N=299

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.030 [0.245] N=2,047	-0.083 [0.273] N=1,817	-0.013 [0.300] N=1,553	-0.237 [0.317] N=1,270	-0.207 [0.412] N=970
WHO weight-for-age z score	-0.091 [0.152] N=2,216	-0.081 [0.165] N=1,979	-0.233 [0.182] N=1,687	-0.319 [0.217] N=1,374	-0.346 [0.251] N=1,042
WHO weight-for-length/height z score, 0-60 months	-0.247 [0.249] N=1,513	-0.245 [0.263] N=1,352	-0.446 [0.312] N=1,145	-0.322 [0.338] N=948	-0.725 [0.436] N=717
WHO BMI-for-age z score	**-0.479** [0.215] N=1,966**	**-0.603*** [0.227] N=1,744**	**-0.724*** [0.267] N=1,493**	**-0.587* [0.311] N=1,221**	**-1.144*** [0.379] N=933**
Stunted: HAZ below -2	0.018 [0.046] N=2,047	0.012 [0.051] N=1,817	-0.021 [0.054] N=1,553	-0.018 [0.063] N=1,270	-0.014 [0.084] N=970
Underweight: WAZ below -2	-0.007 [0.024] N=2,216	-0.009 [0.028] N=1,979	0.008 [0.029] N=1,687	-0.008 [0.032] N=1,374	-0.001 [0.036] N=1,042
Wasted: WLZ/WHZ below -2, 0-60 months	**0.086** [0.039] N=1,513**	0.071 [0.044] N=1,352	**0.121** [0.049] N=1,145**	0.060 [0.050] N=948	**0.126** [0.052] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.055] N=1,513	0.000 [0.059] N=1,352	-0.020 [0.065] N=1,145	-0.022 [0.078] N=948	-0.058 [0.098] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.041] N=1,513	-0.010 [0.043] N=1,352	-0.025 [0.051] N=1,145	-0.016 [0.064] N=948	-0.115 [0.079] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.064] N=1,483	0.058 [0.073] N=1,322	0.072 [0.082] N=1,117	0.001 [0.096] N=925	0.075 [0.116] N=699
BMI-for-age z score below -2	**0.107*** [0.031] N=1,966**	**0.113*** [0.037] N=1,744**	**0.132*** [0.042] N=1,493**	**0.109*** [0.039] N=1,221**	**0.176*** [0.050] N=933**
WHO 2007 BMI overweight above +1, 61+ months	-0.157 [0.137] N=449	**-0.292** [0.144] N=390**	-0.231 [0.158] N=346	-0.183 [0.208] N=273	**-0.607** [0.290] N=210**
WHO 2007 BMI obesity above +2, 61+ months	0.075 [0.120] N=449	-0.050 [0.127] N=390	-0.040 [0.150] N=346	0.035 [0.175] N=273	-0.130 [0.260] N=210
WHO 2007 BMI thinness below -2, 61+ months	0.092 [0.072] N=449	0.129 [0.081] N=390	0.077 [0.084] N=346	0.138 [0.095] N=273	**0.224* [0.115] N=210**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.272	-0.217 [0.192] N=2,387	-0.254 [0.203] N=2,157	-0.270 [0.218] N=1,893	**-0.385* [0.229] N=1,610**	**-0.442* [0.259] N=1,310**
WHO weight-for-age z score	0.183	-0.092 [0.097] N=2,580	-0.082 [0.102] N=2,343	-0.149 [0.107] N=2,051	-0.158 [0.113] N=1,738	-0.169 [0.124] N=1,406
WHO weight-for-length/height z score, 0-60 months	0.366	-0.022 [0.188] N=1,746	-0.000 [0.199] N=1,585	-0.042 [0.221] N=1,378	0.089 [0.232] N=1,181	0.039 [0.280] N=950
WHO BMI-for-age z score	0.356	-0.084 [0.192] N=2,288	-0.110 [0.206] N=2,066	-0.101 [0.226] N=1,815	0.052 [0.242] N=1,543	-0.016 [0.281] N=1,255
Stunted: HAZ below -2	0.215 P=510	0.037 [0.034] N=2,387	0.036 [0.036] N=2,157	0.029 [0.038] N=1,893	0.038 [0.041] N=1,610	0.050 [0.046] N=1,310
Underweight: WAZ below -2	0.053 P=134	0.005 [0.016] N=2,580	0.003 [0.017] N=2,343	0.012 [0.017] N=2,051	0.004 [0.017] N=1,738	0.011 [0.019] N=1,406
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=192	0.031 [0.032] N=1,746	0.020 [0.035] N=1,585	0.037 [0.037] N=1,378	0.002 [0.038] N=1,181	0.014 [0.047] N=950
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=344	0.021 [0.037] N=1,746	0.018 [0.039] N=1,585	0.013 [0.042] N=1,378	0.025 [0.043] N=1,181	0.030 [0.049] N=950
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=164	0.013 [0.028] N=1,746	0.006 [0.029] N=1,585	0.001 [0.033] N=1,378	0.014 [0.036] N=1,181	-0.004 [0.039] N=950
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=498	0.065 [0.041] N=1,710	0.045 [0.045] N=1,549	0.054 [0.045] N=1,344	0.024 [0.048] N=1,152	0.061 [0.053] N=926
BMI-for-age z score below -2	0.119 P=274	0.050 [0.032] N=2,288	0.048 [0.035] N=2,066	0.050 [0.038] N=1,815	0.031 [0.039] N=1,543	0.037 [0.045] N=1,255
WHO 2007 BMI overweight above +1, 61+ months	0.402 P=212	-0.034 [0.082] N=538	-0.092 [0.085] N=479	-0.036 [0.086] N=435	0.013 [0.093] N=362	-0.075 [0.104] N=299
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=118	0.018 [0.080] N=538	-0.053 [0.083] N=479	-0.036 [0.090] N=435	-0.002 [0.097] N=362	-0.076 [0.112] N=299
WHO 2007 BMI thinness below -2, 61+ months	0.122 P=67	0.048 [0.056] N=538	0.058 [0.059] N=479	0.025 [0.060] N=435	0.044 [0.066] N=362	0.052 [0.074] N=299

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.044 [0.242] N=2,047	-0.091 [0.270] N=1,817	-0.027 [0.301] N=1,553	-0.232 [0.317] N=1,270	-0.263 [0.398] N=970
WHO weight-for-age z score	-0.084 [0.149] N=2,216	-0.066 [0.162] N=1,979	-0.219 [0.177] N=1,687	-0.272 [0.208] N=1,374	-0.320 [0.233] N=1,042
WHO weight-for-length/height z score, 0-60 months	-0.219 [0.245] N=1,513	-0.222 [0.257] N=1,352	-0.446 [0.302] N=1,145	-0.313 [0.325] N=948	-0.730 [0.438] N=717
WHO BMI-for-age z score	**-0.477** [0.214] N=1,966**	**-0.604*** [0.225] N=1,744**	**-0.725*** [0.264] N=1,493**	**-0.570* [0.302] N=1,221**	**-1.107*** [0.376] N=933**
Stunted: HAZ below -2	0.023 [0.045] N=2,047	0.016 [0.050] N=1,817	-0.018 [0.054] N=1,553	-0.017 [0.063] N=1,270	0.001 [0.086] N=970
Underweight: WAZ below -2	-0.007 [0.024] N=2,216	-0.010 [0.028] N=1,979	0.008 [0.030] N=1,687	-0.013 [0.032] N=1,374	-0.004 [0.036] N=1,042
Wasted: WLZ/WHZ below -2, 0-60 months	**0.084** [0.038] N=1,513**	**0.072* [0.043] N=1,352**	**0.127*** [0.047] N=1,145**	0.066 [0.048] N=948	**0.136** [0.052] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.054] N=1,513	0.004 [0.057] N=1,352	-0.018 [0.063] N=1,145	-0.019 [0.075] N=948	-0.054 [0.095] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.011 [0.040] N=1,513	-0.008 [0.041] N=1,352	-0.025 [0.050] N=1,145	-0.016 [0.062] N=948	-0.117 [0.078] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.094 [0.062] N=1,483	0.060 [0.071] N=1,322	0.072 [0.080] N=1,117	0.000 [0.091] N=925	0.096 [0.110] N=699
BMI-for-age z score below -2	**0.110*** [0.031] N=1,966**	**0.115*** [0.037] N=1,744**	**0.136*** [0.042] N=1,493**	**0.113*** [0.038] N=1,221**	**0.180*** [0.048] N=933**
WHO 2007 BMI overweight above +1, 61+ months	-0.152 [0.134] N=449	**-0.286** [0.141] N=390**	-0.210 [0.151] N=346	-0.152 [0.195] N=273	**-0.496* [0.272] N=210**
WHO 2007 BMI obesity above +2, 61+ months	0.074 [0.119] N=449	-0.054 [0.127] N=390	-0.042 [0.149] N=346	0.038 [0.175] N=273	-0.097 [0.261] N=210
WHO 2007 BMI thinness below -2, 61+ months	0.113 [0.071] N=449	**0.153* [0.080] N=390**	0.090 [0.084] N=346	0.147 [0.095] N=273	**0.243** [0.121] N=210**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.280	-0.218 [0.193] N=2,237	-0.254 [0.207] N=2,021	-0.257 [0.219] N=1,777	-0.376 [0.231] N=1,507	**-0.500* [0.265] N=1,229**
WHO weight-for-age z score	0.183	-0.079 [0.103] N=2,412	-0.064 [0.107] N=2,189	-0.141 [0.114] N=1,920	-0.156 [0.124] N=1,625	-0.208 [0.132] N=1,319
WHO weight-for-length/height z score, 0-60 months	0.372	0.035 [0.200] N=1,626	0.070 [0.212] N=1,476	-0.014 [0.234] N=1,284	0.098 [0.247] N=1,100	0.035 [0.292] N=888
WHO BMI-for-age z score	0.372	-0.098 [0.199] N=2,144	-0.120 [0.213] N=1,936	-0.151 [0.235] N=1,704	-0.017 [0.250] N=1,446	-0.111 [0.287] N=1,180
Stunted: HAZ below -2	0.217 P=482	0.042 [0.034] N=2,237	0.044 [0.036] N=2,021	0.036 [0.037] N=1,777	0.045 [0.040] N=1,507	0.072 [0.045] N=1,229
Underweight: WAZ below -2	0.053 P=127	0.003 [0.017] N=2,412	0.001 [0.018] N=2,189	0.007 [0.018] N=1,920	0.003 [0.019] N=1,625	0.015 [0.021] N=1,319
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=180	0.030 [0.033] N=1,626	0.016 [0.036] N=1,476	0.034 [0.038] N=1,284	0.005 [0.039] N=1,100	0.021 [0.048] N=888
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=322	0.018 [0.038] N=1,626	0.018 [0.040] N=1,476	0.008 [0.042] N=1,284	0.018 [0.044] N=1,100	0.024 [0.050] N=888
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=156	0.014 [0.028] N=1,626	0.008 [0.029] N=1,476	0.004 [0.032] N=1,284	0.019 [0.036] N=1,100	0.009 [0.040] N=888
CIAF: any stunting, underweight, or wasting, 0-60 months	0.298 P=469	0.073 [0.045] N=1,592	0.056 [0.048] N=1,442	0.067 [0.048] N=1,252	0.035 [0.052] N=1,072	0.092 [0.056] N=864
BMI-for-age z score below -2	0.117 P=251	**0.056* [0.032] N=2,144**	0.053 [0.035] N=1,936	0.059 [0.037] N=1,704	0.045 [0.037] N=1,446	0.059 [0.042] N=1,180
WHO 2007 BMI overweight above +1, 61+ months	0.408 P=206	-0.061 [0.088] N=516	-0.136 [0.090] N=460	-0.075 [0.092] N=419	-0.032 [0.103] N=347	-0.141 [0.113] N=287
WHO 2007 BMI obesity above +2, 61+ months	0.228 P=114	0.010 [0.086] N=516	-0.078 [0.089] N=460	-0.064 [0.097] N=419	-0.038 [0.107] N=347	-0.117 [0.120] N=287
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=62	0.067 [0.060] N=516	0.082 [0.064] N=460	0.054 [0.066] N=419	0.064 [0.072] N=347	0.086 [0.079] N=287

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.005 [0.250] N=1,918	-0.015 [0.283] N=1,702	0.043 [0.318] N=1,458	-0.134 [0.336] N=1,188	-0.271 [0.436] N=910
WHO weight-for-age z score	-0.084 [0.160] N=2,072	-0.050 [0.172] N=1,849	-0.253 [0.190] N=1,580	-0.295 [0.236] N=1,285	-0.378 [0.279] N=979
WHO weight-for-length/height z score, 0-60 months	-0.175 [0.266] N=1,406	-0.174 [0.282] N=1,256	-0.477 [0.336] N=1,064	-0.375 [0.367] N=880	**-0.872* [0.479] N=668**
WHO BMI-for-age z score	**-0.456** [0.228] N=1,840**	**-0.580** [0.242] N=1,632**	**-0.762** [0.292] N=1,400**	**-0.618* [0.340] N=1,142**	**-1.218*** [0.415] N=876**
Stunted: HAZ below -2	0.012 [0.049] N=1,918	0.009 [0.055] N=1,702	-0.024 [0.058] N=1,458	-0.029 [0.066] N=1,188	0.000 [0.091] N=910
Underweight: WAZ below -2	-0.008 [0.025] N=2,072	-0.012 [0.029] N=1,849	0.005 [0.030] N=1,580	-0.010 [0.036] N=1,285	0.001 [0.042] N=979
Wasted: WLZ/WHZ below -2, 0-60 months	**0.078* [0.041] N=1,406**	0.062 [0.045] N=1,256	**0.119** [0.050] N=1,064**	0.057 [0.052] N=880	**0.130** [0.058] N=668**
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.056] N=1,406	-0.013 [0.060] N=1,256	-0.042 [0.067] N=1,064	-0.053 [0.082] N=880	-0.093 [0.105] N=668
Obese: WLZ/WHZ above +3, 0-60 months	-0.003 [0.043] N=1,406	-0.023 [0.044] N=1,256	-0.043 [0.051] N=1,064	-0.034 [0.065] N=880	-0.130 [0.084] N=668
CIAF: any stunting, underweight, or wasting, 0-60 months	0.078 [0.071] N=1,378	0.047 [0.080] N=1,228	0.066 [0.089] N=1,038	-0.030 [0.106] N=858	0.076 [0.134] N=650
BMI-for-age z score below -2	**0.107*** [0.034] N=1,840**	**0.113*** [0.040] N=1,632**	**0.138*** [0.046] N=1,400**	**0.122*** [0.043] N=1,142**	**0.193*** [0.053] N=876**
WHO 2007 BMI overweight above +1, 61+ months	-0.143 [0.144] N=432	**-0.295** [0.149] N=376**	-0.209 [0.164] N=335	-0.124 [0.212] N=263	**-0.510* [0.303] N=203**
WHO 2007 BMI obesity above +2, 61+ months	0.100 [0.126] N=432	-0.038 [0.135] N=376	-0.024 [0.157] N=335	0.086 [0.183] N=263	-0.042 [0.280] N=203
WHO 2007 BMI thinness below -2, 61+ months	0.100 [0.076] N=432	**0.145* [0.085] N=376**	0.093 [0.089] N=335	0.142 [0.102] N=263	**0.230* [0.123] N=203**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.280	-0.222 [0.193] N=2,237	-0.260 [0.207] N=2,021	-0.261 [0.220] N=1,777	-0.375 [0.232] N=1,507	**-0.495* [0.265] N=1,229**
WHO weight-for-age z score	0.183	-0.080 [0.104] N=2,412	-0.066 [0.108] N=2,189	-0.148 [0.115] N=1,920	-0.159 [0.125] N=1,625	-0.207 [0.136] N=1,319
WHO weight-for-length/height z score, 0-60 months	0.372	0.042 [0.201] N=1,626	0.076 [0.213] N=1,476	-0.003 [0.236] N=1,284	0.118 [0.250] N=1,100	0.049 [0.300] N=888
WHO BMI-for-age z score	0.372	-0.095 [0.199] N=2,144	-0.119 [0.214] N=1,936	-0.146 [0.236] N=1,704	-0.012 [0.252] N=1,446	-0.111 [0.289] N=1,180
Stunted: HAZ below -2	0.217 P=482	0.041 [0.034] N=2,237	0.044 [0.036] N=2,021	0.035 [0.037] N=1,777	0.044 [0.040] N=1,507	0.070 [0.045] N=1,229
Underweight: WAZ below -2	0.053 P=127	0.003 [0.017] N=2,412	0.002 [0.019] N=2,189	0.009 [0.019] N=1,920	0.003 [0.019] N=1,625	0.015 [0.022] N=1,319
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=180	0.027 [0.033] N=1,626	0.013 [0.036] N=1,476	0.030 [0.038] N=1,284	-0.000 [0.039] N=1,100	0.015 [0.047] N=888
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=322	0.018 [0.038] N=1,626	0.017 [0.040] N=1,476	0.008 [0.042] N=1,284	0.019 [0.044] N=1,100	0.024 [0.051] N=888
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=156	0.014 [0.028] N=1,626	0.008 [0.029] N=1,476	0.002 [0.032] N=1,284	0.019 [0.036] N=1,100	0.007 [0.040] N=888
CIAF: any stunting, underweight, or wasting, 0-60 months	0.298 P=469	0.067 [0.045] N=1,592	0.048 [0.048] N=1,442	0.060 [0.048] N=1,252	0.028 [0.051] N=1,072	0.082 [0.055] N=864
BMI-for-age z score below -2	0.117 P=251	**0.055* [0.032] N=2,144**	0.052 [0.035] N=1,936	0.057 [0.037] N=1,704	0.043 [0.037] N=1,446	0.058 [0.042] N=1,180
WHO 2007 BMI overweight above +1, 61+ months	0.408 P=206	-0.064 [0.088] N=516	-0.135 [0.090] N=460	-0.079 [0.092] N=419	-0.038 [0.101] N=347	-0.134 [0.111] N=287
WHO 2007 BMI obesity above +2, 61+ months	0.228 P=114	0.008 [0.086] N=516	-0.077 [0.089] N=460	-0.065 [0.096] N=419	-0.039 [0.106] N=347	-0.117 [0.119] N=287
WHO 2007 BMI thinness below -2, 61+ months	0.118 P=62	0.069 [0.058] N=516	0.085 [0.062] N=460	0.053 [0.063] N=419	0.066 [0.068] N=347	0.090 [0.073] N=287

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.014 [0.248] N=1,918	-0.038 [0.278] N=1,702	0.027 [0.312] N=1,458	-0.179 [0.337] N=1,188	-0.329 [0.432] N=910
WHO weight-for-age z score	-0.084 [0.160] N=2,072	-0.053 [0.172] N=1,849	-0.263 [0.190] N=1,580	-0.304 [0.235] N=1,285	-0.415 [0.278] N=979
WHO weight-for-length/height z score, 0-60 months	-0.170 [0.265] N=1,406	-0.166 [0.281] N=1,256	-0.468 [0.335] N=1,064	-0.328 [0.374] N=880	**-0.860* [0.480] N=668**
WHO BMI-for-age z score	**-0.448* [0.228] N=1,840**	**-0.575** [0.242] N=1,632**	**-0.755** [0.290] N=1,400**	**-0.593* [0.343] N=1,142**	**-1.216*** [0.414] N=876**
Stunted: HAZ below -2	0.013 [0.049] N=1,918	0.011 [0.054] N=1,702	-0.022 [0.058] N=1,458	-0.022 [0.067] N=1,188	0.011 [0.090] N=910
Underweight: WAZ below -2	-0.008 [0.025] N=2,072	-0.011 [0.028] N=1,849	0.008 [0.030] N=1,580	-0.006 [0.035] N=1,285	0.012 [0.042] N=979
Wasted: WLZ/WHZ below -2, 0-60 months	**0.077* [0.041] N=1,406**	0.060 [0.045] N=1,256	**0.117** [0.049] N=1,064**	0.051 [0.051] N=880	**0.130** [0.056] N=668**
Overweight: WLZ/WHZ above +2, 0-60 months	0.002 [0.056] N=1,406	-0.011 [0.059] N=1,256	-0.040 [0.067] N=1,064	-0.046 [0.083] N=880	-0.092 [0.104] N=668
Obese: WLZ/WHZ above +3, 0-60 months	-0.002 [0.043] N=1,406	-0.021 [0.044] N=1,256	-0.043 [0.051] N=1,064	-0.028 [0.066] N=880	-0.127 [0.083] N=668
CIAF: any stunting, underweight, or wasting, 0-60 months	0.077 [0.070] N=1,378	0.046 [0.080] N=1,228	0.066 [0.089] N=1,038	-0.025 [0.105] N=858	0.087 [0.134] N=650
BMI-for-age z score below -2	**0.107*** [0.034] N=1,840**	**0.113*** [0.040] N=1,632**	**0.138*** [0.046] N=1,400**	**0.119*** [0.042] N=1,142**	**0.196*** [0.052] N=876**
WHO 2007 BMI overweight above +1, 61+ months	-0.140 [0.143] N=432	**-0.279* [0.149] N=376**	-0.204 [0.165] N=335	-0.118 [0.215] N=263	-0.460 [0.308] N=203
WHO 2007 BMI obesity above +2, 61+ months	0.101 [0.124] N=432	-0.034 [0.134] N=376	-0.021 [0.156] N=335	0.084 [0.181] N=263	-0.050 [0.274] N=203
WHO 2007 BMI thinness below -2, 61+ months	0.105 [0.075] N=432	**0.150* [0.082] N=376**	0.088 [0.086] N=335	0.139 [0.097] N=263	**0.231** [0.115] N=203**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 71 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.