Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-5 Using 10 October 2022 Child Age

The child cohort uses complete birth dates and completed ages 0 to 71 months on 10 October 2022 without age-boundary assignments. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications are available through month 60 only. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.084	**0.328* [0.177] N=2,434**	**0.375** [0.187] N=2,255**	**0.380* [0.200] N=2,031**	**0.501** [0.208] N=1,750**	0.350 [0.238] N=1,440
WHO weight-for-age z score	0.299	-0.052 [0.100] N=2,715	-0.022 [0.109] N=2,515	0.025 [0.110] N=2,266	0.008 [0.120] N=1,946	-0.011 [0.125] N=1,600
WHO weight-for-length/height z score, 0-60 months	0.351	0.038 [0.155] N=2,055	0.031 [0.158] N=1,908	0.039 [0.168] N=1,720	-0.153 [0.185] N=1,474	-0.082 [0.197] N=1,208
WHO BMI-for-age z score	0.360	-0.022 [0.154] N=2,354	0.008 [0.156] N=2,186	0.041 [0.167] N=1,964	-0.206 [0.173] N=1,688	-0.111 [0.187] N=1,389
Stunted: HAZ below -2	0.194 P=460	-0.041 [0.033] N=2,434	-0.045 [0.034] N=2,255	-0.042 [0.035] N=2,031	**-0.064* [0.037] N=1,750**	-0.034 [0.042] N=1,440
Underweight: WAZ below -2	0.054 P=148	0.011 [0.018] N=2,715	0.005 [0.019] N=2,515	-0.003 [0.020] N=2,266	-0.012 [0.023] N=1,946	-0.016 [0.028] N=1,600
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=230	-0.025 [0.026] N=2,055	-0.021 [0.027] N=1,908	-0.025 [0.029] N=1,720	-0.006 [0.031] N=1,474	-0.027 [0.032] N=1,208
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=403	0.010 [0.036] N=2,055	0.006 [0.038] N=1,908	0.010 [0.040] N=1,720	-0.016 [0.045] N=1,474	-0.022 [0.052] N=1,208
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=191	-0.008 [0.028] N=2,055	-0.011 [0.030] N=1,908	0.004 [0.032] N=1,720	-0.013 [0.036] N=1,474	-0.014 [0.042] N=1,208
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=513	-0.044 [0.037] N=1,987	-0.037 [0.039] N=1,845	-0.032 [0.040] N=1,662	-0.053 [0.042] N=1,424	-0.037 [0.046] N=1,167
BMI-for-age z score below -2	0.117 P=277	-0.004 [0.026] N=2,354	-0.009 [0.026] N=2,186	-0.007 [0.029] N=1,964	0.023 [0.030] N=1,688	0.021 [0.032] N=1,389
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=121	-0.070 [0.114] N=298	-0.079 [0.121] N=276	-0.059 [0.130] N=248	-0.085 [0.140] N=222	-0.048 [0.161] N=184
WHO 2007 BMI obesity above +2, 61+ months	0.231 P=69	-0.094 [0.079] N=298	-0.119 [0.082] N=276	-0.101 [0.089] N=248	-0.139 [0.094] N=222	-0.119 [0.115] N=184
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=32	0.012 [0.064] N=298	-0.002 [0.064] N=276	-0.025 [0.064] N=248	-0.040 [0.072] N=222	-0.092 [0.093] N=184

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.134 [0.247] N=2,042	0.173 [0.273] N=1,863	0.159 [0.304] N=1,639	0.435 [0.325] N=1,358	-0.054 [0.442] N=1,048
WHO weight-for-age z score	-0.224 [0.143] N=2,287	-0.200 [0.164] N=2,087	-0.140 [0.171] N=1,838	-0.231 [0.208] N=1,518	-0.352 [0.261] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.052 [0.222] N=1,731	0.051 [0.235] N=1,584	0.027 [0.266] N=1,396	-0.369 [0.325] N=1,150	-0.326 [0.410] N=884
WHO BMI-for-age z score	0.042 [0.230] N=1,975	0.117 [0.240] N=1,807	0.172 [0.279] N=1,585	-0.311 [0.327] N=1,309	-0.128 [0.420] N=1,010
Stunted: HAZ below -2	0.015 [0.047] N=2,042	0.020 [0.051] N=1,863	0.036 [0.054] N=1,639	0.006 [0.064] N=1,358	0.141 [0.087] N=1,048
Underweight: WAZ below -2	0.033 [0.024] N=2,287	0.026 [0.026] N=2,087	0.016 [0.028] N=1,838	0.002 [0.034] N=1,518	0.003 [0.048] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.023 [0.036] N=1,731	-0.017 [0.039] N=1,584	-0.020 [0.044] N=1,396	0.013 [0.055] N=1,150	-0.035 [0.063] N=884
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.044] N=1,731	0.005 [0.047] N=1,584	0.006 [0.052] N=1,396	-0.053 [0.065] N=1,150	-0.105 [0.088] N=884
Obese: WLZ/WHZ above +3, 0-60 months	-0.018 [0.030] N=1,731	-0.022 [0.034] N=1,584	0.004 [0.038] N=1,396	-0.036 [0.045] N=1,150	-0.066 [0.064] N=884
CIAF: any stunting, underweight, or wasting, 0-60 months	0.012 [0.050] N=1,672	0.033 [0.055] N=1,530	0.055 [0.057] N=1,347	0.021 [0.068] N=1,109	0.115 [0.082] N=852
BMI-for-age z score below -2	-0.033 [0.034] N=1,975	-0.051 [0.034] N=1,807	-0.055 [0.040] N=1,585	-0.015 [0.045] N=1,309	-0.038 [0.054] N=1,010
WHO 2007 BMI overweight above +1, 61+ months	-0.111 [0.170] N=240	-0.124 [0.181] N=218	-0.080 [0.198] N=190	-0.156 [0.219] N=164	-0.099 [0.281] N=126
WHO 2007 BMI obesity above +2, 61+ months	-0.056 [0.126] N=240	-0.097 [0.130] N=218	-0.051 [0.152] N=190	-0.134 [0.171] N=164	-0.114 [0.209] N=126
WHO 2007 BMI thinness below -2, 61+ months	-0.006 [0.094] N=240	-0.047 [0.096] N=218	-0.112 [0.098] N=190	-0.174 [0.118] N=164	**-0.356** [0.164] N=126**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.084	**0.329* [0.178] N=2,434**	**0.376** [0.188] N=2,255**	**0.382* [0.201] N=2,031**	**0.508** [0.207] N=1,750**	0.357 [0.237] N=1,440
WHO weight-for-age z score	0.299	-0.054 [0.100] N=2,715	-0.025 [0.109] N=2,515	0.023 [0.110] N=2,266	0.008 [0.119] N=1,946	-0.012 [0.125] N=1,600
WHO weight-for-length/height z score, 0-60 months	0.351	0.038 [0.155] N=2,055	0.031 [0.159] N=1,908	0.037 [0.169] N=1,720	-0.153 [0.186] N=1,474	-0.090 [0.198] N=1,208
WHO BMI-for-age z score	0.360	-0.020 [0.154] N=2,354	0.008 [0.157] N=2,186	0.041 [0.167] N=1,964	-0.203 [0.174] N=1,688	-0.112 [0.187] N=1,389
Stunted: HAZ below -2	0.194 P=460	-0.041 [0.033] N=2,434	-0.044 [0.034] N=2,255	-0.042 [0.036] N=2,031	**-0.064* [0.037] N=1,750**	-0.034 [0.042] N=1,440
Underweight: WAZ below -2	0.054 P=148	0.012 [0.018] N=2,715	0.005 [0.019] N=2,515	-0.002 [0.020] N=2,266	-0.013 [0.023] N=1,946	-0.016 [0.028] N=1,600
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=230	-0.023 [0.026] N=2,055	-0.018 [0.027] N=1,908	-0.021 [0.029] N=1,720	-0.002 [0.031] N=1,474	-0.021 [0.032] N=1,208
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=403	0.011 [0.036] N=2,055	0.008 [0.038] N=1,908	0.013 [0.040] N=1,720	-0.012 [0.045] N=1,474	-0.018 [0.051] N=1,208
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=191	-0.007 [0.028] N=2,055	-0.008 [0.030] N=1,908	0.007 [0.032] N=1,720	-0.010 [0.035] N=1,474	-0.010 [0.041] N=1,208
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=513	-0.041 [0.037] N=1,987	-0.033 [0.038] N=1,845	-0.027 [0.039] N=1,662	-0.048 [0.042] N=1,424	-0.030 [0.045] N=1,167
BMI-for-age z score below -2	0.117 P=277	-0.004 [0.026] N=2,354	-0.008 [0.026] N=2,186	-0.006 [0.029] N=1,964	0.025 [0.030] N=1,688	0.022 [0.032] N=1,389
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=121	-0.075 [0.114] N=298	-0.084 [0.121] N=276	-0.068 [0.130] N=248	-0.093 [0.140] N=222	-0.046 [0.164] N=184
WHO 2007 BMI obesity above +2, 61+ months	0.231 P=69	-0.102 [0.080] N=298	-0.126 [0.082] N=276	-0.115 [0.090] N=248	-0.153 [0.094] N=222	-0.131 [0.113] N=184
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=32	0.020 [0.065] N=298	0.004 [0.064] N=276	-0.014 [0.063] N=248	-0.028 [0.071] N=222	-0.070 [0.092] N=184

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.131 [0.246] N=2,042	0.171 [0.272] N=1,863	0.154 [0.301] N=1,639	0.419 [0.323] N=1,358	-0.112 [0.438] N=1,048
WHO weight-for-age z score	-0.224 [0.143] N=2,287	-0.200 [0.164] N=2,087	-0.140 [0.171] N=1,838	-0.232 [0.208] N=1,518	-0.353 [0.260] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.052 [0.222] N=1,731	0.050 [0.236] N=1,584	0.025 [0.266] N=1,396	-0.370 [0.326] N=1,150	-0.330 [0.411] N=884
WHO BMI-for-age z score	0.042 [0.231] N=1,975	0.117 [0.240] N=1,807	0.173 [0.278] N=1,585	-0.311 [0.327] N=1,309	-0.115 [0.421] N=1,010
Stunted: HAZ below -2	0.015 [0.047] N=2,042	0.020 [0.051] N=1,863	0.036 [0.054] N=1,639	0.007 [0.064] N=1,358	**0.148* [0.087] N=1,048**
Underweight: WAZ below -2	0.033 [0.024] N=2,287	0.026 [0.026] N=2,087	0.016 [0.028] N=1,838	0.002 [0.034] N=1,518	0.003 [0.048] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.023 [0.037] N=1,731	-0.016 [0.039] N=1,584	-0.018 [0.044] N=1,396	0.014 [0.055] N=1,150	-0.034 [0.064] N=884
Overweight: WLZ/WHZ above +2, 0-60 months	0.010 [0.044] N=1,731	0.006 [0.047] N=1,584	0.007 [0.052] N=1,396	-0.052 [0.064] N=1,150	-0.104 [0.087] N=884
Obese: WLZ/WHZ above +3, 0-60 months	-0.018 [0.030] N=1,731	-0.022 [0.034] N=1,584	0.005 [0.038] N=1,396	-0.035 [0.045] N=1,150	-0.065 [0.064] N=884
CIAF: any stunting, underweight, or wasting, 0-60 months	0.012 [0.049] N=1,672	0.033 [0.054] N=1,530	0.055 [0.057] N=1,347	0.020 [0.068] N=1,109	0.114 [0.082] N=852
BMI-for-age z score below -2	-0.033 [0.034] N=1,975	-0.051 [0.034] N=1,807	-0.055 [0.040] N=1,585	-0.016 [0.045] N=1,309	-0.041 [0.054] N=1,010
WHO 2007 BMI overweight above +1, 61+ months	-0.116 [0.173] N=240	-0.130 [0.185] N=218	-0.090 [0.204] N=190	-0.171 [0.226] N=164	-0.027 [0.290] N=126
WHO 2007 BMI obesity above +2, 61+ months	-0.068 [0.128] N=240	-0.109 [0.132] N=218	-0.083 [0.156] N=190	-0.180 [0.175] N=164	-0.134 [0.209] N=126
WHO 2007 BMI thinness below -2, 61+ months	0.007 [0.096] N=240	-0.036 [0.099] N=218	-0.084 [0.100] N=190	-0.129 [0.121] N=164	-0.269 [0.164] N=126

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	**Mean**	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO length/height-for-age z score	-0.084	**0.303* [0.179] N=2,434**	**0.347* [0.188] N=2,255**	**0.350* [0.203] N=2,031**	**0.478** [0.210] N=1,750**	0.307 [0.236] N=1,440
WHO weight-for-age z score	0.299	-0.092 [0.097] N=2,715	-0.060 [0.106] N=2,515	-0.016 [0.107] N=2,266	-0.023 [0.117] N=1,946	-0.050 [0.123] N=1,600
WHO weight-for-length/height z score, 0-60 months	0.351	0.033 [0.158] N=2,055	0.031 [0.162] N=1,908	0.043 [0.171] N=1,720	-0.146 [0.188] N=1,474	-0.087 [0.201] N=1,208
WHO BMI-for-age z score	0.360	-0.021 [0.155] N=2,354	0.011 [0.158] N=2,186	0.047 [0.168] N=1,964	-0.193 [0.174] N=1,688	-0.101 [0.190] N=1,389
Stunted: HAZ below -2	0.194 P=460	-0.036 [0.033] N=2,434	-0.038 [0.034] N=2,255	-0.036 [0.036] N=2,031	-0.059 [0.038] N=1,750	-0.026 [0.042] N=1,440
Underweight: WAZ below -2	0.054 P=148	0.017 [0.018] N=2,715	0.011 [0.019] N=2,515	0.004 [0.020] N=2,266	-0.006 [0.022] N=1,946	-0.007 [0.027] N=1,600
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=230	-0.019 [0.026] N=2,055	-0.013 [0.027] N=1,908	-0.017 [0.029] N=1,720	0.001 [0.031] N=1,474	-0.017 [0.032] N=1,208
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=403	0.015 [0.036] N=2,055	0.013 [0.038] N=1,908	0.017 [0.040] N=1,720	-0.008 [0.045] N=1,474	-0.012 [0.052] N=1,208
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=191	-0.005 [0.028] N=2,055	-0.006 [0.030] N=1,908	0.009 [0.032] N=1,720	-0.008 [0.035] N=1,474	-0.009 [0.041] N=1,208
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=513	-0.031 [0.037] N=1,987	-0.020 [0.038] N=1,845	-0.018 [0.039] N=1,662	-0.041 [0.042] N=1,424	-0.019 [0.046] N=1,167
BMI-for-age z score below -2	0.117 P=277	-0.000 [0.026] N=2,354	-0.004 [0.026] N=2,186	-0.002 [0.029] N=1,964	0.028 [0.030] N=1,688	0.026 [0.032] N=1,389
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=121	-0.071 [0.115] N=298	-0.075 [0.122] N=276	-0.056 [0.131] N=248	-0.081 [0.143] N=222	-0.010 [0.174] N=184
WHO 2007 BMI obesity above +2, 61+ months	0.231 P=69	-0.093 [0.081] N=298	-0.113 [0.083] N=276	-0.095 [0.089] N=248	-0.134 [0.094] N=222	-0.093 [0.117] N=184
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=32	0.033 [0.064] N=298	0.017 [0.064] N=276	0.003 [0.064] N=248	-0.006 [0.071] N=222	-0.052 [0.091] N=184

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.110 [0.246] N=2,042	0.139 [0.271] N=1,863	0.118 [0.301] N=1,639	0.378 [0.324] N=1,358	-0.167 [0.426] N=1,048
WHO weight-for-age z score	**-0.257* [0.139] N=2,287**	-0.231 [0.160] N=2,087	-0.173 [0.165] N=1,838	-0.254 [0.203] N=1,518	-0.388 [0.255] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.046 [0.225] N=1,731	0.052 [0.241] N=1,584	0.035 [0.268] N=1,396	-0.371 [0.328] N=1,150	-0.349 [0.415] N=884
WHO BMI-for-age z score	0.043 [0.232] N=1,975	0.123 [0.243] N=1,807	0.186 [0.278] N=1,585	-0.293 [0.327] N=1,309	-0.103 [0.422] N=1,010
Stunted: HAZ below -2	0.019 [0.047] N=2,042	0.027 [0.051] N=1,863	0.042 [0.054] N=1,639	0.011 [0.064] N=1,358	**0.154* [0.085] N=1,048**
Underweight: WAZ below -2	0.037 [0.023] N=2,287	0.031 [0.026] N=2,087	0.021 [0.028] N=1,838	0.006 [0.034] N=1,518	0.010 [0.048] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.037] N=1,731	-0.010 [0.040] N=1,584	-0.015 [0.045] N=1,396	0.018 [0.056] N=1,150	-0.026 [0.067] N=884
Overweight: WLZ/WHZ above +2, 0-60 months	0.014 [0.045] N=1,731	0.012 [0.048] N=1,584	0.011 [0.053] N=1,396	-0.051 [0.065] N=1,150	-0.095 [0.087] N=884
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.030] N=1,731	-0.019 [0.035] N=1,584	0.006 [0.038] N=1,396	-0.036 [0.045] N=1,150	-0.068 [0.063] N=884
CIAF: any stunting, underweight, or wasting, 0-60 months	0.023 [0.049] N=1,672	0.048 [0.054] N=1,530	0.062 [0.056] N=1,347	0.020 [0.067] N=1,109	0.122 [0.082] N=852
BMI-for-age z score below -2	-0.029 [0.034] N=1,975	-0.047 [0.034] N=1,807	-0.052 [0.041] N=1,585	-0.013 [0.045] N=1,309	-0.035 [0.056] N=1,010
WHO 2007 BMI overweight above +1, 61+ months	-0.123 [0.171] N=240	-0.136 [0.186] N=218	-0.097 [0.204] N=190	-0.192 [0.225] N=164	0.007 [0.292] N=126
WHO 2007 BMI obesity above +2, 61+ months	-0.062 [0.126] N=240	-0.098 [0.133] N=218	-0.069 [0.152] N=190	-0.175 [0.175] N=164	-0.078 [0.220] N=126
WHO 2007 BMI thinness below -2, 61+ months	0.025 [0.094] N=240	-0.022 [0.099] N=218	-0.065 [0.103] N=190	-0.092 [0.118] N=164	-0.259 [0.159] N=126

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.084	0.274 [0.176] N=2,434	**0.320* [0.186] N=2,255**	0.320 [0.200] N=2,031	**0.431** [0.207] N=1,750**	0.255 [0.232] N=1,440
WHO weight-for-age z score	0.299	-0.093 [0.097] N=2,715	-0.069 [0.106] N=2,515	-0.027 [0.108] N=2,266	-0.044 [0.119] N=1,946	-0.083 [0.126] N=1,600
WHO weight-for-length/height z score, 0-60 months	0.351	0.049 [0.157] N=2,055	0.040 [0.161] N=1,908	0.047 [0.171] N=1,720	-0.142 [0.188] N=1,474	-0.092 [0.200] N=1,208
WHO BMI-for-age z score	0.360	-0.005 [0.156] N=2,354	0.022 [0.158] N=2,186	0.059 [0.169] N=1,964	-0.184 [0.178] N=1,688	-0.095 [0.192] N=1,389
Stunted: HAZ below -2	0.194 P=460	-0.030 [0.033] N=2,434	-0.032 [0.034] N=2,255	-0.028 [0.036] N=2,031	-0.047 [0.037] N=1,750	-0.014 [0.041] N=1,440
Underweight: WAZ below -2	0.054 P=148	0.018 [0.018] N=2,715	0.013 [0.019] N=2,515	0.008 [0.020] N=2,266	-0.002 [0.022] N=1,946	-0.002 [0.027] N=1,600
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=230	-0.021 [0.026] N=2,055	-0.015 [0.027] N=1,908	-0.017 [0.029] N=1,720	0.002 [0.031] N=1,474	-0.018 [0.032] N=1,208
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=403	0.018 [0.036] N=2,055	0.016 [0.038] N=1,908	0.019 [0.040] N=1,720	-0.006 [0.045] N=1,474	-0.015 [0.051] N=1,208
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=191	-0.005 [0.028] N=2,055	-0.008 [0.030] N=1,908	0.008 [0.032] N=1,720	-0.008 [0.035] N=1,474	-0.011 [0.041] N=1,208
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=513	-0.024 [0.037] N=1,987	-0.011 [0.038] N=1,845	-0.006 [0.039] N=1,662	-0.027 [0.042] N=1,424	-0.004 [0.045] N=1,167
BMI-for-age z score below -2	0.117 P=277	-0.002 [0.026] N=2,354	-0.006 [0.026] N=2,186	-0.004 [0.029] N=1,964	0.028 [0.030] N=1,688	0.024 [0.033] N=1,389
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=121	-0.063 [0.111] N=298	-0.075 [0.120] N=276	-0.058 [0.129] N=248	-0.068 [0.136] N=222	-0.002 [0.161] N=184
WHO 2007 BMI obesity above +2, 61+ months	0.231 P=69	-0.088 [0.081] N=298	-0.118 [0.083] N=276	-0.106 [0.090] N=248	-0.138 [0.096] N=222	-0.111 [0.115] N=184
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=32	0.023 [0.065] N=298	0.013 [0.066] N=276	-0.015 [0.065] N=248	-0.033 [0.073] N=222	-0.058 [0.091] N=184

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.096 [0.243] N=2,042	0.136 [0.267] N=1,863	0.118 [0.297] N=1,639	0.389 [0.322] N=1,358	-0.189 [0.414] N=1,048
WHO weight-for-age z score	**-0.242* [0.140] N=2,287**	-0.226 [0.161] N=2,087	-0.174 [0.168] N=1,838	-0.250 [0.207] N=1,518	-0.378 [0.258] N=1,172
WHO weight-for-length/height z score, 0-60 months	0.054 [0.224] N=1,731	0.056 [0.239] N=1,584	0.032 [0.270] N=1,396	-0.365 [0.331] N=1,150	-0.324 [0.416] N=884
WHO BMI-for-age z score	0.049 [0.233] N=1,975	0.122 [0.242] N=1,807	0.180 [0.281] N=1,585	-0.294 [0.330] N=1,309	-0.095 [0.422] N=1,010
Stunted: HAZ below -2	0.020 [0.047] N=2,042	0.027 [0.051] N=1,863	0.042 [0.054] N=1,639	0.007 [0.063] N=1,358	**0.153* [0.084] N=1,048**
Underweight: WAZ below -2	0.036 [0.024] N=2,287	0.030 [0.026] N=2,087	0.022 [0.028] N=1,838	0.004 [0.034] N=1,518	0.009 [0.048] N=1,172
Wasted: WLZ/WHZ below -2, 0-60 months	-0.018 [0.037] N=1,731	-0.012 [0.039] N=1,584	-0.015 [0.045] N=1,396	0.019 [0.056] N=1,150	-0.025 [0.068] N=884
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.044] N=1,731	0.012 [0.047] N=1,584	0.012 [0.053] N=1,396	-0.049 [0.065] N=1,150	-0.091 [0.087] N=884
Obese: WLZ/WHZ above +3, 0-60 months	-0.015 [0.031] N=1,731	-0.019 [0.035] N=1,584	0.008 [0.039] N=1,396	-0.034 [0.045] N=1,150	-0.066 [0.062] N=884
CIAF: any stunting, underweight, or wasting, 0-60 months	0.024 [0.050] N=1,672	0.046 [0.055] N=1,530	0.064 [0.057] N=1,347	0.017 [0.068] N=1,109	0.121 [0.084] N=852
BMI-for-age z score below -2	-0.028 [0.034] N=1,975	-0.048 [0.034] N=1,807	-0.052 [0.041] N=1,585	-0.012 [0.046] N=1,309	-0.033 [0.057] N=1,010
WHO 2007 BMI overweight above +1, 61+ months	-0.095 [0.170] N=240	-0.125 [0.188] N=218	-0.092 [0.202] N=190	-0.163 [0.216] N=164	0.035 [0.304] N=126
WHO 2007 BMI obesity above +2, 61+ months	-0.035 [0.130] N=240	-0.088 [0.136] N=218	-0.085 [0.153] N=190	-0.179 [0.173] N=164	-0.110 [0.227] N=126
WHO 2007 BMI thinness below -2, 61+ months	0.018 [0.096] N=240	-0.010 [0.105] N=218	-0.073 [0.102] N=190	-0.117 [0.121] N=164	-0.221 [0.159] N=126

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.092	**0.304* [0.180] N=2,274**	**0.318* [0.191] N=2,111**	0.308 [0.203] N=1,896	**0.395* [0.209] N=1,642**	0.200 [0.233] N=1,347
WHO weight-for-age z score	0.297	-0.032 [0.103] N=2,531	-0.004 [0.110] N=2,349	0.041 [0.113] N=2,111	0.016 [0.124] N=1,821	0.007 [0.131] N=1,496
WHO weight-for-length/height z score, 0-60 months	0.362	0.143 [0.158] N=1,919	0.139 [0.160] N=1,789	0.153 [0.169] N=1,610	-0.027 [0.188] N=1,390	0.078 [0.207] N=1,137
WHO BMI-for-age z score	0.370	0.080 [0.157] N=2,201	0.104 [0.158] N=2,049	0.155 [0.167] N=1,838	-0.078 [0.174] N=1,589	0.036 [0.192] N=1,305
Stunted: HAZ below -2	0.194 P=429	-0.041 [0.035] N=2,274	-0.042 [0.037] N=2,111	-0.037 [0.038] N=1,896	-0.053 [0.040] N=1,642	-0.013 [0.045] N=1,347
Underweight: WAZ below -2	0.053 P=138	0.010 [0.019] N=2,531	0.003 [0.020] N=2,349	-0.003 [0.021] N=2,111	-0.010 [0.024] N=1,821	-0.012 [0.029] N=1,496
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=214	-0.034 [0.027] N=1,919	-0.030 [0.027] N=1,789	-0.030 [0.029] N=1,610	-0.011 [0.031] N=1,390	-0.029 [0.032] N=1,137
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.023 [0.037] N=1,919	0.020 [0.039] N=1,789	0.027 [0.041] N=1,610	-0.000 [0.046] N=1,390	0.003 [0.055] N=1,137
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=184	-0.008 [0.029] N=1,919	-0.008 [0.032] N=1,789	0.011 [0.033] N=1,610	-0.002 [0.037] N=1,390	0.003 [0.043] N=1,137
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=479	-0.044 [0.040] N=1,855	-0.033 [0.041] N=1,729	-0.030 [0.042] N=1,554	-0.051 [0.045] N=1,342	-0.028 [0.050] N=1,097
BMI-for-age z score below -2	0.114 P=253	-0.015 [0.025] N=2,201	-0.020 [0.025] N=2,049	-0.021 [0.028] N=1,838	0.007 [0.029] N=1,589	-0.001 [0.031] N=1,305
WHO 2007 BMI overweight above +1, 61+ months	0.394 P=114	-0.069 [0.122] N=283	-0.070 [0.130] N=261	-0.053 [0.139] N=234	-0.109 [0.146] N=209	-0.063 [0.186] N=173
WHO 2007 BMI obesity above +2, 61+ months	0.218 P=63	-0.105 [0.085] N=283	-0.117 [0.085] N=261	-0.118 [0.094] N=234	**-0.180* [0.095] N=209**	-0.175 [0.124] N=173
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=29	0.026 [0.063] N=283	0.005 [0.065] N=261	-0.018 [0.061] N=234	-0.028 [0.070] N=209	-0.051 [0.092] N=173

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.236 [0.257] N=1,909	0.246 [0.287] N=1,746	0.210 [0.315] N=1,531	0.438 [0.348] N=1,277	-0.132 [0.482] N=982
WHO weight-for-age z score	-0.219 [0.144] N=2,132	-0.200 [0.160] N=1,950	-0.158 [0.173] N=1,712	-0.283 [0.214] N=1,422	-0.368 [0.263] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.049 [0.232] N=1,615	0.026 [0.240] N=1,485	0.010 [0.270] N=1,306	-0.355 [0.341] N=1,086	-0.222 [0.427] N=833
WHO BMI-for-age z score	0.032 [0.240] N=1,847	0.078 [0.245] N=1,695	0.164 [0.281] N=1,484	-0.285 [0.336] N=1,235	-0.051 [0.435] N=951
Stunted: HAZ below -2	-0.006 [0.050] N=1,909	-0.000 [0.054] N=1,746	0.019 [0.058] N=1,531	-0.008 [0.069] N=1,277	0.142 [0.096] N=982
Underweight: WAZ below -2	0.034 [0.025] N=2,132	0.027 [0.027] N=1,950	0.021 [0.030] N=1,712	0.016 [0.036] N=1,422	0.022 [0.048] N=1,097
Wasted: WLZ/WHZ below -2, 0-60 months	-0.035 [0.037] N=1,615	-0.028 [0.038] N=1,485	-0.028 [0.043] N=1,306	-0.004 [0.055] N=1,086	-0.063 [0.059] N=833
Overweight: WLZ/WHZ above +2, 0-60 months	0.009 [0.046] N=1,615	-0.000 [0.049] N=1,485	0.007 [0.054] N=1,306	-0.059 [0.068] N=1,086	-0.084 [0.094] N=833
Obese: WLZ/WHZ above +3, 0-60 months	-0.034 [0.033] N=1,615	-0.038 [0.037] N=1,485	-0.007 [0.040] N=1,306	-0.041 [0.048] N=1,086	-0.061 [0.068] N=833
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.002 [0.053] N=1,560	0.018 [0.057] N=1,434	0.041 [0.062] N=1,259	-0.008 [0.074] N=1,047	0.074 [0.096] N=802
BMI-for-age z score below -2	-0.023 [0.034] N=1,847	-0.041 [0.033] N=1,695	-0.048 [0.039] N=1,484	-0.014 [0.045] N=1,235	-0.051 [0.053] N=951
WHO 2007 BMI overweight above +1, 61+ months	-0.107 [0.186] N=229	-0.104 [0.197] N=207	-0.062 [0.222] N=180	-0.177 [0.247] N=155	-0.014 [0.332] N=119
WHO 2007 BMI obesity above +2, 61+ months	-0.072 [0.132] N=229	-0.086 [0.132] N=207	-0.081 [0.155] N=180	-0.210 [0.171] N=155	-0.232 [0.236] N=119
WHO 2007 BMI thinness below -2, 61+ months	0.049 [0.101] N=229	-0.006 [0.107] N=207	-0.056 [0.105] N=180	-0.096 [0.127] N=155	**-0.277* [0.162] N=119**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.092	**0.312* [0.179] N=2,274**	**0.323* [0.190] N=2,111**	0.309 [0.203] N=1,896	**0.397* [0.209] N=1,642**	0.209 [0.232] N=1,347
WHO weight-for-age z score	0.297	-0.018 [0.103] N=2,531	0.006 [0.111] N=2,349	0.052 [0.113] N=2,111	0.028 [0.123] N=1,821	0.015 [0.131] N=1,496
WHO weight-for-length/height z score, 0-60 months	0.362	0.138 [0.158] N=1,919	0.135 [0.159] N=1,789	0.142 [0.168] N=1,610	-0.037 [0.186] N=1,390	0.058 [0.206] N=1,137
WHO BMI-for-age z score	0.370	0.084 [0.157] N=2,201	0.108 [0.157] N=2,049	0.157 [0.167] N=1,838	-0.076 [0.173] N=1,589	0.029 [0.193] N=1,305
Stunted: HAZ below -2	0.194 P=429	-0.043 [0.035] N=2,274	-0.043 [0.036] N=2,111	-0.038 [0.038] N=1,896	-0.054 [0.040] N=1,642	-0.016 [0.045] N=1,347
Underweight: WAZ below -2	0.053 P=138	0.006 [0.019] N=2,531	0.001 [0.020] N=2,349	-0.006 [0.021] N=2,111	-0.013 [0.024] N=1,821	-0.015 [0.029] N=1,496
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=214	-0.034 [0.027] N=1,919	-0.029 [0.027] N=1,789	-0.029 [0.029] N=1,610	-0.011 [0.031] N=1,390	-0.031 [0.032] N=1,137
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.022 [0.037] N=1,919	0.019 [0.039] N=1,789	0.024 [0.041] N=1,610	-0.004 [0.046] N=1,390	-0.002 [0.055] N=1,137
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=184	-0.008 [0.029] N=1,919	-0.008 [0.031] N=1,789	0.010 [0.033] N=1,610	-0.003 [0.037] N=1,390	0.000 [0.043] N=1,137
CIAF: any stunting, underweight, or wasting, 0-60 months	0.262 P=479	-0.044 [0.040] N=1,855	-0.034 [0.041] N=1,729	-0.028 [0.042] N=1,554	-0.049 [0.045] N=1,342	-0.028 [0.049] N=1,097
BMI-for-age z score below -2	0.114 P=253	-0.015 [0.025] N=2,201	-0.021 [0.025] N=2,049	-0.021 [0.028] N=1,838	0.007 [0.029] N=1,589	-0.001 [0.031] N=1,305
WHO 2007 BMI overweight above +1, 61+ months	0.394 P=114	-0.065 [0.116] N=283	-0.065 [0.125] N=261	-0.059 [0.133] N=234	-0.105 [0.139] N=209	-0.064 [0.177] N=173
WHO 2007 BMI obesity above +2, 61+ months	0.218 P=63	-0.100 [0.082] N=283	-0.116 [0.084] N=261	-0.109 [0.092] N=234	**-0.170* [0.092] N=209**	-0.161 [0.120] N=173
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=29	0.011 [0.063] N=283	-0.006 [0.063] N=261	-0.024 [0.060] N=234	-0.037 [0.067] N=209	-0.072 [0.085] N=173

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.243 [0.256] N=1,909	0.244 [0.287] N=1,746	0.211 [0.314] N=1,531	0.433 [0.349] N=1,277	-0.109 [0.479] N=982
WHO weight-for-age z score	-0.198 [0.143] N=2,132	-0.186 [0.161] N=1,950	-0.137 [0.170] N=1,712	-0.254 [0.209] N=1,422	-0.346 [0.259] N=1,097
WHO weight-for-length/height z score, 0-60 months	0.072 [0.233] N=1,615	0.058 [0.241] N=1,485	0.020 [0.270] N=1,306	-0.348 [0.343] N=1,086	-0.231 [0.435] N=833
WHO BMI-for-age z score	0.053 [0.239] N=1,847	0.105 [0.244] N=1,695	0.177 [0.280] N=1,484	-0.272 [0.334] N=1,235	-0.054 [0.434] N=951
Stunted: HAZ below -2	-0.009 [0.050] N=1,909	-0.001 [0.054] N=1,746	0.015 [0.057] N=1,531	-0.014 [0.068] N=1,277	0.132 [0.094] N=982
Underweight: WAZ below -2	0.030 [0.025] N=2,132	0.025 [0.027] N=1,950	0.017 [0.029] N=1,712	0.010 [0.036] N=1,422	0.018 [0.048] N=1,097
Wasted: WLZ/WHZ below -2, 0-60 months	-0.035 [0.037] N=1,615	-0.030 [0.039] N=1,485	-0.029 [0.044] N=1,306	-0.006 [0.055] N=1,086	-0.068 [0.061] N=833
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.046] N=1,615	0.008 [0.049] N=1,485	0.010 [0.054] N=1,306	-0.055 [0.069] N=1,086	-0.084 [0.096] N=833
Obese: WLZ/WHZ above +3, 0-60 months	-0.032 [0.033] N=1,615	-0.036 [0.037] N=1,485	-0.005 [0.040] N=1,306	-0.042 [0.048] N=1,086	-0.063 [0.068] N=833
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.003 [0.053] N=1,560	0.017 [0.058] N=1,434	0.039 [0.061] N=1,259	-0.016 [0.073] N=1,047	0.063 [0.091] N=802
BMI-for-age z score below -2	-0.026 [0.034] N=1,847	-0.045 [0.033] N=1,695	-0.049 [0.039] N=1,484	-0.016 [0.045] N=1,235	-0.051 [0.053] N=951
WHO 2007 BMI overweight above +1, 61+ months	-0.096 [0.178] N=229	-0.091 [0.191] N=207	-0.060 [0.210] N=180	-0.163 [0.231] N=155	-0.028 [0.332] N=119
WHO 2007 BMI obesity above +2, 61+ months	-0.080 [0.127] N=229	-0.103 [0.128] N=207	-0.085 [0.151] N=180	-0.226 [0.164] N=155	-0.240 [0.236] N=119
WHO 2007 BMI thinness below -2, 61+ months	0.023 [0.100] N=229	-0.025 [0.105] N=207	-0.076 [0.104] N=180	-0.118 [0.124] N=155	**-0.278* [0.163] N=119**

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033	-0.247 [0.205] N=2,349	-0.290 [0.216] N=2,123	-0.347 [0.229] N=1,863	**-0.451* [0.249] N=1,585**	**-0.517* [0.270] N=1,288**
WHO weight-for-age z score	0.329	-0.057 [0.105] N=2,602	-0.050 [0.111] N=2,364	-0.168 [0.117] N=2,068	-0.158 [0.128] N=1,752	-0.145 [0.142] N=1,419
WHO weight-for-length/height z score, 0-60 months	0.389	0.040 [0.182] N=1,912	0.072 [0.187] N=1,733	0.059 [0.207] N=1,512	0.132 [0.223] N=1,287	0.057 [0.266] N=1,037
WHO BMI-for-age z score	0.381	0.074 [0.201] N=2,270	0.076 [0.209] N=2,054	0.066 [0.232] N=1,803	0.173 [0.248] N=1,532	0.137 [0.281] N=1,247
Stunted: HAZ below -2	0.188 P=446	0.013 [0.036] N=2,349	0.021 [0.038] N=2,123	0.008 [0.041] N=1,863	0.025 [0.045] N=1,585	0.021 [0.051] N=1,288
Underweight: WAZ below -2	0.049 P=132	0.011 [0.015] N=2,602	0.006 [0.016] N=2,364	0.016 [0.017] N=2,068	0.015 [0.017] N=1,752	0.007 [0.017] N=1,419
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=208	0.025 [0.028] N=1,912	0.019 [0.029] N=1,733	0.031 [0.032] N=1,512	-0.003 [0.032] N=1,287	0.006 [0.037] N=1,037
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=387	0.035 [0.035] N=1,912	0.039 [0.036] N=1,733	0.034 [0.038] N=1,512	0.045 [0.043] N=1,287	0.036 [0.049] N=1,037
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=192	0.007 [0.028] N=1,912	0.005 [0.029] N=1,733	-0.000 [0.031] N=1,512	0.007 [0.035] N=1,287	-0.011 [0.038] N=1,037
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=487	0.049 [0.038] N=1,850	0.045 [0.040] N=1,674	0.042 [0.042] N=1,460	0.014 [0.044] N=1,243	0.026 [0.047] N=1,000
BMI-for-age z score below -2	0.118 P=267	0.029 [0.031] N=2,270	0.025 [0.033] N=2,054	0.023 [0.036] N=1,803	0.009 [0.037] N=1,532	0.024 [0.043] N=1,247
WHO 2007 BMI overweight above +1, 61+ months	0.405 P=145	-0.010 [0.102] N=362	-0.087 [0.103] N=322	-0.118 [0.109] N=292	-0.045 [0.120] N=247	-0.065 [0.137] N=207
WHO 2007 BMI obesity above +2, 61+ months	0.236 P=82	0.038 [0.097] N=362	-0.044 [0.099] N=322	-0.070 [0.108] N=292	-0.002 [0.118] N=247	-0.046 [0.138] N=207
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=40	0.026 [0.067] N=362	0.037 [0.072] N=322	0.020 [0.074] N=292	0.016 [0.078] N=247	0.011 [0.089] N=207

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.010 [0.264] N=2,015	-0.041 [0.294] N=1,789	-0.133 [0.332] N=1,529	-0.352 [0.395] N=1,251	-0.359 [0.473] N=954
WHO weight-for-age z score	-0.055 [0.147] N=2,239	-0.028 [0.159] N=2,001	**-0.287* [0.169] N=1,705**	-0.302 [0.199] N=1,389	-0.296 [0.243] N=1,056
WHO weight-for-length/height z score, 0-60 months	-0.186 [0.260] N=1,654	-0.172 [0.273] N=1,475	-0.269 [0.321] N=1,254	-0.176 [0.358] N=1,029	-0.675 [0.429] N=779
WHO BMI-for-age z score	-0.334 [0.231] N=1,952	-0.399 [0.244] N=1,736	**-0.515* [0.290] N=1,485**	-0.421 [0.322] N=1,214	**-0.880** [0.378] N=929**
Stunted: HAZ below -2	0.019 [0.046] N=2,015	0.030 [0.050] N=1,789	0.010 [0.055] N=1,529	0.050 [0.065] N=1,251	0.018 [0.085] N=954
Underweight: WAZ below -2	0.002 [0.025] N=2,239	-0.008 [0.027] N=2,001	0.010 [0.031] N=1,705	-0.001 [0.034] N=1,389	-0.034 [0.039] N=1,056
Wasted: WLZ/WHZ below -2, 0-60 months	**0.077** [0.038] N=1,654**	**0.073* [0.041] N=1,475**	**0.109** [0.046] N=1,254**	0.056 [0.050] N=1,029	**0.123** [0.054] N=779**
Overweight: WLZ/WHZ above +2, 0-60 months	0.024 [0.054] N=1,654	0.020 [0.057] N=1,475	0.010 [0.064] N=1,254	0.031 [0.075] N=1,029	-0.036 [0.094] N=779
Obese: WLZ/WHZ above +3, 0-60 months	0.010 [0.041] N=1,654	0.001 [0.043] N=1,475	-0.010 [0.048] N=1,254	0.003 [0.062] N=1,029	-0.084 [0.080] N=779
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.101* [0.060] N=1,601**	0.096 [0.065] N=1,425	0.108 [0.074] N=1,211	0.068 [0.086] N=994	0.091 [0.104] N=751
BMI-for-age z score below -2	**0.078*** [0.030] N=1,952**	**0.080** [0.034] N=1,736**	**0.089** [0.040] N=1,485**	**0.076** [0.038] N=1,214**	**0.159*** [0.043] N=929**
WHO 2007 BMI overweight above +1, 61+ months	-0.109 [0.151] N=301	-0.262 [0.159] N=261	**-0.370** [0.180] N=231**	-0.310 [0.233] N=186	**-0.591* [0.322] N=146**
WHO 2007 BMI obesity above +2, 61+ months	0.013 [0.137] N=301	-0.138 [0.141] N=261	-0.217 [0.167] N=231	-0.120 [0.187] N=186	-0.273 [0.286] N=146
WHO 2007 BMI thinness below -2, 61+ months	0.080 [0.076] N=301	0.130 [0.086] N=261	0.106 [0.088] N=231	0.134 [0.088] N=186	**0.225** [0.108] N=146**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033	-0.249 [0.204] N=2,349	-0.294 [0.216] N=2,123	-0.350 [0.229] N=1,863	**-0.457* [0.248] N=1,585**	**-0.514* [0.268] N=1,288**
WHO weight-for-age z score	0.329	-0.057 [0.105] N=2,602	-0.050 [0.111] N=2,364	-0.168 [0.117] N=2,068	-0.158 [0.128] N=1,752	-0.145 [0.142] N=1,419
WHO weight-for-length/height z score, 0-60 months	0.389	0.045 [0.181] N=1,912	0.075 [0.187] N=1,733	0.061 [0.206] N=1,512	0.133 [0.222] N=1,287	0.053 [0.263] N=1,037
WHO BMI-for-age z score	0.381	0.074 [0.201] N=2,270	0.076 [0.210] N=2,054	0.066 [0.232] N=1,803	0.173 [0.248] N=1,532	0.136 [0.281] N=1,247
Stunted: HAZ below -2	0.188 P=446	0.013 [0.036] N=2,349	0.021 [0.038] N=2,123	0.008 [0.041] N=1,863	0.025 [0.045] N=1,585	0.021 [0.051] N=1,288
Underweight: WAZ below -2	0.049 P=132	0.011 [0.015] N=2,602	0.006 [0.016] N=2,364	0.016 [0.017] N=2,068	0.015 [0.017] N=1,752	0.007 [0.017] N=1,419
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=208	0.023 [0.028] N=1,912	0.017 [0.029] N=1,733	0.029 [0.031] N=1,512	-0.003 [0.032] N=1,287	0.007 [0.036] N=1,037
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=387	0.034 [0.035] N=1,912	0.037 [0.036] N=1,733	0.033 [0.039] N=1,512	0.044 [0.044] N=1,287	0.037 [0.050] N=1,037
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=192	0.006 [0.029] N=1,912	0.004 [0.029] N=1,733	-0.002 [0.032] N=1,512	0.006 [0.036] N=1,287	-0.009 [0.039] N=1,037
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=487	0.047 [0.039] N=1,850	0.043 [0.040] N=1,674	0.040 [0.043] N=1,460	0.013 [0.044] N=1,243	0.027 [0.047] N=1,000
BMI-for-age z score below -2	0.118 P=267	0.029 [0.031] N=2,270	0.024 [0.033] N=2,054	0.023 [0.036] N=1,803	0.009 [0.037] N=1,532	0.025 [0.043] N=1,247
WHO 2007 BMI overweight above +1, 61+ months	0.405 P=145	0.003 [0.103] N=362	-0.077 [0.104] N=322	-0.111 [0.110] N=292	-0.046 [0.121] N=247	-0.057 [0.139] N=207
WHO 2007 BMI obesity above +2, 61+ months	0.236 P=82	0.052 [0.100] N=362	-0.033 [0.103] N=322	-0.062 [0.111] N=292	-0.003 [0.120] N=247	-0.041 [0.141] N=207
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=40	0.020 [0.066] N=362	0.032 [0.072] N=322	0.014 [0.074] N=292	0.018 [0.079] N=247	0.002 [0.090] N=207

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.014 [0.264] N=2,015	-0.049 [0.294] N=1,789	-0.144 [0.330] N=1,529	-0.380 [0.393] N=1,251	-0.384 [0.470] N=954
WHO weight-for-age z score	-0.055 [0.147] N=2,239	-0.028 [0.159] N=2,001	**-0.290* [0.169] N=1,705**	-0.304 [0.199] N=1,389	-0.297 [0.243] N=1,056
WHO weight-for-length/height z score, 0-60 months	-0.172 [0.262] N=1,654	-0.164 [0.276] N=1,475	-0.266 [0.323] N=1,254	-0.172 [0.361] N=1,029	-0.658 [0.434] N=779
WHO BMI-for-age z score	-0.332 [0.233] N=1,952	-0.399 [0.246] N=1,736	**-0.519* [0.291] N=1,485**	-0.424 [0.324] N=1,214	**-0.876** [0.382] N=929**
Stunted: HAZ below -2	0.018 [0.046] N=2,015	0.029 [0.050] N=1,789	0.009 [0.055] N=1,529	0.049 [0.065] N=1,251	0.019 [0.085] N=954
Underweight: WAZ below -2	0.002 [0.025] N=2,239	-0.009 [0.027] N=2,001	0.011 [0.031] N=1,705	-0.001 [0.034] N=1,389	-0.034 [0.039] N=1,056
Wasted: WLZ/WHZ below -2, 0-60 months	**0.071* [0.038] N=1,654**	0.065 [0.041] N=1,475	**0.102** [0.046] N=1,254**	0.048 [0.050] N=1,029	**0.116** [0.054] N=779**
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.054] N=1,654	0.016 [0.057] N=1,475	0.004 [0.064] N=1,254	0.022 [0.076] N=1,029	-0.040 [0.094] N=779
Obese: WLZ/WHZ above +3, 0-60 months	0.006 [0.041] N=1,654	-0.004 [0.043] N=1,475	-0.017 [0.049] N=1,254	-0.005 [0.063] N=1,029	-0.089 [0.081] N=779
CIAF: any stunting, underweight, or wasting, 0-60 months	0.093 [0.060] N=1,601	0.087 [0.065] N=1,425	0.097 [0.075] N=1,211	0.056 [0.087] N=994	0.084 [0.104] N=751
BMI-for-age z score below -2	**0.076** [0.030] N=1,952**	**0.076** [0.034] N=1,736**	**0.086** [0.040] N=1,485**	**0.072* [0.038] N=1,214**	**0.157*** [0.044] N=929**
WHO 2007 BMI overweight above +1, 61+ months	-0.091 [0.151] N=301	-0.251 [0.158] N=261	**-0.361** [0.180] N=231**	-0.316 [0.234] N=186	**-0.568* [0.326] N=146**
WHO 2007 BMI obesity above +2, 61+ months	0.035 [0.145] N=301	-0.121 [0.150] N=261	-0.204 [0.176] N=231	-0.127 [0.195] N=186	-0.250 [0.299] N=146
WHO 2007 BMI thinness below -2, 61+ months	0.075 [0.078] N=301	0.127 [0.087] N=261	0.100 [0.086] N=231	0.137 [0.085] N=186	**0.205* [0.110] N=146**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033	-0.250 [0.201] N=2,349	-0.286 [0.213] N=2,123	-0.333 [0.228] N=1,863	**-0.445*** **[0.243]** **N=1,585**	**-0.502*** **[0.262]** **N=1,288**
WHO weight-for-age z score	0.329	-0.052 [0.105] N=2,602	-0.043 [0.110] N=2,364	-0.152 [0.116] N=2,068	-0.153 [0.129] N=1,752	-0.142 [0.144] N=1,419
WHO weight-for-length/height z score, 0-60 months	0.389	0.054 [0.183] N=1,912	0.081 [0.189] N=1,733	0.073 [0.210] N=1,512	0.136 [0.223] N=1,287	0.055 [0.262] N=1,037
WHO BMI-for-age z score	0.381	0.077 [0.201] N=2,270	0.079 [0.209] N=2,054	0.070 [0.231] N=1,803	0.171 [0.246] N=1,532	0.127 [0.280] N=1,247
Stunted: HAZ below -2	0.188 P=446	0.014 [0.035] N=2,349	0.019 [0.037] N=2,123	0.003 [0.040] N=1,863	0.021 [0.044] N=1,585	0.016 [0.049] N=1,288
Underweight: WAZ below -2	0.049 P=132	0.011 [0.015] N=2,602	0.006 [0.016] N=2,364	0.013 [0.017] N=2,068	0.013 [0.017] N=1,752	0.005 [0.017] N=1,419
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=208	0.023 [0.028] N=1,912	0.016 [0.029] N=1,733	0.027 [0.032] N=1,512	-0.004 [0.032] N=1,287	0.006 [0.036] N=1,037
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=387	0.037 [0.036] N=1,912	0.039 [0.037] N=1,733	0.034 [0.039] N=1,512	0.044 [0.044] N=1,287	0.038 [0.050] N=1,037
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=192	0.008 [0.029] N=1,912	0.005 [0.029] N=1,733	-0.001 [0.032] N=1,512	0.006 [0.036] N=1,287	-0.009 [0.040] N=1,037
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=487	0.051 [0.038] N=1,850	0.044 [0.040] N=1,674	0.036 [0.043] N=1,460	0.010 [0.044] N=1,243	0.023 [0.047] N=1,000
BMI-for-age z score below -2	0.118 P=267	0.028 [0.030] N=2,270	0.023 [0.033] N=2,054	0.021 [0.036] N=1,803	0.007 [0.038] N=1,532	0.023 [0.044] N=1,247
WHO 2007 BMI overweight above +1, 61+ months	0.405 P=145	-0.010 [0.105] N=362	-0.090 [0.106] N=322	-0.127 [0.115] N=292	-0.055 [0.125] N=247	-0.071 [0.148] N=207
WHO 2007 BMI obesity above +2, 61+ months	0.236 P=82	0.031 [0.099] N=362	-0.050 [0.103] N=322	-0.087 [0.113] N=292	-0.030 [0.121] N=247	-0.045 [0.147] N=207
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=40	0.013 [0.065] N=362	0.019 [0.068] N=322	0.007 [0.069] N=292	0.007 [0.075] N=247	-0.014 [0.085] N=207

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.008 [0.262] N=2,015	-0.016 [0.291] N=1,789	-0.082 [0.330] N=1,529	-0.318 [0.386] N=1,251	-0.315 [0.456] N=954
WHO weight-for-age z score	-0.047 [0.146] N=2,239	-0.018 [0.159] N=2,001	-0.258 [0.170] N=1,705	-0.288 [0.199] N=1,389	-0.288 [0.243] N=1,056
WHO weight-for-length/height z score, 0-60 months	-0.173 [0.265] N=1,654	-0.164 [0.280] N=1,475	-0.266 [0.326] N=1,254	-0.188 [0.359] N=1,029	-0.668 [0.430] N=779
WHO BMI-for-age z score	-0.339 [0.232] N=1,952	-0.402 [0.246] N=1,736	**-0.530* [0.289] N=1,485**	-0.447 [0.321] N=1,214	**-0.911** [0.385] N=929**
Stunted: HAZ below -2	0.015 [0.046] N=2,015	0.024 [0.050] N=1,789	-0.003 [0.055] N=1,529	0.037 [0.064] N=1,251	0.006 [0.083] N=954
Underweight: WAZ below -2	-0.000 [0.024] N=2,239	-0.011 [0.027] N=2,001	0.003 [0.031] N=1,705	-0.007 [0.035] N=1,389	-0.040 [0.040] N=1,056
Wasted: WLZ/WHZ below -2, 0-60 months	**0.072* [0.039] N=1,654**	0.064 [0.042] N=1,475	**0.098** [0.048] N=1,254**	0.045 [0.051] N=1,029	**0.114** [0.056] N=779**
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.055] N=1,654	0.018 [0.058] N=1,475	0.000 [0.065] N=1,254	0.016 [0.077] N=1,029	-0.041 [0.095] N=779
Obese: WLZ/WHZ above +3, 0-60 months	0.007 [0.041] N=1,654	-0.004 [0.043] N=1,475	-0.019 [0.049] N=1,254	-0.008 [0.063] N=1,029	-0.090 [0.081] N=779
CIAF: any stunting, underweight, or wasting, 0-60 months	0.096 [0.060] N=1,601	0.085 [0.066] N=1,425	0.087 [0.076] N=1,211	0.045 [0.088] N=994	0.079 [0.104] N=751
BMI-for-age z score below -2	**0.075** [0.030] N=1,952**	**0.073** [0.034] N=1,736**	**0.082** [0.040] N=1,485**	**0.068* [0.039] N=1,214**	**0.152*** [0.045] N=929**
WHO 2007 BMI overweight above +1, 61+ months	-0.127 [0.147] N=301	**-0.290* [0.153] N=261**	**-0.411** [0.174] N=231**	-0.348 [0.225] N=186	**-0.725** [0.309] N=146**
WHO 2007 BMI obesity above +2, 61+ months	0.001 [0.143] N=301	-0.154 [0.146] N=261	-0.242 [0.174] N=231	-0.158 [0.191] N=186	-0.314 [0.303] N=146
WHO 2007 BMI thinness below -2, 61+ months	0.069 [0.076] N=301	0.112 [0.085] N=261	0.091 [0.083] N=231	0.114 [0.083] N=186	0.163 [0.131] N=146

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033	-0.243 [0.202] N=2,349	-0.274 [0.214] N=2,123	-0.320 [0.231] N=1,863	**-0.425* [0.245] N=1,585**	**-0.489* [0.261] N=1,288**
WHO weight-for-age z score	0.329	-0.058 [0.104] N=2,602	-0.044 [0.109] N=2,364	-0.155 [0.115] N=2,068	-0.145 [0.126] N=1,752	-0.143 [0.139] N=1,419
WHO weight-for-length/height z score, 0-60 months	0.389	0.052 [0.182] N=1,912	0.081 [0.188] N=1,733	0.068 [0.208] N=1,512	0.144 [0.223] N=1,287	0.067 [0.265] N=1,037
WHO BMI-for-age z score	0.381	0.070 [0.201] N=2,270	0.073 [0.209] N=2,054	0.065 [0.232] N=1,803	0.171 [0.245] N=1,532	0.127 [0.278] N=1,247
Stunted: HAZ below -2	0.188 P=446	0.013 [0.035] N=2,349	0.018 [0.037] N=2,123	0.003 [0.040] N=1,863	0.020 [0.044] N=1,585	0.017 [0.049] N=1,288
Underweight: WAZ below -2	0.049 P=132	0.011 [0.015] N=2,602	0.005 [0.016] N=2,364	0.014 [0.017] N=2,068	0.013 [0.017] N=1,752	0.006 [0.018] N=1,419
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=208	0.023 [0.028] N=1,912	0.016 [0.029] N=1,733	0.029 [0.032] N=1,512	-0.004 [0.032] N=1,287	0.006 [0.036] N=1,037
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=387	0.035 [0.035] N=1,912	0.038 [0.037] N=1,733	0.034 [0.039] N=1,512	0.045 [0.044] N=1,287	0.039 [0.050] N=1,037
Obese: WLZ/WHZ above +3, 0-60 months	0.097 P=192	0.009 [0.029] N=1,912	0.006 [0.029] N=1,733	0.001 [0.032] N=1,512	0.009 [0.036] N=1,287	-0.006 [0.040] N=1,037
CIAF: any stunting, underweight, or wasting, 0-60 months	0.264 P=487	0.050 [0.038] N=1,850	0.043 [0.039] N=1,674	0.036 [0.042] N=1,460	0.010 [0.043] N=1,243	0.027 [0.046] N=1,000
BMI-for-age z score below -2	0.118 P=267	0.027 [0.031] N=2,270	0.022 [0.033] N=2,054	0.021 [0.036] N=1,803	0.006 [0.037] N=1,532	0.021 [0.043] N=1,247
WHO 2007 BMI overweight above +1, 61+ months	0.405 P=145	-0.005 [0.106] N=362	-0.084 [0.109] N=322	-0.117 [0.115] N=292	-0.038 [0.126] N=247	-0.052 [0.147] N=207
WHO 2007 BMI obesity above +2, 61+ months	0.236 P=82	0.025 [0.104] N=362	-0.061 [0.107] N=322	-0.094 [0.116] N=292	-0.035 [0.126] N=247	-0.079 [0.146] N=207
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=40	0.006 [0.067] N=362	0.015 [0.071] N=322	-0.014 [0.074] N=292	-0.008 [0.078] N=247	-0.049 [0.085] N=207

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.011 [0.261] N=2,015	-0.005 [0.294] N=1,789	-0.055 [0.337] N=1,529	-0.275 [0.392] N=1,251	-0.324 [0.466] N=954
WHO weight-for-age z score	-0.036 [0.143] N=2,239	0.001 [0.155] N=2,001	-0.236 [0.167] N=1,705	-0.239 [0.193] N=1,389	-0.248 [0.232] N=1,056
WHO weight-for-length/height z score, 0-60 months	-0.171 [0.263] N=1,654	-0.167 [0.275] N=1,475	-0.283 [0.318] N=1,254	-0.191 [0.354] N=1,029	-0.658 [0.429] N=779
WHO BMI-for-age z score	-0.344 [0.232] N=1,952	**-0.409* [0.245] N=1,736**	**-0.538* [0.289] N=1,485**	-0.443 [0.316] N=1,214	**-0.884** [0.381] N=929**
Stunted: HAZ below -2	0.015 [0.045] N=2,015	0.023 [0.049] N=1,789	-0.006 [0.054] N=1,529	0.030 [0.064] N=1,251	0.009 [0.083] N=954
Underweight: WAZ below -2	-0.001 [0.024] N=2,239	-0.014 [0.027] N=2,001	0.002 [0.031] N=1,705	-0.009 [0.035] N=1,389	-0.041 [0.040] N=1,056
Wasted: WLZ/WHZ below -2, 0-60 months	**0.072* [0.038] N=1,654**	0.065 [0.041] N=1,475	**0.105** [0.046] N=1,254**	0.050 [0.050] N=1,029	**0.119** [0.054] N=779**
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.054] N=1,654	0.014 [0.058] N=1,475	-0.003 [0.064] N=1,254	0.013 [0.076] N=1,029	-0.042 [0.095] N=779
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.041] N=1,654	-0.003 [0.043] N=1,475	-0.018 [0.049] N=1,254	-0.006 [0.063] N=1,029	-0.089 [0.082] N=779
CIAF: any stunting, underweight, or wasting, 0-60 months	0.095 [0.059] N=1,601	0.082 [0.065] N=1,425	0.082 [0.075] N=1,211	0.037 [0.086] N=994	0.079 [0.101] N=751
BMI-for-age z score below -2	**0.077** [0.030] N=1,952**	**0.075** [0.035] N=1,736**	**0.084** [0.041] N=1,485**	**0.071* [0.039] N=1,214**	**0.155*** [0.044] N=929**
WHO 2007 BMI overweight above +1, 61+ months	-0.118 [0.143] N=301	**-0.275* [0.149] N=261**	**-0.397** [0.164] N=231**	-0.350 [0.210] N=186	**-0.660** [0.273] N=146**
WHO 2007 BMI obesity above +2, 61+ months	0.008 [0.143] N=301	-0.153 [0.147] N=261	-0.247 [0.172] N=231	-0.168 [0.191] N=186	-0.313 [0.295] N=146
WHO 2007 BMI thinness below -2, 61+ months	0.067 [0.081] N=301	0.111 [0.090] N=261	0.065 [0.089] N=231	0.103 [0.089] N=186	0.107 [0.110] N=146

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.045	-0.303 [0.202] N=2,202	-0.336 [0.216] N=1,990	**-0.393*** **[0.228]** **N=1,749**	**-0.511**** **[0.244]** **N=1,483**	**-0.636**** **[0.262]** **N=1,207**
WHO weight-for-age z score	0.330	-0.045 [0.105] N=2,434	-0.022 [0.110] N=2,210	-0.145 [0.114] N=1,937	-0.148 [0.126] N=1,639	-0.173 [0.136] N=1,332
WHO weight-for-length/height z score, 0-60 months	0.394	0.104 [0.193] N=1,788	0.151 [0.201] N=1,621	0.107 [0.221] N=1,415	0.173 [0.236] N=1,203	0.076 [0.277] N=973
WHO BMI-for-age z score	0.396	0.061 [0.211] N=2,129	0.070 [0.221] N=1,927	0.012 [0.245] N=1,694	0.098 [0.260] N=1,437	0.034 [0.296] N=1,173
Stunted: HAZ below -2	0.188 P=419	0.019 [0.034] N=2,202	0.029 [0.036] N=1,990	0.016 [0.038] N=1,749	0.035 [0.041] N=1,483	0.042 [0.045] N=1,207
Underweight: WAZ below -2	0.048 P=123	0.009 [0.016] N=2,434	0.003 [0.017] N=2,210	0.009 [0.017] N=1,937	0.013 [0.018] N=1,639	0.010 [0.019] N=1,332
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=195	0.025 [0.029] N=1,788	0.016 [0.030] N=1,621	0.030 [0.032] N=1,415	0.001 [0.033] N=1,203	0.013 [0.036] N=973
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=364	0.040 [0.036] N=1,788	0.043 [0.038] N=1,621	0.034 [0.039] N=1,415	0.044 [0.043] N=1,203	0.034 [0.049] N=973
Obese: WLZ/WHZ above +3, 0-60 months	0.099 P=184	0.008 [0.029] N=1,788	0.006 [0.029] N=1,621	0.001 [0.031] N=1,415	0.013 [0.035] N=1,203	0.003 [0.038] N=973
CIAF: any stunting, underweight, or wasting, 0-60 months	0.265 P=459	0.057 [0.040] N=1,730	0.052 [0.042] N=1,566	0.048 [0.043] N=1,367	0.020 [0.045] N=1,161	0.041 [0.047] N=937
BMI-for-age z score below -2	0.115 P=243	0.027 [0.030] N=2,129	0.021 [0.032] N=1,927	0.023 [0.035] N=1,694	0.014 [0.035] N=1,437	0.032 [0.040] N=1,173
WHO 2007 BMI overweight above +1, 61+ months	0.414 P=142	-0.046 [0.107] N=347	-0.140 [0.106] N=309	-0.179 [0.114] N=281	-0.134 [0.127] N=237	-0.164 [0.146] N=198
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=80	0.008 [0.108] N=347	-0.089 [0.108] N=309	-0.132 [0.117] N=281	-0.082 [0.126] N=237	-0.119 [0.151] N=198
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=35	0.035 [0.069] N=347	0.047 [0.075] N=309	0.030 [0.077] N=281	0.038 [0.081] N=237	0.033 [0.090] N=198

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.033 [0.259] N=1,890	0.032 [0.291] N=1,678	-0.050 [0.331] N=1,437	-0.237 [0.386] N=1,171	-0.304 [0.441] N=895
WHO weight-for-age z score	-0.043 [0.148] N=2,095	0.006 [0.159] N=1,871	-0.266 [0.167] N=1,598	-0.278 [0.200] N=1,300	-0.296 [0.241] N=993
WHO weight-for-length/height z score, 0-60 months	-0.134 [0.277] N=1,545	-0.115 [0.293] N=1,378	-0.277 [0.343] N=1,172	-0.205 [0.374] N=960	-0.703 [0.455] N=730
WHO BMI-for-age z score	-0.319 [0.242] N=1,829	-0.367 [0.258] N=1,627	**-0.551* [0.307] N=1,394**	-0.469 [0.329] N=1,137	**-0.924** [0.389] N=873**
Stunted: HAZ below -2	0.005 [0.047] N=1,890	0.017 [0.052] N=1,678	-0.004 [0.057] N=1,437	0.026 [0.065] N=1,171	0.006 [0.085] N=895
Underweight: WAZ below -2	-0.002 [0.026] N=2,095	-0.014 [0.029] N=1,871	-0.001 [0.032] N=1,598	-0.007 [0.038] N=1,300	-0.041 [0.043] N=993
Wasted: WLZ/WHZ below -2, 0-60 months	**0.074* [0.040] N=1,545**	0.065 [0.043] N=1,378	**0.102** [0.048] N=1,172**	0.050 [0.054] N=960	**0.109* [0.058] N=730**
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.056] N=1,545	0.014 [0.060] N=1,378	-0.007 [0.066] N=1,172	0.001 [0.077] N=960	-0.058 [0.097] N=730
Obese: WLZ/WHZ above +3, 0-60 months	-0.006 [0.043] N=1,545	-0.018 [0.044] N=1,378	-0.033 [0.051] N=1,172	-0.019 [0.065] N=960	-0.092 [0.085] N=730
CIAF: any stunting, underweight, or wasting, 0-60 months	0.093 [0.065] N=1,496	0.084 [0.071] N=1,332	0.091 [0.079] N=1,133	0.026 [0.096] N=927	0.047 [0.118] N=703
BMI-for-age z score below -2	**0.074** [0.032] N=1,829**	**0.070* [0.036] N=1,627**	**0.083* [0.042] N=1,394**	**0.074* [0.041] N=1,137**	**0.155*** [0.045] N=873**
WHO 2007 BMI overweight above +1, 61+ months	-0.133 [0.151] N=289	**-0.308* [0.160] N=251**	**-0.418** [0.179] N=223**	**-0.388* [0.230] N=179**	**-0.658** [0.314] N=140**
WHO 2007 BMI obesity above +2, 61+ months	-0.001 [0.149] N=289	-0.159 [0.156] N=251	-0.256 [0.179] N=223	-0.157 [0.187] N=179	-0.270 [0.285] N=140
WHO 2007 BMI thinness below -2, 61+ months	0.069 [0.078] N=289	0.120 [0.089] N=251	0.090 [0.087] N=223	0.130 [0.084] N=179	**0.180* [0.105] N=140**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.045	-0.308 [0.202] N=2,202	-0.340 [0.216] N=1,990	**-0.396*** **[0.229]** **N=1,749**	**-0.508**** **[0.247]** **N=1,483**	**-0.627**** **[0.261]** **N=1,207**
WHO weight-for-age z score	0.330	-0.049 [0.105] N=2,434	-0.028 [0.111] N=2,210	-0.157 [0.115] N=1,937	-0.153 [0.128] N=1,639	-0.172 [0.141] N=1,332
WHO weight-for-length/height z score, 0-60 months	0.394	0.106 [0.193] N=1,788	0.149 [0.201] N=1,621	0.108 [0.221] N=1,415	0.179 [0.235] N=1,203	0.080 [0.280] N=973
WHO BMI-for-age z score	0.396	0.063 [0.211] N=2,129	0.071 [0.221] N=1,927	0.012 [0.245] N=1,694	0.097 [0.261] N=1,437	0.030 [0.298] N=1,173
Stunted: HAZ below -2	0.188 P=419	0.019 [0.034] N=2,202	0.028 [0.036] N=1,990	0.016 [0.038] N=1,749	0.034 [0.041] N=1,483	0.039 [0.045] N=1,207
Underweight: WAZ below -2	0.048 P=123	0.009 [0.016] N=2,434	0.004 [0.017] N=2,210	0.012 [0.017] N=1,937	0.014 [0.018] N=1,639	0.009 [0.019] N=1,332
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=195	0.023 [0.029] N=1,788	0.015 [0.030] N=1,621	0.028 [0.032] N=1,415	-0.004 [0.032] N=1,203	0.008 [0.036] N=973
Overweight: WLZ/WHZ above +2, 0-60 months	0.203 P=364	0.038 [0.036] N=1,788	0.041 [0.037] N=1,621	0.031 [0.039] N=1,415	0.041 [0.043] N=1,203	0.031 [0.049] N=973
Obese: WLZ/WHZ above +3, 0-60 months	0.099 P=184	0.007 [0.028] N=1,788	0.005 [0.028] N=1,621	-0.001 [0.031] N=1,415	0.010 [0.034] N=1,203	-0.000 [0.037] N=973
CIAF: any stunting, underweight, or wasting, 0-60 months	0.265 P=459	0.053 [0.040] N=1,730	0.047 [0.041] N=1,566	0.043 [0.043] N=1,367	0.012 [0.044] N=1,161	0.035 [0.046] N=937
BMI-for-age z score below -2	0.115 P=243	0.026 [0.030] N=2,129	0.020 [0.032] N=1,927	0.022 [0.035] N=1,694	0.013 [0.035] N=1,437	0.031 [0.040] N=1,173
WHO 2007 BMI overweight above +1, 61+ months	0.414 P=142	-0.046 [0.106] N=347	-0.145 [0.105] N=309	**-0.189*** **[0.112]** **N=281**	-0.142 [0.125] N=237	-0.165 [0.145] N=198
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=80	0.012 [0.106] N=347	-0.090 [0.107] N=309	-0.129 [0.117] N=281	-0.083 [0.127] N=237	-0.120 [0.151] N=198
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=35	0.026 [0.068] N=347	0.042 [0.074] N=309	0.019 [0.076] N=281	0.030 [0.079] N=237	0.022 [0.086] N=198

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.016 [0.257] N=1,890	0.016 [0.287] N=1,678	-0.064 [0.326] N=1,437	-0.283 [0.387] N=1,171	-0.356 [0.436] N=895
WHO weight-for-age z score	-0.044 [0.148] N=2,095	0.001 [0.161] N=1,871	**-0.282* [0.168] N=1,598**	-0.286 [0.199] N=1,300	-0.326 [0.245] N=993
WHO weight-for-length/height z score, 0-60 months	-0.133 [0.274] N=1,545	-0.113 [0.289] N=1,378	-0.274 [0.339] N=1,172	-0.171 [0.374] N=960	-0.696 [0.446] N=730
WHO BMI-for-age z score	-0.309 [0.241] N=1,829	-0.360 [0.256] N=1,627	**-0.550* [0.303] N=1,394**	-0.446 [0.329] N=1,137	**-0.920** [0.390] N=873**
Stunted: HAZ below -2	0.007 [0.048] N=1,890	0.019 [0.052] N=1,678	-0.001 [0.057] N=1,437	0.036 [0.066] N=1,171	0.021 [0.084] N=895
Underweight: WAZ below -2	-0.002 [0.026] N=2,095	-0.013 [0.029] N=1,871	0.002 [0.032] N=1,598	-0.005 [0.038] N=1,300	-0.033 [0.043] N=993
Wasted: WLZ/WHZ below -2, 0-60 months	**0.074* [0.039] N=1,545**	0.065 [0.042] N=1,378	**0.103** [0.047] N=1,172**	0.046 [0.052] N=960	**0.111* [0.056] N=730**
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.055] N=1,545	0.015 [0.059] N=1,378	-0.007 [0.065] N=1,172	0.005 [0.077] N=960	-0.059 [0.095] N=730
Obese: WLZ/WHZ above +3, 0-60 months	-0.005 [0.042] N=1,545	-0.017 [0.044] N=1,378	-0.032 [0.050] N=1,172	-0.018 [0.065] N=960	-0.094 [0.083] N=730
CIAF: any stunting, underweight, or wasting, 0-60 months	0.094 [0.064] N=1,496	0.087 [0.071] N=1,332	0.097 [0.079] N=1,133	0.036 [0.095] N=927	0.073 [0.116] N=703
BMI-for-age z score below -2	**0.073** [0.032] N=1,829**	**0.071* [0.036] N=1,627**	**0.083** [0.042] N=1,394**	**0.072* [0.040] N=1,137**	**0.158*** [0.044] N=873**
WHO 2007 BMI overweight above +1, 61+ months	-0.122 [0.151] N=289	**-0.296* [0.158] N=251**	**-0.419** [0.178] N=223**	**-0.387* [0.226] N=179**	**-0.654** [0.308] N=140**
WHO 2007 BMI obesity above +2, 61+ months	0.009 [0.148] N=289	-0.152 [0.154] N=251	-0.247 [0.180] N=223	-0.163 [0.190] N=179	-0.293 [0.292] N=140
WHO 2007 BMI thinness below -2, 61+ months	0.064 [0.080] N=289	0.117 [0.089] N=251	0.075 [0.088] N=223	0.117 [0.083] N=179	0.152 [0.104] N=140

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 71 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.