Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-5 Using 12 November 2022 Child Age

The child cohort uses complete birth dates and completed ages 0 to 71 months on 12 November 2022 without age-boundary assignments. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications are available through month 60 only. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.221 [0.173] N=2,405	0.229 [0.181] N=2,229	0.216 [0.194] N=2,007	**0.358* [0.200] N=1,729**	0.207 [0.227] N=1,423
WHO weight-for-age z score	0.076	-0.043 [0.101] N=2,689	-0.015 [0.109] N=2,493	0.019 [0.111] N=2,246	-0.012 [0.122] N=1,927	-0.044 [0.127] N=1,584
WHO weight-for-length/height z score, 0-60 months	0.342	0.051 [0.157] N=2,029	0.029 [0.160] N=1,884	0.004 [0.169] N=1,697	-0.160 [0.186] N=1,454	-0.082 [0.199] N=1,191
WHO BMI-for-age z score	0.344	-0.004 [0.162] N=2,332	0.006 [0.164] N=2,166	0.041 [0.177] N=1,945	-0.189 [0.184] N=1,673	-0.111 [0.203] N=1,376
Stunted: HAZ below -2	0.228 P=531	-0.033 [0.035] N=2,405	-0.042 [0.036] N=2,229	-0.043 [0.038] N=2,007	**-0.084** [0.040] N=1,729**	-0.046 [0.044] N=1,423
Underweight: WAZ below -2	0.074 P=194	0.015 [0.020] N=2,689	0.005 [0.021] N=2,493	-0.005 [0.022] N=2,246	-0.007 [0.025] N=1,927	-0.002 [0.028] N=1,584
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.024 [0.027] N=2,029	-0.016 [0.027] N=1,884	-0.013 [0.030] N=1,697	-0.001 [0.032] N=1,454	-0.018 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.020 [0.037] N=2,029	0.015 [0.040] N=1,884	0.016 [0.041] N=1,697	-0.010 [0.047] N=1,454	-0.021 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.009 [0.028] N=2,029	-0.012 [0.030] N=1,884	0.001 [0.031] N=1,697	-0.011 [0.035] N=1,454	-0.009 [0.041] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.041 [0.039] N=1,969	-0.047 [0.041] N=1,827	-0.042 [0.043] N=1,645	-0.076 [0.046] N=1,410	-0.057 [0.050] N=1,156
BMI-for-age z score below -2	0.119 P=282	-0.014 [0.025] N=2,332	-0.023 [0.025] N=2,166	-0.020 [0.027] N=1,945	-0.000 [0.028] N=1,673	-0.005 [0.031] N=1,376
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=123	-0.070 [0.122] N=303	-0.078 [0.128] N=283	-0.042 [0.135] N=254	-0.109 [0.147] N=226	-0.143 [0.176] N=187
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=70	-0.054 [0.094] N=303	-0.086 [0.096] N=283	-0.034 [0.104] N=254	-0.104 [0.111] N=226	-0.163 [0.137] N=187
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=33	-0.035 [0.061] N=303	-0.076 [0.060] N=283	**-0.108* [0.063] N=254**	-0.090 [0.067] N=226	**-0.175** [0.088] N=187**

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.070 [0.248] N=2,024	0.056 [0.268] N=1,848	0.027 [0.301] N=1,626	0.332 [0.312] N=1,348	-0.070 [0.418] N=1,042
WHO weight-for-age z score	-0.192 [0.146] N=2,270	-0.161 [0.166] N=2,074	-0.113 [0.175] N=1,827	-0.220 [0.215] N=1,508	-0.347 [0.269] N=1,165
WHO weight-for-length/height z score, 0-60 months	0.078 [0.232] N=1,707	0.043 [0.246] N=1,562	-0.032 [0.275] N=1,375	-0.354 [0.338] N=1,132	-0.255 [0.412] N=869
WHO BMI-for-age z score	0.047 [0.244] N=1,961	0.089 [0.254] N=1,795	0.161 [0.298] N=1,574	-0.270 [0.346] N=1,302	-0.060 [0.439] N=1,005
Stunted: HAZ below -2	0.017 [0.050] N=2,024	0.006 [0.055] N=1,848	0.009 [0.059] N=1,626	-0.065 [0.069] N=1,348	0.041 [0.093] N=1,042
Underweight: WAZ below -2	0.029 [0.028] N=2,270	0.012 [0.031] N=2,074	-0.009 [0.033] N=1,827	-0.010 [0.040] N=1,508	0.001 [0.056] N=1,165
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.037] N=1,707	-0.008 [0.040] N=1,562	0.002 [0.046] N=1,375	0.023 [0.056] N=1,132	-0.020 [0.066] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.044] N=1,707	0.014 [0.048] N=1,562	0.012 [0.052] N=1,375	-0.038 [0.065] N=1,132	-0.097 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.031] N=1,707	-0.024 [0.034] N=1,562	-0.004 [0.036] N=1,375	-0.030 [0.045] N=1,132	-0.042 [0.063] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.006 [0.053] N=1,656	-0.001 [0.060] N=1,514	0.016 [0.064] N=1,332	-0.050 [0.076] N=1,097	-0.004 [0.093] N=843
BMI-for-age z score below -2	-0.030 [0.036] N=1,961	-0.051 [0.035] N=1,795	-0.052 [0.041] N=1,574	-0.031 [0.047] N=1,302	-0.067 [0.058] N=1,005
WHO 2007 BMI overweight above +1, 61+ months	-0.034 [0.168] N=252	-0.018 [0.177] N=232	0.045 [0.190] N=203	-0.070 [0.215] N=175	-0.075 [0.290] N=136
WHO 2007 BMI obesity above +2, 61+ months	0.026 [0.125] N=252	-0.011 [0.130] N=232	0.088 [0.148] N=203	-0.022 [0.172] N=175	-0.085 [0.219] N=136
WHO 2007 BMI thinness below -2, 61+ months	-0.047 [0.093] N=252	-0.131 [0.093] N=232	**-0.188* [0.102] N=203**	-0.185 [0.120] N=175	**-0.464** [0.176] N=136**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.222 [0.174] N=2,405	0.230 [0.181] N=2,229	0.218 [0.195] N=2,007	**0.363* [0.200] N=1,729**	0.213 [0.227] N=1,423
WHO weight-for-age z score	0.076	-0.045 [0.101] N=2,689	-0.017 [0.109] N=2,493	0.018 [0.111] N=2,246	-0.011 [0.122] N=1,927	-0.045 [0.126] N=1,584
WHO weight-for-length/height z score, 0-60 months	0.342	0.049 [0.157] N=2,029	0.027 [0.161] N=1,884	0.001 [0.170] N=1,697	-0.161 [0.187] N=1,454	-0.090 [0.200] N=1,191
WHO BMI-for-age z score	0.344	-0.002 [0.163] N=2,332	0.007 [0.165] N=2,166	0.042 [0.178] N=1,945	-0.186 [0.184] N=1,673	-0.112 [0.203] N=1,376
Stunted: HAZ below -2	0.228 P=531	-0.032 [0.035] N=2,405	-0.041 [0.036] N=2,229	-0.042 [0.038] N=2,007	**-0.083** [0.040] N=1,729**	-0.046 [0.044] N=1,423
Underweight: WAZ below -2	0.074 P=194	0.016 [0.020] N=2,689	0.006 [0.021] N=2,493	-0.005 [0.022] N=2,246	-0.007 [0.025] N=1,927	-0.002 [0.028] N=1,584
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.021 [0.026] N=2,029	-0.013 [0.027] N=1,884	-0.009 [0.030] N=1,697	0.002 [0.032] N=1,454	-0.012 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.021 [0.037] N=2,029	0.017 [0.039] N=1,884	0.018 [0.041] N=1,697	-0.007 [0.047] N=1,454	-0.017 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.007 [0.028] N=2,029	-0.009 [0.030] N=1,884	0.003 [0.031] N=1,697	-0.009 [0.035] N=1,454	-0.006 [0.040] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.038 [0.038] N=1,969	-0.042 [0.040] N=1,827	-0.037 [0.042] N=1,645	-0.071 [0.045] N=1,410	-0.050 [0.049] N=1,156
BMI-for-age z score below -2	0.119 P=282	-0.014 [0.025] N=2,332	-0.022 [0.025] N=2,166	-0.019 [0.027] N=1,945	0.000 [0.028] N=1,673	-0.004 [0.031] N=1,376
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=123	-0.079 [0.122] N=303	-0.088 [0.129] N=283	-0.055 [0.136] N=254	-0.121 [0.148] N=226	-0.153 [0.178] N=187
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=70	-0.064 [0.094] N=303	-0.097 [0.097] N=283	-0.051 [0.105] N=254	-0.120 [0.112] N=226	-0.180 [0.136] N=187
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=33	-0.024 [0.061] N=303	-0.065 [0.060] N=283	-0.093 [0.062] N=254	-0.075 [0.065] N=226	**-0.154* [0.084] N=187**

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.066 [0.248] N=2,024	0.052 [0.268] N=1,848	0.021 [0.300] N=1,626	0.319 [0.313] N=1,348	-0.110 [0.421] N=1,042
WHO weight-for-age z score	-0.192 [0.146] N=2,270	-0.161 [0.166] N=2,074	-0.114 [0.175] N=1,827	-0.222 [0.214] N=1,508	-0.349 [0.267] N=1,165
WHO weight-for-length/height z score, 0-60 months	0.078 [0.232] N=1,707	0.041 [0.247] N=1,562	-0.035 [0.276] N=1,375	-0.356 [0.339] N=1,132	-0.262 [0.412] N=869
WHO BMI-for-age z score	0.047 [0.244] N=1,961	0.089 [0.254] N=1,795	0.161 [0.297] N=1,574	-0.271 [0.346] N=1,302	-0.051 [0.439] N=1,005
Stunted: HAZ below -2	0.017 [0.050] N=2,024	0.006 [0.055] N=1,848	0.009 [0.059] N=1,626	-0.065 [0.069] N=1,348	0.044 [0.093] N=1,042
Underweight: WAZ below -2	0.029 [0.028] N=2,270	0.012 [0.031] N=2,074	-0.009 [0.033] N=1,827	-0.010 [0.040] N=1,508	0.001 [0.056] N=1,165
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.037] N=1,707	-0.007 [0.040] N=1,562	0.003 [0.046] N=1,375	0.024 [0.057] N=1,132	-0.017 [0.066] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.044] N=1,707	0.014 [0.047] N=1,562	0.012 [0.052] N=1,375	-0.037 [0.065] N=1,132	-0.096 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.031] N=1,707	-0.024 [0.035] N=1,562	-0.003 [0.037] N=1,375	-0.030 [0.045] N=1,132	-0.041 [0.064] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.005 [0.053] N=1,656	-0.001 [0.059] N=1,514	0.017 [0.063] N=1,332	-0.050 [0.075] N=1,097	-0.003 [0.093] N=843
BMI-for-age z score below -2	-0.030 [0.036] N=1,961	-0.051 [0.035] N=1,795	-0.052 [0.041] N=1,574	-0.032 [0.047] N=1,302	-0.069 [0.058] N=1,005
WHO 2007 BMI overweight above +1, 61+ months	-0.042 [0.172] N=252	-0.029 [0.181] N=232	0.030 [0.196] N=203	-0.086 [0.222] N=175	-0.051 [0.301] N=136
WHO 2007 BMI obesity above +2, 61+ months	0.012 [0.126] N=252	-0.030 [0.132] N=232	0.052 [0.153] N=203	-0.062 [0.177] N=175	-0.135 [0.227] N=136
WHO 2007 BMI thinness below -2, 61+ months	-0.029 [0.094] N=252	-0.111 [0.093] N=232	-0.155 [0.101] N=203	-0.148 [0.117] N=175	**-0.399** [0.166] N=136**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.201 [0.175] N=2,405	0.206 [0.182] N=2,229	0.191 [0.197] N=2,007	0.331 [0.203] N=1,729	0.169 [0.229] N=1,423
WHO weight-for-age z score	0.076	-0.080 [0.098] N=2,689	-0.051 [0.107] N=2,493	-0.020 [0.109] N=2,246	-0.045 [0.120] N=1,927	-0.084 [0.124] N=1,584
WHO weight-for-length/height z score, 0-60 months	0.342	0.043 [0.160] N=2,029	0.027 [0.164] N=1,884	0.006 [0.172] N=1,697	-0.155 [0.189] N=1,454	-0.089 [0.203] N=1,191
WHO BMI-for-age z score	0.344	-0.002 [0.164] N=2,332	0.011 [0.166] N=2,166	0.047 [0.178] N=1,945	-0.176 [0.185] N=1,673	-0.102 [0.206] N=1,376
Stunted: HAZ below -2	0.228 P=531	-0.026 [0.034] N=2,405	-0.033 [0.036] N=2,229	-0.035 [0.038] N=2,007	**-0.075* [0.040] N=1,729**	-0.035 [0.044] N=1,423
Underweight: WAZ below -2	0.074 P=194	0.022 [0.019] N=2,689	0.013 [0.021] N=2,493	0.003 [0.021] N=2,246	0.002 [0.024] N=1,927	0.009 [0.027] N=1,584
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.018 [0.026] N=2,029	-0.009 [0.027] N=1,884	-0.006 [0.030] N=1,697	0.005 [0.032] N=1,454	-0.009 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.024 [0.038] N=2,029	0.021 [0.039] N=1,884	0.022 [0.041] N=1,697	-0.003 [0.047] N=1,454	-0.013 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.007 [0.028] N=2,029	-0.008 [0.030] N=1,884	0.005 [0.031] N=1,697	-0.007 [0.035] N=1,454	-0.006 [0.041] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.026 [0.037] N=1,969	-0.027 [0.039] N=1,827	-0.025 [0.041] N=1,645	-0.062 [0.044] N=1,410	-0.035 [0.049] N=1,156
BMI-for-age z score below -2	0.119 P=282	-0.010 [0.025] N=2,332	-0.018 [0.025] N=2,166	-0.015 [0.028] N=1,945	0.003 [0.028] N=1,673	-0.000 [0.031] N=1,376
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=123	-0.083 [0.123] N=303	-0.094 [0.128] N=283	-0.061 [0.136] N=254	-0.131 [0.149] N=226	-0.156 [0.181] N=187
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=70	-0.049 [0.094] N=303	-0.085 [0.097] N=283	-0.033 [0.104] N=254	-0.112 [0.108] N=226	-0.165 [0.136] N=187
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=33	-0.010 [0.060] N=303	-0.050 [0.059] N=283	-0.076 [0.062] N=254	-0.047 [0.065] N=226	-0.129 [0.085] N=187

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.051 [0.247] N=2,024	0.026 [0.265] N=1,848	-0.008 [0.298] N=1,626	0.271 [0.314] N=1,348	-0.168 [0.413] N=1,042
WHO weight-for-age z score	-0.221 [0.142] N=2,270	-0.190 [0.163] N=2,074	-0.145 [0.171] N=1,827	-0.248 [0.210] N=1,508	-0.387 [0.262] N=1,165
WHO weight-for-length/height z score, 0-60 months	0.071 [0.235] N=1,707	0.044 [0.253] N=1,562	-0.026 [0.277] N=1,375	-0.356 [0.339] N=1,132	-0.275 [0.414] N=869
WHO BMI-for-age z score	0.051 [0.247] N=1,961	0.099 [0.257] N=1,795	0.177 [0.298] N=1,574	-0.245 [0.346] N=1,302	-0.019 [0.440] N=1,005
Stunted: HAZ below -2	0.021 [0.049] N=2,024	0.013 [0.054] N=1,848	0.015 [0.058] N=1,626	-0.058 [0.067] N=1,348	0.055 [0.089] N=1,042
Underweight: WAZ below -2	0.034 [0.027] N=2,270	0.018 [0.031] N=2,074	-0.003 [0.032] N=1,827	-0.006 [0.039] N=1,508	0.010 [0.055] N=1,165
Wasted: WLZ/WHZ below -2, 0-60 months	-0.016 [0.038] N=1,707	-0.001 [0.041] N=1,562	0.006 [0.047] N=1,375	0.028 [0.057] N=1,132	-0.011 [0.068] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.045] N=1,707	0.020 [0.049] N=1,562	0.016 [0.053] N=1,375	-0.035 [0.065] N=1,132	-0.088 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.016 [0.031] N=1,707	-0.022 [0.035] N=1,562	-0.002 [0.037] N=1,375	-0.030 [0.045] N=1,132	-0.044 [0.062] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.019 [0.051] N=1,656	0.018 [0.057] N=1,514	0.026 [0.060] N=1,332	-0.048 [0.072] N=1,097	0.011 [0.087] N=843
BMI-for-age z score below -2	-0.027 [0.036] N=1,961	-0.047 [0.036] N=1,795	-0.049 [0.042] N=1,574	-0.030 [0.047] N=1,302	-0.066 [0.060] N=1,005
WHO 2007 BMI overweight above +1, 61+ months	-0.046 [0.170] N=252	-0.036 [0.181] N=232	0.024 [0.194] N=203	-0.104 [0.221] N=175	-0.043 [0.301] N=136
WHO 2007 BMI obesity above +2, 61+ months	0.024 [0.124] N=252	-0.017 [0.131] N=232	0.072 [0.150] N=203	-0.057 [0.176] N=175	-0.107 [0.233] N=136
WHO 2007 BMI thinness below -2, 61+ months	-0.013 [0.093] N=252	-0.100 [0.093] N=232	-0.142 [0.104] N=203	-0.108 [0.117] N=175	**-0.369** [0.166] N=136**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.373	0.170 [0.172] N=2,405	0.183 [0.180] N=2,229	0.167 [0.194] N=2,007	0.294 [0.201] N=1,729	0.122 [0.225] N=1,423
WHO weight-for-age z score	0.076	-0.081 [0.098] N=2,689	-0.060 [0.107] N=2,493	-0.031 [0.110] N=2,246	-0.065 [0.121] N=1,927	-0.120 [0.127] N=1,584
WHO weight-for-length/height z score, 0-60 months	0.342	0.057 [0.159] N=2,029	0.033 [0.162] N=1,884	0.009 [0.171] N=1,697	-0.151 [0.189] N=1,454	-0.092 [0.201] N=1,191
WHO BMI-for-age z score	0.344	0.012 [0.165] N=2,332	0.019 [0.165] N=2,166	0.059 [0.178] N=1,945	-0.163 [0.187] N=1,673	-0.092 [0.207] N=1,376
Stunted: HAZ below -2	0.228 P=531	-0.021 [0.034] N=2,405	-0.029 [0.036] N=2,229	-0.029 [0.038] N=2,007	-0.065 [0.040] N=1,729	-0.025 [0.043] N=1,423
Underweight: WAZ below -2	0.074 P=194	0.023 [0.019] N=2,689	0.015 [0.021] N=2,493	0.006 [0.021] N=2,246	0.005 [0.024] N=1,927	0.013 [0.027] N=1,584
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.018 [0.026] N=2,029	-0.009 [0.027] N=1,884	-0.005 [0.030] N=1,697	0.006 [0.032] N=1,454	-0.010 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.027 [0.037] N=2,029	0.024 [0.039] N=1,884	0.024 [0.041] N=1,697	-0.000 [0.047] N=1,454	-0.014 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.006 [0.028] N=2,029	-0.010 [0.029] N=1,884	0.003 [0.031] N=1,697	-0.007 [0.034] N=1,454	-0.007 [0.040] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.020 [0.037] N=1,969	-0.021 [0.039] N=1,827	-0.016 [0.041] N=1,645	-0.049 [0.044] N=1,410	-0.024 [0.048] N=1,156
BMI-for-age z score below -2	0.119 P=282	-0.011 [0.026] N=2,332	-0.019 [0.025] N=2,166	-0.016 [0.028] N=1,945	0.004 [0.028] N=1,673	-0.001 [0.031] N=1,376
WHO 2007 BMI overweight above +1, 61+ months	0.399 P=123	-0.068 [0.122] N=303	-0.087 [0.129] N=283	-0.052 [0.136] N=254	-0.107 [0.147] N=226	-0.131 [0.178] N=187
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=70	-0.046 [0.097] N=303	-0.090 [0.100] N=283	-0.044 [0.107] N=254	-0.121 [0.117] N=226	-0.178 [0.144] N=187
WHO 2007 BMI thinness below -2, 61+ months	0.113 P=33	-0.028 [0.060] N=303	-0.060 [0.060] N=283	-0.100 [0.061] N=254	-0.081 [0.063] N=226	**-0.139* [0.080] N=187**

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.033 [0.244] N=2,024	0.025 [0.262] N=1,848	-0.001 [0.294] N=1,626	0.288 [0.310] N=1,348	-0.189 [0.401] N=1,042
WHO weight-for-age z score	-0.208 [0.143] N=2,270	-0.186 [0.163] N=2,074	-0.146 [0.172] N=1,827	-0.243 [0.213] N=1,508	-0.372 [0.264] N=1,165
WHO weight-for-length/height z score, 0-60 months	0.079 [0.234] N=1,707	0.047 [0.249] N=1,562	-0.028 [0.279] N=1,375	-0.352 [0.343] N=1,132	-0.252 [0.415] N=869
WHO BMI-for-age z score	0.058 [0.246] N=1,961	0.097 [0.255] N=1,795	0.171 [0.299] N=1,574	-0.252 [0.348] N=1,302	-0.021 [0.442] N=1,005
Stunted: HAZ below -2	0.023 [0.049] N=2,024	0.013 [0.054] N=1,848	0.014 [0.058] N=1,626	-0.061 [0.067] N=1,348	0.056 [0.088] N=1,042
Underweight: WAZ below -2	0.031 [0.027] N=2,270	0.017 [0.031] N=2,074	-0.002 [0.032] N=1,827	-0.006 [0.039] N=1,508	0.010 [0.054] N=1,165
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.038] N=1,707	-0.004 [0.040] N=1,562	0.006 [0.047] N=1,375	0.029 [0.058] N=1,132	-0.010 [0.069] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.026 [0.045] N=1,707	0.021 [0.048] N=1,562	0.016 [0.053] N=1,375	-0.035 [0.065] N=1,132	-0.087 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.014 [0.031] N=1,707	-0.021 [0.035] N=1,562	-0.001 [0.037] N=1,375	-0.029 [0.045] N=1,132	-0.042 [0.061] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.020 [0.051] N=1,656	0.015 [0.057] N=1,514	0.027 [0.061] N=1,332	-0.049 [0.072] N=1,097	0.011 [0.088] N=843
BMI-for-age z score below -2	-0.027 [0.036] N=1,961	-0.049 [0.036] N=1,795	-0.051 [0.042] N=1,574	-0.029 [0.047] N=1,302	-0.065 [0.060] N=1,005
WHO 2007 BMI overweight above +1, 61+ months	-0.025 [0.169] N=252	-0.028 [0.183] N=232	0.030 [0.195] N=203	-0.083 [0.215] N=175	0.006 [0.313] N=136
WHO 2007 BMI obesity above +2, 61+ months	0.036 [0.127] N=252	-0.021 [0.131] N=232	0.047 [0.148] N=203	-0.072 [0.175] N=175	-0.093 [0.236] N=136
WHO 2007 BMI thinness below -2, 61+ months	-0.028 [0.092] N=252	-0.096 [0.094] N=232	-0.154 [0.098] N=203	-0.143 [0.113] N=175	**-0.352** [0.152] N=136**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.378	0.192 [0.177] N=2,246	0.177 [0.187] N=2,087	0.143 [0.200] N=1,874	0.262 [0.204] N=1,623	0.075 [0.225] N=1,332
WHO weight-for-age z score	0.076	-0.014 [0.103] N=2,506	0.008 [0.111] N=2,328	0.043 [0.115] N=2,092	0.006 [0.126] N=1,803	-0.006 [0.133] N=1,481
WHO weight-for-length/height z score, 0-60 months	0.350	0.171 [0.160] N=1,893	0.147 [0.163] N=1,765	0.130 [0.171] N=1,587	-0.027 [0.192] N=1,370	0.081 [0.214] N=1,120
WHO BMI-for-age z score	0.356	0.096 [0.168] N=2,180	0.099 [0.169] N=2,030	0.159 [0.181] N=1,820	-0.056 [0.191] N=1,575	0.058 [0.214] N=1,293
Stunted: HAZ below -2	0.229 P=497	-0.028 [0.037] N=2,246	-0.036 [0.039] N=2,087	-0.035 [0.040] N=1,874	**-0.071* [0.043] N=1,623**	-0.025 [0.048] N=1,332
Underweight: WAZ below -2	0.074 P=181	0.012 [0.021] N=2,506	0.003 [0.023] N=2,328	-0.007 [0.024] N=2,092	-0.007 [0.027] N=1,803	-0.001 [0.031] N=1,481
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=213	-0.034 [0.027] N=1,893	-0.025 [0.027] N=1,765	-0.019 [0.029] N=1,587	-0.007 [0.032] N=1,370	-0.021 [0.033] N=1,120
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=368	0.033 [0.039] N=1,893	0.029 [0.041] N=1,765	0.034 [0.043] N=1,587	0.007 [0.049] N=1,370	0.006 [0.058] N=1,120
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=179	-0.007 [0.030] N=1,893	-0.007 [0.032] N=1,765	0.010 [0.033] N=1,587	-0.001 [0.036] N=1,370	0.005 [0.043] N=1,120
CIAF: any stunting, underweight, or wasting, 0-60 months	0.315 P=566	-0.039 [0.041] N=1,836	-0.041 [0.043] N=1,711	-0.038 [0.045] N=1,537	-0.074 [0.049] N=1,328	-0.046 [0.053] N=1,086
BMI-for-age z score below -2	0.117 P=258	-0.018 [0.026] N=2,180	-0.026 [0.025] N=2,030	-0.026 [0.028] N=1,820	-0.008 [0.029] N=1,575	-0.018 [0.032] N=1,293
WHO 2007 BMI overweight above +1, 61+ months	0.396 P=116	-0.106 [0.129] N=288	-0.110 [0.137] N=268	-0.074 [0.146] N=240	-0.156 [0.155] N=213	-0.159 [0.193] N=176
WHO 2007 BMI obesity above +2, 61+ months	0.219 P=64	-0.088 [0.100] N=288	-0.114 [0.103] N=268	-0.070 [0.113] N=240	-0.151 [0.118] N=213	-0.205 [0.150] N=176
WHO 2007 BMI thinness below -2, 61+ months	0.110 P=30	-0.013 [0.054] N=288	-0.060 [0.054] N=268	**-0.091* [0.053] N=240**	-0.073 [0.057] N=213	**-0.156** [0.072] N=176**

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.149 [0.258] N=1,891	0.113 [0.285] N=1,732	0.059 [0.319] N=1,519	0.353 [0.344] N=1,268	-0.107 [0.468] N=977
WHO weight-for-age z score	-0.178 [0.145] N=2,116	-0.162 [0.161] N=1,938	-0.129 [0.175] N=1,702	-0.264 [0.215] N=1,413	-0.355 [0.267] N=1,091
WHO weight-for-length/height z score, 0-60 months	0.107 [0.239] N=1,591	0.045 [0.251] N=1,463	-0.020 [0.278] N=1,285	-0.326 [0.347] N=1,068	-0.130 [0.425] N=818
WHO BMI-for-age z score	0.031 [0.253] N=1,834	0.044 [0.261] N=1,684	0.145 [0.303] N=1,474	-0.272 [0.354] N=1,229	0.008 [0.448] N=947
Stunted: HAZ below -2	0.001 [0.052] N=1,891	-0.013 [0.058] N=1,732	-0.010 [0.063] N=1,519	-0.086 [0.075] N=1,268	0.036 [0.102] N=977
Underweight: WAZ below -2	0.026 [0.029] N=2,116	0.011 [0.032] N=1,938	-0.007 [0.035] N=1,702	0.000 [0.042] N=1,413	0.015 [0.059] N=1,091
Wasted: WLZ/WHZ below -2, 0-60 months	-0.036 [0.038] N=1,591	-0.021 [0.040] N=1,463	-0.008 [0.046] N=1,285	0.007 [0.056] N=1,068	-0.047 [0.061] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.045] N=1,591	0.010 [0.048] N=1,463	0.015 [0.052] N=1,285	-0.040 [0.065] N=1,068	-0.071 [0.090] N=818
Obese: WLZ/WHZ above +3, 0-60 months	-0.028 [0.034] N=1,591	-0.034 [0.038] N=1,463	-0.007 [0.040] N=1,285	-0.030 [0.048] N=1,068	-0.032 [0.067] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.012 [0.056] N=1,543	-0.020 [0.062] N=1,418	-0.008 [0.068] N=1,244	-0.091 [0.080] N=1,035	-0.040 [0.102] N=793
BMI-for-age z score below -2	-0.021 [0.036] N=1,834	-0.041 [0.034] N=1,684	-0.045 [0.040] N=1,474	-0.030 [0.046] N=1,229	-0.085 [0.055] N=947
WHO 2007 BMI overweight above +1, 61+ months	-0.079 [0.189] N=241	-0.056 [0.202] N=221	0.005 [0.224] N=193	-0.136 [0.245] N=166	-0.023 [0.331] N=129
WHO 2007 BMI obesity above +2, 61+ months	-0.031 [0.134] N=241	-0.060 [0.143] N=221	0.013 [0.164] N=193	-0.129 [0.185] N=166	-0.217 [0.249] N=129
WHO 2007 BMI thinness below -2, 61+ months	0.012 [0.098] N=241	-0.082 [0.098] N=221	-0.130 [0.104] N=193	-0.132 [0.125] N=166	**-0.456*** [0.167] N=129**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.378	0.199 [0.176] N=2,246	0.183 [0.186] N=2,087	0.145 [0.200] N=1,874	0.267 [0.205] N=1,623	0.084 [0.225] N=1,332
WHO weight-for-age z score	0.076	-0.005 [0.103] N=2,506	0.013 [0.111] N=2,328	0.046 [0.114] N=2,092	0.010 [0.125] N=1,803	-0.004 [0.134] N=1,481
WHO weight-for-length/height z score, 0-60 months	0.350	0.169 [0.160] N=1,893	0.146 [0.162] N=1,765	0.123 [0.170] N=1,587	-0.033 [0.191] N=1,370	0.065 [0.213] N=1,120
WHO BMI-for-age z score	0.356	0.100 [0.167] N=2,180	0.103 [0.168] N=2,030	0.160 [0.180] N=1,820	-0.056 [0.190] N=1,575	0.050 [0.214] N=1,293
Stunted: HAZ below -2	0.229 P=497	-0.029 [0.037] N=2,246	-0.037 [0.039] N=2,087	-0.037 [0.040] N=1,874	**-0.074* [0.043] N=1,623**	-0.028 [0.047] N=1,332
Underweight: WAZ below -2	0.074 P=181	0.009 [0.021] N=2,506	0.001 [0.023] N=2,328	-0.009 [0.024] N=2,092	-0.009 [0.026] N=1,803	-0.003 [0.031] N=1,481
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=213	-0.035 [0.027] N=1,893	-0.025 [0.027] N=1,765	-0.019 [0.029] N=1,587	-0.008 [0.032] N=1,370	-0.024 [0.033] N=1,120
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=368	0.033 [0.039] N=1,893	0.029 [0.040] N=1,765	0.031 [0.043] N=1,587	0.003 [0.049] N=1,370	0.000 [0.057] N=1,120
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=179	-0.007 [0.029] N=1,893	-0.007 [0.031] N=1,765	0.009 [0.033] N=1,587	-0.002 [0.036] N=1,370	0.003 [0.043] N=1,120
CIAF: any stunting, underweight, or wasting, 0-60 months	0.315 P=566	-0.039 [0.041] N=1,836	-0.041 [0.043] N=1,711	-0.037 [0.045] N=1,537	-0.073 [0.049] N=1,328	-0.045 [0.052] N=1,086
BMI-for-age z score below -2	0.117 P=258	-0.019 [0.026] N=2,180	-0.026 [0.025] N=2,030	-0.026 [0.028] N=1,820	-0.008 [0.029] N=1,575	-0.018 [0.032] N=1,293
WHO 2007 BMI overweight above +1, 61+ months	0.396 P=116	-0.095 [0.125] N=288	-0.094 [0.134] N=268	-0.068 [0.142] N=240	-0.143 [0.151] N=213	-0.155 [0.189] N=176
WHO 2007 BMI obesity above +2, 61+ months	0.219 P=64	-0.087 [0.096] N=288	-0.113 [0.100] N=268	-0.064 [0.110] N=240	-0.147 [0.114] N=213	-0.188 [0.147] N=176
WHO 2007 BMI thinness below -2, 61+ months	0.110 P=30	-0.030 [0.054] N=288	-0.073 [0.053] N=268	**-0.100* [0.053] N=240**	-0.082 [0.055] N=213	**-0.168** [0.071] N=176**

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.156 [0.257] N=1,891	0.113 [0.285] N=1,732	0.063 [0.319] N=1,519	0.356 [0.346] N=1,268	-0.070 [0.467] N=977
WHO weight-for-age z score	-0.162 [0.144] N=2,116	-0.151 [0.161] N=1,938	-0.120 [0.172] N=1,702	-0.248 [0.213] N=1,413	-0.331 [0.265] N=1,091
WHO weight-for-length/height z score, 0-60 months	0.132 [0.239] N=1,591	0.078 [0.251] N=1,463	-0.014 [0.279] N=1,285	-0.322 [0.351] N=1,068	-0.146 [0.433] N=818
WHO BMI-for-age z score	0.051 [0.251] N=1,834	0.066 [0.259] N=1,684	0.152 [0.301] N=1,474	-0.263 [0.352] N=1,229	-0.002 [0.448] N=947
Stunted: HAZ below -2	-0.001 [0.052] N=1,891	-0.013 [0.058] N=1,732	-0.013 [0.063] N=1,519	-0.091 [0.074] N=1,268	0.023 [0.101] N=977
Underweight: WAZ below -2	0.022 [0.029] N=2,116	0.008 [0.032] N=1,938	-0.011 [0.034] N=1,702	-0.005 [0.041] N=1,413	0.008 [0.058] N=1,091
Wasted: WLZ/WHZ below -2, 0-60 months	-0.037 [0.038] N=1,591	-0.022 [0.040] N=1,463	-0.008 [0.046] N=1,285	0.005 [0.056] N=1,068	-0.052 [0.062] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.030 [0.045] N=1,591	0.020 [0.048] N=1,463	0.017 [0.052] N=1,285	-0.037 [0.067] N=1,068	-0.073 [0.091] N=818
Obese: WLZ/WHZ above +3, 0-60 months	-0.026 [0.033] N=1,591	-0.032 [0.037] N=1,463	-0.006 [0.039] N=1,285	-0.031 [0.048] N=1,068	-0.035 [0.067] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.012 [0.056] N=1,543	-0.020 [0.063] N=1,418	-0.009 [0.068] N=1,244	-0.098 [0.080] N=1,035	-0.060 [0.099] N=793
BMI-for-age z score below -2	-0.023 [0.036] N=1,834	-0.043 [0.034] N=1,684	-0.046 [0.040] N=1,474	-0.030 [0.046] N=1,229	-0.084 [0.056] N=947
WHO 2007 BMI overweight above +1, 61+ months	-0.065 [0.179] N=241	-0.035 [0.194] N=221	0.014 [0.211] N=193	-0.125 [0.232] N=166	-0.055 [0.332] N=129
WHO 2007 BMI obesity above +2, 61+ months	-0.044 [0.126] N=241	-0.075 [0.134] N=221	0.001 [0.155] N=193	-0.156 [0.174] N=166	-0.229 [0.246] N=129
WHO 2007 BMI thinness below -2, 61+ months	-0.017 [0.098] N=241	-0.105 [0.099] N=221	-0.151 [0.105] N=193	-0.140 [0.122] N=166	**-0.432** [0.170] N=129**

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.300	-0.269 [0.197] N=2,314	-0.302 [0.208] N=2,092	-0.331 [0.219] N=1,835	**-0.424*** **[0.238]** **N=1,560**	**-0.514*** **[0.268]** **N=1,269**
WHO weight-for-age z score	0.123	-0.075 [0.106] N=2,569	-0.063 [0.112] N=2,334	-0.184 [0.116] N=2,042	-0.143 [0.127] N=1,730	-0.143 [0.143] N=1,400
WHO weight-for-length/height z score, 0-60 months	0.376	0.091 [0.182] N=1,886	0.119 [0.189] N=1,707	0.079 [0.210] N=1,489	0.200 [0.223] N=1,268	0.172 [0.264] N=1,022
WHO BMI-for-age z score	0.368	0.064 [0.196] N=2,242	0.052 [0.207] N=2,027	0.030 [0.230] N=1,780	0.201 [0.242] N=1,513	0.163 [0.271] N=1,230
Stunted: HAZ below -2	0.220 P=510	0.046 [0.037] N=2,314	0.046 [0.039] N=2,092	0.043 [0.041] N=1,835	0.057 [0.046] N=1,560	0.063 [0.053] N=1,269
Underweight: WAZ below -2	0.068 P=172	0.000 [0.019] N=2,569	-0.002 [0.020] N=2,334	0.010 [0.020] N=2,042	0.010 [0.021] N=1,730	0.010 [0.022] N=1,400
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.025 [0.029] N=1,886	0.020 [0.030] N=1,707	0.035 [0.032] N=1,489	-0.003 [0.032] N=1,268	0.002 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.047 [0.035] N=1,886	0.048 [0.036] N=1,707	0.043 [0.039] N=1,489	0.058 [0.042] N=1,268	0.061 [0.047] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.018 [0.029] N=1,886	0.014 [0.030] N=1,707	0.011 [0.033] N=1,489	0.024 [0.036] N=1,268	0.010 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.073*** **[0.041]** **N=1,830**	0.064 [0.043] N=1,654	**0.077*** **[0.045]** **N=1,441**	0.044 [0.046] N=1,227	0.059 [0.051] N=989
BMI-for-age z score below -2	0.119 P=268	0.032 [0.032] N=2,242	0.031 [0.035] N=2,027	0.029 [0.038] N=1,780	0.007 [0.039] N=1,513	0.018 [0.044] N=1,230
WHO 2007 BMI overweight above +1, 61+ months	0.410 P=145	-0.098 [0.105] N=359	**-0.204*** **[0.106]** **N=322**	**-0.198*** **[0.110]** **N=292**	-0.113 [0.120] N=247	-0.156 [0.130] N=206
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=83	0.021 [0.097] N=359	-0.071 [0.099] N=322	-0.091 [0.109] N=292	-0.009 [0.119] N=247	-0.060 [0.131] N=206
WHO 2007 BMI thinness below -2, 61+ months	0.111 P=39	0.035 [0.069] N=359	0.042 [0.075] N=322	0.022 [0.075] N=292	0.017 [0.082] N=247	0.012 [0.093] N=206

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.038 [0.257] N=1,984	-0.037 [0.287] N=1,762	-0.049 [0.316] N=1,505	-0.227 [0.362] N=1,230	-0.315 [0.481] N=939
WHO weight-for-age z score	-0.090 [0.150] N=2,209	-0.064 [0.164] N=1,974	**-0.318* [0.174] N=1,682**	-0.299 [0.200] N=1,370	-0.346 [0.251] N=1,040
WHO weight-for-length/height z score, 0-60 months	-0.165 [0.261] N=1,630	-0.164 [0.274] N=1,451	-0.323 [0.322] N=1,233	-0.191 [0.354] N=1,012	-0.566 [0.429] N=766
WHO BMI-for-age z score	-0.328 [0.233] N=1,927	**-0.432* [0.249] N=1,712**	**-0.544* [0.299] N=1,465**	-0.330 [0.336] N=1,198	**-0.720* [0.375] N=915**
Stunted: HAZ below -2	0.028 [0.050] N=1,984	0.026 [0.053] N=1,762	0.011 [0.059] N=1,505	0.041 [0.065] N=1,230	0.041 [0.091] N=939
Underweight: WAZ below -2	-0.002 [0.029] N=2,209	-0.004 [0.033] N=1,974	0.018 [0.035] N=1,682	0.020 [0.041] N=1,370	0.021 [0.048] N=1,040
Wasted: WLZ/WHZ below -2, 0-60 months	**0.081** [0.039] N=1,630**	**0.079* [0.042] N=1,451**	**0.125*** [0.047] N=1,233**	0.067 [0.048] N=1,012	**0.117** [0.055] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.036 [0.052] N=1,630	0.030 [0.056] N=1,451	0.016 [0.063] N=1,233	0.040 [0.071] N=1,012	0.006 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.013 [0.040] N=1,630	0.001 [0.041] N=1,451	-0.010 [0.047] N=1,233	0.008 [0.059] N=1,012	-0.081 [0.077] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.115* [0.063] N=1,582**	0.105 [0.070] N=1,406	**0.145* [0.080] N=1,193**	0.098 [0.088] N=979	0.140 [0.115] N=741
BMI-for-age z score below -2	**0.093*** [0.032] N=1,927**	**0.104*** [0.037] N=1,712**	**0.111** [0.043] N=1,465**	**0.084** [0.041] N=1,198**	**0.138*** [0.047] N=915**
WHO 2007 BMI overweight above +1, 61+ months	-0.232 [0.164] N=297	**-0.460*** [0.166] N=260**	**-0.498*** [0.187] N=230**	**-0.429* [0.255] N=185**	**-0.754** [0.318] N=144**
WHO 2007 BMI obesity above +2, 61+ months	0.031 [0.141] N=297	-0.141 [0.148] N=260	-0.213 [0.183] N=230	-0.074 [0.217] N=185	-0.206 [0.305] N=144
WHO 2007 BMI thinness below -2, 61+ months	0.109 [0.078] N=297	**0.157* [0.088] N=260**	0.115 [0.089] N=230	0.144 [0.093] N=185	**0.205* [0.112] N=144**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.300	-0.270 [0.196] N=2,314	-0.304 [0.208] N=2,092	-0.331 [0.219] N=1,835	**-0.427* [0.238] N=1,560**	**-0.508* [0.267] N=1,269**
WHO weight-for-age z score	0.123	-0.075 [0.106] N=2,569	-0.063 [0.112] N=2,334	-0.184 [0.116] N=2,042	-0.143 [0.128] N=1,730	-0.143 [0.143] N=1,400
WHO weight-for-length/height z score, 0-60 months	0.376	0.097 [0.181] N=1,886	0.122 [0.188] N=1,707	0.082 [0.210] N=1,489	0.201 [0.223] N=1,268	0.169 [0.261] N=1,022
WHO BMI-for-age z score	0.368	0.064 [0.196] N=2,242	0.052 [0.207] N=2,027	0.031 [0.230] N=1,780	0.201 [0.242] N=1,513	0.162 [0.271] N=1,230
Stunted: HAZ below -2	0.220 P=510	0.045 [0.038] N=2,314	0.046 [0.039] N=2,092	0.043 [0.041] N=1,835	0.057 [0.046] N=1,560	0.063 [0.053] N=1,269
Underweight: WAZ below -2	0.068 P=172	0.000 [0.019] N=2,569	-0.002 [0.020] N=2,334	0.010 [0.020] N=2,042	0.010 [0.021] N=1,730	0.010 [0.022] N=1,400
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.028] N=1,886	0.018 [0.030] N=1,707	0.034 [0.032] N=1,489	-0.004 [0.032] N=1,268	0.003 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.046 [0.035] N=1,886	0.047 [0.037] N=1,707	0.041 [0.039] N=1,489	0.057 [0.042] N=1,268	0.062 [0.048] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.016 [0.029] N=1,886	0.012 [0.030] N=1,707	0.009 [0.033] N=1,489	0.023 [0.037] N=1,268	0.011 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.071* [0.042] N=1,830**	0.060 [0.043] N=1,654	0.074 [0.046] N=1,441	0.043 [0.047] N=1,227	0.060 [0.051] N=989
BMI-for-age z score below -2	0.119 P=268	0.032 [0.032] N=2,242	0.031 [0.035] N=2,027	0.030 [0.038] N=1,780	0.007 [0.039] N=1,513	0.018 [0.044] N=1,230
WHO 2007 BMI overweight above +1, 61+ months	0.410 P=145	-0.073 [0.104] N=359	**-0.184* [0.105] N=322**	**-0.183* [0.109] N=292**	-0.108 [0.120] N=247	-0.144 [0.131] N=206
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=83	0.045 [0.101] N=359	-0.052 [0.102] N=322	-0.072 [0.113] N=292	-0.004 [0.121] N=247	-0.051 [0.134] N=206
WHO 2007 BMI thinness below -2, 61+ months	0.111 P=39	0.023 [0.069] N=359	0.032 [0.075] N=322	0.007 [0.076] N=292	0.011 [0.084] N=247	-0.006 [0.096] N=206

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.042 [0.257] N=1,984	-0.046 [0.286] N=1,762	-0.059 [0.314] N=1,505	-0.256 [0.360] N=1,230	-0.341 [0.481] N=939
WHO weight-for-age z score	-0.090 [0.151] N=2,209	-0.065 [0.163] N=1,974	**-0.320* [0.173] N=1,682**	-0.302 [0.200] N=1,370	-0.347 [0.250] N=1,040
WHO weight-for-length/height z score, 0-60 months	-0.147 [0.263] N=1,630	-0.152 [0.277] N=1,451	-0.315 [0.325] N=1,233	-0.181 [0.356] N=1,012	-0.538 [0.436] N=766
WHO BMI-for-age z score	-0.327 [0.234] N=1,927	**-0.432* [0.251] N=1,712**	**-0.546* [0.300] N=1,465**	-0.329 [0.337] N=1,198	**-0.713* [0.380] N=915**
Stunted: HAZ below -2	0.028 [0.050] N=1,984	0.024 [0.053] N=1,762	0.009 [0.058] N=1,505	0.039 [0.065] N=1,230	0.041 [0.091] N=939
Underweight: WAZ below -2	-0.002 [0.029] N=2,209	-0.005 [0.033] N=1,974	0.018 [0.035] N=1,682	0.018 [0.041] N=1,370	0.020 [0.048] N=1,040
Wasted: WLZ/WHZ below -2, 0-60 months	**0.074* [0.039] N=1,630**	**0.071* [0.042] N=1,451**	**0.116** [0.047] N=1,233**	0.057 [0.049] N=1,012	**0.106* [0.056] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.035 [0.052] N=1,630	0.026 [0.056] N=1,451	0.010 [0.063] N=1,233	0.032 [0.071] N=1,012	0.002 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.009 [0.040] N=1,630	-0.005 [0.041] N=1,451	-0.018 [0.047] N=1,233	-0.001 [0.059] N=1,012	-0.087 [0.077] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.107* [0.064] N=1,582**	0.093 [0.070] N=1,406	0.131 [0.081] N=1,193	0.085 [0.090] N=979	0.129 [0.116] N=741
BMI-for-age z score below -2	**0.092*** [0.032] N=1,927**	**0.101*** [0.037] N=1,712**	**0.109** [0.043] N=1,465**	**0.081* [0.042] N=1,198**	**0.137*** [0.048] N=915**
WHO 2007 BMI overweight above +1, 61+ months	-0.196 [0.162] N=297	**-0.432*** [0.163] N=260**	**-0.477** [0.185] N=230**	**-0.425* [0.254] N=185**	**-0.733** [0.322] N=144**
WHO 2007 BMI obesity above +2, 61+ months	0.069 [0.148] N=297	-0.110 [0.154] N=260	-0.179 [0.190] N=230	-0.067 [0.224] N=185	-0.177 [0.317] N=144
WHO 2007 BMI thinness below -2, 61+ months	0.101 [0.082] N=297	0.150 [0.093] N=260	0.099 [0.092] N=230	0.140 [0.092] N=185	0.171 [0.117] N=144

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.300	-0.273 [0.192] N=2,314	-0.297 [0.202] N=2,092	-0.317 [0.216] N=1,835	**-0.414* [0.232] N=1,560**	**-0.494* [0.258] N=1,269**
WHO weight-for-age z score	0.123	-0.073 [0.105] N=2,569	-0.059 [0.111] N=2,334	-0.171 [0.116] N=2,042	-0.140 [0.129] N=1,730	-0.140 [0.145] N=1,400
WHO weight-for-length/height z score, 0-60 months	0.376	0.105 [0.183] N=1,886	0.127 [0.191] N=1,707	0.093 [0.213] N=1,489	0.202 [0.223] N=1,268	0.168 [0.260] N=1,022
WHO BMI-for-age z score	0.368	0.065 [0.196] N=2,242	0.052 [0.207] N=2,027	0.036 [0.228] N=1,780	0.199 [0.241] N=1,513	0.154 [0.271] N=1,230
Stunted: HAZ below -2	0.220 P=510	0.046 [0.036] N=2,314	0.044 [0.038] N=2,092	0.037 [0.040] N=1,835	0.052 [0.044] N=1,560	0.057 [0.050] N=1,269
Underweight: WAZ below -2	0.068 P=172	0.001 [0.019] N=2,569	-0.001 [0.020] N=2,334	0.008 [0.020] N=2,042	0.009 [0.021] N=1,730	0.009 [0.022] N=1,400
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.029] N=1,886	0.016 [0.030] N=1,707	0.031 [0.033] N=1,489	-0.005 [0.033] N=1,268	0.002 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.048 [0.035] N=1,886	0.048 [0.037] N=1,707	0.042 [0.040] N=1,489	0.056 [0.043] N=1,268	0.062 [0.048] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.018 [0.030] N=1,886	0.013 [0.030] N=1,707	0.009 [0.033] N=1,489	0.022 [0.037] N=1,268	0.010 [0.041] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.075* [0.041] N=1,830**	0.060 [0.043] N=1,654	0.070 [0.046] N=1,441	0.040 [0.046] N=1,227	0.057 [0.050] N=989
BMI-for-age z score below -2	0.119 P=268	0.032 [0.032] N=2,242	0.031 [0.035] N=2,027	0.028 [0.038] N=1,780	0.006 [0.039] N=1,513	0.017 [0.045] N=1,230
WHO 2007 BMI overweight above +1, 61+ months	0.410 P=145	-0.083 [0.104] N=359	**-0.194* [0.105] N=322**	**-0.195* [0.110] N=292**	-0.117 [0.118] N=247	-0.173 [0.134] N=206
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=83	0.024 [0.099] N=359	-0.065 [0.102] N=322	-0.095 [0.113] N=292	-0.033 [0.119] N=247	-0.084 [0.134] N=206
WHO 2007 BMI thinness below -2, 61+ months	0.111 P=39	0.022 [0.067] N=359	0.027 [0.072] N=322	0.008 [0.071] N=292	0.006 [0.079] N=247	-0.024 [0.088] N=206

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.030 [0.252] N=1,984	-0.022 [0.279] N=1,762	-0.015 [0.306] N=1,505	-0.202 [0.349] N=1,230	-0.269 [0.454] N=939
WHO weight-for-age z score	-0.088 [0.150] N=2,209	-0.060 [0.163] N=1,974	**-0.294* [0.175] N=1,682**	-0.290 [0.201] N=1,370	-0.334 [0.250] N=1,040
WHO weight-for-length/height z score, 0-60 months	-0.150 [0.264] N=1,630	-0.154 [0.280] N=1,451	-0.312 [0.326] N=1,233	-0.196 [0.353] N=1,012	-0.551 [0.430] N=766
WHO BMI-for-age z score	-0.334 [0.233] N=1,927	**-0.437* [0.250] N=1,712**	**-0.553* [0.297] N=1,465**	-0.350 [0.334] N=1,198	**-0.754** [0.378] N=915**
Stunted: HAZ below -2	0.025 [0.049] N=1,984	0.019 [0.053] N=1,762	-0.002 [0.057] N=1,505	0.026 [0.063] N=1,230	0.023 [0.086] N=939
Underweight: WAZ below -2	-0.003 [0.029] N=2,209	-0.006 [0.033] N=1,974	0.011 [0.036] N=1,682	0.013 [0.041] N=1,370	0.014 [0.048] N=1,040
Wasted: WLZ/WHZ below -2, 0-60 months	**0.075* [0.040] N=1,630**	0.069 [0.043] N=1,451	**0.112** [0.048] N=1,233**	0.054 [0.050] N=1,012	**0.105* [0.057] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.037 [0.053] N=1,630	0.028 [0.057] N=1,451	0.007 [0.064] N=1,233	0.026 [0.073] N=1,012	0.000 [0.091] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.010 [0.040] N=1,630	-0.006 [0.042] N=1,451	-0.019 [0.048] N=1,233	-0.004 [0.060] N=1,012	-0.089 [0.078] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.111* [0.063] N=1,582**	0.092 [0.071] N=1,406	0.119 [0.082] N=1,193	0.072 [0.090] N=979	0.123 [0.112] N=741
BMI-for-age z score below -2	**0.091*** [0.032] N=1,927**	**0.099*** [0.037] N=1,712**	**0.104** [0.043] N=1,465**	**0.077* [0.042] N=1,198**	**0.131*** [0.049] N=915**
WHO 2007 BMI overweight above +1, 61+ months	-0.211 [0.159] N=297	**-0.444*** [0.160] N=260**	**-0.497*** [0.182] N=230**	**-0.424* [0.250] N=185**	**-0.885*** [0.313] N=144**
WHO 2007 BMI obesity above +2, 61+ months	0.048 [0.146] N=297	-0.123 [0.152] N=260	-0.202 [0.189] N=230	-0.085 [0.218] N=185	-0.284 [0.320] N=144
WHO 2007 BMI thinness below -2, 61+ months	0.104 [0.080] N=297	**0.149* [0.090] N=260**	0.103 [0.089] N=230	0.120 [0.092] N=185	0.102 [0.131] N=144

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.300	-0.270 [0.193] N=2,314	-0.291 [0.204] N=2,092	-0.307 [0.218] N=1,835	**-0.400* [0.232] N=1,560**	**-0.481* [0.257] N=1,269**
WHO weight-for-age z score	0.123	-0.078 [0.104] N=2,569	-0.061 [0.110] N=2,334	-0.171 [0.115] N=2,042	-0.130 [0.126] N=1,730	-0.141 [0.141] N=1,400
WHO weight-for-length/height z score, 0-60 months	0.376	0.107 [0.182] N=1,886	0.130 [0.188] N=1,707	0.091 [0.210] N=1,489	0.214 [0.222] N=1,268	0.182 [0.260] N=1,022
WHO BMI-for-age z score	0.368	0.062 [0.196] N=2,242	0.049 [0.207] N=2,027	0.032 [0.228] N=1,780	0.203 [0.239] N=1,513	0.156 [0.268] N=1,230
Stunted: HAZ below -2	0.220 P=510	0.047 [0.037] N=2,314	0.045 [0.038] N=2,092	0.039 [0.041] N=1,835	0.052 [0.045] N=1,560	0.058 [0.051] N=1,269
Underweight: WAZ below -2	0.068 P=172	0.001 [0.018] N=2,569	-0.002 [0.020] N=2,334	0.007 [0.020] N=2,042	0.008 [0.021] N=1,730	0.009 [0.022] N=1,400
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.029] N=1,886	0.017 [0.030] N=1,707	0.034 [0.032] N=1,489	-0.004 [0.032] N=1,268	0.001 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.047 [0.035] N=1,886	0.047 [0.036] N=1,707	0.042 [0.039] N=1,489	0.058 [0.042] N=1,268	0.063 [0.047] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.020 [0.029] N=1,886	0.015 [0.030] N=1,707	0.012 [0.033] N=1,489	0.027 [0.036] N=1,268	0.014 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.074* [0.041] N=1,830**	0.060 [0.043] N=1,654	0.070 [0.045] N=1,441	0.039 [0.046] N=1,227	0.060 [0.050] N=989
BMI-for-age z score below -2	0.119 P=268	0.031 [0.032] N=2,242	0.029 [0.035] N=2,027	0.027 [0.038] N=1,780	0.005 [0.039] N=1,513	0.013 [0.044] N=1,230
WHO 2007 BMI overweight above +1, 61+ months	0.410 P=145	-0.078 [0.105] N=359	**-0.184* [0.107] N=322**	-0.176 [0.112] N=292	-0.091 [0.121] N=247	-0.132 [0.136] N=206
WHO 2007 BMI obesity above +2, 61+ months	0.242 P=83	0.023 [0.103] N=359	-0.072 [0.105] N=322	-0.095 [0.116] N=292	-0.029 [0.124] N=247	-0.090 [0.137] N=206
WHO 2007 BMI thinness below -2, 61+ months	0.111 P=39	0.019 [0.069] N=359	0.026 [0.074] N=322	-0.005 [0.074] N=292	-0.004 [0.080] N=247	-0.055 [0.089] N=206

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033 [0.252] N=1,984	-0.018 [0.282] N=1,762	0.006 [0.312] N=1,505	-0.172 [0.355] N=1,230	-0.275 [0.456] N=939
WHO weight-for-age z score	-0.076 [0.147] N=2,209	-0.041 [0.160] N=1,974	-0.267 [0.172] N=1,682	-0.233 [0.194] N=1,370	-0.295 [0.236] N=1,040
WHO weight-for-length/height z score, 0-60 months	-0.141 [0.262] N=1,630	-0.151 [0.274] N=1,451	-0.322 [0.317] N=1,233	-0.192 [0.347] N=1,012	-0.533 [0.430] N=766
WHO BMI-for-age z score	-0.336 [0.233] N=1,927	**-0.442* [0.249] N=1,712**	**-0.558* [0.296] N=1,465**	-0.337 [0.330] N=1,198	**-0.721* [0.378] N=915**
Stunted: HAZ below -2	0.027 [0.048] N=1,984	0.020 [0.052] N=1,762	-0.003 [0.057] N=1,505	0.022 [0.063] N=1,230	0.029 [0.087] N=939
Underweight: WAZ below -2	-0.006 [0.028] N=2,209	-0.010 [0.033] N=1,974	0.008 [0.036] N=1,682	0.008 [0.041] N=1,370	0.011 [0.049] N=1,040
Wasted: WLZ/WHZ below -2, 0-60 months	**0.075* [0.039] N=1,630**	**0.071* [0.042] N=1,451**	**0.118** [0.047] N=1,233**	0.058 [0.049] N=1,012	**0.109* [0.055] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.035 [0.052] N=1,630	0.024 [0.055] N=1,451	0.005 [0.062] N=1,233	0.024 [0.071] N=1,012	0.001 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.012 [0.040] N=1,630	-0.004 [0.041] N=1,451	-0.017 [0.047] N=1,233	0.001 [0.059] N=1,012	-0.086 [0.078] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.108* [0.062] N=1,582**	0.088 [0.069] N=1,406	0.113 [0.080] N=1,193	0.064 [0.087] N=979	0.126 [0.107] N=741
BMI-for-age z score below -2	**0.093*** [0.032] N=1,927**	**0.100*** [0.037] N=1,712**	**0.107** [0.043] N=1,465**	**0.080* [0.042] N=1,198**	**0.134*** [0.048] N=915**
WHO 2007 BMI overweight above +1, 61+ months	-0.211 [0.157] N=297	**-0.438*** [0.158] N=260**	**-0.472*** [0.177] N=230**	-0.403 [0.245] N=185	**-0.770** [0.296] N=144**
WHO 2007 BMI obesity above +2, 61+ months	0.057 [0.146] N=297	-0.123 [0.152] N=260	-0.196 [0.186] N=230	-0.080 [0.222] N=185	-0.234 [0.323] N=144
WHO 2007 BMI thinness below -2, 61+ months	0.114 [0.084] N=297	**0.161* [0.094] N=260**	0.095 [0.096] N=230	0.130 [0.097] N=185	0.077 [0.119] N=144

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.309	-0.290 [0.196] N=2,167	-0.309 [0.210] N=1,959	-0.313 [0.220] N=1,721	**-0.410* [0.235] N=1,459**	**-0.544** [0.262] N=1,189**
WHO weight-for-age z score	0.125	-0.067 [0.106] N=2,401	-0.042 [0.111] N=2,180	-0.159 [0.114] N=1,911	-0.129 [0.126] N=1,617	-0.166 [0.139] N=1,313
WHO weight-for-length/height z score, 0-60 months	0.378	0.136 [0.192] N=1,762	0.176 [0.200] N=1,595	0.099 [0.221] N=1,392	0.197 [0.233] N=1,184	0.133 [0.270] N=957
WHO BMI-for-age z score	0.382	0.045 [0.202] N=2,101	0.036 [0.214] N=1,900	-0.029 [0.237] N=1,671	0.121 [0.251] N=1,418	0.049 [0.283] N=1,156
Stunted: HAZ below -2	0.220 P=479	0.053 [0.036] N=2,167	0.055 [0.038] N=1,959	0.048 [0.039] N=1,721	0.060 [0.042] N=1,459	0.072 [0.047] N=1,189
Underweight: WAZ below -2	0.067 P=160	-0.002 [0.019] N=2,401	-0.004 [0.020] N=2,180	0.001 [0.020] N=1,911	0.006 [0.021] N=1,617	0.013 [0.023] N=1,313
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=196	0.026 [0.029] N=1,762	0.017 [0.030] N=1,595	0.036 [0.032] N=1,392	0.003 [0.033] N=1,184	0.012 [0.037] N=957
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.046 [0.035] N=1,762	0.048 [0.037] N=1,595	0.036 [0.038] N=1,392	0.047 [0.041] N=1,184	0.046 [0.047] N=957
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=180	0.016 [0.029] N=1,762	0.012 [0.030] N=1,595	0.007 [0.032] N=1,392	0.023 [0.036] N=1,184	0.013 [0.039] N=957
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=529	**0.087** [0.043] N=1,709**	0.074 [0.046] N=1,545	**0.088* [0.047] N=1,347**	0.051 [0.048] N=1,145	0.074 [0.052] N=925
BMI-for-age z score below -2	0.116 P=244	0.039 [0.032] N=2,101	0.038 [0.034] N=1,900	0.039 [0.037] N=1,671	0.023 [0.037] N=1,418	0.035 [0.040] N=1,156
WHO 2007 BMI overweight above +1, 61+ months	0.418 P=141	-0.120 [0.108] N=343	**-0.255** [0.108] N=308**	**-0.249** [0.114] N=280**	-0.191 [0.130] N=236	**-0.254* [0.140] N=197**
WHO 2007 BMI obesity above +2, 61+ months	0.245 P=80	0.011 [0.108] N=343	-0.101 [0.108] N=308	-0.127 [0.120] N=280	-0.058 [0.132] N=236	-0.110 [0.149] N=197
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=34	0.038 [0.071] N=343	0.046 [0.078] N=308	0.019 [0.079] N=280	0.024 [0.087] N=236	0.008 [0.097] N=197

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.001 [0.259] N=1,858	0.037 [0.293] N=1,650	0.059 [0.324] N=1,412	-0.074 [0.361] N=1,150	-0.198 [0.470] N=880
WHO weight-for-age z score	-0.081 [0.153] N=2,065	-0.033 [0.167] N=1,844	**-0.298* [0.176] N=1,575**	-0.267 [0.206] N=1,281	-0.327 [0.261] N=977
WHO weight-for-length/height z score, 0-60 months	-0.108 [0.276] N=1,522	-0.102 [0.294] N=1,355	-0.334 [0.343] N=1,152	-0.232 [0.369] N=944	-0.609 [0.456] N=717
WHO BMI-for-age z score	-0.315 [0.240] N=1,804	-0.414 [0.261] N=1,603	**-0.589* [0.313] N=1,374**	-0.396 [0.342] N=1,121	**-0.819** [0.389] N=859**
Stunted: HAZ below -2	0.019 [0.052] N=1,858	0.016 [0.057] N=1,650	-0.003 [0.062] N=1,412	0.012 [0.066] N=1,150	0.013 [0.093] N=880
Underweight: WAZ below -2	-0.008 [0.029] N=2,065	-0.011 [0.033] N=1,844	0.004 [0.035] N=1,575	0.009 [0.043] N=1,281	0.012 [0.051] N=977
Wasted: WLZ/WHZ below -2, 0-60 months	**0.077* [0.041] N=1,522**	0.068 [0.044] N=1,355	**0.115** [0.048] N=1,152**	0.060 [0.053] N=944	**0.105* [0.058] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.054] N=1,522	0.022 [0.057] N=1,355	-0.004 [0.063] N=1,152	0.004 [0.072] N=944	-0.022 [0.093] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.000 [0.042] N=1,522	-0.016 [0.044] N=1,355	-0.033 [0.049] N=1,152	-0.014 [0.062] N=944	-0.089 [0.083] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.111 [0.069] N=1,477	0.094 [0.077] N=1,313	0.128 [0.087] N=1,115	0.053 [0.100] N=913	0.091 [0.130] N=693
BMI-for-age z score below -2	**0.095*** [0.034] N=1,804**	**0.104*** [0.040] N=1,603**	**0.115** [0.046] N=1,374**	**0.099** [0.044] N=1,121**	**0.149*** [0.048] N=859**
WHO 2007 BMI overweight above +1, 61+ months	-0.227 [0.164] N=284	**-0.487*** [0.167] N=249**	**-0.512*** [0.188] N=221**	**-0.471* [0.258] N=177**	**-0.826** [0.325] N=138**
WHO 2007 BMI obesity above +2, 61+ months	0.051 [0.152] N=284	-0.131 [0.161] N=249	-0.198 [0.195] N=221	-0.045 [0.227] N=177	-0.144 [0.332] N=138
WHO 2007 BMI thinness below -2, 61+ months	0.104 [0.085] N=284	0.154 [0.097] N=249	0.102 [0.096] N=221	0.141 [0.091] N=177	0.148 [0.108] N=138

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.309	-0.296 [0.196] N=2,167	-0.315 [0.211] N=1,959	-0.318 [0.221] N=1,721	**-0.410* [0.237] N=1,459**	**-0.541** [0.262] N=1,189**
WHO weight-for-age z score	0.125	-0.067 [0.106] N=2,401	-0.043 [0.112] N=2,180	-0.164 [0.114] N=1,911	-0.130 [0.127] N=1,617	-0.164 [0.141] N=1,313
WHO weight-for-length/height z score, 0-60 months	0.378	0.143 [0.192] N=1,762	0.179 [0.200] N=1,595	0.105 [0.221] N=1,392	0.213 [0.232] N=1,184	0.155 [0.272] N=957
WHO BMI-for-age z score	0.382	0.048 [0.203] N=2,101	0.038 [0.214] N=1,900	-0.027 [0.237] N=1,671	0.120 [0.252] N=1,418	0.049 [0.284] N=1,156
Stunted: HAZ below -2	0.220 P=479	0.052 [0.036] N=2,167	0.054 [0.038] N=1,959	0.048 [0.039] N=1,721	0.059 [0.042] N=1,459	0.070 [0.047] N=1,189
Underweight: WAZ below -2	0.067 P=160	-0.003 [0.019] N=2,401	-0.004 [0.020] N=2,180	0.002 [0.020] N=1,911	0.006 [0.021] N=1,617	0.012 [0.023] N=1,313
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=196	0.023 [0.029] N=1,762	0.015 [0.030] N=1,595	0.033 [0.032] N=1,392	-0.003 [0.033] N=1,184	0.005 [0.037] N=957
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.045 [0.035] N=1,762	0.046 [0.036] N=1,595	0.034 [0.038] N=1,392	0.045 [0.040] N=1,184	0.045 [0.046] N=957
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=180	0.015 [0.029] N=1,762	0.011 [0.029] N=1,595	0.005 [0.032] N=1,392	0.021 [0.035] N=1,184	0.011 [0.039] N=957
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=529	**0.081* [0.043] N=1,709**	0.069 [0.045] N=1,545	**0.082* [0.047] N=1,347**	0.043 [0.048] N=1,145	0.065 [0.051] N=925
BMI-for-age z score below -2	0.116 P=244	0.038 [0.032] N=2,101	0.037 [0.034] N=1,900	0.037 [0.037] N=1,671	0.021 [0.037] N=1,418	0.034 [0.040] N=1,156
WHO 2007 BMI overweight above +1, 61+ months	0.418 P=141	-0.124 [0.108] N=343	**-0.259** [0.106] N=308**	**-0.260** [0.112] N=280**	-0.195 [0.127] N=236	**-0.252* [0.136] N=197**
WHO 2007 BMI obesity above +2, 61+ months	0.245 P=80	0.012 [0.107] N=343	-0.100 [0.107] N=308	-0.126 [0.119] N=280	-0.063 [0.130] N=236	-0.119 [0.146] N=197
WHO 2007 BMI thinness below -2, 61+ months	0.102 P=34	0.033 [0.071] N=343	0.048 [0.076] N=308	0.011 [0.078] N=280	0.019 [0.083] N=236	0.004 [0.091] N=197

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.016 [0.257] N=1,858	0.022 [0.289] N=1,650	0.042 [0.318] N=1,412	-0.125 [0.361] N=1,150	-0.265 [0.462] N=880
WHO weight-for-age z score	-0.079 [0.153] N=2,065	-0.034 [0.167] N=1,844	**-0.305* [0.176] N=1,575**	-0.276 [0.206] N=1,281	-0.376 [0.260] N=977
WHO weight-for-length/height z score, 0-60 months	-0.108 [0.275] N=1,522	-0.101 [0.291] N=1,355	-0.330 [0.340] N=1,152	-0.201 [0.371] N=944	-0.604 [0.449] N=717
WHO BMI-for-age z score	-0.308 [0.240] N=1,804	-0.407 [0.259] N=1,603	**-0.585* [0.311] N=1,374**	-0.370 [0.346] N=1,121	**-0.805** [0.389] N=859**
Stunted: HAZ below -2	0.022 [0.052] N=1,858	0.019 [0.056] N=1,650	0.002 [0.061] N=1,412	0.023 [0.066] N=1,150	0.033 [0.091] N=880
Underweight: WAZ below -2	-0.009 [0.029] N=2,065	-0.011 [0.033] N=1,844	0.006 [0.035] N=1,575	0.012 [0.042] N=1,281	0.023 [0.050] N=977
Wasted: WLZ/WHZ below -2, 0-60 months	**0.076* [0.040] N=1,522**	0.069 [0.044] N=1,355	**0.116** [0.048] N=1,152**	0.055 [0.051] N=944	**0.106* [0.056] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.053] N=1,522	0.022 [0.056] N=1,355	-0.004 [0.062] N=1,152	0.008 [0.071] N=944	-0.023 [0.090] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.001 [0.042] N=1,522	-0.015 [0.043] N=1,355	-0.033 [0.049] N=1,152	-0.013 [0.062] N=944	-0.091 [0.080] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.112 [0.068] N=1,477	0.097 [0.076] N=1,313	0.133 [0.086] N=1,115	0.063 [0.099] N=913	0.115 [0.127] N=693
BMI-for-age z score below -2	**0.094*** [0.034] N=1,804**	**0.104*** [0.039] N=1,603**	**0.115** [0.045] N=1,374**	**0.096** [0.044] N=1,121**	**0.152*** [0.047] N=859**
WHO 2007 BMI overweight above +1, 61+ months	-0.218 [0.163] N=284	**-0.469*** [0.163] N=249**	**-0.516*** [0.186] N=221**	**-0.459* [0.252] N=177**	**-0.787** [0.308] N=138**
WHO 2007 BMI obesity above +2, 61+ months	0.060 [0.151] N=284	-0.122 [0.159] N=249	-0.189 [0.194] N=221	-0.056 [0.223] N=177	-0.180 [0.320] N=138
WHO 2007 BMI thinness below -2, 61+ months	0.099 [0.086] N=284	0.154 [0.095] N=249	0.085 [0.096] N=221	0.129 [0.088] N=177	0.125 [0.108] N=138

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 71 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.