# Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-6 Using 10 October 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 83 months on 10 October 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

### Father RDD primary analysis

*Specification 1 Monthly child-age fixed effects  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.055 | 0.078 [0.134] N=2,853 | 0.064 [0.137] N=2,640 | 0.021 [0.149] N=2,365 | 0.086 [0.151] N=2,033 | 0.112 [0.166] N=1,679 |
| WHO weight-for-age z score | 0.360 | 0.037 [0.086] N=3,026 | 0.074 [0.095] N=2,793 | 0.123 [0.097] N=2,499 | 0.037 [0.103] N=2,144 | 0.068 [0.117] N=1,769 |
| WHO weight-for-length/height z score, 0-60 months | 0.336 | 0.197 [0.183] N=1,613 | 0.217 [0.189] N=1,488 | 0.182 [0.203] N=1,328 | -0.019 [0.221] N=1,128 | 0.076 [0.232] N=917 |
| WHO BMI-for-age z score | 0.356 | 0.059 [0.149] N=2,731 | 0.067 [0.155] N=2,531 | 0.084 [0.165] N=2,260 | -0.136 [0.178] N=1,943 | -0.107 [0.192] N=1,604 |
| Stunted: HAZ below -2 | 0.180 P=494 | -0.019 [0.027] N=2,853 | -0.013 [0.028] N=2,640 | 0.000 [0.032] N=2,365 | -0.022 [0.033] N=2,033 | -0.011 [0.036] N=1,679 |
| Underweight: WAZ below -2 | 0.046 P=135 | -0.008 [0.014] N=3,026 | -0.012 [0.015] N=2,793 | -0.021 [0.016] N=2,499 | -0.019 [0.018] N=2,144 | -0.030 [0.020] N=1,769 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=179 | -0.025 [0.030] N=1,613 | -0.026 [0.032] N=1,488 | -0.017 [0.035] N=1,328 | -0.003 [0.041] N=1,128 | -0.017 [0.042] N=917 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.194 P=302 | **0.073* [0.041] N=1,613** | **0.075* [0.044] N=1,488** | **0.080* [0.047] N=1,328** | 0.034 [0.052] N=1,128 | 0.043 [0.056] N=917 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.082 P=131 | 0.030 [0.028] N=1,613 | 0.036 [0.030] N=1,488 | 0.045 [0.034] N=1,328 | 0.016 [0.037] N=1,128 | 0.017 [0.044] N=917 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.242 P=375 | -0.030 [0.040] N=1,583 | -0.032 [0.041] N=1,462 | 0.002 [0.045] N=1,306 | -0.009 [0.049] N=1,107 | 0.007 [0.052] N=902 |
| BMI-for-age z score below -2 | 0.114 P=318 | 0.016 [0.028] N=2,731 | 0.017 [0.029] N=2,531 | 0.022 [0.030] N=2,260 | 0.045 [0.032] N=1,943 | 0.052 [0.034] N=1,604 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.374 P=417 | 0.035 [0.050] N=1,111 | 0.023 [0.052] N=1,034 | 0.064 [0.057] N=929 | 0.041 [0.062] N=815 | 0.010 [0.069] N=681 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.213 P=238 | 0.009 [0.043] N=1,111 | -0.012 [0.044] N=1,034 | -0.003 [0.048] N=929 | -0.033 [0.049] N=815 | **-0.098* [0.055] N=681** |
| WHO 2007 BMI thinness below -2, 61+ months | 0.109 P=127 | 0.034 [0.042] N=1,111 | 0.042 [0.043] N=1,034 | 0.021 [0.046] N=929 | 0.032 [0.050] N=815 | 0.034 [0.055] N=681 |

*Specification 1 Monthly child-age fixed effects 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.062 [0.200] N=2,398 | -0.117 [0.212] N=2,185 | -0.252 [0.241] N=1,910 | -0.144 [0.275] N=1,578 | -0.280 [0.360] N=1,224 |
| WHO weight-for-age z score | -0.063 [0.111] N=2,551 | -0.007 [0.128] N=2,318 | 0.057 [0.135] N=2,024 | -0.125 [0.151] N=1,669 | -0.131 [0.202] N=1,294 |
| WHO weight-for-length/height z score, 0-60 months | 0.169 [0.260] N=1,371 | 0.212 [0.280] N=1,246 | 0.120 [0.327] N=1,086 | -0.347 [0.379] N=886 | -0.245 [0.461] N=675 |
| WHO BMI-for-age z score | 0.248 [0.189] N=2,301 | 0.298 [0.200] N=2,101 | 0.376 [0.229] N=1,830 | 0.038 [0.276] N=1,513 | 0.228 [0.353] N=1,174 |
| Stunted: HAZ below -2 | 0.028 [0.039] N=2,398 | 0.047 [0.041] N=2,185 | **0.087* [0.049] N=1,910** | 0.063 [0.056] N=1,578 | **0.140* [0.071] N=1,224** |
| Underweight: WAZ below -2 | 0.015 [0.019] N=2,551 | 0.012 [0.021] N=2,318 | 0.003 [0.024] N=2,024 | 0.012 [0.028] N=1,669 | 0.004 [0.036] N=1,294 |
| Wasted: WLZ/WHZ below -2, 0-60 months | -0.008 [0.039] N=1,371 | -0.011 [0.042] N=1,246 | 0.009 [0.048] N=1,086 | 0.037 [0.063] N=886 | 0.021 [0.066] N=675 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.068 [0.055] N=1,371 | 0.072 [0.060] N=1,246 | 0.082 [0.068] N=1,086 | -0.026 [0.083] N=886 | -0.027 [0.109] N=675 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.048 [0.032] N=1,371 | **0.062* [0.034] N=1,246** | **0.086** [0.040] N=1,086** | 0.028 [0.048] N=886 | 0.042 [0.066] N=675 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.027 [0.054] N=1,344 | 0.027 [0.057] N=1,223 | **0.107* [0.064] N=1,067** | 0.101 [0.078] N=868 | **0.214** [0.090] N=663** |
| BMI-for-age z score below -2 | -0.011 [0.032] N=2,301 | -0.016 [0.034] N=2,101 | -0.014 [0.038] N=1,830 | 0.012 [0.043] N=1,513 | 0.013 [0.050] N=1,174 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.102 [0.073] N=923 | 0.092 [0.080] N=846 | **0.193** [0.092] N=741** | **0.197* [0.106] N=627** | 0.192 [0.138] N=493 |
| WHO 2007 BMI obesity above +2, 61+ months | **0.107* [0.063] N=923** | 0.083 [0.068] N=846 | **0.131* [0.078] N=741** | 0.124 [0.089] N=627 | 0.028 [0.112] N=493 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.013 [0.046] N=923 | 0.023 [0.047] N=846 | -0.020 [0.056] N=741 | -0.017 [0.060] N=627 | -0.034 [0.075] N=493 |

*Specification 2 Monthly fixed effects and female  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.055 | 0.088 [0.134] N=2,853 | 0.073 [0.137] N=2,640 | 0.034 [0.149] N=2,365 | 0.100 [0.149] N=2,033 | 0.122 [0.166] N=1,679 |
| WHO weight-for-age z score | 0.360 | 0.044 [0.086] N=3,026 | 0.080 [0.095] N=2,793 | 0.132 [0.097] N=2,499 | 0.045 [0.103] N=2,144 | 0.071 [0.117] N=1,769 |
| WHO weight-for-length/height z score, 0-60 months | 0.336 | 0.191 [0.184] N=1,613 | 0.210 [0.190] N=1,488 | 0.173 [0.204] N=1,328 | -0.022 [0.223] N=1,128 | 0.069 [0.233] N=917 |
| WHO BMI-for-age z score | 0.356 | 0.057 [0.149] N=2,731 | 0.063 [0.155] N=2,531 | 0.081 [0.167] N=2,260 | -0.137 [0.179] N=1,943 | -0.110 [0.192] N=1,604 |
| Stunted: HAZ below -2 | 0.180 P=494 | -0.019 [0.028] N=2,853 | -0.013 [0.028] N=2,640 | -0.000 [0.032] N=2,365 | -0.022 [0.033] N=2,033 | -0.012 [0.037] N=1,679 |
| Underweight: WAZ below -2 | 0.046 P=135 | -0.008 [0.014] N=3,026 | -0.012 [0.015] N=2,793 | -0.021 [0.016] N=2,499 | -0.019 [0.018] N=2,144 | -0.030 [0.021] N=1,769 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=179 | -0.023 [0.030] N=1,613 | -0.024 [0.032] N=1,488 | -0.014 [0.036] N=1,328 | -0.001 [0.041] N=1,128 | -0.014 [0.042] N=917 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.194 P=302 | **0.073* [0.041] N=1,613** | **0.076* [0.044] N=1,488** | **0.081* [0.047] N=1,328** | 0.036 [0.052] N=1,128 | 0.045 [0.056] N=917 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.082 P=131 | 0.030 [0.028] N=1,613 | 0.037 [0.030] N=1,488 | 0.045 [0.033] N=1,328 | 0.016 [0.037] N=1,128 | 0.017 [0.044] N=917 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.242 P=375 | -0.029 [0.040] N=1,583 | -0.030 [0.041] N=1,462 | 0.005 [0.045] N=1,306 | -0.007 [0.049] N=1,107 | 0.009 [0.052] N=902 |
| BMI-for-age z score below -2 | 0.114 P=318 | 0.017 [0.028] N=2,731 | 0.019 [0.029] N=2,531 | 0.024 [0.030] N=2,260 | 0.047 [0.032] N=1,943 | 0.053 [0.034] N=1,604 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.374 P=417 | 0.035 [0.050] N=1,111 | 0.022 [0.052] N=1,034 | 0.065 [0.057] N=929 | 0.041 [0.062] N=815 | 0.010 [0.069] N=681 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.213 P=238 | 0.012 [0.043] N=1,111 | -0.008 [0.044] N=1,034 | 0.002 [0.048] N=929 | -0.030 [0.050] N=815 | **-0.097* [0.055] N=681** |
| WHO 2007 BMI thinness below -2, 61+ months | 0.109 P=127 | 0.034 [0.042] N=1,111 | 0.042 [0.043] N=1,034 | 0.022 [0.047] N=929 | 0.032 [0.050] N=815 | 0.035 [0.055] N=681 |

*Specification 2 Monthly fixed effects and female 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.050 [0.199] N=2,398 | -0.106 [0.211] N=2,185 | -0.230 [0.241] N=1,910 | -0.121 [0.270] N=1,578 | -0.270 [0.358] N=1,224 |
| WHO weight-for-age z score | -0.053 [0.110] N=2,551 | 0.001 [0.127] N=2,318 | 0.074 [0.134] N=2,024 | -0.111 [0.147] N=1,669 | -0.128 [0.199] N=1,294 |
| WHO weight-for-length/height z score, 0-60 months | 0.158 [0.261] N=1,371 | 0.197 [0.281] N=1,246 | 0.095 [0.327] N=1,086 | -0.361 [0.380] N=886 | -0.283 [0.467] N=675 |
| WHO BMI-for-age z score | 0.245 [0.190] N=2,301 | 0.292 [0.201] N=2,101 | 0.370 [0.231] N=1,830 | 0.034 [0.277] N=1,513 | 0.220 [0.355] N=1,174 |
| Stunted: HAZ below -2 | 0.028 [0.039] N=2,398 | 0.046 [0.041] N=2,185 | **0.087* [0.049] N=1,910** | 0.062 [0.056] N=1,578 | **0.139* [0.071] N=1,224** |
| Underweight: WAZ below -2 | 0.014 [0.019] N=2,551 | 0.012 [0.021] N=2,318 | 0.002 [0.024] N=2,024 | 0.011 [0.028] N=1,669 | 0.004 [0.037] N=1,294 |
| Wasted: WLZ/WHZ below -2, 0-60 months | -0.006 [0.039] N=1,371 | -0.008 [0.042] N=1,246 | 0.015 [0.049] N=1,086 | 0.041 [0.063] N=886 | 0.029 [0.067] N=675 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.068 [0.055] N=1,371 | 0.072 [0.060] N=1,246 | 0.083 [0.068] N=1,086 | -0.024 [0.083] N=886 | -0.022 [0.108] N=675 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.048 [0.032] N=1,371 | **0.062* [0.034] N=1,246** | **0.085** [0.040] N=1,086** | 0.028 [0.048] N=886 | 0.042 [0.066] N=675 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.028 [0.054] N=1,344 | 0.029 [0.058] N=1,223 | **0.110* [0.065] N=1,067** | 0.103 [0.078] N=868 | **0.216** [0.091] N=663** |
| BMI-for-age z score below -2 | -0.010 [0.032] N=2,301 | -0.014 [0.034] N=2,101 | -0.011 [0.038] N=1,830 | 0.014 [0.043] N=1,513 | 0.015 [0.050] N=1,174 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.102 [0.072] N=923 | 0.091 [0.079] N=846 | **0.194** [0.092] N=741** | **0.197* [0.106] N=627** | 0.195 [0.137] N=493 |
| WHO 2007 BMI obesity above +2, 61+ months | **0.112* [0.063] N=923** | 0.088 [0.068] N=846 | **0.140* [0.078] N=741** | 0.127 [0.090] N=627 | 0.024 [0.112] N=493 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.013 [0.046] N=923 | 0.022 [0.047] N=846 | -0.021 [0.056] N=741 | -0.017 [0.061] N=627 | -0.034 [0.075] N=493 |

*Specification 3 Monthly fixed effects female and household income groups  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.055 | 0.053 [0.137] N=2,853 | 0.040 [0.140] N=2,640 | 0.004 [0.154] N=2,365 | 0.068 [0.156] N=2,033 | 0.074 [0.174] N=1,679 |
| WHO weight-for-age z score | 0.360 | 0.007 [0.084] N=3,026 | 0.045 [0.094] N=2,793 | 0.094 [0.096] N=2,499 | 0.009 [0.101] N=2,144 | 0.032 [0.116] N=1,769 |
| WHO weight-for-length/height z score, 0-60 months | 0.336 | 0.164 [0.184] N=1,613 | 0.189 [0.190] N=1,488 | 0.158 [0.201] N=1,328 | -0.029 [0.221] N=1,128 | 0.066 [0.232] N=917 |
| WHO BMI-for-age z score | 0.356 | 0.052 [0.150] N=2,731 | 0.059 [0.157] N=2,531 | 0.078 [0.167] N=2,260 | -0.138 [0.179] N=1,943 | -0.106 [0.195] N=1,604 |
| Stunted: HAZ below -2 | 0.180 P=494 | -0.014 [0.028] N=2,853 | -0.008 [0.028] N=2,640 | 0.005 [0.032] N=2,365 | -0.017 [0.033] N=2,033 | -0.005 [0.038] N=1,679 |
| Underweight: WAZ below -2 | 0.046 P=135 | -0.003 [0.014] N=3,026 | -0.007 [0.014] N=2,793 | -0.015 [0.016] N=2,499 | -0.013 [0.017] N=2,144 | -0.021 [0.020] N=1,769 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=179 | -0.014 [0.030] N=1,613 | -0.014 [0.032] N=1,488 | -0.006 [0.035] N=1,328 | 0.005 [0.041] N=1,128 | -0.007 [0.042] N=917 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.194 P=302 | **0.074* [0.041] N=1,613** | **0.077* [0.044] N=1,488** | **0.082* [0.047] N=1,328** | 0.036 [0.052] N=1,128 | 0.045 [0.056] N=917 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.082 P=131 | 0.031 [0.028] N=1,613 | 0.039 [0.030] N=1,488 | 0.046 [0.033] N=1,328 | 0.017 [0.037] N=1,128 | 0.017 [0.044] N=917 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.242 P=375 | -0.012 [0.039] N=1,583 | -0.011 [0.040] N=1,462 | 0.018 [0.044] N=1,306 | 0.004 [0.047] N=1,107 | 0.026 [0.050] N=902 |
| BMI-for-age z score below -2 | 0.114 P=318 | 0.021 [0.028] N=2,731 | 0.023 [0.029] N=2,531 | 0.028 [0.031] N=2,260 | 0.050 [0.032] N=1,943 | **0.058* [0.034] N=1,604** |
| WHO 2007 BMI overweight above +1, 61+ months | 0.374 P=417 | 0.034 [0.050] N=1,111 | 0.023 [0.053] N=1,034 | 0.067 [0.058] N=929 | 0.046 [0.062] N=815 | 0.011 [0.070] N=681 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.213 P=238 | 0.011 [0.043] N=1,111 | -0.009 [0.045] N=1,034 | 0.002 [0.049] N=929 | -0.033 [0.050] N=815 | **-0.098* [0.057] N=681** |
| WHO 2007 BMI thinness below -2, 61+ months | 0.109 P=127 | 0.033 [0.042] N=1,111 | 0.041 [0.043] N=1,034 | 0.022 [0.047] N=929 | 0.034 [0.050] N=815 | 0.034 [0.055] N=681 |

*Specification 3 Monthly fixed effects female and household income groups 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.083 [0.199] N=2,398 | -0.137 [0.210] N=2,185 | -0.260 [0.241] N=1,910 | -0.157 [0.272] N=1,578 | -0.352 [0.358] N=1,224 |
| WHO weight-for-age z score | -0.087 [0.107] N=2,551 | -0.027 [0.124] N=2,318 | 0.039 [0.129] N=2,024 | -0.147 [0.142] N=1,669 | -0.178 [0.197] N=1,294 |
| WHO weight-for-length/height z score, 0-60 months | 0.123 [0.261] N=1,371 | 0.176 [0.281] N=1,246 | 0.081 [0.324] N=1,086 | -0.386 [0.381] N=886 | -0.297 [0.463] N=675 |
| WHO BMI-for-age z score | 0.239 [0.190] N=2,301 | 0.287 [0.201] N=2,101 | 0.371 [0.229] N=1,830 | 0.037 [0.276] N=1,513 | 0.236 [0.355] N=1,174 |
| Stunted: HAZ below -2 | 0.032 [0.039] N=2,398 | 0.051 [0.041] N=2,185 | **0.091* [0.049] N=1,910** | 0.065 [0.055] N=1,578 | **0.148** [0.071] N=1,224** |
| Underweight: WAZ below -2 | 0.017 [0.019] N=2,551 | 0.016 [0.020] N=2,318 | 0.006 [0.023] N=2,024 | 0.015 [0.027] N=1,669 | 0.013 [0.036] N=1,294 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.005 [0.040] N=1,371 | 0.002 [0.043] N=1,246 | 0.021 [0.049] N=1,086 | 0.048 [0.064] N=886 | 0.037 [0.070] N=675 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.069 [0.055] N=1,371 | 0.074 [0.060] N=1,246 | 0.084 [0.067] N=1,086 | -0.022 [0.082] N=886 | -0.011 [0.104] N=675 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.048 [0.032] N=1,371 | **0.063* [0.034] N=1,246** | **0.085** [0.040] N=1,086** | 0.024 [0.048] N=886 | 0.035 [0.065] N=675 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.044 [0.055] N=1,344 | 0.046 [0.058] N=1,223 | **0.117* [0.065] N=1,067** | 0.100 [0.078] N=868 | **0.224** [0.094] N=663** |
| BMI-for-age z score below -2 | -0.007 [0.033] N=2,301 | -0.011 [0.034] N=2,101 | -0.008 [0.038] N=1,830 | 0.016 [0.043] N=1,513 | 0.019 [0.051] N=1,174 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.102 [0.073] N=923 | 0.092 [0.080] N=846 | **0.198** [0.092] N=741** | **0.197* [0.107] N=627** | 0.202 [0.137] N=493 |
| WHO 2007 BMI obesity above +2, 61+ months | **0.112* [0.063] N=923** | 0.088 [0.069] N=846 | **0.144* [0.079] N=741** | 0.122 [0.089] N=627 | 0.030 [0.114] N=493 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.012 [0.046] N=923 | 0.022 [0.047] N=846 | -0.018 [0.056] N=741 | -0.010 [0.060] N=627 | -0.032 [0.077] N=493 |

*Specification 4 Monthly fixed effects female and per-person income groups  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.055 | 0.041 [0.136] N=2,853 | 0.030 [0.139] N=2,640 | -0.008 [0.153] N=2,365 | 0.046 [0.156] N=2,033 | 0.044 [0.172] N=1,679 |
| WHO weight-for-age z score | 0.360 | 0.017 [0.084] N=3,026 | 0.050 [0.094] N=2,793 | 0.098 [0.098] N=2,499 | 0.011 [0.104] N=2,144 | 0.029 [0.119] N=1,769 |
| WHO weight-for-length/height z score, 0-60 months | 0.336 | 0.197 [0.183] N=1,613 | 0.217 [0.189] N=1,488 | 0.181 [0.202] N=1,328 | -0.012 [0.221] N=1,128 | 0.103 [0.233] N=917 |
| WHO BMI-for-age z score | 0.356 | 0.067 [0.150] N=2,731 | 0.072 [0.156] N=2,531 | 0.089 [0.167] N=2,260 | -0.122 [0.182] N=1,943 | -0.085 [0.195] N=1,604 |
| Stunted: HAZ below -2 | 0.180 P=494 | -0.010 [0.028] N=2,853 | -0.005 [0.029] N=2,640 | 0.008 [0.033] N=2,365 | -0.011 [0.034] N=2,033 | 0.002 [0.038] N=1,679 |
| Underweight: WAZ below -2 | 0.046 P=135 | -0.004 [0.014] N=3,026 | -0.007 [0.015] N=2,793 | -0.015 [0.016] N=2,499 | -0.014 [0.018] N=2,144 | -0.023 [0.020] N=1,769 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=179 | -0.018 [0.030] N=1,613 | -0.019 [0.032] N=1,488 | -0.010 [0.035] N=1,328 | 0.003 [0.040] N=1,128 | -0.013 [0.042] N=917 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.194 P=302 | **0.075* [0.041] N=1,613** | **0.077* [0.043] N=1,488** | **0.081* [0.046] N=1,328** | 0.037 [0.051] N=1,128 | 0.045 [0.055] N=917 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.082 P=131 | 0.034 [0.029] N=1,613 | 0.040 [0.030] N=1,488 | 0.048 [0.034] N=1,328 | 0.021 [0.038] N=1,128 | 0.025 [0.045] N=917 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.242 P=375 | -0.014 [0.039] N=1,583 | -0.016 [0.040] N=1,462 | 0.016 [0.044] N=1,306 | 0.009 [0.047] N=1,107 | 0.030 [0.051] N=902 |
| BMI-for-age z score below -2 | 0.114 P=318 | 0.019 [0.027] N=2,731 | 0.020 [0.028] N=2,531 | 0.024 [0.030] N=2,260 | 0.047 [0.032] N=1,943 | 0.053 [0.033] N=1,604 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.374 P=417 | 0.036 [0.050] N=1,111 | 0.026 [0.053] N=1,034 | 0.068 [0.058] N=929 | 0.046 [0.064] N=815 | 0.017 [0.071] N=681 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.213 P=238 | 0.013 [0.043] N=1,111 | -0.007 [0.045] N=1,034 | 0.003 [0.049] N=929 | -0.030 [0.051] N=815 | -0.093 [0.057] N=681 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.109 P=127 | 0.033 [0.042] N=1,111 | 0.040 [0.043] N=1,034 | 0.021 [0.047] N=929 | 0.033 [0.050] N=815 | 0.034 [0.053] N=681 |

*Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.071 [0.198] N=2,398 | -0.125 [0.210] N=2,185 | -0.246 [0.241] N=1,910 | -0.133 [0.273] N=1,578 | -0.332 [0.356] N=1,224 |
| WHO weight-for-age z score | -0.070 [0.108] N=2,551 | -0.020 [0.125] N=2,318 | 0.047 [0.132] N=2,024 | -0.139 [0.145] N=1,669 | -0.162 [0.192] N=1,294 |
| WHO weight-for-length/height z score, 0-60 months | 0.156 [0.262] N=1,371 | 0.206 [0.280] N=1,246 | 0.100 [0.326] N=1,086 | -0.367 [0.380] N=886 | -0.261 [0.462] N=675 |
| WHO BMI-for-age z score | 0.241 [0.191] N=2,301 | 0.288 [0.202] N=2,101 | 0.366 [0.232] N=1,830 | 0.018 [0.277] N=1,513 | 0.209 [0.352] N=1,174 |
| Stunted: HAZ below -2 | 0.032 [0.039] N=2,398 | 0.051 [0.040] N=2,185 | **0.091* [0.049] N=1,910** | 0.065 [0.056] N=1,578 | **0.151** [0.071] N=1,224** |
| Underweight: WAZ below -2 | 0.016 [0.019] N=2,551 | 0.015 [0.020] N=2,318 | 0.007 [0.024] N=2,024 | 0.016 [0.028] N=1,669 | 0.016 [0.037] N=1,294 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.002 [0.040] N=1,371 | -0.003 [0.043] N=1,246 | 0.019 [0.048] N=1,086 | 0.046 [0.063] N=886 | 0.033 [0.068] N=675 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.074 [0.055] N=1,371 | 0.077 [0.060] N=1,246 | 0.086 [0.068] N=1,086 | -0.020 [0.083] N=886 | -0.003 [0.105] N=675 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.052 [0.032] N=1,371 | **0.066* [0.034] N=1,246** | **0.088** [0.041] N=1,086** | 0.028 [0.048] N=886 | 0.042 [0.063] N=675 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.046 [0.054] N=1,344 | 0.042 [0.057] N=1,223 | **0.119* [0.064] N=1,067** | 0.107 [0.077] N=868 | **0.226** [0.094] N=663** |
| BMI-for-age z score below -2 | -0.006 [0.033] N=2,301 | -0.011 [0.035] N=2,101 | -0.008 [0.039] N=1,830 | 0.020 [0.043] N=1,513 | 0.027 [0.051] N=1,174 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.100 [0.073] N=923 | 0.092 [0.080] N=846 | **0.199** [0.093] N=741** | **0.198* [0.108] N=627** | 0.214 [0.136] N=493 |
| WHO 2007 BMI obesity above +2, 61+ months | **0.109* [0.063] N=923** | 0.087 [0.068] N=846 | **0.136* [0.078] N=741** | 0.112 [0.090] N=627 | 0.026 [0.116] N=493 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.013 [0.046] N=923 | 0.023 [0.048] N=846 | -0.021 [0.056] N=741 | -0.014 [0.062] N=627 | -0.018 [0.073] N=493 |

*Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.064 | 0.043 [0.148] N=2,664 | -0.009 [0.154] N=2,471 | -0.060 [0.165] N=2,207 | -0.031 [0.168] N=1,904 | -0.020 [0.193] N=1,568 |
| WHO weight-for-age z score | 0.353 | 0.054 [0.087] N=2,822 | 0.086 [0.094] N=2,611 | 0.142 [0.099] N=2,330 | 0.050 [0.107] N=2,006 | 0.081 [0.124] N=1,653 |
| WHO weight-for-length/height z score, 0-60 months | 0.352 | **0.315* [0.187] N=1,491** | **0.361* [0.192] N=1,379** | 0.318 [0.204] N=1,227 | 0.116 [0.230] N=1,050 | 0.255 [0.245] N=850 |
| WHO BMI-for-age z score | 0.362 | 0.121 [0.146] N=2,551 | 0.124 [0.149] N=2,370 | 0.158 [0.163] N=2,112 | -0.046 [0.178] N=1,823 | -0.023 [0.195] N=1,501 |
| Stunted: HAZ below -2 | 0.182 P=464 | -0.008 [0.029] N=2,664 | 0.002 [0.029] N=2,471 | 0.017 [0.033] N=2,207 | 0.001 [0.034] N=1,904 | 0.014 [0.039] N=1,568 |
| Underweight: WAZ below -2 | 0.047 P=130 | -0.009 [0.015] N=2,822 | -0.013 [0.015] N=2,611 | -0.020 [0.017] N=2,330 | -0.017 [0.020] N=2,006 | -0.024 [0.023] N=1,653 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=165 | -0.030 [0.032] N=1,491 | -0.034 [0.034] N=1,379 | -0.020 [0.036] N=1,227 | -0.007 [0.043] N=1,050 | -0.020 [0.044] N=850 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.197 P=282 | **0.085* [0.043] N=1,491** | **0.091** [0.045] N=1,379** | **0.094* [0.050] N=1,227** | 0.043 [0.055] N=1,050 | 0.058 [0.062] N=850 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.085 P=125 | 0.036 [0.029] N=1,491 | 0.048 [0.031] N=1,379 | **0.057* [0.034] N=1,227** | 0.027 [0.038] N=1,050 | 0.033 [0.046] N=850 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.243 P=348 | -0.022 [0.046] N=1,463 | -0.020 [0.048] N=1,355 | 0.010 [0.053] N=1,206 | -0.006 [0.056] N=1,030 | 0.020 [0.060] N=835 |
| BMI-for-age z score below -2 | 0.112 P=293 | 0.014 [0.026] N=2,551 | 0.015 [0.026] N=2,370 | 0.018 [0.028] N=2,112 | 0.037 [0.030] N=1,823 | 0.039 [0.031] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.370 P=392 | 0.042 [0.049] N=1,057 | 0.033 [0.051] N=987 | 0.082 [0.056] N=886 | 0.059 [0.060] N=777 | 0.021 [0.068] N=649 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.208 P=222 | 0.011 [0.043] N=1,057 | -0.006 [0.045] N=987 | 0.010 [0.049] N=886 | -0.027 [0.049] N=777 | **-0.106* [0.056] N=649** |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=122 | 0.043 [0.042] N=1,057 | 0.054 [0.042] N=987 | 0.034 [0.046] N=886 | 0.042 [0.049] N=777 | 0.044 [0.055] N=649 |

*Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.021 [0.218] N=2,240 | -0.121 [0.233] N=2,047 | -0.253 [0.264] N=1,783 | -0.188 [0.300] N=1,480 | -0.361 [0.428] N=1,144 |
| WHO weight-for-age z score | -0.059 [0.108] N=2,379 | -0.014 [0.119] N=2,168 | 0.064 [0.133] N=1,887 | -0.128 [0.150] N=1,563 | -0.135 [0.194] N=1,210 |
| WHO weight-for-length/height z score, 0-60 months | 0.198 [0.267] N=1,267 | 0.258 [0.282] N=1,155 | 0.164 [0.315] N=1,003 | -0.281 [0.386] N=826 | -0.112 [0.480] N=626 |
| WHO BMI-for-age z score | 0.264 [0.192] N=2,150 | 0.302 [0.197] N=1,969 | **0.416* [0.225] N=1,711** | 0.123 [0.273] N=1,422 | 0.306 [0.357] N=1,100 |
| Stunted: HAZ below -2 | 0.029 [0.041] N=2,240 | 0.053 [0.043] N=2,047 | **0.096* [0.051] N=1,783** | 0.076 [0.059] N=1,480 | **0.150* [0.078] N=1,144** |
| Underweight: WAZ below -2 | 0.013 [0.020] N=2,379 | 0.011 [0.021] N=2,168 | 0.004 [0.025] N=1,887 | 0.017 [0.029] N=1,563 | 0.018 [0.038] N=1,210 |
| Wasted: WLZ/WHZ below -2, 0-60 months | -0.009 [0.039] N=1,267 | -0.014 [0.041] N=1,155 | 0.013 [0.046] N=1,003 | 0.031 [0.062] N=826 | 0.003 [0.063] N=626 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.078 [0.056] N=1,267 | 0.087 [0.061] N=1,155 | 0.107 [0.068] N=1,003 | -0.007 [0.084] N=826 | 0.008 [0.112] N=626 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.049 [0.034] N=1,267 | **0.070* [0.036] N=1,155** | **0.093** [0.041] N=1,003** | 0.033 [0.051] N=826 | 0.055 [0.073] N=626 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.037 [0.059] N=1,242 | 0.045 [0.061] N=1,134 | **0.121* [0.070] N=985** | 0.091 [0.086] N=809 | **0.200** [0.095] N=614** |
| BMI-for-age z score below -2 | 0.003 [0.032] N=2,150 | 0.000 [0.032] N=1,969 | 0.005 [0.037] N=1,711 | 0.024 [0.041] N=1,422 | 0.017 [0.047] N=1,100 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.102 [0.074] N=878 | 0.097 [0.080] N=808 | **0.209** [0.095] N=707** | **0.207* [0.107] N=598** | 0.188 [0.142] N=470 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.103 [0.064] N=878 | 0.086 [0.069] N=808 | **0.147* [0.079] N=707** | 0.122 [0.088] N=598 | -0.008 [0.114] N=470 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.025 [0.047] N=878 | 0.038 [0.048] N=808 | -0.005 [0.056] N=707 | -0.009 [0.060] N=598 | -0.028 [0.076] N=470 |

*Specification 6 Monthly fixed effects female and any aid (0/1)  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.064 | 0.051 [0.145] N=2,664 | -0.003 [0.150] N=2,471 | -0.062 [0.161] N=2,207 | -0.030 [0.162] N=1,904 | -0.006 [0.182] N=1,568 |
| WHO weight-for-age z score | 0.353 | 0.063 [0.087] N=2,822 | 0.093 [0.094] N=2,611 | 0.151 [0.098] N=2,330 | 0.061 [0.106] N=2,006 | 0.099 [0.122] N=1,653 |
| WHO weight-for-length/height z score, 0-60 months | 0.352 | **0.321* [0.186] N=1,491** | **0.366* [0.192] N=1,379** | 0.321 [0.204] N=1,227 | 0.119 [0.230] N=1,050 | 0.246 [0.247] N=850 |
| WHO BMI-for-age z score | 0.362 | 0.122 [0.146] N=2,551 | 0.125 [0.148] N=2,370 | 0.161 [0.162] N=2,112 | -0.044 [0.177] N=1,823 | -0.018 [0.194] N=1,501 |
| Stunted: HAZ below -2 | 0.182 P=464 | -0.010 [0.029] N=2,664 | 0.001 [0.029] N=2,471 | 0.017 [0.033] N=2,207 | -0.000 [0.034] N=1,904 | 0.010 [0.038] N=1,568 |
| Underweight: WAZ below -2 | 0.047 P=130 | -0.011 [0.015] N=2,822 | -0.015 [0.016] N=2,611 | -0.023 [0.017] N=2,330 | -0.020 [0.020] N=2,006 | -0.029 [0.023] N=1,653 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.109 P=165 | -0.031 [0.031] N=1,491 | -0.035 [0.033] N=1,379 | -0.020 [0.036] N=1,227 | -0.007 [0.043] N=1,050 | -0.022 [0.045] N=850 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.197 P=282 | **0.086** [0.043] N=1,491** | **0.091** [0.045] N=1,379** | **0.096* [0.050] N=1,227** | 0.046 [0.056] N=1,050 | 0.061 [0.062] N=850 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.085 P=125 | 0.037 [0.029] N=1,491 | 0.048 [0.031] N=1,379 | **0.058* [0.034] N=1,227** | 0.029 [0.038] N=1,050 | 0.035 [0.046] N=850 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.243 P=348 | -0.025 [0.045] N=1,463 | -0.022 [0.047] N=1,355 | 0.012 [0.052] N=1,206 | -0.004 [0.055] N=1,030 | 0.018 [0.059] N=835 |
| BMI-for-age z score below -2 | 0.112 P=293 | 0.014 [0.026] N=2,551 | 0.014 [0.026] N=2,370 | 0.019 [0.028] N=2,112 | 0.037 [0.030] N=1,823 | 0.039 [0.031] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.370 P=392 | 0.042 [0.048] N=1,057 | 0.032 [0.051] N=987 | 0.080 [0.055] N=886 | 0.054 [0.059] N=777 | 0.013 [0.067] N=649 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.208 P=222 | 0.009 [0.043] N=1,057 | -0.008 [0.044] N=987 | 0.009 [0.048] N=886 | -0.028 [0.049] N=777 | **-0.106* [0.056] N=649** |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=122 | 0.041 [0.042] N=1,057 | 0.052 [0.042] N=987 | 0.032 [0.045] N=886 | 0.040 [0.049] N=777 | 0.043 [0.054] N=649 |

*Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.009 [0.216] N=2,240 | -0.112 [0.231] N=2,047 | -0.257 [0.263] N=1,783 | -0.198 [0.297] N=1,480 | -0.335 [0.412] N=1,144 |
| WHO weight-for-age z score | -0.047 [0.108] N=2,379 | -0.006 [0.121] N=2,168 | 0.077 [0.132] N=1,887 | -0.106 [0.149] N=1,563 | -0.089 [0.190] N=1,210 |
| WHO weight-for-length/height z score, 0-60 months | 0.230 [0.266] N=1,267 | 0.295 [0.280] N=1,155 | 0.178 [0.313] N=1,003 | -0.253 [0.387] N=826 | -0.094 [0.480] N=626 |
| WHO BMI-for-age z score | 0.270 [0.192] N=2,150 | 0.308 [0.197] N=1,969 | **0.419* [0.224] N=1,711** | 0.129 [0.273] N=1,422 | 0.325 [0.360] N=1,100 |
| Stunted: HAZ below -2 | 0.027 [0.041] N=2,240 | 0.052 [0.043] N=2,047 | **0.094* [0.051] N=1,783** | 0.073 [0.058] N=1,480 | **0.140* [0.077] N=1,144** |
| Underweight: WAZ below -2 | 0.011 [0.020] N=2,379 | 0.010 [0.022] N=2,168 | 0.001 [0.025] N=1,887 | 0.012 [0.030] N=1,563 | 0.006 [0.039] N=1,210 |
| Wasted: WLZ/WHZ below -2, 0-60 months | -0.015 [0.038] N=1,267 | -0.020 [0.041] N=1,155 | 0.011 [0.046] N=1,003 | 0.030 [0.062] N=826 | -0.006 [0.061] N=626 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.083 [0.055] N=1,267 | 0.093 [0.060] N=1,155 | 0.110 [0.067] N=1,003 | 0.003 [0.085] N=826 | 0.024 [0.113] N=626 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.048 [0.033] N=1,267 | **0.068* [0.036] N=1,155** | **0.094** [0.041] N=1,003** | 0.038 [0.051] N=826 | 0.064 [0.074] N=626 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.031 [0.058] N=1,242 | 0.038 [0.061] N=1,134 | **0.118* [0.070] N=985** | 0.089 [0.084] N=809 | **0.177* [0.094] N=614** |
| BMI-for-age z score below -2 | 0.001 [0.032] N=2,150 | -0.002 [0.032] N=1,969 | 0.004 [0.037] N=1,711 | 0.025 [0.041] N=1,422 | 0.017 [0.047] N=1,100 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.104 [0.074] N=878 | 0.097 [0.080] N=808 | **0.207** [0.094] N=707** | **0.206* [0.104] N=598** | 0.183 [0.138] N=470 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.100 [0.063] N=878 | 0.081 [0.068] N=808 | **0.144* [0.079] N=707** | 0.118 [0.088] N=598 | -0.010 [0.114] N=470 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.023 [0.047] N=878 | 0.036 [0.048] N=808 | -0.005 [0.057] N=707 | -0.009 [0.061] N=598 | -0.030 [0.075] N=470 |

**Mother RDD secondary analysis**

*Specification 1 Monthly child-age fixed effects  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.029 | -0.178 [0.184] N=2,845 | -0.203 [0.199] N=2,583 | -0.248 [0.209] N=2,275 | **-0.390* [0.224] N=1,932** | -0.382 [0.249] N=1,569 |
| WHO weight-for-age z score | 0.374 | -0.085 [0.095] N=3,009 | -0.066 [0.099] N=2,744 | -0.080 [0.104] N=2,414 | -0.160 [0.112] N=2,052 | -0.143 [0.126] N=1,658 |
| WHO weight-for-length/height z score, 0-60 months | 0.374 | 0.049 [0.224] N=1,527 | 0.105 [0.234] N=1,391 | 0.092 [0.258] N=1,204 | 0.128 [0.278] N=1,030 | 0.035 [0.324] N=831 |
| WHO BMI-for-age z score | 0.378 | 0.113 [0.174] N=2,725 | 0.153 [0.182] N=2,476 | 0.253 [0.190] N=2,181 | 0.319 [0.209] N=1,850 | 0.289 [0.243] N=1,501 |
| Stunted: HAZ below -2 | 0.177 P=494 | 0.024 [0.033] N=2,845 | 0.025 [0.035] N=2,583 | 0.021 [0.038] N=2,275 | 0.026 [0.042] N=1,932 | 0.026 [0.047] N=1,569 |
| Underweight: WAZ below -2 | 0.040 P=122 | 0.009 [0.013] N=3,009 | 0.005 [0.014] N=2,744 | 0.008 [0.015] N=2,414 | 0.014 [0.015] N=2,052 | 0.022 [0.016] N=1,658 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.107 P=162 | 0.015 [0.037] N=1,527 | 0.001 [0.040] N=1,391 | 0.005 [0.043] N=1,204 | -0.014 [0.044] N=1,030 | -0.001 [0.053] N=831 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.195 P=299 | 0.032 [0.038] N=1,527 | 0.034 [0.039] N=1,391 | 0.027 [0.040] N=1,204 | 0.026 [0.044] N=1,030 | 0.022 [0.049] N=831 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.090 P=141 | 0.035 [0.029] N=1,527 | 0.032 [0.030] N=1,391 | 0.022 [0.032] N=1,204 | 0.021 [0.037] N=1,030 | -0.009 [0.039] N=831 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.246 P=363 | 0.054 [0.045] N=1,497 | 0.045 [0.049] N=1,361 | 0.037 [0.050] N=1,177 | -0.001 [0.053] N=1,007 | 0.026 [0.060] N=811 |
| BMI-for-age z score below -2 | 0.114 P=311 | 0.030 [0.028] N=2,725 | 0.021 [0.029] N=2,476 | 0.009 [0.031] N=2,181 | -0.008 [0.032] N=1,850 | 0.003 [0.038] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.388 P=460 | -0.001 [0.058] N=1,200 | -0.017 [0.062] N=1,087 | 0.002 [0.064] N=979 | 0.012 [0.073] N=821 | -0.030 [0.082] N=664 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.214 P=252 | 0.019 [0.049] N=1,200 | -0.011 [0.052] N=1,087 | 0.003 [0.055] N=979 | 0.030 [0.061] N=821 | 0.021 [0.072] N=664 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=137 | 0.025 [0.037] N=1,200 | 0.025 [0.039] N=1,087 | 0.002 [0.040] N=979 | -0.024 [0.045] N=821 | -0.008 [0.050] N=664 |

*Specification 1 Monthly child-age fixed effects 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.050 [0.226] N=2,439 | 0.038 [0.253] N=2,177 | -0.043 [0.273] N=1,869 | -0.284 [0.294] N=1,526 | -0.122 [0.371] N=1,163 |
| WHO weight-for-age z score | -0.154 [0.140] N=2,584 | -0.126 [0.154] N=2,319 | -0.186 [0.171] N=1,989 | **-0.395** [0.192] N=1,627** | **-0.454* [0.233] N=1,233** |
| WHO weight-for-length/height z score, 0-60 months | -0.084 [0.276] N=1,334 | -0.022 [0.291] N=1,198 | -0.146 [0.347] N=1,011 | -0.205 [0.367] N=837 | **-0.768* [0.457] N=638** |
| WHO BMI-for-age z score | **-0.393** [0.171] N=2,340** | **-0.406** [0.180] N=2,091** | **-0.352* [0.203] N=1,796** | **-0.448* [0.231] N=1,465** | **-1.027*** [0.300] N=1,116** |
| Stunted: HAZ below -2 | 0.024 [0.040] N=2,439 | 0.028 [0.045] N=2,177 | 0.028 [0.048] N=1,869 | 0.037 [0.058] N=1,526 | 0.026 [0.076] N=1,163 |
| Underweight: WAZ below -2 | -0.003 [0.020] N=2,584 | -0.013 [0.022] N=2,319 | -0.011 [0.026] N=1,989 | -0.010 [0.026] N=1,627 | -0.004 [0.030] N=1,233 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.063 [0.041] N=1,334 | 0.045 [0.045] N=1,198 | 0.070 [0.051] N=1,011 | 0.047 [0.053] N=837 | **0.129** [0.060] N=638** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.036 [0.059] N=1,334 | 0.034 [0.063] N=1,198 | 0.011 [0.068] N=1,011 | -0.014 [0.079] N=837 | -0.074 [0.093] N=638 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.053 [0.043] N=1,334 | 0.047 [0.045] N=1,198 | 0.027 [0.052] N=1,011 | 0.025 [0.062] N=837 | -0.080 [0.070] N=638 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.095 [0.062] N=1,309 | 0.086 [0.069] N=1,173 | 0.098 [0.075] N=989 | 0.032 [0.085] N=819 | 0.110 [0.097] N=623 |
| BMI-for-age z score below -2 | **0.086*** [0.025] N=2,340** | **0.078*** [0.028] N=2,091** | **0.067** [0.033] N=1,796** | **0.059* [0.030] N=1,465** | **0.139*** [0.043] N=1,116** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.058 [0.080] N=1,010 | -0.088 [0.090] N=897 | -0.065 [0.097] N=789 | -0.067 [0.120] N=631 | **-0.253* [0.137] N=474** |
| WHO 2007 BMI obesity above +2, 61+ months | -0.058 [0.062] N=1,010 | **-0.122* [0.067] N=897** | -0.118 [0.072] N=789 | -0.115 [0.082] N=631 | **-0.219* [0.111] N=474** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.095** [0.044] N=1,010** | **0.106** [0.048] N=897** | 0.066 [0.052] N=789 | 0.035 [0.061] N=631 | 0.112 [0.075] N=474 |

*Specification 2 Monthly fixed effects and female  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.029 | -0.176 [0.185] N=2,845 | -0.201 [0.200] N=2,583 | -0.247 [0.210] N=2,275 | **-0.387* [0.225] N=1,932** | -0.372 [0.249] N=1,569 |
| WHO weight-for-age z score | 0.374 | -0.081 [0.095] N=3,009 | -0.062 [0.099] N=2,744 | -0.078 [0.105] N=2,414 | -0.158 [0.113] N=2,052 | -0.137 [0.126] N=1,658 |
| WHO weight-for-length/height z score, 0-60 months | 0.374 | 0.059 [0.224] N=1,527 | 0.113 [0.234] N=1,391 | 0.096 [0.258] N=1,204 | 0.131 [0.277] N=1,030 | 0.029 [0.321] N=831 |
| WHO BMI-for-age z score | 0.378 | 0.112 [0.174] N=2,725 | 0.152 [0.182] N=2,476 | 0.254 [0.190] N=2,181 | 0.318 [0.209] N=1,850 | 0.284 [0.243] N=1,501 |
| Stunted: HAZ below -2 | 0.177 P=494 | 0.024 [0.033] N=2,845 | 0.025 [0.035] N=2,583 | 0.021 [0.038] N=2,275 | 0.027 [0.042] N=1,932 | 0.026 [0.047] N=1,569 |
| Underweight: WAZ below -2 | 0.040 P=122 | 0.008 [0.013] N=3,009 | 0.005 [0.014] N=2,744 | 0.008 [0.015] N=2,414 | 0.014 [0.015] N=2,052 | 0.022 [0.015] N=1,658 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.107 P=162 | 0.013 [0.037] N=1,527 | -0.001 [0.040] N=1,391 | 0.003 [0.043] N=1,204 | -0.015 [0.044] N=1,030 | 0.000 [0.052] N=831 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.195 P=299 | 0.032 [0.038] N=1,527 | 0.034 [0.039] N=1,391 | 0.026 [0.040] N=1,204 | 0.026 [0.044] N=1,030 | 0.023 [0.049] N=831 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.090 P=141 | 0.035 [0.029] N=1,527 | 0.032 [0.030] N=1,391 | 0.021 [0.032] N=1,204 | 0.021 [0.037] N=1,030 | -0.009 [0.039] N=831 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.246 P=363 | 0.053 [0.045] N=1,497 | 0.044 [0.049] N=1,361 | 0.036 [0.049] N=1,177 | -0.002 [0.053] N=1,007 | 0.027 [0.060] N=811 |
| BMI-for-age z score below -2 | 0.114 P=311 | 0.031 [0.028] N=2,725 | 0.022 [0.030] N=2,476 | 0.009 [0.031] N=2,181 | -0.007 [0.032] N=1,850 | 0.006 [0.039] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.388 P=460 | 0.002 [0.059] N=1,200 | -0.014 [0.062] N=1,087 | 0.004 [0.064] N=979 | 0.013 [0.073] N=821 | -0.025 [0.081] N=664 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.214 P=252 | 0.026 [0.049] N=1,200 | -0.005 [0.053] N=1,087 | 0.008 [0.056] N=979 | 0.033 [0.061] N=821 | 0.026 [0.072] N=664 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=137 | 0.027 [0.037] N=1,200 | 0.027 [0.039] N=1,087 | 0.002 [0.039] N=979 | -0.023 [0.045] N=821 | -0.008 [0.049] N=664 |

*Specification 2 Monthly fixed effects and female 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.051 [0.226] N=2,439 | 0.039 [0.253] N=2,177 | -0.043 [0.273] N=1,869 | -0.283 [0.295] N=1,526 | -0.103 [0.373] N=1,163 |
| WHO weight-for-age z score | -0.150 [0.141] N=2,584 | -0.121 [0.154] N=2,319 | -0.184 [0.173] N=1,989 | **-0.392** [0.194] N=1,627** | **-0.442* [0.235] N=1,233** |
| WHO weight-for-length/height z score, 0-60 months | -0.065 [0.279] N=1,334 | -0.006 [0.294] N=1,198 | -0.139 [0.347] N=1,011 | -0.192 [0.369] N=837 | -0.754 [0.468] N=638 |
| WHO BMI-for-age z score | **-0.393** [0.171] N=2,340** | **-0.407** [0.180] N=2,091** | **-0.352* [0.203] N=1,796** | **-0.449* [0.231] N=1,465** | **-1.034*** [0.299] N=1,116** |
| Stunted: HAZ below -2 | 0.024 [0.040] N=2,439 | 0.028 [0.045] N=2,177 | 0.028 [0.048] N=1,869 | 0.037 [0.058] N=1,526 | 0.025 [0.075] N=1,163 |
| Underweight: WAZ below -2 | -0.004 [0.020] N=2,584 | -0.013 [0.022] N=2,319 | -0.011 [0.026] N=1,989 | -0.011 [0.026] N=1,627 | -0.005 [0.029] N=1,233 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.059 [0.041] N=1,334 | 0.041 [0.045] N=1,198 | 0.065 [0.052] N=1,011 | 0.040 [0.054] N=837 | **0.125** [0.061] N=638** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.036 [0.059] N=1,334 | 0.033 [0.063] N=1,198 | 0.008 [0.068] N=1,011 | -0.016 [0.079] N=837 | -0.074 [0.093] N=638 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.053 [0.043] N=1,334 | 0.045 [0.045] N=1,198 | 0.024 [0.052] N=1,011 | 0.023 [0.062] N=837 | -0.080 [0.071] N=638 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.092 [0.062] N=1,309 | 0.083 [0.070] N=1,173 | 0.092 [0.075] N=989 | 0.026 [0.086] N=819 | 0.107 [0.099] N=623 |
| BMI-for-age z score below -2 | **0.087*** [0.025] N=2,340** | **0.079*** [0.028] N=2,091** | **0.068** [0.033] N=1,796** | **0.060** [0.030] N=1,465** | **0.144*** [0.043] N=1,116** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.054 [0.081] N=1,010 | -0.085 [0.091] N=897 | -0.063 [0.098] N=789 | -0.068 [0.121] N=631 | **-0.244* [0.140] N=474** |
| WHO 2007 BMI obesity above +2, 61+ months | -0.049 [0.064] N=1,010 | -0.114 [0.069] N=897 | -0.111 [0.074] N=789 | -0.110 [0.085] N=631 | **-0.211* [0.113] N=474** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.098** [0.044] N=1,010** | **0.109** [0.048] N=897** | 0.066 [0.052] N=789 | 0.036 [0.062] N=631 | 0.114 [0.076] N=474 |

*Specification 3 Monthly fixed effects female and household income groups  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.029 | -0.173 [0.183] N=2,845 | -0.190 [0.197] N=2,583 | -0.229 [0.207] N=2,275 | **-0.374* [0.220] N=1,932** | -0.352 [0.244] N=1,569 |
| WHO weight-for-age z score | 0.374 | -0.086 [0.094] N=3,009 | -0.064 [0.099] N=2,744 | -0.071 [0.104] N=2,414 | -0.158 [0.112] N=2,052 | -0.132 [0.126] N=1,658 |
| WHO weight-for-length/height z score, 0-60 months | 0.374 | 0.056 [0.224] N=1,527 | 0.112 [0.234] N=1,391 | 0.110 [0.260] N=1,204 | 0.146 [0.279] N=1,030 | 0.055 [0.328] N=831 |
| WHO BMI-for-age z score | 0.378 | 0.106 [0.174] N=2,725 | 0.145 [0.182] N=2,476 | 0.252 [0.190] N=2,181 | 0.312 [0.210] N=1,850 | 0.278 [0.245] N=1,501 |
| Stunted: HAZ below -2 | 0.177 P=494 | 0.023 [0.033] N=2,845 | 0.023 [0.035] N=2,583 | 0.019 [0.038] N=2,275 | 0.024 [0.042] N=1,932 | 0.021 [0.047] N=1,569 |
| Underweight: WAZ below -2 | 0.040 P=122 | 0.009 [0.013] N=3,009 | 0.005 [0.014] N=2,744 | 0.007 [0.015] N=2,414 | 0.013 [0.014] N=2,052 | 0.021 [0.015] N=1,658 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.107 P=162 | 0.017 [0.038] N=1,527 | 0.002 [0.040] N=1,391 | 0.005 [0.043] N=1,204 | -0.013 [0.045] N=1,030 | 0.002 [0.053] N=831 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.195 P=299 | 0.033 [0.038] N=1,527 | 0.035 [0.039] N=1,391 | 0.031 [0.041] N=1,204 | 0.030 [0.044] N=1,030 | 0.029 [0.049] N=831 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.090 P=141 | 0.037 [0.030] N=1,527 | 0.033 [0.030] N=1,391 | 0.024 [0.033] N=1,204 | 0.024 [0.037] N=1,030 | -0.005 [0.040] N=831 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.246 P=363 | 0.061 [0.044] N=1,497 | 0.049 [0.049] N=1,361 | 0.039 [0.049] N=1,177 | 0.001 [0.053] N=1,007 | 0.027 [0.059] N=811 |
| BMI-for-age z score below -2 | 0.114 P=311 | 0.031 [0.027] N=2,725 | 0.021 [0.029] N=2,476 | 0.008 [0.031] N=2,181 | -0.007 [0.032] N=1,850 | 0.005 [0.039] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.388 P=460 | -0.007 [0.059] N=1,200 | -0.022 [0.063] N=1,087 | -0.006 [0.065] N=979 | 0.008 [0.073] N=821 | -0.025 [0.081] N=664 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.214 P=252 | 0.017 [0.049] N=1,200 | -0.013 [0.053] N=1,087 | -0.003 [0.055] N=979 | 0.024 [0.061] N=821 | 0.025 [0.071] N=664 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=137 | 0.022 [0.036] N=1,200 | 0.024 [0.038] N=1,087 | 0.002 [0.039] N=979 | -0.022 [0.045] N=821 | -0.007 [0.050] N=664 |

*Specification 3 Monthly fixed effects female and household income groups 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.058 [0.227] N=2,439 | 0.055 [0.253] N=2,177 | -0.004 [0.274] N=1,869 | -0.249 [0.294] N=1,526 | -0.038 [0.378] N=1,163 |
| WHO weight-for-age z score | -0.157 [0.140] N=2,584 | -0.125 [0.154] N=2,319 | -0.173 [0.173] N=1,989 | **-0.395** [0.192] N=1,627** | **-0.431* [0.238] N=1,233** |
| WHO weight-for-length/height z score, 0-60 months | -0.080 [0.281] N=1,334 | -0.018 [0.296] N=1,198 | -0.133 [0.348] N=1,011 | -0.193 [0.367] N=837 | -0.756 [0.465] N=638 |
| WHO BMI-for-age z score | **-0.406** [0.171] N=2,340** | **-0.420** [0.182] N=2,091** | **-0.363* [0.205] N=1,796** | **-0.470** [0.234] N=1,465** | **-1.055*** [0.309] N=1,116** |
| Stunted: HAZ below -2 | 0.022 [0.041] N=2,439 | 0.024 [0.046] N=2,177 | 0.021 [0.049] N=1,869 | 0.030 [0.059] N=1,526 | 0.011 [0.078] N=1,163 |
| Underweight: WAZ below -2 | -0.004 [0.020] N=2,584 | -0.015 [0.022] N=2,319 | -0.014 [0.026] N=1,989 | -0.012 [0.026] N=1,627 | -0.007 [0.028] N=1,233 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.064 [0.043] N=1,334 | 0.043 [0.047] N=1,198 | 0.065 [0.053] N=1,011 | 0.041 [0.056] N=837 | **0.127** [0.061] N=638** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.036 [0.060] N=1,334 | 0.031 [0.064] N=1,198 | 0.008 [0.069] N=1,011 | -0.018 [0.079] N=837 | -0.072 [0.097] N=638 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.053 [0.044] N=1,334 | 0.044 [0.046] N=1,198 | 0.024 [0.053] N=1,011 | 0.022 [0.063] N=837 | -0.083 [0.073] N=638 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.100 [0.064] N=1,309 | 0.085 [0.072] N=1,173 | 0.087 [0.076] N=989 | 0.019 [0.088] N=819 | 0.095 [0.101] N=623 |
| BMI-for-age z score below -2 | **0.086*** [0.025] N=2,340** | **0.077*** [0.028] N=2,091** | **0.065** [0.033] N=1,796** | **0.059* [0.030] N=1,465** | **0.141*** [0.043] N=1,116** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.071 [0.081] N=1,010 | -0.098 [0.092] N=897 | -0.080 [0.099] N=789 | -0.083 [0.122] N=631 | **-0.255* [0.142] N=474** |
| WHO 2007 BMI obesity above +2, 61+ months | -0.062 [0.064] N=1,010 | **-0.124* [0.069] N=897** | **-0.125* [0.075] N=789** | -0.126 [0.085] N=631 | **-0.213* [0.113] N=474** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.092** [0.044] N=1,010** | **0.104** [0.047] N=897** | 0.064 [0.052] N=789 | 0.038 [0.062] N=631 | 0.115 [0.075] N=474 |

*Specification 4 Monthly fixed effects female and per-person income groups  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.029 | -0.177 [0.183] N=2,845 | -0.193 [0.197] N=2,583 | -0.241 [0.208] N=2,275 | **-0.377* [0.222] N=1,932** | -0.365 [0.246] N=1,569 |
| WHO weight-for-age z score | 0.374 | -0.086 [0.091] N=3,009 | -0.063 [0.095] N=2,744 | -0.077 [0.100] N=2,414 | -0.159 [0.107] N=2,052 | -0.150 [0.119] N=1,658 |
| WHO weight-for-length/height z score, 0-60 months | 0.374 | 0.063 [0.225] N=1,527 | 0.117 [0.234] N=1,391 | 0.107 [0.260] N=1,204 | 0.148 [0.277] N=1,030 | 0.030 [0.324] N=831 |
| WHO BMI-for-age z score | 0.378 | 0.110 [0.174] N=2,725 | 0.150 [0.182] N=2,476 | 0.253 [0.190] N=2,181 | 0.316 [0.209] N=1,850 | 0.270 [0.244] N=1,501 |
| Stunted: HAZ below -2 | 0.177 P=494 | 0.025 [0.032] N=2,845 | 0.025 [0.035] N=2,583 | 0.022 [0.038] N=2,275 | 0.027 [0.042] N=1,932 | 0.025 [0.047] N=1,569 |
| Underweight: WAZ below -2 | 0.040 P=122 | 0.009 [0.013] N=3,009 | 0.005 [0.014] N=2,744 | 0.008 [0.015] N=2,414 | 0.014 [0.014] N=2,052 | 0.022 [0.015] N=1,658 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.107 P=162 | 0.016 [0.037] N=1,527 | 0.002 [0.040] N=1,391 | 0.007 [0.043] N=1,204 | -0.014 [0.044] N=1,030 | -0.000 [0.052] N=831 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.195 P=299 | 0.036 [0.039] N=1,527 | 0.038 [0.040] N=1,391 | 0.035 [0.041] N=1,204 | 0.033 [0.044] N=1,030 | 0.024 [0.049] N=831 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.090 P=141 | 0.038 [0.029] N=1,527 | 0.035 [0.030] N=1,391 | 0.024 [0.032] N=1,204 | 0.024 [0.037] N=1,030 | -0.008 [0.040] N=831 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.246 P=363 | 0.064 [0.044] N=1,497 | 0.053 [0.048] N=1,361 | 0.043 [0.048] N=1,177 | 0.004 [0.051] N=1,007 | 0.027 [0.057] N=811 |
| BMI-for-age z score below -2 | 0.114 P=311 | 0.031 [0.028] N=2,725 | 0.021 [0.029] N=2,476 | 0.009 [0.031] N=2,181 | -0.008 [0.032] N=1,850 | 0.003 [0.038] N=1,501 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.388 P=460 | -0.003 [0.059] N=1,200 | -0.018 [0.064] N=1,087 | 0.001 [0.065] N=979 | 0.012 [0.073] N=821 | -0.025 [0.082] N=664 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.214 P=252 | 0.019 [0.049] N=1,200 | -0.010 [0.052] N=1,087 | 0.001 [0.055] N=979 | 0.024 [0.060] N=821 | 0.016 [0.070] N=664 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=137 | 0.023 [0.036] N=1,200 | 0.025 [0.038] N=1,087 | 0.000 [0.039] N=979 | -0.025 [0.044] N=821 | -0.005 [0.048] N=664 |

*Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.048 [0.225] N=2,439 | 0.047 [0.251] N=2,177 | -0.019 [0.271] N=1,869 | -0.252 [0.293] N=1,526 | -0.085 [0.371] N=1,163 |
| WHO weight-for-age z score | -0.146 [0.137] N=2,584 | -0.112 [0.150] N=2,319 | -0.162 [0.167] N=1,989 | **-0.366* [0.185] N=1,627** | **-0.425* [0.225] N=1,233** |
| WHO weight-for-length/height z score, 0-60 months | -0.061 [0.280] N=1,334 | -0.006 [0.293] N=1,198 | -0.137 [0.344] N=1,011 | -0.184 [0.362] N=837 | -0.753 [0.470] N=638 |
| WHO BMI-for-age z score | **-0.391** [0.171] N=2,340** | **-0.406** [0.180] N=2,091** | **-0.358* [0.203] N=1,796** | **-0.459* [0.232] N=1,465** | **-1.045*** [0.311] N=1,116** |
| Stunted: HAZ below -2 | 0.025 [0.040] N=2,439 | 0.027 [0.045] N=2,177 | 0.022 [0.048] N=1,869 | 0.029 [0.059] N=1,526 | 0.017 [0.077] N=1,163 |
| Underweight: WAZ below -2 | -0.003 [0.020] N=2,584 | -0.014 [0.022] N=2,319 | -0.013 [0.026] N=1,989 | -0.014 [0.026] N=1,627 | -0.008 [0.029] N=1,233 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.065 [0.041] N=1,334 | 0.046 [0.045] N=1,198 | 0.073 [0.051] N=1,011 | 0.045 [0.053] N=837 | **0.130** [0.059] N=638** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.042 [0.059] N=1,334 | 0.037 [0.063] N=1,198 | 0.015 [0.067] N=1,011 | -0.011 [0.078] N=837 | -0.070 [0.094] N=638 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.057 [0.043] N=1,334 | 0.048 [0.046] N=1,198 | 0.025 [0.052] N=1,011 | 0.022 [0.062] N=837 | -0.083 [0.072] N=638 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | **0.107* [0.061] N=1,309** | 0.092 [0.070] N=1,173 | 0.093 [0.074] N=989 | 0.022 [0.085] N=819 | 0.102 [0.096] N=623 |
| BMI-for-age z score below -2 | **0.089*** [0.025] N=2,340** | **0.080*** [0.028] N=2,091** | **0.069** [0.033] N=1,796** | **0.061** [0.030] N=1,465** | **0.146*** [0.043] N=1,116** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.069 [0.081] N=1,010 | -0.096 [0.092] N=897 | -0.079 [0.099] N=789 | -0.084 [0.120] N=631 | **-0.253* [0.143] N=474** |
| WHO 2007 BMI obesity above +2, 61+ months | -0.060 [0.064] N=1,010 | **-0.120* [0.069] N=897** | -0.120 [0.075] N=789 | -0.118 [0.083] N=631 | **-0.213* [0.112] N=474** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.092** [0.043] N=1,010** | **0.106** [0.047] N=897** | 0.063 [0.051] N=789 | 0.033 [0.060] N=631 | 0.113 [0.072] N=474 |

*Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.042 | -0.176 [0.189] N=2,667 | -0.184 [0.204] N=2,422 | -0.238 [0.214] N=2,137 | **-0.374* [0.225] N=1,810** | **-0.425* [0.251] N=1,472** |
| WHO weight-for-age z score | 0.374 | -0.055 [0.103] N=2,816 | -0.028 [0.108] N=2,568 | -0.049 [0.116] N=2,264 | -0.126 [0.126] N=1,921 | -0.151 [0.140] N=1,556 |
| WHO weight-for-length/height z score, 0-60 months | 0.379 | 0.112 [0.237] N=1,419 | 0.182 [0.247] N=1,293 | 0.110 [0.272] N=1,120 | 0.141 [0.289] N=958 | 0.028 [0.334] N=778 |
| WHO BMI-for-age z score | 0.389 | 0.099 [0.176] N=2,555 | 0.145 [0.184] N=2,322 | 0.215 [0.192] N=2,049 | 0.270 [0.208] N=1,735 | 0.219 [0.241] N=1,411 |
| Stunted: HAZ below -2 | 0.179 P=467 | 0.028 [0.032] N=2,667 | 0.030 [0.035] N=2,422 | 0.027 [0.037] N=2,137 | 0.033 [0.040] N=1,810 | 0.042 [0.044] N=1,472 |
| Underweight: WAZ below -2 | 0.041 P=117 | 0.008 [0.014] N=2,816 | 0.004 [0.015] N=2,568 | 0.003 [0.015] N=2,264 | 0.013 [0.016] N=1,921 | 0.026 [0.016] N=1,556 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.108 P=152 | 0.015 [0.038] N=1,419 | -0.002 [0.041] N=1,293 | 0.009 [0.043] N=1,120 | -0.006 [0.045] N=958 | 0.009 [0.052] N=778 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.196 P=280 | 0.031 [0.041] N=1,419 | 0.036 [0.042] N=1,293 | 0.026 [0.043] N=1,120 | 0.024 [0.046] N=958 | 0.021 [0.053] N=778 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.091 P=134 | 0.040 [0.030] N=1,419 | 0.038 [0.031] N=1,293 | 0.028 [0.033] N=1,120 | 0.034 [0.038] N=958 | 0.013 [0.042] N=778 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.249 P=341 | 0.070 [0.050] N=1,391 | 0.059 [0.055] N=1,265 | 0.056 [0.055] N=1,095 | 0.019 [0.059] N=936 | 0.063 [0.065] N=758 |
| BMI-for-age z score below -2 | 0.113 P=287 | 0.028 [0.027] N=2,555 | 0.017 [0.028] N=2,322 | 0.008 [0.029] N=2,049 | -0.004 [0.029] N=1,735 | 0.011 [0.033] N=1,411 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.391 P=440 | -0.007 [0.060] N=1,142 | -0.024 [0.063] N=1,035 | 0.001 [0.066] N=934 | 0.013 [0.073] N=781 | -0.017 [0.081] N=630 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.217 P=243 | 0.013 [0.050] N=1,142 | -0.021 [0.054] N=1,035 | -0.009 [0.056] N=934 | 0.014 [0.062] N=781 | 0.011 [0.073] N=630 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=129 | 0.027 [0.038] N=1,142 | 0.028 [0.040] N=1,035 | 0.002 [0.040] N=934 | -0.028 [0.045] N=781 | -0.013 [0.049] N=630 |

*Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.119 [0.233] N=2,286 | 0.145 [0.261] N=2,041 | 0.039 [0.286] N=1,756 | -0.145 [0.301] N=1,429 | -0.076 [0.371] N=1,091 |
| WHO weight-for-age z score | -0.127 [0.149] N=2,418 | -0.089 [0.162] N=2,170 | -0.177 [0.184] N=1,866 | **-0.364* [0.203] N=1,523** | **-0.474* [0.242] N=1,158** |
| WHO weight-for-length/height z score, 0-60 months | -0.044 [0.308] N=1,237 | 0.023 [0.325] N=1,111 | -0.220 [0.386] N=938 | -0.322 [0.418] N=776 | -0.917 [0.553] N=596 |
| WHO BMI-for-age z score | **-0.401** [0.182] N=2,192** | **-0.410** [0.193] N=1,959** | **-0.399* [0.218] N=1,686** | **-0.501** [0.249] N=1,372** | **-1.123*** [0.321] N=1,048** |
| Stunted: HAZ below -2 | 0.016 [0.043] N=2,286 | 0.019 [0.048] N=2,041 | 0.019 [0.052] N=1,756 | 0.019 [0.061] N=1,429 | 0.022 [0.074] N=1,091 |
| Underweight: WAZ below -2 | -0.006 [0.021] N=2,418 | -0.018 [0.024] N=2,170 | -0.021 [0.027] N=1,866 | -0.017 [0.028] N=1,523 | -0.009 [0.032] N=1,158 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.062 [0.044] N=1,237 | 0.039 [0.047] N=1,111 | 0.074 [0.054] N=938 | 0.054 [0.059] N=776 | **0.138** [0.069] N=596** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.024 [0.063] N=1,237 | 0.022 [0.066] N=1,111 | -0.012 [0.071] N=938 | -0.049 [0.087] N=776 | -0.102 [0.110] N=596 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.040 [0.046] N=1,237 | 0.030 [0.048] N=1,111 | 0.005 [0.055] N=938 | 0.003 [0.066] N=776 | -0.092 [0.079] N=596 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.093 [0.069] N=1,214 | 0.082 [0.078] N=1,088 | 0.094 [0.085] N=918 | 0.005 [0.098] N=759 | 0.107 [0.119] N=581 |
| BMI-for-age z score below -2 | **0.088*** [0.027] N=2,192** | **0.078*** [0.029] N=1,959** | **0.070** [0.034] N=1,686** | **0.065** [0.032] N=1,372** | **0.160*** [0.044] N=1,048** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.060 [0.081] N=961 | -0.094 [0.091] N=854 | -0.058 [0.099] N=753 | -0.045 [0.121] N=600 | -0.193 [0.140] N=449 |
| WHO 2007 BMI obesity above +2, 61+ months | -0.055 [0.068] N=961 | **-0.124* [0.073] N=854** | -0.117 [0.078] N=753 | -0.110 [0.089] N=600 | **-0.205* [0.116] N=449** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.102** [0.047] N=961** | **0.115** [0.051] N=854** | 0.069 [0.054] N=753 | 0.029 [0.064] N=600 | 0.108 [0.081] N=449 |

*Specification 6 Monthly fixed effects female and any aid (0/1)  No donut hole*

| Outcome | Mean | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|---|
| WHO length/height-for-age z score | -0.042 | -0.178 [0.189] N=2,667 | -0.185 [0.204] N=2,422 | -0.237 [0.214] N=2,137 | -0.374 [0.227] N=1,810 | -0.417 [0.253] N=1,472 |
| WHO weight-for-age z score | 0.374 | -0.058 [0.104] N=2,816 | -0.031 [0.109] N=2,568 | -0.056 [0.116] N=2,264 | -0.131 [0.125] N=1,921 | -0.150 [0.140] N=1,556 |
| WHO weight-for-length/height z score, 0-60 months | 0.379 | 0.121 [0.238] N=1,419 | 0.188 [0.249] N=1,293 | 0.126 [0.274] N=1,120 | 0.156 [0.294] N=958 | 0.035 [0.347] N=778 |
| WHO BMI-for-age z score | 0.389 | 0.100 [0.176] N=2,555 | 0.146 [0.184] N=2,322 | 0.216 [0.192] N=2,049 | 0.273 [0.208] N=1,735 | 0.220 [0.242] N=1,411 |
| Stunted: HAZ below -2 | 0.179 P=467 | 0.027 [0.032] N=2,667 | 0.029 [0.035] N=2,422 | 0.026 [0.037] N=2,137 | 0.033 [0.040] N=1,810 | 0.041 [0.045] N=1,472 |
| Underweight: WAZ below -2 | 0.041 P=117 | 0.008 [0.014] N=2,816 | 0.004 [0.014] N=2,568 | 0.005 [0.015] N=2,264 | 0.014 [0.016] N=1,921 | 0.026 [0.016] N=1,556 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.108 P=152 | 0.011 [0.038] N=1,419 | -0.006 [0.041] N=1,293 | 0.002 [0.043] N=1,120 | -0.013 [0.045] N=958 | 0.000 [0.053] N=778 |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.196 P=280 | 0.030 [0.040] N=1,419 | 0.034 [0.041] N=1,293 | 0.024 [0.042] N=1,120 | 0.023 [0.046] N=958 | 0.018 [0.053] N=778 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.091 P=134 | 0.039 [0.030] N=1,419 | 0.036 [0.030] N=1,293 | 0.026 [0.033] N=1,120 | 0.032 [0.038] N=958 | 0.009 [0.042] N=778 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.249 P=341 | 0.064 [0.049] N=1,391 | 0.052 [0.054] N=1,265 | 0.046 [0.055] N=1,095 | 0.010 [0.059] N=936 | 0.051 [0.066] N=758 |
| BMI-for-age z score below -2 | 0.113 P=287 | 0.028 [0.027] N=2,555 | 0.016 [0.028] N=2,322 | 0.007 [0.029] N=2,049 | -0.006 [0.029] N=1,735 | 0.009 [0.034] N=1,411 |
| WHO 2007 BMI overweight above +1, 61+ months | 0.391 P=440 | -0.012 [0.060] N=1,142 | -0.028 [0.064] N=1,035 | -0.003 [0.066] N=934 | 0.010 [0.074] N=781 | -0.024 [0.082] N=630 |
| WHO 2007 BMI obesity above +2, 61+ months | 0.217 P=243 | 0.011 [0.050] N=1,142 | -0.023 [0.054] N=1,035 | -0.011 [0.056] N=934 | 0.012 [0.062] N=781 | 0.008 [0.072] N=630 |
| WHO 2007 BMI thinness below -2, 61+ months | 0.110 P=129 | 0.027 [0.038] N=1,142 | 0.028 [0.039] N=1,035 | 0.000 [0.039] N=934 | -0.030 [0.044] N=781 | -0.017 [0.047] N=630 |

*Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole*

| Outcome | 8 years | 7 years | 6 years | 5 years | 4 years |
|---|---|---|---|---|---|
| WHO length/height-for-age z score | 0.106 [0.231] N=2,286 | 0.136 [0.257] N=2,041 | 0.036 [0.283] N=1,756 | -0.168 [0.302] N=1,429 | -0.085 [0.373] N=1,091 |
| WHO weight-for-age z score | -0.129 [0.148] N=2,418 | -0.090 [0.162] N=2,170 | -0.183 [0.184] N=1,866 | **-0.366* [0.203] N=1,523** | **-0.480* [0.246] N=1,158** |
| WHO weight-for-length/height z score, 0-60 months | -0.034 [0.307] N=1,237 | 0.028 [0.324] N=1,111 | -0.206 [0.384] N=938 | -0.294 [0.419] N=776 | **-0.943* [0.537] N=596** |
| WHO BMI-for-age z score | **-0.397** [0.182] N=2,192** | **-0.408** [0.192] N=1,959** | **-0.398* [0.216] N=1,686** | **-0.487* [0.250] N=1,372** | **-1.118*** [0.320] N=1,048** |
| Stunted: HAZ below -2 | 0.017 [0.043] N=2,286 | 0.020 [0.048] N=2,041 | 0.020 [0.052] N=1,756 | 0.024 [0.061] N=1,429 | 0.027 [0.074] N=1,091 |
| Underweight: WAZ below -2 | -0.006 [0.021] N=2,418 | -0.017 [0.023] N=2,170 | -0.019 [0.027] N=1,866 | -0.016 [0.028] N=1,523 | -0.003 [0.031] N=1,158 |
| Wasted: WLZ/WHZ below -2, 0-60 months | 0.059 [0.044] N=1,237 | 0.037 [0.047] N=1,111 | 0.070 [0.054] N=938 | 0.050 [0.057] N=776 | **0.136** [0.067] N=596** |
| Overweight: WLZ/WHZ above +2, 0-60 months | 0.026 [0.062] N=1,237 | 0.022 [0.066] N=1,111 | -0.011 [0.071] N=938 | -0.044 [0.087] N=776 | -0.106 [0.107] N=596 |
| Obese: WLZ/WHZ above +3, 0-60 months | 0.041 [0.045] N=1,237 | 0.030 [0.047] N=1,111 | 0.005 [0.054] N=938 | 0.006 [0.065] N=776 | -0.094 [0.076] N=596 |
| CIAF: any stunting, underweight, or wasting, 0-60 months | 0.092 [0.069] N=1,214 | 0.081 [0.078] N=1,088 | 0.092 [0.085] N=918 | 0.012 [0.098] N=759 | 0.119 [0.120] N=581 |
| BMI-for-age z score below -2 | **0.088*** [0.026] N=2,192** | **0.079*** [0.029] N=1,959** | **0.070** [0.034] N=1,686** | **0.065** [0.031] N=1,372** | **0.161*** [0.044] N=1,048** |
| WHO 2007 BMI overweight above +1, 61+ months | -0.063 [0.082] N=961 | -0.094 [0.091] N=854 | -0.057 [0.099] N=753 | -0.040 [0.122] N=600 | -0.192 [0.142] N=449 |
| WHO 2007 BMI obesity above +2, 61+ months | -0.055 [0.068] N=961 | **-0.124* [0.073] N=854** | -0.118 [0.078] N=753 | -0.110 [0.088] N=600 | **-0.204* [0.115] N=449** |
| WHO 2007 BMI thinness below -2, 61+ months | **0.105** [0.047] N=961** | **0.119** [0.050] N=854** | 0.069 [0.053] N=753 | 0.030 [0.063] N=600 | 0.102 [0.077] N=449 |

*Notes:* Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 83 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.