Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-6 Using 16 December 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 83 months on 16 December 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.457	0.063 [0.145] N=2,793	0.023 [0.154] N=2,587	-0.015 [0.169] N=2,321	0.030 [0.171] N=1,992	0.068 [0.194] N=1,645
WHO weight-for-age z score	0.071	0.045 [0.093] N=2,970	0.078 [0.100] N=2,743	0.118 [0.104] N=2,459	0.058 [0.115] N=2,108	0.088 [0.127] N=1,739
WHO weight-for-length/height z score, 0-60 months	0.330	0.269 [0.197] N=1,583	0.218 [0.205] N=1,464	0.104 [0.220] N=1,312	-0.019 [0.246] N=1,119	0.132 [0.263] N=912
WHO BMI-for-age z score	0.353	0.095 [0.154] N=2,673	0.107 [0.161] N=2,476	0.119 [0.175] N=2,215	-0.087 [0.195] N=1,903	-0.013 [0.216] N=1,571
Stunted: HAZ below -2	0.227 P=619	-0.018 [0.027] N=2,793	-0.015 [0.028] N=2,587	-0.001 [0.031] N=2,321	-0.021 [0.031] N=1,992	-0.003 [0.034] N=1,645
Underweight: WAZ below -2	0.059 P=171	0.000 [0.015] N=2,970	-0.004 [0.016] N=2,743	-0.016 [0.017] N=2,459	-0.010 [0.019] N=2,108	-0.012 [0.022] N=1,739
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=176	-0.043 [0.032] N=1,583	-0.035 [0.033] N=1,464	-0.014 [0.035] N=1,312	-0.008 [0.042] N=1,119	-0.032 [0.044] N=912
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=292	0.072 [0.044] N=1,583	0.062 [0.047] N=1,464	0.049 [0.051] N=1,312	0.017 [0.057] N=1,119	0.031 [0.062] N=912
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=126	0.027 [0.031] N=1,583	0.026 [0.032] N=1,464	0.030 [0.036] N=1,312	0.021 [0.039] N=1,119	0.021 [0.046] N=912
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=481	-0.033 [0.043] N=1,555	-0.038 [0.046] N=1,440	0.000 [0.049] N=1,291	-0.015 [0.052] N=1,100	0.001 [0.057] N=899
BMI-for-age z score below -2	0.113 P=310	-0.003 [0.027] N=2,673	-0.002 [0.027] N=2,476	0.002 [0.029] N=2,215	0.022 [0.032] N=1,903	0.019 [0.035] N=1,571
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=406	0.033 [0.056] N=1,086	0.039 [0.059] N=1,008	0.090 [0.065] N=903	0.048 [0.072] N=785	0.027 [0.084] N=654
WHO 2007 BMI obesity above +2, 61+ months	0.207 P=228	0.000 [0.045] N=1,086	0.001 [0.046] N=1,008	0.025 [0.048] N=903	-0.013 [0.053] N=785	-0.075 [0.060] N=654
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=118	0.035 [0.043] N=1,086	0.023 [0.044] N=1,008	-0.009 [0.049] N=903	0.014 [0.053] N=785	0.022 [0.061] N=654

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.054 [0.205] N=2,351	-0.162 [0.221] N=2,145	-0.295 [0.260] N=1,879	-0.225 [0.288] N=1,550	-0.241 [0.379] N=1,203
WHO weight-for-age z score	-0.048 [0.121] N=2,504	-0.004 [0.135] N=2,277	0.044 [0.145] N=1,993	-0.112 [0.174] N=1,642	-0.095 [0.219] N=1,273
WHO weight-for-length/height z score, 0-60 months	0.201 [0.287] N=1,345	0.102 [0.307] N=1,226	-0.161 [0.353] N=1,074	-0.517 [0.418] N=881	-0.471 [0.548] N=674
WHO BMI-for-age z score	0.266 [0.202] N=2,254	0.325 [0.212] N=2,057	0.402 [0.245] N=1,796	0.029 [0.302] N=1,484	0.264 [0.379] N=1,152
Stunted: HAZ below -2	0.001 [0.041] N=2,351	0.009 [0.044] N=2,145	0.040 [0.054] N=1,879	-0.004 [0.060] N=1,550	0.052 [0.076] N=1,203
Underweight: WAZ below -2	0.013 [0.021] N=2,504	0.007 [0.023] N=2,277	-0.014 [0.025] N=1,993	0.004 [0.029] N=1,642	0.016 [0.040] N=1,273
Wasted: WLZ/WHZ below -2, 0-60 months	-0.033 [0.040] N=1,345	-0.019 [0.041] N=1,226	0.020 [0.046] N=1,074	0.039 [0.061] N=881	-0.011 [0.067] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.064 [0.059] N=1,345	0.046 [0.066] N=1,226	0.017 [0.075] N=1,074	-0.059 [0.090] N=881	-0.082 [0.122] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.038] N=1,345	0.026 [0.041] N=1,226	0.031 [0.048] N=1,074	0.014 [0.060] N=881	0.003 [0.090] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.022 [0.060] N=1,322	-0.033 [0.068] N=1,207	0.030 [0.078] N=1,058	-0.015 [0.095] N=867	0.008 [0.121] N=666
BMI-for-age z score below -2	-0.025 [0.033] N=2,254	-0.031 [0.031] N=2,057	-0.036 [0.037] N=1,796	-0.001 [0.041] N=1,484	-0.023 [0.052] N=1,152
WHO 2007 BMI overweight above +1, 61+ months	0.086 [0.074] N=905	0.111 [0.080] N=827	**0.220** [0.091] N=722**	0.166 [0.110] N=604	0.179 [0.149] N=473
WHO 2007 BMI obesity above +2, 61+ months	0.095 [0.064] N=905	0.110 [0.069] N=827	**0.183** [0.077] N=722**	0.153 [0.095] N=604	0.066 [0.121] N=473
WHO 2007 BMI thinness below -2, 61+ months	0.016 [0.049] N=905	-0.016 [0.050] N=827	-0.088 [0.058] N=722	-0.066 [0.066] N=604	-0.088 [0.087] N=473

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.457	0.067 [0.144] N=2,793	0.027 [0.153] N=2,587	-0.006 [0.168] N=2,321	0.037 [0.169] N=1,992	0.070 [0.192] N=1,645
WHO weight-for-age z score	0.071	0.048 [0.093] N=2,970	0.081 [0.100] N=2,743	0.124 [0.104] N=2,459	0.061 [0.115] N=2,108	0.088 [0.127] N=1,739
WHO weight-for-length/height z score, 0-60 months	0.330	0.263 [0.198] N=1,583	0.209 [0.206] N=1,464	0.090 [0.223] N=1,312	-0.027 [0.249] N=1,119	0.121 [0.266] N=912
WHO BMI-for-age z score	0.353	0.095 [0.155] N=2,673	0.105 [0.161] N=2,476	0.116 [0.176] N=2,215	-0.088 [0.196] N=1,903	-0.013 [0.215] N=1,571
Stunted: HAZ below -2	0.227 P=619	-0.017 [0.027] N=2,793	-0.014 [0.028] N=2,587	-0.001 [0.031] N=2,321	-0.021 [0.031] N=1,992	-0.004 [0.034] N=1,645
Underweight: WAZ below -2	0.059 P=171	0.000 [0.015] N=2,970	-0.004 [0.016] N=2,743	-0.016 [0.017] N=2,459	-0.010 [0.019] N=2,108	-0.012 [0.022] N=1,739
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=176	-0.041 [0.032] N=1,583	-0.032 [0.033] N=1,464	-0.010 [0.035] N=1,312	-0.005 [0.042] N=1,119	-0.028 [0.044] N=912
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=292	0.072 [0.044] N=1,583	0.062 [0.047] N=1,464	0.050 [0.051] N=1,312	0.018 [0.057] N=1,119	0.033 [0.062] N=912
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=126	0.027 [0.030] N=1,583	0.027 [0.032] N=1,464	0.031 [0.035] N=1,312	0.022 [0.039] N=1,119	0.021 [0.046] N=912
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=481	-0.032 [0.043] N=1,555	-0.035 [0.045] N=1,440	0.004 [0.049] N=1,291	-0.012 [0.052] N=1,100	0.004 [0.057] N=899
BMI-for-age z score below -2	0.113 P=310	-0.002 [0.027] N=2,673	-0.001 [0.027] N=2,476	0.004 [0.029] N=2,215	0.023 [0.032] N=1,903	0.019 [0.035] N=1,571
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=406	0.033 [0.056] N=1,086	0.039 [0.059] N=1,008	0.090 [0.065] N=903	0.047 [0.072] N=785	0.026 [0.084] N=654
WHO 2007 BMI obesity above +2, 61+ months	0.207 P=228	0.001 [0.045] N=1,086	0.002 [0.046] N=1,008	0.027 [0.049] N=903	-0.014 [0.053] N=785	-0.078 [0.060] N=654
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=118	0.035 [0.043] N=1,086	0.023 [0.044] N=1,008	-0.009 [0.049] N=903	0.014 [0.052] N=785	0.022 [0.060] N=654

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.050 [0.204] N=2,351	-0.156 [0.220] N=2,145	-0.277 [0.260] N=1,879	-0.213 [0.287] N=1,550	-0.252 [0.379] N=1,203
WHO weight-for-age z score	-0.045 [0.120] N=2,504	-0.002 [0.134] N=2,277	0.053 [0.144] N=1,993	-0.107 [0.171] N=1,642	-0.101 [0.217] N=1,273
WHO weight-for-length/height z score, 0-60 months	0.192 [0.289] N=1,345	0.083 [0.311] N=1,226	-0.196 [0.356] N=1,074	-0.552 [0.424] N=881	-0.524 [0.554] N=674
WHO BMI-for-age z score	0.265 [0.202] N=2,254	0.321 [0.212] N=2,057	0.394 [0.246] N=1,796	0.027 [0.304] N=1,484	0.266 [0.380] N=1,152
Stunted: HAZ below -2	0.001 [0.041] N=2,351	0.009 [0.044] N=2,145	0.040 [0.054] N=1,879	-0.004 [0.060] N=1,550	0.053 [0.076] N=1,203
Underweight: WAZ below -2	0.013 [0.021] N=2,504	0.007 [0.023] N=2,277	-0.014 [0.025] N=1,993	0.004 [0.029] N=1,642	0.016 [0.040] N=1,273
Wasted: WLZ/WHZ below -2, 0-60 months	-0.031 [0.041] N=1,345	-0.015 [0.041] N=1,226	0.027 [0.046] N=1,074	0.046 [0.062] N=881	-0.001 [0.069] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.063 [0.059] N=1,345	0.046 [0.066] N=1,226	0.017 [0.075] N=1,074	-0.058 [0.091] N=881	-0.079 [0.122] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.038] N=1,345	0.026 [0.041] N=1,226	0.030 [0.048] N=1,074	0.013 [0.060] N=881	0.002 [0.091] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.021 [0.060] N=1,322	-0.030 [0.068] N=1,207	0.036 [0.078] N=1,058	-0.010 [0.094] N=867	0.013 [0.121] N=666
BMI-for-age z score below -2	-0.025 [0.033] N=2,254	-0.029 [0.031] N=2,057	-0.033 [0.037] N=1,796	0.000 [0.041] N=1,484	-0.024 [0.052] N=1,152
WHO 2007 BMI overweight above +1, 61+ months	0.086 [0.074] N=905	0.111 [0.080] N=827	**0.220** [0.092] N=722**	0.165 [0.111] N=604	0.177 [0.150] N=473
WHO 2007 BMI obesity above +2, 61+ months	0.095 [0.064] N=905	0.110 [0.068] N=827	**0.186** [0.076] N=722**	0.149 [0.095] N=604	0.049 [0.120] N=473
WHO 2007 BMI thinness below -2, 61+ months	0.016 [0.049] N=905	-0.016 [0.050] N=827	-0.089 [0.058] N=722	-0.064 [0.066] N=604	-0.086 [0.086] N=473

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.457	0.031 [0.146] N=2,793	-0.006 [0.155] N=2,587	-0.037 [0.170] N=2,321	0.006 [0.173] N=1,992	0.018 [0.196] N=1,645
WHO weight-for-age z score	0.071	0.009 [0.091] N=2,970	0.045 [0.099] N=2,743	0.085 [0.103] N=2,459	0.026 [0.112] N=2,108	0.047 [0.125] N=1,739
WHO weight-for-length/height z score, 0-60 months	0.330	0.236 [0.199] N=1,583	0.188 [0.207] N=1,464	0.078 [0.221] N=1,312	-0.034 [0.248] N=1,119	0.124 [0.267] N=912
WHO BMI-for-age z score	0.353	0.087 [0.155] N=2,673	0.100 [0.161] N=2,476	0.113 [0.176] N=2,215	-0.088 [0.195] N=1,903	-0.011 [0.217] N=1,571
Stunted: HAZ below -2	0.227 P=619	-0.010 [0.027] N=2,793	-0.006 [0.027] N=2,587	0.007 [0.031] N=2,321	-0.013 [0.031] N=1,992	0.010 [0.035] N=1,645
Underweight: WAZ below -2	0.059 P=171	0.007 [0.015] N=2,970	0.003 [0.016] N=2,743	-0.008 [0.016] N=2,459	-0.003 [0.018] N=2,108	-0.002 [0.021] N=1,739
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=176	-0.031 [0.032] N=1,583	-0.022 [0.033] N=1,464	-0.003 [0.035] N=1,312	0.001 [0.042] N=1,119	-0.023 [0.044] N=912
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=292	0.072 [0.045] N=1,583	0.064 [0.047] N=1,464	0.052 [0.051] N=1,312	0.019 [0.057] N=1,119	0.032 [0.063] N=912
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=126	0.029 [0.030] N=1,583	0.030 [0.032] N=1,464	0.033 [0.035] N=1,312	0.023 [0.038] N=1,119	0.022 [0.046] N=912
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=481	-0.007 [0.041] N=1,555	-0.009 [0.043] N=1,440	0.022 [0.046] N=1,291	0.004 [0.048] N=1,100	0.032 [0.054] N=899
BMI-for-age z score below -2	0.113 P=310	0.003 [0.027] N=2,673	0.005 [0.027] N=2,476	0.008 [0.030] N=2,215	0.027 [0.032] N=1,903	0.025 [0.035] N=1,571
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=406	0.033 [0.057] N=1,086	0.040 [0.060] N=1,008	0.094 [0.066] N=903	0.055 [0.073] N=785	0.029 [0.085] N=654
WHO 2007 BMI obesity above +2, 61+ months	0.207 P=228	-0.001 [0.045] N=1,086	-0.001 [0.046] N=1,008	0.026 [0.049] N=903	-0.015 [0.053] N=785	-0.079 [0.060] N=654
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=118	0.035 [0.042] N=1,086	0.022 [0.043] N=1,008	-0.009 [0.049] N=903	0.016 [0.052] N=785	0.020 [0.060] N=654

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.087 [0.203] N=2,351	-0.191 [0.218] N=2,145	-0.312 [0.258] N=1,879	-0.254 [0.287] N=1,550	-0.331 [0.378] N=1,203
WHO weight-for-age z score	-0.082 [0.117] N=2,504	-0.032 [0.132] N=2,277	0.019 [0.141] N=1,993	-0.133 [0.166] N=1,642	-0.133 [0.213] N=1,273
WHO weight-for-length/height z score, 0-60 months	0.161 [0.287] N=1,345	0.066 [0.311] N=1,226	-0.208 [0.352] N=1,074	-0.578 [0.421] N=881	-0.534 [0.557] N=674
WHO BMI-for-age z score	0.260 [0.202] N=2,254	0.322 [0.212] N=2,057	0.401 [0.244] N=1,796	0.047 [0.301] N=1,484	0.305 [0.382] N=1,152
Stunted: HAZ below -2	0.008 [0.040] N=2,351	0.016 [0.043] N=2,145	0.046 [0.053] N=1,879	0.002 [0.059] N=1,550	0.070 [0.075] N=1,203
Underweight: WAZ below -2	0.018 [0.020] N=2,504	0.012 [0.022] N=2,277	-0.009 [0.024] N=1,993	0.006 [0.027] N=1,642	0.023 [0.038] N=1,273
Wasted: WLZ/WHZ below -2, 0-60 months	-0.019 [0.041] N=1,345	-0.003 [0.042] N=1,226	0.033 [0.046] N=1,074	0.053 [0.062] N=881	0.006 [0.070] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.065 [0.059] N=1,345	0.050 [0.066] N=1,226	0.018 [0.074] N=1,074	-0.061 [0.090] N=881	-0.075 [0.123] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.030 [0.038] N=1,345	0.029 [0.041] N=1,226	0.030 [0.048] N=1,074	0.010 [0.059] N=881	-0.006 [0.091] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.003 [0.058] N=1,322	-0.005 [0.067] N=1,207	0.045 [0.076] N=1,058	-0.010 [0.092] N=867	0.030 [0.121] N=666
BMI-for-age z score below -2	-0.020 [0.033] N=2,254	-0.025 [0.032] N=2,057	-0.031 [0.037] N=1,796	0.001 [0.042] N=1,484	-0.021 [0.053] N=1,152
WHO 2007 BMI overweight above +1, 61+ months	0.087 [0.075] N=905	0.113 [0.081] N=827	**0.230** [0.092] N=722**	**0.184* [0.110] N=604**	0.196 [0.147] N=473
WHO 2007 BMI obesity above +2, 61+ months	0.096 [0.064] N=905	0.110 [0.069] N=827	**0.191** [0.076] N=722**	0.155 [0.094] N=604	0.059 [0.121] N=473
WHO 2007 BMI thinness below -2, 61+ months	0.015 [0.049] N=905	-0.016 [0.049] N=827	-0.090 [0.058] N=722	-0.062 [0.065] N=604	-0.094 [0.088] N=473

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.457	0.025 [0.145] N=2,793	-0.011 [0.154] N=2,587	-0.042 [0.170] N=2,321	-0.008 [0.173] N=1,992	-0.000 [0.197] N=1,645
WHO weight-for-age z score	0.071	0.018 [0.091] N=2,970	0.050 [0.099] N=2,743	0.088 [0.104] N=2,459	0.026 [0.114] N=2,108	0.039 [0.126] N=1,739
WHO weight-for-length/height z score, 0-60 months	0.330	0.271 [0.198] N=1,583	0.220 [0.206] N=1,464	0.105 [0.222] N=1,312	-0.014 [0.248] N=1,119	0.160 [0.267] N=912
WHO BMI-for-age z score	0.353	0.103 [0.156] N=2,673	0.111 [0.162] N=2,476	0.120 [0.177] N=2,215	-0.080 [0.198] N=1,903	-0.000 [0.220] N=1,571
Stunted: HAZ below -2	0.227 P=619	-0.008 [0.027] N=2,793	-0.005 [0.028] N=2,587	0.008 [0.031] N=2,321	-0.009 [0.031] N=1,992	0.013 [0.035] N=1,645
Underweight: WAZ below -2	0.059 P=171	0.006 [0.015] N=2,970	0.003 [0.016] N=2,743	-0.008 [0.017] N=2,459	-0.002 [0.019] N=2,108	-0.002 [0.021] N=1,739
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=176	-0.035 [0.031] N=1,583	-0.027 [0.033] N=1,464	-0.007 [0.035] N=1,312	-0.002 [0.042] N=1,119	-0.027 [0.045] N=912
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=292	**0.075* [0.044] N=1,583**	0.065 [0.047] N=1,464	0.053 [0.050] N=1,312	0.022 [0.056] N=1,119	0.035 [0.061] N=912
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=126	0.033 [0.031] N=1,583	0.033 [0.032] N=1,464	0.036 [0.036] N=1,312	0.029 [0.039] N=1,119	0.032 [0.047] N=912
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=481	-0.008 [0.040] N=1,555	-0.012 [0.043] N=1,440	0.022 [0.046] N=1,291	0.012 [0.048] N=1,100	0.036 [0.053] N=899
BMI-for-age z score below -2	0.113 P=310	0.000 [0.027] N=2,673	0.001 [0.027] N=2,476	0.005 [0.029] N=2,215	0.025 [0.031] N=1,903	0.021 [0.035] N=1,571
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=406	0.035 [0.057] N=1,086	0.041 [0.060] N=1,008	0.094 [0.066] N=903	0.051 [0.074] N=785	0.035 [0.086] N=654
WHO 2007 BMI obesity above +2, 61+ months	0.207 P=228	0.001 [0.045] N=1,086	0.000 [0.047] N=1,008	0.026 [0.049] N=903	-0.014 [0.053] N=785	-0.076 [0.061] N=654
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=118	0.033 [0.042] N=1,086	0.021 [0.043] N=1,008	-0.010 [0.048] N=903	0.016 [0.052] N=785	0.019 [0.058] N=654

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.080 [0.202] N=2,351	-0.185 [0.218] N=2,145	-0.304 [0.257] N=1,879	-0.237 [0.287] N=1,550	-0.323 [0.377] N=1,203
WHO weight-for-age z score	-0.069 [0.119] N=2,504	-0.028 [0.133] N=2,277	0.020 [0.144] N=1,993	-0.134 [0.170] N=1,642	-0.125 [0.212] N=1,273
WHO weight-for-length/height z score, 0-60 months	0.192 [0.287] N=1,345	0.099 [0.308] N=1,226	-0.183 [0.354] N=1,074	-0.559 [0.419] N=881	-0.512 [0.545] N=674
WHO BMI-for-age z score	0.263 [0.203] N=2,254	0.320 [0.212] N=2,057	0.394 [0.245] N=1,796	0.025 [0.302] N=1,484	0.297 [0.377] N=1,152
Stunted: HAZ below -2	0.008 [0.040] N=2,351	0.016 [0.043] N=2,145	0.046 [0.054] N=1,879	0.001 [0.059] N=1,550	0.071 [0.076] N=1,203
Underweight: WAZ below -2	0.017 [0.021] N=2,504	0.012 [0.022] N=2,277	-0.008 [0.025] N=1,993	0.009 [0.028] N=1,642	0.026 [0.039] N=1,273
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.040] N=1,345	-0.009 [0.041] N=1,226	0.030 [0.045] N=1,074	0.051 [0.061] N=881	0.007 [0.068] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.071 [0.060] N=1,345	0.053 [0.066] N=1,226	0.022 [0.075] N=1,074	-0.056 [0.090] N=881	-0.065 [0.121] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.036 [0.038] N=1,345	0.034 [0.041] N=1,226	0.035 [0.048] N=1,074	0.014 [0.059] N=881	0.001 [0.087] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.003 [0.057] N=1,322	-0.010 [0.065] N=1,207	0.046 [0.074] N=1,058	-0.000 [0.089] N=867	0.034 [0.117] N=666
BMI-for-age z score below -2	-0.019 [0.033] N=2,254	-0.025 [0.032] N=2,057	-0.029 [0.037] N=1,796	0.007 [0.041] N=1,484	-0.013 [0.052] N=1,152
WHO 2007 BMI overweight above +1, 61+ months	0.084 [0.075] N=905	0.112 [0.081] N=827	**0.227** [0.091] N=722**	0.177 [0.110] N=604	0.218 [0.145] N=473
WHO 2007 BMI obesity above +2, 61+ months	0.093 [0.064] N=905	0.107 [0.068] N=827	**0.184** [0.075] N=722**	0.147 [0.094] N=604	0.067 [0.121] N=473
WHO 2007 BMI thinness below -2, 61+ months	0.017 [0.049] N=905	-0.013 [0.050] N=827	-0.086 [0.057] N=722	-0.061 [0.064] N=604	-0.084 [0.083] N=473

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.461	0.032 [0.155] N=2,609	-0.029 [0.164] N=2,421	-0.067 [0.177] N=2,165	-0.045 [0.178] N=1,865	-0.012 [0.207] N=1,536
WHO weight-for-age z score	0.070	0.054 [0.094] N=2,768	0.076 [0.099] N=2,562	0.114 [0.107] N=2,290	0.044 [0.120] N=1,970	0.086 [0.134] N=1,623
WHO weight-for-length/height z score, 0-60 months	0.347	**0.381* [0.200] N=1,466**	0.338 [0.208] N=1,360	0.197 [0.221] N=1,216	0.080 [0.253] N=1,045	0.280 [0.273] N=848
WHO BMI-for-age z score	0.362	0.147 [0.152] N=2,496	0.151 [0.157] N=2,317	0.164 [0.174] N=2,068	-0.024 [0.196] N=1,784	0.068 [0.222] N=1,469
Stunted: HAZ below -2	0.229 P=580	-0.007 [0.029] N=2,609	0.001 [0.030] N=2,421	0.017 [0.033] N=2,165	-0.001 [0.033] N=1,865	0.016 [0.039] N=1,536
Underweight: WAZ below -2	0.061 P=164	0.001 [0.016] N=2,768	-0.003 [0.017] N=2,562	-0.013 [0.018] N=2,290	-0.007 [0.020] N=1,970	-0.007 [0.024] N=1,623
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=162	-0.051 [0.033] N=1,466	-0.043 [0.034] N=1,360	-0.014 [0.036] N=1,216	-0.008 [0.044] N=1,045	-0.032 [0.047] N=848
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=273	**0.077* [0.046] N=1,466**	0.071 [0.049] N=1,360	0.055 [0.054] N=1,216	0.017 [0.060] N=1,045	0.037 [0.068] N=848
Obese: WLZ/WHZ above +3, 0-60 months	0.083 P=121	0.032 [0.032] N=1,466	0.036 [0.033] N=1,360	0.041 [0.037] N=1,216	0.028 [0.040] N=1,045	0.032 [0.048] N=848
CIAF: any stunting, underweight, or wasting, 0-60 months	0.314 P=446	-0.022 [0.049] N=1,441	-0.024 [0.052] N=1,338	0.018 [0.056] N=1,196	-0.004 [0.060] N=1,027	0.019 [0.069] N=835
BMI-for-age z score below -2	0.112 P=285	-0.003 [0.028] N=2,496	-0.001 [0.027] N=2,317	0.003 [0.030] N=2,068	0.020 [0.033] N=1,784	0.011 [0.037] N=1,469
WHO 2007 BMI overweight above +1, 61+ months	0.369 P=382	0.039 [0.056] N=1,029	0.046 [0.058] N=957	0.099 [0.064] N=855	0.053 [0.070] N=743	0.033 [0.083] N=619
WHO 2007 BMI obesity above +2, 61+ months	0.204 P=214	0.001 [0.044] N=1,029	0.004 [0.046] N=957	0.031 [0.049] N=855	-0.017 [0.052] N=743	-0.095 [0.060] N=619
WHO 2007 BMI thinness below -2, 61+ months	0.108 P=113	0.046 [0.042] N=1,029	0.035 [0.043] N=957	0.002 [0.047] N=855	0.022 [0.052] N=743	0.021 [0.060] N=619

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.028 [0.224] N=2,197	-0.153 [0.241] N=2,009	-0.284 [0.281] N=1,753	-0.232 [0.315] N=1,453	-0.275 [0.428] N=1,124
WHO weight-for-age z score	-0.042 [0.120] N=2,334	-0.017 [0.130] N=2,128	0.020 [0.146] N=1,856	-0.163 [0.175] N=1,536	-0.151 [0.218] N=1,189
WHO weight-for-length/height z score, 0-60 months	0.248 [0.291] N=1,246	0.140 [0.311] N=1,140	-0.169 [0.347] N=996	-0.539 [0.423] N=825	-0.451 [0.564] N=628
WHO BMI-for-age z score	0.270 [0.200] N=2,106	0.309 [0.207] N=1,927	0.385 [0.239] N=1,678	0.026 [0.297] N=1,394	0.273 [0.388] N=1,079
Stunted: HAZ below -2	0.004 [0.043] N=2,197	0.017 [0.045] N=2,009	0.052 [0.057] N=1,753	0.010 [0.063] N=1,453	0.062 [0.080] N=1,124
Underweight: WAZ below -2	0.013 [0.022] N=2,334	0.007 [0.023] N=2,128	-0.011 [0.026] N=1,856	0.010 [0.031] N=1,536	0.028 [0.042] N=1,189
Wasted: WLZ/WHZ below -2, 0-60 months	-0.043 [0.040] N=1,246	-0.028 [0.040] N=1,140	0.023 [0.043] N=996	0.039 [0.060] N=825	-0.010 [0.066] N=628
Overweight: WLZ/WHZ above +2, 0-60 months	0.068 [0.059] N=1,246	0.055 [0.066] N=1,140	0.028 [0.075] N=996	-0.060 [0.092] N=825	-0.067 [0.129] N=628
Obese: WLZ/WHZ above +3, 0-60 months	0.028 [0.040] N=1,246	0.030 [0.044] N=1,140	0.033 [0.052] N=996	0.007 [0.064] N=825	0.001 [0.101] N=628
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.015 [0.064] N=1,226	-0.022 [0.071] N=1,123	0.052 [0.082] N=981	-0.007 [0.099] N=812	0.042 [0.127] N=620
BMI-for-age z score below -2	-0.020 [0.033] N=2,106	-0.024 [0.030] N=1,927	-0.024 [0.036] N=1,678	0.005 [0.041] N=1,394	-0.029 [0.052] N=1,079
WHO 2007 BMI overweight above +1, 61+ months	0.086 [0.076] N=857	0.113 [0.081] N=785	**0.223** [0.093] N=683**	0.156 [0.109] N=571	0.159 [0.157] N=447
WHO 2007 BMI obesity above +2, 61+ months	0.089 [0.064] N=857	0.108 [0.068] N=785	**0.186** [0.077] N=683**	0.133 [0.094] N=571	-0.000 [0.122] N=447
WHO 2007 BMI thinness below -2, 61+ months	0.033 [0.050] N=857	0.004 [0.050] N=785	-0.071 [0.058] N=683	-0.052 [0.067] N=571	-0.080 [0.090] N=447

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.461	0.042 [0.152] N=2,609	-0.019 [0.161] N=2,421	-0.068 [0.174] N=2,165	-0.041 [0.174] N=1,865	0.002 [0.200] N=1,536
WHO weight-for-age z score	0.070	0.065 [0.094] N=2,768	0.085 [0.100] N=2,562	0.123 [0.107] N=2,290	0.056 [0.120] N=1,970	0.102 [0.134] N=1,623
WHO weight-for-length/height z score, 0-60 months	0.347	**0.388* [0.201] N=1,466**	**0.348* [0.209] N=1,360**	0.203 [0.222] N=1,216	0.091 [0.253] N=1,045	0.283 [0.273] N=848
WHO BMI-for-age z score	0.362	0.148 [0.152] N=2,496	0.151 [0.157] N=2,317	0.165 [0.174] N=2,068	-0.023 [0.196] N=1,784	0.067 [0.221] N=1,469
Stunted: HAZ below -2	0.229 P=580	-0.009 [0.028] N=2,609	-0.001 [0.029] N=2,421	0.015 [0.033] N=2,165	-0.004 [0.032] N=1,865	0.010 [0.038] N=1,536
Underweight: WAZ below -2	0.061 P=164	-0.002 [0.016] N=2,768	-0.005 [0.017] N=2,562	-0.016 [0.018] N=2,290	-0.010 [0.020] N=1,970	-0.013 [0.024] N=1,623
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=162	-0.053 [0.032] N=1,466	-0.044 [0.034] N=1,360	-0.015 [0.036] N=1,216	-0.010 [0.044] N=1,045	-0.038 [0.047] N=848
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=273	**0.079* [0.047] N=1,466**	0.073 [0.050] N=1,360	0.058 [0.055] N=1,216	0.021 [0.061] N=1,045	0.041 [0.068] N=848
Obese: WLZ/WHZ above +3, 0-60 months	0.083 P=121	0.033 [0.032] N=1,466	0.037 [0.033] N=1,360	0.042 [0.037] N=1,216	0.031 [0.041] N=1,045	0.034 [0.048] N=848
CIAF: any stunting, underweight, or wasting, 0-60 months	0.314 P=446	-0.024 [0.049] N=1,441	-0.025 [0.052] N=1,338	0.017 [0.056] N=1,196	-0.005 [0.060] N=1,027	0.008 [0.067] N=835
BMI-for-age z score below -2	0.112 P=285	-0.004 [0.028] N=2,496	-0.002 [0.027] N=2,317	0.003 [0.030] N=2,068	0.019 [0.033] N=1,784	0.011 [0.037] N=1,469
WHO 2007 BMI overweight above +1, 61+ months	0.369 P=382	0.041 [0.055] N=1,029	0.048 [0.057] N=957	0.100 [0.063] N=855	0.053 [0.068] N=743	0.028 [0.081] N=619
WHO 2007 BMI obesity above +2, 61+ months	0.204 P=214	-0.001 [0.044] N=1,029	0.001 [0.046] N=957	0.030 [0.049] N=855	-0.017 [0.052] N=743	-0.095 [0.060] N=619
WHO 2007 BMI thinness below -2, 61+ months	0.108 P=113	0.045 [0.042] N=1,029	0.033 [0.042] N=957	0.002 [0.047] N=855	0.021 [0.052] N=743	0.022 [0.060] N=619

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.013 [0.222] N=2,197	-0.139 [0.239] N=2,009	-0.284 [0.280] N=1,753	-0.235 [0.314] N=1,453	-0.246 [0.423] N=1,124
WHO weight-for-age z score	-0.029 [0.119] N=2,334	-0.008 [0.131] N=2,128	0.033 [0.146] N=1,856	-0.143 [0.176] N=1,536	-0.118 [0.218] N=1,189
WHO weight-for-length/height z score, 0-60 months	0.267 [0.292] N=1,246	0.170 [0.315] N=1,140	-0.159 [0.351] N=996	-0.510 [0.428] N=825	-0.424 [0.573] N=628
WHO BMI-for-age z score	0.277 [0.200] N=2,106	0.314 [0.207] N=1,927	0.388 [0.239] N=1,678	0.031 [0.298] N=1,394	0.284 [0.394] N=1,079
Stunted: HAZ below -2	0.001 [0.042] N=2,197	0.015 [0.045] N=2,009	0.050 [0.056] N=1,753	0.007 [0.062] N=1,453	0.052 [0.080] N=1,124
Underweight: WAZ below -2	0.010 [0.022] N=2,334	0.005 [0.024] N=2,128	-0.014 [0.027] N=1,856	0.005 [0.031] N=1,536	0.018 [0.043] N=1,189
Wasted: WLZ/WHZ below -2, 0-60 months	-0.046 [0.039] N=1,246	-0.032 [0.040] N=1,140	0.021 [0.043] N=996	0.034 [0.060] N=825	-0.029 [0.063] N=628
Overweight: WLZ/WHZ above +2, 0-60 months	0.072 [0.059] N=1,246	0.061 [0.067] N=1,140	0.031 [0.076] N=996	-0.052 [0.093] N=825	-0.057 [0.130] N=628
Obese: WLZ/WHZ above +3, 0-60 months	0.027 [0.040] N=1,246	0.029 [0.045] N=1,140	0.035 [0.052] N=996	0.013 [0.064] N=825	0.010 [0.099] N=628
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.014 [0.063] N=1,226	-0.020 [0.071] N=1,123	0.049 [0.082] N=981	-0.011 [0.099] N=812	0.001 [0.121] N=620
BMI-for-age z score below -2	-0.021 [0.033] N=2,106	-0.025 [0.030] N=1,927	-0.024 [0.036] N=1,678	0.005 [0.042] N=1,394	-0.030 [0.053] N=1,079
WHO 2007 BMI overweight above +1, 61+ months	0.089 [0.075] N=857	0.115 [0.081] N=785	**0.225** [0.093] N=683**	0.159 [0.108] N=571	0.159 [0.152] N=447
WHO 2007 BMI obesity above +2, 61+ months	0.085 [0.063] N=857	0.101 [0.067] N=785	**0.183** [0.076] N=683**	0.129 [0.092] N=571	-0.003 [0.121] N=447
WHO 2007 BMI thinness below -2, 61+ months	0.030 [0.050] N=857	0.002 [0.049] N=785	-0.070 [0.058] N=683	-0.051 [0.066] N=571	-0.083 [0.090] N=447

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.409	-0.217 [0.181] N=2,767	-0.227 [0.193] N=2,514	-0.261 [0.203] N=2,211	**-0.372* [0.217] N=1,880**	**-0.420* [0.251] N=1,533**
WHO weight-for-age z score	0.092	-0.107 [0.100] N=2,939	-0.090 [0.104] N=2,681	-0.121 [0.112] N=2,354	-0.193 [0.119] N=2,001	-0.202 [0.135] N=1,620
WHO weight-for-length/height z score, 0-60 months	0.365	-0.029 [0.210] N=1,494	-0.017 [0.220] N=1,357	-0.065 [0.247] N=1,175	-0.026 [0.256] N=1,006	-0.061 [0.297] N=806
WHO BMI-for-age z score	0.376	0.063 [0.165] N=2,657	0.065 [0.172] N=2,414	0.106 [0.182] N=2,123	0.147 [0.196] N=1,804	0.131 [0.225] N=1,467
Stunted: HAZ below -2	0.225 P=616	0.048 [0.033] N=2,767	0.047 [0.035] N=2,514	0.046 [0.037] N=2,211	0.064 [0.039] N=1,880	0.074 [0.045] N=1,533
Underweight: WAZ below -2	0.060 P=172	0.010 [0.017] N=2,939	0.011 [0.018] N=2,681	0.015 [0.020] N=2,354	0.025 [0.022] N=2,001	**0.040* [0.024] N=1,620**
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=160	0.034 [0.037] N=1,494	0.024 [0.040] N=1,357	0.034 [0.042] N=1,175	0.008 [0.044] N=1,006	-0.001 [0.054] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=288	0.031 [0.041] N=1,494	0.028 [0.043] N=1,357	0.016 [0.045] N=1,175	0.013 [0.048] N=1,006	0.004 [0.053] N=806
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=137	0.037 [0.029] N=1,494	0.030 [0.030] N=1,357	0.021 [0.033] N=1,175	0.011 [0.037] N=1,006	-0.022 [0.036] N=806
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=465	**0.081* [0.047] N=1,463**	0.062 [0.051] N=1,326	0.067 [0.052] N=1,147	0.047 [0.056] N=981	0.066 [0.062] N=786
BMI-for-age z score below -2	0.111 P=298	0.029 [0.026] N=2,657	0.025 [0.029] N=2,414	0.022 [0.030] N=2,123	0.010 [0.031] N=1,804	0.012 [0.035] N=1,467
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=448	0.005 [0.056] N=1,166	-0.016 [0.059] N=1,061	0.007 [0.061] N=951	0.017 [0.068] N=800	-0.022 [0.075] N=657
WHO 2007 BMI obesity above +2, 61+ months	0.217 P=250	0.002 [0.049] N=1,166	-0.032 [0.052] N=1,061	-0.018 [0.055] N=951	0.011 [0.060] N=800	0.010 [0.070] N=657
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=124	0.020 [0.034] N=1,166	0.020 [0.036] N=1,061	0.009 [0.037] N=951	0.004 [0.040] N=800	0.030 [0.045] N=657

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.014 [0.225] N=2,371	0.007 [0.250] N=2,118	-0.025 [0.271] N=1,815	-0.272 [0.302] N=1,484	-0.185 [0.396] N=1,137
WHO weight-for-age z score	-0.121 [0.148] N=2,522	-0.083 [0.161] N=2,264	-0.153 [0.186] N=1,937	**-0.360* [0.208] N=1,584**	**-0.454* [0.263] N=1,203**
WHO weight-for-length/height z score, 0-60 months	-0.051 [0.273] N=1,306	-0.041 [0.286] N=1,169	-0.135 [0.349] N=987	-0.104 [0.361] N=818	-0.400 [0.451] N=618
WHO BMI-for-age z score	**-0.323* [0.181] N=2,279**	**-0.389** [0.189] N=2,036**	**-0.385* [0.218] N=1,745**	-0.420 [0.262] N=1,426	**-0.952*** [0.319] N=1,089**
Stunted: HAZ below -2	0.024 [0.045] N=2,371	0.019 [0.050] N=2,118	0.007 [0.053] N=1,815	0.043 [0.060] N=1,484	0.036 [0.085] N=1,137
Underweight: WAZ below -2	-0.020 [0.024] N=2,522	-0.024 [0.027] N=2,264	-0.024 [0.032] N=1,937	-0.015 [0.038] N=1,584	0.002 [0.045] N=1,203
Wasted: WLZ/WHZ below -2, 0-60 months	**0.084** [0.040] N=1,306**	0.070 [0.043] N=1,169	**0.092* [0.048] N=987**	0.054 [0.047] N=818	0.065 [0.057] N=618
Overweight: WLZ/WHZ above +2, 0-60 months	0.058 [0.059] N=1,306	0.052 [0.063] N=1,169	0.040 [0.070] N=987	0.034 [0.080] N=818	-0.019 [0.100] N=618
Obese: WLZ/WHZ above +3, 0-60 months	0.057 [0.044] N=1,306	0.046 [0.047] N=1,169	0.034 [0.056] N=987	0.020 [0.070] N=818	-0.112 [0.081] N=618
CIAF: any stunting, underweight, or wasting, 0-60 months	0.101 [0.066] N=1,281	0.073 [0.075] N=1,144	0.071 [0.079] N=965	0.036 [0.091] N=799	0.059 [0.115] N=604
BMI-for-age z score below -2	**0.083*** [0.026] N=2,279**	**0.083*** [0.031] N=2,036**	**0.081** [0.037] N=1,745**	**0.068** [0.034] N=1,426**	**0.128*** [0.042] N=1,089**
WHO 2007 BMI overweight above +1, 61+ months	-0.053 [0.082] N=976	-0.102 [0.090] N=871	-0.078 [0.098] N=761	-0.083 [0.121] N=610	**-0.287* [0.153] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.056 [0.064] N=976	**-0.134** [0.067] N=871**	**-0.133* [0.073] N=761**	-0.102 [0.085] N=610	-0.178 [0.126] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.077* [0.043] N=976**	**0.085* [0.047] N=871**	0.072 [0.052] N=761	0.080 [0.057] N=610	**0.204*** [0.060] N=467**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.409	-0.214 [0.181] N=2,767	-0.223 [0.193] N=2,514	-0.259 [0.203] N=2,211	**-0.369* [0.218] N=1,880**	-0.410 [0.253] N=1,533
WHO weight-for-age z score	0.092	-0.103 [0.101] N=2,939	-0.085 [0.104] N=2,681	-0.116 [0.113] N=2,354	-0.190 [0.120] N=2,001	-0.196 [0.136] N=1,620
WHO weight-for-length/height z score, 0-60 months	0.365	-0.022 [0.210] N=1,494	-0.012 [0.220] N=1,357	-0.063 [0.246] N=1,175	-0.024 [0.256] N=1,006	-0.065 [0.295] N=806
WHO BMI-for-age z score	0.376	0.060 [0.165] N=2,657	0.062 [0.173] N=2,414	0.106 [0.182] N=2,123	0.145 [0.197] N=1,804	0.124 [0.227] N=1,467
Stunted: HAZ below -2	0.225 P=616	0.049 [0.033] N=2,767	0.048 [0.035] N=2,514	0.048 [0.037] N=2,211	**0.066* [0.039] N=1,880**	**0.075* [0.044] N=1,533**
Underweight: WAZ below -2	0.060 P=172	0.010 [0.017] N=2,939	0.011 [0.018] N=2,681	0.015 [0.020] N=2,354	0.025 [0.022] N=2,001	**0.040* [0.024] N=1,620**
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=160	0.032 [0.037] N=1,494	0.022 [0.040] N=1,357	0.032 [0.042] N=1,175	0.007 [0.044] N=1,006	0.000 [0.054] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=288	0.030 [0.041] N=1,494	0.027 [0.043] N=1,357	0.015 [0.046] N=1,175	0.012 [0.048] N=1,006	0.005 [0.053] N=806
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=137	0.036 [0.029] N=1,494	0.029 [0.030] N=1,357	0.019 [0.033] N=1,175	0.010 [0.038] N=1,006	-0.021 [0.036] N=806
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=465	**0.079* [0.047] N=1,463**	0.060 [0.051] N=1,326	0.064 [0.052] N=1,147	0.047 [0.056] N=981	0.068 [0.062] N=786
BMI-for-age z score below -2	0.111 P=298	0.031 [0.026] N=2,657	0.027 [0.029] N=2,414	0.023 [0.031] N=2,123	0.011 [0.031] N=1,804	0.015 [0.036] N=1,467
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=448	0.011 [0.056] N=1,166	-0.012 [0.059] N=1,061	0.010 [0.062] N=951	0.020 [0.067] N=800	-0.013 [0.074] N=657
WHO 2007 BMI obesity above +2, 61+ months	0.217 P=250	0.010 [0.050] N=1,166	-0.025 [0.053] N=1,061	-0.013 [0.055] N=951	0.015 [0.061] N=800	0.016 [0.070] N=657
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=124	0.022 [0.034] N=1,166	0.023 [0.036] N=1,061	0.009 [0.037] N=951	0.004 [0.040] N=800	0.029 [0.044] N=657

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.011 [0.225] N=2,371	0.009 [0.250] N=2,118	-0.024 [0.270] N=1,815	-0.269 [0.303] N=1,484	-0.168 [0.399] N=1,137
WHO weight-for-age z score	-0.117 [0.149] N=2,522	-0.078 [0.162] N=2,264	-0.148 [0.187] N=1,937	**-0.356* [0.210] N=1,584**	**-0.444* [0.266] N=1,203**
WHO weight-for-length/height z score, 0-60 months	-0.030 [0.273] N=1,306	-0.025 [0.285] N=1,169	-0.128 [0.347] N=987	-0.091 [0.360] N=818	-0.375 [0.458] N=618
WHO BMI-for-age z score	**-0.326* [0.181] N=2,279**	**-0.393** [0.189] N=2,036**	**-0.385* [0.218] N=1,745**	-0.423 [0.262] N=1,426	**-0.966*** [0.318] N=1,089**
Stunted: HAZ below -2	0.024 [0.045] N=2,371	0.019 [0.050] N=2,118	0.008 [0.053] N=1,815	0.043 [0.060] N=1,484	0.036 [0.085] N=1,137
Underweight: WAZ below -2	-0.020 [0.024] N=2,522	-0.025 [0.027] N=2,264	-0.024 [0.033] N=1,937	-0.016 [0.038] N=1,584	0.002 [0.045] N=1,203
Wasted: WLZ/WHZ below -2, 0-60 months	**0.079** [0.040] N=1,306**	0.065 [0.043] N=1,169	**0.087* [0.048] N=987**	0.047 [0.047] N=818	0.058 [0.058] N=618
Overweight: WLZ/WHZ above +2, 0-60 months	0.058 [0.059] N=1,306	0.051 [0.063] N=1,169	0.037 [0.069] N=987	0.031 [0.079] N=818	-0.021 [0.099] N=618
Obese: WLZ/WHZ above +3, 0-60 months	0.056 [0.044] N=1,306	0.044 [0.047] N=1,169	0.029 [0.056] N=987	0.016 [0.070] N=818	-0.115 [0.081] N=618
CIAF: any stunting, underweight, or wasting, 0-60 months	0.095 [0.067] N=1,281	0.066 [0.075] N=1,144	0.062 [0.080] N=965	0.029 [0.092] N=799	0.055 [0.115] N=604
BMI-for-age z score below -2	**0.084*** [0.026] N=2,279**	**0.085*** [0.031] N=2,036**	**0.081** [0.037] N=1,745**	**0.069** [0.034] N=1,426**	**0.131*** [0.041] N=1,089**
WHO 2007 BMI overweight above +1, 61+ months	-0.046 [0.083] N=976	-0.097 [0.091] N=871	-0.075 [0.100] N=761	-0.082 [0.122] N=610	**-0.273* [0.157] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.047 [0.067] N=976	**-0.126* [0.069] N=871**	**-0.128* [0.075] N=761**	-0.101 [0.086] N=610	-0.172 [0.128] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.081* [0.043] N=976**	**0.091* [0.047] N=871**	0.072 [0.052] N=761	0.082 [0.056] N=610	**0.205*** [0.060] N=467**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.409	-0.209 [0.178] N=2,767	-0.211 [0.189] N=2,514	-0.244 [0.198] N=2,211	**-0.355* [0.212] N=1,880**	-0.389 [0.243] N=1,533
WHO weight-for-age z score	0.092	-0.104 [0.099] N=2,939	-0.084 [0.103] N=2,681	-0.107 [0.112] N=2,354	-0.189 [0.118] N=2,001	-0.190 [0.135] N=1,620
WHO weight-for-length/height z score, 0-60 months	0.365	-0.023 [0.209] N=1,494	-0.012 [0.219] N=1,357	-0.048 [0.247] N=1,175	-0.012 [0.257] N=1,006	-0.036 [0.301] N=806
WHO BMI-for-age z score	0.376	0.056 [0.166] N=2,657	0.057 [0.174] N=2,414	0.107 [0.182] N=2,123	0.139 [0.198] N=1,804	0.118 [0.228] N=1,467
Stunted: HAZ below -2	0.225 P=616	0.047 [0.033] N=2,767	0.045 [0.035] N=2,514	0.045 [0.036] N=2,211	0.062 [0.038] N=1,880	0.071 [0.043] N=1,533
Underweight: WAZ below -2	0.060 P=172	0.010 [0.017] N=2,939	0.010 [0.018] N=2,681	0.013 [0.020] N=2,354	0.024 [0.022] N=2,001	**0.040* [0.023] N=1,620**
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=160	0.036 [0.037] N=1,494	0.024 [0.040] N=1,357	0.034 [0.043] N=1,175	0.010 [0.044] N=1,006	0.003 [0.055] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=288	0.032 [0.041] N=1,494	0.028 [0.043] N=1,357	0.020 [0.046] N=1,175	0.017 [0.048] N=1,006	0.012 [0.053] N=806
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=137	0.039 [0.029] N=1,494	0.031 [0.030] N=1,357	0.022 [0.034] N=1,175	0.013 [0.038] N=1,006	-0.016 [0.038] N=806
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=465	**0.087* [0.046] N=1,463**	0.065 [0.051] N=1,326	0.066 [0.052] N=1,147	0.049 [0.055] N=981	0.074 [0.061] N=786
BMI-for-age z score below -2	0.111 P=298	0.030 [0.026] N=2,657	0.026 [0.029] N=2,414	0.021 [0.030] N=2,123	0.010 [0.031] N=1,804	0.014 [0.036] N=1,467
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=448	-0.001 [0.057] N=1,166	-0.022 [0.061] N=1,061	-0.001 [0.062] N=951	0.008 [0.068] N=800	-0.025 [0.074] N=657
WHO 2007 BMI obesity above +2, 61+ months	0.217 P=250	-0.002 [0.050] N=1,166	-0.036 [0.053] N=1,061	-0.025 [0.055] N=951	0.001 [0.060] N=800	0.005 [0.069] N=657
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=124	0.017 [0.034] N=1,166	0.020 [0.035] N=1,061	0.008 [0.037] N=951	0.005 [0.040] N=800	0.028 [0.045] N=657

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.001 [0.226] N=2,371	0.031 [0.250] N=2,118	0.009 [0.271] N=1,815	-0.233 [0.299] N=1,484	-0.096 [0.390] N=1,137
WHO weight-for-age z score	-0.123 [0.148] N=2,522	-0.085 [0.163] N=2,264	-0.142 [0.188] N=1,937	**-0.368* [0.207] N=1,584**	**-0.453* [0.268] N=1,203**
WHO weight-for-length/height z score, 0-60 months	-0.049 [0.274] N=1,306	-0.041 [0.288] N=1,169	-0.129 [0.351] N=987	-0.112 [0.367] N=818	-0.393 [0.468] N=618
WHO BMI-for-age z score	**-0.344* [0.180] N=2,279**	**-0.410** [0.189] N=2,036**	**-0.402* [0.218] N=1,745**	**-0.453* [0.263] N=1,426**	**-1.004*** [0.327] N=1,089**
Stunted: HAZ below -2	0.021 [0.045] N=2,371	0.014 [0.050] N=2,118	0.001 [0.053] N=1,815	0.034 [0.061] N=1,484	0.019 [0.084] N=1,137
Underweight: WAZ below -2	-0.021 [0.024] N=2,522	-0.025 [0.027] N=2,264	-0.026 [0.032] N=1,937	-0.016 [0.037] N=1,584	0.002 [0.044] N=1,203
Wasted: WLZ/WHZ below -2, 0-60 months	**0.083** [0.041] N=1,306**	0.067 [0.044] N=1,169	**0.086* [0.049] N=987**	0.048 [0.048] N=818	0.061 [0.058] N=618
Overweight: WLZ/WHZ above +2, 0-60 months	0.057 [0.059] N=1,306	0.049 [0.064] N=1,169	0.036 [0.071] N=987	0.025 [0.082] N=818	-0.024 [0.105] N=618
Obese: WLZ/WHZ above +3, 0-60 months	0.054 [0.046] N=1,306	0.042 [0.049] N=1,169	0.028 [0.058] N=987	0.012 [0.073] N=818	-0.119 [0.086] N=618
CIAF: any stunting, underweight, or wasting, 0-60 months	0.101 [0.068] N=1,281	0.067 [0.077] N=1,144	0.055 [0.081] N=965	0.021 [0.093] N=799	0.048 [0.114] N=604
BMI-for-age z score below -2	**0.083*** [0.025] N=2,279**	**0.083*** [0.031] N=2,036**	**0.079** [0.036] N=1,745**	**0.066* [0.034] N=1,426**	**0.125*** [0.041] N=1,089**
WHO 2007 BMI overweight above +1, 61+ months	-0.064 [0.084] N=976	-0.112 [0.093] N=871	-0.090 [0.102] N=761	-0.098 [0.125] N=610	**-0.304* [0.159] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.064 [0.067] N=976	**-0.142** [0.070] N=871**	**-0.145* [0.075] N=761**	-0.121 [0.087] N=610	-0.197 [0.131] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.075* [0.043] N=976**	**0.086* [0.046] N=871**	0.072 [0.051] N=761	0.081 [0.056] N=610	**0.199*** [0.061] N=467**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.409	-0.214 [0.177] N=2,767	-0.217 [0.188] N=2,514	-0.257 [0.199] N=2,211	**-0.366* [0.213] N=1,880**	**-0.419* [0.245] N=1,533**
WHO weight-for-age z score	0.092	-0.106 [0.097] N=2,939	-0.086 [0.100] N=2,681	-0.114 [0.108] N=2,354	**-0.188* [0.113] N=2,001**	-0.207 [0.128] N=1,620
WHO weight-for-length/height z score, 0-60 months	0.365	-0.011 [0.209] N=1,494	-0.002 [0.218] N=1,357	-0.041 [0.245] N=1,175	0.010 [0.252] N=1,006	-0.035 [0.293] N=806
WHO BMI-for-age z score	0.376	0.054 [0.164] N=2,657	0.055 [0.172] N=2,414	0.105 [0.181] N=2,123	0.150 [0.196] N=1,804	0.121 [0.226] N=1,467
Stunted: HAZ below -2	0.225 P=616	0.049 [0.033] N=2,767	0.046 [0.035] N=2,514	0.048 [0.036] N=2,211	**0.066* [0.038] N=1,880**	**0.074* [0.043] N=1,533**
Underweight: WAZ below -2	0.060 P=172	0.011 [0.016] N=2,939	0.011 [0.018] N=2,681	0.015 [0.020] N=2,354	0.025 [0.021] N=2,001	**0.041* [0.023] N=1,620**
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=160	0.035 [0.037] N=1,494	0.024 [0.040] N=1,357	0.035 [0.042] N=1,175	0.007 [0.043] N=1,006	-0.003 [0.054] N=806
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=288	0.036 [0.041] N=1,494	0.032 [0.042] N=1,357	0.025 [0.045] N=1,175	0.021 [0.047] N=1,006	0.009 [0.052] N=806
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=137	0.042 [0.029] N=1,494	0.034 [0.030] N=1,357	0.024 [0.033] N=1,175	0.016 [0.037] N=1,006	-0.016 [0.037] N=806
CIAF: any stunting, underweight, or wasting, 0-60 months	0.319 P=465	**0.086* [0.046] N=1,463**	0.065 [0.050] N=1,326	0.070 [0.051] N=1,147	0.051 [0.054] N=981	0.066 [0.060] N=786
BMI-for-age z score below -2	0.111 P=298	0.031 [0.026] N=2,657	0.026 [0.029] N=2,414	0.022 [0.030] N=2,123	0.009 [0.031] N=1,804	0.012 [0.035] N=1,467
WHO 2007 BMI overweight above +1, 61+ months	0.388 P=448	0.003 [0.056] N=1,166	-0.018 [0.060] N=1,061	0.006 [0.061] N=951	0.016 [0.067] N=800	-0.017 [0.074] N=657
WHO 2007 BMI obesity above +2, 61+ months	0.217 P=250	-0.000 [0.049] N=1,166	-0.035 [0.052] N=1,061	-0.023 [0.054] N=951	0.004 [0.059] N=800	0.001 [0.069] N=657
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=124	0.020 [0.034] N=1,166	0.022 [0.035] N=1,061	0.008 [0.037] N=951	0.003 [0.039] N=800	0.028 [0.044] N=657

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.008 [0.222] N=2,371	0.024 [0.245] N=2,118	0.001 [0.268] N=1,815	-0.232 [0.295] N=1,484	-0.135 [0.380] N=1,137
WHO weight-for-age z score	-0.111 [0.145] N=2,522	-0.068 [0.158] N=2,264	-0.125 [0.182] N=1,937	-0.323 [0.198] N=1,584	**-0.428* [0.250] N=1,203**
WHO weight-for-length/height z score, 0-60 months	-0.025 [0.273] N=1,306	-0.025 [0.285] N=1,169	-0.120 [0.346] N=987	-0.068 [0.356] N=818	-0.335 [0.458] N=618
WHO BMI-for-age z score	**-0.330* [0.180] N=2,279**	**-0.398** [0.188] N=2,036**	**-0.390* [0.216] N=1,745**	-0.421 [0.258] N=1,426	**-0.975*** [0.323] N=1,089**
Stunted: HAZ below -2	0.025 [0.045] N=2,371	0.017 [0.050] N=2,118	0.003 [0.053] N=1,815	0.036 [0.060] N=1,484	0.029 [0.083] N=1,137
Underweight: WAZ below -2	-0.020 [0.023] N=2,522	-0.025 [0.027] N=2,264	-0.026 [0.032] N=1,937	-0.019 [0.036] N=1,584	0.001 [0.043] N=1,203
Wasted: WLZ/WHZ below -2, 0-60 months	**0.084** [0.040] N=1,306**	0.070 [0.043] N=1,169	**0.093* [0.048] N=987**	0.049 [0.046] N=818	0.059 [0.056] N=618
Overweight: WLZ/WHZ above +2, 0-60 months	0.063 [0.059] N=1,306	0.054 [0.063] N=1,169	0.044 [0.070] N=987	0.037 [0.080] N=818	-0.013 [0.099] N=618
Obese: WLZ/WHZ above +3, 0-60 months	0.061 [0.045] N=1,306	0.047 [0.047] N=1,169	0.030 [0.057] N=987	0.016 [0.071] N=818	-0.112 [0.084] N=618
CIAF: any stunting, underweight, or wasting, 0-60 months	0.107 [0.066] N=1,281	0.072 [0.075] N=1,144	0.063 [0.079] N=965	0.027 [0.087] N=799	0.068 [0.106] N=604
BMI-for-age z score below -2	**0.086*** [0.026] N=2,279**	**0.085*** [0.032] N=2,036**	**0.082** [0.037] N=1,745**	**0.068** [0.034] N=1,426**	**0.131*** [0.041] N=1,089**
WHO 2007 BMI overweight above +1, 61+ months	-0.062 [0.083] N=976	-0.108 [0.091] N=871	-0.087 [0.099] N=761	-0.094 [0.121] N=610	**-0.296* [0.155] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.059 [0.065] N=976	**-0.135* [0.069] N=871**	**-0.138* [0.073] N=761**	-0.109 [0.083] N=610	-0.189 [0.125] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.076* [0.043] N=976**	**0.087* [0.047] N=871**	0.069 [0.051] N=761	0.076 [0.056] N=610	**0.189*** [0.061] N=467**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.417	-0.176 [0.181] N=2,593	-0.153 [0.195] N=2,357	-0.193 [0.204] N=2,077	-0.288 [0.222] N=1,761	-0.390 [0.255] N=1,439
WHO weight-for-age z score	0.097	-0.075 [0.110] N=2,747	-0.046 [0.114] N=2,506	-0.084 [0.124] N=2,205	-0.152 [0.133] N=1,871	-0.198 [0.151] N=1,519
WHO weight-for-length/height z score, 0-60 months	0.370	0.053 [0.220] N=1,391	0.078 [0.230] N=1,264	-0.013 [0.254] N=1,095	0.009 [0.261] N=937	-0.046 [0.296] N=754
WHO BMI-for-age z score	0.390	0.047 [0.165] N=2,489	0.049 [0.174] N=2,262	0.065 [0.180] N=1,993	0.097 [0.193] N=1,690	0.066 [0.221] N=1,378
Stunted: HAZ below -2	0.228 P=582	0.047 [0.033] N=2,593	0.043 [0.035] N=2,357	0.041 [0.036] N=2,077	0.057 [0.037] N=1,761	**0.074* [0.041] N=1,439**
Underweight: WAZ below -2	0.062 P=165	0.007 [0.019] N=2,747	0.007 [0.020] N=2,506	0.008 [0.022] N=2,205	0.024 [0.024] N=1,871	**0.047* [0.027] N=1,519**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=150	0.033 [0.038] N=1,391	0.018 [0.040] N=1,264	0.034 [0.042] N=1,095	0.013 [0.044] N=937	0.006 [0.054] N=754
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=270	0.029 [0.042] N=1,391	0.028 [0.044] N=1,264	0.017 [0.046] N=1,095	0.011 [0.048] N=937	0.005 [0.053] N=754
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=131	0.041 [0.030] N=1,391	0.035 [0.030] N=1,264	0.026 [0.033] N=1,095	0.022 [0.039] N=937	-0.001 [0.038] N=754
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=437	**0.085* [0.049] N=1,363**	0.061 [0.054] N=1,236	0.068 [0.055] N=1,070	0.051 [0.059] N=914	0.084 [0.067] N=735
BMI-for-age z score below -2	0.110 P=274	0.035 [0.026] N=2,489	0.031 [0.028] N=2,262	0.028 [0.029] N=1,993	0.020 [0.028] N=1,690	0.024 [0.032] N=1,378
WHO 2007 BMI overweight above +1, 61+ months	0.393 P=428	-0.002 [0.057] N=1,104	-0.028 [0.060] N=1,005	0.004 [0.063] N=903	0.021 [0.068] N=757	-0.009 [0.073] N=622
WHO 2007 BMI obesity above +2, 61+ months	0.222 P=242	-0.008 [0.050] N=1,104	-0.050 [0.053] N=1,005	-0.039 [0.055] N=903	-0.014 [0.060] N=757	-0.004 [0.070] N=622
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=116	0.026 [0.035] N=1,104	0.027 [0.037] N=1,005	0.013 [0.037] N=903	0.002 [0.040] N=757	0.030 [0.044] N=622

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.040 [0.239] N=2,220	0.114 [0.266] N=1,984	0.055 [0.296] N=1,704	-0.174 [0.335] N=1,388	-0.175 [0.445] N=1,066
WHO weight-for-age z score	-0.091 [0.158] N=2,357	-0.033 [0.171] N=2,116	-0.124 [0.198] N=1,815	-0.316 [0.222] N=1,481	-0.467 [0.281] N=1,129
WHO weight-for-length/height z score, 0-60 months	0.036 [0.301] N=1,213	0.058 [0.318] N=1,086	-0.132 [0.386] N=917	-0.137 [0.410] N=759	-0.440 [0.552] N=576
WHO BMI-for-age z score	**-0.328* [0.190] N=2,133**	**-0.394** [0.198] N=1,906**	**-0.414* [0.228] N=1,637**	-0.442 [0.275] N=1,334	**-1.017*** [0.336] N=1,022**
Stunted: HAZ below -2	0.018 [0.047] N=2,220	0.007 [0.053] N=1,984	-0.007 [0.056] N=1,704	0.025 [0.061] N=1,388	0.022 [0.081] N=1,066
Underweight: WAZ below -2	-0.029 [0.026] N=2,357	-0.036 [0.029] N=2,116	-0.038 [0.034] N=1,815	-0.022 [0.041] N=1,481	0.004 [0.048] N=1,129
Wasted: WLZ/WHZ below -2, 0-60 months	**0.081* [0.042] N=1,213**	0.059 [0.045] N=1,086	**0.089* [0.050] N=917**	0.056 [0.053] N=759	0.060 [0.067] N=576
Overweight: WLZ/WHZ above +2, 0-60 months	0.050 [0.062] N=1,213	0.043 [0.066] N=1,086	0.024 [0.073] N=917	0.009 [0.086] N=759	-0.040 [0.114] N=576
Obese: WLZ/WHZ above +3, 0-60 months	0.046 [0.048] N=1,213	0.034 [0.050] N=1,086	0.014 [0.059] N=917	0.004 [0.075] N=759	-0.124 [0.090] N=576
CIAF: any stunting, underweight, or wasting, 0-60 months	0.096 [0.074] N=1,190	0.056 [0.084] N=1,063	0.049 [0.090] N=897	0.009 [0.104] N=741	0.051 [0.128] N=562
BMI-for-age z score below -2	**0.092*** [0.027] N=2,133**	**0.093*** [0.033] N=1,906**	**0.091** [0.038] N=1,637**	**0.088** [0.034] N=1,334**	**0.157*** [0.042] N=1,022**
WHO 2007 BMI overweight above +1, 61+ months	-0.057 [0.085] N=925	-0.113 [0.092] N=826	-0.071 [0.103] N=724	-0.050 [0.125] N=578	-0.218 [0.159] N=443
WHO 2007 BMI obesity above +2, 61+ months	-0.056 [0.070] N=925	**-0.142* [0.073] N=826**	**-0.142* [0.078] N=724**	-0.105 [0.090] N=578	-0.159 [0.131] N=443
WHO 2007 BMI thinness below -2, 61+ months	**0.088* [0.046] N=925**	**0.100** [0.050] N=826**	0.079 [0.054] N=724	0.074 [0.058] N=578	**0.197*** [0.064] N=443**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.417	-0.173 [0.182] N=2,593	-0.151 [0.195] N=2,357	-0.187 [0.205] N=2,077	-0.277 [0.223] N=1,761	-0.372 [0.257] N=1,439
WHO weight-for-age z score	0.097	-0.078 [0.110] N=2,747	-0.048 [0.115] N=2,506	-0.090 [0.125] N=2,205	-0.154 [0.134] N=1,871	-0.196 [0.153] N=1,519
WHO weight-for-length/height z score, 0-60 months	0.370	0.061 [0.220] N=1,391	0.085 [0.230] N=1,264	-0.002 [0.254] N=1,095	0.026 [0.263] N=937	-0.033 [0.307] N=754
WHO BMI-for-age z score	0.390	0.047 [0.165] N=2,489	0.050 [0.174] N=2,262	0.066 [0.180] N=1,993	0.101 [0.194] N=1,690	0.068 [0.223] N=1,378
Stunted: HAZ below -2	0.228 P=582	0.045 [0.033] N=2,593	0.041 [0.035] N=2,357	0.039 [0.036] N=2,077	0.055 [0.037] N=1,761	**0.073* [0.042] N=1,439**
Underweight: WAZ below -2	0.062 P=165	0.008 [0.019] N=2,747	0.007 [0.020] N=2,506	0.010 [0.022] N=2,205	0.025 [0.024] N=1,871	**0.047* [0.027] N=1,519**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=150	0.030 [0.038] N=1,391	0.015 [0.040] N=1,264	0.029 [0.042] N=1,095	0.007 [0.044] N=937	-0.002 [0.055] N=754
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=270	0.030 [0.042] N=1,391	0.028 [0.043] N=1,264	0.017 [0.046] N=1,095	0.012 [0.047] N=937	0.006 [0.053] N=754
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=131	0.041 [0.029] N=1,391	0.035 [0.030] N=1,264	0.025 [0.033] N=1,095	0.021 [0.038] N=937	-0.002 [0.038] N=754
CIAF: any stunting, underweight, or wasting, 0-60 months	0.322 P=437	0.080 [0.048] N=1,363	0.054 [0.053] N=1,236	0.059 [0.054] N=1,070	0.042 [0.058] N=914	0.071 [0.064] N=735
BMI-for-age z score below -2	0.110 P=274	0.034 [0.026] N=2,489	0.030 [0.028] N=2,262	0.027 [0.029] N=1,993	0.018 [0.029] N=1,690	0.023 [0.032] N=1,378
WHO 2007 BMI overweight above +1, 61+ months	0.393 P=428	-0.008 [0.057] N=1,104	-0.034 [0.060] N=1,005	-0.002 [0.063] N=903	0.013 [0.069] N=757	-0.016 [0.073] N=622
WHO 2007 BMI obesity above +2, 61+ months	0.222 P=242	-0.010 [0.050] N=1,104	-0.052 [0.053] N=1,005	-0.042 [0.055] N=903	-0.017 [0.060] N=757	-0.007 [0.069] N=622
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=116	0.025 [0.034] N=1,104	0.027 [0.036] N=1,005	0.008 [0.036] N=903	-0.003 [0.039] N=757	0.023 [0.043] N=622

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.027 [0.238] N=2,220	0.101 [0.265] N=1,984	0.051 [0.296] N=1,704	-0.194 [0.338] N=1,388	-0.183 [0.448] N=1,066
WHO weight-for-age z score	-0.092 [0.157] N=2,357	-0.034 [0.172] N=2,116	-0.128 [0.199] N=1,815	-0.315 [0.223] N=1,481	-0.472 [0.284] N=1,129
WHO weight-for-length/height z score, 0-60 months	0.039 [0.299] N=1,213	0.055 [0.315] N=1,086	-0.125 [0.383] N=917	-0.101 [0.404] N=759	-0.452 [0.534] N=576
WHO BMI-for-age z score	**-0.323* [0.190] N=2,133**	**-0.392* [0.199] N=1,906**	**-0.412* [0.227] N=1,637**	-0.427 [0.274] N=1,334	**-1.014*** [0.333] N=1,022**
Stunted: HAZ below -2	0.019 [0.047] N=2,220	0.009 [0.053] N=1,984	-0.006 [0.056] N=1,704	0.029 [0.062] N=1,388	0.031 [0.082] N=1,066
Underweight: WAZ below -2	-0.028 [0.025] N=2,357	-0.035 [0.029] N=2,116	-0.035 [0.033] N=1,815	-0.018 [0.040] N=1,481	0.012 [0.047] N=1,129
Wasted: WLZ/WHZ below -2, 0-60 months	**0.079* [0.041] N=1,213**	0.058 [0.045] N=1,086	**0.085* [0.050] N=917**	0.050 [0.052] N=759	0.058 [0.066] N=576
Overweight: WLZ/WHZ above +2, 0-60 months	0.050 [0.061] N=1,213	0.041 [0.066] N=1,086	0.024 [0.073] N=917	0.014 [0.085] N=759	-0.041 [0.111] N=576
Obese: WLZ/WHZ above +3, 0-60 months	0.047 [0.047] N=1,213	0.034 [0.049] N=1,086	0.014 [0.058] N=917	0.007 [0.075] N=759	-0.124 [0.088] N=576
CIAF: any stunting, underweight, or wasting, 0-60 months	0.094 [0.073] N=1,190	0.053 [0.084] N=1,063	0.045 [0.090] N=897	0.011 [0.104] N=741	0.049 [0.127] N=562
BMI-for-age z score below -2	**0.092*** [0.027] N=2,133**	**0.093*** [0.033] N=1,906**	**0.091** [0.038] N=1,637**	**0.086** [0.035] N=1,334**	**0.158*** [0.042] N=1,022**
WHO 2007 BMI overweight above +1, 61+ months	-0.059 [0.085] N=925	-0.112 [0.093] N=826	-0.071 [0.103] N=724	-0.051 [0.127] N=578	-0.217 [0.160] N=443
WHO 2007 BMI obesity above +2, 61+ months	-0.056 [0.070] N=925	**-0.143* [0.073] N=826**	**-0.142* [0.078] N=724**	-0.107 [0.089] N=578	-0.162 [0.131] N=443
WHO 2007 BMI thinness below -2, 61+ months	**0.090** [0.045] N=925**	**0.103** [0.049] N=826**	0.077 [0.053] N=724	0.072 [0.057] N=578	**0.193*** [0.062] N=443**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 83 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.