Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Ages 0-6 Using 12 November 2022 Child Age

The earliest-born eligible child in each household is selected within completed ages 0 to 83 months on 12 November 2022. Ages outside this range are excluded. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications apply through month 60. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.267	0.069 [0.146] N=2,830	0.052 [0.153] N=2,620	0.029 [0.169] N=2,347	0.090 [0.174] N=2,016	0.111 [0.200] N=1,665
WHO weight-for-age z score	0.204	0.055 [0.088] N=3,006	0.103 [0.096] N=2,776	0.155 [0.099] N=2,485	0.064 [0.105] N=2,131	0.085 [0.120] N=1,759
WHO weight-for-length/height z score, 0-60 months	0.337	0.209 [0.187] N=1,601	0.194 [0.193] N=1,476	0.098 [0.206] N=1,320	-0.044 [0.233] N=1,123	0.112 [0.247] N=916
WHO BMI-for-age z score	0.349	0.101 [0.152] N=2,708	0.115 [0.158] N=2,509	0.125 [0.172] N=2,241	-0.078 [0.190] N=1,926	-0.020 [0.209] N=1,591
Stunted: HAZ below -2	0.199 P=544	-0.012 [0.029] N=2,830	-0.015 [0.030] N=2,620	-0.008 [0.033] N=2,347	-0.033 [0.034] N=2,016	-0.020 [0.039] N=1,665
Underweight: WAZ below -2	0.054 P=156	-0.014 [0.015] N=3,006	-0.020 [0.016] N=2,776	**-0.030* [0.017] N=2,485**	-0.021 [0.019] N=2,131	-0.025 [0.021] N=1,759
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=179	-0.023 [0.032] N=1,601	-0.017 [0.033] N=1,476	0.002 [0.036] N=1,320	0.011 [0.042] N=1,123	-0.008 [0.045] N=916
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=299	**0.079* [0.042] N=1,601**	**0.076* [0.044] N=1,476**	0.072 [0.048] N=1,320	0.035 [0.054] N=1,123	0.053 [0.060] N=916
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=128	0.027 [0.030] N=1,601	0.031 [0.031] N=1,476	0.033 [0.034] N=1,320	0.019 [0.038] N=1,123	0.025 [0.045] N=916
CIAF: any stunting, underweight, or wasting, 0-60 months	0.275 P=425	-0.030 [0.044] N=1,575	-0.043 [0.045] N=1,453	-0.005 [0.049] N=1,300	-0.020 [0.051] N=1,104	0.003 [0.057] N=903
BMI-for-age z score below -2	0.113 P=315	0.001 [0.027] N=2,708	0.001 [0.026] N=2,509	0.003 [0.028] N=2,241	0.022 [0.031] N=1,926	0.027 [0.033] N=1,591
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=415	0.043 [0.050] N=1,101	0.043 [0.053] N=1,027	0.088 [0.058] N=920	0.055 [0.065] N=804	0.025 [0.072] N=670
WHO 2007 BMI obesity above +2, 61+ months	0.210 P=235	0.014 [0.044] N=1,101	0.006 [0.046] N=1,027	0.025 [0.048] N=920	-0.010 [0.050] N=804	-0.074 [0.056] N=670
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=121	0.022 [0.043] N=1,101	0.016 [0.044] N=1,027	-0.016 [0.049] N=920	0.004 [0.053] N=804	0.017 [0.060] N=670

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.052 [0.204] N=2,378	-0.106 [0.216] N=2,168	-0.216 [0.251] N=1,895	-0.107 [0.277] N=1,564	-0.215 [0.371] N=1,213
WHO weight-for-age z score	-0.019 [0.115] N=2,531	0.058 [0.131] N=2,301	0.140 [0.140] N=2,010	-0.039 [0.160] N=1,656	-0.016 [0.215] N=1,284
WHO weight-for-length/height z score, 0-60 months	0.159 [0.264] N=1,359	0.109 [0.279] N=1,234	-0.107 [0.317] N=1,078	-0.465 [0.386] N=881	-0.327 [0.477] N=674
WHO BMI-for-age z score	0.273 [0.195] N=2,279	0.330 [0.206] N=2,080	**0.417* [0.239] N=1,812**	0.081 [0.287] N=1,497	0.352 [0.362] N=1,162
Stunted: HAZ below -2	0.033 [0.041] N=2,378	0.031 [0.044] N=2,168	0.060 [0.053] N=1,895	0.018 [0.060] N=1,564	0.083 [0.076] N=1,213
Underweight: WAZ below -2	-0.002 [0.022] N=2,531	-0.012 [0.023] N=2,301	-0.029 [0.027] N=2,010	-0.008 [0.032] N=1,656	-0.004 [0.043] N=1,284
Wasted: WLZ/WHZ below -2, 0-60 months	-0.009 [0.040] N=1,359	0.004 [0.040] N=1,234	0.042 [0.045] N=1,078	0.058 [0.061] N=881	0.024 [0.064] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.067 [0.054] N=1,359	0.058 [0.058] N=1,234	0.051 [0.067] N=1,078	-0.041 [0.083] N=881	-0.045 [0.113] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.036] N=1,359	0.045 [0.039] N=1,234	0.053 [0.045] N=1,078	0.023 [0.057] N=881	0.035 [0.083] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.010 [0.064] N=1,336	-0.015 [0.068] N=1,214	0.067 [0.079] N=1,061	0.031 [0.093] N=865	0.111 [0.108] N=664
BMI-for-age z score below -2	-0.024 [0.032] N=2,279	-0.031 [0.032] N=2,080	-0.038 [0.036] N=1,812	-0.018 [0.040] N=1,497	-0.032 [0.050] N=1,162
WHO 2007 BMI overweight above +1, 61+ months	0.106 [0.073] N=915	0.123 [0.082] N=841	**0.232** [0.093] N=734**	**0.204* [0.114] N=618**	0.218 [0.148] N=484
WHO 2007 BMI obesity above +2, 61+ months	**0.118* [0.064] N=915**	**0.120* [0.070] N=841**	**0.191** [0.078] N=734**	**0.166* [0.095] N=618**	0.107 [0.120] N=484
WHO 2007 BMI thinness below -2, 61+ months	-0.002 [0.046] N=915	-0.021 [0.047] N=841	-0.090 [0.056] N=734	-0.071 [0.062] N=618	-0.094 [0.085] N=484

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.267	0.076 [0.145] N=2,830	0.059 [0.152] N=2,620	0.041 [0.168] N=2,347	0.103 [0.172] N=2,016	0.119 [0.199] N=1,665
WHO weight-for-age z score	0.204	0.060 [0.088] N=3,006	0.108 [0.096] N=2,776	**0.164* [0.098] N=2,485**	0.072 [0.105] N=2,131	0.089 [0.120] N=1,759
WHO weight-for-length/height z score, 0-60 months	0.337	0.203 [0.188] N=1,601	0.186 [0.195] N=1,476	0.088 [0.208] N=1,320	-0.048 [0.235] N=1,123	0.105 [0.249] N=916
WHO BMI-for-age z score	0.349	0.100 [0.153] N=2,708	0.112 [0.158] N=2,509	0.122 [0.174] N=2,241	-0.080 [0.192] N=1,926	-0.023 [0.209] N=1,591
Stunted: HAZ below -2	0.199 P=544	-0.012 [0.029] N=2,830	-0.015 [0.030] N=2,620	-0.007 [0.033] N=2,347	-0.033 [0.034] N=2,016	-0.020 [0.039] N=1,665
Underweight: WAZ below -2	0.054 P=156	-0.014 [0.015] N=3,006	-0.020 [0.016] N=2,776	**-0.030* [0.017] N=2,485**	-0.021 [0.019] N=2,131	-0.025 [0.022] N=1,759
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=179	-0.021 [0.032] N=1,601	-0.014 [0.033] N=1,476	0.006 [0.036] N=1,320	0.013 [0.043] N=1,123	-0.005 [0.046] N=916
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=299	**0.079* [0.042] N=1,601**	**0.077* [0.044] N=1,476**	0.074 [0.048] N=1,320	0.037 [0.054] N=1,123	0.054 [0.060] N=916
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=128	0.027 [0.030] N=1,601	0.032 [0.031] N=1,476	0.033 [0.034] N=1,320	0.019 [0.038] N=1,123	0.025 [0.045] N=916
CIAF: any stunting, underweight, or wasting, 0-60 months	0.275 P=425	-0.029 [0.044] N=1,575	-0.041 [0.045] N=1,453	-0.001 [0.048] N=1,300	-0.018 [0.051] N=1,104	0.006 [0.057] N=903
BMI-for-age z score below -2	0.113 P=315	0.002 [0.027] N=2,708	0.002 [0.026] N=2,509	0.006 [0.028] N=2,241	0.024 [0.031] N=1,926	0.028 [0.033] N=1,591
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=415	0.042 [0.050] N=1,101	0.042 [0.053] N=1,027	0.088 [0.058] N=920	0.055 [0.064] N=804	0.024 [0.072] N=670
WHO 2007 BMI obesity above +2, 61+ months	0.210 P=235	0.016 [0.044] N=1,101	0.009 [0.045] N=1,027	0.029 [0.048] N=920	-0.008 [0.051] N=804	-0.073 [0.056] N=670
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=121	0.022 [0.043] N=1,101	0.016 [0.044] N=1,027	-0.016 [0.050] N=920	0.004 [0.053] N=804	0.018 [0.060] N=670

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.046 [0.203] N=2,378	-0.099 [0.216] N=2,168	-0.198 [0.251] N=1,895	-0.093 [0.275] N=1,564	-0.220 [0.371] N=1,213
WHO weight-for-age z score	-0.015 [0.114] N=2,531	0.061 [0.130] N=2,301	0.150 [0.139] N=2,010	-0.032 [0.157] N=1,656	-0.019 [0.213] N=1,284
WHO weight-for-length/height z score, 0-60 months	0.147 [0.266] N=1,359	0.091 [0.282] N=1,234	-0.134 [0.320] N=1,078	-0.485 [0.389] N=881	-0.374 [0.487] N=674
WHO BMI-for-age z score	0.271 [0.195] N=2,279	0.325 [0.206] N=2,080	**0.410* [0.240] N=1,812**	0.077 [0.288] N=1,497	0.349 [0.363] N=1,162
Stunted: HAZ below -2	0.033 [0.041] N=2,378	0.031 [0.044] N=2,168	0.060 [0.053] N=1,895	0.018 [0.060] N=1,564	0.083 [0.076] N=1,213
Underweight: WAZ below -2	-0.002 [0.022] N=2,531	-0.012 [0.023] N=2,301	-0.029 [0.027] N=2,010	-0.008 [0.032] N=1,656	-0.004 [0.043] N=1,284
Wasted: WLZ/WHZ below -2, 0-60 months	-0.007 [0.040] N=1,359	0.008 [0.040] N=1,234	0.048 [0.045] N=1,078	0.063 [0.062] N=881	0.034 [0.066] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.066 [0.054] N=1,359	0.058 [0.059] N=1,234	0.052 [0.067] N=1,078	-0.039 [0.083] N=881	-0.041 [0.113] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.036] N=1,359	0.045 [0.038] N=1,234	0.053 [0.045] N=1,078	0.023 [0.057] N=881	0.034 [0.083] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.011 [0.064] N=1,336	-0.012 [0.068] N=1,214	0.073 [0.078] N=1,061	0.035 [0.093] N=865	0.117 [0.107] N=664
BMI-for-age z score below -2	-0.024 [0.032] N=2,279	-0.030 [0.032] N=2,080	-0.035 [0.036] N=1,812	-0.016 [0.040] N=1,497	-0.031 [0.050] N=1,162
WHO 2007 BMI overweight above +1, 61+ months	0.106 [0.073] N=915	0.122 [0.082] N=841	**0.232** [0.093] N=734**	**0.204* [0.114] N=618**	0.222 [0.148] N=484
WHO 2007 BMI obesity above +2, 61+ months	**0.119* [0.064] N=915**	**0.121* [0.070] N=841**	**0.195** [0.078] N=734**	**0.166* [0.095] N=618**	0.099 [0.120] N=484
WHO 2007 BMI thinness below -2, 61+ months	-0.002 [0.046] N=915	-0.021 [0.047] N=841	-0.090 [0.056] N=734	-0.071 [0.062] N=618	-0.094 [0.085] N=484

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.267	0.041 [0.148] N=2,830	0.026 [0.155] N=2,620	0.011 [0.173] N=2,347	0.076 [0.177] N=2,016	0.074 [0.206] N=1,665
WHO weight-for-age z score	0.204	0.022 [0.086] N=3,006	0.072 [0.094] N=2,776	0.123 [0.097] N=2,485	0.037 [0.103] N=2,131	0.050 [0.118] N=1,759
WHO weight-for-length/height z score, 0-60 months	0.337	0.174 [0.189] N=1,601	0.166 [0.195] N=1,476	0.074 [0.207] N=1,320	-0.056 [0.234] N=1,123	0.099 [0.250] N=916
WHO BMI-for-age z score	0.349	0.091 [0.154] N=2,708	0.105 [0.160] N=2,509	0.115 [0.174] N=2,241	-0.084 [0.191] N=1,926	-0.026 [0.212] N=1,591
Stunted: HAZ below -2	0.199 P=544	-0.004 [0.029] N=2,830	-0.007 [0.030] N=2,620	-0.000 [0.034] N=2,347	-0.026 [0.034] N=2,016	-0.009 [0.040] N=1,665
Underweight: WAZ below -2	0.054 P=156	-0.008 [0.015] N=3,006	-0.013 [0.015] N=2,776	-0.022 [0.017] N=2,485	-0.014 [0.018] N=2,131	-0.016 [0.021] N=1,759
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=179	-0.012 [0.032] N=1,601	-0.004 [0.033] N=1,476	0.013 [0.035] N=1,320	0.019 [0.042] N=1,123	0.001 [0.045] N=916
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=299	**0.078* [0.042] N=1,601**	**0.078* [0.044] N=1,476**	0.074 [0.048] N=1,320	0.036 [0.054] N=1,123	0.051 [0.060] N=916
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=128	0.028 [0.030] N=1,601	0.034 [0.031] N=1,476	0.034 [0.033] N=1,320	0.019 [0.038] N=1,123	0.024 [0.045] N=916
CIAF: any stunting, underweight, or wasting, 0-60 months	0.275 P=425	-0.007 [0.042] N=1,575	-0.016 [0.043] N=1,453	0.017 [0.046] N=1,300	-0.003 [0.049] N=1,104	0.029 [0.055] N=903
BMI-for-age z score below -2	0.113 P=315	0.007 [0.026] N=2,708	0.007 [0.026] N=2,509	0.010 [0.029] N=2,241	0.028 [0.031] N=1,926	0.034 [0.033] N=1,591
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=415	0.042 [0.051] N=1,101	0.042 [0.054] N=1,027	0.090 [0.059] N=920	0.059 [0.065] N=804	0.024 [0.073] N=670
WHO 2007 BMI obesity above +2, 61+ months	0.210 P=235	0.016 [0.044] N=1,101	0.006 [0.046] N=1,027	0.029 [0.049] N=920	-0.009 [0.051] N=804	-0.072 [0.058] N=670
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=121	0.022 [0.043] N=1,101	0.015 [0.044] N=1,027	-0.015 [0.050] N=920	0.006 [0.053] N=804	0.018 [0.060] N=670

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.084 [0.202] N=2,378	-0.135 [0.213] N=2,168	-0.237 [0.250] N=1,895	-0.132 [0.275] N=1,564	-0.307 [0.371] N=1,213
WHO weight-for-age z score	-0.054 [0.111] N=2,531	0.027 [0.127] N=2,301	0.106 [0.134] N=2,010	-0.069 [0.152] N=1,656	-0.068 [0.208] N=1,284
WHO weight-for-length/height z score, 0-60 months	0.113 [0.266] N=1,359	0.073 [0.284] N=1,234	-0.147 [0.318] N=1,078	-0.505 [0.387] N=881	-0.387 [0.486] N=674
WHO BMI-for-age z score	0.265 [0.196] N=2,279	0.321 [0.207] N=2,080	**0.411* [0.239] N=1,812**	0.086 [0.287] N=1,497	0.374 [0.365] N=1,162
Stunted: HAZ below -2	0.040 [0.040] N=2,378	0.037 [0.043] N=2,168	0.067 [0.052] N=1,895	0.022 [0.058] N=1,564	0.097 [0.074] N=1,213
Underweight: WAZ below -2	0.003 [0.021] N=2,531	-0.006 [0.023] N=2,301	-0.023 [0.025] N=2,010	-0.004 [0.031] N=1,656	0.005 [0.041] N=1,284
Wasted: WLZ/WHZ below -2, 0-60 months	0.004 [0.040] N=1,359	0.019 [0.041] N=1,234	0.054 [0.045] N=1,078	0.069 [0.062] N=881	0.041 [0.068] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.066 [0.054] N=1,359	0.061 [0.059] N=1,234	0.053 [0.066] N=1,078	-0.039 [0.082] N=881	-0.034 [0.112] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.040 [0.036] N=1,359	0.047 [0.038] N=1,234	0.053 [0.045] N=1,078	0.021 [0.056] N=881	0.026 [0.083] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.033 [0.062] N=1,336	0.011 [0.066] N=1,214	0.083 [0.076] N=1,061	0.034 [0.089] N=865	0.131 [0.106] N=664
BMI-for-age z score below -2	-0.019 [0.032] N=2,279	-0.026 [0.032] N=2,080	-0.032 [0.036] N=1,812	-0.014 [0.041] N=1,497	-0.027 [0.051] N=1,162
WHO 2007 BMI overweight above +1, 61+ months	0.107 [0.073] N=915	0.124 [0.082] N=841	**0.239** [0.092] N=734**	**0.212* [0.114] N=618**	0.237 [0.146] N=484
WHO 2007 BMI obesity above +2, 61+ months	**0.121* [0.064] N=915**	**0.120* [0.071] N=841**	**0.199** [0.078] N=734**	**0.166* [0.094] N=618**	0.109 [0.122] N=484
WHO 2007 BMI thinness below -2, 61+ months	-0.003 [0.045] N=915	-0.022 [0.047] N=841	-0.090 [0.055] N=734	-0.068 [0.061] N=618	-0.097 [0.086] N=484

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.267	0.032 [0.147] N=2,830	0.020 [0.154] N=2,620	0.004 [0.172] N=2,347	0.058 [0.176] N=2,016	0.046 [0.205] N=1,665
WHO weight-for-age z score	0.204	0.031 [0.086] N=3,006	0.077 [0.094] N=2,776	0.127 [0.098] N=2,485	0.036 [0.105] N=2,131	0.041 [0.119] N=1,759
WHO weight-for-length/height z score, 0-60 months	0.337	0.209 [0.188] N=1,601	0.195 [0.195] N=1,476	0.099 [0.208] N=1,320	-0.036 [0.234] N=1,123	0.139 [0.250] N=916
WHO BMI-for-age z score	0.349	0.109 [0.154] N=2,708	0.120 [0.159] N=2,509	0.128 [0.175] N=2,241	-0.069 [0.194] N=1,926	-0.005 [0.213] N=1,591
Stunted: HAZ below -2	0.199 P=544	-0.001 [0.029] N=2,830	-0.005 [0.030] N=2,620	0.002 [0.034] N=2,347	-0.021 [0.034] N=2,016	-0.004 [0.040] N=1,665
Underweight: WAZ below -2	0.054 P=156	-0.009 [0.015] N=3,006	-0.014 [0.016] N=2,776	-0.023 [0.017] N=2,485	-0.014 [0.019] N=2,131	-0.017 [0.021] N=1,759
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=179	-0.016 [0.032] N=1,601	-0.010 [0.033] N=1,476	0.008 [0.036] N=1,320	0.016 [0.042] N=1,123	-0.005 [0.046] N=916
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=299	**0.081* [0.041] N=1,601**	**0.078* [0.043] N=1,476**	0.074 [0.047] N=1,320	0.038 [0.053] N=1,123	0.053 [0.059] N=916
Obese: WLZ/WHZ above +3, 0-60 months	0.081 P=128	0.032 [0.030] N=1,601	0.036 [0.031] N=1,476	0.037 [0.034] N=1,320	0.024 [0.039] N=1,123	0.033 [0.046] N=916
CIAF: any stunting, underweight, or wasting, 0-60 months	0.275 P=425	-0.008 [0.042] N=1,575	-0.020 [0.043] N=1,453	0.016 [0.047] N=1,300	0.003 [0.049] N=1,104	0.032 [0.055] N=903
BMI-for-age z score below -2	0.113 P=315	0.004 [0.026] N=2,708	0.003 [0.026] N=2,509	0.006 [0.028] N=2,241	0.025 [0.030] N=1,926	0.029 [0.033] N=1,591
WHO 2007 BMI overweight above +1, 61+ months	0.374 P=415	0.046 [0.051] N=1,101	0.048 [0.054] N=1,027	0.094 [0.060] N=920	0.061 [0.067] N=804	0.036 [0.074] N=670
WHO 2007 BMI obesity above +2, 61+ months	0.210 P=235	0.019 [0.044] N=1,101	0.011 [0.046] N=1,027	0.031 [0.049] N=920	-0.006 [0.052] N=804	-0.066 [0.059] N=670
WHO 2007 BMI thinness below -2, 61+ months	0.107 P=121	0.021 [0.043] N=1,101	0.014 [0.044] N=1,027	-0.016 [0.049] N=920	0.005 [0.053] N=804	0.016 [0.058] N=670

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.075 [0.201] N=2,378	-0.125 [0.213] N=2,168	-0.223 [0.249] N=1,895	-0.115 [0.275] N=1,564	-0.305 [0.370] N=1,213
WHO weight-for-age z score	-0.039 [0.112] N=2,531	0.034 [0.128] N=2,301	0.112 [0.137] N=2,010	-0.067 [0.156] N=1,656	-0.063 [0.205] N=1,284
WHO weight-for-length/height z score, 0-60 months	0.145 [0.267] N=1,359	0.102 [0.282] N=1,234	-0.122 [0.321] N=1,078	-0.488 [0.386] N=881	-0.349 [0.481] N=674
WHO BMI-for-age z score	0.270 [0.196] N=2,279	0.323 [0.207] N=2,080	**0.408* [0.240] N=1,812**	0.067 [0.287] N=1,497	0.362 [0.362] N=1,162
Stunted: HAZ below -2	0.040 [0.040] N=2,378	0.037 [0.043] N=2,168	0.066 [0.052] N=1,895	0.022 [0.058] N=1,564	0.098 [0.074] N=1,213
Underweight: WAZ below -2	0.002 [0.021] N=2,531	-0.007 [0.023] N=2,301	-0.023 [0.026] N=2,010	-0.001 [0.032] N=1,656	0.010 [0.042] N=1,284
Wasted: WLZ/WHZ below -2, 0-60 months	0.001 [0.040] N=1,359	0.012 [0.041] N=1,234	0.050 [0.045] N=1,078	0.068 [0.061] N=881	0.039 [0.067] N=674
Overweight: WLZ/WHZ above +2, 0-60 months	0.073 [0.054] N=1,359	0.063 [0.059] N=1,234	0.056 [0.067] N=1,078	-0.035 [0.082] N=881	-0.022 [0.110] N=674
Obese: WLZ/WHZ above +3, 0-60 months	0.045 [0.036] N=1,359	0.051 [0.039] N=1,234	0.056 [0.045] N=1,078	0.025 [0.056] N=881	0.038 [0.080] N=674
CIAF: any stunting, underweight, or wasting, 0-60 months	0.032 [0.061] N=1,336	0.005 [0.065] N=1,214	0.082 [0.075] N=1,061	0.042 [0.088] N=865	0.134 [0.103] N=664
BMI-for-age z score below -2	-0.019 [0.032] N=2,279	-0.027 [0.032] N=2,080	-0.032 [0.036] N=1,812	-0.010 [0.041] N=1,497	-0.021 [0.050] N=1,162
WHO 2007 BMI overweight above +1, 61+ months	0.105 [0.073] N=915	0.123 [0.082] N=841	**0.238** [0.093] N=734**	**0.210* [0.115] N=618**	**0.261* [0.144] N=484**
WHO 2007 BMI obesity above +2, 61+ months	**0.119* [0.065] N=915**	**0.120* [0.071] N=841**	**0.193** [0.078] N=734**	0.157 [0.095] N=618	0.114 [0.123] N=484
WHO 2007 BMI thinness below -2, 61+ months	-0.001 [0.045] N=915	-0.018 [0.048] N=841	-0.087 [0.056] N=734	-0.068 [0.062] N=618	-0.083 [0.081] N=484

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.273	0.050 [0.155] N=2,644	0.001 [0.162] N=2,453	-0.024 [0.176] N=2,191	0.010 [0.180] N=1,889	0.025 [0.210] N=1,556
WHO weight-for-age z score	0.200	0.072 [0.090] N=2,803	0.110 [0.096] N=2,594	0.165 [0.102] N=2,316	0.069 [0.109] N=1,993	0.097 [0.127] N=1,643
WHO weight-for-length/height z score, 0-60 months	0.355	**0.344* [0.190] N=1,480**	**0.346* [0.197] N=1,368**	0.231 [0.208] N=1,220	0.088 [0.242] N=1,046	0.274 [0.263] N=850
WHO BMI-for-age z score	0.357	0.164 [0.152] N=2,530	0.169 [0.156] N=2,349	0.192 [0.172] N=2,094	0.006 [0.194] N=1,807	0.065 [0.218] N=1,489
Stunted: HAZ below -2	0.201 P=513	0.000 [0.029] N=2,644	0.002 [0.030] N=2,453	0.010 [0.034] N=2,191	-0.011 [0.034] N=1,889	0.002 [0.040] N=1,556
Underweight: WAZ below -2	0.055 P=149	-0.014 [0.016] N=2,803	-0.019 [0.017] N=2,594	-0.028 [0.018] N=2,316	-0.019 [0.020] N=1,993	-0.021 [0.024] N=1,643
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=165	-0.030 [0.032] N=1,480	-0.024 [0.034] N=1,368	-0.001 [0.036] N=1,220	0.008 [0.044] N=1,046	-0.009 [0.047] N=850
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=279	**0.090** [0.044] N=1,480**	**0.093** [0.046] N=1,368**	**0.086* [0.050] N=1,220**	0.046 [0.058] N=1,046	0.067 [0.067] N=850
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=123	0.034 [0.031] N=1,480	0.043 [0.032] N=1,368	0.045 [0.035] N=1,220	0.029 [0.040] N=1,046	0.039 [0.048] N=850
CIAF: any stunting, underweight, or wasting, 0-60 months	0.278 P=396	-0.015 [0.049] N=1,456	-0.024 [0.051] N=1,347	0.013 [0.055] N=1,201	-0.009 [0.058] N=1,028	0.020 [0.067] N=837
BMI-for-age z score below -2	0.112 P=290	0.004 [0.027] N=2,530	0.005 [0.027] N=2,349	0.007 [0.029] N=2,094	0.022 [0.032] N=1,807	0.023 [0.035] N=1,489
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=391	0.047 [0.049] N=1,047	0.046 [0.052] N=979	**0.100* [0.057] N=876**	0.067 [0.062] N=765	0.038 [0.072] N=637
WHO 2007 BMI obesity above +2, 61+ months	0.206 P=220	0.013 [0.044] N=1,047	0.005 [0.045] N=979	0.033 [0.049] N=876	-0.007 [0.051] N=765	-0.078 [0.058] N=637
WHO 2007 BMI thinness below -2, 61+ months	0.109 P=116	0.031 [0.042] N=1,047	0.028 [0.044] N=979	-0.004 [0.048] N=876	0.013 [0.052] N=765	0.022 [0.060] N=637

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.001 [0.220] N=2,222	-0.100 [0.235] N=2,031	-0.199 [0.270] N=1,769	-0.114 [0.304] N=1,467	-0.251 [0.428] N=1,134
WHO weight-for-age z score	-0.011 [0.113] N=2,360	0.044 [0.125] N=2,151	0.131 [0.141] N=1,873	-0.054 [0.162] N=1,550	-0.022 [0.212] N=1,200
WHO weight-for-length/height z score, 0-60 months	0.209 [0.269] N=1,256	0.166 [0.281] N=1,144	-0.075 [0.308] N=996	-0.431 [0.392] N=822	-0.248 [0.507] N=626
WHO BMI-for-age z score	0.292 [0.195] N=2,130	0.328 [0.201] N=1,949	**0.435* [0.230] N=1,694**	0.128 [0.284] N=1,407	0.407 [0.371] N=1,089
Stunted: HAZ below -2	0.035 [0.042] N=2,222	0.039 [0.044] N=2,031	0.071 [0.054] N=1,769	0.030 [0.061] N=1,467	0.092 [0.081] N=1,134
Underweight: WAZ below -2	-0.003 [0.023] N=2,360	-0.012 [0.024] N=2,151	-0.026 [0.028] N=1,873	-0.004 [0.034] N=1,550	0.002 [0.044] N=1,200
Wasted: WLZ/WHZ below -2, 0-60 months	-0.015 [0.039] N=1,256	0.001 [0.039] N=1,144	0.046 [0.042] N=996	0.059 [0.059] N=822	0.022 [0.062] N=626
Overweight: WLZ/WHZ above +2, 0-60 months	0.075 [0.054] N=1,256	0.074 [0.059] N=1,144	0.072 [0.066] N=996	-0.025 [0.084] N=822	-0.011 [0.119] N=626
Obese: WLZ/WHZ above +3, 0-60 months	0.042 [0.038] N=1,256	0.055 [0.042] N=1,144	0.062 [0.048] N=996	0.028 [0.060] N=822	0.045 [0.092] N=626
CIAF: any stunting, underweight, or wasting, 0-60 months	0.020 [0.067] N=1,235	0.005 [0.071] N=1,126	0.088 [0.083] N=980	0.031 [0.096] N=807	0.123 [0.110] N=616
BMI-for-age z score below -2	-0.012 [0.032] N=2,130	-0.016 [0.030] N=1,949	-0.020 [0.034] N=1,694	-0.005 [0.039] N=1,407	-0.030 [0.048] N=1,089
WHO 2007 BMI overweight above +1, 61+ months	0.102 [0.075] N=870	0.118 [0.082] N=802	**0.236** [0.095] N=699**	**0.201* [0.114] N=588**	0.223 [0.154] N=460
WHO 2007 BMI obesity above +2, 61+ months	0.107 [0.065] N=870	0.109 [0.070] N=802	**0.194** [0.079] N=699**	0.151 [0.093] N=588	0.072 [0.123] N=460
WHO 2007 BMI thinness below -2, 61+ months	0.011 [0.046] N=870	-0.003 [0.048] N=802	-0.073 [0.055] N=699	-0.061 [0.062] N=588	-0.090 [0.087] N=460

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.273	0.058 [0.152] N=2,644	0.008 [0.159] N=2,453	-0.026 [0.173] N=2,191	0.009 [0.176] N=1,889	0.034 [0.202] N=1,556
WHO weight-for-age z score	0.200	0.080 [0.090] N=2,803	0.115 [0.096] N=2,594	**0.172* [0.101] N=2,316**	0.079 [0.109] N=1,993	0.113 [0.126] N=1,643
WHO weight-for-length/height z score, 0-60 months	0.355	**0.350* [0.190] N=1,480**	**0.352* [0.197] N=1,368**	0.234 [0.209] N=1,220	0.095 [0.242] N=1,046	0.276 [0.264] N=850
WHO BMI-for-age z score	0.357	0.165 [0.152] N=2,530	0.168 [0.155] N=2,349	0.194 [0.172] N=2,094	0.008 [0.193] N=1,807	0.069 [0.217] N=1,489
Stunted: HAZ below -2	0.201 P=513	-0.002 [0.029] N=2,644	0.000 [0.030] N=2,453	0.009 [0.033] N=2,191	-0.012 [0.034] N=1,889	-0.002 [0.039] N=1,556
Underweight: WAZ below -2	0.055 P=149	-0.016 [0.016] N=2,803	-0.021 [0.017] N=2,594	**-0.030* [0.018] N=2,316**	-0.022 [0.020] N=1,993	-0.027 [0.024] N=1,643
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=165	-0.032 [0.032] N=1,480	-0.026 [0.033] N=1,368	-0.001 [0.036] N=1,220	0.007 [0.044] N=1,046	-0.015 [0.048] N=850
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=279	**0.091** [0.044] N=1,480**	**0.093** [0.046] N=1,368**	**0.088* [0.051] N=1,220**	0.048 [0.059] N=1,046	0.071 [0.067] N=850
Obese: WLZ/WHZ above +3, 0-60 months	0.084 P=123	0.035 [0.031] N=1,480	0.043 [0.032] N=1,368	0.046 [0.035] N=1,220	0.031 [0.040] N=1,046	0.041 [0.048] N=850
CIAF: any stunting, underweight, or wasting, 0-60 months	0.278 P=396	-0.018 [0.048] N=1,456	-0.026 [0.050] N=1,347	0.013 [0.054] N=1,201	-0.008 [0.058] N=1,028	0.016 [0.065] N=837
BMI-for-age z score below -2	0.112 P=290	0.004 [0.027] N=2,530	0.004 [0.027] N=2,349	0.007 [0.029] N=2,094	0.022 [0.032] N=1,807	0.022 [0.035] N=1,489
WHO 2007 BMI overweight above +1, 61+ months	0.371 P=391	0.048 [0.049] N=1,047	0.047 [0.052] N=979	**0.100* [0.056] N=876**	0.065 [0.061] N=765	0.031 [0.070] N=637
WHO 2007 BMI obesity above +2, 61+ months	0.206 P=220	0.012 [0.043] N=1,047	0.003 [0.045] N=979	0.033 [0.049] N=876	-0.009 [0.050] N=765	-0.079 [0.058] N=637
WHO 2007 BMI thinness below -2, 61+ months	0.109 P=116	0.030 [0.042] N=1,047	0.026 [0.043] N=979	-0.005 [0.048] N=876	0.011 [0.052] N=765	0.022 [0.059] N=637

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.012 [0.219] N=2,222	-0.090 [0.234] N=2,031	-0.200 [0.269] N=1,769	-0.119 [0.301] N=1,467	-0.226 [0.416] N=1,134
WHO weight-for-age z score	-0.000 [0.114] N=2,360	0.050 [0.127] N=2,151	0.142 [0.140] N=1,873	-0.035 [0.163] N=1,550	0.014 [0.210] N=1,200
WHO weight-for-length/height z score, 0-60 months	0.235 [0.268] N=1,256	0.198 [0.282] N=1,144	-0.065 [0.308] N=996	-0.403 [0.393] N=822	-0.214 [0.510] N=626
WHO BMI-for-age z score	0.298 [0.194] N=2,130	0.332 [0.201] N=1,949	**0.437* [0.230] N=1,694**	0.131 [0.284] N=1,407	0.419 [0.374] N=1,089
Stunted: HAZ below -2	0.033 [0.042] N=2,222	0.038 [0.044] N=2,031	0.069 [0.053] N=1,769	0.027 [0.061] N=1,467	0.082 [0.080] N=1,134
Underweight: WAZ below -2	-0.005 [0.023] N=2,360	-0.013 [0.024] N=2,151	-0.029 [0.028] N=1,873	-0.009 [0.034] N=1,550	-0.010 [0.045] N=1,200
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.038] N=1,256	-0.002 [0.038] N=1,144	0.045 [0.042] N=996	0.056 [0.059] N=822	0.005 [0.060] N=626
Overweight: WLZ/WHZ above +2, 0-60 months	0.081 [0.053] N=1,256	0.080 [0.058] N=1,144	0.074 [0.066] N=996	-0.017 [0.084] N=822	0.001 [0.121] N=626
Obese: WLZ/WHZ above +3, 0-60 months	0.042 [0.038] N=1,256	0.053 [0.041] N=1,144	0.063 [0.048] N=996	0.033 [0.060] N=822	0.054 [0.092] N=626
CIAF: any stunting, underweight, or wasting, 0-60 months	0.018 [0.066] N=1,235	0.004 [0.070] N=1,126	0.086 [0.082] N=980	0.029 [0.095] N=807	0.097 [0.106] N=616
BMI-for-age z score below -2	-0.013 [0.032] N=2,130	-0.018 [0.030] N=1,949	-0.020 [0.034] N=1,694	-0.004 [0.039] N=1,407	-0.030 [0.048] N=1,089
WHO 2007 BMI overweight above +1, 61+ months	0.104 [0.074] N=870	0.120 [0.082] N=802	**0.238** [0.094] N=699**	**0.204* [0.113] N=588**	0.219 [0.151] N=460
WHO 2007 BMI obesity above +2, 61+ months	0.103 [0.064] N=870	0.103 [0.069] N=802	**0.191** [0.078] N=699**	0.147 [0.092] N=588	0.070 [0.122] N=460
WHO 2007 BMI thinness below -2, 61+ months	0.010 [0.046] N=870	-0.005 [0.047] N=802	-0.073 [0.055] N=699	-0.061 [0.062] N=588	-0.092 [0.086] N=460

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.227	-0.262 [0.178] N=2,814	-0.300 [0.193] N=2,553	-0.333 [0.202] N=2,248	**-0.439** [0.220] N=1,911**	**-0.456* [0.251] N=1,554**
WHO weight-for-age z score	0.224	-0.133 [0.099] N=2,981	-0.108 [0.104] N=2,716	-0.130 [0.111] N=2,388	**-0.201* [0.119] N=2,030**	-0.202 [0.136] N=1,641
WHO weight-for-length/height z score, 0-60 months	0.367	0.031 [0.214] N=1,520	0.081 [0.225] N=1,385	0.049 [0.248] N=1,199	0.103 [0.261] N=1,026	0.047 [0.309] N=828
WHO BMI-for-age z score	0.372	0.082 [0.168] N=2,700	0.114 [0.177] N=2,449	0.182 [0.187] N=2,156	0.247 [0.205] N=1,832	0.192 [0.240] N=1,487
Stunted: HAZ below -2	0.196 P=544	0.035 [0.033] N=2,814	0.036 [0.035] N=2,553	0.034 [0.037] N=2,248	0.047 [0.041] N=1,911	0.051 [0.046] N=1,554
Underweight: WAZ below -2	0.050 P=145	0.013 [0.015] N=2,981	0.014 [0.016] N=2,716	0.015 [0.017] N=2,388	0.021 [0.017] N=2,030	**0.038* [0.019] N=1,641**
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=164	0.015 [0.037] N=1,520	0.003 [0.039] N=1,385	0.013 [0.042] N=1,199	-0.011 [0.043] N=1,026	-0.002 [0.053] N=828
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=296	0.027 [0.040] N=1,520	0.029 [0.041] N=1,385	0.021 [0.044] N=1,199	0.021 [0.045] N=1,026	0.027 [0.051] N=828
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=140	0.033 [0.029] N=1,520	0.030 [0.029] N=1,385	0.026 [0.033] N=1,199	0.030 [0.036] N=1,026	0.006 [0.037] N=828
CIAF: any stunting, underweight, or wasting, 0-60 months	0.281 P=414	0.055 [0.046] N=1,492	0.040 [0.050] N=1,357	0.045 [0.052] N=1,173	0.016 [0.056] N=1,004	0.052 [0.065] N=809
BMI-for-age z score below -2	0.111 P=303	0.035 [0.027] N=2,700	0.028 [0.030] N=2,449	0.024 [0.031] N=2,156	0.009 [0.032] N=1,832	0.024 [0.038] N=1,487
WHO 2007 BMI overweight above +1, 61+ months	0.392 P=458	0.003 [0.058] N=1,181	-0.015 [0.061] N=1,067	0.017 [0.064] N=961	0.016 [0.073] N=809	-0.032 [0.080] N=656
WHO 2007 BMI obesity above +2, 61+ months	0.216 P=253	0.021 [0.049] N=1,181	-0.005 [0.053] N=1,067	0.018 [0.056] N=961	0.048 [0.061] N=809	0.042 [0.071] N=656
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=127	0.024 [0.035] N=1,181	0.019 [0.037] N=1,067	0.005 [0.038] N=961	-0.009 [0.043] N=809	0.018 [0.048] N=656

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.069 [0.218] N=2,412	-0.109 [0.248] N=2,151	-0.140 [0.259] N=1,846	-0.332 [0.284] N=1,509	-0.216 [0.373] N=1,152
WHO weight-for-age z score	-0.168 [0.148] N=2,559	-0.123 [0.162] N=2,294	-0.197 [0.182] N=1,966	**-0.397* [0.206] N=1,608**	**-0.486* [0.254] N=1,219**
WHO weight-for-length/height z score, 0-60 months	-0.067 [0.274] N=1,326	-0.004 [0.288] N=1,191	-0.171 [0.345] N=1,005	-0.173 [0.365] N=832	-0.602 [0.463] N=634
WHO BMI-for-age z score	**-0.353** [0.176] N=2,317**	**-0.372** [0.182] N=2,066**	**-0.345* [0.206] N=1,773**	-0.389 [0.250] N=1,449	**-0.968*** [0.303] N=1,104**
Stunted: HAZ below -2	0.019 [0.043] N=2,412	0.018 [0.049] N=2,151	0.004 [0.052] N=1,846	0.023 [0.060] N=1,509	0.013 [0.082] N=1,152
Underweight: WAZ below -2	-0.003 [0.022] N=2,559	-0.004 [0.024] N=2,294	-0.003 [0.027] N=1,966	0.001 [0.028] N=1,608	0.031 [0.035] N=1,219
Wasted: WLZ/WHZ below -2, 0-60 months	0.061 [0.041] N=1,326	0.046 [0.045] N=1,191	0.084 [0.051] N=1,005	0.046 [0.049] N=832	**0.110* [0.056] N=634**
Overweight: WLZ/WHZ above +2, 0-60 months	0.042 [0.060] N=1,326	0.040 [0.064] N=1,191	0.016 [0.071] N=1,005	-0.006 [0.084] N=832	-0.042 [0.101] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.047 [0.042] N=1,326	0.037 [0.044] N=1,191	0.025 [0.052] N=1,005	0.028 [0.062] N=832	-0.080 [0.074] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.085 [0.062] N=1,303	0.067 [0.070] N=1,168	0.082 [0.078] N=984	0.026 [0.088] N=815	0.113 [0.107] N=620
BMI-for-age z score below -2	**0.087*** [0.025] N=2,317**	**0.083*** [0.030] N=2,066**	**0.081** [0.035] N=1,773**	**0.065** [0.032] N=1,449**	**0.143*** [0.041] N=1,104**
WHO 2007 BMI overweight above +1, 61+ months	-0.050 [0.083] N=992	-0.087 [0.093] N=878	-0.041 [0.101] N=772	-0.055 [0.124] N=620	**-0.252* [0.144] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.049 [0.063] N=992	-0.108 [0.067] N=878	-0.090 [0.073] N=772	-0.072 [0.082] N=620	-0.154 [0.113] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.078* [0.044] N=992**	0.079 [0.049] N=878	0.051 [0.054] N=772	0.038 [0.063] N=620	**0.144** [0.069] N=467**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.227	-0.259 [0.178] N=2,814	-0.297 [0.193] N=2,553	-0.332 [0.202] N=2,248	**-0.436* [0.221] N=1,911**	**-0.448* [0.253] N=1,554**
WHO weight-for-age z score	0.224	-0.129 [0.099] N=2,981	-0.103 [0.104] N=2,716	-0.127 [0.111] N=2,388	**-0.198* [0.119] N=2,030**	-0.197 [0.136] N=1,641
WHO weight-for-length/height z score, 0-60 months	0.367	0.039 [0.214] N=1,520	0.088 [0.225] N=1,385	0.052 [0.248] N=1,199	0.105 [0.261] N=1,026	0.041 [0.308] N=828
WHO BMI-for-age z score	0.372	0.080 [0.169] N=2,700	0.112 [0.177] N=2,449	0.182 [0.187] N=2,156	0.246 [0.206] N=1,832	0.189 [0.240] N=1,487
Stunted: HAZ below -2	0.196 P=544	0.035 [0.033] N=2,814	0.037 [0.035] N=2,553	0.034 [0.037] N=2,248	0.047 [0.040] N=1,911	0.052 [0.046] N=1,554
Underweight: WAZ below -2	0.050 P=145	0.013 [0.015] N=2,981	0.013 [0.016] N=2,716	0.015 [0.017] N=2,388	0.021 [0.017] N=2,030	**0.038* [0.019] N=1,641**
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=164	0.014 [0.037] N=1,520	0.001 [0.039] N=1,385	0.011 [0.041] N=1,199	-0.012 [0.042] N=1,026	-0.000 [0.052] N=828
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=296	0.027 [0.040] N=1,520	0.029 [0.041] N=1,385	0.020 [0.043] N=1,199	0.021 [0.046] N=1,026	0.027 [0.051] N=828
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=140	0.033 [0.029] N=1,520	0.029 [0.029] N=1,385	0.025 [0.033] N=1,199	0.029 [0.036] N=1,026	0.006 [0.037] N=828
CIAF: any stunting, underweight, or wasting, 0-60 months	0.281 P=414	0.054 [0.046] N=1,492	0.038 [0.050] N=1,357	0.043 [0.052] N=1,173	0.016 [0.057] N=1,004	0.053 [0.065] N=809
BMI-for-age z score below -2	0.111 P=303	0.036 [0.027] N=2,700	0.029 [0.030] N=2,449	0.025 [0.032] N=2,156	0.010 [0.032] N=1,832	0.025 [0.038] N=1,487
WHO 2007 BMI overweight above +1, 61+ months	0.392 P=458	0.007 [0.058] N=1,181	-0.012 [0.061] N=1,067	0.019 [0.064] N=961	0.018 [0.072] N=809	-0.027 [0.079] N=656
WHO 2007 BMI obesity above +2, 61+ months	0.216 P=253	0.029 [0.050] N=1,181	0.002 [0.054] N=1,067	0.023 [0.057] N=961	0.052 [0.062] N=809	0.046 [0.071] N=656
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=127	0.026 [0.035] N=1,181	0.022 [0.037] N=1,067	0.006 [0.038] N=961	-0.009 [0.043] N=809	0.017 [0.048] N=656

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.067 [0.218] N=2,412	-0.107 [0.247] N=2,151	-0.139 [0.258] N=1,846	-0.331 [0.284] N=1,509	-0.202 [0.375] N=1,152
WHO weight-for-age z score	-0.164 [0.148] N=2,559	-0.119 [0.163] N=2,294	-0.193 [0.183] N=1,966	**-0.394* [0.208] N=1,608**	**-0.476* [0.257] N=1,219**
WHO weight-for-length/height z score, 0-60 months	-0.043 [0.277] N=1,326	0.018 [0.291] N=1,191	-0.156 [0.347] N=1,005	-0.152 [0.367] N=832	-0.579 [0.475] N=634
WHO BMI-for-age z score	**-0.355** [0.176] N=2,317**	**-0.374** [0.182] N=2,066**	**-0.345* [0.206] N=1,773**	-0.390 [0.250] N=1,449	**-0.975*** [0.302] N=1,104**
Stunted: HAZ below -2	0.019 [0.043] N=2,412	0.018 [0.049] N=2,151	0.004 [0.052] N=1,846	0.023 [0.060] N=1,509	0.013 [0.082] N=1,152
Underweight: WAZ below -2	-0.003 [0.022] N=2,559	-0.005 [0.024] N=2,294	-0.003 [0.027] N=1,966	0.001 [0.028] N=1,608	0.032 [0.034] N=1,219
Wasted: WLZ/WHZ below -2, 0-60 months	0.056 [0.042] N=1,326	0.041 [0.045] N=1,191	0.077 [0.052] N=1,005	0.038 [0.050] N=832	**0.103* [0.058] N=634**
Overweight: WLZ/WHZ above +2, 0-60 months	0.042 [0.060] N=1,326	0.040 [0.064] N=1,191	0.014 [0.071] N=1,005	-0.008 [0.083] N=832	-0.043 [0.101] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.046 [0.042] N=1,326	0.036 [0.044] N=1,191	0.022 [0.052] N=1,005	0.026 [0.062] N=832	-0.082 [0.074] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.081 [0.062] N=1,303	0.061 [0.071] N=1,168	0.073 [0.079] N=984	0.019 [0.089] N=815	0.109 [0.108] N=620
BMI-for-age z score below -2	**0.088*** [0.025] N=2,317**	**0.084*** [0.030] N=2,066**	**0.082** [0.036] N=1,773**	**0.066** [0.032] N=1,449**	**0.146*** [0.041] N=1,104**
WHO 2007 BMI overweight above +1, 61+ months	-0.044 [0.083] N=992	-0.082 [0.094] N=878	-0.039 [0.102] N=772	-0.054 [0.125] N=620	-0.239 [0.147] N=467
WHO 2007 BMI obesity above +2, 61+ months	-0.037 [0.065] N=992	-0.097 [0.070] N=878	-0.083 [0.076] N=772	-0.067 [0.085] N=620	-0.144 [0.116] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.083* [0.045] N=992**	**0.086* [0.049] N=878**	0.051 [0.054] N=772	0.040 [0.063] N=620	**0.145** [0.069] N=467**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.227	-0.255 [0.176] N=2,814	-0.284 [0.190] N=2,553	-0.313 [0.198] N=2,248	**-0.423* [0.216] N=1,911**	**-0.424* [0.245] N=1,554**
WHO weight-for-age z score	0.224	-0.132 [0.098] N=2,981	-0.103 [0.104] N=2,716	-0.117 [0.110] N=2,388	**-0.198* [0.117] N=2,030**	-0.188 [0.134] N=1,641
WHO weight-for-length/height z score, 0-60 months	0.367	0.039 [0.214] N=1,520	0.090 [0.225] N=1,385	0.077 [0.249] N=1,199	0.130 [0.262] N=1,026	0.084 [0.313] N=828
WHO BMI-for-age z score	0.372	0.076 [0.169] N=2,700	0.108 [0.178] N=2,449	0.184 [0.187] N=2,156	0.241 [0.206] N=1,832	0.184 [0.241] N=1,487
Stunted: HAZ below -2	0.196 P=544	0.035 [0.033] N=2,814	0.034 [0.035] N=2,553	0.032 [0.037] N=2,248	0.045 [0.040] N=1,911	0.047 [0.045] N=1,554
Underweight: WAZ below -2	0.050 P=145	0.013 [0.015] N=2,981	0.013 [0.016] N=2,716	0.014 [0.017] N=2,388	0.020 [0.017] N=2,030	**0.037** [0.018] N=1,641**
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=164	0.018 [0.037] N=1,520	0.004 [0.039] N=1,385	0.013 [0.042] N=1,199	-0.009 [0.043] N=1,026	0.001 [0.053] N=828
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=296	0.029 [0.040] N=1,520	0.030 [0.041] N=1,385	0.027 [0.043] N=1,199	0.028 [0.045] N=1,026	0.037 [0.051] N=828
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=140	0.036 [0.029] N=1,520	0.031 [0.030] N=1,385	0.029 [0.033] N=1,199	0.034 [0.037] N=1,026	0.013 [0.038] N=828
CIAF: any stunting, underweight, or wasting, 0-60 months	0.281 P=414	0.065 [0.046] N=1,492	0.046 [0.050] N=1,357	0.048 [0.051] N=1,173	0.022 [0.057] N=1,004	0.057 [0.065] N=809
BMI-for-age z score below -2	0.111 P=303	0.035 [0.027] N=2,700	0.028 [0.029] N=2,449	0.023 [0.031] N=2,156	0.009 [0.032] N=1,832	0.024 [0.038] N=1,487
WHO 2007 BMI overweight above +1, 61+ months	0.392 P=458	-0.003 [0.058] N=1,181	-0.019 [0.063] N=1,067	0.009 [0.065] N=961	0.010 [0.073] N=809	-0.030 [0.080] N=656
WHO 2007 BMI obesity above +2, 61+ months	0.216 P=253	0.017 [0.050] N=1,181	-0.008 [0.054] N=1,067	0.011 [0.057] N=961	0.038 [0.061] N=809	0.041 [0.070] N=656
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=127	0.021 [0.034] N=1,181	0.017 [0.036] N=1,067	0.004 [0.038] N=961	-0.010 [0.043] N=809	0.014 [0.048] N=656

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.056 [0.219] N=2,412	-0.083 [0.247] N=2,151	-0.096 [0.257] N=1,846	-0.289 [0.281] N=1,509	-0.113 [0.375] N=1,152
WHO weight-for-age z score	-0.172 [0.148] N=2,559	-0.123 [0.163] N=2,294	-0.182 [0.184] N=1,966	**-0.403* [0.206] N=1,608**	**-0.473* [0.261] N=1,219**
WHO weight-for-length/height z score, 0-60 months	-0.056 [0.279] N=1,326	0.008 [0.294] N=1,191	-0.145 [0.348] N=1,005	-0.154 [0.368] N=832	-0.574 [0.476] N=634
WHO BMI-for-age z score	**-0.367** [0.175] N=2,317**	**-0.385** [0.183] N=2,066**	**-0.356* [0.206] N=1,773**	-0.418 [0.252] N=1,449	**-1.010*** [0.311] N=1,104**
Stunted: HAZ below -2	0.016 [0.044] N=2,412	0.012 [0.050] N=2,151	-0.004 [0.053] N=1,846	0.013 [0.062] N=1,509	-0.007 [0.084] N=1,152
Underweight: WAZ below -2	-0.003 [0.021] N=2,559	-0.005 [0.024] N=2,294	-0.006 [0.027] N=1,966	0.000 [0.028] N=1,608	0.029 [0.033] N=1,219
Wasted: WLZ/WHZ below -2, 0-60 months	0.062 [0.042] N=1,326	0.043 [0.046] N=1,191	0.077 [0.052] N=1,005	0.039 [0.051] N=832	**0.104* [0.057] N=634**
Overweight: WLZ/WHZ above +2, 0-60 months	0.043 [0.061] N=1,326	0.039 [0.065] N=1,191	0.017 [0.072] N=1,005	-0.008 [0.084] N=832	-0.040 [0.105] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.046 [0.043] N=1,326	0.035 [0.045] N=1,191	0.022 [0.053] N=1,005	0.025 [0.063] N=832	-0.083 [0.077] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.063] N=1,303	0.064 [0.072] N=1,168	0.067 [0.080] N=984	0.013 [0.091] N=815	0.095 [0.108] N=620
BMI-for-age z score below -2	**0.087*** [0.025] N=2,317**	**0.080*** [0.030] N=2,066**	**0.078** [0.035] N=1,773**	**0.062* [0.032] N=1,449**	**0.141*** [0.041] N=1,104**
WHO 2007 BMI overweight above +1, 61+ months	-0.062 [0.084] N=992	-0.096 [0.096] N=878	-0.057 [0.104] N=772	-0.072 [0.127] N=620	**-0.261* [0.149] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.054 [0.065] N=992	-0.111 [0.071] N=878	-0.100 [0.076] N=772	-0.088 [0.085] N=620	-0.156 [0.117] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.077* [0.043] N=992**	**0.079* [0.047] N=878**	0.049 [0.053] N=772	0.040 [0.062] N=620	**0.138** [0.069] N=467**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.227	-0.260 [0.176] N=2,814	-0.290 [0.189] N=2,553	-0.326 [0.199] N=2,248	**-0.427* [0.217] N=1,911**	**-0.442* [0.245] N=1,554**
WHO weight-for-age z score	0.224	-0.134 [0.096] N=2,981	-0.104 [0.101] N=2,716	-0.125 [0.106] N=2,388	**-0.197* [0.112] N=2,030**	-0.208 [0.127] N=1,641
WHO weight-for-length/height z score, 0-60 months	0.367	0.047 [0.214] N=1,520	0.099 [0.225] N=1,385	0.076 [0.248] N=1,199	0.133 [0.259] N=1,026	0.055 [0.308] N=828
WHO BMI-for-age z score	0.372	0.077 [0.168] N=2,700	0.109 [0.176] N=2,449	0.181 [0.186] N=2,156	0.246 [0.205] N=1,832	0.176 [0.240] N=1,487
Stunted: HAZ below -2	0.196 P=544	0.036 [0.032] N=2,814	0.036 [0.035] N=2,553	0.034 [0.037] N=2,248	0.046 [0.040] N=1,911	0.049 [0.045] N=1,554
Underweight: WAZ below -2	0.050 P=145	0.013 [0.015] N=2,981	0.013 [0.016] N=2,716	0.015 [0.016] N=2,388	0.021 [0.017] N=2,030	**0.038** [0.018] N=1,641**
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=164	0.017 [0.037] N=1,520	0.004 [0.039] N=1,385	0.015 [0.042] N=1,199	-0.010 [0.042] N=1,026	-0.001 [0.053] N=828
Overweight: WLZ/WHZ above +2, 0-60 months	0.194 P=296	0.033 [0.040] N=1,520	0.034 [0.041] N=1,385	0.031 [0.043] N=1,199	0.030 [0.045] N=1,026	0.030 [0.051] N=828
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=140	0.038 [0.029] N=1,520	0.033 [0.029] N=1,385	0.030 [0.033] N=1,199	0.035 [0.036] N=1,026	0.010 [0.038] N=828
CIAF: any stunting, underweight, or wasting, 0-60 months	0.281 P=414	0.066 [0.045] N=1,492	0.048 [0.050] N=1,357	0.051 [0.051] N=1,173	0.024 [0.056] N=1,004	0.055 [0.063] N=809
BMI-for-age z score below -2	0.111 P=303	0.035 [0.027] N=2,700	0.027 [0.030] N=2,449	0.023 [0.031] N=2,156	0.007 [0.032] N=1,832	0.021 [0.037] N=1,487
WHO 2007 BMI overweight above +1, 61+ months	0.392 P=458	0.000 [0.058] N=1,181	-0.017 [0.062] N=1,067	0.015 [0.064] N=961	0.015 [0.072] N=809	-0.027 [0.080] N=656
WHO 2007 BMI obesity above +2, 61+ months	0.216 P=253	0.018 [0.049] N=1,181	-0.008 [0.053] N=1,067	0.012 [0.056] N=961	0.038 [0.060] N=809	0.030 [0.070] N=656
WHO 2007 BMI thinness below -2, 61+ months	0.106 P=127	0.022 [0.035] N=1,181	0.019 [0.036] N=1,067	0.003 [0.038] N=961	-0.012 [0.043] N=809	0.015 [0.047] N=656

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.069 [0.215] N=2,412	-0.096 [0.243] N=2,151	-0.112 [0.254] N=1,846	-0.289 [0.277] N=1,509	-0.161 [0.358] N=1,152
WHO weight-for-age z score	-0.162 [0.144] N=2,559	-0.110 [0.159] N=2,294	-0.172 [0.178] N=1,966	**-0.367* [0.198] N=1,608**	**-0.459* [0.246] N=1,219**
WHO weight-for-length/height z score, 0-60 months	-0.035 [0.278] N=1,326	0.025 [0.290] N=1,191	-0.144 [0.343] N=1,005	-0.133 [0.362] N=832	-0.568 [0.479] N=634
WHO BMI-for-age z score	**-0.354** [0.175] N=2,317**	**-0.375** [0.181] N=2,066**	**-0.354* [0.205] N=1,773**	-0.399 [0.249] N=1,449	**-0.991*** [0.310] N=1,104**
Stunted: HAZ below -2	0.020 [0.043] N=2,412	0.016 [0.049] N=2,151	-0.003 [0.053] N=1,846	0.013 [0.062] N=1,509	0.001 [0.083] N=1,152
Underweight: WAZ below -2	-0.002 [0.021] N=2,559	-0.005 [0.024] N=2,294	-0.005 [0.027] N=1,966	-0.002 [0.028] N=1,608	0.028 [0.033] N=1,219
Wasted: WLZ/WHZ below -2, 0-60 months	0.062 [0.041] N=1,326	0.045 [0.045] N=1,191	0.084 [0.051] N=1,005	0.042 [0.049] N=832	**0.107* [0.056] N=634**
Overweight: WLZ/WHZ above +2, 0-60 months	0.049 [0.060] N=1,326	0.045 [0.064] N=1,191	0.025 [0.069] N=1,005	0.001 [0.081] N=832	-0.037 [0.102] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.051 [0.043] N=1,326	0.039 [0.044] N=1,191	0.024 [0.052] N=1,005	0.027 [0.062] N=832	-0.082 [0.077] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.097 [0.061] N=1,303	0.070 [0.070] N=1,168	0.073 [0.078] N=984	0.016 [0.086] N=815	0.108 [0.100] N=620
BMI-for-age z score below -2	**0.090*** [0.025] N=2,317**	**0.084*** [0.030] N=2,066**	**0.082** [0.035] N=1,773**	**0.065** [0.032] N=1,449**	**0.146*** [0.041] N=1,104**
WHO 2007 BMI overweight above +1, 61+ months	-0.059 [0.084] N=992	-0.094 [0.095] N=878	-0.056 [0.103] N=772	-0.073 [0.124] N=620	**-0.252* [0.148] N=467**
WHO 2007 BMI obesity above +2, 61+ months	-0.049 [0.065] N=992	-0.106 [0.070] N=878	-0.096 [0.075] N=772	-0.080 [0.082] N=620	-0.156 [0.114] N=467
WHO 2007 BMI thinness below -2, 61+ months	**0.077* [0.044] N=992**	**0.081* [0.048] N=878**	0.048 [0.053] N=772	0.035 [0.062] N=620	**0.137* [0.069] N=467**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.237	-0.249 [0.181] N=2,639	-0.263 [0.197] N=2,395	-0.295 [0.204] N=2,113	**-0.392* [0.223] N=1,791**	**-0.455* [0.256] N=1,459**
WHO weight-for-age z score	0.227	-0.111 [0.106] N=2,789	-0.076 [0.112] N=2,541	-0.105 [0.121] N=2,239	-0.166 [0.131] N=1,900	-0.203 [0.147] N=1,540
WHO weight-for-length/height z score, 0-60 months	0.373	0.087 [0.225] N=1,413	0.152 [0.237] N=1,288	0.062 [0.259] N=1,116	0.094 [0.272] N=955	0.023 [0.317] N=775
WHO BMI-for-age z score	0.386	0.062 [0.167] N=2,532	0.097 [0.176] N=2,297	0.143 [0.185] N=2,026	0.209 [0.200] N=1,718	0.142 [0.234] N=1,398
Stunted: HAZ below -2	0.199 P=516	0.041 [0.032] N=2,639	0.042 [0.034] N=2,395	0.039 [0.035] N=2,113	0.051 [0.038] N=1,791	0.062 [0.042] N=1,459
Underweight: WAZ below -2	0.051 P=138	0.011 [0.016] N=2,789	0.012 [0.017] N=2,541	0.010 [0.018] N=2,239	0.020 [0.019] N=1,900	**0.044** [0.021] N=1,540**
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=154	0.015 [0.037] N=1,413	-0.001 [0.039] N=1,288	0.016 [0.042] N=1,116	-0.001 [0.043] N=955	0.009 [0.052] N=775
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=277	0.024 [0.042] N=1,413	0.028 [0.043] N=1,288	0.018 [0.046] N=1,116	0.013 [0.049] N=955	0.019 [0.056] N=775
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=134	0.039 [0.029] N=1,413	0.036 [0.030] N=1,288	0.032 [0.033] N=1,116	0.040 [0.038] N=955	0.022 [0.041] N=775
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=390	0.071 [0.050] N=1,387	0.053 [0.055] N=1,262	0.065 [0.057] N=1,092	0.038 [0.062] N=934	0.086 [0.070] N=756
BMI-for-age z score below -2	0.110 P=279	0.039 [0.026] N=2,532	0.032 [0.029] N=2,297	0.030 [0.030] N=2,026	0.019 [0.029] N=1,718	0.036 [0.033] N=1,398
WHO 2007 BMI overweight above +1, 61+ months	0.396 P=439	-0.002 [0.059] N=1,123	-0.021 [0.062] N=1,015	0.019 [0.065] N=916	0.028 [0.074] N=768	-0.010 [0.080] N=622
WHO 2007 BMI obesity above +2, 61+ months	0.220 P=245	0.014 [0.052] N=1,123	-0.015 [0.055] N=1,015	0.007 [0.058] N=916	0.042 [0.062] N=768	0.041 [0.073] N=622
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=119	0.028 [0.036] N=1,123	0.025 [0.037] N=1,015	0.007 [0.038] N=916	-0.013 [0.043] N=768	0.015 [0.047] N=622

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.022 [0.227] N=2,261	-0.009 [0.257] N=2,017	-0.054 [0.274] N=1,735	-0.203 [0.300] N=1,413	-0.163 [0.389] N=1,081
WHO weight-for-age z score	-0.154 [0.155] N=2,394	-0.093 [0.169] N=2,146	-0.197 [0.193] N=1,844	**-0.371* [0.217] N=1,505**	**-0.501* [0.261] N=1,145**
WHO weight-for-length/height z score, 0-60 months	-0.008 [0.307] N=1,230	0.066 [0.326] N=1,105	-0.228 [0.390] N=933	-0.299 [0.431] N=772	-0.770 [0.576] N=592
WHO BMI-for-age z score	**-0.364** [0.184] N=2,171**	**-0.384** [0.193] N=1,936**	**-0.398* [0.218] N=1,665**	**-0.445* [0.264] N=1,357**	**-1.072*** [0.319] N=1,037**
Stunted: HAZ below -2	0.017 [0.046] N=2,261	0.014 [0.052] N=2,017	0.001 [0.055] N=1,735	0.014 [0.063] N=1,413	0.013 [0.081] N=1,081
Underweight: WAZ below -2	-0.008 [0.023] N=2,394	-0.010 [0.025] N=2,146	-0.013 [0.028] N=1,844	-0.003 [0.030] N=1,505	0.034 [0.036] N=1,145
Wasted: WLZ/WHZ below -2, 0-60 months	0.057 [0.044] N=1,230	0.035 [0.048] N=1,105	0.083 [0.053] N=933	0.052 [0.054] N=772	**0.121* [0.063] N=592**
Overweight: WLZ/WHZ above +2, 0-60 months	0.032 [0.064] N=1,230	0.029 [0.067] N=1,105	-0.006 [0.076] N=933	-0.045 [0.095] N=772	-0.072 [0.120] N=592
Obese: WLZ/WHZ above +3, 0-60 months	0.037 [0.045] N=1,230	0.026 [0.047] N=1,105	0.004 [0.055] N=933	0.008 [0.067] N=772	-0.089 [0.085] N=592
CIAF: any stunting, underweight, or wasting, 0-60 months	0.084 [0.070] N=1,209	0.058 [0.081] N=1,084	0.077 [0.091] N=914	0.006 [0.103] N=756	0.125 [0.129] N=578
BMI-for-age z score below -2	**0.093*** [0.027] N=2,171**	**0.089*** [0.032] N=1,936**	**0.090** [0.037] N=1,665**	**0.082** [0.033] N=1,357**	**0.173*** [0.041] N=1,037**
WHO 2007 BMI overweight above +1, 61+ months	-0.051 [0.084] N=944	-0.092 [0.095] N=836	-0.032 [0.104] N=737	-0.019 [0.126] N=589	-0.180 [0.150] N=443
WHO 2007 BMI obesity above +2, 61+ months	-0.047 [0.068] N=944	-0.110 [0.074] N=836	-0.092 [0.079] N=737	-0.056 [0.087] N=589	-0.124 [0.118] N=443
WHO 2007 BMI thinness below -2, 61+ months	**0.090* [0.048] N=944**	**0.096* [0.052] N=836**	0.058 [0.057] N=737	0.036 [0.066] N=589	**0.142* [0.077] N=443**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.237	-0.247 [0.181] N=2,639	-0.260 [0.197] N=2,395	-0.289 [0.205] N=2,113	**-0.383* [0.225] N=1,791**	**-0.440* [0.259] N=1,459**
WHO weight-for-age z score	0.227	-0.114 [0.107] N=2,789	-0.077 [0.113] N=2,541	-0.111 [0.122] N=2,239	-0.168 [0.132] N=1,900	-0.200 [0.149] N=1,540
WHO weight-for-length/height z score, 0-60 months	0.373	0.099 [0.226] N=1,413	0.162 [0.238] N=1,288	0.084 [0.262] N=1,116	0.123 [0.278] N=955	0.047 [0.330] N=775
WHO BMI-for-age z score	0.386	0.062 [0.168] N=2,532	0.097 [0.176] N=2,297	0.143 [0.185] N=2,026	0.211 [0.202] N=1,718	0.142 [0.236] N=1,398
Stunted: HAZ below -2	0.199 P=516	0.040 [0.032] N=2,639	0.040 [0.034] N=2,395	0.037 [0.035] N=2,113	0.050 [0.038] N=1,791	0.061 [0.043] N=1,459
Underweight: WAZ below -2	0.051 P=138	0.012 [0.016] N=2,789	0.012 [0.017] N=2,541	0.012 [0.018] N=2,239	0.021 [0.019] N=1,900	**0.044** [0.021] N=1,540**
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=154	0.011 [0.038] N=1,413	-0.006 [0.040] N=1,288	0.009 [0.042] N=1,116	-0.010 [0.043] N=955	-0.002 [0.053] N=775
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=277	0.024 [0.041] N=1,413	0.027 [0.043] N=1,288	0.018 [0.045] N=1,116	0.014 [0.048] N=955	0.018 [0.056] N=775
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=134	0.038 [0.029] N=1,413	0.035 [0.030] N=1,288	0.030 [0.033] N=1,116	0.038 [0.037] N=955	0.020 [0.040] N=775
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=390	0.066 [0.050] N=1,387	0.046 [0.055] N=1,262	0.054 [0.057] N=1,092	0.026 [0.062] N=934	0.073 [0.070] N=756
BMI-for-age z score below -2	0.110 P=279	0.039 [0.026] N=2,532	0.031 [0.029] N=2,297	0.029 [0.030] N=2,026	0.016 [0.029] N=1,718	0.033 [0.033] N=1,398
WHO 2007 BMI overweight above +1, 61+ months	0.396 P=439	-0.007 [0.059] N=1,123	-0.026 [0.063] N=1,015	0.014 [0.065] N=916	0.022 [0.074] N=768	-0.018 [0.080] N=622
WHO 2007 BMI obesity above +2, 61+ months	0.220 P=245	0.012 [0.051] N=1,123	-0.017 [0.055] N=1,015	0.004 [0.058] N=916	0.037 [0.062] N=768	0.037 [0.072] N=622
WHO 2007 BMI thinness below -2, 61+ months	0.105 P=119	0.028 [0.035] N=1,123	0.025 [0.036] N=1,015	0.004 [0.037] N=916	-0.017 [0.041] N=768	0.007 [0.044] N=622

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.033 [0.225] N=2,261	-0.016 [0.254] N=2,017	-0.050 [0.272] N=1,735	-0.211 [0.305] N=1,413	-0.167 [0.394] N=1,081
WHO weight-for-age z score	-0.156 [0.154] N=2,394	-0.094 [0.169] N=2,146	-0.201 [0.194] N=1,844	**-0.369* [0.218] N=1,505**	**-0.506* [0.263] N=1,145**
WHO weight-for-length/height z score, 0-60 months	0.000 [0.305] N=1,230	0.070 [0.324] N=1,105	-0.215 [0.388] N=933	-0.260 [0.429] N=772	-0.787 [0.557] N=592
WHO BMI-for-age z score	**-0.360* [0.184] N=2,171**	**-0.382** [0.192] N=1,936**	**-0.397* [0.216] N=1,665**	-0.430 [0.265] N=1,357	**-1.066*** [0.317] N=1,037**
Stunted: HAZ below -2	0.018 [0.046] N=2,261	0.015 [0.052] N=2,017	0.001 [0.055] N=1,735	0.016 [0.063] N=1,413	0.019 [0.082] N=1,081
Underweight: WAZ below -2	-0.006 [0.022] N=2,394	-0.009 [0.025] N=2,146	-0.011 [0.028] N=1,844	-0.000 [0.030] N=1,505	0.041 [0.036] N=1,145
Wasted: WLZ/WHZ below -2, 0-60 months	0.054 [0.044] N=1,230	0.033 [0.048] N=1,105	0.079 [0.053] N=933	0.044 [0.053] N=772	**0.117* [0.061] N=592**
Overweight: WLZ/WHZ above +2, 0-60 months	0.033 [0.063] N=1,230	0.029 [0.067] N=1,105	-0.005 [0.075] N=933	-0.039 [0.094] N=772	-0.078 [0.117] N=592
Obese: WLZ/WHZ above +3, 0-60 months	0.038 [0.045] N=1,230	0.026 [0.046] N=1,105	0.005 [0.054] N=933	0.011 [0.066] N=772	-0.091 [0.082] N=592
CIAF: any stunting, underweight, or wasting, 0-60 months	0.084 [0.069] N=1,209	0.058 [0.080] N=1,084	0.075 [0.090] N=914	0.009 [0.102] N=756	0.137 [0.128] N=578
BMI-for-age z score below -2	**0.093*** [0.027] N=2,171**	**0.089*** [0.032] N=1,936**	**0.090** [0.037] N=1,665**	**0.080** [0.033] N=1,357**	**0.173*** [0.040] N=1,037**
WHO 2007 BMI overweight above +1, 61+ months	-0.053 [0.085] N=944	-0.092 [0.095] N=836	-0.032 [0.104] N=737	-0.018 [0.126] N=589	-0.176 [0.149] N=443
WHO 2007 BMI obesity above +2, 61+ months	-0.047 [0.068] N=944	-0.111 [0.073] N=836	-0.093 [0.079] N=737	-0.059 [0.087] N=589	-0.126 [0.118] N=443
WHO 2007 BMI thinness below -2, 61+ months	**0.094** [0.047] N=944**	**0.100** [0.050] N=836**	0.057 [0.055] N=737	0.034 [0.064] N=589	**0.134* [0.073] N=443**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 83 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.