Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 0-6 Using 12 November 2022 Child Age

The child cohort uses complete birth dates and completed ages 0 to 83 months on 12 November 2022 without age-boundary assignments. WHO 2006 covers months 0-60 and WHO 2007 covers months 61 onward. Weight-for-length/height classifications are available through month 60 only. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.348	0.207 [0.158] N=2,706	0.222 [0.166] N=2,496	0.262 [0.179] N=2,231	**0.398** [0.181] N=1,921**	**0.364** [0.205] N=1,587**
WHO weight-for-age z score	0.098	-0.043 [0.096] N=2,998	-0.019 [0.105] N=2,767	0.035 [0.104] N=2,476	-0.024 [0.114] N=2,125	-0.027 [0.121] N=1,752
WHO weight-for-length/height z score, 0-60 months	0.342	0.051 [0.157] N=2,029	0.029 [0.160] N=1,884	0.004 [0.169] N=1,697	-0.160 [0.186] N=1,454	-0.082 [0.199] N=1,191
WHO BMI-for-age z score	0.346	0.012 [0.147] N=2,626	-0.003 [0.150] N=2,425	0.007 [0.164] N=2,163	-0.250 [0.168] N=1,861	-0.222 [0.182] N=1,537
Stunted: HAZ below -2	0.219 P=573	-0.025 [0.033] N=2,706	-0.036 [0.035] N=2,496	-0.040 [0.036] N=2,231	**-0.078** [0.039] N=1,921**	-0.061 [0.044] N=1,587
Underweight: WAZ below -2	0.072 P=211	0.020 [0.019] N=2,998	0.012 [0.020] N=2,767	0.000 [0.021] N=2,476	0.000 [0.023] N=2,125	0.001 [0.026] N=1,752
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.024 [0.027] N=2,029	-0.016 [0.027] N=1,884	-0.013 [0.030] N=1,697	-0.001 [0.032] N=1,454	-0.018 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.020 [0.037] N=2,029	0.015 [0.040] N=1,884	0.016 [0.041] N=1,697	-0.010 [0.047] N=1,454	-0.021 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.009 [0.028] N=2,029	-0.012 [0.030] N=1,884	0.001 [0.031] N=1,697	-0.011 [0.035] N=1,454	-0.009 [0.041] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.041 [0.039] N=1,969	-0.047 [0.041] N=1,827	-0.042 [0.043] N=1,645	-0.076 [0.046] N=1,410	-0.057 [0.050] N=1,156
BMI-for-age z score below -2	0.116 P=310	-0.006 [0.026] N=2,626	-0.009 [0.026] N=2,425	-0.010 [0.029] N=2,163	0.012 [0.030] N=1,861	0.010 [0.033] N=1,537
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=228	-0.031 [0.073] N=597	-0.040 [0.078] N=542	-0.033 [0.082] N=472	-0.090 [0.087] N=414	-0.142 [0.101] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=138	-0.024 [0.061] N=597	-0.044 [0.064] N=542	-0.018 [0.073] N=472	-0.082 [0.075] N=414	**-0.166** [0.083] N=348**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=61	0.031 [0.070] N=597	0.021 [0.074] N=542	-0.010 [0.078] N=472	0.018 [0.081] N=414	-0.005 [0.104] N=348

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.085 [0.216] N=2,270	-0.118 [0.234] N=2,060	-0.092 [0.270] N=1,795	0.147 [0.274] N=1,485	-0.084 [0.362] N=1,151
WHO weight-for-age z score	-0.203 [0.135] N=2,521	-0.177 [0.156] N=2,290	-0.088 [0.161] N=1,999	-0.245 [0.195] N=1,648	-0.317 [0.245] N=1,275
WHO weight-for-length/height z score, 0-60 months	0.078 [0.232] N=1,707	0.043 [0.246] N=1,562	-0.032 [0.275] N=1,375	-0.354 [0.338] N=1,132	-0.255 [0.412] N=869
WHO BMI-for-age z score	0.164 [0.220] N=2,200	0.188 [0.232] N=1,999	0.254 [0.278] N=1,737	-0.191 [0.314] N=1,435	-0.049 [0.399] N=1,111
Stunted: HAZ below -2	0.042 [0.045] N=2,270	0.034 [0.050] N=2,060	0.035 [0.054] N=1,795	-0.024 [0.063] N=1,485	0.056 [0.084] N=1,151
Underweight: WAZ below -2	0.035 [0.026] N=2,521	0.021 [0.029] N=2,290	-0.004 [0.031] N=1,999	0.000 [0.037] N=1,648	0.004 [0.050] N=1,275
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.037] N=1,707	-0.008 [0.040] N=1,562	0.002 [0.046] N=1,375	0.023 [0.056] N=1,132	-0.020 [0.066] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.044] N=1,707	0.014 [0.048] N=1,562	0.012 [0.052] N=1,375	-0.038 [0.065] N=1,132	-0.097 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.031] N=1,707	-0.024 [0.034] N=1,562	-0.004 [0.036] N=1,375	-0.030 [0.045] N=1,132	-0.042 [0.063] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.006 [0.053] N=1,656	-0.001 [0.060] N=1,514	0.016 [0.064] N=1,332	-0.050 [0.076] N=1,097	-0.004 [0.093] N=843
BMI-for-age z score below -2	-0.022 [0.034] N=2,200	-0.035 [0.033] N=1,999	-0.045 [0.039] N=1,737	-0.015 [0.043] N=1,435	-0.039 [0.054] N=1,111
WHO 2007 BMI overweight above +1, 61+ months	0.044 [0.120] N=491	0.071 [0.130] N=436	0.114 [0.138] N=366	0.051 [0.162] N=308	-0.018 [0.221] N=242
WHO 2007 BMI obesity above +2, 61+ months	0.080 [0.093] N=491	0.082 [0.101] N=436	0.165 [0.116] N=366	0.093 [0.132] N=308	-0.045 [0.158] N=242
WHO 2007 BMI thinness below -2, 61+ months	-0.005 [0.066] N=491	-0.039 [0.071] N=436	-0.102 [0.079] N=366	-0.059 [0.089] N=308	-0.191 [0.142] N=242

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.348	0.209 [0.158] N=2,706	0.224 [0.166] N=2,496	0.266 [0.179] N=2,231	**0.403** [0.179] N=1,921**	**0.369* [0.203] N=1,587**
WHO weight-for-age z score	0.098	-0.043 [0.096] N=2,998	-0.020 [0.105] N=2,767	0.035 [0.104] N=2,476	-0.022 [0.114] N=2,125	-0.027 [0.121] N=1,752
WHO weight-for-length/height z score, 0-60 months	0.342	0.049 [0.157] N=2,029	0.027 [0.161] N=1,884	0.001 [0.170] N=1,697	-0.161 [0.187] N=1,454	-0.090 [0.200] N=1,191
WHO BMI-for-age z score	0.346	0.014 [0.148] N=2,626	-0.002 [0.150] N=2,425	0.007 [0.165] N=2,163	-0.248 [0.169] N=1,861	-0.222 [0.182] N=1,537
Stunted: HAZ below -2	0.219 P=573	-0.025 [0.033] N=2,706	-0.035 [0.035] N=2,496	-0.039 [0.036] N=2,231	**-0.078** [0.039] N=1,921**	-0.061 [0.044] N=1,587
Underweight: WAZ below -2	0.072 P=211	0.021 [0.019] N=2,998	0.012 [0.020] N=2,767	0.001 [0.021] N=2,476	0.000 [0.023] N=2,125	0.001 [0.026] N=1,752
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.021 [0.026] N=2,029	-0.013 [0.027] N=1,884	-0.009 [0.030] N=1,697	0.002 [0.032] N=1,454	-0.012 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.021 [0.037] N=2,029	0.017 [0.039] N=1,884	0.018 [0.041] N=1,697	-0.007 [0.047] N=1,454	-0.017 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.007 [0.028] N=2,029	-0.009 [0.030] N=1,884	0.003 [0.031] N=1,697	-0.009 [0.035] N=1,454	-0.006 [0.040] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.038 [0.038] N=1,969	-0.042 [0.040] N=1,827	-0.037 [0.042] N=1,645	-0.071 [0.045] N=1,410	-0.050 [0.049] N=1,156
BMI-for-age z score below -2	0.116 P=310	-0.005 [0.026] N=2,626	-0.008 [0.026] N=2,425	-0.009 [0.028] N=2,163	0.013 [0.030] N=1,861	0.011 [0.033] N=1,537
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=228	-0.032 [0.073] N=597	-0.042 [0.078] N=542	-0.035 [0.082] N=472	-0.096 [0.087] N=414	-0.148 [0.102] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=138	-0.026 [0.062] N=597	-0.047 [0.065] N=542	-0.021 [0.074] N=472	-0.091 [0.077] N=414	**-0.178** [0.085] N=348**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=61	0.033 [0.070] N=597	0.022 [0.074] N=542	-0.009 [0.078] N=472	0.020 [0.080] N=414	-0.004 [0.101] N=348

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.088 [0.215] N=2,270	-0.120 [0.233] N=2,060	-0.093 [0.268] N=1,795	0.132 [0.273] N=1,485	-0.131 [0.364] N=1,151
WHO weight-for-age z score	-0.203 [0.135] N=2,521	-0.177 [0.156] N=2,290	-0.088 [0.161] N=1,999	-0.249 [0.194] N=1,648	-0.324 [0.243] N=1,275
WHO weight-for-length/height z score, 0-60 months	0.078 [0.232] N=1,707	0.041 [0.247] N=1,562	-0.035 [0.276] N=1,375	-0.356 [0.339] N=1,132	-0.262 [0.412] N=869
WHO BMI-for-age z score	0.164 [0.220] N=2,200	0.188 [0.232] N=1,999	0.253 [0.278] N=1,737	-0.193 [0.314] N=1,435	-0.042 [0.399] N=1,111
Stunted: HAZ below -2	0.042 [0.045] N=2,270	0.034 [0.050] N=2,060	0.035 [0.054] N=1,795	-0.024 [0.063] N=1,485	0.060 [0.084] N=1,151
Underweight: WAZ below -2	0.035 [0.026] N=2,521	0.021 [0.029] N=2,290	-0.004 [0.031] N=1,999	0.001 [0.037] N=1,648	0.004 [0.050] N=1,275
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.037] N=1,707	-0.007 [0.040] N=1,562	0.003 [0.046] N=1,375	0.024 [0.057] N=1,132	-0.017 [0.066] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.044] N=1,707	0.014 [0.047] N=1,562	0.012 [0.052] N=1,375	-0.037 [0.065] N=1,132	-0.096 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.031] N=1,707	-0.024 [0.035] N=1,562	-0.003 [0.037] N=1,375	-0.030 [0.045] N=1,132	-0.041 [0.064] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.005 [0.053] N=1,656	-0.001 [0.059] N=1,514	0.017 [0.063] N=1,332	-0.050 [0.075] N=1,097	-0.003 [0.093] N=843
BMI-for-age z score below -2	-0.022 [0.034] N=2,200	-0.035 [0.033] N=1,999	-0.045 [0.039] N=1,737	-0.016 [0.043] N=1,435	-0.041 [0.054] N=1,111
WHO 2007 BMI overweight above +1, 61+ months	0.044 [0.121] N=491	0.070 [0.131] N=436	0.110 [0.140] N=366	0.040 [0.164] N=308	-0.014 [0.224] N=242
WHO 2007 BMI obesity above +2, 61+ months	0.078 [0.092] N=491	0.077 [0.100] N=436	0.156 [0.116] N=366	0.066 [0.131] N=308	-0.089 [0.157] N=242
WHO 2007 BMI thinness below -2, 61+ months	-0.002 [0.065] N=491	-0.035 [0.070] N=436	-0.096 [0.078] N=366	-0.042 [0.085] N=308	-0.156 [0.134] N=242

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.348	0.195 [0.161] N=2,706	0.210 [0.168] N=2,496	0.248 [0.183] N=2,231	**0.387** [0.185] N=1,921**	0.334 [0.209] N=1,587
WHO weight-for-age z score	0.098	-0.073 [0.093] N=2,998	-0.046 [0.103] N=2,767	0.003 [0.103] N=2,476	-0.049 [0.113] N=2,125	-0.059 [0.120] N=1,752
WHO weight-for-length/height z score, 0-60 months	0.342	0.043 [0.160] N=2,029	0.027 [0.164] N=1,884	0.006 [0.172] N=1,697	-0.155 [0.189] N=1,454	-0.089 [0.203] N=1,191
WHO BMI-for-age z score	0.346	0.009 [0.148] N=2,626	-0.004 [0.151] N=2,425	0.006 [0.165] N=2,163	-0.246 [0.169] N=1,861	-0.221 [0.185] N=1,537
Stunted: HAZ below -2	0.219 P=573	-0.021 [0.033] N=2,706	-0.031 [0.035] N=2,496	-0.034 [0.037] N=2,231	**-0.073* [0.039] N=1,921**	-0.052 [0.045] N=1,587
Underweight: WAZ below -2	0.072 P=211	0.026 [0.018] N=2,998	0.018 [0.019] N=2,767	0.007 [0.020] N=2,476	0.007 [0.023] N=2,125	0.010 [0.025] N=1,752
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.018 [0.026] N=2,029	-0.009 [0.027] N=1,884	-0.006 [0.030] N=1,697	0.005 [0.032] N=1,454	-0.009 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.024 [0.038] N=2,029	0.021 [0.039] N=1,884	0.022 [0.041] N=1,697	-0.003 [0.047] N=1,454	-0.013 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.007 [0.028] N=2,029	-0.008 [0.030] N=1,884	0.005 [0.031] N=1,697	-0.007 [0.035] N=1,454	-0.006 [0.041] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.026 [0.037] N=1,969	-0.027 [0.039] N=1,827	-0.025 [0.041] N=1,645	-0.062 [0.044] N=1,410	-0.035 [0.049] N=1,156
BMI-for-age z score below -2	0.116 P=310	-0.003 [0.026] N=2,626	-0.006 [0.026] N=2,425	-0.006 [0.029] N=2,163	0.015 [0.030] N=1,861	0.013 [0.033] N=1,537
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=228	-0.032 [0.074] N=597	-0.040 [0.078] N=542	-0.032 [0.083] N=472	-0.094 [0.089] N=414	-0.147 [0.105] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=138	-0.028 [0.062] N=597	-0.050 [0.066] N=542	-0.023 [0.075] N=472	-0.091 [0.078] N=414	**-0.174** [0.086] N=348**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=61	0.031 [0.071] N=597	0.019 [0.074] N=542	-0.008 [0.078] N=472	0.021 [0.080] N=414	-0.003 [0.102] N=348

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.099 [0.216] N=2,270	-0.137 [0.232] N=2,060	-0.118 [0.268] N=1,795	0.100 [0.276] N=1,485	-0.205 [0.359] N=1,151
WHO weight-for-age z score	**-0.231* [0.131] N=2,521**	-0.202 [0.153] N=2,290	-0.120 [0.159] N=1,999	-0.277 [0.190] N=1,648	-0.362 [0.239] N=1,275
WHO weight-for-length/height z score, 0-60 months	0.071 [0.235] N=1,707	0.044 [0.253] N=1,562	-0.026 [0.277] N=1,375	-0.356 [0.339] N=1,132	-0.275 [0.414] N=869
WHO BMI-for-age z score	0.160 [0.222] N=2,200	0.188 [0.234] N=1,999	0.259 [0.278] N=1,737	-0.183 [0.313] N=1,435	-0.024 [0.401] N=1,111
Stunted: HAZ below -2	0.045 [0.044] N=2,270	0.038 [0.049] N=2,060	0.040 [0.053] N=1,795	-0.021 [0.062] N=1,485	0.075 [0.081] N=1,151
Underweight: WAZ below -2	0.040 [0.025] N=2,521	0.025 [0.029] N=2,290	0.002 [0.030] N=1,999	0.005 [0.037] N=1,648	0.013 [0.049] N=1,275
Wasted: WLZ/WHZ below -2, 0-60 months	-0.016 [0.038] N=1,707	-0.001 [0.041] N=1,562	0.006 [0.047] N=1,375	0.028 [0.057] N=1,132	-0.011 [0.068] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.025 [0.045] N=1,707	0.020 [0.049] N=1,562	0.016 [0.053] N=1,375	-0.035 [0.065] N=1,132	-0.088 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.016 [0.031] N=1,707	-0.022 [0.035] N=1,562	-0.002 [0.037] N=1,375	-0.030 [0.045] N=1,132	-0.044 [0.062] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.019 [0.051] N=1,656	0.018 [0.057] N=1,514	0.026 [0.060] N=1,332	-0.048 [0.072] N=1,097	0.011 [0.087] N=843
BMI-for-age z score below -2	-0.020 [0.034] N=2,200	-0.033 [0.033] N=1,999	-0.043 [0.039] N=1,737	-0.016 [0.043] N=1,435	-0.037 [0.055] N=1,111
WHO 2007 BMI overweight above +1, 61+ months	0.046 [0.119] N=491	0.080 [0.131] N=436	0.113 [0.139] N=366	0.036 [0.164] N=308	-0.012 [0.223] N=242
WHO 2007 BMI obesity above +2, 61+ months	0.077 [0.091] N=491	0.080 [0.099] N=436	0.153 [0.113] N=366	0.059 [0.129] N=308	-0.086 [0.158] N=242
WHO 2007 BMI thinness below -2, 61+ months	-0.000 [0.064] N=491	-0.037 [0.070] N=436	-0.093 [0.078] N=366	-0.030 [0.085] N=308	-0.156 [0.139] N=242

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.348	0.177 [0.159] N=2,706	0.198 [0.167] N=2,496	0.239 [0.182] N=2,231	**0.365* [0.184] N=1,921**	0.313 [0.207] N=1,587
WHO weight-for-age z score	0.098	-0.074 [0.093] N=2,998	-0.056 [0.103] N=2,767	-0.007 [0.104] N=2,476	-0.067 [0.114] N=2,125	-0.084 [0.122] N=1,752
WHO weight-for-length/height z score, 0-60 months	0.342	0.057 [0.159] N=2,029	0.033 [0.162] N=1,884	0.009 [0.171] N=1,697	-0.151 [0.189] N=1,454	-0.092 [0.201] N=1,191
WHO BMI-for-age z score	0.346	0.021 [0.149] N=2,626	0.002 [0.152] N=2,425	0.012 [0.166] N=2,163	-0.238 [0.172] N=1,861	-0.214 [0.187] N=1,537
Stunted: HAZ below -2	0.219 P=573	-0.017 [0.033] N=2,706	-0.027 [0.035] N=2,496	-0.030 [0.037] N=2,231	**-0.066* [0.039] N=1,921**	-0.046 [0.044] N=1,587
Underweight: WAZ below -2	0.072 P=211	0.026 [0.018] N=2,998	0.020 [0.019] N=2,767	0.010 [0.020] N=2,476	0.011 [0.022] N=2,125	0.014 [0.025] N=1,752
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=229	-0.018 [0.026] N=2,029	-0.009 [0.027] N=1,884	-0.005 [0.030] N=1,697	0.006 [0.032] N=1,454	-0.010 [0.033] N=1,191
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=394	0.027 [0.037] N=2,029	0.024 [0.039] N=1,884	0.024 [0.041] N=1,697	-0.000 [0.047] N=1,454	-0.014 [0.053] N=1,191
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=186	-0.006 [0.028] N=2,029	-0.010 [0.029] N=1,884	0.003 [0.031] N=1,697	-0.007 [0.034] N=1,454	-0.007 [0.040] N=1,191
CIAF: any stunting, underweight, or wasting, 0-60 months	0.313 P=605	-0.020 [0.037] N=1,969	-0.021 [0.039] N=1,827	-0.016 [0.041] N=1,645	-0.049 [0.044] N=1,410	-0.024 [0.048] N=1,156
BMI-for-age z score below -2	0.116 P=310	-0.003 [0.026] N=2,626	-0.007 [0.026] N=2,425	-0.007 [0.029] N=2,163	0.015 [0.030] N=1,861	0.013 [0.034] N=1,537
WHO 2007 BMI overweight above +1, 61+ months	0.372 P=228	-0.033 [0.073] N=597	-0.044 [0.078] N=542	-0.036 [0.083] N=472	-0.094 [0.089] N=414	-0.144 [0.105] N=348
WHO 2007 BMI obesity above +2, 61+ months	0.226 P=138	-0.031 [0.063] N=597	-0.053 [0.066] N=542	-0.028 [0.075] N=472	-0.095 [0.079] N=414	**-0.171* [0.088] N=348**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=61	0.031 [0.071] N=597	0.022 [0.073] N=542	-0.010 [0.077] N=472	0.017 [0.079] N=414	0.004 [0.093] N=348

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.106 [0.213] N=2,270	-0.135 [0.229] N=2,060	-0.110 [0.265] N=1,795	0.114 [0.274] N=1,485	-0.214 [0.348] N=1,151
WHO weight-for-age z score	**-0.218* [0.131] N=2,521**	-0.201 [0.153] N=2,290	-0.123 [0.160] N=1,999	-0.274 [0.194] N=1,648	-0.346 [0.238] N=1,275
WHO weight-for-length/height z score, 0-60 months	0.079 [0.234] N=1,707	0.047 [0.249] N=1,562	-0.028 [0.279] N=1,375	-0.352 [0.343] N=1,132	-0.252 [0.415] N=869
WHO BMI-for-age z score	0.170 [0.221] N=2,200	0.192 [0.233] N=1,999	0.259 [0.279] N=1,737	-0.185 [0.316] N=1,435	-0.014 [0.401] N=1,111
Stunted: HAZ below -2	0.046 [0.044] N=2,270	0.038 [0.049] N=2,060	0.040 [0.053] N=1,795	-0.023 [0.061] N=1,485	0.072 [0.079] N=1,151
Underweight: WAZ below -2	0.038 [0.025] N=2,521	0.025 [0.029] N=2,290	0.004 [0.030] N=1,999	0.007 [0.037] N=1,648	0.015 [0.049] N=1,275
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.038] N=1,707	-0.004 [0.040] N=1,562	0.006 [0.047] N=1,375	0.029 [0.058] N=1,132	-0.010 [0.069] N=869
Overweight: WLZ/WHZ above +2, 0-60 months	0.026 [0.045] N=1,707	0.021 [0.048] N=1,562	0.016 [0.053] N=1,375	-0.035 [0.065] N=1,132	-0.087 [0.085] N=869
Obese: WLZ/WHZ above +3, 0-60 months	-0.014 [0.031] N=1,707	-0.021 [0.035] N=1,562	-0.001 [0.037] N=1,375	-0.029 [0.045] N=1,132	-0.042 [0.061] N=869
CIAF: any stunting, underweight, or wasting, 0-60 months	0.020 [0.051] N=1,656	0.015 [0.057] N=1,514	0.027 [0.061] N=1,332	-0.049 [0.072] N=1,097	0.011 [0.088] N=843
BMI-for-age z score below -2	-0.020 [0.034] N=2,200	-0.034 [0.033] N=1,999	-0.044 [0.039] N=1,737	-0.015 [0.043] N=1,435	-0.038 [0.055] N=1,111
WHO 2007 BMI overweight above +1, 61+ months	0.045 [0.117] N=491	0.071 [0.129] N=436	0.102 [0.137] N=366	0.023 [0.161] N=308	-0.005 [0.229] N=242
WHO 2007 BMI obesity above +2, 61+ months	0.066 [0.092] N=491	0.066 [0.098] N=436	0.133 [0.112] N=366	0.039 [0.129] N=308	-0.077 [0.162] N=242
WHO 2007 BMI thinness below -2, 61+ months	-0.000 [0.066] N=491	-0.025 [0.072] N=436	-0.088 [0.078] N=366	-0.032 [0.086] N=308	-0.112 [0.128] N=242

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	0.187 [0.159] N=2,528	0.178 [0.169] N=2,338	0.198 [0.182] N=2,084	**0.303* [0.181] N=1,802**	0.233 [0.200] N=1,484
WHO weight-for-age z score	0.096	-0.013 [0.099] N=2,796	0.000 [0.107] N=2,586	0.058 [0.109] N=2,308	-0.011 [0.119] N=1,988	-0.005 [0.126] N=1,637
WHO weight-for-length/height z score, 0-60 months	0.350	0.171 [0.160] N=1,893	0.147 [0.163] N=1,765	0.130 [0.171] N=1,587	-0.027 [0.192] N=1,370	0.081 [0.214] N=1,120
WHO BMI-for-age z score	0.353	0.104 [0.151] N=2,455	0.079 [0.152] N=2,273	0.115 [0.167] N=2,024	-0.128 [0.174] N=1,750	-0.076 [0.191] N=1,442
Stunted: HAZ below -2	0.220 P=536	-0.020 [0.034] N=2,528	-0.027 [0.036] N=2,338	-0.030 [0.037] N=2,084	-0.061 [0.039] N=1,802	-0.034 [0.044] N=1,484
Underweight: WAZ below -2	0.072 P=198	0.017 [0.020] N=2,796	0.010 [0.021] N=2,586	-0.002 [0.022] N=2,308	0.001 [0.025] N=1,988	0.005 [0.029] N=1,637
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=213	-0.034 [0.027] N=1,893	-0.025 [0.027] N=1,765	-0.019 [0.029] N=1,587	-0.007 [0.032] N=1,370	-0.021 [0.033] N=1,120
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=368	0.033 [0.039] N=1,893	0.029 [0.041] N=1,765	0.034 [0.043] N=1,587	0.007 [0.049] N=1,370	0.006 [0.058] N=1,120
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=179	-0.007 [0.030] N=1,893	-0.007 [0.032] N=1,765	0.010 [0.033] N=1,587	-0.001 [0.036] N=1,370	0.005 [0.043] N=1,120
CIAF: any stunting, underweight, or wasting, 0-60 months	0.315 P=566	-0.039 [0.041] N=1,836	-0.041 [0.043] N=1,711	-0.038 [0.045] N=1,537	-0.074 [0.049] N=1,328	-0.046 [0.053] N=1,086
BMI-for-age z score below -2	0.115 P=285	-0.010 [0.027] N=2,455	-0.011 [0.027] N=2,273	-0.014 [0.030] N=2,024	0.006 [0.032] N=1,750	-0.001 [0.035] N=1,442
WHO 2007 BMI overweight above +1, 61+ months	0.363 P=211	-0.033 [0.079] N=563	-0.036 [0.084] N=511	-0.026 [0.090] N=444	-0.099 [0.094] N=388	-0.154 [0.114] N=325
WHO 2007 BMI obesity above +2, 61+ months	0.218 P=127	-0.042 [0.066] N=563	-0.053 [0.069] N=511	-0.023 [0.081] N=444	-0.103 [0.083] N=388	**-0.199** [0.096] N=325**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=57	0.032 [0.072] N=563	0.023 [0.077] N=511	-0.009 [0.081] N=444	0.022 [0.085] N=388	-0.019 [0.106] N=325

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.006 [0.227] N=2,121	-0.066 [0.251] N=1,931	-0.057 [0.288] N=1,677	0.146 [0.304] N=1,395	-0.129 [0.412] N=1,077
WHO weight-for-age z score	-0.180 [0.137] N=2,351	-0.175 [0.153] N=2,141	-0.094 [0.163] N=1,863	-0.284 [0.196] N=1,543	-0.326 [0.240] N=1,192
WHO weight-for-length/height z score, 0-60 months	0.107 [0.239] N=1,591	0.045 [0.251] N=1,463	-0.020 [0.278] N=1,285	-0.326 [0.347] N=1,068	-0.130 [0.425] N=818
WHO BMI-for-age z score	0.165 [0.228] N=2,057	0.162 [0.238] N=1,875	0.258 [0.283] N=1,626	-0.169 [0.322] N=1,352	0.026 [0.408] N=1,044
Stunted: HAZ below -2	0.025 [0.047] N=2,121	0.016 [0.052] N=1,931	0.018 [0.057] N=1,677	-0.039 [0.068] N=1,395	0.058 [0.091] N=1,077
Underweight: WAZ below -2	0.031 [0.027] N=2,351	0.018 [0.030] N=2,141	-0.003 [0.033] N=1,863	0.011 [0.039] N=1,543	0.022 [0.053] N=1,192
Wasted: WLZ/WHZ below -2, 0-60 months	-0.036 [0.038] N=1,591	-0.021 [0.040] N=1,463	-0.008 [0.046] N=1,285	0.007 [0.056] N=1,068	-0.047 [0.061] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.045] N=1,591	0.010 [0.048] N=1,463	0.015 [0.052] N=1,285	-0.040 [0.065] N=1,068	-0.071 [0.090] N=818
Obese: WLZ/WHZ above +3, 0-60 months	-0.028 [0.034] N=1,591	-0.034 [0.038] N=1,463	-0.007 [0.040] N=1,285	-0.030 [0.048] N=1,068	-0.032 [0.067] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.012 [0.056] N=1,543	-0.020 [0.062] N=1,418	-0.008 [0.068] N=1,244	-0.091 [0.080] N=1,035	-0.040 [0.102] N=793
BMI-for-age z score below -2	-0.016 [0.034] N=2,057	-0.026 [0.032] N=1,875	-0.039 [0.037] N=1,626	-0.015 [0.042] N=1,352	-0.055 [0.051] N=1,044
WHO 2007 BMI overweight above +1, 61+ months	0.044 [0.130] N=464	0.087 [0.139] N=412	0.125 [0.152] N=345	0.032 [0.174] N=289	-0.031 [0.247] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.042 [0.095] N=464	0.061 [0.101] N=412	0.142 [0.119] N=345	0.038 [0.127] N=289	-0.151 [0.171] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.016 [0.069] N=464	-0.021 [0.076] N=412	-0.088 [0.083] N=345	-0.043 [0.093] N=289	-0.195 [0.137] N=226

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.353	0.192 [0.158] N=2,528	0.181 [0.167] N=2,338	0.199 [0.181] N=2,084	**0.310* [0.181] N=1,802**	0.254 [0.197] N=1,484
WHO weight-for-age z score	0.096	-0.007 [0.098] N=2,796	0.004 [0.106] N=2,586	0.061 [0.108] N=2,308	-0.005 [0.118] N=1,988	0.007 [0.126] N=1,637
WHO weight-for-length/height z score, 0-60 months	0.350	0.169 [0.160] N=1,893	0.146 [0.162] N=1,765	0.123 [0.170] N=1,587	-0.033 [0.191] N=1,370	0.065 [0.213] N=1,120
WHO BMI-for-age z score	0.353	0.106 [0.151] N=2,455	0.080 [0.152] N=2,273	0.116 [0.166] N=2,024	-0.128 [0.174] N=1,750	-0.083 [0.192] N=1,442
Stunted: HAZ below -2	0.220 P=536	-0.021 [0.033] N=2,528	-0.028 [0.035] N=2,338	-0.031 [0.037] N=2,084	-0.063 [0.039] N=1,802	-0.039 [0.044] N=1,484
Underweight: WAZ below -2	0.072 P=198	0.015 [0.020] N=2,796	0.008 [0.021] N=2,586	-0.003 [0.022] N=2,308	-0.001 [0.025] N=1,988	0.001 [0.029] N=1,637
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=213	-0.035 [0.027] N=1,893	-0.025 [0.027] N=1,765	-0.019 [0.029] N=1,587	-0.008 [0.032] N=1,370	-0.024 [0.033] N=1,120
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=368	0.033 [0.039] N=1,893	0.029 [0.040] N=1,765	0.031 [0.043] N=1,587	0.003 [0.049] N=1,370	0.000 [0.057] N=1,120
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=179	-0.007 [0.029] N=1,893	-0.007 [0.031] N=1,765	0.009 [0.033] N=1,587	-0.002 [0.036] N=1,370	0.003 [0.043] N=1,120
CIAF: any stunting, underweight, or wasting, 0-60 months	0.315 P=566	-0.039 [0.041] N=1,836	-0.041 [0.043] N=1,711	-0.037 [0.045] N=1,537	-0.073 [0.049] N=1,328	-0.045 [0.052] N=1,086
BMI-for-age z score below -2	0.115 P=285	-0.010 [0.027] N=2,455	-0.011 [0.027] N=2,273	-0.014 [0.030] N=2,024	0.006 [0.032] N=1,750	-0.000 [0.036] N=1,442
WHO 2007 BMI overweight above +1, 61+ months	0.363 P=211	-0.027 [0.078] N=563	-0.032 [0.083] N=511	-0.026 [0.088] N=444	-0.097 [0.092] N=388	-0.156 [0.113] N=325
WHO 2007 BMI obesity above +2, 61+ months	0.218 P=127	-0.042 [0.065] N=563	-0.056 [0.068] N=511	-0.023 [0.080] N=444	-0.104 [0.082] N=388	**-0.190** [0.094] N=325**
WHO 2007 BMI thinness below -2, 61+ months	0.104 P=57	0.029 [0.073] N=563	0.021 [0.077] N=511	-0.011 [0.083] N=444	0.021 [0.086] N=388	-0.008 [0.109] N=325

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.004 [0.226] N=2,121	-0.070 [0.250] N=1,931	-0.059 [0.288] N=1,677	0.149 [0.307] N=1,395	-0.079 [0.409] N=1,077
WHO weight-for-age z score	-0.170 [0.136] N=2,351	-0.170 [0.152] N=2,141	-0.088 [0.160] N=1,863	-0.271 [0.194] N=1,543	-0.291 [0.236] N=1,192
WHO weight-for-length/height z score, 0-60 months	0.132 [0.239] N=1,591	0.078 [0.251] N=1,463	-0.014 [0.279] N=1,285	-0.322 [0.351] N=1,068	-0.146 [0.433] N=818
WHO BMI-for-age z score	0.178 [0.227] N=2,057	0.172 [0.236] N=1,875	0.262 [0.281] N=1,626	-0.166 [0.319] N=1,352	0.015 [0.407] N=1,044
Stunted: HAZ below -2	0.025 [0.047] N=2,121	0.018 [0.052] N=1,931	0.017 [0.057] N=1,677	-0.043 [0.067] N=1,395	0.043 [0.089] N=1,077
Underweight: WAZ below -2	0.029 [0.027] N=2,351	0.017 [0.030] N=2,141	-0.005 [0.032] N=1,863	0.007 [0.038] N=1,543	0.012 [0.052] N=1,192
Wasted: WLZ/WHZ below -2, 0-60 months	-0.037 [0.038] N=1,591	-0.022 [0.040] N=1,463	-0.008 [0.046] N=1,285	0.005 [0.056] N=1,068	-0.052 [0.062] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.030 [0.045] N=1,591	0.020 [0.048] N=1,463	0.017 [0.052] N=1,285	-0.037 [0.067] N=1,068	-0.073 [0.091] N=818
Obese: WLZ/WHZ above +3, 0-60 months	-0.026 [0.033] N=1,591	-0.032 [0.037] N=1,463	-0.006 [0.039] N=1,285	-0.031 [0.048] N=1,068	-0.035 [0.067] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.012 [0.056] N=1,543	-0.020 [0.063] N=1,418	-0.009 [0.068] N=1,244	-0.098 [0.080] N=1,035	-0.060 [0.099] N=793
BMI-for-age z score below -2	-0.017 [0.034] N=2,057	-0.027 [0.032] N=1,875	-0.038 [0.037] N=1,626	-0.015 [0.042] N=1,352	-0.054 [0.052] N=1,044
WHO 2007 BMI overweight above +1, 61+ months	0.053 [0.126] N=464	0.093 [0.137] N=412	0.130 [0.148] N=345	0.046 [0.168] N=289	-0.026 [0.247] N=226
WHO 2007 BMI obesity above +2, 61+ months	0.041 [0.092] N=464	0.055 [0.099] N=412	0.138 [0.115] N=345	0.027 [0.125] N=289	-0.147 [0.171] N=226
WHO 2007 BMI thinness below -2, 61+ months	0.004 [0.068] N=464	-0.028 [0.075] N=412	-0.096 [0.084] N=345	-0.042 [0.091] N=289	-0.190 [0.143] N=226

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.271	-0.258 [0.180] N=2,705	-0.286 [0.192] N=2,452	**-0.334*** **[0.200]** **N=2,164**	**-0.413*** **[0.215]** **N=1,845**	**-0.488**** **[0.243]** **N=1,499**
WHO weight-for-age z score	0.150	-0.074 [0.098] N=2,967	-0.042 [0.105] N=2,702	-0.131 [0.111] N=2,378	-0.102 [0.120] N=2,021	-0.089 [0.135] N=1,634
WHO weight-for-length/height z score, 0-60 months	0.376	0.091 [0.182] N=1,886	0.119 [0.189] N=1,707	0.079 [0.210] N=1,489	0.200 [0.223] N=1,268	0.172 [0.264] N=1,022
WHO BMI-for-age z score	0.374	0.112 [0.172] N=2,624	0.133 [0.179] N=2,380	0.145 [0.197] N=2,102	0.264 [0.209] N=1,791	0.243 [0.234] N=1,453
Stunted: HAZ below -2	0.209 P=562	0.044 [0.034] N=2,705	0.044 [0.036] N=2,452	0.044 [0.038] N=2,164	0.063 [0.042] N=1,845	0.067 [0.049] N=1,499
Underweight: WAZ below -2	0.066 P=191	-0.004 [0.017] N=2,967	-0.005 [0.018] N=2,702	0.002 [0.018] N=2,378	0.007 [0.018] N=2,021	0.010 [0.019] N=1,634
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.025 [0.029] N=1,886	0.020 [0.030] N=1,707	0.035 [0.032] N=1,489	-0.003 [0.032] N=1,268	0.002 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.047 [0.035] N=1,886	0.048 [0.036] N=1,707	0.043 [0.039] N=1,489	0.058 [0.042] N=1,268	0.061 [0.047] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.018 [0.029] N=1,886	0.014 [0.030] N=1,707	0.011 [0.033] N=1,489	0.024 [0.036] N=1,268	0.010 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.073*** **[0.041]** **N=1,830**	0.064 [0.043] N=1,654	**0.077*** **[0.045]** **N=1,441**	0.044 [0.046] N=1,227	0.059 [0.051] N=989
BMI-for-age z score below -2	0.115 P=302	0.034 [0.029] N=2,624	0.032 [0.031] N=2,380	0.027 [0.034] N=2,102	0.009 [0.036] N=1,791	0.018 [0.041] N=1,453
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=281	-0.002 [0.069] N=741	-0.039 [0.071] N=675	-0.041 [0.076] N=614	-0.035 [0.082] N=525	-0.040 [0.090] N=429
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=166	0.018 [0.062] N=741	-0.006 [0.066] N=675	-0.003 [0.071] N=614	0.021 [0.075] N=525	0.020 [0.087] N=429
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=73	0.036 [0.047] N=741	0.041 [0.050] N=675	0.024 [0.051] N=614	0.011 [0.055] N=525	-0.000 [0.064] N=429

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.050 [0.228] N=2,319	-0.046 [0.254] N=2,066	-0.090 [0.269] N=1,778	-0.254 [0.307] N=1,459	-0.288 [0.408] N=1,113
WHO weight-for-age z score	-0.169 [0.136] N=2,550	-0.119 [0.150] N=2,285	**-0.328** [0.158] N=1,961**	**-0.356** [0.175] N=1,604**	**-0.404* [0.221] N=1,217**
WHO weight-for-length/height z score, 0-60 months	-0.165 [0.261] N=1,630	-0.164 [0.274] N=1,451	-0.323 [0.322] N=1,233	-0.191 [0.354] N=1,012	-0.566 [0.429] N=766
WHO BMI-for-age z score	-0.300 [0.198] N=2,253	-0.337 [0.206] N=2,009	-0.398 [0.245] N=1,731	-0.294 [0.275] N=1,420	**-0.633** [0.315] N=1,082**
Stunted: HAZ below -2	0.021 [0.043] N=2,319	0.016 [0.048] N=2,066	0.005 [0.051] N=1,778	0.045 [0.056] N=1,459	0.034 [0.079] N=1,113
Underweight: WAZ below -2	-0.003 [0.026] N=2,550	-0.005 [0.030] N=2,285	0.008 [0.032] N=1,961	0.022 [0.036] N=1,604	0.035 [0.041] N=1,217
Wasted: WLZ/WHZ below -2, 0-60 months	**0.081** [0.039] N=1,630**	**0.079* [0.042] N=1,451**	**0.125*** [0.047] N=1,233**	0.067 [0.048] N=1,012	**0.117** [0.055] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.036 [0.052] N=1,630	0.030 [0.056] N=1,451	0.016 [0.063] N=1,233	0.040 [0.071] N=1,012	0.006 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.013 [0.040] N=1,630	0.001 [0.041] N=1,451	-0.010 [0.047] N=1,233	0.008 [0.059] N=1,012	-0.081 [0.077] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.115* [0.063] N=1,582**	0.105 [0.070] N=1,406	**0.145* [0.080] N=1,193**	0.098 [0.088] N=979	0.140 [0.115] N=741
BMI-for-age z score below -2	**0.093*** [0.027] N=2,253**	**0.099*** [0.031] N=2,009**	**0.101*** [0.036] N=1,731**	**0.083** [0.034] N=1,420**	**0.138*** [0.041] N=1,082**
WHO 2007 BMI overweight above +1, 61+ months	-0.074 [0.099] N=623	-0.153 [0.107] N=557	-0.191 [0.118] N=496	-0.224 [0.138] N=407	**-0.326* [0.179] N=311**
WHO 2007 BMI obesity above +2, 61+ months	-0.058 [0.079] N=623	-0.119 [0.083] N=557	-0.149 [0.094] N=496	-0.140 [0.099] N=407	-0.201 [0.150] N=311
WHO 2007 BMI thinness below -2, 61+ months	**0.098* [0.054] N=623**	**0.126** [0.057] N=557**	**0.103* [0.061] N=496**	0.104 [0.065] N=407	**0.132* [0.075] N=311**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.271	-0.258 [0.180] N=2,705	-0.287 [0.192] N=2,452	**-0.335* [0.201] N=2,164**	**-0.415* [0.216] N=1,845**	**-0.481* [0.243] N=1,499**
WHO weight-for-age z score	0.150	-0.074 [0.098] N=2,967	-0.041 [0.105] N=2,702	-0.131 [0.112] N=2,378	-0.102 [0.120] N=2,021	-0.087 [0.136] N=1,634
WHO weight-for-length/height z score, 0-60 months	0.376	0.097 [0.181] N=1,886	0.122 [0.188] N=1,707	0.082 [0.210] N=1,489	0.201 [0.223] N=1,268	0.169 [0.261] N=1,022
WHO BMI-for-age z score	0.374	0.112 [0.172] N=2,624	0.133 [0.179] N=2,380	0.146 [0.197] N=2,102	0.264 [0.209] N=1,791	0.243 [0.234] N=1,453
Stunted: HAZ below -2	0.209 P=562	0.044 [0.034] N=2,705	0.044 [0.036] N=2,452	0.044 [0.038] N=2,164	0.063 [0.042] N=1,845	0.067 [0.049] N=1,499
Underweight: WAZ below -2	0.066 P=191	-0.003 [0.017] N=2,967	-0.005 [0.018] N=2,702	0.002 [0.018] N=2,378	0.007 [0.018] N=2,021	0.011 [0.019] N=1,634
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.028] N=1,886	0.018 [0.030] N=1,707	0.034 [0.032] N=1,489	-0.004 [0.032] N=1,268	0.003 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.046 [0.035] N=1,886	0.047 [0.037] N=1,707	0.041 [0.039] N=1,489	0.057 [0.042] N=1,268	0.062 [0.048] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.016 [0.029] N=1,886	0.012 [0.030] N=1,707	0.009 [0.033] N=1,489	0.023 [0.037] N=1,268	0.011 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.071* [0.042] N=1,830**	0.060 [0.043] N=1,654	0.074 [0.046] N=1,441	0.043 [0.047] N=1,227	0.060 [0.051] N=989
BMI-for-age z score below -2	0.115 P=302	0.035 [0.029] N=2,624	0.032 [0.031] N=2,380	0.027 [0.034] N=2,102	0.009 [0.036] N=1,791	0.019 [0.041] N=1,453
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=281	0.008 [0.070] N=741	-0.030 [0.071] N=675	-0.035 [0.075] N=614	-0.033 [0.082] N=525	-0.033 [0.091] N=429
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=166	0.028 [0.063] N=741	0.004 [0.067] N=675	0.004 [0.072] N=614	0.023 [0.077] N=525	0.024 [0.089] N=429
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=73	0.036 [0.047] N=741	0.042 [0.050] N=675	0.023 [0.051] N=614	0.011 [0.056] N=525	-0.001 [0.064] N=429

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.052 [0.228] N=2,319	-0.051 [0.254] N=2,066	-0.097 [0.269] N=1,778	-0.268 [0.309] N=1,459	-0.292 [0.408] N=1,113
WHO weight-for-age z score	-0.168 [0.136] N=2,550	-0.120 [0.150] N=2,285	**-0.329** [0.158] N=1,961**	**-0.359** [0.176] N=1,604**	**-0.401* [0.221] N=1,217**
WHO weight-for-length/height z score, 0-60 months	-0.147 [0.263] N=1,630	-0.152 [0.277] N=1,451	-0.315 [0.325] N=1,233	-0.181 [0.356] N=1,012	-0.538 [0.436] N=766
WHO BMI-for-age z score	-0.300 [0.198] N=2,253	-0.338 [0.207] N=2,009	-0.400 [0.246] N=1,731	-0.295 [0.275] N=1,420	**-0.633** [0.316] N=1,082**
Stunted: HAZ below -2	0.021 [0.044] N=2,319	0.016 [0.048] N=2,066	0.004 [0.051] N=1,778	0.044 [0.057] N=1,459	0.034 [0.079] N=1,113
Underweight: WAZ below -2	-0.003 [0.026] N=2,550	-0.005 [0.030] N=2,285	0.008 [0.032] N=1,961	0.021 [0.035] N=1,604	0.035 [0.041] N=1,217
Wasted: WLZ/WHZ below -2, 0-60 months	**0.074* [0.039] N=1,630**	**0.071* [0.042] N=1,451**	**0.116** [0.047] N=1,233**	0.057 [0.049] N=1,012	**0.106* [0.056] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.035 [0.052] N=1,630	0.026 [0.056] N=1,451	0.010 [0.063] N=1,233	0.032 [0.071] N=1,012	0.002 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.009 [0.040] N=1,630	-0.005 [0.041] N=1,451	-0.018 [0.047] N=1,233	-0.001 [0.059] N=1,012	-0.087 [0.077] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.107* [0.064] N=1,582**	0.093 [0.070] N=1,406	0.131 [0.081] N=1,193	0.085 [0.090] N=979	0.129 [0.116] N=741
BMI-for-age z score below -2	**0.093*** [0.028] N=2,253**	**0.098*** [0.031] N=2,009**	**0.100*** [0.036] N=1,731**	**0.080** [0.035] N=1,420**	**0.138*** [0.041] N=1,082**
WHO 2007 BMI overweight above +1, 61+ months	-0.053 [0.099] N=623	-0.132 [0.106] N=557	-0.177 [0.117] N=496	-0.215 [0.138] N=407	-0.296 [0.183] N=311
WHO 2007 BMI obesity above +2, 61+ months	-0.037 [0.082] N=623	-0.097 [0.086] N=557	-0.132 [0.096] N=496	-0.128 [0.102] N=407	-0.179 [0.156] N=311
WHO 2007 BMI thinness below -2, 61+ months	**0.105* [0.055] N=623**	**0.134** [0.059] N=557**	**0.105* [0.062] N=496**	0.109 [0.066] N=407	**0.137* [0.077] N=311**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.271	-0.254 [0.176] N=2,705	-0.275 [0.187] N=2,452	-0.319 [0.197] N=2,164	**-0.396* [0.209] N=1,845**	**-0.461* [0.234] N=1,499**
WHO weight-for-age z score	0.150	-0.073 [0.098] N=2,967	-0.038 [0.105] N=2,702	-0.123 [0.112] N=2,378	-0.100 [0.121] N=2,021	-0.085 [0.137] N=1,634
WHO weight-for-length/height z score, 0-60 months	0.376	0.105 [0.183] N=1,886	0.127 [0.191] N=1,707	0.093 [0.213] N=1,489	0.202 [0.223] N=1,268	0.168 [0.260] N=1,022
WHO BMI-for-age z score	0.374	0.107 [0.171] N=2,624	0.128 [0.179] N=2,380	0.143 [0.195] N=2,102	0.254 [0.208] N=1,791	0.232 [0.234] N=1,453
Stunted: HAZ below -2	0.209 P=562	0.043 [0.033] N=2,705	0.041 [0.035] N=2,452	0.039 [0.037] N=2,164	0.058 [0.041] N=1,845	0.061 [0.047] N=1,499
Underweight: WAZ below -2	0.066 P=191	-0.003 [0.017] N=2,967	-0.005 [0.018] N=2,702	-0.000 [0.018] N=2,378	0.006 [0.018] N=2,021	0.009 [0.019] N=1,634
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.029] N=1,886	0.016 [0.030] N=1,707	0.031 [0.033] N=1,489	-0.005 [0.033] N=1,268	0.002 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.048 [0.035] N=1,886	0.048 [0.037] N=1,707	0.042 [0.040] N=1,489	0.056 [0.043] N=1,268	0.062 [0.048] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.018 [0.030] N=1,886	0.013 [0.030] N=1,707	0.009 [0.033] N=1,489	0.022 [0.037] N=1,268	0.010 [0.041] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.075* [0.041] N=1,830**	0.060 [0.043] N=1,654	0.070 [0.046] N=1,441	0.040 [0.046] N=1,227	0.057 [0.050] N=989
BMI-for-age z score below -2	0.115 P=302	0.034 [0.029] N=2,624	0.031 [0.031] N=2,380	0.027 [0.034] N=2,102	0.008 [0.036] N=1,791	0.018 [0.041] N=1,453
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=281	0.003 [0.070] N=741	-0.031 [0.072] N=675	-0.038 [0.076] N=614	-0.040 [0.083] N=525	-0.030 [0.091] N=429
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=166	0.019 [0.063] N=741	-0.002 [0.067] N=675	-0.004 [0.072] N=614	0.010 [0.077] N=525	0.014 [0.088] N=429
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=73	0.035 [0.046] N=741	0.038 [0.048] N=675	0.025 [0.050] N=614	0.011 [0.055] N=525	-0.009 [0.062] N=429

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.039 [0.226] N=2,319	-0.028 [0.251] N=2,066	-0.062 [0.265] N=1,778	-0.219 [0.300] N=1,459	-0.235 [0.390] N=1,113
WHO weight-for-age z score	-0.167 [0.136] N=2,550	-0.117 [0.151] N=2,285	**-0.313* [0.161] N=1,961**	**-0.354** [0.176] N=1,604**	**-0.399* [0.222] N=1,217**
WHO weight-for-length/height z score, 0-60 months	-0.150 [0.264] N=1,630	-0.154 [0.280] N=1,451	-0.312 [0.326] N=1,233	-0.196 [0.353] N=1,012	-0.551 [0.430] N=766
WHO BMI-for-age z score	-0.309 [0.197] N=2,253	**-0.345* [0.207] N=2,009**	**-0.407* [0.243] N=1,731**	-0.318 [0.273] N=1,420	**-0.661** [0.315] N=1,082**
Stunted: HAZ below -2	0.018 [0.043] N=2,319	0.011 [0.047] N=2,066	-0.003 [0.051] N=1,778	0.034 [0.056] N=1,459	0.022 [0.078] N=1,113
Underweight: WAZ below -2	-0.004 [0.026] N=2,550	-0.006 [0.029] N=2,285	0.003 [0.032] N=1,961	0.018 [0.036] N=1,604	0.031 [0.042] N=1,217
Wasted: WLZ/WHZ below -2, 0-60 months	**0.075* [0.040] N=1,630**	0.069 [0.043] N=1,451	**0.112** [0.048] N=1,233**	0.054 [0.050] N=1,012	**0.105* [0.057] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.037 [0.053] N=1,630	0.028 [0.057] N=1,451	0.007 [0.064] N=1,233	0.026 [0.073] N=1,012	0.000 [0.091] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.010 [0.040] N=1,630	-0.006 [0.042] N=1,451	-0.019 [0.048] N=1,233	-0.004 [0.060] N=1,012	-0.089 [0.078] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.111* [0.063] N=1,582**	0.092 [0.071] N=1,406	0.119 [0.082] N=1,193	0.072 [0.090] N=979	0.123 [0.112] N=741
BMI-for-age z score below -2	**0.091*** [0.028] N=2,253**	**0.095*** [0.031] N=2,009**	**0.097*** [0.036] N=1,731**	**0.078** [0.036] N=1,420**	**0.133*** [0.042] N=1,082**
WHO 2007 BMI overweight above +1, 61+ months	-0.065 [0.099] N=623	-0.138 [0.108] N=557	-0.184 [0.118] N=496	-0.228 [0.142] N=407	-0.312 [0.189] N=311
WHO 2007 BMI obesity above +2, 61+ months	-0.047 [0.081] N=623	-0.102 [0.087] N=557	-0.137 [0.095] N=496	-0.138 [0.101] N=407	-0.199 [0.153] N=311
WHO 2007 BMI thinness below -2, 61+ months	**0.105* [0.053] N=623**	**0.130** [0.057] N=557**	**0.111* [0.062] N=496**	**0.113* [0.066] N=407**	0.125 [0.078] N=311

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.271	-0.249 [0.175] N=2,705	-0.265 [0.187] N=2,452	-0.309 [0.198] N=2,164	**-0.380* [0.209] N=1,845**	**-0.447* [0.234] N=1,499**
WHO weight-for-age z score	0.150	-0.073 [0.096] N=2,967	-0.037 [0.103] N=2,702	-0.120 [0.109] N=2,378	-0.089 [0.117] N=2,021	-0.084 [0.132] N=1,634
WHO weight-for-length/height z score, 0-60 months	0.376	0.107 [0.182] N=1,886	0.130 [0.188] N=1,707	0.091 [0.210] N=1,489	0.214 [0.222] N=1,268	0.182 [0.260] N=1,022
WHO BMI-for-age z score	0.374	0.106 [0.171] N=2,624	0.127 [0.179] N=2,380	0.139 [0.197] N=2,102	0.258 [0.209] N=1,791	0.229 [0.234] N=1,453
Stunted: HAZ below -2	0.209 P=562	0.043 [0.033] N=2,705	0.041 [0.035] N=2,452	0.039 [0.038] N=2,164	0.057 [0.041] N=1,845	0.060 [0.048] N=1,499
Underweight: WAZ below -2	0.066 P=191	-0.004 [0.017] N=2,967	-0.006 [0.018] N=2,702	-0.001 [0.018] N=2,378	0.005 [0.018] N=2,021	0.008 [0.019] N=1,634
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=209	0.023 [0.029] N=1,886	0.017 [0.030] N=1,707	0.034 [0.032] N=1,489	-0.004 [0.032] N=1,268	0.001 [0.037] N=1,022
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=378	0.047 [0.035] N=1,886	0.047 [0.036] N=1,707	0.042 [0.039] N=1,489	0.058 [0.042] N=1,268	0.063 [0.047] N=1,022
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=188	0.020 [0.029] N=1,886	0.015 [0.030] N=1,707	0.012 [0.033] N=1,489	0.027 [0.036] N=1,268	0.014 [0.040] N=1,022
CIAF: any stunting, underweight, or wasting, 0-60 months	0.310 P=563	**0.074* [0.041] N=1,830**	0.060 [0.043] N=1,654	0.070 [0.045] N=1,441	0.039 [0.046] N=1,227	0.060 [0.050] N=989
BMI-for-age z score below -2	0.115 P=302	0.033 [0.029] N=2,624	0.030 [0.031] N=2,380	0.026 [0.034] N=2,102	0.006 [0.035] N=1,791	0.014 [0.041] N=1,453
WHO 2007 BMI overweight above +1, 61+ months	0.379 P=281	0.000 [0.071] N=741	-0.036 [0.073] N=675	-0.040 [0.078] N=614	-0.038 [0.084] N=525	-0.035 [0.093] N=429
WHO 2007 BMI obesity above +2, 61+ months	0.227 P=166	0.013 [0.063] N=741	-0.013 [0.067] N=675	-0.013 [0.072] N=614	-0.001 [0.076] N=525	-0.010 [0.087] N=429
WHO 2007 BMI thinness below -2, 61+ months	0.099 P=73	0.031 [0.046] N=741	0.036 [0.048] N=675	0.020 [0.049] N=614	0.004 [0.053] N=525	-0.022 [0.060] N=429

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.045 [0.223] N=2,319	-0.032 [0.250] N=2,066	-0.055 [0.265] N=1,778	-0.205 [0.297] N=1,459	-0.238 [0.385] N=1,113
WHO weight-for-age z score	-0.153 [0.133] N=2,550	-0.100 [0.147] N=2,285	**-0.288* [0.158] N=1,961**	**-0.305* [0.171] N=1,604**	**-0.356* [0.211] N=1,217**
WHO weight-for-length/height z score, 0-60 months	-0.141 [0.262] N=1,630	-0.151 [0.274] N=1,451	-0.322 [0.317] N=1,233	-0.192 [0.347] N=1,012	-0.533 [0.430] N=766
WHO BMI-for-age z score	-0.303 [0.197] N=2,253	**-0.341* [0.206] N=2,009**	**-0.413* [0.244] N=1,731**	-0.307 [0.272] N=1,420	**-0.648** [0.316] N=1,082**
Stunted: HAZ below -2	0.019 [0.043] N=2,319	0.011 [0.047] N=2,066	-0.005 [0.050] N=1,778	0.030 [0.055] N=1,459	0.023 [0.077] N=1,113
Underweight: WAZ below -2	-0.006 [0.026] N=2,550	-0.009 [0.029] N=2,285	0.001 [0.033] N=1,961	0.013 [0.036] N=1,604	0.026 [0.043] N=1,217
Wasted: WLZ/WHZ below -2, 0-60 months	**0.075* [0.039] N=1,630**	**0.071* [0.042] N=1,451**	**0.118** [0.047] N=1,233**	0.058 [0.049] N=1,012	**0.109* [0.055] N=766**
Overweight: WLZ/WHZ above +2, 0-60 months	0.035 [0.052] N=1,630	0.024 [0.055] N=1,451	0.005 [0.062] N=1,233	0.024 [0.071] N=1,012	0.001 [0.090] N=766
Obese: WLZ/WHZ above +3, 0-60 months	0.012 [0.040] N=1,630	-0.004 [0.041] N=1,451	-0.017 [0.047] N=1,233	0.001 [0.059] N=1,012	-0.086 [0.078] N=766
CIAF: any stunting, underweight, or wasting, 0-60 months	**0.108* [0.062] N=1,582**	0.088 [0.069] N=1,406	0.113 [0.080] N=1,193	0.064 [0.087] N=979	0.126 [0.107] N=741
BMI-for-age z score below -2	**0.093*** [0.028] N=2,253**	**0.097*** [0.031] N=2,009**	**0.099*** [0.036] N=1,731**	**0.079** [0.035] N=1,420**	**0.138*** [0.041] N=1,082**
WHO 2007 BMI overweight above +1, 61+ months	-0.073 [0.099] N=623	-0.146 [0.108] N=557	-0.195 [0.118] N=496	**-0.235* [0.138] N=407**	**-0.324* [0.183] N=311**
WHO 2007 BMI obesity above +2, 61+ months	-0.045 [0.080] N=623	-0.105 [0.085] N=557	-0.139 [0.093] N=496	-0.135 [0.097] N=407	-0.196 [0.149] N=311
WHO 2007 BMI thinness below -2, 61+ months	**0.100* [0.054] N=623**	**0.131** [0.057] N=557**	**0.108* [0.061] N=496**	**0.111* [0.067] N=407**	0.115 [0.075] N=311

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.282	-0.256 [0.183] N=2,536	-0.263 [0.197] N=2,300	-0.300 [0.204] N=2,032	**-0.373* [0.219] N=1,727**	**-0.482* [0.247] N=1,405**
WHO weight-for-age z score	0.154	-0.056 [0.100] N=2,777	-0.015 [0.108] N=2,529	-0.100 [0.114] N=2,229	-0.072 [0.123] N=1,891	-0.088 [0.137] N=1,533
WHO weight-for-length/height z score, 0-60 months	0.378	0.136 [0.192] N=1,762	0.176 [0.200] N=1,595	0.099 [0.221] N=1,392	0.197 [0.233] N=1,184	0.133 [0.270] N=957
WHO BMI-for-age z score	0.390	0.092 [0.177] N=2,462	0.116 [0.184] N=2,234	0.104 [0.202] N=1,975	0.211 [0.216] N=1,679	0.163 [0.244] N=1,365
Stunted: HAZ below -2	0.209 P=528	0.048 [0.033] N=2,536	0.049 [0.035] N=2,300	0.048 [0.035] N=2,032	**0.064* [0.038] N=1,727**	**0.073* [0.043] N=1,405**
Underweight: WAZ below -2	0.066 P=179	-0.006 [0.017] N=2,777	-0.008 [0.018] N=2,529	-0.006 [0.018] N=2,229	0.004 [0.018] N=1,891	0.014 [0.019] N=1,533
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=196	0.026 [0.029] N=1,762	0.017 [0.030] N=1,595	0.036 [0.032] N=1,392	0.003 [0.033] N=1,184	0.012 [0.037] N=957
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.046 [0.035] N=1,762	0.048 [0.037] N=1,595	0.036 [0.038] N=1,392	0.047 [0.041] N=1,184	0.046 [0.047] N=957
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=180	0.016 [0.029] N=1,762	0.012 [0.030] N=1,595	0.007 [0.032] N=1,392	0.023 [0.036] N=1,184	0.013 [0.039] N=957
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=529	**0.087** [0.043] N=1,709**	0.074 [0.046] N=1,545	**0.088* [0.047] N=1,347**	0.051 [0.048] N=1,145	0.074 [0.052] N=925
BMI-for-age z score below -2	0.112 P=275	0.040 [0.029] N=2,462	0.038 [0.030] N=2,234	0.035 [0.033] N=1,975	0.021 [0.034] N=1,679	0.033 [0.038] N=1,365
WHO 2007 BMI overweight above +1, 61+ months	0.384 P=270	-0.013 [0.071] N=704	-0.051 [0.073] N=642	-0.043 [0.077] N=584	-0.042 [0.085] N=497	-0.034 [0.095] N=406
WHO 2007 BMI obesity above +2, 61+ months	0.234 P=162	0.006 [0.067] N=704	-0.017 [0.070] N=642	-0.017 [0.076] N=584	0.003 [0.081] N=497	0.013 [0.096] N=406
WHO 2007 BMI thinness below -2, 61+ months	0.094 P=65	0.044 [0.047] N=704	0.052 [0.049] N=642	0.034 [0.051] N=584	0.021 [0.055] N=497	0.014 [0.062] N=406

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.008 [0.230] N=2,174	0.036 [0.258] N=1,938	-0.001 [0.273] N=1,670	-0.125 [0.306] N=1,365	-0.162 [0.396] N=1,043
WHO weight-for-age z score	-0.160 [0.139] N=2,387	-0.098 [0.154] N=2,139	**-0.315* [0.163] N=1,839**	**-0.328* [0.178] N=1,501**	**-0.396* [0.218] N=1,143**
WHO weight-for-length/height z score, 0-60 months	-0.108 [0.276] N=1,522	-0.102 [0.294] N=1,355	-0.334 [0.343] N=1,152	-0.232 [0.369] N=944	-0.609 [0.456] N=717
WHO BMI-for-age z score	-0.306 [0.200] N=2,112	-0.340 [0.210] N=1,884	**-0.430* [0.248] N=1,625**	-0.331 [0.268] N=1,329	**-0.723** [0.300] N=1,015**
Stunted: HAZ below -2	0.012 [0.046] N=2,174	0.007 [0.050] N=1,938	-0.004 [0.053] N=1,670	0.023 [0.056] N=1,365	0.010 [0.078] N=1,043
Underweight: WAZ below -2	-0.010 [0.026] N=2,387	-0.012 [0.030] N=2,139	-0.006 [0.032] N=1,839	0.013 [0.036] N=1,501	0.030 [0.042] N=1,143
Wasted: WLZ/WHZ below -2, 0-60 months	**0.077* [0.041] N=1,522**	0.068 [0.044] N=1,355	**0.115** [0.048] N=1,152**	0.060 [0.053] N=944	**0.105* [0.058] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.054] N=1,522	0.022 [0.057] N=1,355	-0.004 [0.063] N=1,152	0.004 [0.072] N=944	-0.022 [0.093] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.000 [0.042] N=1,522	-0.016 [0.044] N=1,355	-0.033 [0.049] N=1,152	-0.014 [0.062] N=944	-0.089 [0.083] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.111 [0.069] N=1,477	0.094 [0.077] N=1,313	0.128 [0.087] N=1,115	0.053 [0.100] N=913	0.091 [0.130] N=693
BMI-for-age z score below -2	**0.096*** [0.029] N=2,112**	**0.102*** [0.032] N=1,884**	**0.105*** [0.038] N=1,625**	**0.092** [0.037] N=1,329**	**0.151*** [0.041] N=1,015**
WHO 2007 BMI overweight above +1, 61+ months	-0.079 [0.099] N=592	-0.162 [0.106] N=530	-0.181 [0.115] N=472	-0.211 [0.136] N=385	-0.274 [0.184] N=294
WHO 2007 BMI obesity above +2, 61+ months	-0.061 [0.085] N=592	-0.118 [0.089] N=530	-0.150 [0.097] N=472	-0.135 [0.104] N=385	-0.185 [0.158] N=294
WHO 2007 BMI thinness below -2, 61+ months	**0.108* [0.055] N=592**	**0.142** [0.058] N=530**	**0.112* [0.061] N=472**	0.105 [0.064] N=385	**0.126* [0.076] N=294**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	**Mean**	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO length/height-for-age z score	-0.282	-0.263 [0.182] N=2,536	-0.268 [0.196] N=2,300	-0.304 [0.203] N=2,032	**-0.373* [0.220] N=1,727**	**-0.473* [0.247] N=1,405**
WHO weight-for-age z score	0.154	-0.059 [0.100] N=2,777	-0.017 [0.108] N=2,529	-0.105 [0.114] N=2,229	-0.073 [0.123] N=1,891	-0.083 [0.138] N=1,533
WHO weight-for-length/height z score, 0-60 months	0.378	0.143 [0.192] N=1,762	0.179 [0.200] N=1,595	0.105 [0.221] N=1,392	0.213 [0.232] N=1,184	0.155 [0.272] N=957
WHO BMI-for-age z score	0.390	0.093 [0.176] N=2,462	0.118 [0.184] N=2,234	0.105 [0.202] N=1,975	0.211 [0.217] N=1,679	0.161 [0.244] N=1,365
Stunted: HAZ below -2	0.209 P=528	0.048 [0.033] N=2,536	0.048 [0.035] N=2,300	0.047 [0.036] N=2,032	**0.063* [0.038] N=1,727**	0.071 [0.043] N=1,405
Underweight: WAZ below -2	0.066 P=179	-0.006 [0.017] N=2,777	-0.008 [0.018] N=2,529	-0.005 [0.018] N=2,229	0.004 [0.018] N=1,891	0.013 [0.019] N=1,533
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=196	0.023 [0.029] N=1,762	0.015 [0.030] N=1,595	0.033 [0.032] N=1,392	-0.003 [0.033] N=1,184	0.005 [0.037] N=957
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=354	0.045 [0.035] N=1,762	0.046 [0.036] N=1,595	0.034 [0.038] N=1,392	0.045 [0.040] N=1,184	0.045 [0.046] N=957
Obese: WLZ/WHZ above +3, 0-60 months	0.098 P=180	0.015 [0.029] N=1,762	0.011 [0.029] N=1,595	0.005 [0.032] N=1,392	0.021 [0.035] N=1,184	0.011 [0.039] N=957
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=529	**0.081* [0.043] N=1,709**	0.069 [0.045] N=1,545	**0.082* [0.047] N=1,347**	0.043 [0.048] N=1,145	0.065 [0.051] N=925
BMI-for-age z score below -2	0.112 P=275	0.039 [0.028] N=2,462	0.037 [0.030] N=2,234	0.034 [0.033] N=1,975	0.019 [0.034] N=1,679	0.031 [0.038] N=1,365
WHO 2007 BMI overweight above +1, 61+ months	0.384 P=270	-0.015 [0.071] N=704	-0.054 [0.072] N=642	-0.050 [0.076] N=584	-0.046 [0.084] N=497	-0.034 [0.093] N=406
WHO 2007 BMI obesity above +2, 61+ months	0.234 P=162	0.008 [0.066] N=704	-0.017 [0.070] N=642	-0.016 [0.075] N=584	0.002 [0.079] N=497	0.015 [0.093] N=406
WHO 2007 BMI thinness below -2, 61+ months	0.094 P=65	0.040 [0.046] N=704	0.050 [0.048] N=642	0.025 [0.049] N=584	0.010 [0.053] N=497	-0.002 [0.059] N=406

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.031 [0.226] N=2,174	0.017 [0.253] N=1,938	-0.021 [0.267] N=1,670	-0.167 [0.306] N=1,365	-0.205 [0.394] N=1,043
WHO weight-for-age z score	-0.163 [0.138] N=2,387	-0.100 [0.154] N=2,139	**-0.321* [0.163] N=1,839**	**-0.332* [0.178] N=1,501**	**-0.414* [0.219] N=1,143**
WHO weight-for-length/height z score, 0-60 months	-0.108 [0.275] N=1,522	-0.101 [0.291] N=1,355	-0.330 [0.340] N=1,152	-0.201 [0.371] N=944	-0.604 [0.449] N=717
WHO BMI-for-age z score	-0.299 [0.199] N=2,112	-0.333 [0.209] N=1,884	**-0.427* [0.247] N=1,625**	-0.307 [0.271] N=1,329	**-0.706** [0.306] N=1,015**
Stunted: HAZ below -2	0.016 [0.045] N=2,174	0.011 [0.050] N=1,938	-0.000 [0.053] N=1,670	0.032 [0.057] N=1,365	0.024 [0.077] N=1,043
Underweight: WAZ below -2	-0.009 [0.026] N=2,387	-0.012 [0.030] N=2,139	-0.004 [0.032] N=1,839	0.015 [0.036] N=1,501	0.036 [0.042] N=1,143
Wasted: WLZ/WHZ below -2, 0-60 months	**0.076* [0.040] N=1,522**	0.069 [0.044] N=1,355	**0.116** [0.048] N=1,152**	0.055 [0.051] N=944	**0.106* [0.056] N=717**
Overweight: WLZ/WHZ above +2, 0-60 months	0.034 [0.053] N=1,522	0.022 [0.056] N=1,355	-0.004 [0.062] N=1,152	0.008 [0.071] N=944	-0.023 [0.090] N=717
Obese: WLZ/WHZ above +3, 0-60 months	0.001 [0.042] N=1,522	-0.015 [0.043] N=1,355	-0.033 [0.049] N=1,152	-0.013 [0.062] N=944	-0.091 [0.080] N=717
CIAF: any stunting, underweight, or wasting, 0-60 months	0.112 [0.068] N=1,477	0.097 [0.076] N=1,313	0.133 [0.086] N=1,115	0.063 [0.099] N=913	0.115 [0.127] N=693
BMI-for-age z score below -2	**0.095*** [0.029] N=2,112**	**0.102*** [0.032] N=1,884**	**0.103*** [0.037] N=1,625**	**0.087** [0.036] N=1,329**	**0.149*** [0.040] N=1,015**
WHO 2007 BMI overweight above +1, 61+ months	-0.077 [0.098] N=592	-0.157 [0.105] N=530	-0.181 [0.115] N=472	-0.206 [0.136] N=385	-0.258 [0.184] N=294
WHO 2007 BMI obesity above +2, 61+ months	-0.055 [0.084] N=592	-0.115 [0.088] N=530	-0.144 [0.098] N=472	-0.135 [0.103] N=385	-0.181 [0.158] N=294
WHO 2007 BMI thinness below -2, 61+ months	**0.105* [0.055] N=592**	**0.140** [0.058] N=530**	0.099 [0.061] N=472	0.092 [0.061] N=385	0.101 [0.074] N=294

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 0 to 83 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.