Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Oldest Child

Age reference: 10 October 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,527	-0.012 [0.021] N=1,618	0.006 [0.021] N=1,493	-0.004 [0.023] N=1,340	-0.012 [0.025] N=1,140	-0.015 [0.030] N=936
Vegetables consumed at least weekly	0.902 P=1,457	0.009 [0.027] N=1,618	0.014 [0.028] N=1,493	-0.008 [0.029] N=1,340	-0.015 [0.029] N=1,140	-0.049 [0.033] N=936
Meat fish or poultry consumed at least weekly	0.803 P=1,307	0.070 [0.044] N=1,618	**0.088* [0.046] N=1,493**	0.072 [0.050] N=1,340	0.056 [0.056] N=1,140	0.035 [0.062] N=936
Legumes consumed at least weekly	0.803 P=1,301	-0.058 [0.038] N=1,618	**-0.066* [0.040] N=1,493**	**-0.082** [0.040] N=1,340**	**-0.079* [0.045] N=1,140**	**-0.146*** [0.050] N=936**
Animal products consumed at least weekly	0.919 P=1,487	0.028 [0.028] N=1,618	0.027 [0.029] N=1,493	0.013 [0.032] N=1,340	0.018 [0.036] N=1,140	0.004 [0.039] N=936
Sweets consumed at least weekly	0.723 P=1,166	-0.018 [0.038] N=1,618	0.004 [0.040] N=1,493	0.017 [0.042] N=1,340	0.021 [0.045] N=1,140	0.031 [0.050] N=936
Snack foods consumed at least weekly	0.594 P=952	-0.037 [0.042] N=1,618	-0.021 [0.044] N=1,493	-0.015 [0.048] N=1,340	-0.023 [0.054] N=1,140	-0.047 [0.058] N=936
Packaged soft drinks consumed at least weekly	0.370 P=588	0.007 [0.050] N=1,618	0.022 [0.053] N=1,493	0.019 [0.058] N=1,340	0.042 [0.064] N=1,140	0.025 [0.074] N=936
Grains consumed at least weekly	0.928 P=1,499	0.002 [0.027] N=1,618	0.016 [0.029] N=1,493	-0.003 [0.030] N=1,340	0.006 [0.033] N=1,140	-0.012 [0.038] N=936
Child was ever breastfed	0.963 P=1,554	0.005 [0.017] N=1,619	0.009 [0.018] N=1,494	0.013 [0.020] N=1,340	0.013 [0.022] N=1,140	0.006 [0.024] N=936

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.032] N=1,378	0.033 [0.031] N=1,253	0.020 [0.033] N=1,100	0.005 [0.040] N=900	0.020 [0.054] N=696
Vegetables consumed at least weekly	0.051 [0.039] N=1,378	0.071 [0.043] N=1,253	0.041 [0.047] N=1,100	0.031 [0.056] N=900	-0.040 [0.069] N=696
Meat fish or poultry consumed at least weekly	0.064 [0.057] N=1,378	0.094 [0.058] N=1,253	0.068 [0.067] N=1,100	0.021 [0.078] N=900	-0.041 [0.090] N=696
Legumes consumed at least weekly	0.011 [0.058] N=1,378	0.007 [0.064] N=1,253	0.001 [0.067] N=1,100	0.031 [0.084] N=900	-0.101 [0.109] N=696
Animal products consumed at least weekly	0.039 [0.035] N=1,378	0.039 [0.037] N=1,253	0.013 [0.045] N=1,100	0.026 [0.057] N=900	-0.010 [0.074] N=696
Sweets consumed at least weekly	-0.048 [0.051] N=1,378	-0.015 [0.055] N=1,253	0.008 [0.062] N=1,100	0.015 [0.071] N=900	0.056 [0.093] N=696
Snack foods consumed at least weekly	-0.056 [0.056] N=1,378	-0.033 [0.061] N=1,253	-0.022 [0.073] N=1,100	-0.038 [0.091] N=900	-0.132 [0.101] N=696
Packaged soft drinks consumed at least weekly	-0.033 [0.060] N=1,378	-0.011 [0.065] N=1,253	-0.018 [0.073] N=1,100	0.009 [0.089] N=900	-0.056 [0.112] N=696
Grains consumed at least weekly	0.037 [0.043] N=1,378	0.072 [0.048] N=1,253	0.047 [0.053] N=1,100	0.091 [0.063] N=900	0.081 [0.092] N=696
Child was ever breastfed	0.017 [0.026] N=1,379	0.025 [0.029] N=1,254	0.035 [0.035] N=1,100	0.040 [0.046] N=900	0.048 [0.059] N=696

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,527	-0.012 [0.021] N=1,618	0.006 [0.021] N=1,493	-0.003 [0.023] N=1,340	-0.012 [0.025] N=1,140	-0.014 [0.030] N=936
Vegetables consumed at least weekly	0.902 P=1,457	0.009 [0.027] N=1,618	0.014 [0.028] N=1,493	-0.008 [0.029] N=1,340	-0.015 [0.029] N=1,140	-0.048 [0.033] N=936
Meat fish or poultry consumed at least weekly	0.803 P=1,307	0.070 [0.044] N=1,618	**0.088* [0.046] N=1,493**	0.072 [0.051] N=1,340	0.057 [0.056] N=1,140	0.036 [0.063] N=936
Legumes consumed at least weekly	0.803 P=1,301	-0.058 [0.038] N=1,618	**-0.066* [0.040] N=1,493**	**-0.082** [0.040] N=1,340**	**-0.079* [0.046] N=1,140**	**-0.146*** [0.050] N=936**
Animal products consumed at least weekly	0.919 P=1,487	0.027 [0.028] N=1,618	0.027 [0.029] N=1,493	0.013 [0.032] N=1,340	0.018 [0.036] N=1,140	0.005 [0.039] N=936
Sweets consumed at least weekly	0.723 P=1,166	-0.018 [0.038] N=1,618	0.004 [0.040] N=1,493	0.017 [0.042] N=1,340	0.022 [0.045] N=1,140	0.031 [0.050] N=936
Snack foods consumed at least weekly	0.594 P=952	-0.038 [0.043] N=1,618	-0.021 [0.045] N=1,493	-0.015 [0.049] N=1,340	-0.022 [0.054] N=1,140	-0.047 [0.058] N=936
Packaged soft drinks consumed at least weekly	0.370 P=588	0.007 [0.050] N=1,618	0.022 [0.053] N=1,493	0.019 [0.058] N=1,340	0.042 [0.064] N=1,140	0.024 [0.074] N=936
Grains consumed at least weekly	0.928 P=1,499	0.002 [0.027] N=1,618	0.016 [0.029] N=1,493	-0.003 [0.030] N=1,340	0.006 [0.032] N=1,140	-0.013 [0.038] N=936
Child was ever breastfed	0.963 P=1,554	0.005 [0.017] N=1,619	0.009 [0.018] N=1,494	0.014 [0.020] N=1,340	0.013 [0.022] N=1,140	0.006 [0.024] N=936

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.032] N=1,378	0.034 [0.031] N=1,253	0.021 [0.033] N=1,100	0.005 [0.040] N=900	0.022 [0.054] N=696
Vegetables consumed at least weekly	0.051 [0.039] N=1,378	0.071 [0.043] N=1,253	0.042 [0.047] N=1,100	0.033 [0.056] N=900	-0.037 [0.068] N=696
Meat fish or poultry consumed at least weekly	0.064 [0.057] N=1,378	0.093 [0.058] N=1,253	0.069 [0.067] N=1,100	0.023 [0.079] N=900	-0.035 [0.089] N=696
Legumes consumed at least weekly	0.011 [0.058] N=1,378	0.007 [0.064] N=1,253	0.002 [0.067] N=1,100	0.034 [0.084] N=900	-0.098 [0.109] N=696
Animal products consumed at least weekly	0.039 [0.035] N=1,378	0.042 [0.037] N=1,253	0.018 [0.045] N=1,100	0.031 [0.057] N=900	-0.006 [0.073] N=696
Sweets consumed at least weekly	-0.048 [0.051] N=1,378	-0.015 [0.055] N=1,253	0.007 [0.062] N=1,100	0.015 [0.070] N=900	0.055 [0.093] N=696
Snack foods consumed at least weekly	-0.055 [0.056] N=1,378	-0.030 [0.061] N=1,253	-0.017 [0.073] N=1,100	-0.032 [0.092] N=900	-0.125 [0.101] N=696
Packaged soft drinks consumed at least weekly	-0.033 [0.060] N=1,378	-0.011 [0.065] N=1,253	-0.019 [0.073] N=1,100	0.008 [0.089] N=900	-0.060 [0.112] N=696
Grains consumed at least weekly	0.037 [0.043] N=1,378	0.073 [0.048] N=1,253	0.047 [0.053] N=1,100	0.090 [0.063] N=900	0.079 [0.091] N=696
Child was ever breastfed	0.017 [0.026] N=1,379	0.026 [0.030] N=1,254	0.036 [0.035] N=1,100	0.040 [0.046] N=900	0.049 [0.059] N=696

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,527	-0.021 [0.022] N=1,618	-0.003 [0.022] N=1,493	-0.012 [0.023] N=1,340	-0.021 [0.026] N=1,140	-0.024 [0.031] N=936
Vegetables consumed at least weekly	0.902 P=1,457	0.002 [0.027] N=1,618	0.006 [0.028] N=1,493	-0.015 [0.029] N=1,340	-0.022 [0.030] N=1,140	**-0.059* [0.033] N=936**
Meat fish or poultry consumed at least weekly	0.803 P=1,307	0.046 [0.043] N=1,618	0.059 [0.045] N=1,493	0.047 [0.050] N=1,340	0.031 [0.055] N=1,140	0.006 [0.062] N=936
Legumes consumed at least weekly	0.803 P=1,301	**-0.066* [0.038] N=1,618**	**-0.076* [0.040] N=1,493**	**-0.091** [0.040] N=1,340**	**-0.086* [0.046] N=1,140**	**-0.152*** [0.051] N=936**
Animal products consumed at least weekly	0.919 P=1,487	0.022 [0.028] N=1,618	0.020 [0.029] N=1,493	0.007 [0.033] N=1,340	0.012 [0.037] N=1,140	-0.005 [0.040] N=936
Sweets consumed at least weekly	0.723 P=1,166	-0.025 [0.038] N=1,618	-0.005 [0.040] N=1,493	0.007 [0.042] N=1,340	0.012 [0.045] N=1,140	0.018 [0.050] N=936
Snack foods consumed at least weekly	0.594 P=952	-0.036 [0.042] N=1,618	-0.019 [0.045] N=1,493	-0.015 [0.047] N=1,340	-0.022 [0.053] N=1,140	-0.044 [0.058] N=936
Packaged soft drinks consumed at least weekly	0.370 P=588	0.019 [0.049] N=1,618	0.039 [0.052] N=1,493	0.033 [0.056] N=1,340	0.057 [0.063] N=1,140	0.044 [0.072] N=936
Grains consumed at least weekly	0.928 P=1,499	-0.006 [0.027] N=1,618	0.008 [0.029] N=1,493	-0.009 [0.030] N=1,340	-0.003 [0.032] N=1,140	-0.020 [0.037] N=936
Child was ever breastfed	0.963 P=1,554	0.000 [0.017] N=1,619	0.003 [0.018] N=1,494	0.007 [0.019] N=1,340	0.007 [0.022] N=1,140	0.001 [0.024] N=936

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.031] N=1,378	0.025 [0.030] N=1,253	0.015 [0.032] N=1,100	0.008 [0.039] N=900	0.027 [0.053] N=696
Vegetables consumed at least weekly	0.047 [0.039] N=1,378	0.065 [0.043] N=1,253	0.039 [0.047] N=1,100	0.037 [0.055] N=900	-0.034 [0.067] N=696
Meat fish or poultry consumed at least weekly	0.047 [0.056] N=1,378	0.066 [0.057] N=1,253	0.050 [0.067] N=1,100	0.008 [0.078] N=900	-0.049 [0.093] N=696
Legumes consumed at least weekly	0.005 [0.058] N=1,378	-0.004 [0.064] N=1,253	-0.006 [0.067] N=1,100	0.034 [0.085] N=900	-0.097 [0.110] N=696
Animal products consumed at least weekly	0.036 [0.035] N=1,378	0.036 [0.037] N=1,253	0.013 [0.046] N=1,100	0.032 [0.058] N=900	-0.004 [0.073] N=696
Sweets consumed at least weekly	-0.053 [0.052] N=1,378	-0.021 [0.055] N=1,253	0.001 [0.063] N=1,100	0.014 [0.070] N=900	0.056 [0.092] N=696
Snack foods consumed at least weekly	-0.056 [0.055] N=1,378	-0.030 [0.060] N=1,253	-0.020 [0.070] N=1,100	-0.033 [0.087] N=900	-0.118 [0.097] N=696
Packaged soft drinks consumed at least weekly	-0.023 [0.059] N=1,378	0.008 [0.064] N=1,253	-0.005 [0.071] N=1,100	0.024 [0.088] N=900	-0.040 [0.108] N=696
Grains consumed at least weekly	0.030 [0.042] N=1,378	0.067 [0.047] N=1,253	0.042 [0.053] N=1,100	0.085 [0.062] N=900	0.069 [0.089] N=696
Child was ever breastfed	0.014 [0.026] N=1,379	0.021 [0.030] N=1,254	0.032 [0.035] N=1,100	0.038 [0.047] N=900	0.043 [0.058] N=696

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,527	-0.022 [0.022] N=1,618	-0.005 [0.022] N=1,493	-0.015 [0.024] N=1,340	-0.023 [0.027] N=1,140	-0.024 [0.032] N=936
Vegetables consumed at least weekly	0.902 P=1,457	0.002 [0.027] N=1,618	0.006 [0.028] N=1,493	-0.015 [0.029] N=1,340	-0.023 [0.030] N=1,140	**-0.058* [0.034] N=936**
Meat fish or poultry consumed at least weekly	0.803 P=1,307	0.041 [0.043] N=1,618	0.052 [0.044] N=1,493	0.037 [0.049] N=1,340	0.022 [0.054] N=1,140	-0.006 [0.061] N=936
Legumes consumed at least weekly	0.803 P=1,301	**-0.067* [0.039] N=1,618**	**-0.079* [0.041] N=1,493**	**-0.098** [0.041] N=1,340**	**-0.089* [0.047] N=1,140**	**-0.155*** [0.052] N=936**
Animal products consumed at least weekly	0.919 P=1,487	0.020 [0.028] N=1,618	0.020 [0.029] N=1,493	0.006 [0.033] N=1,340	0.010 [0.037] N=1,140	-0.005 [0.040] N=936
Sweets consumed at least weekly	0.723 P=1,166	-0.024 [0.038] N=1,618	-0.006 [0.039] N=1,493	0.002 [0.041] N=1,340	0.010 [0.044] N=1,140	0.015 [0.050] N=936
Snack foods consumed at least weekly	0.594 P=952	-0.037 [0.042] N=1,618	-0.021 [0.044] N=1,493	-0.020 [0.048] N=1,340	-0.023 [0.054] N=1,140	-0.047 [0.058] N=936
Packaged soft drinks consumed at least weekly	0.370 P=588	0.026 [0.049] N=1,618	0.042 [0.052] N=1,493	0.037 [0.057] N=1,340	0.065 [0.063] N=1,140	0.057 [0.072] N=936
Grains consumed at least weekly	0.928 P=1,499	-0.006 [0.026] N=1,618	0.008 [0.028] N=1,493	-0.011 [0.030] N=1,340	-0.003 [0.032] N=1,140	-0.020 [0.039] N=936
Child was ever breastfed	0.963 P=1,554	-0.000 [0.017] N=1,619	0.003 [0.018] N=1,494	0.006 [0.020] N=1,340	0.006 [0.022] N=1,140	0.000 [0.024] N=936

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.009 [0.031] N=1,378	0.025 [0.031] N=1,253	0.014 [0.033] N=1,100	0.004 [0.040] N=900	0.020 [0.054] N=696
Vegetables consumed at least weekly	0.048 [0.039] N=1,378	0.065 [0.043] N=1,253	0.037 [0.047] N=1,100	0.032 [0.056] N=900	-0.047 [0.068] N=696
Meat fish or poultry consumed at least weekly	0.043 [0.057] N=1,378	0.064 [0.058] N=1,253	0.048 [0.068] N=1,100	0.012 [0.079] N=900	-0.056 [0.098] N=696
Legumes consumed at least weekly	0.010 [0.058] N=1,378	0.002 [0.064] N=1,253	-0.001 [0.065] N=1,100	0.051 [0.083] N=900	-0.088 [0.113] N=696
Animal products consumed at least weekly	0.034 [0.035] N=1,378	0.037 [0.038] N=1,253	0.014 [0.046] N=1,100	0.027 [0.058] N=900	-0.010 [0.076] N=696
Sweets consumed at least weekly	-0.048 [0.051] N=1,378	-0.019 [0.054] N=1,253	0.004 [0.059] N=1,100	0.032 [0.069] N=900	0.069 [0.092] N=696
Snack foods consumed at least weekly	-0.050 [0.055] N=1,378	-0.027 [0.059] N=1,253	-0.017 [0.068] N=1,100	-0.012 [0.087] N=900	-0.097 [0.095] N=696
Packaged soft drinks consumed at least weekly	-0.016 [0.059] N=1,378	0.007 [0.064] N=1,253	-0.004 [0.072] N=1,100	0.034 [0.088] N=900	-0.018 [0.107] N=696
Grains consumed at least weekly	0.029 [0.041] N=1,378	0.064 [0.045] N=1,253	0.040 [0.050] N=1,100	0.085 [0.059] N=900	0.065 [0.083] N=696
Child was ever breastfed	0.015 [0.027] N=1,379	0.022 [0.030] N=1,254	0.032 [0.036] N=1,100	0.039 [0.048] N=900	0.046 [0.060] N=696

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.944 P=1,425	-0.021 [0.023] N=1,511	-0.006 [0.023] N=1,401	-0.013 [0.024] N=1,255	-0.020 [0.027] N=1,071	-0.024 [0.031] N=881
Vegetables consumed at least weekly	0.903 P=1,362	-0.002 [0.028] N=1,511	0.000 [0.029] N=1,401	-0.023 [0.030] N=1,255	-0.026 [0.030] N=1,071	-0.056 [0.034] N=881
Meat fish or poultry consumed at least weekly	0.802 P=1,217	0.055 [0.044] N=1,511	0.062 [0.045] N=1,401	0.041 [0.049] N=1,255	0.027 [0.054] N=1,071	-0.002 [0.060] N=881
Legumes consumed at least weekly	0.798 P=1,208	**-0.070* [0.038] N=1,511**	**-0.082** [0.039] N=1,401**	**-0.099** [0.038] N=1,255**	**-0.089** [0.044] N=1,071**	**-0.157*** [0.049] N=881**
Animal products consumed at least weekly	0.919 P=1,390	0.028 [0.029] N=1,511	0.027 [0.030] N=1,401	0.014 [0.033] N=1,255	0.014 [0.037] N=1,071	0.003 [0.041] N=881
Sweets consumed at least weekly	0.724 P=1,088	0.000 [0.038] N=1,511	0.017 [0.040] N=1,401	0.021 [0.042] N=1,255	0.033 [0.045] N=1,071	0.041 [0.050] N=881
Snack foods consumed at least weekly	0.594 P=888	-0.039 [0.045] N=1,511	-0.021 [0.048] N=1,401	-0.020 [0.051] N=1,255	-0.023 [0.058] N=1,071	-0.031 [0.064] N=881
Packaged soft drinks consumed at least weekly	0.376 P=558	0.036 [0.049] N=1,511	0.053 [0.052] N=1,401	0.040 [0.057] N=1,255	0.063 [0.062] N=1,071	0.063 [0.072] N=881
Grains consumed at least weekly	0.927 P=1,399	-0.002 [0.028] N=1,511	0.010 [0.030] N=1,401	-0.013 [0.031] N=1,255	-0.001 [0.034] N=1,071	-0.016 [0.040] N=881
Child was ever breastfed	0.963 P=1,452	0.003 [0.018] N=1,512	0.009 [0.019] N=1,402	0.014 [0.021] N=1,255	0.017 [0.024] N=1,071	0.007 [0.026] N=881

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.006 [0.034] N=1,286	0.020 [0.034] N=1,176	0.009 [0.037] N=1,030	-0.004 [0.045] N=846	0.017 [0.059] N=656
Vegetables consumed at least weekly	0.042 [0.041] N=1,286	0.053 [0.045] N=1,176	0.018 [0.049] N=1,030	0.016 [0.057] N=846	-0.036 [0.068] N=656
Meat fish or poultry consumed at least weekly	0.071 [0.058] N=1,286	0.081 [0.059] N=1,176	0.055 [0.068] N=1,030	0.016 [0.080] N=846	-0.033 [0.091] N=656
Legumes consumed at least weekly	0.008 [0.060] N=1,286	-0.008 [0.065] N=1,176	-0.020 [0.067] N=1,030	0.015 [0.084] N=846	-0.116 [0.109] N=656
Animal products consumed at least weekly	0.043 [0.037] N=1,286	0.043 [0.040] N=1,176	0.022 [0.048] N=1,030	0.030 [0.060] N=846	0.004 [0.075] N=656
Sweets consumed at least weekly	-0.028 [0.052] N=1,286	-0.006 [0.057] N=1,176	-0.003 [0.064] N=1,030	0.019 [0.074] N=846	0.031 [0.101] N=656
Snack foods consumed at least weekly	-0.065 [0.057] N=1,286	-0.042 [0.063] N=1,176	-0.046 [0.073] N=1,030	-0.061 [0.092] N=846	-0.153 [0.101] N=656
Packaged soft drinks consumed at least weekly	-0.006 [0.061] N=1,286	0.013 [0.064] N=1,176	-0.016 [0.072] N=1,030	0.005 [0.088] N=846	-0.032 [0.114] N=656
Grains consumed at least weekly	0.038 [0.045] N=1,286	0.068 [0.050] N=1,176	0.033 [0.054] N=1,030	0.086 [0.065] N=846	0.091 [0.095] N=656
Child was ever breastfed	0.010 [0.026] N=1,287	0.020 [0.029] N=1,177	0.029 [0.035] N=1,030	0.038 [0.047] N=846	0.034 [0.059] N=656

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.944 P=1,425	-0.021 [0.023] N=1,511	-0.007 [0.022] N=1,401	-0.014 [0.024] N=1,255	-0.022 [0.026] N=1,071	-0.025 [0.030] N=881
Vegetables consumed at least weekly	0.903 P=1,362	-0.003 [0.028] N=1,511	-0.000 [0.029] N=1,401	-0.025 [0.030] N=1,255	-0.027 [0.030] N=1,071	**-0.057* [0.034] N=881**
Meat fish or poultry consumed at least weekly	0.802 P=1,217	0.058 [0.045] N=1,511	0.065 [0.046] N=1,401	0.047 [0.051] N=1,255	0.036 [0.056] N=1,071	0.016 [0.063] N=881
Legumes consumed at least weekly	0.798 P=1,208	**-0.072* [0.038] N=1,511**	**-0.084** [0.039] N=1,401**	**-0.102*** [0.039] N=1,255**	**-0.096** [0.044] N=1,071**	**-0.169*** [0.050] N=881**
Animal products consumed at least weekly	0.919 P=1,390	0.027 [0.029] N=1,511	0.026 [0.030] N=1,401	0.014 [0.033] N=1,255	0.015 [0.037] N=1,071	0.004 [0.040] N=881
Sweets consumed at least weekly	0.724 P=1,088	0.000 [0.038] N=1,511	0.016 [0.040] N=1,401	0.023 [0.041] N=1,255	0.035 [0.044] N=1,071	0.045 [0.048] N=881
Snack foods consumed at least weekly	0.594 P=888	-0.040 [0.045] N=1,511	-0.023 [0.047] N=1,401	-0.021 [0.051] N=1,255	-0.025 [0.057] N=1,071	-0.039 [0.062] N=881
Packaged soft drinks consumed at least weekly	0.376 P=558	0.034 [0.050] N=1,511	0.050 [0.052] N=1,401	0.038 [0.057] N=1,255	0.060 [0.063] N=1,071	0.050 [0.073] N=881
Grains consumed at least weekly	0.927 P=1,399	-0.003 [0.028] N=1,511	0.009 [0.030] N=1,401	-0.014 [0.031] N=1,255	-0.003 [0.034] N=1,071	-0.020 [0.040] N=881
Child was ever breastfed	0.963 P=1,452	0.002 [0.018] N=1,512	0.007 [0.019] N=1,402	0.012 [0.021] N=1,255	0.014 [0.024] N=1,071	0.004 [0.026] N=881

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.034] N=1,286	0.020 [0.033] N=1,176	0.011 [0.036] N=1,030	-0.003 [0.043] N=846	0.014 [0.056] N=656
Vegetables consumed at least weekly	0.041 [0.041] N=1,286	0.053 [0.045] N=1,176	0.020 [0.049] N=1,030	0.016 [0.058] N=846	-0.039 [0.067] N=656
Meat fish or poultry consumed at least weekly	0.067 [0.058] N=1,286	0.077 [0.058] N=1,176	0.052 [0.067] N=1,030	0.017 [0.079] N=846	-0.022 [0.090] N=656
Legumes consumed at least weekly	0.005 [0.060] N=1,286	-0.010 [0.066] N=1,176	-0.016 [0.067] N=1,030	0.018 [0.086] N=846	-0.121 [0.112] N=656
Animal products consumed at least weekly	0.040 [0.037] N=1,286	0.041 [0.039] N=1,176	0.020 [0.048] N=1,030	0.025 [0.059] N=846	-0.002 [0.076] N=656
Sweets consumed at least weekly	-0.038 [0.052] N=1,286	-0.019 [0.057] N=1,176	-0.011 [0.063] N=1,030	0.014 [0.074] N=846	0.037 [0.099] N=656
Snack foods consumed at least weekly	-0.074 [0.056] N=1,286	-0.054 [0.062] N=1,176	-0.053 [0.073] N=1,030	-0.065 [0.092] N=846	-0.148 [0.101] N=656
Packaged soft drinks consumed at least weekly	-0.011 [0.061] N=1,286	0.009 [0.065] N=1,176	-0.017 [0.073] N=1,030	0.003 [0.088] N=846	-0.040 [0.112] N=656
Grains consumed at least weekly	0.037 [0.045] N=1,286	0.069 [0.050] N=1,176	0.038 [0.056] N=1,030	0.089 [0.067] N=846	0.089 [0.096] N=656
Child was ever breastfed	0.012 [0.028] N=1,287	0.024 [0.032] N=1,177	0.033 [0.038] N=1,030	0.042 [0.049] N=846	0.041 [0.063] N=656

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,368	-0.013 [0.027] N=1,446	-0.014 [0.029] N=1,316	-0.017 [0.032] N=1,150	-0.017 [0.032] N=964	-0.011 [0.039] N=770
Vegetables consumed at least weekly	0.904 P=1,304	0.003 [0.034] N=1,446	0.012 [0.036] N=1,316	0.019 [0.039] N=1,150	0.016 [0.042] N=964	0.019 [0.049] N=770
Meat fish or poultry consumed at least weekly	0.789 P=1,152	0.068 [0.042] N=1,446	**0.081* [0.044] N=1,316**	**0.081* [0.048] N=1,150**	**0.123** [0.048] N=964**	**0.126** [0.056] N=770**
Legumes consumed at least weekly	0.801 P=1,165	-0.035 [0.046] N=1,446	-0.017 [0.049] N=1,316	-0.022 [0.053] N=1,150	-0.030 [0.058] N=964	-0.008 [0.069] N=770
Animal products consumed at least weekly	0.916 P=1,326	0.000 [0.034] N=1,446	0.005 [0.036] N=1,316	0.013 [0.038] N=1,150	-0.000 [0.040] N=964	0.030 [0.045] N=770
Sweets consumed at least weekly	0.731 P=1,059	0.022 [0.052] N=1,446	0.027 [0.054] N=1,316	0.010 [0.058] N=1,150	0.012 [0.062] N=964	-0.029 [0.072] N=770
Snack foods consumed at least weekly	0.592 P=852	-0.008 [0.059] N=1,446	-0.001 [0.062] N=1,316	0.006 [0.065] N=1,150	0.028 [0.070] N=964	-0.012 [0.080] N=770
Packaged soft drinks consumed at least weekly	0.355 P=509	-0.007 [0.048] N=1,446	-0.002 [0.050] N=1,316	0.036 [0.052] N=1,150	0.055 [0.056] N=964	0.050 [0.064] N=770
Grains consumed at least weekly	0.929 P=1,342	0.042 [0.037] N=1,446	0.048 [0.039] N=1,316	0.054 [0.042] N=1,150	0.060 [0.046] N=964	0.064 [0.052] N=770
Child was ever breastfed	0.962 P=1,391	-0.007 [0.024] N=1,447	-0.006 [0.024] N=1,317	-0.005 [0.025] N=1,151	-0.002 [0.025] N=965	-0.012 [0.028] N=771

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.049 [0.034] N=1,259	-0.058 [0.038] N=1,129	-0.073 [0.045] N=963	-0.069 [0.047] N=777	-0.075 [0.071] N=583
Vegetables consumed at least weekly	-0.049 [0.039] N=1,259	-0.043 [0.044] N=1,129	-0.043 [0.049] N=963	-0.044 [0.055] N=777	-0.051 [0.078] N=583
Meat fish or poultry consumed at least weekly	-0.002 [0.058] N=1,259	0.005 [0.062] N=1,129	-0.020 [0.069] N=963	0.053 [0.071] N=777	0.023 [0.092] N=583
Legumes consumed at least weekly	-0.063 [0.055] N=1,259	-0.037 [0.060] N=1,129	-0.051 [0.071] N=963	-0.047 [0.079] N=777	0.026 [0.097] N=583
Animal products consumed at least weekly	-0.054 [0.039] N=1,259	-0.046 [0.044] N=1,129	-0.044 [0.047] N=963	**-0.089* [0.050] N=777**	-0.054 [0.063] N=583
Sweets consumed at least weekly	0.032 [0.068] N=1,259	0.042 [0.074] N=1,129	0.013 [0.083] N=963	0.018 [0.096] N=777	-0.096 [0.121] N=583
Snack foods consumed at least weekly	0.020 [0.087] N=1,259	0.030 [0.094] N=1,129	0.048 [0.106] N=963	0.115 [0.124] N=777	0.056 [0.164] N=583
Packaged soft drinks consumed at least weekly	-0.042 [0.078] N=1,259	-0.036 [0.085] N=1,129	0.028 [0.095] N=963	0.063 [0.111] N=777	0.060 [0.142] N=583
Grains consumed at least weekly	0.015 [0.038] N=1,259	0.016 [0.041] N=1,129	0.019 [0.047] N=963	0.013 [0.058] N=777	-0.026 [0.079] N=583
Child was ever breastfed	0.024 [0.045] N=1,260	0.028 [0.047] N=1,130	0.037 [0.054] N=964	0.053 [0.060] N=778	0.053 [0.078] N=584

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,368	-0.013 [0.027] N=1,446	-0.014 [0.029] N=1,316	-0.017 [0.032] N=1,150	-0.017 [0.032] N=964	-0.011 [0.039] N=770
Vegetables consumed at least weekly	0.904 P=1,304	0.004 [0.034] N=1,446	0.012 [0.036] N=1,316	0.019 [0.038] N=1,150	0.016 [0.042] N=964	0.019 [0.049] N=770
Meat fish or poultry consumed at least weekly	0.789 P=1,152	0.069 [0.042] N=1,446	**0.082* [0.044] N=1,316**	**0.082* [0.048] N=1,150**	**0.123** [0.048] N=964**	**0.126** [0.057] N=770**
Legumes consumed at least weekly	0.801 P=1,165	-0.035 [0.046] N=1,446	-0.017 [0.049] N=1,316	-0.023 [0.053] N=1,150	-0.030 [0.058] N=964	-0.007 [0.068] N=770
Animal products consumed at least weekly	0.916 P=1,326	0.002 [0.034] N=1,446	0.007 [0.035] N=1,316	0.014 [0.038] N=1,150	-0.001 [0.040] N=964	0.029 [0.045] N=770
Sweets consumed at least weekly	0.731 P=1,059	0.024 [0.052] N=1,446	0.028 [0.055] N=1,316	0.010 [0.058] N=1,150	0.012 [0.063] N=964	-0.029 [0.072] N=770
Snack foods consumed at least weekly	0.592 P=852	-0.006 [0.059] N=1,446	-0.000 [0.062] N=1,316	0.007 [0.065] N=1,150	0.027 [0.071] N=964	-0.013 [0.081] N=770
Packaged soft drinks consumed at least weekly	0.355 P=509	-0.007 [0.048] N=1,446	-0.002 [0.050] N=1,316	0.036 [0.052] N=1,150	0.055 [0.056] N=964	0.049 [0.064] N=770
Grains consumed at least weekly	0.929 P=1,342	0.042 [0.037] N=1,446	0.048 [0.039] N=1,316	0.054 [0.042] N=1,150	0.060 [0.046] N=964	0.064 [0.053] N=770
Child was ever breastfed	0.962 P=1,391	-0.007 [0.024] N=1,447	-0.006 [0.024] N=1,317	-0.005 [0.025] N=1,151	-0.002 [0.025] N=965	-0.012 [0.027] N=771

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.049 [0.034] N=1,259	-0.058 [0.038] N=1,129	-0.074 [0.045] N=963	-0.070 [0.047] N=777	-0.075 [0.072] N=583
Vegetables consumed at least weekly	-0.048 [0.039] N=1,259	-0.042 [0.044] N=1,129	-0.042 [0.050] N=963	-0.043 [0.055] N=777	-0.051 [0.078] N=583
Meat fish or poultry consumed at least weekly	0.001 [0.058] N=1,259	0.008 [0.062] N=1,129	-0.017 [0.070] N=963	0.057 [0.072] N=777	0.031 [0.095] N=583
Legumes consumed at least weekly	-0.064 [0.055] N=1,259	-0.039 [0.060] N=1,129	-0.053 [0.071] N=963	-0.050 [0.079] N=777	0.018 [0.096] N=583
Animal products consumed at least weekly	-0.049 [0.039] N=1,259	-0.040 [0.043] N=1,129	-0.039 [0.047] N=963	**-0.085* [0.050] N=777**	-0.046 [0.063] N=583
Sweets consumed at least weekly	0.035 [0.069] N=1,259	0.046 [0.074] N=1,129	0.015 [0.083] N=963	0.018 [0.095] N=777	-0.093 [0.121] N=583
Snack foods consumed at least weekly	0.028 [0.088] N=1,259	0.036 [0.095] N=1,129	0.052 [0.106] N=963	0.118 [0.124] N=777	0.066 [0.167] N=583
Packaged soft drinks consumed at least weekly	-0.041 [0.078] N=1,259	-0.036 [0.084] N=1,129	0.027 [0.095] N=963	0.062 [0.111] N=777	0.067 [0.141] N=583
Grains consumed at least weekly	0.015 [0.038] N=1,259	0.016 [0.041] N=1,129	0.017 [0.047] N=963	0.010 [0.057] N=777	-0.030 [0.079] N=583
Child was ever breastfed	0.025 [0.045] N=1,260	0.029 [0.047] N=1,130	0.037 [0.054] N=964	0.053 [0.060] N=778	0.055 [0.079] N=584

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,368	-0.014 [0.027] N=1,446	-0.015 [0.029] N=1,316	-0.016 [0.032] N=1,150	-0.017 [0.033] N=964	-0.011 [0.040] N=770
Vegetables consumed at least weekly	0.904 P=1,304	0.003 [0.033] N=1,446	0.012 [0.035] N=1,316	0.020 [0.038] N=1,150	0.016 [0.042] N=964	0.019 [0.049] N=770
Meat fish or poultry consumed at least weekly	0.789 P=1,152	0.063 [0.041] N=1,446	**0.079* [0.043] N=1,316**	**0.085* [0.046] N=1,150**	**0.123*** [0.047] N=964**	**0.127** [0.055] N=770**
Legumes consumed at least weekly	0.801 P=1,165	-0.037 [0.046] N=1,446	-0.018 [0.050] N=1,316	-0.022 [0.054] N=1,150	-0.030 [0.058] N=964	-0.006 [0.069] N=770
Animal products consumed at least weekly	0.916 P=1,326	0.000 [0.034] N=1,446	0.005 [0.036] N=1,316	0.015 [0.037] N=1,150	-0.001 [0.040] N=964	0.030 [0.045] N=770
Sweets consumed at least weekly	0.731 P=1,059	0.023 [0.052] N=1,446	0.027 [0.054] N=1,316	0.011 [0.058] N=1,150	0.011 [0.062] N=964	-0.029 [0.071] N=770
Snack foods consumed at least weekly	0.592 P=852	-0.003 [0.059] N=1,446	0.001 [0.063] N=1,316	0.007 [0.066] N=1,150	0.026 [0.070] N=964	-0.013 [0.081] N=770
Packaged soft drinks consumed at least weekly	0.355 P=509	-0.005 [0.049] N=1,446	-0.002 [0.051] N=1,316	0.034 [0.053] N=1,150	0.053 [0.057] N=964	0.048 [0.066] N=770
Grains consumed at least weekly	0.929 P=1,342	0.040 [0.037] N=1,446	0.047 [0.039] N=1,316	0.055 [0.042] N=1,150	0.060 [0.046] N=964	0.065 [0.052] N=770
Child was ever breastfed	0.962 P=1,391	-0.008 [0.024] N=1,447	-0.007 [0.024] N=1,317	-0.004 [0.025] N=1,151	-0.002 [0.025] N=965	-0.011 [0.027] N=771

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.047 [0.034] N=1,259	-0.055 [0.037] N=1,129	-0.068 [0.044] N=963	-0.066 [0.048] N=777	-0.070 [0.073] N=583
Vegetables consumed at least weekly	-0.047 [0.039] N=1,259	-0.040 [0.044] N=1,129	-0.038 [0.049] N=963	-0.040 [0.055] N=777	-0.048 [0.080] N=583
Meat fish or poultry consumed at least weekly	0.001 [0.056] N=1,259	0.014 [0.059] N=1,129	-0.001 [0.066] N=963	0.069 [0.071] N=777	0.043 [0.091] N=583
Legumes consumed at least weekly	-0.061 [0.054] N=1,259	-0.038 [0.060] N=1,129	-0.052 [0.071] N=963	-0.048 [0.079] N=777	0.016 [0.097] N=583
Animal products consumed at least weekly	-0.050 [0.039] N=1,259	-0.040 [0.043] N=1,129	-0.037 [0.047] N=963	**-0.087* [0.051] N=777**	-0.050 [0.063] N=583
Sweets consumed at least weekly	0.032 [0.068] N=1,259	0.042 [0.074] N=1,129	0.015 [0.082] N=963	0.013 [0.094] N=777	-0.103 [0.120] N=583
Snack foods consumed at least weekly	0.031 [0.088] N=1,259	0.037 [0.096] N=1,129	0.048 [0.106] N=963	0.111 [0.124] N=777	0.060 [0.170] N=583
Packaged soft drinks consumed at least weekly	-0.051 [0.080] N=1,259	-0.050 [0.087] N=1,129	0.007 [0.098] N=963	0.037 [0.113] N=777	0.045 [0.147] N=583
Grains consumed at least weekly	0.013 [0.038] N=1,259	0.015 [0.040] N=1,129	0.020 [0.045] N=963	0.008 [0.056] N=777	-0.035 [0.076] N=583
Child was ever breastfed	0.025 [0.045] N=1,260	0.029 [0.047] N=1,130	0.040 [0.054] N=964	0.058 [0.060] N=778	0.057 [0.079] N=584

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,368	-0.014 [0.027] N=1,446	-0.014 [0.029] N=1,316	-0.015 [0.032] N=1,150	-0.020 [0.032] N=964	-0.015 [0.038] N=770
Vegetables consumed at least weekly	0.904 P=1,304	0.002 [0.035] N=1,446	0.012 [0.036] N=1,316	0.020 [0.039] N=1,150	0.014 [0.042] N=964	0.019 [0.049] N=770
Meat fish or poultry consumed at least weekly	0.789 P=1,152	0.061 [0.043] N=1,446	**0.079* [0.045] N=1,316**	**0.082* [0.048] N=1,150**	**0.116** [0.048] N=964**	**0.117** [0.056] N=770**
Legumes consumed at least weekly	0.801 P=1,165	-0.033 [0.046] N=1,446	-0.014 [0.050] N=1,316	-0.017 [0.053] N=1,150	-0.026 [0.057] N=964	-0.001 [0.068] N=770
Animal products consumed at least weekly	0.916 P=1,326	0.007 [0.034] N=1,446	0.013 [0.036] N=1,316	0.022 [0.038] N=1,150	0.003 [0.041] N=964	0.034 [0.046] N=770
Sweets consumed at least weekly	0.731 P=1,059	0.030 [0.053] N=1,446	0.033 [0.056] N=1,316	0.013 [0.059] N=1,150	0.012 [0.064] N=964	-0.030 [0.074] N=770
Snack foods consumed at least weekly	0.592 P=852	0.005 [0.058] N=1,446	0.008 [0.062] N=1,316	0.012 [0.065] N=1,150	0.031 [0.070] N=964	-0.006 [0.080] N=770
Packaged soft drinks consumed at least weekly	0.355 P=509	0.006 [0.048] N=1,446	0.005 [0.050] N=1,316	0.039 [0.052] N=1,150	0.058 [0.057] N=964	0.054 [0.066] N=770
Grains consumed at least weekly	0.929 P=1,342	0.039 [0.036] N=1,446	0.045 [0.038] N=1,316	0.050 [0.041] N=1,150	0.055 [0.046] N=964	0.060 [0.052] N=770
Child was ever breastfed	0.962 P=1,391	-0.007 [0.024] N=1,447	-0.006 [0.024] N=1,317	-0.005 [0.025] N=1,151	-0.003 [0.026] N=965	-0.013 [0.029] N=771

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.050 [0.034] N=1,259	-0.057 [0.037] N=1,129	-0.070 [0.044] N=963	-0.072 [0.047] N=777	-0.083 [0.071] N=583
Vegetables consumed at least weekly	-0.050 [0.040] N=1,259	-0.041 [0.044] N=1,129	-0.038 [0.050] N=963	-0.041 [0.054] N=777	-0.047 [0.075] N=583
Meat fish or poultry consumed at least weekly	-0.004 [0.057] N=1,259	0.008 [0.061] N=1,129	-0.010 [0.067] N=963	0.050 [0.074] N=777	0.007 [0.094] N=583
Legumes consumed at least weekly	-0.061 [0.054] N=1,259	-0.035 [0.060] N=1,129	-0.042 [0.071] N=963	-0.038 [0.078] N=777	0.034 [0.093] N=583
Animal products consumed at least weekly	-0.043 [0.037] N=1,259	-0.033 [0.042] N=1,129	-0.031 [0.045] N=963	**-0.084* [0.050] N=777**	-0.047 [0.060] N=583
Sweets consumed at least weekly	0.041 [0.068] N=1,259	0.050 [0.074] N=1,129	0.016 [0.082] N=963	0.013 [0.095] N=777	-0.106 [0.117] N=583
Snack foods consumed at least weekly	0.036 [0.085] N=1,259	0.039 [0.093] N=1,129	0.049 [0.102] N=963	0.106 [0.120] N=777	0.055 [0.159] N=583
Packaged soft drinks consumed at least weekly	-0.030 [0.078] N=1,259	-0.035 [0.084] N=1,129	0.019 [0.095] N=963	0.053 [0.110] N=777	0.071 [0.143] N=583
Grains consumed at least weekly	0.011 [0.037] N=1,259	0.012 [0.040] N=1,129	0.012 [0.045] N=963	0.000 [0.056] N=777	-0.041 [0.076] N=583
Child was ever breastfed	0.025 [0.045] N=1,260	0.030 [0.047] N=1,130	0.039 [0.055] N=964	0.054 [0.061] N=778	0.054 [0.081] N=584

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,277	-0.012 [0.029] N=1,350	-0.013 [0.031] N=1,227	-0.010 [0.035] N=1,076	-0.015 [0.035] N=901	-0.011 [0.041] N=723
Vegetables consumed at least weekly	0.906 P=1,221	0.009 [0.034] N=1,350	0.018 [0.036] N=1,227	0.031 [0.039] N=1,076	0.024 [0.044] N=901	0.013 [0.051] N=723
Meat fish or poultry consumed at least weekly	0.788 P=1,074	0.069 [0.044] N=1,350	**0.081* [0.046] N=1,227**	**0.085* [0.050] N=1,076**	**0.102** [0.051] N=901**	**0.100* [0.059] N=723**
Legumes consumed at least weekly	0.797 P=1,084	-0.035 [0.048] N=1,350	-0.015 [0.051] N=1,227	-0.018 [0.056] N=1,076	-0.029 [0.061] N=901	-0.013 [0.072] N=723
Animal products consumed at least weekly	0.917 P=1,240	0.002 [0.035] N=1,350	0.013 [0.037] N=1,227	0.022 [0.040] N=1,076	0.010 [0.042] N=901	0.041 [0.047] N=723
Sweets consumed at least weekly	0.738 P=997	0.023 [0.053] N=1,350	0.026 [0.056] N=1,227	0.003 [0.061] N=1,076	-0.001 [0.066] N=901	-0.038 [0.078] N=723
Snack foods consumed at least weekly	0.597 P=802	0.005 [0.060] N=1,350	0.005 [0.064] N=1,227	0.002 [0.068] N=1,076	0.021 [0.072] N=901	-0.011 [0.083] N=723
Packaged soft drinks consumed at least weekly	0.365 P=490	-0.015 [0.050] N=1,350	-0.016 [0.052] N=1,227	0.022 [0.054] N=1,076	0.042 [0.060] N=901	0.036 [0.069] N=723
Grains consumed at least weekly	0.929 P=1,253	0.048 [0.038] N=1,350	0.053 [0.040] N=1,227	0.058 [0.043] N=1,076	0.066 [0.048] N=901	0.068 [0.054] N=723
Child was ever breastfed	0.963 P=1,302	-0.010 [0.022] N=1,351	-0.010 [0.023] N=1,228	-0.010 [0.024] N=1,077	-0.012 [0.024] N=902	-0.025 [0.027] N=724

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.044 [0.037] N=1,174	-0.052 [0.041] N=1,051	-0.054 [0.050] N=900	-0.057 [0.051] N=725	-0.059 [0.073] N=547
Vegetables consumed at least weekly	-0.043 [0.039] N=1,174	-0.038 [0.043] N=1,051	-0.026 [0.050] N=900	-0.029 [0.058] N=725	-0.073 [0.074] N=547
Meat fish or poultry consumed at least weekly	0.020 [0.064] N=1,174	0.029 [0.068] N=1,051	0.020 [0.076] N=900	0.051 [0.080] N=725	0.018 [0.103] N=547
Legumes consumed at least weekly	-0.051 [0.056] N=1,174	-0.023 [0.060] N=1,051	-0.024 [0.073] N=900	-0.023 [0.082] N=725	0.036 [0.095] N=547
Animal products consumed at least weekly	-0.050 [0.041] N=1,174	-0.034 [0.045] N=1,051	-0.026 [0.049] N=900	-0.063 [0.054] N=725	-0.017 [0.064] N=547
Sweets consumed at least weekly	0.033 [0.072] N=1,174	0.040 [0.077] N=1,051	-0.006 [0.086] N=900	-0.015 [0.098] N=725	-0.126 [0.127] N=547
Snack foods consumed at least weekly	0.044 [0.092] N=1,174	0.045 [0.099] N=1,051	0.042 [0.113] N=900	0.098 [0.129] N=725	0.070 [0.173] N=547
Packaged soft drinks consumed at least weekly	-0.058 [0.078] N=1,174	-0.061 [0.085] N=1,051	-0.005 [0.095] N=900	0.026 [0.112] N=725	0.002 [0.143] N=547
Grains consumed at least weekly	0.028 [0.042] N=1,174	0.030 [0.045] N=1,051	0.033 [0.053] N=900	0.028 [0.062] N=725	-0.012 [0.082] N=547
Child was ever breastfed	0.026 [0.042] N=1,175	0.028 [0.045] N=1,052	0.035 [0.051] N=901	0.043 [0.055] N=726	0.027 [0.074] N=548

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.949 P=1,277	-0.012 [0.029] N=1,350	-0.012 [0.031] N=1,227	-0.010 [0.035] N=1,076	-0.016 [0.035] N=901	-0.012 [0.041] N=723
Vegetables consumed at least weekly	0.906 P=1,221	0.008 [0.034] N=1,350	0.016 [0.036] N=1,227	0.029 [0.039] N=1,076	0.021 [0.044] N=901	0.011 [0.051] N=723
Meat fish or poultry consumed at least weekly	0.788 P=1,074	0.070 [0.044] N=1,350	**0.084* [0.045] N=1,227**	**0.087* [0.049] N=1,076**	**0.105** [0.050] N=901**	**0.102* [0.058] N=723**
Legumes consumed at least weekly	0.797 P=1,084	-0.035 [0.048] N=1,350	-0.015 [0.051] N=1,227	-0.017 [0.056] N=1,076	-0.032 [0.061] N=901	-0.018 [0.072] N=723
Animal products consumed at least weekly	0.917 P=1,240	0.002 [0.035] N=1,350	0.012 [0.037] N=1,227	0.021 [0.040] N=1,076	0.008 [0.042] N=901	0.039 [0.047] N=723
Sweets consumed at least weekly	0.738 P=997	0.023 [0.054] N=1,350	0.026 [0.057] N=1,227	0.003 [0.062] N=1,076	-0.001 [0.067] N=901	-0.039 [0.079] N=723
Snack foods consumed at least weekly	0.597 P=802	0.006 [0.060] N=1,350	0.005 [0.063] N=1,227	0.003 [0.067] N=1,076	0.021 [0.072] N=901	-0.011 [0.083] N=723
Packaged soft drinks consumed at least weekly	0.365 P=490	-0.015 [0.050] N=1,350	-0.016 [0.052] N=1,227	0.022 [0.054] N=1,076	0.041 [0.060] N=901	0.033 [0.070] N=723
Grains consumed at least weekly	0.929 P=1,253	0.048 [0.038] N=1,350	0.053 [0.040] N=1,227	0.058 [0.043] N=1,076	0.064 [0.048] N=901	0.067 [0.054] N=723
Child was ever breastfed	0.963 P=1,302	-0.009 [0.023] N=1,351	-0.011 [0.023] N=1,228	-0.009 [0.024] N=1,077	-0.008 [0.024] N=902	-0.021 [0.028] N=724

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.037] N=1,174	-0.049 [0.041] N=1,051	-0.053 [0.050] N=900	-0.059 [0.052] N=725	-0.063 [0.075] N=547
Vegetables consumed at least weekly	-0.044 [0.039] N=1,174	-0.040 [0.044] N=1,051	-0.028 [0.051] N=900	-0.033 [0.058] N=725	-0.076 [0.075] N=547
Meat fish or poultry consumed at least weekly	0.021 [0.064] N=1,174	0.032 [0.068] N=1,051	0.019 [0.076] N=900	0.052 [0.080] N=725	0.018 [0.102] N=547
Legumes consumed at least weekly	-0.050 [0.056] N=1,174	-0.021 [0.061] N=1,051	-0.024 [0.073] N=900	-0.031 [0.083] N=725	0.030 [0.094] N=547
Animal products consumed at least weekly	-0.050 [0.041] N=1,174	-0.034 [0.046] N=1,051	-0.027 [0.050] N=900	-0.066 [0.054] N=725	-0.022 [0.064] N=547
Sweets consumed at least weekly	0.035 [0.072] N=1,174	0.042 [0.078] N=1,051	-0.001 [0.086] N=900	-0.009 [0.099] N=725	-0.117 [0.127] N=547
Snack foods consumed at least weekly	0.045 [0.091] N=1,174	0.044 [0.099] N=1,051	0.042 [0.112] N=900	0.104 [0.128] N=725	0.071 [0.173] N=547
Packaged soft drinks consumed at least weekly	-0.058 [0.079] N=1,174	-0.064 [0.086] N=1,051	-0.003 [0.095] N=900	0.030 [0.111] N=725	0.008 [0.143] N=547
Grains consumed at least weekly	0.027 [0.042] N=1,174	0.030 [0.045] N=1,051	0.031 [0.052] N=900	0.024 [0.062] N=725	-0.018 [0.081] N=547
Child was ever breastfed	0.026 [0.043] N=1,175	0.025 [0.045] N=1,052	0.035 [0.052] N=901	0.050 [0.057] N=726	0.033 [0.077] N=548

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.