Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Oldest Child

Age reference: 10 October 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.120	**0.405* [0.237] N=1,422**	**0.448* [0.253] N=1,317**	0.441 [0.278] N=1,182	0.438 [0.282] N=1,009	0.149 [0.307] N=834
WHO weight-for-age z score	0.256	-0.059 [0.132] N=1,609	-0.002 [0.139] N=1,485	-0.009 [0.145] N=1,331	-0.041 [0.158] N=1,133	-0.216 [0.174] N=930
WHO weight-for-length/height z score, 0-60 months	0.371	-0.002 [0.227] N=1,396	-0.049 [0.240] N=1,299	-0.147 [0.273] N=1,167	-0.185 [0.304] N=997	-0.160 [0.342] N=820
WHO BMI-for-age z score	0.335	-0.009 [0.245] N=1,364	-0.003 [0.263] N=1,269	-0.124 [0.300] N=1,136	-0.241 [0.337] N=969	-0.164 [0.374] N=800
Stunted: HAZ below -2	0.205 P=279	-0.058 [0.041] N=1,422	-0.069 [0.043] N=1,317	-0.042 [0.046] N=1,182	-0.027 [0.048] N=1,009	-0.002 [0.056] N=834
Underweight: WAZ below -2	0.051 P=87	0.017 [0.019] N=1,609	0.006 [0.020] N=1,485	0.006 [0.021] N=1,331	0.009 [0.024] N=1,133	0.029 [0.031] N=930
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=162	-0.030 [0.036] N=1,396	-0.022 [0.036] N=1,299	-0.020 [0.039] N=1,167	-0.015 [0.042] N=997	-0.032 [0.044] N=820
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=266	-0.007 [0.043] N=1,396	-0.017 [0.046] N=1,299	-0.024 [0.051] N=1,167	-0.019 [0.059] N=997	-0.013 [0.065] N=820
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=126	0.002 [0.031] N=1,396	-0.009 [0.032] N=1,299	-0.008 [0.035] N=1,167	-0.011 [0.039] N=997	-0.009 [0.042] N=820
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=380	**-0.080* [0.043] N=1,354**	**-0.084** [0.042] N=1,260**	-0.066 [0.045] N=1,132	-0.047 [0.046] N=965	-0.043 [0.052] N=796
BMI-for-age z score below -2	0.123 P=173	-0.012 [0.040] N=1,364	-0.006 [0.041] N=1,269	0.010 [0.045] N=1,136	0.044 [0.049] N=969	0.035 [0.053] N=800

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.362 [0.335] N=1,203	0.424 [0.378] N=1,098	0.496 [0.440] N=963	0.621 [0.473] N=790	-0.235 [0.619] N=615
WHO weight-for-age z score	-0.168 [0.182] N=1,370	-0.088 [0.197] N=1,246	-0.117 [0.211] N=1,092	-0.205 [0.267] N=894	**-0.887** [0.348] N=691**
WHO weight-for-length/height z score, 0-60 months	0.080 [0.247] N=1,178	0.030 [0.272] N=1,081	-0.217 [0.348] N=949	-0.330 [0.409] N=779	-0.404 [0.497] N=602
WHO BMI-for-age z score	0.144 [0.264] N=1,154	0.203 [0.295] N=1,059	-0.074 [0.381] N=926	-0.369 [0.453] N=759	-0.293 [0.525] N=590
Stunted: HAZ below -2	-0.022 [0.056] N=1,203	-0.035 [0.061] N=1,098	0.016 [0.066] N=963	0.054 [0.073] N=790	0.174 [0.106] N=615
Underweight: WAZ below -2	0.031 [0.024] N=1,370	0.015 [0.027] N=1,246	0.015 [0.029] N=1,092	0.029 [0.036] N=894	**0.118** [0.056] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.028 [0.049] N=1,178	-0.014 [0.051] N=1,081	-0.004 [0.059] N=949	0.006 [0.071] N=779	-0.027 [0.089] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.051] N=1,178	-0.015 [0.055] N=1,081	-0.039 [0.064] N=949	-0.037 [0.081] N=779	-0.044 [0.088] N=602
Obese: WLZ/WHZ above +3, 0-60 months	0.020 [0.036] N=1,178	0.005 [0.039] N=1,081	0.014 [0.046] N=949	0.015 [0.057] N=779	0.024 [0.065] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.025 [0.064] N=1,144	-0.025 [0.066] N=1,050	0.021 [0.076] N=922	0.082 [0.086] N=755	0.166 [0.129] N=586
BMI-for-age z score below -2	-0.043 [0.048] N=1,154	-0.040 [0.050] N=1,059	-0.010 [0.060] N=926	0.052 [0.073] N=759	0.031 [0.090] N=590

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.120	**0.405* [0.237] N=1,422**	**0.448* [0.253] N=1,317**	0.441 [0.278] N=1,182	0.439 [0.282] N=1,009	0.150 [0.307] N=834
WHO weight-for-age z score	0.256	-0.061 [0.131] N=1,609	-0.003 [0.139] N=1,485	-0.009 [0.145] N=1,331	-0.041 [0.158] N=1,133	-0.215 [0.173] N=930
WHO weight-for-length/height z score, 0-60 months	0.371	-0.006 [0.227] N=1,396	-0.054 [0.240] N=1,299	-0.150 [0.273] N=1,167	-0.190 [0.303] N=997	-0.166 [0.340] N=820
WHO BMI-for-age z score	0.335	-0.012 [0.245] N=1,364	-0.009 [0.262] N=1,269	-0.127 [0.300] N=1,136	-0.248 [0.336] N=969	-0.174 [0.371] N=800
Stunted: HAZ below -2	0.205 P=279	-0.058 [0.041] N=1,422	-0.069 [0.043] N=1,317	-0.042 [0.046] N=1,182	-0.027 [0.048] N=1,009	-0.002 [0.056] N=834
Underweight: WAZ below -2	0.051 P=87	0.017 [0.019] N=1,609	0.006 [0.020] N=1,485	0.006 [0.021] N=1,331	0.008 [0.024] N=1,133	0.029 [0.031] N=930
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=162	-0.029 [0.036] N=1,396	-0.020 [0.036] N=1,299	-0.019 [0.039] N=1,167	-0.014 [0.042] N=997	-0.030 [0.044] N=820
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=266	-0.007 [0.043] N=1,396	-0.017 [0.046] N=1,299	-0.024 [0.051] N=1,167	-0.019 [0.059] N=997	-0.013 [0.065] N=820
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=126	0.002 [0.031] N=1,396	-0.009 [0.032] N=1,299	-0.007 [0.035] N=1,167	-0.011 [0.039] N=997	-0.009 [0.042] N=820
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=380	**-0.078* [0.043] N=1,354**	**-0.083* [0.042] N=1,260**	-0.066 [0.045] N=1,132	-0.046 [0.046] N=965	-0.041 [0.052] N=796
BMI-for-age z score below -2	0.123 P=173	-0.011 [0.040] N=1,364	-0.004 [0.041] N=1,269	0.011 [0.044] N=1,136	0.046 [0.048] N=969	0.038 [0.052] N=800

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.361 [0.334] N=1,203	0.418 [0.376] N=1,098	0.486 [0.438] N=963	0.597 [0.467] N=790	-0.270 [0.614] N=615
WHO weight-for-age z score	-0.168 [0.181] N=1,370	-0.083 [0.196] N=1,246	-0.109 [0.210] N=1,092	-0.197 [0.267] N=894	**-0.867** [0.348] N=691**
WHO weight-for-length/height z score, 0-60 months	0.075 [0.247] N=1,178	0.025 [0.272] N=1,081	-0.216 [0.347] N=949	-0.333 [0.408] N=779	-0.408 [0.504] N=602
WHO BMI-for-age z score	0.142 [0.265] N=1,154	0.202 [0.295] N=1,059	-0.067 [0.379] N=926	-0.367 [0.453] N=759	-0.299 [0.537] N=590
Stunted: HAZ below -2	-0.022 [0.056] N=1,203	-0.035 [0.061] N=1,098	0.016 [0.066] N=963	0.056 [0.072] N=790	**0.178* [0.106] N=615**
Underweight: WAZ below -2	0.031 [0.024] N=1,370	0.015 [0.027] N=1,246	0.015 [0.029] N=1,092	0.028 [0.035] N=894	**0.116** [0.055] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.026 [0.049] N=1,178	-0.012 [0.051] N=1,081	-0.005 [0.059] N=949	0.007 [0.071] N=779	-0.026 [0.090] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.051] N=1,178	-0.016 [0.055] N=1,081	-0.039 [0.064] N=949	-0.037 [0.081] N=779	-0.044 [0.088] N=602
Obese: WLZ/WHZ above +3, 0-60 months	0.021 [0.037] N=1,178	0.005 [0.040] N=1,081	0.014 [0.047] N=949	0.015 [0.057] N=779	0.024 [0.065] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.024 [0.064] N=1,144	-0.025 [0.066] N=1,050	0.018 [0.076] N=922	0.081 [0.086] N=755	0.165 [0.129] N=586
BMI-for-age z score below -2	-0.043 [0.048] N=1,154	-0.040 [0.050] N=1,059	-0.011 [0.060] N=926	0.051 [0.074] N=759	0.032 [0.093] N=590

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.120	0.370 [0.237] N=1,422	0.391 [0.252] N=1,317	0.399 [0.276] N=1,182	0.398 [0.278] N=1,009	0.076 [0.301] N=834
WHO weight-for-age z score	0.256	-0.109 [0.126] N=1,609	-0.054 [0.134] N=1,485	-0.058 [0.141] N=1,331	-0.086 [0.153] N=1,133	-0.273 [0.167] N=930
WHO weight-for-length/height z score, 0-60 months	0.371	-0.033 [0.227] N=1,396	-0.074 [0.242] N=1,299	-0.165 [0.272] N=1,167	-0.201 [0.301] N=997	-0.190 [0.337] N=820
WHO BMI-for-age z score	0.335	-0.023 [0.245] N=1,364	-0.016 [0.263] N=1,269	-0.133 [0.298] N=1,136	-0.252 [0.332] N=969	-0.178 [0.368] N=800
Stunted: HAZ below -2	0.205 P=279	-0.051 [0.041] N=1,422	-0.057 [0.043] N=1,317	-0.033 [0.045] N=1,182	-0.019 [0.048] N=1,009	0.011 [0.055] N=834
Underweight: WAZ below -2	0.051 P=87	0.022 [0.018] N=1,609	0.013 [0.019] N=1,485	0.013 [0.020] N=1,331	0.016 [0.023] N=1,133	0.038 [0.030] N=930
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=162	-0.022 [0.035] N=1,396	-0.011 [0.036] N=1,299	-0.011 [0.038] N=1,167	-0.008 [0.041] N=997	-0.025 [0.044] N=820
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=266	-0.006 [0.044] N=1,396	-0.013 [0.046] N=1,299	-0.020 [0.052] N=1,167	-0.016 [0.059] N=997	-0.011 [0.065] N=820
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=126	0.004 [0.031] N=1,396	-0.004 [0.032] N=1,299	-0.004 [0.035] N=1,167	-0.009 [0.039] N=997	-0.009 [0.042] N=820
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=380	-0.062 [0.043] N=1,354	-0.060 [0.043] N=1,260	-0.047 [0.045] N=1,132	-0.030 [0.046] N=965	-0.019 [0.052] N=796
BMI-for-age z score below -2	0.123 P=173	-0.006 [0.039] N=1,364	0.004 [0.041] N=1,269	0.018 [0.044] N=1,136	0.050 [0.048] N=969	0.041 [0.052] N=800

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.334 [0.335] N=1,203	0.348 [0.379] N=1,098	0.440 [0.441] N=963	0.572 [0.467] N=790	-0.365 [0.604] N=615
WHO weight-for-age z score	-0.200 [0.177] N=1,370	-0.121 [0.192] N=1,246	-0.136 [0.208] N=1,092	-0.209 [0.269] N=894	**-0.905** [0.348] N=691**
WHO weight-for-length/height z score, 0-60 months	0.068 [0.252] N=1,178	0.035 [0.281] N=1,081	-0.196 [0.354] N=949	-0.329 [0.415] N=779	-0.448 [0.517] N=602
WHO BMI-for-age z score	0.141 [0.269] N=1,154	0.212 [0.302] N=1,059	-0.054 [0.384] N=926	-0.383 [0.460] N=759	-0.327 [0.553] N=590
Stunted: HAZ below -2	-0.018 [0.057] N=1,203	-0.020 [0.062] N=1,098	0.024 [0.067] N=963	0.056 [0.072] N=790	**0.186* [0.105] N=615**
Underweight: WAZ below -2	0.035 [0.024] N=1,370	0.022 [0.026] N=1,246	0.020 [0.028] N=1,092	0.033 [0.035] N=894	**0.122** [0.055] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.020 [0.048] N=1,178	-0.003 [0.051] N=1,081	0.001 [0.059] N=949	0.016 [0.071] N=779	-0.009 [0.092] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	0.003 [0.052] N=1,178	-0.007 [0.056] N=1,081	-0.033 [0.065] N=949	-0.032 [0.081] N=779	-0.036 [0.086] N=602
Obese: WLZ/WHZ above +3, 0-60 months	0.023 [0.037] N=1,178	0.013 [0.040] N=1,081	0.017 [0.048] N=949	0.015 [0.057] N=779	0.018 [0.065] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.011 [0.064] N=1,144	-0.001 [0.066] N=1,050	0.033 [0.075] N=922	0.088 [0.086] N=755	0.196 [0.124] N=586
BMI-for-age z score below -2	-0.037 [0.048] N=1,154	-0.029 [0.050] N=1,059	-0.004 [0.060] N=926	0.062 [0.075] N=759	0.046 [0.096] N=590

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.120	0.294 [0.234] N=1,422	0.317 [0.250] N=1,317	0.331 [0.275] N=1,182	0.312 [0.277] N=1,009	-0.026 [0.301] N=834
WHO weight-for-age z score	0.256	-0.128 [0.128] N=1,609	-0.076 [0.136] N=1,485	-0.076 [0.143] N=1,331	-0.116 [0.159] N=1,133	**-0.312* [0.175] N=930**
WHO weight-for-length/height z score, 0-60 months	0.371	0.008 [0.224] N=1,396	-0.034 [0.238] N=1,299	-0.140 [0.271] N=1,167	-0.179 [0.305] N=997	-0.159 [0.342] N=820
WHO BMI-for-age z score	0.335	0.018 [0.243] N=1,364	0.030 [0.260] N=1,269	-0.099 [0.297] N=1,136	-0.218 [0.337] N=969	-0.139 [0.374] N=800
Stunted: HAZ below -2	0.205 P=279	-0.036 [0.041] N=1,422	-0.040 [0.043] N=1,317	-0.014 [0.045] N=1,182	0.004 [0.047] N=1,009	0.037 [0.055] N=834
Underweight: WAZ below -2	0.051 P=87	0.025 [0.018] N=1,609	0.016 [0.020] N=1,485	0.016 [0.021] N=1,331	0.019 [0.023] N=1,133	0.043 [0.030] N=930
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=162	-0.025 [0.035] N=1,396	-0.016 [0.036] N=1,299	-0.014 [0.039] N=1,167	-0.010 [0.042] N=997	-0.031 [0.044] N=820
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=266	0.001 [0.043] N=1,396	-0.008 [0.046] N=1,299	-0.018 [0.051] N=1,167	-0.012 [0.060] N=997	-0.011 [0.065] N=820
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=126	0.007 [0.031] N=1,396	-0.003 [0.032] N=1,299	-0.004 [0.035] N=1,167	-0.007 [0.040] N=997	-0.006 [0.043] N=820
CIAF: any stunting, underweight, or wasting, 0-60 months	0.285 P=380	-0.049 [0.044] N=1,354	-0.048 [0.044] N=1,260	-0.031 [0.047] N=1,132	-0.008 [0.049] N=965	0.005 [0.055] N=796
BMI-for-age z score below -2	0.123 P=173	-0.008 [0.039] N=1,364	-0.001 [0.041] N=1,269	0.015 [0.045] N=1,136	0.049 [0.049] N=969	0.036 [0.053] N=800

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.290 [0.336] N=1,203	0.323 [0.376] N=1,098	0.430 [0.440] N=963	0.542 [0.464] N=790	-0.462 [0.601] N=615
WHO weight-for-age z score	-0.227 [0.179] N=1,370	-0.150 [0.194] N=1,246	-0.165 [0.210] N=1,092	-0.252 [0.272] N=894	**-0.919** [0.350] N=691**
WHO weight-for-length/height z score, 0-60 months	0.062 [0.250] N=1,178	0.026 [0.274] N=1,081	-0.213 [0.350] N=949	-0.317 [0.413] N=779	-0.347 [0.522] N=602
WHO BMI-for-age z score	0.128 [0.268] N=1,154	0.207 [0.298] N=1,059	-0.061 [0.384] N=926	-0.346 [0.455] N=759	-0.226 [0.552] N=590
Stunted: HAZ below -2	-0.013 [0.057] N=1,203	-0.019 [0.063] N=1,098	0.021 [0.068] N=963	0.051 [0.075] N=790	**0.188* [0.108] N=615**
Underweight: WAZ below -2	0.038 [0.024] N=1,370	0.023 [0.027] N=1,246	0.022 [0.029] N=1,092	0.035 [0.036] N=894	**0.124** [0.057] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.016 [0.049] N=1,178	-0.004 [0.051] N=1,081	0.002 [0.059] N=949	0.016 [0.072] N=779	-0.010 [0.093] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	0.005 [0.052] N=1,178	-0.007 [0.055] N=1,081	-0.032 [0.064] N=949	-0.024 [0.080] N=779	-0.015 [0.086] N=602
Obese: WLZ/WHZ above +3, 0-60 months	0.027 [0.038] N=1,178	0.013 [0.040] N=1,081	0.019 [0.048] N=949	0.021 [0.058] N=779	0.035 [0.067] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.000 [0.065] N=1,144	0.003 [0.068] N=1,050	0.037 [0.077] N=922	0.093 [0.089] N=755	0.212 [0.129] N=586
BMI-for-age z score below -2	-0.032 [0.048] N=1,154	-0.032 [0.051] N=1,059	-0.006 [0.061] N=926	0.059 [0.075] N=759	0.045 [0.096] N=590

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.130	**0.420* [0.248] N=1,329**	0.417 [0.268] N=1,237	0.417 [0.289] N=1,107	0.408 [0.292] N=949	0.150 [0.310] N=784
WHO weight-for-age z score	0.255	-0.018 [0.135] N=1,502	0.044 [0.140] N=1,393	0.043 [0.147] N=1,246	0.026 [0.158] N=1,064	-0.103 [0.183] N=875
WHO weight-for-length/height z score, 0-60 months	0.380	0.107 [0.228] N=1,303	0.072 [0.240] N=1,220	-0.041 [0.271] N=1,094	-0.065 [0.300] N=939	0.004 [0.335] N=772
WHO BMI-for-age z score	0.346	0.120 [0.248] N=1,272	0.111 [0.263] N=1,190	-0.007 [0.298] N=1,064	-0.091 [0.335] N=912	0.038 [0.369] N=752
Stunted: HAZ below -2	0.205 P=261	-0.049 [0.045] N=1,329	-0.056 [0.047] N=1,237	-0.030 [0.049] N=1,107	-0.013 [0.052] N=949	0.016 [0.059] N=784
Underweight: WAZ below -2	0.052 P=83	0.015 [0.020] N=1,502	0.004 [0.021] N=1,393	0.003 [0.023] N=1,246	0.005 [0.025] N=1,064	0.021 [0.034] N=875
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=151	-0.037 [0.037] N=1,303	-0.031 [0.037] N=1,220	-0.026 [0.039] N=1,094	-0.023 [0.043] N=939	-0.041 [0.045] N=772
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=249	0.003 [0.046] N=1,303	-0.007 [0.048] N=1,220	-0.014 [0.054] N=1,094	-0.008 [0.062] N=939	0.006 [0.069] N=772
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=121	-0.001 [0.033] N=1,303	-0.009 [0.034] N=1,220	-0.008 [0.037] N=1,094	-0.007 [0.041] N=939	0.002 [0.044] N=772
CIAF: any stunting, underweight, or wasting, 0-60 months	0.286 P=356	-0.066 [0.046] N=1,265	-0.071 [0.045] N=1,184	-0.057 [0.049] N=1,061	-0.041 [0.051] N=909	-0.032 [0.057] N=749
BMI-for-age z score below -2	0.122 P=160	-0.026 [0.037] N=1,272	-0.022 [0.037] N=1,190	-0.007 [0.040] N=1,064	0.020 [0.044] N=912	0.005 [0.047] N=752

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.398 [0.338] N=1,124	0.407 [0.388] N=1,032	0.488 [0.448] N=902	0.597 [0.472] N=744	-0.126 [0.630] N=579
WHO weight-for-age z score	-0.209 [0.188] N=1,278	-0.134 [0.200] N=1,169	-0.164 [0.218] N=1,022	-0.241 [0.271] N=840	**-0.846** [0.375] N=651**
WHO weight-for-length/height z score, 0-60 months	0.062 [0.255] N=1,099	-0.005 [0.272] N=1,016	-0.288 [0.338] N=890	-0.357 [0.421] N=735	-0.375 [0.512] N=568
WHO BMI-for-age z score	0.150 [0.268] N=1,076	0.143 [0.287] N=994	-0.132 [0.363] N=868	-0.365 [0.458] N=716	-0.230 [0.532] N=556
Stunted: HAZ below -2	-0.019 [0.060] N=1,124	-0.031 [0.066] N=1,032	0.016 [0.071] N=902	0.054 [0.078] N=744	0.152 [0.117] N=579
Underweight: WAZ below -2	0.038 [0.025] N=1,278	0.022 [0.027] N=1,169	0.022 [0.029] N=1,022	0.037 [0.033] N=840	**0.118** [0.056] N=651**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.028 [0.050] N=1,099	-0.015 [0.050] N=1,016	-0.004 [0.057] N=890	-0.008 [0.071] N=735	-0.063 [0.089] N=568
Overweight: WLZ/WHZ above +2, 0-60 months	-0.003 [0.055] N=1,099	-0.025 [0.057] N=1,016	-0.050 [0.068] N=890	-0.057 [0.088] N=735	-0.046 [0.096] N=568
Obese: WLZ/WHZ above +3, 0-60 months	0.004 [0.039] N=1,099	-0.008 [0.043] N=1,016	-0.006 [0.049] N=890	-0.000 [0.060] N=735	0.010 [0.067] N=568
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.005 [0.069] N=1,069	-0.010 [0.068] N=988	0.028 [0.079] N=865	0.063 [0.091] N=713	0.113 [0.138] N=553
BMI-for-age z score below -2	-0.035 [0.048] N=1,076	-0.029 [0.047] N=994	-0.001 [0.057] N=868	0.048 [0.074] N=716	0.016 [0.092] N=556

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.130	0.406 [0.247] N=1,329	0.407 [0.266] N=1,237	0.397 [0.291] N=1,107	0.376 [0.299] N=949	0.099 [0.327] N=784
WHO weight-for-age z score	0.255	-0.028 [0.134] N=1,502	0.035 [0.139] N=1,393	0.030 [0.146] N=1,246	0.009 [0.157] N=1,064	-0.137 [0.183] N=875
WHO weight-for-length/height z score, 0-60 months	0.380	0.090 [0.230] N=1,303	0.052 [0.242] N=1,220	-0.058 [0.275] N=1,094	-0.080 [0.308] N=939	-0.018 [0.349] N=772
WHO BMI-for-age z score	0.346	0.104 [0.249] N=1,272	0.090 [0.263] N=1,190	-0.027 [0.301] N=1,064	-0.114 [0.344] N=912	-0.006 [0.386] N=752
Stunted: HAZ below -2	0.205 P=261	-0.048 [0.044] N=1,329	-0.055 [0.047] N=1,237	-0.028 [0.049] N=1,107	-0.010 [0.052] N=949	0.015 [0.059] N=784
Underweight: WAZ below -2	0.052 P=83	0.016 [0.020] N=1,502	0.004 [0.021] N=1,393	0.004 [0.023] N=1,246	0.006 [0.025] N=1,064	0.023 [0.034] N=875
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=151	-0.035 [0.037] N=1,303	-0.028 [0.037] N=1,220	-0.026 [0.040] N=1,094	-0.025 [0.045] N=939	-0.047 [0.048] N=772
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=249	0.002 [0.046] N=1,303	-0.008 [0.048] N=1,220	-0.015 [0.055] N=1,094	-0.012 [0.063] N=939	0.002 [0.069] N=772
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=121	-0.001 [0.033] N=1,303	-0.009 [0.034] N=1,220	-0.008 [0.037] N=1,094	-0.007 [0.041] N=939	-0.000 [0.045] N=772
CIAF: any stunting, underweight, or wasting, 0-60 months	0.286 P=356	-0.063 [0.046] N=1,265	-0.068 [0.045] N=1,184	-0.055 [0.048] N=1,061	-0.041 [0.050] N=909	-0.040 [0.056] N=749
BMI-for-age z score below -2	0.122 P=160	-0.023 [0.037] N=1,272	-0.019 [0.037] N=1,190	-0.006 [0.041] N=1,064	0.021 [0.045] N=912	0.004 [0.049] N=752

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.432 [0.341] N=1,124	0.434 [0.388] N=1,032	0.505 [0.445] N=902	0.596 [0.484] N=744	-0.176 [0.661] N=579
WHO weight-for-age z score	-0.185 [0.186] N=1,278	-0.105 [0.198] N=1,169	-0.150 [0.217] N=1,022	-0.241 [0.271] N=840	**-0.869** [0.379] N=651**
WHO weight-for-length/height z score, 0-60 months	0.078 [0.253] N=1,099	0.026 [0.270] N=1,016	-0.257 [0.339] N=890	-0.354 [0.425] N=735	-0.382 [0.518] N=568
WHO BMI-for-age z score	0.174 [0.266] N=1,076	0.183 [0.285] N=994	-0.095 [0.360] N=868	-0.338 [0.457] N=716	-0.226 [0.539] N=556
Stunted: HAZ below -2	-0.024 [0.060] N=1,124	-0.033 [0.066] N=1,032	0.015 [0.070] N=902	0.050 [0.078] N=744	0.151 [0.116] N=579
Underweight: WAZ below -2	0.037 [0.025] N=1,278	0.020 [0.027] N=1,169	0.021 [0.029] N=1,022	0.038 [0.033] N=840	**0.121** [0.055] N=651**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.029 [0.049] N=1,099	-0.018 [0.049] N=1,016	-0.006 [0.056] N=890	-0.007 [0.069] N=735	-0.064 [0.086] N=568
Overweight: WLZ/WHZ above +2, 0-60 months	-0.004 [0.055] N=1,099	-0.022 [0.058] N=1,016	-0.046 [0.069] N=890	-0.052 [0.089] N=735	-0.048 [0.096] N=568
Obese: WLZ/WHZ above +3, 0-60 months	0.002 [0.039] N=1,099	-0.009 [0.043] N=1,016	-0.004 [0.050] N=890	0.001 [0.061] N=735	0.014 [0.069] N=568
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.009 [0.068] N=1,069	-0.013 [0.067] N=988	0.027 [0.077] N=865	0.063 [0.089] N=713	0.108 [0.134] N=553
BMI-for-age z score below -2	-0.037 [0.048] N=1,076	-0.033 [0.047] N=994	-0.004 [0.057] N=868	0.046 [0.073] N=716	0.012 [0.091] N=556

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.018	-0.174 [0.306] N=1,279	-0.273 [0.309] N=1,155	-0.253 [0.324] N=1,010	-0.471 [0.363] N=855	-0.492 [0.403] N=687
WHO weight-for-age z score	0.264	0.081 [0.145] N=1,439	0.132 [0.151] N=1,310	0.017 [0.154] N=1,144	0.076 [0.159] N=960	0.125 [0.184] N=768
WHO weight-for-length/height z score, 0-60 months	0.374	0.228 [0.204] N=1,263	0.229 [0.212] N=1,144	0.197 [0.227] N=1,002	0.392 [0.245] N=844	0.249 [0.274] N=680
WHO BMI-for-age z score	0.303	0.174 [0.237] N=1,226	0.205 [0.249] N=1,111	0.123 [0.265] N=974	0.279 [0.292] N=821	0.257 [0.324] N=665
Stunted: HAZ below -2	0.192 P=246	0.009 [0.050] N=1,279	0.019 [0.051] N=1,155	0.012 [0.053] N=1,010	0.029 [0.059] N=855	0.015 [0.065] N=687
Underweight: WAZ below -2	0.048 P=73	0.018 [0.022] N=1,439	0.007 [0.023] N=1,310	0.013 [0.025] N=1,144	0.003 [0.026] N=960	-0.016 [0.028] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=142	0.013 [0.036] N=1,263	0.006 [0.038] N=1,144	0.025 [0.039] N=1,002	-0.042 [0.039] N=844	-0.030 [0.041] N=680
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=246	**0.102** [0.045] N=1,263**	**0.095** [0.046] N=1,144**	**0.099** [0.050] N=1,002**	**0.119** [0.057] N=844**	**0.107* [0.062] N=680**
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=123	0.046 [0.037] N=1,263	0.041 [0.038] N=1,144	0.029 [0.042] N=1,002	0.038 [0.048] N=844	-0.001 [0.048] N=680
CIAF: any stunting, underweight, or wasting, 0-60 months	0.276 P=339	0.028 [0.054] N=1,225	0.017 [0.055] N=1,106	0.038 [0.058] N=969	-0.006 [0.062] N=819	-0.001 [0.065] N=659
BMI-for-age z score below -2	0.125 P=152	0.017 [0.042] N=1,226	0.009 [0.045] N=1,111	0.021 [0.049] N=974	-0.026 [0.049] N=821	-0.015 [0.051] N=665

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.068 [0.413] N=1,107	-0.066 [0.439] N=983	-0.014 [0.500] N=838	-0.528 [0.635] N=683	-0.526 [0.763] N=515
WHO weight-for-age z score	0.058 [0.209] N=1,253	0.139 [0.222] N=1,124	-0.076 [0.235] N=958	0.033 [0.255] N=774	0.143 [0.334] N=582
WHO weight-for-length/height z score, 0-60 months	0.078 [0.280] N=1,095	0.081 [0.299] N=976	0.039 [0.342] N=834	0.496 [0.385] N=676	0.084 [0.457] N=512
WHO BMI-for-age z score	-0.177 [0.301] N=1,067	-0.148 [0.327] N=952	-0.317 [0.370] N=815	-0.010 [0.449] N=662	-0.212 [0.529] N=506
Stunted: HAZ below -2	0.034 [0.081] N=1,107	0.050 [0.087] N=983	0.050 [0.097] N=838	0.110 [0.122] N=683	0.075 [0.147] N=515
Underweight: WAZ below -2	0.027 [0.034] N=1,253	0.009 [0.036] N=1,124	0.018 [0.041] N=958	-0.006 [0.045] N=774	-0.056 [0.047] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	0.075 [0.047] N=1,095	0.068 [0.052] N=976	**0.115* [0.060] N=834**	-0.018 [0.063] N=676	0.017 [0.069] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.123** [0.055] N=1,095**	**0.111* [0.058] N=976**	**0.131** [0.066] N=834**	**0.182** [0.081] N=676**	0.127 [0.090] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.096** [0.041] N=1,095**	**0.089** [0.041] N=976**	**0.083* [0.048] N=834**	**0.114* [0.065] N=676**	0.001 [0.076] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.101 [0.084] N=1,061	0.084 [0.090] N=942	0.149 [0.101] N=805	0.078 [0.121] N=655	0.078 [0.138] N=495
BMI-for-age z score below -2	**0.090** [0.045] N=1,067**	0.080 [0.052] N=952	**0.116* [0.062] N=815**	0.025 [0.061] N=662	0.070 [0.070] N=506

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.018	-0.181 [0.304] N=1,279	-0.279 [0.308] N=1,155	-0.257 [0.323] N=1,010	-0.476 [0.361] N=855	-0.495 [0.402] N=687
WHO weight-for-age z score	0.264	0.084 [0.145] N=1,439	0.134 [0.151] N=1,310	0.019 [0.153] N=1,144	0.075 [0.160] N=960	0.121 [0.184] N=768
WHO weight-for-length/height z score, 0-60 months	0.374	0.237 [0.204] N=1,263	0.237 [0.212] N=1,144	0.206 [0.227] N=1,002	0.395 [0.245] N=844	0.251 [0.273] N=680
WHO BMI-for-age z score	0.303	0.183 [0.237] N=1,226	0.212 [0.249] N=1,111	0.129 [0.265] N=974	0.280 [0.293] N=821	0.254 [0.326] N=665
Stunted: HAZ below -2	0.192 P=246	0.010 [0.050] N=1,279	0.019 [0.051] N=1,155	0.012 [0.053] N=1,010	0.030 [0.059] N=855	0.015 [0.065] N=687
Underweight: WAZ below -2	0.048 P=73	0.018 [0.022] N=1,439	0.007 [0.023] N=1,310	0.013 [0.025] N=1,144	0.003 [0.026] N=960	-0.015 [0.028] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=142	0.009 [0.036] N=1,263	0.002 [0.038] N=1,144	0.021 [0.040] N=1,002	-0.043 [0.039] N=844	-0.031 [0.041] N=680
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=246	**0.102** [0.045] N=1,263**	**0.096** [0.047] N=1,144**	**0.099* [0.050] N=1,002**	**0.118** [0.057] N=844**	**0.107* [0.062] N=680**
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=123	0.045 [0.037] N=1,263	0.040 [0.038] N=1,144	0.027 [0.042] N=1,002	0.037 [0.048] N=844	-0.001 [0.049] N=680
CIAF: any stunting, underweight, or wasting, 0-60 months	0.276 P=339	0.025 [0.054] N=1,225	0.013 [0.055] N=1,106	0.033 [0.058] N=969	-0.007 [0.062] N=819	-0.001 [0.066] N=659
BMI-for-age z score below -2	0.125 P=152	0.014 [0.042] N=1,226	0.006 [0.045] N=1,111	0.018 [0.048] N=974	-0.026 [0.049] N=821	-0.015 [0.051] N=665

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.058 [0.412] N=1,107	-0.075 [0.438] N=983	-0.012 [0.500] N=838	-0.549 [0.631] N=683	-0.560 [0.766] N=515
WHO weight-for-age z score	0.066 [0.206] N=1,253	0.147 [0.219] N=1,124	-0.064 [0.230] N=958	0.048 [0.250] N=774	0.178 [0.327] N=582
WHO weight-for-length/height z score, 0-60 months	0.103 [0.284] N=1,095	0.106 [0.301] N=976	0.073 [0.344] N=834	0.535 [0.387] N=676	0.188 [0.476] N=512
WHO BMI-for-age z score	-0.143 [0.304] N=1,067	-0.113 [0.328] N=952	-0.276 [0.369] N=815	0.035 [0.450] N=662	-0.102 [0.544] N=506
Stunted: HAZ below -2	0.034 [0.082] N=1,107	0.048 [0.087] N=983	0.046 [0.098] N=838	0.110 [0.123] N=683	0.076 [0.149] N=515
Underweight: WAZ below -2	0.026 [0.034] N=1,253	0.008 [0.036] N=1,124	0.018 [0.042] N=958	-0.007 [0.045] N=774	-0.059 [0.048] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	0.067 [0.048] N=1,095	0.057 [0.053] N=976	**0.101* [0.061] N=834**	-0.030 [0.064] N=676	-0.005 [0.072] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.125** [0.056] N=1,095**	**0.113* [0.059] N=976**	**0.132* [0.067] N=834**	**0.182** [0.082] N=676**	0.133 [0.093] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.094** [0.041] N=1,095**	**0.088** [0.041] N=976**	**0.079* [0.048] N=834**	**0.110* [0.064] N=676**	-0.002 [0.076] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.086] N=1,061	0.070 [0.091] N=942	0.128 [0.103] N=805	0.062 [0.123] N=655	0.055 [0.142] N=495
BMI-for-age z score below -2	**0.080* [0.046] N=1,067**	0.067 [0.052] N=952	**0.102* [0.060] N=815**	0.012 [0.061] N=662	0.049 [0.072] N=506

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.018	-0.190 [0.301] N=1,279	-0.278 [0.305] N=1,155	-0.253 [0.321] N=1,010	-0.473 [0.354] N=855	-0.501 [0.392] N=687
WHO weight-for-age z score	0.264	0.075 [0.146] N=1,439	0.131 [0.152] N=1,310	0.024 [0.155] N=1,144	0.077 [0.160] N=960	0.123 [0.185] N=768
WHO weight-for-length/height z score, 0-60 months	0.374	0.230 [0.206] N=1,263	0.232 [0.214] N=1,144	0.210 [0.228] N=1,002	0.393 [0.244] N=844	0.253 [0.273] N=680
WHO BMI-for-age z score	0.303	0.173 [0.239] N=1,226	0.201 [0.251] N=1,111	0.129 [0.266] N=974	0.274 [0.292] N=821	0.256 [0.324] N=665
Stunted: HAZ below -2	0.192 P=246	0.011 [0.049] N=1,279	0.019 [0.051] N=1,155	0.011 [0.053] N=1,010	0.029 [0.058] N=855	0.015 [0.063] N=687
Underweight: WAZ below -2	0.048 P=73	0.019 [0.023] N=1,439	0.007 [0.023] N=1,310	0.012 [0.025] N=1,144	0.003 [0.026] N=960	-0.015 [0.028] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=142	0.011 [0.037] N=1,263	0.003 [0.038] N=1,144	0.020 [0.040] N=1,002	-0.043 [0.040] N=844	-0.032 [0.042] N=680
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=246	**0.102** [0.045] N=1,263**	**0.095** [0.047] N=1,144**	**0.099** [0.050] N=1,002**	**0.118** [0.057] N=844**	**0.107* [0.062] N=680**
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=123	0.045 [0.037] N=1,263	0.040 [0.037] N=1,144	0.028 [0.042] N=1,002	0.037 [0.048] N=844	-0.001 [0.049] N=680
CIAF: any stunting, underweight, or wasting, 0-60 months	0.276 P=339	0.029 [0.054] N=1,225	0.014 [0.055] N=1,106	0.033 [0.058] N=969	-0.008 [0.062] N=819	-0.003 [0.066] N=659
BMI-for-age z score below -2	0.125 P=152	0.017 [0.043] N=1,226	0.008 [0.045] N=1,111	0.019 [0.049] N=974	-0.025 [0.049] N=821	-0.013 [0.052] N=665

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.071 [0.409] N=1,107	-0.061 [0.437] N=983	0.033 [0.494] N=838	-0.501 [0.627] N=683	-0.485 [0.749] N=515
WHO weight-for-age z score	0.048 [0.210] N=1,253	0.138 [0.223] N=1,124	-0.057 [0.236] N=958	0.052 [0.257] N=774	0.170 [0.337] N=582
WHO weight-for-length/height z score, 0-60 months	0.071 [0.284] N=1,095	0.089 [0.303] N=976	0.054 [0.343] N=834	0.505 [0.386] N=676	0.145 [0.475] N=512
WHO BMI-for-age z score	-0.167 [0.305] N=1,067	-0.127 [0.330] N=952	-0.299 [0.369] N=815	0.005 [0.447] N=662	-0.130 [0.541] N=506
Stunted: HAZ below -2	0.027 [0.081] N=1,107	0.041 [0.087] N=983	0.035 [0.097] N=838	0.094 [0.122] N=683	0.058 [0.149] N=515
Underweight: WAZ below -2	0.025 [0.035] N=1,253	0.005 [0.037] N=1,124	0.012 [0.042] N=958	-0.013 [0.046] N=774	-0.066 [0.051] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.049] N=1,095	0.056 [0.054] N=976	0.101 [0.061] N=834	-0.030 [0.064] N=676	-0.005 [0.073] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.125** [0.056] N=1,095**	**0.113* [0.059] N=976**	**0.130* [0.067] N=834**	**0.178** [0.083] N=676**	0.128 [0.096] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.091** [0.041] N=1,095**	**0.085** [0.042] N=976**	0.077 [0.048] N=834	0.104 [0.065] N=676	-0.012 [0.077] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.089 [0.086] N=1,061	0.067 [0.092] N=942	0.123 [0.105] N=805	0.052 [0.125] N=655	0.048 [0.145] N=495
BMI-for-age z score below -2	**0.080* [0.047] N=1,067**	0.067 [0.053] N=952	**0.103* [0.060] N=815**	0.012 [0.061] N=662	0.047 [0.073] N=506

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.018	-0.152 [0.302] N=1,279	-0.236 [0.307] N=1,155	-0.208 [0.325] N=1,010	-0.459 [0.360] N=855	-0.489 [0.402] N=687
WHO weight-for-age z score	0.264	0.078 [0.147] N=1,439	0.133 [0.154] N=1,310	0.031 [0.157] N=1,144	0.081 [0.163] N=960	0.131 [0.186] N=768
WHO weight-for-length/height z score, 0-60 months	0.374	0.195 [0.205] N=1,263	0.199 [0.212] N=1,144	0.173 [0.226] N=1,002	0.381 [0.245] N=844	0.253 [0.276] N=680
WHO BMI-for-age z score	0.303	0.135 [0.236] N=1,226	0.166 [0.247] N=1,111	0.087 [0.261] N=974	0.252 [0.289] N=821	0.257 [0.325] N=665
Stunted: HAZ below -2	0.192 P=246	0.005 [0.049] N=1,279	0.011 [0.051] N=1,155	0.003 [0.054] N=1,010	0.025 [0.059] N=855	0.012 [0.064] N=687
Underweight: WAZ below -2	0.048 P=73	0.018 [0.023] N=1,439	0.006 [0.023] N=1,310	0.010 [0.025] N=1,144	0.001 [0.027] N=960	-0.021 [0.029] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=142	0.015 [0.036] N=1,263	0.007 [0.038] N=1,144	0.027 [0.039] N=1,002	-0.040 [0.039] N=844	-0.032 [0.042] N=680
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=246	**0.097** [0.044] N=1,263**	**0.090* [0.046] N=1,144**	**0.096* [0.050] N=1,002**	**0.118** [0.056] N=844**	**0.108* [0.061] N=680**
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=123	0.045 [0.036] N=1,263	0.041 [0.037] N=1,144	0.028 [0.041] N=1,002	0.039 [0.047] N=844	0.001 [0.048] N=680
CIAF: any stunting, underweight, or wasting, 0-60 months	0.276 P=339	0.028 [0.054] N=1,225	0.014 [0.055] N=1,106	0.032 [0.058] N=969	-0.007 [0.062] N=819	-0.000 [0.064] N=659
BMI-for-age z score below -2	0.125 P=152	0.017 [0.042] N=1,226	0.009 [0.045] N=1,111	0.021 [0.049] N=974	-0.025 [0.049] N=821	-0.018 [0.052] N=665

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.095 [0.403] N=1,107	-0.022 [0.429] N=983	0.072 [0.491] N=838	-0.526 [0.627] N=683	-0.635 [0.741] N=515
WHO weight-for-age z score	0.068 [0.204] N=1,253	0.157 [0.218] N=1,124	-0.026 [0.228] N=958	0.072 [0.248] N=774	0.195 [0.313] N=582
WHO weight-for-length/height z score, 0-60 months	0.053 [0.283] N=1,095	0.060 [0.299] N=976	0.019 [0.338] N=834	0.507 [0.391] N=676	0.226 [0.497] N=512
WHO BMI-for-age z score	-0.209 [0.303] N=1,067	-0.177 [0.325] N=952	-0.355 [0.362] N=815	-0.024 [0.446] N=662	-0.058 [0.562] N=506
Stunted: HAZ below -2	0.025 [0.080] N=1,107	0.034 [0.086] N=983	0.026 [0.098] N=838	0.091 [0.125] N=683	0.069 [0.150] N=515
Underweight: WAZ below -2	0.026 [0.035] N=1,253	0.006 [0.036] N=1,124	0.011 [0.043] N=958	-0.013 [0.046] N=774	-0.072 [0.050] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	0.073 [0.048] N=1,095	0.065 [0.053] N=976	**0.113* [0.060] N=834**	-0.026 [0.063] N=676	-0.010 [0.072] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.118** [0.056] N=1,095**	**0.106* [0.059] N=976**	**0.125* [0.067] N=834**	**0.175** [0.084] N=676**	0.135 [0.097] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.093** [0.041] N=1,095**	**0.087** [0.042] N=976**	0.077 [0.048] N=834	0.107 [0.066] N=676	-0.005 [0.078] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.086] N=1,061	0.067 [0.093] N=942	0.120 [0.106] N=805	0.048 [0.127] N=655	0.051 [0.144] N=495
BMI-for-age z score below -2	**0.084* [0.046] N=1,067**	0.073 [0.052] N=952	**0.112* [0.060] N=815**	0.016 [0.060] N=662	0.041 [0.072] N=506

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.029	-0.243 [0.300] N=1,195	-0.352 [0.307] N=1,078	-0.348 [0.323] N=947	**-0.589* [0.354] N=799**	**-0.669* [0.392] N=644**
WHO weight-for-age z score	0.256	0.128 [0.151] N=1,343	0.200 [0.157] N=1,221	0.055 [0.161] N=1,070	0.094 [0.168] N=897	0.088 [0.187] N=721
WHO weight-for-length/height z score, 0-60 months	0.377	0.348 [0.219] N=1,179	0.372 [0.232] N=1,067	0.310 [0.249] N=937	**0.472* [0.269] N=788**	0.322 [0.292] N=638
WHO BMI-for-age z score	0.304	0.222 [0.251] N=1,144	0.277 [0.268] N=1,036	0.164 [0.286] N=911	0.277 [0.312] N=767	0.221 [0.347] N=624
Stunted: HAZ below -2	0.194 P=231	0.017 [0.052] N=1,195	0.033 [0.053] N=1,078	0.023 [0.056] N=947	0.042 [0.060] N=799	0.040 [0.066] N=644
Underweight: WAZ below -2	0.049 P=70	0.021 [0.023] N=1,343	0.008 [0.024] N=1,221	0.012 [0.026] N=1,070	0.004 [0.028] N=897	-0.008 [0.031] N=721
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=133	0.010 [0.038] N=1,179	0.002 [0.040] N=1,067	0.024 [0.041] N=937	-0.033 [0.042] N=788	-0.019 [0.044] N=638
Overweight: WLZ/WHZ above +2, 0-60 months	0.191 P=230	**0.120** [0.047] N=1,179**	**0.116** [0.049] N=1,067**	**0.118** [0.053] N=937**	**0.133** [0.060] N=788**	**0.122* [0.064] N=638**
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=118	0.057 [0.037] N=1,179	0.052 [0.038] N=1,067	0.041 [0.043] N=937	0.057 [0.049] N=788	0.022 [0.049] N=638
CIAF: any stunting, underweight, or wasting, 0-60 months	0.279 P=320	0.034 [0.057] N=1,145	0.024 [0.058] N=1,033	0.046 [0.061] N=908	0.008 [0.064] N=765	0.032 [0.066] N=618
BMI-for-age z score below -2	0.125 P=142	0.013 [0.043] N=1,144	0.001 [0.046] N=1,036	0.013 [0.049] N=911	-0.021 [0.051] N=767	-0.005 [0.053] N=624

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.172 [0.407] N=1,033	0.031 [0.440] N=916	0.078 [0.504] N=785	-0.446 [0.624] N=637	-0.437 [0.733] N=482
WHO weight-for-age z score	0.163 [0.211] N=1,168	0.271 [0.221] N=1,046	0.014 [0.235] N=895	0.143 [0.255] N=722	0.223 [0.320] N=546
WHO weight-for-length/height z score, 0-60 months	0.201 [0.303] N=1,020	0.235 [0.325] N=908	0.137 [0.371] N=778	0.545 [0.421] N=629	0.211 [0.519] N=479
WHO BMI-for-age z score	-0.095 [0.327] N=993	-0.031 [0.359] N=885	-0.251 [0.404] N=760	0.035 [0.488] N=616	-0.145 [0.599] N=473
Stunted: HAZ below -2	0.017 [0.087] N=1,033	0.040 [0.094] N=916	0.031 [0.104] N=785	0.081 [0.129] N=637	0.054 [0.160] N=482
Underweight: WAZ below -2	0.029 [0.036] N=1,168	0.006 [0.038] N=1,046	0.013 [0.044] N=895	-0.013 [0.049] N=722	-0.058 [0.052] N=546
Wasted: WLZ/WHZ below -2, 0-60 months	0.064 [0.052] N=1,020	0.050 [0.056] N=908	0.097 [0.063] N=778	-0.028 [0.069] N=629	-0.004 [0.080] N=479
Overweight: WLZ/WHZ above +2, 0-60 months	**0.141** [0.061] N=1,020**	**0.130** [0.065] N=908**	**0.144** [0.073] N=778**	**0.179* [0.092] N=629**	0.141 [0.106] N=479
Obese: WLZ/WHZ above +3, 0-60 months	**0.090** [0.044] N=1,020**	**0.082* [0.045] N=908**	0.071 [0.053] N=778	0.100 [0.071] N=629	-0.014 [0.085] N=479
CIAF: any stunting, underweight, or wasting, 0-60 months	0.077 [0.094] N=990	0.060 [0.101] N=878	0.118 [0.113] N=753	0.034 [0.136] N=610	0.044 [0.162] N=463
BMI-for-age z score below -2	**0.088* [0.049] N=993**	0.071 [0.056] N=885	**0.105* [0.062] N=760**	0.022 [0.065] N=616	0.069 [0.076] N=473

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.029	-0.273 [0.307] N=1,195	-0.381 [0.315] N=1,078	-0.370 [0.330] N=947	**-0.634* [0.362] N=799**	**-0.709* [0.404] N=644**
WHO weight-for-age z score	0.256	0.127 [0.152] N=1,343	0.195 [0.158] N=1,221	0.053 [0.161] N=1,070	0.088 [0.168] N=897	0.085 [0.190] N=721
WHO weight-for-length/height z score, 0-60 months	0.377	**0.367* [0.221] N=1,179**	0.382 [0.233] N=1,067	0.322 [0.250] N=937	**0.497* [0.268] N=788**	0.348 [0.296] N=638
WHO BMI-for-age z score	0.304	0.243 [0.257] N=1,144	0.287 [0.273] N=1,036	0.177 [0.291] N=911	0.303 [0.317] N=767	0.247 [0.357] N=624
Stunted: HAZ below -2	0.194 P=231	0.021 [0.052] N=1,195	0.034 [0.053] N=1,078	0.023 [0.055] N=947	0.045 [0.060] N=799	0.044 [0.067] N=644
Underweight: WAZ below -2	0.049 P=70	0.022 [0.023] N=1,343	0.008 [0.024] N=1,221	0.012 [0.026] N=1,070	0.005 [0.028] N=897	-0.007 [0.031] N=721
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=133	0.006 [0.039] N=1,179	-0.002 [0.041] N=1,067	0.019 [0.042] N=937	-0.039 [0.043] N=788	-0.025 [0.045] N=638
Overweight: WLZ/WHZ above +2, 0-60 months	0.191 P=230	**0.121*** [0.046] N=1,179**	**0.117** [0.048] N=1,067**	**0.118** [0.052] N=937**	**0.135** [0.059] N=788**	**0.125* [0.064] N=638**
Obese: WLZ/WHZ above +3, 0-60 months	0.096 P=118	0.057 [0.037] N=1,179	0.052 [0.037] N=1,067	0.041 [0.042] N=937	0.057 [0.049] N=788	0.023 [0.049] N=638
CIAF: any stunting, underweight, or wasting, 0-60 months	0.279 P=320	0.032 [0.056] N=1,145	0.021 [0.057] N=1,033	0.042 [0.060] N=908	0.005 [0.063] N=765	0.031 [0.065] N=618
BMI-for-age z score below -2	0.125 P=142	0.009 [0.045] N=1,144	-0.002 [0.048] N=1,036	0.010 [0.051] N=911	-0.027 [0.053] N=767	-0.009 [0.056] N=624

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.129 [0.417] N=1,033	-0.003 [0.447] N=916	0.045 [0.512] N=785	-0.558 [0.637] N=637	-0.607 [0.769] N=482
WHO weight-for-age z score	0.154 [0.214] N=1,168	0.256 [0.226] N=1,046	0.001 [0.239] N=895	0.118 [0.261] N=722	0.182 [0.337] N=546
WHO weight-for-length/height z score, 0-60 months	0.201 [0.302] N=1,020	0.213 [0.321] N=908	0.121 [0.368] N=778	0.579 [0.417] N=629	0.204 [0.508] N=479
WHO BMI-for-age z score	-0.094 [0.325] N=993	-0.056 [0.353] N=885	-0.267 [0.400] N=760	0.066 [0.481] N=616	-0.128 [0.590] N=473
Stunted: HAZ below -2	0.021 [0.087] N=1,033	0.040 [0.093] N=916	0.030 [0.104] N=785	0.091 [0.129] N=637	0.068 [0.160] N=482
Underweight: WAZ below -2	0.029 [0.036] N=1,168	0.008 [0.038] N=1,046	0.015 [0.044] N=895	-0.009 [0.049] N=722	-0.051 [0.054] N=546
Wasted: WLZ/WHZ below -2, 0-60 months	0.064 [0.051] N=1,020	0.054 [0.056] N=908	0.099 [0.063] N=778	-0.033 [0.068] N=629	-0.005 [0.079] N=479
Overweight: WLZ/WHZ above +2, 0-60 months	**0.143** [0.060] N=1,020**	**0.130** [0.064] N=908**	**0.144** [0.072] N=778**	**0.186** [0.091] N=629**	0.143 [0.105] N=479
Obese: WLZ/WHZ above +3, 0-60 months	**0.091** [0.044] N=1,020**	**0.080* [0.044] N=908**	0.069 [0.051] N=778	0.104 [0.070] N=629	-0.008 [0.085] N=479
CIAF: any stunting, underweight, or wasting, 0-60 months	0.080 [0.094] N=990	0.062 [0.101] N=878	0.119 [0.112] N=753	0.035 [0.135] N=610	0.048 [0.160] N=463
BMI-for-age z score below -2	**0.089* [0.048] N=993**	0.076 [0.055] N=885	**0.109* [0.061] N=760**	0.018 [0.062] N=616	0.067 [0.073] N=473

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.