Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,505	-0.001 [0.023] N=1,603	0.015 [0.023] N=1,478	0.006 [0.025] N=1,333	0.001 [0.027] N=1,134	-0.006 [0.032] N=930
Vegetables consumed at least weekly	0.900 P=1,442	0.015 [0.028] N=1,603	0.017 [0.028] N=1,478	-0.001 [0.029] N=1,333	0.002 [0.029] N=1,134	-0.038 [0.033] N=930
Meat fish or poultry consumed at least weekly	0.797 P=1,286	**0.080* [0.044] N=1,603**	**0.098** [0.046] N=1,478**	0.080 [0.050] N=1,333	0.063 [0.055] N=1,134	0.056 [0.063] N=930
Legumes consumed at least weekly	0.795 P=1,280	-0.053 [0.040] N=1,603	-0.060 [0.041] N=1,478	-0.069 [0.042] N=1,333	-0.071 [0.046] N=1,134	**-0.121** [0.052] N=930**
Animal products consumed at least weekly	0.912 P=1,463	0.034 [0.030] N=1,603	0.037 [0.032] N=1,478	0.026 [0.035] N=1,333	0.024 [0.037] N=1,134	0.012 [0.041] N=930
Sweets consumed at least weekly	0.683 P=1,092	-0.016 [0.039] N=1,603	0.003 [0.042] N=1,478	0.009 [0.044] N=1,333	-0.001 [0.047] N=1,134	0.021 [0.054] N=930
Snack foods consumed at least weekly	0.559 P=887	-0.043 [0.041] N=1,603	-0.031 [0.044] N=1,478	-0.013 [0.047] N=1,333	-0.019 [0.053] N=1,134	-0.031 [0.057] N=930
Packaged soft drinks consumed at least weekly	0.341 P=536	0.004 [0.050] N=1,603	0.012 [0.054] N=1,478	0.012 [0.057] N=1,333	0.027 [0.061] N=1,134	0.015 [0.071] N=930
Grains consumed at least weekly	0.914 P=1,464	0.013 [0.028] N=1,603	0.026 [0.030] N=1,478	0.008 [0.031] N=1,333	0.015 [0.033] N=1,134	0.003 [0.040] N=930
Child was ever breastfed	0.964 P=1,542	0.009 [0.016] N=1,604	0.012 [0.017] N=1,479	0.018 [0.018] N=1,333	0.024 [0.021] N=1,134	0.013 [0.021] N=930

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.018 [0.035] N=1,365	0.050 [0.035] N=1,240	0.040 [0.038] N=1,095	0.039 [0.043] N=896	0.045 [0.061] N=692
Vegetables consumed at least weekly	0.052 [0.041] N=1,365	0.063 [0.044] N=1,240	0.041 [0.048] N=1,095	0.063 [0.055] N=896	-0.026 [0.075] N=692
Meat fish or poultry consumed at least weekly	0.078 [0.056] N=1,365	**0.108* [0.060] N=1,240**	0.080 [0.067] N=1,095	0.024 [0.076] N=896	-0.015 [0.091] N=692
Legumes consumed at least weekly	0.019 [0.060] N=1,365	0.015 [0.063] N=1,240	0.020 [0.067] N=1,095	0.045 [0.082] N=896	-0.027 [0.114] N=692
Animal products consumed at least weekly	0.043 [0.041] N=1,365	0.050 [0.044] N=1,240	0.031 [0.052] N=1,095	0.040 [0.061] N=896	0.001 [0.076] N=692
Sweets consumed at least weekly	-0.032 [0.053] N=1,365	-0.003 [0.057] N=1,240	0.001 [0.066] N=1,095	-0.026 [0.074] N=896	0.047 [0.100] N=692
Snack foods consumed at least weekly	-0.052 [0.052] N=1,365	-0.036 [0.056] N=1,240	-0.007 [0.067] N=1,095	-0.021 [0.084] N=896	-0.057 [0.098] N=692
Packaged soft drinks consumed at least weekly	-0.020 [0.059] N=1,365	-0.011 [0.066] N=1,240	-0.011 [0.070] N=1,095	0.014 [0.082] N=896	-0.037 [0.099] N=692
Grains consumed at least weekly	0.049 [0.046] N=1,365	0.079 [0.052] N=1,240	0.057 [0.056] N=1,095	0.095 [0.068] N=896	0.109 [0.101] N=692
Child was ever breastfed	0.017 [0.026] N=1,366	0.023 [0.030] N=1,241	0.034 [0.036] N=1,095	0.050 [0.046] N=896	0.045 [0.060] N=692

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,505	-0.001 [0.023] N=1,603	0.016 [0.023] N=1,478	0.007 [0.025] N=1,333	0.002 [0.027] N=1,134	-0.005 [0.032] N=930
Vegetables consumed at least weekly	0.900 P=1,442	0.015 [0.028] N=1,603	0.017 [0.028] N=1,478	-0.000 [0.029] N=1,333	0.002 [0.029] N=1,134	-0.038 [0.033] N=930
Meat fish or poultry consumed at least weekly	0.797 P=1,286	**0.080* [0.044] N=1,603**	**0.098** [0.046] N=1,478**	0.081 [0.050] N=1,333	0.064 [0.056] N=1,134	0.057 [0.064] N=930
Legumes consumed at least weekly	0.795 P=1,280	-0.053 [0.040] N=1,603	-0.060 [0.041] N=1,478	-0.068 [0.042] N=1,333	-0.070 [0.046] N=1,134	**-0.121** [0.052] N=930**
Animal products consumed at least weekly	0.912 P=1,463	0.034 [0.030] N=1,603	0.037 [0.031] N=1,478	0.028 [0.035] N=1,333	0.025 [0.037] N=1,134	0.012 [0.041] N=930
Sweets consumed at least weekly	0.683 P=1,092	-0.016 [0.039] N=1,603	0.004 [0.042] N=1,478	0.009 [0.044] N=1,333	-0.000 [0.047] N=1,134	0.021 [0.054] N=930
Snack foods consumed at least weekly	0.559 P=887	-0.043 [0.042] N=1,603	-0.030 [0.044] N=1,478	-0.012 [0.048] N=1,333	-0.018 [0.053] N=1,134	-0.031 [0.058] N=930
Packaged soft drinks consumed at least weekly	0.341 P=536	0.004 [0.050] N=1,603	0.012 [0.054] N=1,478	0.012 [0.057] N=1,333	0.027 [0.061] N=1,134	0.014 [0.071] N=930
Grains consumed at least weekly	0.914 P=1,464	0.013 [0.028] N=1,603	0.026 [0.030] N=1,478	0.008 [0.031] N=1,333	0.014 [0.032] N=1,134	0.003 [0.040] N=930
Child was ever breastfed	0.964 P=1,542	0.009 [0.016] N=1,604	0.012 [0.017] N=1,479	0.018 [0.018] N=1,333	0.024 [0.021] N=1,134	0.013 [0.021] N=930

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.018 [0.035] N=1,365	0.051 [0.035] N=1,240	0.041 [0.038] N=1,095	0.040 [0.044] N=896	0.047 [0.062] N=692
Vegetables consumed at least weekly	0.052 [0.041] N=1,365	0.064 [0.044] N=1,240	0.041 [0.048] N=1,095	0.064 [0.055] N=896	-0.026 [0.075] N=692
Meat fish or poultry consumed at least weekly	0.077 [0.056] N=1,365	**0.108* [0.060] N=1,240**	0.081 [0.067] N=1,095	0.026 [0.076] N=896	-0.010 [0.091] N=692
Legumes consumed at least weekly	0.019 [0.060] N=1,365	0.015 [0.063] N=1,240	0.023 [0.067] N=1,095	0.048 [0.083] N=896	-0.025 [0.114] N=692
Animal products consumed at least weekly	0.043 [0.041] N=1,365	0.052 [0.044] N=1,240	0.035 [0.052] N=1,095	0.044 [0.060] N=896	0.004 [0.075] N=692
Sweets consumed at least weekly	-0.033 [0.053] N=1,365	-0.003 [0.057] N=1,240	0.000 [0.066] N=1,095	-0.026 [0.074] N=896	0.047 [0.100] N=692
Snack foods consumed at least weekly	-0.051 [0.053] N=1,365	-0.033 [0.057] N=1,240	-0.000 [0.068] N=1,095	-0.016 [0.085] N=896	-0.051 [0.098] N=692
Packaged soft drinks consumed at least weekly	-0.020 [0.059] N=1,365	-0.011 [0.066] N=1,240	-0.012 [0.070] N=1,095	0.014 [0.082] N=896	-0.039 [0.099] N=692
Grains consumed at least weekly	0.048 [0.046] N=1,365	0.080 [0.052] N=1,240	0.057 [0.056] N=1,095	0.094 [0.067] N=896	0.108 [0.101] N=692
Child was ever breastfed	0.017 [0.026] N=1,366	0.024 [0.030] N=1,241	0.035 [0.036] N=1,095	0.050 [0.046] N=896	0.046 [0.060] N=692

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,505	-0.009 [0.023] N=1,603	0.006 [0.023] N=1,478	-0.002 [0.025] N=1,333	-0.006 [0.028] N=1,134	-0.014 [0.034] N=930
Vegetables consumed at least weekly	0.900 P=1,442	0.009 [0.027] N=1,603	0.010 [0.028] N=1,478	-0.006 [0.029] N=1,333	-0.003 [0.029] N=1,134	-0.044 [0.033] N=930
Meat fish or poultry consumed at least weekly	0.797 P=1,286	0.058 [0.043] N=1,603	0.071 [0.045] N=1,478	0.059 [0.050] N=1,333	0.044 [0.054] N=1,134	0.035 [0.062] N=930
Legumes consumed at least weekly	0.795 P=1,280	-0.060 [0.040] N=1,603	**-0.069* [0.041] N=1,478**	**-0.076* [0.042] N=1,333**	-0.076 [0.046] N=1,134	**-0.125** [0.052] N=930**
Animal products consumed at least weekly	0.912 P=1,463	0.029 [0.031] N=1,603	0.031 [0.032] N=1,478	0.021 [0.035] N=1,333	0.020 [0.038] N=1,134	0.005 [0.042] N=930
Sweets consumed at least weekly	0.683 P=1,092	-0.023 [0.039] N=1,603	-0.004 [0.042] N=1,478	0.001 [0.044] N=1,333	-0.007 [0.047] N=1,134	0.013 [0.053] N=930
Snack foods consumed at least weekly	0.559 P=887	-0.042 [0.042] N=1,603	-0.030 [0.044] N=1,478	-0.013 [0.047] N=1,333	-0.019 [0.052] N=1,134	-0.031 [0.057] N=930
Packaged soft drinks consumed at least weekly	0.341 P=536	0.016 [0.048] N=1,603	0.029 [0.052] N=1,478	0.026 [0.055] N=1,333	0.039 [0.059] N=1,134	0.028 [0.066] N=930
Grains consumed at least weekly	0.914 P=1,464	0.007 [0.028] N=1,603	0.020 [0.030] N=1,478	0.003 [0.031] N=1,333	0.008 [0.032] N=1,134	-0.002 [0.041] N=930
Child was ever breastfed	0.964 P=1,542	0.005 [0.016] N=1,604	0.006 [0.017] N=1,479	0.013 [0.018] N=1,333	0.020 [0.020] N=1,134	0.011 [0.021] N=930

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.011 [0.034] N=1,365	0.042 [0.034] N=1,240	0.036 [0.037] N=1,095	0.046 [0.043] N=896	0.057 [0.062] N=692
Vegetables consumed at least weekly	0.048 [0.041] N=1,365	0.058 [0.044] N=1,240	0.038 [0.048] N=1,095	0.069 [0.054] N=896	-0.019 [0.075] N=692
Meat fish or poultry consumed at least weekly	0.059 [0.055] N=1,365	0.080 [0.058] N=1,240	0.062 [0.067] N=1,095	0.019 [0.076] N=896	-0.008 [0.094] N=692
Legumes consumed at least weekly	0.012 [0.060] N=1,365	0.005 [0.064] N=1,240	0.015 [0.068] N=1,095	0.051 [0.083] N=896	-0.019 [0.116] N=692
Animal products consumed at least weekly	0.039 [0.041] N=1,365	0.046 [0.044] N=1,240	0.031 [0.052] N=1,095	0.049 [0.061] N=896	0.015 [0.075] N=692
Sweets consumed at least weekly	-0.036 [0.054] N=1,365	-0.008 [0.059] N=1,240	-0.003 [0.067] N=1,095	-0.023 [0.075] N=896	0.055 [0.103] N=692
Snack foods consumed at least weekly	-0.048 [0.052] N=1,365	-0.031 [0.056] N=1,240	0.002 [0.066] N=1,095	-0.011 [0.082] N=896	-0.045 [0.096] N=692
Packaged soft drinks consumed at least weekly	-0.009 [0.058] N=1,365	0.008 [0.063] N=1,240	0.004 [0.068] N=1,095	0.027 [0.080] N=896	-0.032 [0.094] N=692
Grains consumed at least weekly	0.041 [0.046] N=1,365	0.073 [0.051] N=1,240	0.052 [0.056] N=1,095	0.092 [0.067] N=896	0.106 [0.100] N=692
Child was ever breastfed	0.014 [0.026] N=1,366	0.019 [0.030] N=1,241	0.032 [0.036] N=1,095	0.051 [0.046] N=896	0.043 [0.060] N=692

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,505	-0.010 [0.023] N=1,603	0.006 [0.023] N=1,478	-0.003 [0.025] N=1,333	-0.007 [0.027] N=1,134	-0.013 [0.033] N=930
Vegetables consumed at least weekly	0.900 P=1,442	0.009 [0.027] N=1,603	0.011 [0.028] N=1,478	-0.005 [0.029] N=1,333	-0.003 [0.029] N=1,134	-0.041 [0.033] N=930
Meat fish or poultry consumed at least weekly	0.797 P=1,286	0.053 [0.043] N=1,603	0.066 [0.045] N=1,478	0.050 [0.049] N=1,333	0.035 [0.054] N=1,134	0.025 [0.062] N=930
Legumes consumed at least weekly	0.795 P=1,280	-0.063 [0.041] N=1,603	**-0.074* [0.042] N=1,478**	**-0.084* [0.043] N=1,333**	**-0.082* [0.048] N=1,134**	**-0.129** [0.053] N=930**
Animal products consumed at least weekly	0.912 P=1,463	0.027 [0.031] N=1,603	0.031 [0.032] N=1,478	0.021 [0.035] N=1,333	0.017 [0.038] N=1,134	0.003 [0.041] N=930
Sweets consumed at least weekly	0.683 P=1,092	-0.022 [0.039] N=1,603	-0.005 [0.041] N=1,478	-0.003 [0.043] N=1,333	-0.010 [0.046] N=1,134	0.009 [0.054] N=930
Snack foods consumed at least weekly	0.559 P=887	-0.040 [0.042] N=1,603	-0.028 [0.044] N=1,478	-0.015 [0.047] N=1,333	-0.019 [0.052] N=1,134	-0.033 [0.056] N=930
Packaged soft drinks consumed at least weekly	0.341 P=536	0.022 [0.048] N=1,603	0.031 [0.052] N=1,478	0.028 [0.055] N=1,333	0.044 [0.058] N=1,134	0.036 [0.065] N=930
Grains consumed at least weekly	0.914 P=1,464	0.007 [0.028] N=1,603	0.020 [0.029] N=1,478	0.002 [0.031] N=1,333	0.008 [0.032] N=1,134	-0.002 [0.041] N=930
Child was ever breastfed	0.964 P=1,542	0.004 [0.016] N=1,604	0.006 [0.017] N=1,479	0.012 [0.018] N=1,333	0.019 [0.021] N=1,134	0.009 [0.021] N=930

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.035] N=1,365	0.042 [0.034] N=1,240	0.034 [0.038] N=1,095	0.039 [0.044] N=896	0.044 [0.063] N=692
Vegetables consumed at least weekly	0.047 [0.041] N=1,365	0.057 [0.044] N=1,240	0.036 [0.048] N=1,095	0.061 [0.055] N=896	-0.036 [0.076] N=692
Meat fish or poultry consumed at least weekly	0.054 [0.056] N=1,365	0.075 [0.059] N=1,240	0.056 [0.068] N=1,095	0.014 [0.078] N=896	-0.022 [0.098] N=692
Legumes consumed at least weekly	0.015 [0.060] N=1,365	0.007 [0.063] N=1,240	0.016 [0.066] N=1,095	0.061 [0.082] N=896	-0.016 [0.120] N=692
Animal products consumed at least weekly	0.038 [0.041] N=1,365	0.049 [0.044] N=1,240	0.031 [0.052] N=1,095	0.041 [0.060] N=896	0.000 [0.079] N=692
Sweets consumed at least weekly	-0.031 [0.053] N=1,365	-0.006 [0.057] N=1,240	-0.002 [0.065] N=1,095	-0.013 [0.075] N=896	0.064 [0.104] N=692
Snack foods consumed at least weekly	-0.042 [0.052] N=1,365	-0.027 [0.055] N=1,240	0.005 [0.064] N=1,095	0.006 [0.081] N=896	-0.022 [0.095] N=692
Packaged soft drinks consumed at least weekly	-0.000 [0.058] N=1,365	0.010 [0.063] N=1,240	0.007 [0.068] N=1,095	0.036 [0.081] N=896	-0.015 [0.093] N=692
Grains consumed at least weekly	0.041 [0.044] N=1,365	0.071 [0.049] N=1,240	0.050 [0.054] N=1,095	0.088 [0.063] N=896	0.095 [0.094] N=692
Child was ever breastfed	0.015 [0.026] N=1,366	0.020 [0.030] N=1,241	0.031 [0.036] N=1,095	0.051 [0.048] N=896	0.047 [0.061] N=692

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,397	-0.009 [0.025] N=1,488	0.002 [0.025] N=1,378	-0.006 [0.026] N=1,241	-0.009 [0.029] N=1,060	-0.020 [0.034] N=869
Vegetables consumed at least weekly	0.901 P=1,340	0.001 [0.029] N=1,488	-0.000 [0.029] N=1,378	-0.022 [0.030] N=1,241	-0.013 [0.030] N=1,060	-0.053 [0.033] N=869
Meat fish or poultry consumed at least weekly	0.795 P=1,190	0.066 [0.042] N=1,488	**0.073* [0.043] N=1,378**	0.048 [0.047] N=1,241	0.030 [0.051] N=1,060	0.011 [0.058] N=869
Legumes consumed at least weekly	0.791 P=1,182	**-0.069* [0.040] N=1,488**	**-0.082** [0.041] N=1,378**	**-0.096** [0.040] N=1,241**	**-0.092** [0.045] N=1,060**	**-0.147*** [0.050] N=869**
Animal products consumed at least weekly	0.912 P=1,360	0.031 [0.031] N=1,488	0.033 [0.032] N=1,378	0.022 [0.034] N=1,241	0.014 [0.037] N=1,060	-0.000 [0.040] N=869
Sweets consumed at least weekly	0.687 P=1,018	-0.000 [0.041] N=1,488	0.014 [0.043] N=1,378	0.013 [0.045] N=1,241	0.010 [0.049] N=1,060	0.028 [0.058] N=869
Snack foods consumed at least weekly	0.561 P=826	-0.045 [0.045] N=1,488	-0.033 [0.048] N=1,378	-0.018 [0.051] N=1,241	-0.022 [0.058] N=1,060	-0.021 [0.065] N=869
Packaged soft drinks consumed at least weekly	0.348 P=508	0.032 [0.050] N=1,488	0.042 [0.054] N=1,378	0.031 [0.057] N=1,241	0.043 [0.060] N=1,060	0.043 [0.069] N=869
Grains consumed at least weekly	0.914 P=1,359	0.006 [0.029] N=1,488	0.015 [0.030] N=1,378	-0.009 [0.031] N=1,241	0.002 [0.033] N=1,060	-0.009 [0.039] N=869
Child was ever breastfed	0.965 P=1,432	0.006 [0.017] N=1,489	0.011 [0.018] N=1,379	0.018 [0.020] N=1,241	0.029 [0.023] N=1,060	0.017 [0.024] N=869

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.015 [0.037] N=1,265	0.037 [0.037] N=1,155	0.025 [0.040] N=1,018	0.027 [0.047] N=837	0.027 [0.063] N=646
Vegetables consumed at least weekly	0.039 [0.043] N=1,265	0.041 [0.047] N=1,155	0.007 [0.051] N=1,018	0.040 [0.057] N=837	-0.042 [0.071] N=646
Meat fish or poultry consumed at least weekly	0.094 [0.057] N=1,265	**0.105* [0.060] N=1,155**	0.069 [0.067] N=1,018	0.020 [0.077] N=837	-0.017 [0.090] N=646
Legumes consumed at least weekly	0.013 [0.061] N=1,265	-0.002 [0.064] N=1,155	-0.009 [0.065] N=1,018	0.025 [0.082] N=837	-0.061 [0.111] N=646
Animal products consumed at least weekly	0.045 [0.043] N=1,265	0.051 [0.046] N=1,155	0.030 [0.054] N=1,018	0.035 [0.062] N=837	-0.004 [0.078] N=646
Sweets consumed at least weekly	-0.012 [0.055] N=1,265	0.006 [0.060] N=1,155	-0.010 [0.067] N=1,018	-0.026 [0.076] N=837	0.024 [0.105] N=646
Snack foods consumed at least weekly	-0.060 [0.054] N=1,265	-0.045 [0.058] N=1,155	-0.022 [0.068] N=1,018	-0.042 [0.086] N=837	-0.080 [0.099] N=646
Packaged soft drinks consumed at least weekly	0.009 [0.060] N=1,265	0.020 [0.065] N=1,155	-0.004 [0.068] N=1,018	0.014 [0.079] N=837	-0.011 [0.101] N=646
Grains consumed at least weekly	0.046 [0.047] N=1,265	0.069 [0.053] N=1,155	0.030 [0.056] N=1,018	0.078 [0.067] N=837	0.094 [0.099] N=646
Child was ever breastfed	0.010 [0.027] N=1,266	0.020 [0.031] N=1,156	0.030 [0.036] N=1,018	0.049 [0.046] N=837	0.038 [0.062] N=646

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.940 P=1,397	-0.011 [0.024] N=1,488	0.001 [0.024] N=1,378	-0.007 [0.025] N=1,241	-0.011 [0.028] N=1,060	-0.022 [0.033] N=869
Vegetables consumed at least weekly	0.901 P=1,340	-0.000 [0.028] N=1,488	-0.001 [0.029] N=1,378	-0.023 [0.030] N=1,241	-0.014 [0.030] N=1,060	-0.052 [0.033] N=869
Meat fish or poultry consumed at least weekly	0.795 P=1,190	0.066 [0.043] N=1,488	**0.075* [0.044] N=1,378**	0.052 [0.048] N=1,241	0.036 [0.053] N=1,060	0.025 [0.060] N=869
Legumes consumed at least weekly	0.791 P=1,182	**-0.072* [0.040] N=1,488**	**-0.084** [0.040] N=1,378**	**-0.099** [0.040] N=1,241**	**-0.098** [0.045] N=1,060**	**-0.158*** [0.050] N=869**
Animal products consumed at least weekly	0.912 P=1,360	0.030 [0.030] N=1,488	0.032 [0.032] N=1,378	0.021 [0.034] N=1,241	0.015 [0.036] N=1,060	0.001 [0.040] N=869
Sweets consumed at least weekly	0.687 P=1,018	-0.001 [0.041] N=1,488	0.012 [0.043] N=1,378	0.014 [0.045] N=1,241	0.010 [0.048] N=1,060	0.029 [0.056] N=869
Snack foods consumed at least weekly	0.561 P=826	-0.047 [0.045] N=1,488	-0.036 [0.047] N=1,378	-0.020 [0.051] N=1,241	-0.025 [0.057] N=1,060	-0.029 [0.063] N=869
Packaged soft drinks consumed at least weekly	0.348 P=508	0.030 [0.050] N=1,488	0.039 [0.054] N=1,378	0.030 [0.057] N=1,241	0.040 [0.061] N=1,060	0.035 [0.071] N=869
Grains consumed at least weekly	0.914 P=1,359	0.005 [0.029] N=1,488	0.014 [0.030] N=1,378	-0.010 [0.031] N=1,241	-0.000 [0.032] N=1,060	-0.015 [0.038] N=869
Child was ever breastfed	0.965 P=1,432	0.005 [0.017] N=1,489	0.010 [0.018] N=1,379	0.016 [0.020] N=1,241	0.026 [0.022] N=1,060	0.013 [0.022] N=869

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.013 [0.036] N=1,265	0.036 [0.036] N=1,155	0.027 [0.039] N=1,018	0.027 [0.046] N=837	0.026 [0.062] N=646
Vegetables consumed at least weekly	0.036 [0.043] N=1,265	0.040 [0.046] N=1,155	0.008 [0.050] N=1,018	0.040 [0.057] N=837	-0.040 [0.070] N=646
Meat fish or poultry consumed at least weekly	0.085 [0.056] N=1,265	0.098 [0.059] N=1,155	0.066 [0.066] N=1,018	0.021 [0.077] N=837	-0.008 [0.090] N=646
Legumes consumed at least weekly	0.010 [0.061] N=1,265	-0.004 [0.065] N=1,155	-0.006 [0.067] N=1,018	0.029 [0.084] N=837	-0.065 [0.113] N=646
Animal products consumed at least weekly	0.042 [0.042] N=1,265	0.048 [0.046] N=1,155	0.030 [0.053] N=1,018	0.031 [0.062] N=837	-0.005 [0.079] N=646
Sweets consumed at least weekly	-0.023 [0.055] N=1,265	-0.008 [0.060] N=1,155	-0.018 [0.067] N=1,018	-0.030 [0.077] N=837	0.027 [0.105] N=646
Snack foods consumed at least weekly	-0.068 [0.053] N=1,265	-0.055 [0.057] N=1,155	-0.027 [0.068] N=1,018	-0.043 [0.086] N=837	-0.080 [0.099] N=646
Packaged soft drinks consumed at least weekly	0.002 [0.060] N=1,265	0.012 [0.065] N=1,155	-0.006 [0.069] N=1,018	0.012 [0.080] N=837	-0.016 [0.100] N=646
Grains consumed at least weekly	0.044 [0.048] N=1,265	0.069 [0.053] N=1,155	0.035 [0.058] N=1,018	0.081 [0.070] N=837	0.091 [0.100] N=646
Child was ever breastfed	0.012 [0.028] N=1,266	0.021 [0.032] N=1,156	0.031 [0.038] N=1,018	0.053 [0.049] N=837	0.038 [0.066] N=646

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,345	0.001 [0.028] N=1,427	0.003 [0.030] N=1,293	-0.001 [0.033] N=1,132	0.006 [0.034] N=950	0.023 [0.041] N=767
Vegetables consumed at least weekly	0.902 P=1,284	0.016 [0.035] N=1,427	0.030 [0.037] N=1,293	0.035 [0.039] N=1,132	0.041 [0.043] N=950	0.051 [0.049] N=767
Meat fish or poultry consumed at least weekly	0.783 P=1,128	**0.084** [0.041] N=1,427**	**0.091** [0.042] N=1,293**	**0.083* [0.046] N=1,132**	**0.126*** [0.047] N=950**	**0.136** [0.055] N=767**
Legumes consumed at least weekly	0.795 P=1,141	-0.032 [0.047] N=1,427	-0.006 [0.049] N=1,293	-0.018 [0.053] N=1,132	-0.041 [0.058] N=950	-0.005 [0.071] N=767
Animal products consumed at least weekly	0.912 P=1,303	0.001 [0.035] N=1,427	0.010 [0.037] N=1,293	0.021 [0.039] N=1,132	0.005 [0.041] N=950	0.041 [0.047] N=767
Sweets consumed at least weekly	0.687 P=983	0.010 [0.051] N=1,427	0.013 [0.054] N=1,293	0.005 [0.058] N=1,132	0.000 [0.061] N=950	-0.052 [0.069] N=767
Snack foods consumed at least weekly	0.557 P=789	-0.028 [0.051] N=1,427	-0.021 [0.055] N=1,293	-0.019 [0.058] N=1,132	-0.013 [0.063] N=950	-0.055 [0.073] N=767
Packaged soft drinks consumed at least weekly	0.326 P=459	-0.020 [0.047] N=1,427	-0.017 [0.048] N=1,293	0.021 [0.050] N=1,132	0.029 [0.056] N=950	0.029 [0.063] N=767
Grains consumed at least weekly	0.914 P=1,303	0.032 [0.042] N=1,427	0.041 [0.044] N=1,293	0.050 [0.047] N=1,132	0.056 [0.052] N=950	0.072 [0.059] N=767
Child was ever breastfed	0.964 P=1,375	-0.001 [0.023] N=1,428	0.002 [0.023] N=1,294	0.006 [0.024] N=1,133	0.001 [0.024] N=951	-0.018 [0.028] N=768

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.036] N=1,236	-0.049 [0.038] N=1,102	-0.067 [0.047] N=941	-0.048 [0.049] N=759	-0.032 [0.076] N=576
Vegetables consumed at least weekly	-0.042 [0.041] N=1,236	-0.030 [0.045] N=1,102	-0.033 [0.050] N=941	-0.020 [0.054] N=759	-0.011 [0.073] N=576
Meat fish or poultry consumed at least weekly	0.015 [0.059] N=1,236	0.019 [0.063] N=1,102	-0.023 [0.072] N=941	0.051 [0.075] N=759	0.036 [0.101] N=576
Legumes consumed at least weekly	-0.041 [0.054] N=1,236	-0.002 [0.059] N=1,102	-0.023 [0.066] N=941	-0.049 [0.070] N=759	0.040 [0.090] N=576
Animal products consumed at least weekly	-0.065 [0.043] N=1,236	-0.053 [0.048] N=1,102	-0.044 [0.051] N=941	**-0.089* [0.052] N=759**	-0.046 [0.065] N=576
Sweets consumed at least weekly	0.015 [0.069] N=1,236	0.022 [0.074] N=1,102	0.007 [0.083] N=941	0.005 [0.099] N=759	-0.142 [0.128] N=576
Snack foods consumed at least weekly	-0.028 [0.075] N=1,236	-0.016 [0.082] N=1,102	-0.010 [0.090] N=941	0.020 [0.109] N=759	-0.085 [0.145] N=576
Packaged soft drinks consumed at least weekly	-0.064 [0.075] N=1,236	-0.062 [0.081] N=1,102	-0.005 [0.091] N=941	0.006 [0.113] N=759	-0.024 [0.143] N=576
Grains consumed at least weekly	-0.027 [0.043] N=1,236	-0.027 [0.046] N=1,102	-0.026 [0.051] N=941	-0.040 [0.062] N=759	-0.080 [0.079] N=576
Child was ever breastfed	0.030 [0.045] N=1,237	0.040 [0.046] N=1,103	0.058 [0.050] N=942	0.055 [0.057] N=760	0.028 [0.075] N=577

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,345	0.001 [0.028] N=1,427	0.003 [0.030] N=1,293	-0.001 [0.033] N=1,132	0.006 [0.034] N=950	0.023 [0.041] N=767
Vegetables consumed at least weekly	0.902 P=1,284	0.016 [0.035] N=1,427	0.030 [0.036] N=1,293	0.036 [0.039] N=1,132	0.040 [0.042] N=950	0.050 [0.049] N=767
Meat fish or poultry consumed at least weekly	0.783 P=1,128	**0.084** [0.041] N=1,427**	**0.091** [0.042] N=1,293**	**0.083* [0.046] N=1,132**	**0.126*** [0.047] N=950**	**0.135** [0.055] N=767**
Legumes consumed at least weekly	0.795 P=1,141	-0.032 [0.047] N=1,427	-0.006 [0.049] N=1,293	-0.018 [0.053] N=1,132	-0.041 [0.058] N=950	-0.005 [0.071] N=767
Animal products consumed at least weekly	0.912 P=1,303	0.003 [0.035] N=1,427	0.011 [0.037] N=1,293	0.022 [0.039] N=1,132	0.004 [0.041] N=950	0.039 [0.047] N=767
Sweets consumed at least weekly	0.687 P=983	0.011 [0.051] N=1,427	0.013 [0.054] N=1,293	0.005 [0.058] N=1,132	-0.000 [0.061] N=950	-0.053 [0.069] N=767
Snack foods consumed at least weekly	0.557 P=789	-0.026 [0.051] N=1,427	-0.020 [0.055] N=1,293	-0.018 [0.058] N=1,132	-0.014 [0.063] N=950	-0.056 [0.073] N=767
Packaged soft drinks consumed at least weekly	0.326 P=459	-0.019 [0.046] N=1,427	-0.016 [0.048] N=1,293	0.021 [0.050] N=1,132	0.028 [0.056] N=950	0.026 [0.062] N=767
Grains consumed at least weekly	0.914 P=1,303	0.032 [0.042] N=1,427	0.041 [0.044] N=1,293	0.050 [0.047] N=1,132	0.056 [0.053] N=950	0.072 [0.059] N=767
Child was ever breastfed	0.964 P=1,375	-0.001 [0.023] N=1,428	0.002 [0.023] N=1,294	0.006 [0.024] N=1,133	0.001 [0.024] N=951	-0.018 [0.028] N=768

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.036] N=1,236	-0.048 [0.039] N=1,102	-0.067 [0.048] N=941	-0.048 [0.049] N=759	-0.031 [0.076] N=576
Vegetables consumed at least weekly	-0.041 [0.041] N=1,236	-0.029 [0.045] N=1,102	-0.032 [0.050] N=941	-0.019 [0.053] N=759	-0.010 [0.072] N=576
Meat fish or poultry consumed at least weekly	0.018 [0.059] N=1,236	0.023 [0.063] N=1,102	-0.020 [0.072] N=941	0.054 [0.075] N=759	0.041 [0.101] N=576
Legumes consumed at least weekly	-0.040 [0.054] N=1,236	-0.002 [0.059] N=1,102	-0.024 [0.066] N=941	-0.049 [0.070] N=759	0.037 [0.090] N=576
Animal products consumed at least weekly	-0.061 [0.041] N=1,236	-0.048 [0.046] N=1,102	-0.040 [0.050] N=941	**-0.086* [0.052] N=759**	-0.041 [0.063] N=576
Sweets consumed at least weekly	0.017 [0.069] N=1,236	0.024 [0.074] N=1,102	0.008 [0.083] N=941	0.005 [0.099] N=759	-0.142 [0.128] N=576
Snack foods consumed at least weekly	-0.022 [0.076] N=1,236	-0.011 [0.082] N=1,102	-0.006 [0.090] N=941	0.022 [0.110] N=759	-0.079 [0.147] N=576
Packaged soft drinks consumed at least weekly	-0.062 [0.074] N=1,236	-0.062 [0.080] N=1,102	-0.004 [0.091] N=941	0.007 [0.113] N=759	-0.018 [0.141] N=576
Grains consumed at least weekly	-0.028 [0.043] N=1,236	-0.028 [0.046] N=1,102	-0.028 [0.051] N=941	-0.042 [0.062] N=759	-0.084 [0.080] N=576
Child was ever breastfed	0.032 [0.045] N=1,237	0.041 [0.046] N=1,103	0.059 [0.050] N=942	0.056 [0.057] N=760	0.030 [0.075] N=577

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,345	-0.001 [0.028] N=1,427	0.002 [0.029] N=1,293	-0.000 [0.033] N=1,132	0.007 [0.034] N=950	0.024 [0.041] N=767
Vegetables consumed at least weekly	0.902 P=1,284	0.015 [0.034] N=1,427	0.029 [0.036] N=1,293	0.036 [0.039] N=1,132	0.041 [0.042] N=950	0.051 [0.049] N=767
Meat fish or poultry consumed at least weekly	0.783 P=1,128	**0.076* [0.040] N=1,427**	**0.087** [0.041] N=1,293**	**0.085* [0.044] N=1,132**	**0.127*** [0.045] N=950**	**0.136** [0.052] N=767**
Legumes consumed at least weekly	0.795 P=1,141	-0.034 [0.047] N=1,427	-0.008 [0.050] N=1,293	-0.017 [0.053] N=1,132	-0.040 [0.058] N=950	-0.003 [0.071] N=767
Animal products consumed at least weekly	0.912 P=1,303	0.001 [0.034] N=1,427	0.010 [0.037] N=1,293	0.023 [0.039] N=1,132	0.004 [0.041] N=950	0.040 [0.047] N=767
Sweets consumed at least weekly	0.687 P=983	0.011 [0.051] N=1,427	0.013 [0.054] N=1,293	0.006 [0.057] N=1,132	-0.001 [0.061] N=950	-0.053 [0.069] N=767
Snack foods consumed at least weekly	0.557 P=789	-0.023 [0.052] N=1,427	-0.018 [0.056] N=1,293	-0.018 [0.058] N=1,132	-0.015 [0.063] N=950	-0.057 [0.074] N=767
Packaged soft drinks consumed at least weekly	0.326 P=459	-0.014 [0.048] N=1,427	-0.014 [0.050] N=1,293	0.020 [0.052] N=1,132	0.025 [0.058] N=950	0.026 [0.064] N=767
Grains consumed at least weekly	0.914 P=1,303	0.029 [0.041] N=1,427	0.039 [0.044] N=1,293	0.050 [0.047] N=1,132	0.056 [0.052] N=950	0.072 [0.059] N=767
Child was ever breastfed	0.964 P=1,375	-0.002 [0.023] N=1,428	0.001 [0.023] N=1,294	0.006 [0.024] N=1,133	0.001 [0.024] N=951	-0.019 [0.028] N=768

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.036] N=1,236	-0.047 [0.038] N=1,102	-0.062 [0.046] N=941	-0.047 [0.050] N=759	-0.027 [0.077] N=576
Vegetables consumed at least weekly	-0.041 [0.041] N=1,236	-0.028 [0.044] N=1,102	-0.028 [0.049] N=941	-0.017 [0.053] N=759	-0.005 [0.073] N=576
Meat fish or poultry consumed at least weekly	0.013 [0.057] N=1,236	0.023 [0.060] N=1,102	-0.009 [0.068] N=941	0.063 [0.074] N=759	0.053 [0.096] N=576
Legumes consumed at least weekly	-0.038 [0.054] N=1,236	0.001 [0.059] N=1,102	-0.021 [0.067] N=941	-0.045 [0.070] N=759	0.038 [0.093] N=576
Animal products consumed at least weekly	-0.062 [0.041] N=1,236	-0.049 [0.046] N=1,102	-0.039 [0.050] N=941	**-0.090* [0.053] N=759**	-0.043 [0.062] N=576
Sweets consumed at least weekly	0.016 [0.069] N=1,236	0.022 [0.074] N=1,102	0.006 [0.082] N=941	-0.002 [0.100] N=759	-0.149 [0.128] N=576
Snack foods consumed at least weekly	-0.015 [0.076] N=1,236	-0.007 [0.084] N=1,102	-0.009 [0.091] N=941	0.013 [0.110] N=759	-0.083 [0.149] N=576
Packaged soft drinks consumed at least weekly	-0.064 [0.075] N=1,236	-0.067 [0.082] N=1,102	-0.021 [0.094] N=941	-0.020 [0.115] N=759	-0.035 [0.143] N=576
Grains consumed at least weekly	-0.032 [0.042] N=1,236	-0.031 [0.045] N=1,102	-0.027 [0.050] N=941	-0.047 [0.061] N=759	-0.089 [0.079] N=576
Child was ever breastfed	0.030 [0.045] N=1,237	0.040 [0.046] N=1,103	0.060 [0.050] N=942	0.059 [0.057] N=760	0.033 [0.075] N=577

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,345	-0.001 [0.028] N=1,427	0.001 [0.029] N=1,293	-0.002 [0.033] N=1,132	0.000 [0.033] N=950	0.015 [0.040] N=767
Vegetables consumed at least weekly	0.902 P=1,284	0.013 [0.036] N=1,427	0.028 [0.037] N=1,293	0.033 [0.040] N=1,132	0.035 [0.043] N=950	0.046 [0.049] N=767
Meat fish or poultry consumed at least weekly	0.783 P=1,128	**0.077* [0.041] N=1,427**	**0.086** [0.042] N=1,293**	**0.080* [0.045] N=1,132**	**0.115** [0.047] N=950**	**0.116** [0.054] N=767**
Legumes consumed at least weekly	0.795 P=1,141	-0.029 [0.047] N=1,427	-0.003 [0.050] N=1,293	-0.015 [0.053] N=1,132	-0.041 [0.057] N=950	-0.004 [0.071] N=767
Animal products consumed at least weekly	0.912 P=1,303	0.007 [0.034] N=1,427	0.017 [0.036] N=1,293	0.031 [0.039] N=1,132	0.008 [0.041] N=950	0.042 [0.047] N=767
Sweets consumed at least weekly	0.687 P=983	0.019 [0.052] N=1,427	0.019 [0.055] N=1,293	0.009 [0.059] N=1,132	0.001 [0.063] N=950	-0.052 [0.072] N=767
Snack foods consumed at least weekly	0.557 P=789	-0.012 [0.052] N=1,427	-0.009 [0.055] N=1,293	-0.009 [0.058] N=1,132	-0.008 [0.063] N=950	-0.047 [0.074] N=767
Packaged soft drinks consumed at least weekly	0.326 P=459	-0.003 [0.048] N=1,427	-0.005 [0.050] N=1,293	0.029 [0.052] N=1,132	0.036 [0.058] N=950	0.039 [0.065] N=767
Grains consumed at least weekly	0.914 P=1,303	0.029 [0.041] N=1,427	0.037 [0.043] N=1,293	0.047 [0.047] N=1,132	0.052 [0.052] N=950	0.068 [0.059] N=767
Child was ever breastfed	0.964 P=1,375	-0.001 [0.023] N=1,428	0.002 [0.023] N=1,294	0.006 [0.024] N=1,133	-0.001 [0.025] N=951	-0.021 [0.029] N=768

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.045 [0.035] N=1,236	-0.051 [0.037] N=1,102	-0.066 [0.046] N=941	-0.057 [0.050] N=759	-0.050 [0.076] N=576
Vegetables consumed at least weekly	-0.045 [0.041] N=1,236	-0.031 [0.045] N=1,102	-0.033 [0.050] N=941	-0.023 [0.053] N=759	-0.014 [0.071] N=576
Meat fish or poultry consumed at least weekly	0.014 [0.056] N=1,236	0.020 [0.059] N=1,102	-0.018 [0.066] N=941	0.041 [0.074] N=759	0.001 [0.093] N=576
Legumes consumed at least weekly	-0.035 [0.054] N=1,236	0.003 [0.059] N=1,102	-0.015 [0.067] N=941	-0.045 [0.070] N=759	0.046 [0.088] N=576
Animal products consumed at least weekly	-0.056 [0.039] N=1,236	-0.042 [0.044] N=1,102	-0.030 [0.047] N=941	**-0.084* [0.050] N=759**	-0.044 [0.058] N=576
Sweets consumed at least weekly	0.027 [0.069] N=1,236	0.032 [0.074] N=1,102	0.013 [0.083] N=941	0.003 [0.100] N=759	-0.146 [0.125] N=576
Snack foods consumed at least weekly	-0.007 [0.074] N=1,236	-0.001 [0.083] N=1,102	0.000 [0.090] N=941	0.022 [0.107] N=759	-0.071 [0.141] N=576
Packaged soft drinks consumed at least weekly	-0.046 [0.075] N=1,236	-0.052 [0.081] N=1,102	-0.001 [0.093] N=941	0.010 [0.114] N=759	0.012 [0.144] N=576
Grains consumed at least weekly	-0.032 [0.041] N=1,236	-0.035 [0.044] N=1,102	-0.033 [0.049] N=941	-0.055 [0.061] N=759	-0.103 [0.077] N=576
Child was ever breastfed	0.031 [0.044] N=1,237	0.043 [0.046] N=1,103	0.060 [0.050] N=942	0.056 [0.057] N=760	0.029 [0.077] N=577

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,249	0.005 [0.029] N=1,324	0.007 [0.031] N=1,198	0.011 [0.035] N=1,054	0.008 [0.035] N=885	0.023 [0.042] N=717
Vegetables consumed at least weekly	0.904 P=1,195	0.021 [0.035] N=1,324	0.035 [0.037] N=1,198	0.049 [0.039] N=1,054	0.045 [0.043] N=885	0.038 [0.050] N=717
Meat fish or poultry consumed at least weekly	0.782 P=1,045	**0.081* [0.044] N=1,324**	**0.086* [0.045] N=1,198**	**0.085* [0.049] N=1,054**	**0.102** [0.051] N=885**	**0.110* [0.059] N=717**
Legumes consumed at least weekly	0.793 P=1,057	-0.028 [0.049] N=1,324	-0.000 [0.052] N=1,198	-0.009 [0.056] N=1,054	-0.044 [0.061] N=885	-0.009 [0.076] N=717
Animal products consumed at least weekly	0.914 P=1,212	0.007 [0.036] N=1,324	0.020 [0.039] N=1,198	0.037 [0.041] N=1,054	0.013 [0.042] N=885	0.047 [0.047] N=717
Sweets consumed at least weekly	0.698 P=925	0.015 [0.053] N=1,324	0.018 [0.056] N=1,198	0.004 [0.061] N=1,054	-0.010 [0.066] N=885	-0.059 [0.076] N=717
Snack foods consumed at least weekly	0.565 P=742	-0.015 [0.052] N=1,324	-0.013 [0.056] N=1,198	-0.022 [0.059] N=1,054	-0.020 [0.065] N=885	-0.046 [0.076] N=717
Packaged soft drinks consumed at least weekly	0.338 P=442	-0.024 [0.047] N=1,324	-0.028 [0.049] N=1,198	0.007 [0.051] N=1,054	0.018 [0.058] N=885	0.015 [0.064] N=717
Grains consumed at least weekly	0.914 P=1,210	0.044 [0.043] N=1,324	0.053 [0.045] N=1,198	0.063 [0.049] N=1,054	0.065 [0.055] N=885	0.077 [0.060] N=717
Child was ever breastfed	0.966 P=1,280	-0.007 [0.021] N=1,325	-0.007 [0.021] N=1,199	-0.003 [0.022] N=1,055	-0.013 [0.022] N=886	-0.030 [0.027] N=718

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.031 [0.038] N=1,143	-0.034 [0.041] N=1,017	-0.033 [0.052] N=873	-0.035 [0.053] N=704	-0.020 [0.078] N=536
Vegetables consumed at least weekly	-0.027 [0.041] N=1,143	-0.013 [0.045] N=1,017	0.003 [0.051] N=873	0.003 [0.056] N=704	-0.023 [0.070] N=536
Meat fish or poultry consumed at least weekly	0.037 [0.067] N=1,143	0.044 [0.071] N=1,017	0.021 [0.080] N=873	0.045 [0.085] N=704	0.025 [0.110] N=536
Legumes consumed at least weekly	-0.018 [0.056] N=1,143	0.027 [0.060] N=1,017	0.017 [0.069] N=873	-0.023 [0.074] N=704	0.069 [0.095] N=536
Animal products consumed at least weekly	-0.057 [0.046] N=1,143	-0.034 [0.051] N=1,017	-0.013 [0.053] N=873	-0.069 [0.056] N=704	-0.023 [0.066] N=536
Sweets consumed at least weekly	0.026 [0.072] N=1,143	0.033 [0.078] N=1,017	0.001 [0.087] N=873	-0.022 [0.102] N=704	-0.181 [0.133] N=536
Snack foods consumed at least weekly	-0.002 [0.077] N=1,143	0.005 [0.085] N=1,017	-0.014 [0.093] N=873	0.006 [0.112] N=704	-0.057 [0.151] N=536
Packaged soft drinks consumed at least weekly	-0.076 [0.074] N=1,143	-0.085 [0.079] N=1,017	-0.034 [0.090] N=873	-0.026 [0.114] N=704	-0.074 [0.145] N=536
Grains consumed at least weekly	-0.003 [0.045] N=1,143	0.001 [0.049] N=1,017	0.008 [0.056] N=873	-0.015 [0.068] N=704	-0.068 [0.087] N=536
Child was ever breastfed	0.025 [0.041] N=1,144	0.030 [0.042] N=1,018	0.046 [0.044] N=874	0.033 [0.049] N=705	0.006 [0.070] N=537

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,249	0.006 [0.029] N=1,324	0.008 [0.031] N=1,198	0.012 [0.035] N=1,054	0.008 [0.035] N=885	0.022 [0.042] N=717
Vegetables consumed at least weekly	0.904 P=1,195	0.021 [0.035] N=1,324	0.034 [0.037] N=1,198	0.048 [0.039] N=1,054	0.042 [0.043] N=885	0.036 [0.050] N=717
Meat fish or poultry consumed at least weekly	0.782 P=1,045	**0.084* [0.043] N=1,324**	**0.091** [0.044] N=1,198**	**0.089* [0.048] N=1,054**	**0.106** [0.050] N=885**	**0.112* [0.057] N=717**
Legumes consumed at least weekly	0.793 P=1,057	-0.029 [0.049] N=1,324	-0.001 [0.052] N=1,198	-0.011 [0.056] N=1,054	-0.049 [0.061] N=885	-0.013 [0.076] N=717
Animal products consumed at least weekly	0.914 P=1,212	0.006 [0.036] N=1,324	0.020 [0.039] N=1,198	0.036 [0.041] N=1,054	0.011 [0.041] N=885	0.045 [0.047] N=717
Sweets consumed at least weekly	0.698 P=925	0.015 [0.053] N=1,324	0.018 [0.056] N=1,198	0.004 [0.061] N=1,054	-0.011 [0.066] N=885	-0.060 [0.077] N=717
Snack foods consumed at least weekly	0.565 P=742	-0.014 [0.051] N=1,324	-0.012 [0.055] N=1,198	-0.020 [0.058] N=1,054	-0.019 [0.064] N=885	-0.047 [0.076] N=717
Packaged soft drinks consumed at least weekly	0.338 P=442	-0.024 [0.047] N=1,324	-0.028 [0.049] N=1,198	0.008 [0.050] N=1,054	0.017 [0.058] N=885	0.012 [0.065] N=717
Grains consumed at least weekly	0.914 P=1,210	0.043 [0.043] N=1,324	0.052 [0.045] N=1,198	0.061 [0.049] N=1,054	0.063 [0.054] N=885	0.074 [0.060] N=717
Child was ever breastfed	0.966 P=1,280	-0.006 [0.022] N=1,325	-0.006 [0.022] N=1,199	0.000 [0.022] N=1,055	-0.007 [0.023] N=886	-0.025 [0.028] N=718

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.031 [0.039] N=1,143	-0.033 [0.042] N=1,017	-0.032 [0.052] N=873	-0.036 [0.054] N=704	-0.022 [0.082] N=536
Vegetables consumed at least weekly	-0.028 [0.041] N=1,143	-0.014 [0.046] N=1,017	0.002 [0.051] N=873	-0.002 [0.056] N=704	-0.026 [0.072] N=536
Meat fish or poultry consumed at least weekly	0.040 [0.066] N=1,143	0.049 [0.070] N=1,017	0.024 [0.080] N=873	0.049 [0.085] N=704	0.027 [0.111] N=536
Legumes consumed at least weekly	-0.019 [0.057] N=1,143	0.027 [0.061] N=1,017	0.016 [0.069] N=873	-0.033 [0.074] N=704	0.061 [0.094] N=536
Animal products consumed at least weekly	-0.057 [0.046] N=1,143	-0.034 [0.052] N=1,017	-0.014 [0.054] N=873	-0.073 [0.056] N=704	-0.031 [0.067] N=536
Sweets consumed at least weekly	0.028 [0.073] N=1,143	0.037 [0.078] N=1,017	0.008 [0.088] N=873	-0.011 [0.104] N=704	-0.164 [0.135] N=536
Snack foods consumed at least weekly	-0.001 [0.077] N=1,143	0.004 [0.085] N=1,017	-0.012 [0.093] N=873	0.013 [0.112] N=704	-0.057 [0.151] N=536
Packaged soft drinks consumed at least weekly	-0.075 [0.074] N=1,143	-0.084 [0.080] N=1,017	-0.028 [0.090] N=873	-0.018 [0.113] N=704	-0.063 [0.145] N=536
Grains consumed at least weekly	-0.004 [0.045] N=1,143	-0.000 [0.048] N=1,017	0.005 [0.055] N=873	-0.019 [0.068] N=704	-0.075 [0.086] N=536
Child was ever breastfed	0.026 [0.042] N=1,144	0.028 [0.043] N=1,018	0.048 [0.046] N=874	0.042 [0.052] N=705	0.012 [0.074] N=537

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.