Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Oldest Child

Age reference: 16 December 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.506	0.217 [0.244] N=1,404	0.186 [0.264] N=1,301	0.199 [0.294] N=1,174	0.304 [0.298] N=1,002	0.095 [0.320] N=824
WHO weight-for-age z score	-0.019	-0.049 [0.134] N=1,594	-0.017 [0.143] N=1,470	-0.002 [0.153] N=1,324	-0.003 [0.165] N=1,127	-0.159 [0.184] N=924
WHO weight-for-length/height z score, 0-60 months	0.354	0.097 [0.226] N=1,378	0.041 [0.241] N=1,281	-0.037 [0.272] N=1,156	-0.113 [0.304] N=986	-0.032 [0.342] N=809
WHO BMI-for-age z score	0.357	0.220 [0.252] N=1,344	0.245 [0.271] N=1,250	0.154 [0.304] N=1,126	-0.015 [0.344] N=960	0.072 [0.392] N=791
Stunted: HAZ below -2	0.259 P=352	-0.065 [0.042] N=1,404	-0.070 [0.045] N=1,301	-0.056 [0.048] N=1,174	-0.061 [0.051] N=1,002	-0.015 [0.055] N=824
Underweight: WAZ below -2	0.072 P=117	0.031 [0.022] N=1,594	0.027 [0.024] N=1,470	0.025 [0.026] N=1,324	0.032 [0.029] N=1,127	**0.060* [0.035] N=924**
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=159	-0.036 [0.036] N=1,378	-0.029 [0.036] N=1,281	-0.030 [0.037] N=1,156	-0.019 [0.042] N=986	-0.043 [0.044] N=809
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=260	0.011 [0.044] N=1,378	0.006 [0.047] N=1,281	-0.014 [0.052] N=1,156	-0.021 [0.059] N=986	-0.021 [0.066] N=809
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=119	0.016 [0.027] N=1,378	0.008 [0.028] N=1,281	0.010 [0.031] N=1,156	0.014 [0.032] N=986	0.026 [0.036] N=809
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=454	**-0.102** [0.046] N=1,335**	**-0.104** [0.047] N=1,242**	**-0.099** [0.049] N=1,121**	**-0.098* [0.052] N=956**	-0.072 [0.058] N=785
BMI-for-age z score below -2	0.120 P=166	-0.030 [0.039] N=1,344	-0.030 [0.039] N=1,250	-0.020 [0.042] N=1,126	0.013 [0.046] N=960	-0.003 [0.050] N=791

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.177 [0.348] N=1,187	0.074 [0.399] N=1,084	0.127 [0.485] N=957	0.425 [0.517] N=785	-0.141 [0.657] N=607
WHO weight-for-age z score	-0.154 [0.185] N=1,357	-0.137 [0.203] N=1,233	-0.135 [0.227] N=1,087	-0.161 [0.284] N=890	**-0.742** [0.356] N=687**
WHO weight-for-length/height z score, 0-60 months	0.129 [0.267] N=1,164	0.046 [0.295] N=1,067	-0.146 [0.368] N=942	-0.366 [0.429] N=772	-0.331 [0.504] N=595
WHO BMI-for-age z score	0.384 [0.285] N=1,137	0.471 [0.315] N=1,043	0.304 [0.389] N=919	-0.056 [0.456] N=753	0.088 [0.551] N=584
Stunted: HAZ below -2	-0.090 [0.061] N=1,187	-0.098 [0.068] N=1,084	-0.089 [0.077] N=957	-0.131 [0.086] N=785	-0.058 [0.114] N=607
Underweight: WAZ below -2	0.014 [0.031] N=1,357	0.008 [0.034] N=1,233	-0.000 [0.039] N=1,087	0.010 [0.049] N=890	0.094 [0.065] N=687
Wasted: WLZ/WHZ below -2, 0-60 months	-0.041 [0.051] N=1,164	-0.029 [0.053] N=1,067	-0.032 [0.060] N=942	-0.012 [0.076] N=772	-0.078 [0.091] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.052] N=1,164	0.007 [0.057] N=1,067	-0.039 [0.066] N=942	-0.059 [0.081] N=772	-0.103 [0.093] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.036] N=1,164	-0.006 [0.040] N=1,067	-0.004 [0.048] N=942	0.003 [0.055] N=772	0.032 [0.068] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.113 [0.072] N=1,129	-0.114 [0.077] N=1,036	-0.116 [0.087] N=915	-0.138 [0.103] N=750	-0.102 [0.144] N=579
BMI-for-age z score below -2	-0.060 [0.052] N=1,137	-0.065 [0.054] N=1,043	-0.055 [0.061] N=919	0.002 [0.077] N=753	-0.057 [0.095] N=584

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.506	0.217 [0.244] N=1,404	0.186 [0.264] N=1,301	0.199 [0.294] N=1,174	0.305 [0.297] N=1,002	0.098 [0.319] N=824
WHO weight-for-age z score	-0.019	-0.051 [0.133] N=1,594	-0.016 [0.142] N=1,470	0.002 [0.152] N=1,324	0.000 [0.164] N=1,127	-0.157 [0.181] N=924
WHO weight-for-length/height z score, 0-60 months	0.354	0.092 [0.226] N=1,378	0.036 [0.241] N=1,281	-0.038 [0.271] N=1,156	-0.116 [0.302] N=986	-0.040 [0.339] N=809
WHO BMI-for-age z score	0.357	0.217 [0.252] N=1,344	0.241 [0.270] N=1,250	0.154 [0.303] N=1,126	-0.018 [0.342] N=960	0.062 [0.388] N=791
Stunted: HAZ below -2	0.259 P=352	-0.065 [0.042] N=1,404	-0.070 [0.045] N=1,301	-0.057 [0.048] N=1,174	-0.061 [0.051] N=1,002	-0.015 [0.055] N=824
Underweight: WAZ below -2	0.072 P=117	0.031 [0.022] N=1,594	0.027 [0.024] N=1,470	0.024 [0.026] N=1,324	0.032 [0.030] N=1,127	**0.060* [0.035] N=924**
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=159	-0.035 [0.036] N=1,378	-0.027 [0.036] N=1,281	-0.030 [0.038] N=1,156	-0.019 [0.042] N=986	-0.041 [0.044] N=809
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=260	0.011 [0.044] N=1,378	0.005 [0.047] N=1,281	-0.014 [0.052] N=1,156	-0.021 [0.058] N=986	-0.022 [0.066] N=809
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=119	0.016 [0.027] N=1,378	0.008 [0.028] N=1,281	0.010 [0.031] N=1,156	0.014 [0.032] N=986	0.026 [0.036] N=809
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=454	**-0.100** [0.045] N=1,335**	**-0.102** [0.046] N=1,242**	**-0.099** [0.049] N=1,121**	**-0.096* [0.051] N=956**	-0.067 [0.057] N=785
BMI-for-age z score below -2	0.120 P=166	-0.030 [0.039] N=1,344	-0.029 [0.039] N=1,250	-0.020 [0.042] N=1,126	0.014 [0.046] N=960	0.000 [0.049] N=791

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.174 [0.348] N=1,187	0.067 [0.398] N=1,084	0.115 [0.484] N=957	0.400 [0.515] N=785	-0.170 [0.647] N=607
WHO weight-for-age z score	-0.153 [0.182] N=1,357	-0.127 [0.200] N=1,233	-0.117 [0.223] N=1,087	-0.145 [0.282] N=890	**-0.712** [0.353] N=687**
WHO weight-for-length/height z score, 0-60 months	0.123 [0.266] N=1,164	0.041 [0.294] N=1,067	-0.141 [0.365] N=942	-0.367 [0.426] N=772	-0.336 [0.501] N=595
WHO BMI-for-age z score	0.383 [0.285] N=1,137	0.470 [0.315] N=1,043	0.317 [0.387] N=919	-0.054 [0.455] N=753	0.075 [0.550] N=584
Stunted: HAZ below -2	-0.091 [0.061] N=1,187	-0.098 [0.068] N=1,084	-0.090 [0.077] N=957	-0.129 [0.086] N=785	-0.056 [0.114] N=607
Underweight: WAZ below -2	0.013 [0.030] N=1,357	0.006 [0.033] N=1,233	-0.004 [0.038] N=1,087	0.007 [0.049] N=890	0.089 [0.065] N=687
Wasted: WLZ/WHZ below -2, 0-60 months	-0.040 [0.052] N=1,164	-0.027 [0.054] N=1,067	-0.034 [0.061] N=942	-0.012 [0.077] N=772	-0.077 [0.092] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.015 [0.051] N=1,164	0.006 [0.057] N=1,067	-0.039 [0.066] N=942	-0.059 [0.080] N=772	-0.104 [0.093] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.008 [0.036] N=1,164	-0.006 [0.041] N=1,067	-0.004 [0.048] N=942	0.003 [0.055] N=772	0.033 [0.068] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.112 [0.072] N=1,129	-0.114 [0.077] N=1,036	-0.120 [0.087] N=915	-0.140 [0.104] N=750	-0.101 [0.146] N=579
BMI-for-age z score below -2	-0.060 [0.053] N=1,137	-0.065 [0.054] N=1,043	-0.058 [0.062] N=919	0.002 [0.078] N=753	-0.055 [0.097] N=584

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.506	0.191 [0.245] N=1,404	0.144 [0.266] N=1,301	0.171 [0.295] N=1,174	0.286 [0.296] N=1,002	0.057 [0.315] N=824
WHO weight-for-age z score	-0.019	-0.091 [0.127] N=1,594	-0.057 [0.136] N=1,470	-0.035 [0.145] N=1,324	-0.026 [0.156] N=1,127	-0.186 [0.169] N=924
WHO weight-for-length/height z score, 0-60 months	0.354	0.071 [0.228] N=1,378	0.022 [0.245] N=1,281	-0.045 [0.272] N=1,156	-0.116 [0.299] N=986	-0.050 [0.335] N=809
WHO BMI-for-age z score	0.357	0.207 [0.254] N=1,344	0.232 [0.273] N=1,250	0.148 [0.303] N=1,126	-0.017 [0.339] N=960	0.056 [0.385] N=791
Stunted: HAZ below -2	0.259 P=352	-0.057 [0.041] N=1,404	-0.056 [0.044] N=1,301	-0.048 [0.047] N=1,174	-0.056 [0.049] N=1,002	-0.005 [0.053] N=824
Underweight: WAZ below -2	0.072 P=117	**0.038* [0.022] N=1,594**	0.036 [0.023] N=1,470	0.032 [0.025] N=1,324	0.039 [0.028] N=1,127	**0.068** [0.033] N=924**
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=159	-0.029 [0.036] N=1,378	-0.019 [0.036] N=1,281	-0.023 [0.038] N=1,156	-0.015 [0.042] N=986	-0.038 [0.044] N=809
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=260	0.012 [0.044] N=1,378	0.008 [0.048] N=1,281	-0.011 [0.052] N=1,156	-0.018 [0.058] N=986	-0.020 [0.066] N=809
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=119	0.018 [0.027] N=1,378	0.013 [0.029] N=1,281	0.013 [0.031] N=1,156	0.016 [0.032] N=986	0.027 [0.037] N=809
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=454	**-0.083* [0.045] N=1,335**	**-0.078* [0.045] N=1,242**	**-0.081* [0.048] N=1,121**	**-0.085* [0.050] N=956**	-0.051 [0.056] N=785
BMI-for-age z score below -2	0.120 P=166	-0.024 [0.039] N=1,344	-0.020 [0.040] N=1,250	-0.012 [0.042] N=1,126	0.018 [0.046] N=960	0.003 [0.050] N=791

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.147 [0.351] N=1,187	0.006 [0.402] N=1,084	0.072 [0.485] N=957	0.393 [0.511] N=785	-0.169 [0.637] N=607
WHO weight-for-age z score	-0.187 [0.178] N=1,357	-0.166 [0.196] N=1,233	-0.147 [0.221] N=1,087	-0.148 [0.281] N=890	**-0.709** [0.350] N=687**
WHO weight-for-length/height z score, 0-60 months	0.119 [0.271] N=1,164	0.048 [0.303] N=1,067	-0.119 [0.371] N=942	-0.368 [0.427] N=772	-0.362 [0.502] N=595
WHO BMI-for-age z score	0.384 [0.289] N=1,137	0.478 [0.322] N=1,043	0.329 [0.391] N=919	-0.074 [0.454] N=753	0.052 [0.551] N=584
Stunted: HAZ below -2	-0.084 [0.061] N=1,187	-0.081 [0.068] N=1,084	-0.080 [0.077] N=957	-0.133 [0.085] N=785	-0.060 [0.109] N=607
Underweight: WAZ below -2	0.019 [0.029] N=1,357	0.016 [0.033] N=1,233	0.003 [0.037] N=1,087	0.009 [0.049] N=890	0.086 [0.066] N=687
Wasted: WLZ/WHZ below -2, 0-60 months	-0.036 [0.051] N=1,164	-0.020 [0.054] N=1,067	-0.030 [0.062] N=942	-0.007 [0.078] N=772	-0.070 [0.094] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.018 [0.052] N=1,164	0.013 [0.058] N=1,067	-0.033 [0.067] N=942	-0.057 [0.079] N=772	-0.101 [0.090] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.010 [0.037] N=1,164	0.001 [0.042] N=1,067	-0.000 [0.050] N=942	0.003 [0.056] N=772	0.029 [0.068] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.099 [0.071] N=1,129	-0.089 [0.076] N=1,036	-0.106 [0.086] N=915	-0.140 [0.103] N=750	-0.098 [0.141] N=579
BMI-for-age z score below -2	-0.055 [0.052] N=1,137	-0.054 [0.054] N=1,043	-0.051 [0.063] N=919	0.009 [0.079] N=753	-0.049 [0.098] N=584

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.506	0.148 [0.243] N=1,404	0.114 [0.264] N=1,301	0.140 [0.294] N=1,174	0.243 [0.297] N=1,002	0.014 [0.315] N=824
WHO weight-for-age z score	-0.019	-0.110 [0.127] N=1,594	-0.078 [0.136] N=1,470	-0.055 [0.146] N=1,324	-0.060 [0.159] N=1,127	-0.231 [0.174] N=924
WHO weight-for-length/height z score, 0-60 months	0.354	0.092 [0.224] N=1,378	0.033 [0.240] N=1,281	-0.045 [0.269] N=1,156	-0.118 [0.302] N=986	-0.044 [0.337] N=809
WHO BMI-for-age z score	0.357	0.233 [0.251] N=1,344	0.255 [0.270] N=1,250	0.167 [0.301] N=1,126	-0.000 [0.341] N=960	0.081 [0.386] N=791
Stunted: HAZ below -2	0.259 P=352	-0.047 [0.041] N=1,404	-0.047 [0.044] N=1,301	-0.036 [0.046] N=1,174	-0.041 [0.048] N=1,002	0.006 [0.052] N=824
Underweight: WAZ below -2	0.072 P=117	**0.041* [0.022] N=1,594**	**0.040* [0.023] N=1,470**	0.036 [0.025] N=1,324	0.044 [0.029] N=1,127	**0.074** [0.034] N=924**
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=159	-0.030 [0.036] N=1,378	-0.021 [0.036] N=1,281	-0.024 [0.038] N=1,156	-0.016 [0.043] N=986	-0.042 [0.044] N=809
Overweight: WLZ/WHZ above +2, 0-60 months	0.195 P=260	0.015 [0.044] N=1,378	0.010 [0.047] N=1,281	-0.011 [0.052] N=1,156	-0.018 [0.059] N=986	-0.022 [0.067] N=809
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=119	0.020 [0.027] N=1,378	0.013 [0.028] N=1,281	0.011 [0.031] N=1,156	0.014 [0.033] N=986	0.027 [0.037] N=809
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=454	-0.073 [0.046] N=1,335	-0.070 [0.046] N=1,242	-0.068 [0.049] N=1,121	-0.066 [0.052] N=956	-0.038 [0.058] N=785
BMI-for-age z score below -2	0.120 P=166	-0.025 [0.039] N=1,344	-0.023 [0.040] N=1,250	-0.013 [0.043] N=1,126	0.018 [0.047] N=960	-0.000 [0.050] N=791

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.111 [0.354] N=1,187	-0.009 [0.403] N=1,084	0.073 [0.491] N=957	0.384 [0.514] N=785	-0.216 [0.629] N=607
WHO weight-for-age z score	-0.219 [0.181] N=1,357	-0.196 [0.196] N=1,233	-0.178 [0.221] N=1,087	-0.187 [0.284] N=890	**-0.733** [0.353] N=687**
WHO weight-for-length/height z score, 0-60 months	0.114 [0.269] N=1,164	0.039 [0.296] N=1,067	-0.135 [0.367] N=942	-0.344 [0.428] N=772	-0.278 [0.504] N=595
WHO BMI-for-age z score	0.379 [0.290] N=1,137	0.478 [0.319] N=1,043	0.335 [0.390] N=919	-0.028 [0.458] N=753	0.144 [0.558] N=584
Stunted: HAZ below -2	-0.079 [0.062] N=1,187	-0.080 [0.069] N=1,084	-0.082 [0.077] N=957	-0.133 [0.086] N=785	-0.051 [0.113] N=607
Underweight: WAZ below -2	0.023 [0.030] N=1,357	0.018 [0.033] N=1,233	0.007 [0.038] N=1,087	0.013 [0.050] N=890	0.091 [0.068] N=687
Wasted: WLZ/WHZ below -2, 0-60 months	-0.032 [0.052] N=1,164	-0.022 [0.054] N=1,067	-0.031 [0.062] N=942	-0.011 [0.079] N=772	-0.075 [0.094] N=595
Overweight: WLZ/WHZ above +2, 0-60 months	0.020 [0.052] N=1,164	0.012 [0.057] N=1,067	-0.034 [0.065] N=942	-0.050 [0.078] N=772	-0.081 [0.089] N=595
Obese: WLZ/WHZ above +3, 0-60 months	0.014 [0.037] N=1,164	0.002 [0.041] N=1,067	0.001 [0.049] N=942	0.009 [0.055] N=772	0.040 [0.068] N=595
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.088 [0.073] N=1,129	-0.088 [0.077] N=1,036	-0.104 [0.087] N=915	-0.135 [0.104] N=750	-0.083 [0.144] N=579
BMI-for-age z score below -2	-0.054 [0.053] N=1,137	-0.062 [0.054] N=1,043	-0.059 [0.063] N=919	-0.001 [0.079] N=753	-0.058 [0.098] N=584

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.515	0.161 [0.250] N=1,309	0.127 [0.271] N=1,219	0.151 [0.298] N=1,098	0.242 [0.293] N=943	0.079 [0.300] N=774
WHO weight-for-age z score	-0.014	-0.000 [0.137] N=1,480	0.031 [0.143] N=1,371	0.052 [0.154] N=1,233	0.048 [0.164] N=1,054	-0.071 [0.190] N=864
WHO weight-for-length/height z score, 0-60 months	0.360	0.232 [0.220] N=1,281	0.183 [0.233] N=1,198	0.089 [0.264] N=1,080	0.017 [0.289] N=927	0.124 [0.320] N=759
WHO BMI-for-age z score	0.365	0.366 [0.245] N=1,249	0.367 [0.260] N=1,168	0.272 [0.291] N=1,052	0.136 [0.325] N=903	0.263 [0.361] N=742
Stunted: HAZ below -2	0.260 P=328	-0.053 [0.045] N=1,309	-0.059 [0.047] N=1,219	-0.047 [0.049] N=1,098	-0.054 [0.053] N=943	-0.011 [0.057] N=774
Underweight: WAZ below -2	0.072 P=109	0.024 [0.023] N=1,480	0.021 [0.025] N=1,371	0.017 [0.027] N=1,233	0.024 [0.030] N=1,054	0.049 [0.038] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=147	-0.046 [0.036] N=1,281	-0.038 [0.035] N=1,198	-0.037 [0.037] N=1,080	-0.028 [0.042] N=927	-0.047 [0.044] N=759
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=241	0.027 [0.045] N=1,281	0.023 [0.049] N=1,198	0.000 [0.054] N=1,080	-0.009 [0.060] N=927	-0.002 [0.069] N=759
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=114	0.018 [0.029] N=1,281	0.014 [0.031] N=1,198	0.017 [0.034] N=1,080	0.020 [0.034] N=927	0.039 [0.039] N=759
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=422	**-0.093* [0.049] N=1,243**	**-0.098** [0.049] N=1,163**	**-0.097* [0.052] N=1,048**	**-0.102* [0.057] N=900**	-0.071 [0.064] N=737
BMI-for-age z score below -2	0.118 P=152	-0.039 [0.038] N=1,249	-0.037 [0.038] N=1,168	-0.028 [0.041] N=1,052	0.001 [0.045] N=903	-0.012 [0.049] N=742

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.139 [0.363] N=1,105	0.046 [0.417] N=1,015	0.109 [0.504] N=894	0.399 [0.536] N=739	-0.064 [0.672] N=570
WHO weight-for-age z score	-0.151 [0.193] N=1,258	-0.140 [0.205] N=1,149	-0.141 [0.233] N=1,011	-0.197 [0.287] N=832	**-0.717* [0.400] N=642**
WHO weight-for-length/height z score, 0-60 months	0.175 [0.265] N=1,080	0.080 [0.288] N=997	-0.151 [0.357] N=879	-0.360 [0.429] N=726	-0.242 [0.507] N=558
WHO BMI-for-age z score	0.435 [0.290] N=1,055	0.456 [0.314] N=974	0.274 [0.382] N=858	-0.054 [0.463] N=709	0.192 [0.562] N=548
Stunted: HAZ below -2	-0.086 [0.064] N=1,105	-0.101 [0.070] N=1,015	-0.096 [0.078] N=894	-0.143 [0.088] N=739	-0.078 [0.116] N=570
Underweight: WAZ below -2	0.010 [0.031] N=1,258	0.005 [0.034] N=1,149	-0.003 [0.039] N=1,011	0.013 [0.046] N=832	0.093 [0.071] N=642
Wasted: WLZ/WHZ below -2, 0-60 months	-0.053 [0.052] N=1,080	-0.041 [0.052] N=997	-0.044 [0.060] N=879	-0.038 [0.077] N=726	-0.123 [0.091] N=558
Overweight: WLZ/WHZ above +2, 0-60 months	0.022 [0.053] N=1,080	0.007 [0.058] N=997	-0.045 [0.068] N=879	-0.081 [0.084] N=726	-0.104 [0.100] N=558
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.040] N=1,080	-0.010 [0.045] N=997	-0.010 [0.053] N=879	-0.007 [0.059] N=726	0.030 [0.072] N=558
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.113 [0.074] N=1,050	-0.125 [0.077] N=970	-0.141 [0.086] N=855	**-0.187* [0.102] N=707**	-0.167 [0.146] N=544
BMI-for-age z score below -2	-0.065 [0.052] N=1,055	-0.067 [0.052] N=974	-0.059 [0.060] N=858	-0.013 [0.077] N=709	-0.086 [0.095] N=548

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.515	0.148 [0.251] N=1,309	0.115 [0.271] N=1,219	0.128 [0.301] N=1,098	0.210 [0.302] N=943	0.012 [0.321] N=774
WHO weight-for-age z score	-0.014	-0.005 [0.135] N=1,480	0.027 [0.142] N=1,371	0.041 [0.153] N=1,233	0.030 [0.163] N=1,054	-0.107 [0.189] N=864
WHO weight-for-length/height z score, 0-60 months	0.360	0.220 [0.221] N=1,281	0.169 [0.234] N=1,198	0.078 [0.266] N=1,080	0.007 [0.296] N=927	0.117 [0.334] N=759
WHO BMI-for-age z score	0.365	0.354 [0.246] N=1,249	0.354 [0.261] N=1,168	0.266 [0.295] N=1,052	0.127 [0.334] N=903	0.251 [0.378] N=742
Stunted: HAZ below -2	0.260 P=328	-0.050 [0.045] N=1,309	-0.056 [0.047] N=1,219	-0.044 [0.049] N=1,098	-0.049 [0.053] N=943	-0.000 [0.057] N=774
Underweight: WAZ below -2	0.072 P=109	0.026 [0.023] N=1,480	0.022 [0.025] N=1,371	0.019 [0.027] N=1,233	0.029 [0.030] N=1,054	0.058 [0.038] N=864
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=147	-0.044 [0.036] N=1,281	-0.036 [0.036] N=1,198	-0.037 [0.038] N=1,080	-0.029 [0.044] N=927	-0.054 [0.047] N=759
Overweight: WLZ/WHZ above +2, 0-60 months	0.196 P=241	0.027 [0.045] N=1,281	0.022 [0.049] N=1,198	-0.000 [0.054] N=1,080	-0.011 [0.061] N=927	-0.003 [0.069] N=759
Obese: WLZ/WHZ above +3, 0-60 months	0.090 P=114	0.018 [0.029] N=1,281	0.015 [0.031] N=1,198	0.017 [0.033] N=1,080	0.020 [0.034] N=927	0.039 [0.039] N=759
CIAF: any stunting, underweight, or wasting, 0-60 months	0.344 P=422	**-0.089* [0.049] N=1,243**	**-0.095* [0.049] N=1,163**	**-0.095* [0.052] N=1,048**	**-0.100* [0.056] N=900**	-0.071 [0.062] N=737
BMI-for-age z score below -2	0.118 P=152	-0.036 [0.038] N=1,249	-0.035 [0.039] N=1,168	-0.028 [0.042] N=1,052	0.001 [0.047] N=903	-0.017 [0.052] N=742

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.160 [0.363] N=1,105	0.065 [0.417] N=1,015	0.132 [0.499] N=894	0.397 [0.541] N=739	-0.103 [0.698] N=570
WHO weight-for-age z score	-0.123 [0.191] N=1,258	-0.108 [0.203] N=1,149	-0.120 [0.231] N=1,011	-0.184 [0.286] N=832	**-0.741* [0.398] N=642**
WHO weight-for-length/height z score, 0-60 months	0.196 [0.260] N=1,080	0.111 [0.284] N=997	-0.117 [0.356] N=879	-0.348 [0.429] N=726	-0.268 [0.505] N=558
WHO BMI-for-age z score	0.448 [0.284] N=1,055	0.478 [0.307] N=974	0.297 [0.378] N=858	-0.044 [0.457] N=709	0.154 [0.554] N=548
Stunted: HAZ below -2	-0.085 [0.064] N=1,105	-0.097 [0.071] N=1,015	-0.097 [0.078] N=894	-0.141 [0.089] N=739	-0.075 [0.119] N=570
Underweight: WAZ below -2	0.007 [0.031] N=1,258	0.001 [0.034] N=1,149	-0.008 [0.039] N=1,011	0.009 [0.046] N=832	0.097 [0.069] N=642
Wasted: WLZ/WHZ below -2, 0-60 months	-0.053 [0.051] N=1,080	-0.041 [0.052] N=997	-0.046 [0.060] N=879	-0.038 [0.076] N=726	-0.124 [0.090] N=558
Overweight: WLZ/WHZ above +2, 0-60 months	0.023 [0.053] N=1,080	0.012 [0.058] N=997	-0.040 [0.069] N=879	-0.075 [0.084] N=726	-0.106 [0.099] N=558
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.040] N=1,080	-0.010 [0.045] N=997	-0.008 [0.053] N=879	-0.006 [0.059] N=726	0.030 [0.073] N=558
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.109 [0.074] N=1,050	-0.120 [0.077] N=970	-0.138 [0.086] N=855	**-0.181* [0.103] N=707**	-0.166 [0.145] N=544
BMI-for-age z score below -2	-0.062 [0.051] N=1,055	-0.066 [0.051] N=974	-0.059 [0.060] N=858	-0.012 [0.077] N=709	-0.086 [0.095] N=548

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.406	-0.212 [0.282] N=1,262	-0.281 [0.290] N=1,133	-0.296 [0.301] N=991	-0.527 [0.334] N=839	-0.599 [0.374] N=686
WHO weight-for-age z score	-0.015	0.034 [0.150] N=1,419	0.057 [0.157] N=1,286	-0.060 [0.156] N=1,125	-0.008 [0.165] N=945	-0.020 [0.184] N=764
WHO weight-for-length/height z score, 0-60 months	0.352	0.164 [0.205] N=1,245	0.076 [0.216] N=1,121	0.019 [0.229] N=983	0.236 [0.246] N=830	0.086 [0.268] N=679
WHO BMI-for-age z score	0.323	0.142 [0.232] N=1,209	0.068 [0.242] N=1,088	-0.031 [0.255] N=955	0.109 [0.279] N=805	0.072 [0.299] N=663
Stunted: HAZ below -2	0.243 P=310	0.030 [0.057] N=1,262	0.023 [0.058] N=1,133	0.024 [0.060] N=991	0.053 [0.065] N=839	0.072 [0.070] N=686
Underweight: WAZ below -2	0.070 P=101	-0.007 [0.031] N=1,419	-0.016 [0.033] N=1,286	0.003 [0.035] N=1,125	-0.004 [0.036] N=945	-0.005 [0.039] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=142	0.022 [0.036] N=1,245	0.026 [0.039] N=1,121	0.043 [0.041] N=983	-0.021 [0.041] N=830	-0.007 [0.047] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.090* [0.045] N=1,245**	0.075 [0.047] N=1,121	0.070 [0.050] N=983	0.088 [0.057] N=830	0.061 [0.058] N=679
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=115	0.052 [0.037] N=1,245	0.036 [0.038] N=1,121	0.028 [0.042] N=983	0.034 [0.049] N=830	-0.013 [0.048] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.334 P=406	0.051 [0.061] N=1,205	0.035 [0.061] N=1,081	0.057 [0.064] N=947	0.030 [0.066] N=801	0.075 [0.071] N=656
BMI-for-age z score below -2	0.122 P=148	0.026 [0.041] N=1,209	0.029 [0.045] N=1,088	0.033 [0.049] N=955	-0.017 [0.049] N=805	-0.013 [0.054] N=663

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.094 [0.393] N=1,088	-0.181 [0.420] N=959	-0.203 [0.473] N=817	-0.783 [0.571] N=665	-1.108 [0.698] N=512
WHO weight-for-age z score	0.056 [0.210] N=1,229	0.077 [0.231] N=1,096	-0.139 [0.230] N=935	-0.037 [0.251] N=755	0.005 [0.328] N=574
WHO weight-for-length/height z score, 0-60 months	0.079 [0.306] N=1,071	-0.093 [0.327] N=947	-0.207 [0.371] N=809	0.299 [0.423] N=656	-0.097 [0.508] N=505
WHO BMI-for-age z score	-0.166 [0.336] N=1,046	-0.317 [0.357] N=925	-0.525 [0.403] N=792	-0.273 [0.478] N=642	-0.437 [0.554] N=500
Stunted: HAZ below -2	0.043 [0.090] N=1,088	0.025 [0.094] N=959	0.032 [0.105] N=817	0.119 [0.129] N=665	0.208 [0.153] N=512
Underweight: WAZ below -2	-0.021 [0.045] N=1,229	-0.036 [0.049] N=1,096	-0.005 [0.055] N=935	-0.034 [0.056] N=755	-0.061 [0.062] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	**0.079* [0.047] N=1,071**	**0.091* [0.053] N=947**	**0.132** [0.058] N=809**	0.000 [0.058] N=656	0.034 [0.071] N=505
Overweight: WLZ/WHZ above +2, 0-60 months	**0.142** [0.059] N=1,071**	**0.119* [0.061] N=947**	**0.124* [0.067] N=809**	**0.190** [0.081] N=656**	0.128 [0.090] N=505
Obese: WLZ/WHZ above +3, 0-60 months	**0.114*** [0.040] N=1,071**	**0.086** [0.040] N=947**	**0.082* [0.047] N=809**	**0.111* [0.065] N=656**	-0.017 [0.065] N=505
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.094] N=1,037	0.062 [0.098] N=913	0.110 [0.111] N=779	0.070 [0.129] N=633	0.185 [0.150] N=488
BMI-for-age z score below -2	**0.119** [0.047] N=1,046**	**0.132** [0.056] N=925**	**0.152** [0.066] N=792**	0.052 [0.064] N=642	0.069 [0.078] N=500

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.406	-0.216 [0.282] N=1,262	-0.284 [0.290] N=1,133	-0.295 [0.301] N=991	-0.528 [0.333] N=839	-0.599 [0.374] N=686
WHO weight-for-age z score	-0.015	0.037 [0.149] N=1,419	0.060 [0.156] N=1,286	-0.057 [0.154] N=1,125	-0.011 [0.165] N=945	-0.027 [0.183] N=764
WHO weight-for-length/height z score, 0-60 months	0.352	0.171 [0.205] N=1,245	0.083 [0.216] N=1,121	0.025 [0.229] N=983	0.237 [0.246] N=830	0.087 [0.267] N=679
WHO BMI-for-age z score	0.323	0.146 [0.233] N=1,209	0.071 [0.243] N=1,088	-0.030 [0.255] N=955	0.109 [0.279] N=805	0.069 [0.301] N=663
Stunted: HAZ below -2	0.243 P=310	0.029 [0.057] N=1,262	0.021 [0.058] N=1,133	0.021 [0.059] N=991	0.053 [0.065] N=839	0.072 [0.070] N=686
Underweight: WAZ below -2	0.070 P=101	-0.008 [0.031] N=1,419	-0.017 [0.033] N=1,286	0.002 [0.035] N=1,125	-0.003 [0.036] N=945	-0.004 [0.039] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=142	0.020 [0.037] N=1,245	0.022 [0.039] N=1,121	0.039 [0.041] N=983	-0.021 [0.041] N=830	-0.007 [0.047] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.090** [0.045] N=1,245**	0.075 [0.047] N=1,121	0.069 [0.050] N=983	0.088 [0.057] N=830	0.061 [0.059] N=679
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=115	0.052 [0.037] N=1,245	0.035 [0.038] N=1,121	0.027 [0.042] N=983	0.033 [0.049] N=830	-0.013 [0.048] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.334 P=406	0.046 [0.061] N=1,205	0.027 [0.061] N=1,081	0.049 [0.063] N=947	0.029 [0.067] N=801	0.075 [0.071] N=656
BMI-for-age z score below -2	0.122 P=148	0.024 [0.041] N=1,209	0.027 [0.045] N=1,088	0.031 [0.048] N=955	-0.016 [0.049] N=805	-0.013 [0.054] N=663

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.102 [0.392] N=1,088	-0.188 [0.419] N=959	-0.200 [0.473] N=817	-0.793 [0.568] N=665	-1.129 [0.695] N=512
WHO weight-for-age z score	0.069 [0.206] N=1,229	0.089 [0.226] N=1,096	-0.125 [0.223] N=935	-0.024 [0.244] N=755	0.033 [0.315] N=574
WHO weight-for-length/height z score, 0-60 months	0.096 [0.308] N=1,071	-0.075 [0.329] N=947	-0.190 [0.372] N=809	0.319 [0.428] N=656	-0.022 [0.527] N=505
WHO BMI-for-age z score	-0.145 [0.339] N=1,046	-0.294 [0.359] N=925	-0.505 [0.404] N=792	-0.254 [0.483] N=642	-0.376 [0.573] N=500
Stunted: HAZ below -2	0.040 [0.090] N=1,088	0.020 [0.094] N=959	0.024 [0.106] N=817	0.116 [0.129] N=665	0.207 [0.152] N=512
Underweight: WAZ below -2	-0.024 [0.045] N=1,229	-0.038 [0.048] N=1,096	-0.008 [0.054] N=935	-0.036 [0.055] N=755	-0.065 [0.061] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.073 [0.047] N=1,071	0.081 [0.054] N=947	**0.121** [0.057] N=809**	-0.008 [0.058] N=656	0.018 [0.073] N=505
Overweight: WLZ/WHZ above +2, 0-60 months	**0.144** [0.059] N=1,071**	**0.120* [0.061] N=947**	**0.123* [0.067] N=809**	**0.188** [0.081] N=656**	0.130 [0.091] N=505
Obese: WLZ/WHZ above +3, 0-60 months	**0.113*** [0.040] N=1,071**	**0.084** [0.040] N=947**	**0.080* [0.047] N=809**	**0.108* [0.064] N=656**	-0.019 [0.065] N=505
CIAF: any stunting, underweight, or wasting, 0-60 months	0.077 [0.095] N=1,037	0.042 [0.098] N=913	0.085 [0.111] N=779	0.054 [0.130] N=633	0.161 [0.151] N=488
BMI-for-age z score below -2	**0.113** [0.048] N=1,046**	**0.123** [0.057] N=925**	**0.143** [0.065] N=792**	0.046 [0.064] N=642	0.060 [0.080] N=500

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.406	-0.227 [0.278] N=1,262	-0.284 [0.286] N=1,133	-0.294 [0.298] N=991	-0.523 [0.327] N=839	**-0.619* [0.366] N=686**
WHO weight-for-age z score	-0.015	0.025 [0.151] N=1,419	0.053 [0.157] N=1,286	-0.054 [0.156] N=1,125	-0.007 [0.165] N=945	-0.024 [0.182] N=764
WHO weight-for-length/height z score, 0-60 months	0.352	0.160 [0.206] N=1,245	0.073 [0.217] N=1,121	0.022 [0.230] N=983	0.229 [0.245] N=830	0.093 [0.266] N=679
WHO BMI-for-age z score	0.323	0.131 [0.234] N=1,209	0.057 [0.245] N=1,088	-0.036 [0.256] N=955	0.103 [0.278] N=805	0.078 [0.299] N=663
Stunted: HAZ below -2	0.243 P=310	0.031 [0.057] N=1,262	0.020 [0.057] N=1,133	0.018 [0.059] N=991	0.049 [0.064] N=839	0.073 [0.068] N=686
Underweight: WAZ below -2	0.070 P=101	-0.006 [0.031] N=1,419	-0.016 [0.033] N=1,286	0.002 [0.035] N=1,125	-0.004 [0.035] N=945	-0.004 [0.038] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=142	0.022 [0.037] N=1,245	0.023 [0.040] N=1,121	0.039 [0.042] N=983	-0.023 [0.042] N=830	-0.009 [0.047] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.090** [0.045] N=1,245**	0.073 [0.047] N=1,121	0.068 [0.050] N=983	0.085 [0.057] N=830	0.060 [0.059] N=679
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=115	0.052 [0.037] N=1,245	0.035 [0.038] N=1,121	0.026 [0.042] N=983	0.031 [0.049] N=830	-0.014 [0.049] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.334 P=406	0.052 [0.061] N=1,205	0.030 [0.062] N=1,081	0.048 [0.064] N=947	0.025 [0.068] N=801	0.074 [0.071] N=656
BMI-for-age z score below -2	0.122 P=148	0.027 [0.042] N=1,209	0.029 [0.045] N=1,088	0.032 [0.049] N=955	-0.016 [0.050] N=805	-0.012 [0.055] N=663

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.106 [0.386] N=1,088	-0.186 [0.414] N=959	-0.185 [0.464] N=817	-0.767 [0.558] N=665	-1.118 [0.681] N=512
WHO weight-for-age z score	0.044 [0.210] N=1,229	0.069 [0.230] N=1,096	-0.128 [0.231] N=935	-0.024 [0.256] N=755	0.030 [0.324] N=574
WHO weight-for-length/height z score, 0-60 months	0.064 [0.306] N=1,071	-0.102 [0.328] N=947	-0.214 [0.371] N=809	0.286 [0.425] N=656	-0.040 [0.520] N=505
WHO BMI-for-age z score	-0.171 [0.337] N=1,046	-0.316 [0.358] N=925	-0.533 [0.400] N=792	-0.291 [0.475] N=642	-0.387 [0.565] N=500
Stunted: HAZ below -2	0.039 [0.089] N=1,088	0.016 [0.093] N=959	0.017 [0.104] N=817	0.105 [0.127] N=665	0.194 [0.148] N=512
Underweight: WAZ below -2	-0.023 [0.046] N=1,229	-0.038 [0.049] N=1,096	-0.011 [0.056] N=935	-0.042 [0.057] N=755	-0.071 [0.062] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,071	0.082 [0.054] N=947	**0.122** [0.059] N=809**	-0.010 [0.059] N=656	0.013 [0.073] N=505
Overweight: WLZ/WHZ above +2, 0-60 months	**0.144** [0.059] N=1,071**	**0.119* [0.061] N=947**	**0.121* [0.068] N=809**	**0.183** [0.082] N=656**	0.127 [0.091] N=505
Obese: WLZ/WHZ above +3, 0-60 months	**0.109*** [0.041] N=1,071**	**0.080* [0.041] N=947**	0.076 [0.048] N=809	0.101 [0.065] N=656	-0.026 [0.064] N=505
CIAF: any stunting, underweight, or wasting, 0-60 months	0.080 [0.094] N=1,037	0.042 [0.099] N=913	0.082 [0.111] N=779	0.045 [0.130] N=633	0.148 [0.149] N=488
BMI-for-age z score below -2	**0.112** [0.049] N=1,046**	**0.122** [0.057] N=925**	**0.141** [0.066] N=792**	0.040 [0.065] N=642	0.051 [0.081] N=500

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.406	-0.207 [0.276] N=1,262	-0.262 [0.284] N=1,133	-0.272 [0.296] N=991	-0.519 [0.325] N=839	**-0.623* [0.369] N=686**
WHO weight-for-age z score	-0.015	0.029 [0.153] N=1,419	0.053 [0.160] N=1,286	-0.054 [0.159] N=1,125	-0.014 [0.169] N=945	-0.036 [0.186] N=764
WHO weight-for-length/height z score, 0-60 months	0.352	0.151 [0.206] N=1,245	0.065 [0.217] N=1,121	0.014 [0.229] N=983	0.235 [0.245] N=830	0.104 [0.268] N=679
WHO BMI-for-age z score	0.323	0.122 [0.233] N=1,209	0.047 [0.243] N=1,088	-0.049 [0.254] N=955	0.098 [0.277] N=805	0.088 [0.301] N=663
Stunted: HAZ below -2	0.243 P=310	0.028 [0.056] N=1,262	0.017 [0.057] N=1,133	0.018 [0.059] N=991	0.052 [0.064] N=839	0.077 [0.070] N=686
Underweight: WAZ below -2	0.070 P=101	-0.007 [0.032] N=1,419	-0.016 [0.033] N=1,286	0.001 [0.036] N=1,125	-0.004 [0.036] N=945	-0.006 [0.040] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=142	0.023 [0.037] N=1,245	0.025 [0.039] N=1,121	0.042 [0.041] N=983	-0.020 [0.041] N=830	-0.008 [0.047] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.090** [0.045] N=1,245**	0.074 [0.047] N=1,121	0.071 [0.050] N=983	0.091 [0.056] N=830	0.066 [0.057] N=679
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=115	0.054 [0.037] N=1,245	0.038 [0.037] N=1,121	0.030 [0.041] N=983	0.037 [0.048] N=830	-0.008 [0.047] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.334 P=406	0.050 [0.061] N=1,205	0.030 [0.062] N=1,081	0.052 [0.064] N=947	0.034 [0.067] N=801	0.084 [0.071] N=656
BMI-for-age z score below -2	0.122 P=148	0.024 [0.041] N=1,209	0.026 [0.045] N=1,088	0.031 [0.049] N=955	-0.018 [0.049] N=805	-0.018 [0.054] N=663

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.088 [0.378] N=1,088	-0.156 [0.405] N=959	-0.145 [0.458] N=817	-0.783 [0.544] N=665	**-1.227* [0.652] N=512**
WHO weight-for-age z score	0.070 [0.204] N=1,229	0.088 [0.226] N=1,096	-0.101 [0.224] N=935	-0.026 [0.244] N=755	0.009 [0.301] N=574
WHO weight-for-length/height z score, 0-60 months	0.065 [0.306] N=1,071	-0.120 [0.327] N=947	-0.238 [0.368] N=809	0.276 [0.429] N=656	0.002 [0.548] N=505
WHO BMI-for-age z score	-0.189 [0.336] N=1,046	-0.350 [0.358] N=925	-0.569 [0.399] N=792	-0.309 [0.480] N=642	-0.332 [0.592] N=500
Stunted: HAZ below -2	0.035 [0.086] N=1,088	0.009 [0.091] N=959	0.012 [0.103] N=817	0.113 [0.126] N=665	0.227 [0.146] N=512
Underweight: WAZ below -2	-0.023 [0.044] N=1,229	-0.037 [0.048] N=1,096	-0.011 [0.055] N=935	-0.036 [0.055] N=755	-0.067 [0.060] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	**0.078* [0.047] N=1,071**	**0.090* [0.053] N=947**	**0.132** [0.057] N=809**	-0.003 [0.058] N=656	0.018 [0.072] N=505
Overweight: WLZ/WHZ above +2, 0-60 months	**0.142** [0.059] N=1,071**	**0.116* [0.061] N=947**	**0.120* [0.068] N=809**	**0.183** [0.082] N=656**	0.139 [0.094] N=505
Obese: WLZ/WHZ above +3, 0-60 months	**0.113*** [0.041] N=1,071**	**0.083** [0.041] N=947**	0.077 [0.047] N=809	0.102 [0.064] N=656	-0.020 [0.067] N=505
CIAF: any stunting, underweight, or wasting, 0-60 months	0.078 [0.093] N=1,037	0.041 [0.098] N=913	0.083 [0.110] N=779	0.054 [0.128] N=633	0.185 [0.147] N=488
BMI-for-age z score below -2	**0.116** [0.048] N=1,046**	**0.128** [0.057] N=925**	**0.150** [0.065] N=792**	0.049 [0.064] N=642	0.057 [0.081] N=500

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.414	-0.228 [0.290] N=1,176	-0.295 [0.307] N=1,055	-0.285 [0.318] N=928	-0.538 [0.348] N=784	**-0.641* [0.382] N=642**
WHO weight-for-age z score	-0.014	0.073 [0.158] N=1,317	0.115 [0.168] N=1,192	-0.032 [0.165] N=1,048	-0.013 [0.178] N=881	-0.069 [0.190] N=715
WHO weight-for-length/height z score, 0-60 months	0.355	0.250 [0.224] N=1,157	0.176 [0.240] N=1,041	0.074 [0.250] N=916	0.254 [0.268] N=773	0.075 [0.282] N=634
WHO BMI-for-age z score	0.324	0.162 [0.253] N=1,124	0.103 [0.269] N=1,011	-0.044 [0.277] N=891	0.063 [0.300] N=751	-0.011 [0.315] N=620
Stunted: HAZ below -2	0.244 P=288	0.019 [0.060] N=1,176	0.012 [0.061] N=1,055	0.004 [0.063] N=928	0.031 [0.067] N=784	0.056 [0.072] N=642
Underweight: WAZ below -2	0.070 P=94	-0.015 [0.032] N=1,317	-0.027 [0.034] N=1,192	-0.008 [0.036] N=1,048	-0.011 [0.037] N=881	-0.005 [0.040] N=715
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=132	0.022 [0.039] N=1,157	0.022 [0.042] N=1,041	0.043 [0.043] N=916	-0.008 [0.044] N=773	0.013 [0.049] N=634
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=222	**0.096** [0.046] N=1,157**	**0.080* [0.048] N=1,041**	0.068 [0.051] N=916	0.080 [0.057] N=773	0.051 [0.058] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=110	0.061 [0.037] N=1,157	0.043 [0.038] N=1,041	0.033 [0.041] N=916	0.043 [0.048] N=773	-0.001 [0.047] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=378	0.041 [0.065] N=1,122	0.019 [0.066] N=1,006	0.038 [0.068] N=885	0.018 [0.071] N=747	0.079 [0.076] N=613
BMI-for-age z score below -2	0.122 P=137	0.033 [0.045] N=1,124	0.032 [0.049] N=1,011	0.037 [0.051] N=891	-0.000 [0.053] N=751	0.010 [0.057] N=620

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.043 [0.402] N=1,010	-0.118 [0.439] N=889	-0.117 [0.501] N=762	-0.773 [0.588] N=618	-1.157 [0.697] N=476
WHO weight-for-age z score	0.172 [0.214] N=1,137	0.236 [0.233] N=1,012	-0.018 [0.229] N=868	0.069 [0.254] N=701	0.099 [0.304] N=535
WHO weight-for-length/height z score, 0-60 months	0.189 [0.340] N=992	0.048 [0.370] N=876	-0.123 [0.415] N=751	0.342 [0.469] N=608	-0.004 [0.584] N=469
WHO BMI-for-age z score	-0.114 [0.374] N=968	-0.234 [0.409] N=855	-0.507 [0.454] N=735	-0.277 [0.527] N=595	-0.445 [0.627] N=464
Stunted: HAZ below -2	0.020 [0.095] N=1,010	-0.000 [0.100] N=889	-0.008 [0.112] N=762	0.066 [0.133] N=618	0.163 [0.153] N=476
Underweight: WAZ below -2	-0.041 [0.046] N=1,137	-0.063 [0.050] N=1,012	-0.038 [0.055] N=868	-0.068 [0.055] N=701	-0.093 [0.058] N=535
Wasted: WLZ/WHZ below -2, 0-60 months	0.070 [0.051] N=992	0.072 [0.059] N=876	**0.116* [0.061] N=751**	0.003 [0.066] N=608	0.039 [0.081] N=469
Overweight: WLZ/WHZ above +2, 0-60 months	**0.153** [0.063] N=992**	**0.127* [0.066] N=876**	**0.124* [0.073] N=751**	**0.176** [0.088] N=608**	0.122 [0.104] N=469
Obese: WLZ/WHZ above +3, 0-60 months	**0.115** [0.045] N=992**	**0.083* [0.046] N=876**	0.076 [0.052] N=751	0.108 [0.070] N=608	-0.025 [0.074] N=469
CIAF: any stunting, underweight, or wasting, 0-60 months	0.060 [0.103] N=962	0.015 [0.109] N=846	0.056 [0.122] N=725	0.011 [0.143] N=587	0.149 [0.167] N=453
BMI-for-age z score below -2	**0.121** [0.054] N=968**	**0.125* [0.064] N=855**	**0.144** [0.071] N=735**	0.062 [0.073] N=595	0.096 [0.088] N=464

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.414	-0.250 [0.296] N=1,176	-0.321 [0.312] N=1,055	-0.313 [0.323] N=928	-0.584 [0.356] N=784	**-0.700* [0.396] N=642**
WHO weight-for-age z score	-0.014	0.072 [0.158] N=1,317	0.108 [0.169] N=1,192	-0.037 [0.165] N=1,048	-0.023 [0.178] N=881	-0.082 [0.193] N=715
WHO weight-for-length/height z score, 0-60 months	0.355	0.274 [0.225] N=1,157	0.194 [0.240] N=1,041	0.097 [0.251] N=916	0.284 [0.266] N=773	0.108 [0.283] N=634
WHO BMI-for-age z score	0.324	0.189 [0.256] N=1,124	0.124 [0.272] N=1,011	-0.015 [0.281] N=891	0.097 [0.302] N=751	0.027 [0.320] N=620
Stunted: HAZ below -2	0.244 P=288	0.021 [0.059] N=1,176	0.014 [0.061] N=1,055	0.005 [0.062] N=928	0.034 [0.067] N=784	0.061 [0.072] N=642
Underweight: WAZ below -2	0.070 P=94	-0.015 [0.032] N=1,317	-0.026 [0.034] N=1,192	-0.008 [0.036] N=1,048	-0.010 [0.037] N=881	-0.003 [0.040] N=715
Wasted: WLZ/WHZ below -2, 0-60 months	0.112 P=132	0.018 [0.040] N=1,157	0.017 [0.043] N=1,041	0.037 [0.044] N=916	-0.016 [0.045] N=773	0.005 [0.049] N=634
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=222	**0.098** [0.045] N=1,157**	**0.081* [0.047] N=1,041**	0.068 [0.050] N=916	0.082 [0.057] N=773	0.056 [0.058] N=634
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=110	**0.061* [0.036] N=1,157**	0.042 [0.037] N=1,041	0.032 [0.040] N=916	0.043 [0.048] N=773	0.000 [0.046] N=634
CIAF: any stunting, underweight, or wasting, 0-60 months	0.335 P=378	0.039 [0.064] N=1,122	0.016 [0.064] N=1,006	0.032 [0.066] N=885	0.013 [0.069] N=747	0.075 [0.074] N=613
BMI-for-age z score below -2	0.122 P=137	0.028 [0.045] N=1,124	0.026 [0.049] N=1,011	0.031 [0.052] N=891	-0.008 [0.054] N=751	0.003 [0.058] N=620

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.077 [0.408] N=1,010	-0.150 [0.441] N=889	-0.156 [0.501] N=762	-0.885 [0.599] N=618	**-1.344* [0.714] N=476**
WHO weight-for-age z score	0.164 [0.216] N=1,137	0.219 [0.237] N=1,012	-0.037 [0.233] N=868	0.040 [0.258] N=701	0.050 [0.317] N=535
WHO weight-for-length/height z score, 0-60 months	0.196 [0.338] N=992	0.042 [0.364] N=876	-0.127 [0.410] N=751	0.374 [0.463] N=608	0.018 [0.569] N=469
WHO BMI-for-age z score	-0.101 [0.374] N=968	-0.233 [0.404] N=855	-0.498 [0.452] N=735	-0.242 [0.526] N=595	-0.383 [0.627] N=464
Stunted: HAZ below -2	0.025 [0.095] N=1,010	0.001 [0.099] N=889	-0.007 [0.111] N=762	0.078 [0.132] N=618	0.182 [0.152] N=476
Underweight: WAZ below -2	-0.040 [0.046] N=1,137	-0.061 [0.050] N=1,012	-0.035 [0.055] N=868	-0.062 [0.055] N=701	-0.082 [0.059] N=535
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.051] N=992	0.072 [0.058] N=876	**0.114* [0.061] N=751**	-0.005 [0.064] N=608	0.029 [0.079] N=469
Overweight: WLZ/WHZ above +2, 0-60 months	**0.156** [0.063] N=992**	**0.129** [0.065] N=876**	**0.124* [0.072] N=751**	**0.182** [0.087] N=608**	0.129 [0.103] N=469
Obese: WLZ/WHZ above +3, 0-60 months	**0.115** [0.045] N=992**	**0.082* [0.045] N=876**	0.074 [0.050] N=751	0.108 [0.069] N=608	-0.022 [0.072] N=469
CIAF: any stunting, underweight, or wasting, 0-60 months	0.063 [0.103] N=962	0.016 [0.107] N=846	0.053 [0.120] N=725	0.010 [0.141] N=587	0.145 [0.164] N=453
BMI-for-age z score below -2	**0.121** [0.053] N=968**	**0.125** [0.063] N=855**	**0.143** [0.071] N=735**	0.057 [0.072] N=595	0.086 [0.086] N=464

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.