Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Oldest Child

Age reference: 12 November 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,522	-0.013 [0.022] N=1,618	0.002 [0.022] N=1,494	-0.007 [0.023] N=1,344	-0.009 [0.026] N=1,147	-0.011 [0.031] N=940
Vegetables consumed at least weekly	0.901 P=1,455	0.009 [0.026] N=1,618	0.011 [0.027] N=1,494	-0.008 [0.028] N=1,344	-0.011 [0.029] N=1,147	-0.043 [0.033] N=940
Meat fish or poultry consumed at least weekly	0.799 P=1,298	**0.078* [0.043] N=1,618**	**0.091** [0.044] N=1,494**	0.073 [0.049] N=1,344	0.057 [0.054] N=1,147	0.039 [0.060] N=940
Legumes consumed at least weekly	0.798 P=1,293	-0.056 [0.040] N=1,618	-0.067 [0.041] N=1,494	**-0.074* [0.043] N=1,344**	-0.069 [0.048] N=1,147	**-0.124** [0.055] N=940**
Animal products consumed at least weekly	0.916 P=1,482	0.029 [0.030] N=1,618	0.032 [0.030] N=1,494	0.019 [0.034] N=1,344	0.020 [0.038] N=1,147	0.009 [0.043] N=940
Sweets consumed at least weekly	0.703 P=1,133	-0.025 [0.038] N=1,618	-0.003 [0.040] N=1,494	0.008 [0.042] N=1,344	0.005 [0.045] N=1,147	0.018 [0.051] N=940
Snack foods consumed at least weekly	0.575 P=924	-0.038 [0.043] N=1,618	-0.024 [0.046] N=1,494	-0.015 [0.049] N=1,344	-0.027 [0.055] N=1,147	-0.043 [0.060] N=940
Packaged soft drinks consumed at least weekly	0.354 P=564	0.021 [0.049] N=1,618	0.030 [0.053] N=1,494	0.029 [0.056] N=1,344	0.055 [0.062] N=1,147	0.046 [0.071] N=940
Grains consumed at least weekly	0.918 P=1,484	0.002 [0.027] N=1,618	0.012 [0.029] N=1,494	-0.009 [0.029] N=1,344	-0.007 [0.032] N=1,147	-0.012 [0.038] N=940
Child was ever breastfed	0.965 P=1,557	0.009 [0.016] N=1,619	0.013 [0.017] N=1,495	0.019 [0.019] N=1,344	0.024 [0.021] N=1,147	0.017 [0.021] N=940

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.032] N=1,379	0.027 [0.031] N=1,255	0.015 [0.033] N=1,105	0.014 [0.039] N=908	0.033 [0.056] N=701
Vegetables consumed at least weekly	0.048 [0.040] N=1,379	0.061 [0.043] N=1,255	0.033 [0.047] N=1,105	0.038 [0.056] N=908	-0.025 [0.075] N=701
Meat fish or poultry consumed at least weekly	0.073 [0.055] N=1,379	0.094 [0.057] N=1,255	0.068 [0.066] N=1,105	0.020 [0.076] N=908	-0.040 [0.087] N=701
Legumes consumed at least weekly	0.007 [0.060] N=1,379	-0.005 [0.064] N=1,255	0.008 [0.068] N=1,105	0.043 [0.083] N=908	-0.046 [0.113] N=701
Animal products consumed at least weekly	0.037 [0.038] N=1,379	0.043 [0.040] N=1,255	0.023 [0.047] N=1,105	0.032 [0.057] N=908	0.011 [0.074] N=701
Sweets consumed at least weekly	-0.054 [0.052] N=1,379	-0.020 [0.056] N=1,255	-0.002 [0.062] N=1,105	-0.020 [0.071] N=908	0.037 [0.094] N=701
Snack foods consumed at least weekly	-0.058 [0.055] N=1,379	-0.040 [0.061] N=1,255	-0.023 [0.073] N=1,105	-0.054 [0.089] N=908	-0.122 [0.100] N=701
Packaged soft drinks consumed at least weekly	-0.022 [0.061] N=1,379	-0.015 [0.067] N=1,255	-0.023 [0.072] N=1,105	0.008 [0.087] N=908	-0.034 [0.111] N=701
Grains consumed at least weekly	0.040 [0.044] N=1,379	0.068 [0.049] N=1,255	0.034 [0.053] N=1,105	0.061 [0.063] N=908	0.086 [0.094] N=701
Child was ever breastfed	0.016 [0.026] N=1,380	0.025 [0.030] N=1,256	0.036 [0.035] N=1,105	0.049 [0.046] N=908	0.054 [0.058] N=701

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,522	-0.013 [0.022] N=1,618	0.002 [0.022] N=1,494	-0.006 [0.023] N=1,344	-0.009 [0.025] N=1,147	-0.010 [0.031] N=940
Vegetables consumed at least weekly	0.901 P=1,455	0.009 [0.026] N=1,618	0.011 [0.027] N=1,494	-0.008 [0.028] N=1,344	-0.010 [0.028] N=1,147	-0.042 [0.033] N=940
Meat fish or poultry consumed at least weekly	0.799 P=1,298	**0.078* [0.043] N=1,618**	**0.091** [0.044] N=1,494**	0.074 [0.049] N=1,344	0.059 [0.054] N=1,147	0.040 [0.061] N=940
Legumes consumed at least weekly	0.798 P=1,293	-0.056 [0.040] N=1,618	-0.067 [0.041] N=1,494	**-0.073* [0.043] N=1,344**	-0.069 [0.048] N=1,147	**-0.123** [0.055] N=940**
Animal products consumed at least weekly	0.916 P=1,482	0.029 [0.030] N=1,618	0.033 [0.030] N=1,494	0.021 [0.034] N=1,344	0.021 [0.037] N=1,147	0.010 [0.042] N=940
Sweets consumed at least weekly	0.703 P=1,133	-0.025 [0.038] N=1,618	-0.003 [0.040] N=1,494	0.008 [0.042] N=1,344	0.005 [0.045] N=1,147	0.018 [0.051] N=940
Snack foods consumed at least weekly	0.575 P=924	-0.038 [0.044] N=1,618	-0.023 [0.046] N=1,494	-0.014 [0.050] N=1,344	-0.026 [0.056] N=1,147	-0.042 [0.060] N=940
Packaged soft drinks consumed at least weekly	0.354 P=564	0.021 [0.049] N=1,618	0.030 [0.053] N=1,494	0.029 [0.056] N=1,344	0.055 [0.062] N=1,147	0.046 [0.071] N=940
Grains consumed at least weekly	0.918 P=1,484	0.002 [0.027] N=1,618	0.012 [0.029] N=1,494	-0.010 [0.029] N=1,344	-0.007 [0.031] N=1,147	-0.012 [0.038] N=940
Child was ever breastfed	0.965 P=1,557	0.009 [0.016] N=1,619	0.013 [0.017] N=1,495	0.020 [0.019] N=1,344	0.024 [0.021] N=1,147	0.017 [0.021] N=940

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.002 [0.032] N=1,379	0.028 [0.031] N=1,255	0.016 [0.033] N=1,105	0.014 [0.039] N=908	0.034 [0.056] N=701
Vegetables consumed at least weekly	0.048 [0.040] N=1,379	0.061 [0.043] N=1,255	0.034 [0.047] N=1,105	0.039 [0.056] N=908	-0.025 [0.075] N=701
Meat fish or poultry consumed at least weekly	0.073 [0.055] N=1,379	0.093 [0.057] N=1,255	0.069 [0.066] N=1,105	0.023 [0.076] N=908	-0.033 [0.086] N=701
Legumes consumed at least weekly	0.008 [0.060] N=1,379	-0.005 [0.064] N=1,255	0.010 [0.068] N=1,105	0.046 [0.083] N=908	-0.044 [0.113] N=701
Animal products consumed at least weekly	0.038 [0.038] N=1,379	0.047 [0.039] N=1,255	0.029 [0.047] N=1,105	0.036 [0.057] N=908	0.014 [0.074] N=701
Sweets consumed at least weekly	-0.055 [0.052] N=1,379	-0.020 [0.056] N=1,255	-0.003 [0.063] N=1,105	-0.021 [0.070] N=908	0.036 [0.094] N=701
Snack foods consumed at least weekly	-0.057 [0.056] N=1,379	-0.036 [0.061] N=1,255	-0.016 [0.073] N=1,105	-0.047 [0.090] N=908	-0.115 [0.100] N=701
Packaged soft drinks consumed at least weekly	-0.022 [0.061] N=1,379	-0.015 [0.067] N=1,255	-0.024 [0.072] N=1,105	0.007 [0.087] N=908	-0.037 [0.111] N=701
Grains consumed at least weekly	0.040 [0.044] N=1,379	0.068 [0.049] N=1,255	0.034 [0.052] N=1,105	0.059 [0.063] N=908	0.084 [0.092] N=701
Child was ever breastfed	0.017 [0.026] N=1,380	0.026 [0.030] N=1,256	0.037 [0.035] N=1,105	0.048 [0.046] N=908	0.056 [0.058] N=701

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,522	-0.021 [0.022] N=1,618	-0.007 [0.022] N=1,494	-0.015 [0.024] N=1,344	-0.017 [0.026] N=1,147	-0.019 [0.032] N=940
Vegetables consumed at least weekly	0.901 P=1,455	0.004 [0.026] N=1,618	0.004 [0.027] N=1,494	-0.014 [0.028] N=1,344	-0.016 [0.028] N=1,147	-0.049 [0.033] N=940
Meat fish or poultry consumed at least weekly	0.799 P=1,298	0.054 [0.042] N=1,618	0.060 [0.043] N=1,494	0.049 [0.048] N=1,344	0.035 [0.052] N=1,147	0.014 [0.058] N=940
Legumes consumed at least weekly	0.798 P=1,293	-0.064 [0.041] N=1,618	**-0.077* [0.042] N=1,494**	**-0.082* [0.043] N=1,344**	-0.075 [0.048] N=1,147	**-0.128** [0.056] N=940**
Animal products consumed at least weekly	0.916 P=1,482	0.024 [0.030] N=1,618	0.026 [0.031] N=1,494	0.014 [0.035] N=1,344	0.015 [0.038] N=1,147	0.000 [0.043] N=940
Sweets consumed at least weekly	0.703 P=1,133	-0.032 [0.038] N=1,618	-0.011 [0.040] N=1,494	-0.001 [0.042] N=1,344	-0.003 [0.045] N=1,147	0.007 [0.051] N=940
Snack foods consumed at least weekly	0.575 P=924	-0.037 [0.043] N=1,618	-0.021 [0.046] N=1,494	-0.014 [0.048] N=1,344	-0.025 [0.054] N=1,147	-0.040 [0.059] N=940
Packaged soft drinks consumed at least weekly	0.354 P=564	0.033 [0.048] N=1,618	0.048 [0.051] N=1,494	0.043 [0.054] N=1,344	0.068 [0.060] N=1,147	0.063 [0.068] N=940
Grains consumed at least weekly	0.918 P=1,484	-0.006 [0.027] N=1,618	0.004 [0.029] N=1,494	-0.016 [0.029] N=1,344	-0.015 [0.031] N=1,147	-0.019 [0.038] N=940
Child was ever breastfed	0.965 P=1,557	0.005 [0.016] N=1,619	0.008 [0.017] N=1,495	0.014 [0.018] N=1,344	0.019 [0.020] N=1,147	0.013 [0.021] N=940

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.009 [0.031] N=1,379	0.019 [0.030] N=1,255	0.011 [0.033] N=1,105	0.020 [0.038] N=908	0.042 [0.055] N=701
Vegetables consumed at least weekly	0.045 [0.039] N=1,379	0.056 [0.042] N=1,255	0.030 [0.047] N=1,105	0.043 [0.055] N=908	-0.020 [0.074] N=701
Meat fish or poultry consumed at least weekly	0.054 [0.054] N=1,379	0.064 [0.056] N=1,255	0.049 [0.066] N=1,105	0.012 [0.076] N=908	-0.040 [0.090] N=701
Legumes consumed at least weekly	0.001 [0.059] N=1,379	-0.016 [0.064] N=1,255	0.003 [0.068] N=1,105	0.048 [0.084] N=908	-0.039 [0.115] N=701
Animal products consumed at least weekly	0.035 [0.037] N=1,379	0.041 [0.039] N=1,255	0.024 [0.047] N=1,105	0.040 [0.057] N=908	0.021 [0.073] N=701
Sweets consumed at least weekly	-0.059 [0.052] N=1,379	-0.025 [0.057] N=1,255	-0.007 [0.064] N=1,105	-0.019 [0.070] N=908	0.042 [0.094] N=701
Snack foods consumed at least weekly	-0.056 [0.055] N=1,379	-0.034 [0.061] N=1,255	-0.015 [0.070] N=1,105	-0.043 [0.086] N=908	-0.105 [0.096] N=701
Packaged soft drinks consumed at least weekly	-0.011 [0.059] N=1,379	0.005 [0.065] N=1,255	-0.010 [0.070] N=1,105	0.022 [0.086] N=908	-0.022 [0.107] N=701
Grains consumed at least weekly	0.032 [0.043] N=1,379	0.061 [0.049] N=1,255	0.028 [0.052] N=1,105	0.054 [0.062] N=908	0.078 [0.091] N=701
Child was ever breastfed	0.014 [0.026] N=1,380	0.021 [0.030] N=1,256	0.033 [0.035] N=1,105	0.048 [0.046] N=908	0.051 [0.058] N=701

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,522	-0.023 [0.022] N=1,618	-0.009 [0.022] N=1,494	-0.018 [0.024] N=1,344	-0.020 [0.027] N=1,147	-0.020 [0.032] N=940
Vegetables consumed at least weekly	0.901 P=1,455	0.003 [0.026] N=1,618	0.005 [0.027] N=1,494	-0.013 [0.028] N=1,344	-0.016 [0.029] N=1,147	-0.048 [0.033] N=940
Meat fish or poultry consumed at least weekly	0.799 P=1,298	0.050 [0.042] N=1,618	0.055 [0.043] N=1,494	0.040 [0.047] N=1,344	0.026 [0.052] N=1,147	0.002 [0.057] N=940
Legumes consumed at least weekly	0.798 P=1,293	-0.066 [0.041] N=1,618	**-0.082* [0.043] N=1,494**	**-0.090** [0.044] N=1,344**	-0.080 [0.050] N=1,147	**-0.132** [0.057] N=940**
Animal products consumed at least weekly	0.916 P=1,482	0.022 [0.030] N=1,618	0.026 [0.031] N=1,494	0.014 [0.034] N=1,344	0.012 [0.038] N=1,147	-0.001 [0.042] N=940
Sweets consumed at least weekly	0.703 P=1,133	-0.031 [0.038] N=1,618	-0.012 [0.040] N=1,494	-0.005 [0.041] N=1,344	-0.006 [0.045] N=1,147	0.004 [0.051] N=940
Snack foods consumed at least weekly	0.575 P=924	-0.038 [0.043] N=1,618	-0.024 [0.046] N=1,494	-0.019 [0.048] N=1,344	-0.027 [0.054] N=1,147	-0.044 [0.059] N=940
Packaged soft drinks consumed at least weekly	0.354 P=564	0.038 [0.048] N=1,618	0.049 [0.051] N=1,494	0.046 [0.054] N=1,344	0.074 [0.060] N=1,147	0.072 [0.067] N=940
Grains consumed at least weekly	0.918 P=1,484	-0.005 [0.027] N=1,618	0.005 [0.028] N=1,494	-0.017 [0.029] N=1,344	-0.016 [0.031] N=1,147	-0.019 [0.039] N=940
Child was ever breastfed	0.965 P=1,557	0.004 [0.016] N=1,619	0.008 [0.017] N=1,495	0.013 [0.019] N=1,344	0.018 [0.021] N=1,147	0.012 [0.021] N=940

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.010 [0.031] N=1,379	0.018 [0.030] N=1,255	0.010 [0.033] N=1,105	0.014 [0.039] N=908	0.033 [0.056] N=701
Vegetables consumed at least weekly	0.044 [0.040] N=1,379	0.054 [0.043] N=1,255	0.029 [0.047] N=1,105	0.038 [0.057] N=908	-0.033 [0.076] N=701
Meat fish or poultry consumed at least weekly	0.050 [0.055] N=1,379	0.060 [0.057] N=1,255	0.045 [0.067] N=1,105	0.010 [0.078] N=908	-0.051 [0.094] N=701
Legumes consumed at least weekly	0.005 [0.059] N=1,379	-0.013 [0.064] N=1,255	0.006 [0.066] N=1,105	0.063 [0.082] N=908	-0.028 [0.119] N=701
Animal products consumed at least weekly	0.034 [0.037] N=1,379	0.043 [0.040] N=1,255	0.025 [0.047] N=1,105	0.034 [0.057] N=908	0.012 [0.078] N=701
Sweets consumed at least weekly	-0.054 [0.052] N=1,379	-0.024 [0.055] N=1,255	-0.007 [0.061] N=1,105	-0.009 [0.069] N=908	0.047 [0.094] N=701
Snack foods consumed at least weekly	-0.049 [0.054] N=1,379	-0.032 [0.059] N=1,255	-0.013 [0.068] N=1,105	-0.028 [0.085] N=908	-0.090 [0.095] N=701
Packaged soft drinks consumed at least weekly	-0.004 [0.059] N=1,379	0.005 [0.065] N=1,255	-0.008 [0.071] N=1,105	0.029 [0.087] N=908	-0.007 [0.108] N=701
Grains consumed at least weekly	0.032 [0.042] N=1,379	0.059 [0.047] N=1,255	0.026 [0.050] N=1,105	0.052 [0.059] N=908	0.070 [0.085] N=701
Child was ever breastfed	0.014 [0.026] N=1,380	0.022 [0.030] N=1,256	0.033 [0.036] N=1,105	0.048 [0.047] N=908	0.054 [0.059] N=701

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,417	-0.021 [0.023] N=1,507	-0.009 [0.023] N=1,398	-0.017 [0.024] N=1,256	-0.018 [0.027] N=1,077	-0.023 [0.032] N=883
Vegetables consumed at least weekly	0.902 P=1,356	-0.002 [0.027] N=1,507	-0.003 [0.028] N=1,398	-0.026 [0.029] N=1,256	-0.023 [0.029] N=1,077	-0.053 [0.032] N=883
Meat fish or poultry consumed at least weekly	0.797 P=1,205	0.067 [0.041] N=1,507	**0.070* [0.042] N=1,398**	0.046 [0.046] N=1,256	0.029 [0.050] N=1,077	-0.003 [0.056] N=883
Legumes consumed at least weekly	0.794 P=1,198	-0.066 [0.040] N=1,507	**-0.082** [0.041] N=1,398**	**-0.093** [0.042] N=1,256**	**-0.082* [0.048] N=1,077**	**-0.140** [0.054] N=883**
Animal products consumed at least weekly	0.916 P=1,382	0.029 [0.030] N=1,507	0.033 [0.031] N=1,398	0.020 [0.034] N=1,256	0.014 [0.038] N=1,077	0.001 [0.042] N=883
Sweets consumed at least weekly	0.705 P=1,058	-0.008 [0.039] N=1,507	0.010 [0.041] N=1,398	0.013 [0.043] N=1,256	0.014 [0.046] N=1,077	0.023 [0.052] N=883
Snack foods consumed at least weekly	0.577 P=862	-0.038 [0.047] N=1,507	-0.022 [0.050] N=1,398	-0.016 [0.053] N=1,256	-0.026 [0.059] N=1,077	-0.028 [0.066] N=883
Packaged soft drinks consumed at least weekly	0.361 P=535	0.048 [0.049] N=1,507	0.060 [0.052] N=1,398	0.047 [0.056] N=1,256	0.072 [0.061] N=1,077	0.078 [0.071] N=883
Grains consumed at least weekly	0.918 P=1,382	-0.002 [0.028] N=1,507	0.006 [0.030] N=1,398	-0.023 [0.030] N=1,256	-0.017 [0.032] N=1,077	-0.021 [0.038] N=883
Child was ever breastfed	0.965 P=1,451	0.006 [0.017] N=1,508	0.012 [0.019] N=1,399	0.019 [0.020] N=1,256	0.027 [0.023] N=1,077	0.018 [0.023] N=883

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.006 [0.034] N=1,283	0.014 [0.034] N=1,174	0.002 [0.036] N=1,032	0.002 [0.043] N=853	0.019 [0.058] N=659
Vegetables consumed at least weekly	0.038 [0.041] N=1,283	0.042 [0.046] N=1,174	0.003 [0.050] N=1,032	0.016 [0.058] N=853	-0.037 [0.072] N=659
Meat fish or poultry consumed at least weekly	0.087 [0.056] N=1,283	0.089 [0.058] N=1,174	0.061 [0.066] N=1,032	0.017 [0.077] N=853	-0.047 [0.084] N=659
Legumes consumed at least weekly	0.006 [0.061] N=1,283	-0.018 [0.064] N=1,174	-0.016 [0.066] N=1,032	0.023 [0.084] N=853	-0.073 [0.109] N=659
Animal products consumed at least weekly	0.043 [0.039] N=1,283	0.049 [0.042] N=1,174	0.031 [0.049] N=1,032	0.032 [0.059] N=853	0.012 [0.075] N=659
Sweets consumed at least weekly	-0.034 [0.053] N=1,283	-0.008 [0.058] N=1,174	-0.009 [0.063] N=1,032	-0.016 [0.072] N=853	0.012 [0.099] N=659
Snack foods consumed at least weekly	-0.063 [0.056] N=1,283	-0.043 [0.063] N=1,174	-0.036 [0.073] N=1,032	-0.072 [0.090] N=853	-0.141 [0.100] N=659
Packaged soft drinks consumed at least weekly	0.004 [0.062] N=1,283	0.013 [0.066] N=1,174	-0.016 [0.071] N=1,032	0.007 [0.085] N=853	-0.006 [0.115] N=659
Grains consumed at least weekly	0.042 [0.045] N=1,283	0.064 [0.050] N=1,174	0.012 [0.053] N=1,032	0.047 [0.062] N=853	0.077 [0.091] N=659
Child was ever breastfed	0.010 [0.027] N=1,284	0.022 [0.030] N=1,175	0.031 [0.036] N=1,032	0.046 [0.045] N=853	0.042 [0.059] N=659

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,417	-0.022 [0.022] N=1,507	-0.011 [0.022] N=1,398	-0.018 [0.024] N=1,256	-0.019 [0.027] N=1,077	-0.024 [0.031] N=883
Vegetables consumed at least weekly	0.902 P=1,356	-0.003 [0.027] N=1,507	-0.004 [0.028] N=1,398	-0.027 [0.029] N=1,256	-0.024 [0.029] N=1,077	**-0.054* [0.032] N=883**
Meat fish or poultry consumed at least weekly	0.797 P=1,205	0.068 [0.042] N=1,507	**0.071* [0.043] N=1,398**	0.052 [0.047] N=1,256	0.038 [0.052] N=1,077	0.014 [0.058] N=883
Legumes consumed at least weekly	0.794 P=1,198	**-0.069* [0.040] N=1,507**	**-0.085** [0.041] N=1,398**	**-0.095** [0.042] N=1,256**	**-0.088* [0.047] N=1,077**	**-0.152*** [0.054] N=883**
Animal products consumed at least weekly	0.916 P=1,382	0.028 [0.030] N=1,507	0.032 [0.031] N=1,398	0.019 [0.034] N=1,256	0.015 [0.037] N=1,077	0.003 [0.041] N=883
Sweets consumed at least weekly	0.705 P=1,058	-0.009 [0.039] N=1,507	0.009 [0.041] N=1,398	0.015 [0.042] N=1,256	0.016 [0.045] N=1,077	0.027 [0.051] N=883
Snack foods consumed at least weekly	0.577 P=862	-0.040 [0.046] N=1,507	-0.025 [0.049] N=1,398	-0.018 [0.052] N=1,256	-0.029 [0.059] N=1,077	-0.038 [0.064] N=883
Packaged soft drinks consumed at least weekly	0.361 P=535	0.046 [0.049] N=1,507	0.057 [0.052] N=1,398	0.047 [0.056] N=1,256	0.069 [0.061] N=1,077	0.068 [0.071] N=883
Grains consumed at least weekly	0.918 P=1,382	-0.003 [0.028] N=1,507	0.005 [0.030] N=1,398	-0.025 [0.030] N=1,256	-0.019 [0.033] N=1,077	-0.026 [0.038] N=883
Child was ever breastfed	0.965 P=1,451	0.005 [0.017] N=1,508	0.011 [0.018] N=1,399	0.017 [0.020] N=1,256	0.024 [0.022] N=1,077	0.015 [0.023] N=883

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.033] N=1,283	0.013 [0.032] N=1,174	0.004 [0.035] N=1,032	0.003 [0.042] N=853	0.018 [0.056] N=659
Vegetables consumed at least weekly	0.036 [0.041] N=1,283	0.043 [0.045] N=1,174	0.006 [0.050] N=1,032	0.016 [0.059] N=853	-0.039 [0.071] N=659
Meat fish or poultry consumed at least weekly	0.080 [0.056] N=1,283	0.083 [0.057] N=1,174	0.057 [0.066] N=1,032	0.018 [0.077] N=853	-0.034 [0.085] N=659
Legumes consumed at least weekly	0.002 [0.061] N=1,283	-0.021 [0.065] N=1,174	-0.013 [0.068] N=1,032	0.026 [0.086] N=853	-0.081 [0.112] N=659
Animal products consumed at least weekly	0.039 [0.039] N=1,283	0.046 [0.041] N=1,174	0.030 [0.049] N=1,032	0.029 [0.059] N=853	0.009 [0.075] N=659
Sweets consumed at least weekly	-0.045 [0.053] N=1,283	-0.023 [0.058] N=1,174	-0.018 [0.064] N=1,032	-0.020 [0.072] N=853	0.021 [0.099] N=659
Snack foods consumed at least weekly	-0.072 [0.056] N=1,283	-0.054 [0.062] N=1,174	-0.041 [0.073] N=1,032	-0.074 [0.090] N=853	-0.136 [0.100] N=659
Packaged soft drinks consumed at least weekly	-0.002 [0.062] N=1,283	0.007 [0.066] N=1,174	-0.019 [0.072] N=1,032	0.005 [0.086] N=853	-0.015 [0.113] N=659
Grains consumed at least weekly	0.041 [0.045] N=1,283	0.065 [0.051] N=1,174	0.017 [0.055] N=1,032	0.049 [0.065] N=853	0.074 [0.092] N=659
Child was ever breastfed	0.011 [0.028] N=1,284	0.023 [0.031] N=1,175	0.032 [0.038] N=1,032	0.050 [0.048] N=853	0.047 [0.062] N=659

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,360	-0.013 [0.027] N=1,441	-0.014 [0.029] N=1,312	-0.019 [0.032] N=1,151	-0.020 [0.033] N=967	-0.013 [0.039] N=781
Vegetables consumed at least weekly	0.903 P=1,299	0.005 [0.033] N=1,441	0.017 [0.035] N=1,312	0.020 [0.038] N=1,151	0.013 [0.042] N=967	0.014 [0.048] N=781
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.077* [0.041] N=1,441**	**0.081* [0.043] N=1,312**	0.076 [0.047] N=1,151	**0.108** [0.047] N=967**	**0.103* [0.056] N=781**
Legumes consumed at least weekly	0.797 P=1,154	-0.046 [0.046] N=1,441	-0.027 [0.049] N=1,312	-0.036 [0.053] N=1,151	-0.053 [0.058] N=967	-0.026 [0.069] N=781
Animal products consumed at least weekly	0.915 P=1,319	-0.004 [0.034] N=1,441	0.005 [0.036] N=1,312	0.013 [0.038] N=1,151	-0.005 [0.041] N=967	0.028 [0.047] N=781
Sweets consumed at least weekly	0.706 P=1,020	0.007 [0.050] N=1,441	0.011 [0.052] N=1,312	0.002 [0.056] N=1,151	0.004 [0.060] N=967	-0.043 [0.069] N=781
Snack foods consumed at least weekly	0.572 P=822	-0.019 [0.057] N=1,441	-0.011 [0.060] N=1,312	-0.005 [0.064] N=1,151	0.013 [0.068] N=967	-0.030 [0.078] N=781
Packaged soft drinks consumed at least weekly	0.338 P=483	-0.002 [0.048] N=1,441	-0.003 [0.050] N=1,312	0.036 [0.052] N=1,151	0.050 [0.058] N=967	0.032 [0.065] N=781
Grains consumed at least weekly	0.919 P=1,323	0.027 [0.038] N=1,441	0.034 [0.040] N=1,312	0.036 [0.043] N=1,151	0.043 [0.048] N=967	0.049 [0.054] N=781
Child was ever breastfed	0.963 P=1,388	-0.007 [0.023] N=1,442	-0.005 [0.023] N=1,313	-0.004 [0.024] N=1,152	-0.002 [0.024] N=968	-0.020 [0.027] N=782

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.050 [0.034] N=1,253	-0.057 [0.037] N=1,124	**-0.080* [0.043] N=963**	-0.075 [0.046] N=779	-0.083 [0.067] N=593
Vegetables consumed at least weekly	-0.053 [0.038] N=1,253	-0.043 [0.043] N=1,124	-0.052 [0.047] N=963	-0.064 [0.053] N=779	-0.087 [0.066] N=593
Meat fish or poultry consumed at least weekly	0.019 [0.057] N=1,253	0.013 [0.061] N=1,124	-0.022 [0.068] N=963	0.032 [0.071] N=779	-0.012 [0.091] N=593
Legumes consumed at least weekly	-0.070 [0.053] N=1,253	-0.046 [0.058] N=1,124	-0.066 [0.069] N=963	-0.083 [0.076] N=779	-0.009 [0.093] N=593
Animal products consumed at least weekly	**-0.064* [0.038] N=1,253**	-0.050 [0.043] N=1,124	-0.046 [0.045] N=963	**-0.093* [0.047] N=779**	-0.051 [0.060] N=593
Sweets consumed at least weekly	0.007 [0.066] N=1,253	0.019 [0.070] N=1,124	0.003 [0.079] N=963	0.008 [0.092] N=779	-0.123 [0.117] N=593
Snack foods consumed at least weekly	-0.006 [0.086] N=1,253	0.009 [0.092] N=1,124	0.018 [0.104] N=963	0.073 [0.121] N=779	-0.014 [0.158] N=593
Packaged soft drinks consumed at least weekly	-0.040 [0.078] N=1,253	-0.044 [0.083] N=1,124	0.018 [0.095] N=963	0.043 [0.114] N=779	0.001 [0.145] N=593
Grains consumed at least weekly	-0.007 [0.039] N=1,253	-0.005 [0.042] N=1,124	-0.014 [0.048] N=963	-0.025 [0.059] N=779	-0.060 [0.075] N=593
Child was ever breastfed	0.024 [0.043] N=1,254	0.031 [0.045] N=1,125	0.038 [0.050] N=964	0.051 [0.055] N=780	0.030 [0.070] N=594

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,360	-0.013 [0.027] N=1,441	-0.013 [0.029] N=1,312	-0.019 [0.032] N=1,151	-0.020 [0.033] N=967	-0.013 [0.039] N=781
Vegetables consumed at least weekly	0.903 P=1,299	0.006 [0.033] N=1,441	0.017 [0.035] N=1,312	0.020 [0.038] N=1,151	0.012 [0.042] N=967	0.013 [0.048] N=781
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.078* [0.041] N=1,441**	**0.082* [0.043] N=1,312**	0.077 [0.047] N=1,151	**0.108** [0.047] N=967**	**0.102* [0.056] N=781**
Legumes consumed at least weekly	0.797 P=1,154	-0.046 [0.046] N=1,441	-0.027 [0.049] N=1,312	-0.036 [0.053] N=1,151	-0.053 [0.058] N=967	-0.025 [0.069] N=781
Animal products consumed at least weekly	0.915 P=1,319	-0.003 [0.033] N=1,441	0.007 [0.036] N=1,312	0.013 [0.038] N=1,151	-0.006 [0.040] N=967	0.026 [0.046] N=781
Sweets consumed at least weekly	0.706 P=1,020	0.008 [0.050] N=1,441	0.012 [0.052] N=1,312	0.002 [0.056] N=1,151	0.004 [0.060] N=967	-0.044 [0.069] N=781
Snack foods consumed at least weekly	0.572 P=822	-0.016 [0.057] N=1,441	-0.009 [0.060] N=1,312	-0.004 [0.064] N=1,151	0.012 [0.068] N=967	-0.032 [0.079] N=781
Packaged soft drinks consumed at least weekly	0.338 P=483	-0.001 [0.047] N=1,441	-0.003 [0.049] N=1,312	0.036 [0.052] N=1,151	0.050 [0.058] N=967	0.030 [0.065] N=781
Grains consumed at least weekly	0.919 P=1,323	0.027 [0.038] N=1,441	0.034 [0.040] N=1,312	0.036 [0.043] N=1,151	0.044 [0.048] N=967	0.050 [0.054] N=781
Child was ever breastfed	0.963 P=1,388	-0.006 [0.023] N=1,442	-0.004 [0.023] N=1,313	-0.003 [0.024] N=1,152	-0.002 [0.024] N=968	-0.020 [0.027] N=782

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.049 [0.034] N=1,253	-0.056 [0.037] N=1,124	**-0.080* [0.044] N=963**	-0.076 [0.046] N=779	-0.084 [0.067] N=593
Vegetables consumed at least weekly	-0.051 [0.038] N=1,253	-0.042 [0.043] N=1,124	-0.052 [0.047] N=963	-0.063 [0.053] N=779	-0.086 [0.066] N=593
Meat fish or poultry consumed at least weekly	0.022 [0.057] N=1,253	0.017 [0.061] N=1,124	-0.018 [0.068] N=963	0.035 [0.072] N=779	-0.007 [0.093] N=593
Legumes consumed at least weekly	-0.070 [0.052] N=1,253	-0.047 [0.058] N=1,124	-0.067 [0.069] N=963	-0.084 [0.075] N=779	-0.012 [0.092] N=593
Animal products consumed at least weekly	-0.059 [0.037] N=1,253	-0.045 [0.042] N=1,124	-0.042 [0.045] N=963	**-0.090* [0.047] N=779**	-0.046 [0.058] N=593
Sweets consumed at least weekly	0.009 [0.066] N=1,253	0.021 [0.070] N=1,124	0.004 [0.079] N=963	0.007 [0.092] N=779	-0.122 [0.117] N=593
Snack foods consumed at least weekly	0.001 [0.086] N=1,253	0.014 [0.092] N=1,124	0.022 [0.104] N=963	0.074 [0.122] N=779	-0.009 [0.161] N=593
Packaged soft drinks consumed at least weekly	-0.039 [0.077] N=1,253	-0.044 [0.083] N=1,124	0.018 [0.094] N=963	0.043 [0.114] N=779	0.006 [0.144] N=593
Grains consumed at least weekly	-0.008 [0.039] N=1,253	-0.007 [0.042] N=1,124	-0.016 [0.048] N=963	-0.027 [0.059] N=779	-0.065 [0.075] N=593
Child was ever breastfed	0.025 [0.043] N=1,254	0.032 [0.045] N=1,125	0.039 [0.050] N=964	0.052 [0.055] N=780	0.032 [0.070] N=594

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,360	-0.014 [0.027] N=1,441	-0.014 [0.029] N=1,312	-0.019 [0.032] N=1,151	-0.020 [0.034] N=967	-0.013 [0.040] N=781
Vegetables consumed at least weekly	0.903 P=1,299	0.005 [0.033] N=1,441	0.017 [0.035] N=1,312	0.021 [0.037] N=1,151	0.013 [0.042] N=967	0.013 [0.048] N=781
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.071* [0.040] N=1,441**	**0.078* [0.042] N=1,312**	**0.078* [0.045] N=1,151**	**0.109** [0.045] N=967**	**0.105* [0.053] N=781**
Legumes consumed at least weekly	0.797 P=1,154	-0.047 [0.046] N=1,441	-0.029 [0.049] N=1,312	-0.037 [0.054] N=1,151	-0.054 [0.058] N=967	-0.024 [0.069] N=781
Animal products consumed at least weekly	0.915 P=1,319	-0.005 [0.033] N=1,441	0.005 [0.036] N=1,312	0.013 [0.038] N=1,151	-0.006 [0.041] N=967	0.026 [0.047] N=781
Sweets consumed at least weekly	0.706 P=1,020	0.007 [0.050] N=1,441	0.011 [0.052] N=1,312	0.002 [0.056] N=1,151	0.003 [0.060] N=967	-0.045 [0.069] N=781
Snack foods consumed at least weekly	0.572 P=822	-0.013 [0.058] N=1,441	-0.007 [0.061] N=1,312	-0.004 [0.064] N=1,151	0.011 [0.069] N=967	-0.034 [0.079] N=781
Packaged soft drinks consumed at least weekly	0.338 P=483	0.001 [0.049] N=1,441	-0.002 [0.051] N=1,312	0.033 [0.054] N=1,151	0.047 [0.059] N=967	0.027 [0.067] N=781
Grains consumed at least weekly	0.919 P=1,323	0.025 [0.038] N=1,441	0.032 [0.040] N=1,312	0.037 [0.043] N=1,151	0.043 [0.049] N=967	0.050 [0.055] N=781
Child was ever breastfed	0.963 P=1,388	-0.007 [0.023] N=1,442	-0.005 [0.023] N=1,313	-0.003 [0.024] N=1,152	-0.001 [0.024] N=968	-0.020 [0.027] N=782

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.047 [0.034] N=1,253	-0.053 [0.036] N=1,124	**-0.074* [0.042] N=963**	-0.072 [0.046] N=779	-0.076 [0.068] N=593
Vegetables consumed at least weekly	-0.051 [0.038] N=1,253	-0.041 [0.043] N=1,124	-0.048 [0.046] N=963	-0.062 [0.052] N=779	-0.081 [0.067] N=593
Meat fish or poultry consumed at least weekly	0.022 [0.054] N=1,253	0.022 [0.058] N=1,124	-0.003 [0.063] N=963	0.052 [0.070] N=779	0.016 [0.087] N=593
Legumes consumed at least weekly	-0.067 [0.052] N=1,253	-0.045 [0.057] N=1,124	-0.067 [0.069] N=963	-0.083 [0.074] N=779	-0.011 [0.091] N=593
Animal products consumed at least weekly	-0.061 [0.037] N=1,253	-0.046 [0.042] N=1,124	-0.042 [0.044] N=963	**-0.094** [0.047] N=779**	-0.047 [0.058] N=593
Sweets consumed at least weekly	0.005 [0.066] N=1,253	0.017 [0.070] N=1,124	0.001 [0.078] N=963	-0.002 [0.092] N=779	-0.130 [0.117] N=593
Snack foods consumed at least weekly	0.006 [0.087] N=1,253	0.016 [0.093] N=1,124	0.017 [0.105] N=963	0.065 [0.123] N=779	-0.017 [0.163] N=593
Packaged soft drinks consumed at least weekly	-0.048 [0.079] N=1,253	-0.057 [0.084] N=1,124	-0.005 [0.096] N=963	0.013 [0.116] N=779	-0.019 [0.146] N=593
Grains consumed at least weekly	-0.011 [0.038] N=1,253	-0.009 [0.041] N=1,124	-0.014 [0.047] N=963	-0.030 [0.058] N=779	-0.067 [0.073] N=593
Child was ever breastfed	0.025 [0.043] N=1,254	0.031 [0.044] N=1,125	0.041 [0.051] N=964	0.058 [0.055] N=780	0.038 [0.071] N=594

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.946 P=1,360	-0.014 [0.027] N=1,441	-0.013 [0.029] N=1,312	-0.018 [0.032] N=1,151	-0.023 [0.033] N=967	-0.018 [0.039] N=781
Vegetables consumed at least weekly	0.903 P=1,299	0.004 [0.034] N=1,441	0.018 [0.036] N=1,312	0.021 [0.038] N=1,151	0.010 [0.042] N=967	0.012 [0.048] N=781
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.073* [0.042] N=1,441**	**0.080* [0.043] N=1,312**	**0.077* [0.046] N=1,151**	**0.101** [0.047] N=967**	0.091 [0.055] N=781
Legumes consumed at least weekly	0.797 P=1,154	-0.042 [0.046] N=1,441	-0.023 [0.049] N=1,312	-0.031 [0.053] N=1,151	-0.050 [0.057] N=967	-0.020 [0.068] N=781
Animal products consumed at least weekly	0.915 P=1,319	0.004 [0.033] N=1,441	0.013 [0.036] N=1,312	0.022 [0.038] N=1,151	-0.001 [0.041] N=967	0.030 [0.047] N=781
Sweets consumed at least weekly	0.706 P=1,020	0.014 [0.051] N=1,441	0.017 [0.053] N=1,312	0.005 [0.058] N=1,151	0.004 [0.062] N=967	-0.045 [0.072] N=781
Snack foods consumed at least weekly	0.572 P=822	-0.005 [0.057] N=1,441	-0.001 [0.060] N=1,312	0.002 [0.064] N=1,151	0.017 [0.068] N=967	-0.024 [0.079] N=781
Packaged soft drinks consumed at least weekly	0.338 P=483	0.012 [0.048] N=1,441	0.005 [0.050] N=1,312	0.040 [0.053] N=1,151	0.054 [0.058] N=967	0.036 [0.067] N=781
Grains consumed at least weekly	0.919 P=1,323	0.025 [0.038] N=1,441	0.031 [0.039] N=1,312	0.034 [0.043] N=1,151	0.040 [0.048] N=967	0.048 [0.054] N=781
Child was ever breastfed	0.963 P=1,388	-0.006 [0.023] N=1,442	-0.004 [0.023] N=1,313	-0.003 [0.024] N=1,152	-0.002 [0.024] N=968	-0.021 [0.029] N=782

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.049 [0.033] N=1,253	-0.056 [0.035] N=1,124	**-0.076* [0.042] N=963**	-0.078 [0.047] N=779	-0.091 [0.067] N=593
Vegetables consumed at least weekly	-0.053 [0.039] N=1,253	-0.041 [0.043] N=1,124	-0.048 [0.047] N=963	-0.062 [0.053] N=779	-0.082 [0.066] N=593
Meat fish or poultry consumed at least weekly	0.021 [0.055] N=1,253	0.020 [0.059] N=1,124	-0.009 [0.064] N=963	0.034 [0.072] N=779	-0.028 [0.090] N=593
Legumes consumed at least weekly	-0.064 [0.052] N=1,253	-0.040 [0.058] N=1,124	-0.055 [0.069] N=963	-0.072 [0.074] N=779	0.002 [0.090] N=593
Animal products consumed at least weekly	-0.052 [0.035] N=1,253	-0.038 [0.040] N=1,124	-0.032 [0.042] N=963	**-0.087* [0.046] N=779**	-0.047 [0.055] N=593
Sweets consumed at least weekly	0.016 [0.066] N=1,253	0.025 [0.070] N=1,124	0.005 [0.078] N=963	0.001 [0.092] N=779	-0.135 [0.114] N=593
Snack foods consumed at least weekly	0.011 [0.084] N=1,253	0.018 [0.091] N=1,124	0.021 [0.101] N=963	0.066 [0.117] N=779	-0.013 [0.154] N=593
Packaged soft drinks consumed at least weekly	-0.027 [0.077] N=1,253	-0.042 [0.083] N=1,124	0.012 [0.094] N=963	0.033 [0.112] N=779	0.008 [0.145] N=593
Grains consumed at least weekly	-0.011 [0.038] N=1,253	-0.011 [0.041] N=1,124	-0.020 [0.046] N=963	-0.035 [0.057] N=779	-0.076 [0.073] N=593
Child was ever breastfed	0.025 [0.043] N=1,254	0.033 [0.044] N=1,125	0.040 [0.050] N=964	0.054 [0.055] N=780	0.034 [0.072] N=594

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,266	-0.010 [0.029] N=1,341	-0.009 [0.031] N=1,219	-0.008 [0.035] N=1,073	-0.017 [0.036] N=902	-0.013 [0.042] N=731
Vegetables consumed at least weekly	0.905 P=1,212	0.010 [0.033] N=1,341	0.023 [0.036] N=1,219	0.032 [0.039] N=1,073	0.015 [0.043] N=902	-0.002 [0.049] N=731
Meat fish or poultry consumed at least weekly	0.783 P=1,059	**0.079* [0.044] N=1,341**	**0.083* [0.046] N=1,219**	**0.085* [0.050] N=1,073**	**0.091* [0.052] N=902**	0.082 [0.060] N=731
Legumes consumed at least weekly	0.794 P=1,072	-0.040 [0.048] N=1,341	-0.019 [0.051] N=1,219	-0.024 [0.056] N=1,073	-0.053 [0.061] N=902	-0.030 [0.074] N=731
Animal products consumed at least weekly	0.916 P=1,230	0.001 [0.035] N=1,341	0.016 [0.038] N=1,219	0.028 [0.039] N=1,073	0.006 [0.042] N=902	0.037 [0.047] N=731
Sweets consumed at least weekly	0.716 P=961	0.008 [0.052] N=1,341	0.012 [0.054] N=1,219	-0.007 [0.059] N=1,073	-0.011 [0.064] N=902	-0.056 [0.075] N=731
Snack foods consumed at least weekly	0.580 P=774	-0.005 [0.059] N=1,341	-0.003 [0.062] N=1,219	-0.010 [0.066] N=1,073	0.007 [0.070] N=902	-0.028 [0.082] N=731
Packaged soft drinks consumed at least weekly	0.349 P=465	-0.010 [0.048] N=1,341	-0.017 [0.050] N=1,219	0.018 [0.053] N=1,073	0.033 [0.060] N=902	0.013 [0.069] N=731
Grains consumed at least weekly	0.919 P=1,232	0.038 [0.039] N=1,341	0.046 [0.041] N=1,219	0.048 [0.044] N=1,073	0.052 [0.050] N=902	0.055 [0.056] N=731
Child was ever breastfed	0.966 P=1,296	-0.013 [0.021] N=1,342	-0.014 [0.021] N=1,220	-0.015 [0.022] N=1,074	-0.017 [0.022] N=903	-0.033 [0.028] N=732

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.040 [0.037] N=1,163	-0.045 [0.040] N=1,041	-0.051 [0.049] N=895	-0.062 [0.050] N=724	-0.072 [0.069] N=553
Vegetables consumed at least weekly	-0.042 [0.038] N=1,163	-0.031 [0.043] N=1,041	-0.024 [0.048] N=895	-0.045 [0.055] N=724	**-0.104* [0.062] N=553**
Meat fish or poultry consumed at least weekly	0.043 [0.065] N=1,163	0.041 [0.069] N=1,041	0.027 [0.077] N=895	0.037 [0.082] N=724	-0.013 [0.101] N=553
Legumes consumed at least weekly	-0.051 [0.054] N=1,163	-0.022 [0.058] N=1,041	-0.026 [0.070] N=895	-0.053 [0.079] N=724	0.015 [0.095] N=553
Animal products consumed at least weekly	-0.056 [0.041] N=1,163	-0.032 [0.045] N=1,041	-0.018 [0.047] N=895	-0.067 [0.050] N=724	-0.018 [0.059] N=553
Sweets consumed at least weekly	0.008 [0.069] N=1,163	0.018 [0.073] N=1,041	-0.022 [0.082] N=895	-0.036 [0.094] N=724	-0.171 [0.123] N=553
Snack foods consumed at least weekly	0.018 [0.089] N=1,163	0.025 [0.096] N=1,041	0.007 [0.110] N=895	0.057 [0.126] N=724	0.007 [0.166] N=553
Packaged soft drinks consumed at least weekly	-0.057 [0.077] N=1,163	-0.071 [0.083] N=1,041	-0.020 [0.095] N=895	-0.002 [0.117] N=724	-0.064 [0.151] N=553
Grains consumed at least weekly	0.014 [0.041] N=1,163	0.019 [0.044] N=1,041	0.014 [0.051] N=895	-0.002 [0.062] N=724	-0.045 [0.076] N=553
Child was ever breastfed	0.018 [0.040] N=1,164	0.020 [0.041] N=1,042	0.022 [0.045] N=896	0.031 [0.047] N=725	0.006 [0.066] N=554

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,266	-0.010 [0.029] N=1,341	-0.008 [0.031] N=1,219	-0.008 [0.035] N=1,073	-0.018 [0.036] N=902	-0.015 [0.041] N=731
Vegetables consumed at least weekly	0.905 P=1,212	0.009 [0.033] N=1,341	0.021 [0.036] N=1,219	0.030 [0.039] N=1,073	0.012 [0.043] N=902	-0.003 [0.050] N=731
Meat fish or poultry consumed at least weekly	0.783 P=1,059	**0.081* [0.044] N=1,341**	**0.086* [0.046] N=1,219**	**0.086* [0.049] N=1,073**	**0.094* [0.050] N=902**	0.083 [0.059] N=731
Legumes consumed at least weekly	0.794 P=1,072	-0.042 [0.048] N=1,341	-0.020 [0.051] N=1,219	-0.026 [0.056] N=1,073	-0.056 [0.061] N=902	-0.034 [0.074] N=731
Animal products consumed at least weekly	0.916 P=1,230	0.000 [0.035] N=1,341	0.016 [0.038] N=1,219	0.026 [0.039] N=1,073	0.004 [0.042] N=902	0.035 [0.047] N=731
Sweets consumed at least weekly	0.716 P=961	0.007 [0.052] N=1,341	0.012 [0.054] N=1,219	-0.007 [0.059] N=1,073	-0.011 [0.065] N=902	-0.056 [0.076] N=731
Snack foods consumed at least weekly	0.580 P=774	-0.004 [0.058] N=1,341	-0.002 [0.061] N=1,219	-0.008 [0.065] N=1,073	0.007 [0.070] N=902	-0.028 [0.082] N=731
Packaged soft drinks consumed at least weekly	0.349 P=465	-0.010 [0.048] N=1,341	-0.017 [0.050] N=1,219	0.018 [0.053] N=1,073	0.033 [0.060] N=902	0.011 [0.070] N=731
Grains consumed at least weekly	0.919 P=1,232	0.037 [0.039] N=1,341	0.045 [0.041] N=1,219	0.046 [0.044] N=1,073	0.051 [0.050] N=902	0.053 [0.056] N=731
Child was ever breastfed	0.966 P=1,296	-0.011 [0.021] N=1,342	-0.012 [0.022] N=1,220	-0.011 [0.023] N=1,074	-0.012 [0.023] N=903	-0.028 [0.028] N=732

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.040 [0.037] N=1,163	-0.044 [0.041] N=1,041	-0.051 [0.049] N=895	-0.064 [0.052] N=724	-0.076 [0.072] N=553
Vegetables consumed at least weekly	-0.043 [0.038] N=1,163	-0.033 [0.043] N=1,041	-0.027 [0.048] N=895	-0.050 [0.056] N=724	**-0.109* [0.063] N=553**
Meat fish or poultry consumed at least weekly	0.045 [0.064] N=1,163	0.045 [0.069] N=1,041	0.027 [0.076] N=895	0.038 [0.082] N=724	-0.012 [0.101] N=553
Legumes consumed at least weekly	-0.053 [0.054] N=1,163	-0.021 [0.058] N=1,041	-0.028 [0.070] N=895	-0.061 [0.079] N=724	0.010 [0.093] N=553
Animal products consumed at least weekly	-0.056 [0.041] N=1,163	-0.032 [0.046] N=1,041	-0.019 [0.048] N=895	-0.070 [0.050] N=724	-0.025 [0.060] N=553
Sweets consumed at least weekly	0.010 [0.070] N=1,163	0.021 [0.074] N=1,041	-0.015 [0.083] N=895	-0.026 [0.095] N=724	-0.160 [0.123] N=553
Snack foods consumed at least weekly	0.018 [0.089] N=1,163	0.024 [0.096] N=1,041	0.010 [0.109] N=895	0.062 [0.125] N=724	0.002 [0.166] N=553
Packaged soft drinks consumed at least weekly	-0.057 [0.077] N=1,163	-0.072 [0.083] N=1,041	-0.016 [0.094] N=895	0.005 [0.115] N=724	-0.056 [0.150] N=553
Grains consumed at least weekly	0.014 [0.041] N=1,163	0.018 [0.044] N=1,041	0.012 [0.051] N=895	-0.005 [0.062] N=724	-0.052 [0.076] N=553
Child was ever breastfed	0.020 [0.041] N=1,164	0.019 [0.042] N=1,042	0.028 [0.047] N=896	0.037 [0.050] N=725	0.011 [0.068] N=554

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.