Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Oldest Child

Age reference: 12 November 2022. Any-child household eligibility; one oldest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.326	0.241 [0.238] N=1,417	0.275 [0.255] N=1,313	0.263 [0.284] N=1,181	0.334 [0.279] N=1,014	0.080 [0.301] N=835
WHO weight-for-age z score	0.118	-0.078 [0.130] N=1,608	-0.032 [0.137] N=1,485	-0.043 [0.144] N=1,334	-0.075 [0.157] N=1,139	-0.252 [0.175] N=933
WHO weight-for-length/height z score, 0-60 months	0.375	0.044 [0.226] N=1,395	-0.017 [0.238] N=1,299	-0.107 [0.272] N=1,170	-0.175 [0.303] N=1,003	-0.121 [0.342] N=823
WHO BMI-for-age z score	0.350	0.129 [0.245] N=1,360	0.148 [0.263] N=1,267	0.042 [0.300] N=1,138	-0.120 [0.338] N=975	-0.005 [0.379] N=803
Stunted: HAZ below -2	0.231 P=311	-0.044 [0.043] N=1,417	-0.059 [0.046] N=1,313	-0.035 [0.049] N=1,181	-0.038 [0.052] N=1,014	-0.004 [0.060] N=835
Underweight: WAZ below -2	0.067 P=110	0.008 [0.021] N=1,608	0.001 [0.021] N=1,485	-0.003 [0.023] N=1,334	0.010 [0.026] N=1,139	0.037 [0.032] N=933
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=160	-0.032 [0.036] N=1,395	-0.023 [0.035] N=1,299	-0.023 [0.038] N=1,170	-0.011 [0.042] N=1,003	-0.029 [0.044] N=823
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=267	0.001 [0.045] N=1,395	-0.007 [0.048] N=1,299	-0.017 [0.054] N=1,170	-0.019 [0.061] N=1,003	-0.019 [0.069] N=823
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=124	0.005 [0.028] N=1,395	-0.005 [0.030] N=1,299	-0.005 [0.033] N=1,170	-0.004 [0.036] N=1,003	0.008 [0.039] N=823
CIAF: any stunting, underweight, or wasting, 0-60 months	0.316 P=415	-0.075 [0.047] N=1,351	**-0.084* [0.047] N=1,257**	-0.065 [0.050] N=1,131	-0.058 [0.052] N=970	-0.031 [0.059] N=798
BMI-for-age z score below -2	0.120 P=168	-0.030 [0.037] N=1,360	-0.030 [0.038] N=1,267	-0.018 [0.041] N=1,138	0.017 [0.045] N=975	0.002 [0.049] N=803

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.222 [0.354] N=1,199	0.272 [0.400] N=1,095	0.323 [0.476] N=963	0.626 [0.506] N=796	-0.079 [0.677] N=617
WHO weight-for-age z score	-0.163 [0.180] N=1,370	-0.110 [0.195] N=1,247	-0.140 [0.210] N=1,096	-0.206 [0.270] N=901	**-0.849** [0.349] N=695**
WHO weight-for-length/height z score, 0-60 months	0.111 [0.257] N=1,178	0.019 [0.281] N=1,082	-0.209 [0.357] N=953	-0.416 [0.411] N=786	-0.424 [0.510] N=606
WHO BMI-for-age z score	0.265 [0.272] N=1,150	0.336 [0.306] N=1,057	0.106 [0.387] N=928	-0.283 [0.450] N=765	-0.061 [0.545] N=593
Stunted: HAZ below -2	-0.024 [0.058] N=1,199	-0.048 [0.065] N=1,095	-0.011 [0.070] N=963	-0.029 [0.078] N=796	0.076 [0.107] N=617
Underweight: WAZ below -2	0.004 [0.030] N=1,370	-0.008 [0.032] N=1,247	-0.019 [0.036] N=1,096	0.005 [0.045] N=901	0.103 [0.064] N=695
Wasted: WLZ/WHZ below -2, 0-60 months	-0.037 [0.050] N=1,178	-0.021 [0.051] N=1,082	-0.019 [0.057] N=953	0.004 [0.072] N=786	-0.044 [0.087] N=606
Overweight: WLZ/WHZ above +2, 0-60 months	-0.002 [0.052] N=1,178	-0.018 [0.056] N=1,082	-0.048 [0.066] N=953	-0.066 [0.082] N=786	-0.106 [0.095] N=606
Obese: WLZ/WHZ above +3, 0-60 months	0.004 [0.036] N=1,178	-0.014 [0.040] N=1,082	-0.015 [0.047] N=953	-0.018 [0.058] N=786	0.003 [0.069] N=606
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.053 [0.071] N=1,142	-0.064 [0.074] N=1,048	-0.028 [0.082] N=922	-0.022 [0.098] N=761	0.079 [0.135] N=589
BMI-for-age z score below -2	-0.057 [0.049] N=1,150	-0.061 [0.051] N=1,057	-0.043 [0.058] N=928	0.024 [0.075] N=765	-0.025 [0.093] N=593

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.326	0.241 [0.238] N=1,417	0.276 [0.255] N=1,313	0.264 [0.284] N=1,181	0.331 [0.279] N=1,014	0.079 [0.301] N=835
WHO weight-for-age z score	0.118	-0.078 [0.129] N=1,608	-0.031 [0.136] N=1,485	-0.039 [0.143] N=1,334	-0.072 [0.157] N=1,139	-0.249 [0.173] N=933
WHO weight-for-length/height z score, 0-60 months	0.375	0.041 [0.225] N=1,395	-0.019 [0.238] N=1,299	-0.106 [0.272] N=1,170	-0.175 [0.302] N=1,003	-0.121 [0.341] N=823
WHO BMI-for-age z score	0.350	0.128 [0.245] N=1,360	0.146 [0.263] N=1,267	0.043 [0.300] N=1,138	-0.119 [0.338] N=975	-0.007 [0.377] N=803
Stunted: HAZ below -2	0.231 P=311	-0.044 [0.043] N=1,417	-0.059 [0.046] N=1,313	-0.036 [0.048] N=1,181	-0.038 [0.052] N=1,014	-0.004 [0.060] N=835
Underweight: WAZ below -2	0.067 P=110	0.008 [0.020] N=1,608	0.000 [0.021] N=1,485	-0.004 [0.023] N=1,334	0.009 [0.026] N=1,139	0.036 [0.032] N=933
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=160	-0.031 [0.036] N=1,395	-0.022 [0.035] N=1,299	-0.023 [0.037] N=1,170	-0.011 [0.041] N=1,003	-0.029 [0.044] N=823
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=267	0.001 [0.045] N=1,395	-0.008 [0.048] N=1,299	-0.017 [0.054] N=1,170	-0.019 [0.061] N=1,003	-0.019 [0.069] N=823
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=124	0.005 [0.028] N=1,395	-0.005 [0.030] N=1,299	-0.005 [0.032] N=1,170	-0.004 [0.036] N=1,003	0.008 [0.039] N=823
CIAF: any stunting, underweight, or wasting, 0-60 months	0.316 P=415	-0.074 [0.047] N=1,351	**-0.084* [0.047] N=1,257**	-0.067 [0.049] N=1,131	-0.059 [0.052] N=970	-0.031 [0.058] N=798
BMI-for-age z score below -2	0.120 P=168	-0.030 [0.037] N=1,360	-0.029 [0.038] N=1,267	-0.019 [0.041] N=1,138	0.017 [0.045] N=975	0.003 [0.048] N=803

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.218 [0.353] N=1,199	0.264 [0.398] N=1,095	0.310 [0.472] N=963	0.591 [0.501] N=796	-0.124 [0.668] N=617
WHO weight-for-age z score	-0.158 [0.178] N=1,370	-0.097 [0.193] N=1,247	-0.123 [0.208] N=1,096	-0.189 [0.269] N=901	**-0.817** [0.348] N=695**
WHO weight-for-length/height z score, 0-60 months	0.109 [0.257] N=1,178	0.019 [0.280] N=1,082	-0.203 [0.355] N=953	-0.410 [0.410] N=786	-0.412 [0.512] N=606
WHO BMI-for-age z score	0.267 [0.272] N=1,150	0.340 [0.306] N=1,057	0.119 [0.385] N=928	-0.271 [0.450] N=765	-0.047 [0.549] N=593
Stunted: HAZ below -2	-0.024 [0.058] N=1,199	-0.048 [0.065] N=1,095	-0.012 [0.070] N=963	-0.027 [0.078] N=796	0.078 [0.107] N=617
Underweight: WAZ below -2	0.003 [0.029] N=1,370	-0.010 [0.031] N=1,247	-0.022 [0.035] N=1,096	0.002 [0.044] N=901	0.099 [0.064] N=695
Wasted: WLZ/WHZ below -2, 0-60 months	-0.036 [0.050] N=1,178	-0.021 [0.051] N=1,082	-0.022 [0.058] N=953	0.003 [0.073] N=786	-0.047 [0.088] N=606
Overweight: WLZ/WHZ above +2, 0-60 months	-0.002 [0.052] N=1,178	-0.018 [0.056] N=1,082	-0.049 [0.067] N=953	-0.066 [0.082] N=786	-0.106 [0.095] N=606
Obese: WLZ/WHZ above +3, 0-60 months	0.004 [0.036] N=1,178	-0.014 [0.040] N=1,082	-0.016 [0.047] N=953	-0.018 [0.058] N=786	0.002 [0.069] N=606
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.054 [0.071] N=1,142	-0.067 [0.074] N=1,048	-0.035 [0.082] N=922	-0.027 [0.098] N=761	0.074 [0.136] N=589
BMI-for-age z score below -2	-0.057 [0.049] N=1,150	-0.062 [0.051] N=1,057	-0.047 [0.059] N=928	0.021 [0.076] N=765	-0.028 [0.095] N=593

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.326	0.208 [0.238] N=1,417	0.220 [0.255] N=1,313	0.224 [0.283] N=1,181	0.300 [0.276] N=1,014	0.014 [0.295] N=835
WHO weight-for-age z score	0.118	-0.128 [0.123] N=1,608	-0.084 [0.130] N=1,485	-0.089 [0.138] N=1,334	-0.114 [0.151] N=1,139	**-0.299* [0.164] N=933**
WHO weight-for-length/height z score, 0-60 months	0.375	0.018 [0.227] N=1,395	-0.034 [0.240] N=1,299	-0.116 [0.272] N=1,170	-0.182 [0.299] N=1,003	-0.139 [0.337] N=823
WHO BMI-for-age z score	0.350	0.116 [0.246] N=1,360	0.139 [0.265] N=1,267	0.037 [0.299] N=1,138	-0.122 [0.333] N=975	-0.015 [0.374] N=803
Stunted: HAZ below -2	0.231 P=311	-0.035 [0.042] N=1,417	-0.043 [0.045] N=1,313	-0.025 [0.047] N=1,181	-0.030 [0.050] N=1,014	0.011 [0.058] N=835
Underweight: WAZ below -2	0.067 P=110	0.015 [0.019] N=1,608	0.011 [0.020] N=1,485	0.006 [0.022] N=1,334	0.019 [0.024] N=1,139	0.047 [0.030] N=933
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=160	-0.025 [0.035] N=1,395	-0.014 [0.035] N=1,299	-0.016 [0.037] N=1,170	-0.007 [0.041] N=1,003	-0.025 [0.044] N=823
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=267	0.002 [0.045] N=1,395	-0.004 [0.048] N=1,299	-0.014 [0.054] N=1,170	-0.016 [0.061] N=1,003	-0.018 [0.069] N=823
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=124	0.006 [0.029] N=1,395	-0.001 [0.030] N=1,299	-0.002 [0.033] N=1,170	-0.003 [0.036] N=1,003	0.008 [0.039] N=823
CIAF: any stunting, underweight, or wasting, 0-60 months	0.316 P=415	-0.055 [0.046] N=1,351	-0.056 [0.046] N=1,257	-0.045 [0.048] N=1,131	-0.043 [0.051] N=970	-0.007 [0.058] N=798
BMI-for-age z score below -2	0.120 P=168	-0.024 [0.037] N=1,360	-0.021 [0.038] N=1,267	-0.011 [0.041] N=1,138	0.020 [0.045] N=975	0.005 [0.048] N=803

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.191 [0.355] N=1,199	0.193 [0.400] N=1,095	0.262 [0.475] N=963	0.580 [0.499] N=796	-0.172 [0.659] N=617
WHO weight-for-age z score	-0.195 [0.172] N=1,370	-0.143 [0.188] N=1,247	-0.156 [0.204] N=1,096	-0.204 [0.269] N=901	**-0.839** [0.346] N=695**
WHO weight-for-length/height z score, 0-60 months	0.101 [0.262] N=1,178	0.026 [0.289] N=1,082	-0.185 [0.362] N=953	-0.421 [0.413] N=786	-0.443 [0.516] N=606
WHO BMI-for-age z score	0.265 [0.276] N=1,150	0.351 [0.313] N=1,057	0.130 [0.389] N=928	-0.295 [0.452] N=765	-0.070 [0.556] N=593
Stunted: HAZ below -2	-0.019 [0.058] N=1,199	-0.032 [0.065] N=1,095	-0.003 [0.071] N=963	-0.033 [0.077] N=796	0.082 [0.104] N=617
Underweight: WAZ below -2	0.009 [0.028] N=1,370	0.001 [0.030] N=1,247	-0.014 [0.034] N=1,096	0.005 [0.044] N=901	0.099 [0.064] N=695
Wasted: WLZ/WHZ below -2, 0-60 months	-0.030 [0.049] N=1,178	-0.012 [0.051] N=1,082	-0.017 [0.058] N=953	0.009 [0.075] N=786	-0.036 [0.090] N=606
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.052] N=1,178	-0.009 [0.058] N=1,082	-0.043 [0.068] N=953	-0.063 [0.081] N=786	-0.099 [0.092] N=606
Obese: WLZ/WHZ above +3, 0-60 months	0.006 [0.037] N=1,178	-0.008 [0.040] N=1,082	-0.013 [0.048] N=953	-0.020 [0.058] N=786	-0.002 [0.069] N=606
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.039 [0.069] N=1,142	-0.038 [0.073] N=1,048	-0.020 [0.080] N=922	-0.025 [0.097] N=761	0.095 [0.126] N=589
BMI-for-age z score below -2	-0.050 [0.049] N=1,150	-0.050 [0.051] N=1,057	-0.039 [0.059] N=928	0.028 [0.077] N=765	-0.019 [0.096] N=593

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.326	0.139 [0.236] N=1,417	0.156 [0.254] N=1,313	0.166 [0.284] N=1,181	0.221 [0.278] N=1,014	-0.090 [0.295] N=835
WHO weight-for-age z score	0.118	-0.145 [0.125] N=1,608	-0.104 [0.132] N=1,485	-0.105 [0.140] N=1,334	-0.144 [0.156] N=1,139	**-0.340** [0.170] N=933**
WHO weight-for-length/height z score, 0-60 months	0.375	0.049 [0.223] N=1,395	-0.009 [0.236] N=1,299	-0.104 [0.270] N=1,170	-0.168 [0.302] N=1,003	-0.116 [0.340] N=823
WHO BMI-for-age z score	0.350	0.154 [0.244] N=1,360	0.177 [0.262] N=1,267	0.065 [0.299] N=1,138	-0.092 [0.337] N=975	0.023 [0.376] N=803
Stunted: HAZ below -2	0.231 P=311	-0.020 [0.042] N=1,417	-0.028 [0.046] N=1,313	-0.009 [0.048] N=1,181	-0.009 [0.050] N=1,014	0.033 [0.058] N=835
Underweight: WAZ below -2	0.067 P=110	0.018 [0.020] N=1,608	0.013 [0.021] N=1,485	0.009 [0.022] N=1,334	0.022 [0.026] N=1,139	0.051 [0.031] N=933
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=160	-0.027 [0.035] N=1,395	-0.017 [0.036] N=1,299	-0.017 [0.038] N=1,170	-0.008 [0.042] N=1,003	-0.030 [0.044] N=823
Overweight: WLZ/WHZ above +2, 0-60 months	0.199 P=267	0.008 [0.045] N=1,395	0.001 [0.048] N=1,299	-0.012 [0.054] N=1,170	-0.013 [0.061] N=1,003	-0.018 [0.070] N=823
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=124	0.008 [0.029] N=1,395	-0.001 [0.030] N=1,299	-0.004 [0.033] N=1,170	-0.003 [0.037] N=1,003	0.009 [0.039] N=823
CIAF: any stunting, underweight, or wasting, 0-60 months	0.316 P=415	-0.042 [0.047] N=1,351	-0.043 [0.048] N=1,257	-0.030 [0.051] N=1,131	-0.020 [0.054] N=970	0.012 [0.060] N=798
BMI-for-age z score below -2	0.120 P=168	-0.026 [0.037] N=1,360	-0.025 [0.038] N=1,267	-0.013 [0.042] N=1,138	0.020 [0.046] N=975	0.001 [0.049] N=803

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.155 [0.354] N=1,199	0.179 [0.397] N=1,095	0.264 [0.475] N=963	0.562 [0.494] N=796	-0.241 [0.638] N=617
WHO weight-for-age z score	-0.224 [0.175] N=1,370	-0.170 [0.190] N=1,247	-0.186 [0.206] N=1,096	-0.242 [0.273] N=901	**-0.864** [0.349] N=695**
WHO weight-for-length/height z score, 0-60 months	0.099 [0.258] N=1,178	0.022 [0.281] N=1,082	-0.195 [0.357] N=953	-0.391 [0.413] N=786	-0.360 [0.521] N=606
WHO BMI-for-age z score	0.259 [0.274] N=1,150	0.350 [0.308] N=1,057	0.133 [0.388] N=928	-0.252 [0.451] N=765	0.008 [0.559] N=593
Stunted: HAZ below -2	-0.013 [0.059] N=1,199	-0.029 [0.066] N=1,095	-0.005 [0.071] N=963	-0.031 [0.078] N=796	0.086 [0.105] N=617
Underweight: WAZ below -2	0.013 [0.029] N=1,370	0.003 [0.031] N=1,247	-0.011 [0.035] N=1,096	0.010 [0.047] N=901	0.108 [0.067] N=695
Wasted: WLZ/WHZ below -2, 0-60 months	-0.028 [0.050] N=1,178	-0.014 [0.052] N=1,082	-0.018 [0.058] N=953	0.007 [0.075] N=786	-0.039 [0.091] N=606
Overweight: WLZ/WHZ above +2, 0-60 months	0.004 [0.052] N=1,178	-0.008 [0.056] N=1,082	-0.042 [0.066] N=953	-0.056 [0.080] N=786	-0.082 [0.092] N=606
Obese: WLZ/WHZ above +3, 0-60 months	0.011 [0.037] N=1,178	-0.006 [0.040] N=1,082	-0.010 [0.048] N=953	-0.012 [0.058] N=786	0.011 [0.069] N=606
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.027 [0.070] N=1,142	-0.033 [0.074] N=1,048	-0.016 [0.082] N=922	-0.017 [0.099] N=761	0.112 [0.131] N=589
BMI-for-age z score below -2	-0.049 [0.049] N=1,150	-0.057 [0.051] N=1,057	-0.045 [0.060] N=928	0.022 [0.077] N=765	-0.024 [0.096] N=593

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.327	0.253 [0.244] N=1,320	0.257 [0.267] N=1,230	0.251 [0.292] N=1,104	0.293 [0.282] N=954	0.053 [0.293] N=784
WHO weight-for-age z score	0.122	-0.022 [0.134] N=1,498	0.027 [0.138] N=1,390	0.019 [0.146] N=1,247	-0.006 [0.158] N=1,070	-0.135 [0.186] N=877
WHO weight-for-length/height z score, 0-60 months	0.382	0.172 [0.227] N=1,299	0.124 [0.238] N=1,217	0.020 [0.271] N=1,095	-0.028 [0.297] N=945	0.063 [0.336] N=774
WHO BMI-for-age z score	0.361	0.264 [0.248] N=1,266	0.269 [0.263] N=1,186	0.166 [0.298] N=1,065	0.055 [0.330] N=919	0.232 [0.365] N=755
Stunted: HAZ below -2	0.233 P=290	-0.032 [0.045] N=1,320	-0.043 [0.049] N=1,230	-0.021 [0.052] N=1,104	-0.025 [0.054] N=954	0.012 [0.062] N=784
Underweight: WAZ below -2	0.066 P=102	-0.001 [0.022] N=1,498	-0.007 [0.023] N=1,390	-0.012 [0.024] N=1,247	-0.001 [0.027] N=1,070	0.022 [0.036] N=877
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=149	-0.042 [0.037] N=1,299	-0.034 [0.036] N=1,217	-0.032 [0.038] N=1,095	-0.024 [0.042] N=945	-0.041 [0.044] N=774
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=248	0.014 [0.047] N=1,299	0.007 [0.050] N=1,217	-0.005 [0.057] N=1,095	-0.006 [0.065] N=945	-0.001 [0.075] N=774
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=119	0.006 [0.031] N=1,299	-0.001 [0.032] N=1,217	-0.001 [0.035] N=1,095	0.001 [0.038] N=945	0.018 [0.042] N=774
CIAF: any stunting, underweight, or wasting, 0-60 months	0.318 P=388	-0.066 [0.050] N=1,258	-0.074 [0.050] N=1,178	-0.060 [0.053] N=1,058	-0.062 [0.056] N=914	-0.034 [0.064] N=750
BMI-for-age z score below -2	0.120 P=156	-0.035 [0.038] N=1,266	-0.035 [0.038] N=1,186	-0.025 [0.041] N=1,065	0.004 [0.045] N=919	-0.012 [0.049] N=755

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.284 [0.352] N=1,116	0.306 [0.411] N=1,026	0.362 [0.486] N=900	0.635 [0.508] N=750	-0.036 [0.685] N=580
WHO weight-for-age z score	-0.172 [0.187] N=1,275	-0.124 [0.197] N=1,167	-0.163 [0.217] N=1,024	-0.238 [0.272] N=847	**-0.815** [0.388] N=654**
WHO weight-for-length/height z score, 0-60 months	0.115 [0.256] N=1,096	0.012 [0.272] N=1,014	-0.257 [0.342] N=892	-0.426 [0.410] N=742	-0.347 [0.513] N=571
WHO BMI-for-age z score	0.270 [0.273] N=1,070	0.278 [0.300] N=990	0.039 [0.375] N=869	-0.285 [0.453] N=723	0.047 [0.552] N=559
Stunted: HAZ below -2	-0.022 [0.061] N=1,116	-0.046 [0.068] N=1,026	-0.011 [0.074] N=900	-0.036 [0.083] N=750	0.071 [0.117] N=580
Underweight: WAZ below -2	0.000 [0.030] N=1,275	-0.010 [0.032] N=1,167	-0.020 [0.036] N=1,024	0.006 [0.043] N=847	0.104 [0.070] N=654
Wasted: WLZ/WHZ below -2, 0-60 months	-0.039 [0.050] N=1,096	-0.023 [0.050] N=1,014	-0.020 [0.056] N=892	-0.009 [0.073] N=742	-0.077 [0.086] N=571
Overweight: WLZ/WHZ above +2, 0-60 months	-0.000 [0.054] N=1,096	-0.021 [0.057] N=1,014	-0.059 [0.069] N=892	-0.084 [0.087] N=742	-0.105 [0.103] N=571
Obese: WLZ/WHZ above +3, 0-60 months	-0.005 [0.040] N=1,096	-0.018 [0.044] N=1,014	-0.024 [0.051] N=892	-0.029 [0.061] N=742	-0.003 [0.073] N=571
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.046 [0.074] N=1,063	-0.056 [0.076] N=983	-0.031 [0.085] N=863	-0.055 [0.099] N=719	0.030 [0.143] N=555
BMI-for-age z score below -2	-0.051 [0.048] N=1,070	-0.052 [0.048] N=990	-0.038 [0.056] N=869	0.018 [0.074] N=723	-0.042 [0.092] N=559

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.327	0.242 [0.244] N=1,320	0.247 [0.266] N=1,230	0.231 [0.293] N=1,104	0.262 [0.289] N=954	-0.008 [0.310] N=784
WHO weight-for-age z score	0.122	-0.029 [0.133] N=1,498	0.020 [0.137] N=1,390	0.007 [0.145] N=1,247	-0.023 [0.157] N=1,070	-0.167 [0.184] N=877
WHO weight-for-length/height z score, 0-60 months	0.382	0.160 [0.228] N=1,299	0.109 [0.239] N=1,217	0.007 [0.273] N=1,095	-0.040 [0.304] N=945	0.056 [0.348] N=774
WHO BMI-for-age z score	0.361	0.253 [0.248] N=1,266	0.255 [0.263] N=1,186	0.153 [0.300] N=1,065	0.038 [0.338] N=919	0.208 [0.380] N=755
Stunted: HAZ below -2	0.233 P=290	-0.030 [0.046] N=1,320	-0.042 [0.049] N=1,230	-0.019 [0.052] N=1,104	-0.021 [0.055] N=954	0.019 [0.063] N=784
Underweight: WAZ below -2	0.066 P=102	0.001 [0.021] N=1,498	-0.006 [0.022] N=1,390	-0.010 [0.024] N=1,247	0.003 [0.027] N=1,070	0.029 [0.035] N=877
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=149	-0.040 [0.037] N=1,299	-0.032 [0.036] N=1,217	-0.032 [0.039] N=1,095	-0.025 [0.044] N=945	-0.049 [0.047] N=774
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=248	0.014 [0.048] N=1,299	0.006 [0.050] N=1,217	-0.006 [0.057] N=1,095	-0.010 [0.065] N=945	-0.002 [0.075] N=774
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=119	0.005 [0.030] N=1,299	-0.001 [0.032] N=1,217	-0.001 [0.035] N=1,095	0.001 [0.038] N=945	0.018 [0.041] N=774
CIAF: any stunting, underweight, or wasting, 0-60 months	0.318 P=388	-0.061 [0.050] N=1,258	-0.069 [0.050] N=1,178	-0.056 [0.053] N=1,058	-0.058 [0.057] N=914	-0.031 [0.064] N=750
BMI-for-age z score below -2	0.120 P=156	-0.032 [0.038] N=1,266	-0.032 [0.038] N=1,186	-0.024 [0.042] N=1,065	0.005 [0.047] N=919	-0.016 [0.051] N=755

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.308 [0.353] N=1,116	0.321 [0.411] N=1,026	0.375 [0.482] N=900	0.626 [0.520] N=750	-0.083 [0.716] N=580
WHO weight-for-age z score	-0.146 [0.186] N=1,275	-0.094 [0.195] N=1,167	-0.149 [0.214] N=1,024	-0.237 [0.273] N=847	**-0.844** [0.389] N=654**
WHO weight-for-length/height z score, 0-60 months	0.141 [0.254] N=1,096	0.050 [0.271] N=1,014	-0.224 [0.343] N=892	-0.419 [0.415] N=742	-0.368 [0.514] N=571
WHO BMI-for-age z score	0.296 [0.269] N=1,070	0.316 [0.296] N=990	0.068 [0.371] N=869	-0.272 [0.450] N=723	0.011 [0.549] N=559
Stunted: HAZ below -2	-0.023 [0.061] N=1,116	-0.044 [0.068] N=1,026	-0.010 [0.073] N=900	-0.036 [0.082] N=750	0.070 [0.116] N=580
Underweight: WAZ below -2	-0.001 [0.030] N=1,275	-0.013 [0.032] N=1,167	-0.024 [0.035] N=1,024	0.004 [0.042] N=847	0.107 [0.068] N=654
Wasted: WLZ/WHZ below -2, 0-60 months	-0.040 [0.049] N=1,096	-0.025 [0.049] N=1,014	-0.022 [0.055] N=892	-0.008 [0.072] N=742	-0.080 [0.083] N=571
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.054] N=1,096	-0.016 [0.057] N=1,014	-0.053 [0.069] N=892	-0.079 [0.087] N=742	-0.108 [0.102] N=571
Obese: WLZ/WHZ above +3, 0-60 months	-0.006 [0.040] N=1,096	-0.018 [0.044] N=1,014	-0.022 [0.052] N=892	-0.027 [0.062] N=742	-0.002 [0.073] N=571
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.045 [0.074] N=1,063	-0.056 [0.075] N=983	-0.031 [0.084] N=863	-0.052 [0.099] N=719	0.027 [0.142] N=555
BMI-for-age z score below -2	-0.051 [0.048] N=1,070	-0.054 [0.048] N=990	-0.039 [0.056] N=869	0.018 [0.074] N=723	-0.044 [0.091] N=559

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.228	-0.217 [0.296] N=1,273	-0.279 [0.300] N=1,151	-0.251 [0.312] N=1,009	-0.472 [0.350] N=855	-0.539 [0.384] N=700
WHO weight-for-age z score	0.128	0.089 [0.146] N=1,433	0.127 [0.152] N=1,305	0.013 [0.153] N=1,144	0.080 [0.161] N=962	0.088 [0.183] N=778
WHO weight-for-length/height z score, 0-60 months	0.371	0.272 [0.213] N=1,259	0.221 [0.222] N=1,140	0.144 [0.234] N=1,001	0.362 [0.249] N=845	0.210 [0.271] N=691
WHO BMI-for-age z score	0.312	0.166 [0.239] N=1,220	0.120 [0.249] N=1,104	-0.010 [0.264] N=971	0.167 [0.289] N=820	0.141 [0.316] N=675
Stunted: HAZ below -2	0.218 P=277	0.010 [0.052] N=1,273	0.005 [0.052] N=1,151	0.003 [0.055] N=1,009	0.032 [0.061] N=855	0.031 [0.068] N=700
Underweight: WAZ below -2	0.058 P=87	0.004 [0.027] N=1,433	-0.006 [0.027] N=1,305	0.003 [0.030] N=1,144	0.002 [0.032] N=962	-0.008 [0.034] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=140	0.015 [0.036] N=1,259	0.014 [0.038] N=1,140	0.036 [0.041] N=1,001	-0.035 [0.039] N=845	-0.020 [0.042] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=245	**0.113** [0.047] N=1,259**	**0.100** [0.048] N=1,140**	**0.098* [0.051] N=1,001**	**0.111* [0.058] N=845**	0.096 [0.061] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=119	0.055 [0.039] N=1,259	0.043 [0.039] N=1,140	0.034 [0.043] N=1,001	0.041 [0.049] N=845	0.002 [0.048] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.302 P=367	0.016 [0.058] N=1,218	-0.004 [0.058] N=1,100	0.021 [0.062] N=965	-0.012 [0.063] N=817	0.016 [0.069] N=670
BMI-for-age z score below -2	0.120 P=146	0.029 [0.043] N=1,220	0.028 [0.047] N=1,104	0.037 [0.052] N=971	-0.019 [0.050] N=820	-0.015 [0.053] N=675

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.053 [0.415] N=1,100	-0.123 [0.432] N=978	-0.081 [0.478] N=836	-0.574 [0.593] N=682	-0.798 [0.726] N=527
WHO weight-for-age z score	0.083 [0.210] N=1,246	0.134 [0.224] N=1,118	-0.073 [0.230] N=957	0.059 [0.249] N=775	0.063 [0.329] N=591
WHO weight-for-length/height z score, 0-60 months	0.161 [0.307] N=1,089	0.057 [0.326] N=970	-0.079 [0.367] N=831	0.381 [0.413] N=675	-0.041 [0.490] N=521
WHO BMI-for-age z score	-0.170 [0.324] N=1,060	-0.270 [0.344] N=944	-0.522 [0.385] N=811	-0.214 [0.461] N=660	-0.382 [0.539] N=515
Stunted: HAZ below -2	0.023 [0.084] N=1,100	0.008 [0.087] N=978	0.017 [0.097] N=836	0.091 [0.122] N=682	0.100 [0.145] N=527
Underweight: WAZ below -2	0.013 [0.035] N=1,246	-0.002 [0.037] N=1,118	0.016 [0.043] N=957	0.010 [0.046] N=775	-0.018 [0.048] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.077 [0.047] N=1,089	0.081 [0.053] N=970	**0.135** [0.060] N=831**	-0.001 [0.059] N=675	0.037 [0.066] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.156*** [0.058] N=1,089**	**0.133** [0.059] N=970**	**0.149** [0.066] N=831**	**0.191** [0.080] N=675**	0.142 [0.089] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.115*** [0.041] N=1,089**	**0.093** [0.040] N=970**	**0.095** [0.047] N=831**	**0.118* [0.064] N=675**	0.000 [0.071] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.066 [0.088] N=1,052	0.031 [0.093] N=934	0.097 [0.105] N=799	0.033 [0.120] N=651	0.087 [0.141] N=504
BMI-for-age z score below -2	**0.127*** [0.046] N=1,060**	**0.129** [0.054] N=944**	**0.164** [0.066] N=811**	0.053 [0.060] N=660	0.066 [0.069] N=515

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.228	-0.226 [0.295] N=1,273	-0.285 [0.298] N=1,151	-0.255 [0.311] N=1,009	-0.475 [0.348] N=855	-0.540 [0.383] N=700
WHO weight-for-age z score	0.128	0.092 [0.146] N=1,433	0.129 [0.152] N=1,305	0.016 [0.152] N=1,144	0.076 [0.162] N=962	0.081 [0.185] N=778
WHO weight-for-length/height z score, 0-60 months	0.371	0.279 [0.214] N=1,259	0.227 [0.222] N=1,140	0.149 [0.234] N=1,001	0.362 [0.250] N=845	0.207 [0.272] N=691
WHO BMI-for-age z score	0.312	0.170 [0.239] N=1,220	0.124 [0.248] N=1,104	-0.008 [0.264] N=971	0.166 [0.290] N=820	0.137 [0.318] N=675
Stunted: HAZ below -2	0.218 P=277	0.010 [0.052] N=1,273	0.004 [0.052] N=1,151	0.001 [0.055] N=1,009	0.032 [0.061] N=855	0.031 [0.068] N=700
Underweight: WAZ below -2	0.058 P=87	0.003 [0.027] N=1,433	-0.007 [0.027] N=1,305	0.002 [0.030] N=1,144	0.003 [0.032] N=962	-0.006 [0.034] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=140	0.012 [0.036] N=1,259	0.010 [0.039] N=1,140	0.032 [0.041] N=1,001	-0.035 [0.040] N=845	-0.019 [0.043] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=245	**0.113** [0.047] N=1,259**	**0.100** [0.048] N=1,140**	**0.097* [0.051] N=1,001**	**0.111* [0.058] N=845**	0.096 [0.061] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=119	0.054 [0.039] N=1,259	0.042 [0.038] N=1,140	0.033 [0.043] N=1,001	0.041 [0.049] N=845	0.003 [0.048] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.302 P=367	0.011 [0.058] N=1,218	-0.011 [0.058] N=1,100	0.014 [0.063] N=965	-0.013 [0.064] N=817	0.016 [0.070] N=670
BMI-for-age z score below -2	0.120 P=146	0.027 [0.043] N=1,220	0.025 [0.046] N=1,104	0.035 [0.051] N=971	-0.018 [0.051] N=820	-0.015 [0.054] N=675

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.062 [0.414] N=1,100	-0.128 [0.430] N=978	-0.077 [0.478] N=836	-0.584 [0.591] N=682	-0.813 [0.727] N=527
WHO weight-for-age z score	0.095 [0.207] N=1,246	0.146 [0.220] N=1,118	-0.059 [0.224] N=957	0.070 [0.245] N=775	0.088 [0.322] N=591
WHO weight-for-length/height z score, 0-60 months	0.180 [0.310] N=1,089	0.077 [0.329] N=970	-0.058 [0.370] N=831	0.401 [0.417] N=675	0.024 [0.512] N=521
WHO BMI-for-age z score	-0.147 [0.326] N=1,060	-0.246 [0.344] N=944	-0.497 [0.385] N=811	-0.191 [0.464] N=660	-0.320 [0.559] N=515
Stunted: HAZ below -2	0.023 [0.084] N=1,100	0.005 [0.087] N=978	0.012 [0.097] N=836	0.090 [0.123] N=682	0.102 [0.146] N=527
Underweight: WAZ below -2	0.011 [0.035] N=1,246	-0.004 [0.037] N=1,118	0.013 [0.043] N=957	0.008 [0.045] N=775	-0.021 [0.048] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.070 [0.048] N=1,089	0.070 [0.054] N=970	**0.123** [0.061] N=831**	-0.009 [0.060] N=675	0.022 [0.068] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.157*** [0.058] N=1,089**	**0.134** [0.059] N=970**	**0.148** [0.067] N=831**	**0.188** [0.080] N=675**	0.144 [0.091] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.113*** [0.042] N=1,089**	**0.092** [0.040] N=970**	**0.092* [0.047] N=831**	**0.114* [0.063] N=675**	-0.002 [0.071] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.053 [0.089] N=1,052	0.012 [0.094] N=934	0.073 [0.107] N=799	0.018 [0.122] N=651	0.067 [0.143] N=504
BMI-for-age z score below -2	**0.120** [0.047] N=1,060**	**0.119** [0.055] N=944**	**0.154** [0.065] N=811**	0.046 [0.061] N=660	0.057 [0.072] N=515

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.228	-0.234 [0.290] N=1,273	-0.282 [0.294] N=1,151	-0.250 [0.306] N=1,009	-0.467 [0.340] N=855	-0.545 [0.375] N=700
WHO weight-for-age z score	0.128	0.081 [0.148] N=1,433	0.125 [0.154] N=1,305	0.018 [0.155] N=1,144	0.082 [0.164] N=962	0.086 [0.187] N=778
WHO weight-for-length/height z score, 0-60 months	0.371	0.273 [0.214] N=1,259	0.224 [0.223] N=1,140	0.155 [0.235] N=1,001	0.361 [0.249] N=845	0.209 [0.271] N=691
WHO BMI-for-age z score	0.312	0.162 [0.240] N=1,220	0.119 [0.250] N=1,104	-0.003 [0.265] N=971	0.165 [0.289] N=820	0.141 [0.317] N=675
Stunted: HAZ below -2	0.218 P=277	0.012 [0.051] N=1,273	0.004 [0.051] N=1,151	0.001 [0.054] N=1,009	0.031 [0.059] N=855	0.030 [0.066] N=700
Underweight: WAZ below -2	0.058 P=87	0.004 [0.027] N=1,433	-0.006 [0.028] N=1,305	0.002 [0.030] N=1,144	0.003 [0.032] N=962	-0.007 [0.034] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=140	0.013 [0.037] N=1,259	0.010 [0.039] N=1,140	0.031 [0.042] N=1,001	-0.036 [0.040] N=845	-0.022 [0.043] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=245	**0.114** [0.047] N=1,259**	**0.099** [0.048] N=1,140**	**0.098* [0.051] N=1,001**	**0.110* [0.057] N=845**	0.094 [0.062] N=691
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=119	0.055 [0.038] N=1,259	0.042 [0.038] N=1,140	0.033 [0.043] N=1,001	0.040 [0.049] N=845	0.001 [0.049] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.302 P=367	0.016 [0.058] N=1,218	-0.009 [0.058] N=1,100	0.014 [0.063] N=965	-0.015 [0.065] N=817	0.012 [0.071] N=670
BMI-for-age z score below -2	0.120 P=146	0.029 [0.044] N=1,220	0.026 [0.047] N=1,104	0.035 [0.052] N=971	-0.019 [0.051] N=820	-0.014 [0.054] N=675

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.052 [0.408] N=1,100	-0.114 [0.426] N=978	-0.041 [0.467] N=836	-0.514 [0.582] N=682	-0.725 [0.709] N=527
WHO weight-for-age z score	0.070 [0.213] N=1,246	0.129 [0.226] N=1,118	-0.061 [0.234] N=957	0.075 [0.256] N=775	0.099 [0.335] N=591
WHO weight-for-length/height z score, 0-60 months	0.154 [0.308] N=1,089	0.068 [0.328] N=970	-0.065 [0.368] N=831	0.373 [0.415] N=675	-0.005 [0.504] N=521
WHO BMI-for-age z score	-0.168 [0.324] N=1,060	-0.254 [0.344] N=944	-0.513 [0.382] N=811	-0.234 [0.457] N=660	-0.351 [0.549] N=515
Stunted: HAZ below -2	0.019 [0.083] N=1,100	0.000 [0.086] N=978	0.006 [0.095] N=836	0.073 [0.119] N=682	0.080 [0.142] N=527
Underweight: WAZ below -2	0.010 [0.036] N=1,246	-0.006 [0.038] N=1,118	0.009 [0.044] N=957	0.001 [0.047] N=775	-0.030 [0.050] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.071 [0.048] N=1,089	0.068 [0.055] N=970	**0.121* [0.062] N=831**	-0.012 [0.061] N=675	0.017 [0.070] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.159*** [0.059] N=1,089**	**0.136** [0.060] N=970**	**0.149** [0.066] N=831**	**0.184** [0.081] N=675**	0.139 [0.091] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.111*** [0.042] N=1,089**	**0.089** [0.040] N=970**	**0.090* [0.048] N=831**	**0.108* [0.064] N=675**	-0.008 [0.070] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.055 [0.090] N=1,052	0.011 [0.095] N=934	0.071 [0.109] N=799	0.004 [0.122] N=651	0.048 [0.142] N=504
BMI-for-age z score below -2	**0.117** [0.048] N=1,060**	**0.115** [0.056] N=944**	**0.151** [0.066] N=811**	0.039 [0.062] N=660	0.049 [0.074] N=515

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.228	-0.195 [0.292] N=1,273	-0.244 [0.296] N=1,151	-0.212 [0.311] N=1,009	-0.458 [0.348] N=855	-0.544 [0.386] N=700
WHO weight-for-age z score	0.128	0.088 [0.149] N=1,433	0.127 [0.156] N=1,305	0.024 [0.156] N=1,144	0.078 [0.166] N=962	0.083 [0.187] N=778
WHO weight-for-length/height z score, 0-60 months	0.371	0.250 [0.212] N=1,259	0.199 [0.219] N=1,140	0.126 [0.230] N=1,001	0.355 [0.245] N=845	0.220 [0.269] N=691
WHO BMI-for-age z score	0.312	0.135 [0.236] N=1,220	0.090 [0.245] N=1,104	-0.040 [0.258] N=971	0.144 [0.283] N=820	0.150 [0.316] N=675
Stunted: HAZ below -2	0.218 P=277	0.005 [0.051] N=1,273	-0.003 [0.051] N=1,151	-0.006 [0.054] N=1,009	0.029 [0.060] N=855	0.032 [0.066] N=700
Underweight: WAZ below -2	0.058 P=87	0.002 [0.027] N=1,433	-0.008 [0.028] N=1,305	-0.002 [0.031] N=1,144	-0.001 [0.033] N=962	-0.014 [0.036] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=140	0.015 [0.036] N=1,259	0.013 [0.039] N=1,140	0.035 [0.041] N=1,001	-0.034 [0.039] N=845	-0.022 [0.043] N=691
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=245	**0.110** [0.046] N=1,259**	**0.096** [0.047] N=1,140**	**0.096* [0.050] N=1,001**	**0.112** [0.055] N=845**	**0.099* [0.058] N=691**
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=119	0.056 [0.038] N=1,259	0.044 [0.037] N=1,140	0.035 [0.042] N=1,001	0.044 [0.047] N=845	0.006 [0.046] N=691
CIAF: any stunting, underweight, or wasting, 0-60 months	0.302 P=367	0.013 [0.058] N=1,218	-0.010 [0.058] N=1,100	0.014 [0.063] N=965	-0.010 [0.064] N=817	0.022 [0.069] N=670
BMI-for-age z score below -2	0.120 P=146	0.026 [0.043] N=1,220	0.024 [0.047] N=1,104	0.035 [0.051] N=971	-0.020 [0.051] N=820	-0.021 [0.054] N=675

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.015 [0.402] N=1,100	-0.072 [0.418] N=978	-0.006 [0.464] N=836	-0.565 [0.579] N=682	-0.888 [0.699] N=527
WHO weight-for-age z score	0.103 [0.206] N=1,246	0.155 [0.221] N=1,118	-0.023 [0.224] N=957	0.087 [0.244] N=775	0.097 [0.307] N=591
WHO weight-for-length/height z score, 0-60 months	0.144 [0.307] N=1,089	0.039 [0.325] N=970	-0.100 [0.360] N=831	0.381 [0.414] N=675	0.070 [0.524] N=521
WHO BMI-for-age z score	-0.195 [0.322] N=1,060	-0.292 [0.340] N=944	-0.557 [0.375] N=811	-0.237 [0.455] N=660	-0.271 [0.571] N=515
Stunted: HAZ below -2	0.012 [0.082] N=1,100	-0.010 [0.085] N=978	-0.006 [0.096] N=836	0.073 [0.121] N=682	0.101 [0.144] N=527
Underweight: WAZ below -2	0.009 [0.035] N=1,246	-0.007 [0.037] N=1,118	0.005 [0.044] N=957	0.001 [0.046] N=775	-0.035 [0.049] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.074 [0.048] N=1,089	0.074 [0.054] N=970	**0.130** [0.060] N=831**	-0.008 [0.060] N=675	0.015 [0.068] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.152*** [0.058] N=1,089**	**0.128** [0.058] N=970**	**0.143** [0.065] N=831**	**0.183** [0.080] N=675**	0.150 [0.091] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.112*** [0.042] N=1,089**	**0.090** [0.040] N=970**	**0.089* [0.047] N=831**	**0.111* [0.063] N=675**	-0.002 [0.072] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.050 [0.089] N=1,052	0.006 [0.095] N=934	0.062 [0.109] N=799	0.003 [0.122] N=651	0.068 [0.142] N=504
BMI-for-age z score below -2	**0.120** [0.047] N=1,060**	**0.120** [0.055] N=944**	**0.158** [0.065] N=811**	0.046 [0.061] N=660	0.050 [0.073] N=515

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.229	-0.245 [0.290] N=1,185	-0.314 [0.298] N=1,070	-0.274 [0.308] N=942	-0.535 [0.339] N=797	**-0.670* [0.371] N=654**
WHO weight-for-age z score	0.128	0.136 [0.153] N=1,334	0.188 [0.159] N=1,213	0.054 [0.161] N=1,067	0.098 [0.169] N=898	0.056 [0.186] N=729
WHO weight-for-length/height z score, 0-60 months	0.375	0.349 [0.230] N=1,172	0.302 [0.243] N=1,060	0.184 [0.255] N=933	0.382 [0.272] N=787	0.225 [0.290] N=646
WHO BMI-for-age z score	0.315	0.176 [0.253] N=1,136	0.135 [0.267] N=1,027	-0.028 [0.282] N=906	0.133 [0.306] N=765	0.088 [0.334] N=632
Stunted: HAZ below -2	0.219 P=258	0.016 [0.054] N=1,185	0.016 [0.055] N=1,070	0.006 [0.057] N=942	0.033 [0.062] N=797	0.043 [0.069] N=654
Underweight: WAZ below -2	0.058 P=80	-0.001 [0.027] N=1,334	-0.013 [0.028] N=1,213	-0.008 [0.031] N=1,067	-0.005 [0.033] N=898	-0.009 [0.035] N=729
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=131	0.013 [0.038] N=1,172	0.010 [0.041] N=1,060	0.037 [0.042] N=933	-0.024 [0.042] N=787	-0.006 [0.045] N=646
Overweight: WLZ/WHZ above +2, 0-60 months	0.191 P=228	**0.123** [0.048] N=1,172**	**0.109** [0.049] N=1,060**	**0.101* [0.052] N=933**	**0.110* [0.058] N=787**	0.093 [0.062] N=646
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=114	**0.064* [0.038] N=1,172**	0.050 [0.038] N=1,060	0.037 [0.042] N=933	0.050 [0.048] N=787	0.012 [0.048] N=646
CIAF: any stunting, underweight, or wasting, 0-60 months	0.305 P=344	0.022 [0.060] N=1,134	0.003 [0.061] N=1,023	0.028 [0.065] N=900	-0.004 [0.066] N=761	0.043 [0.072] N=626
BMI-for-age z score below -2	0.121 P=137	0.036 [0.045] N=1,136	0.032 [0.049] N=1,027	0.043 [0.052] N=906	-0.005 [0.052] N=765	0.004 [0.055] N=632

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.112 [0.411] N=1,021	0.051 [0.434] N=906	0.131 [0.478] N=778	-0.385 [0.567] N=633	-0.634 [0.674] N=490
WHO weight-for-age z score	0.199 [0.215] N=1,157	0.275 [0.226] N=1,036	0.039 [0.235] N=890	0.177 [0.256] N=721	0.149 [0.320] N=552
WHO weight-for-length/height z score, 0-60 months	0.246 [0.335] N=1,011	0.150 [0.360] N=899	-0.058 [0.399] N=772	0.384 [0.451] N=626	0.001 [0.561] N=485
WHO BMI-for-age z score	-0.146 [0.351] N=983	-0.244 [0.379] N=874	-0.571 [0.420] N=753	-0.260 [0.499] N=612	-0.439 [0.614] N=479
Stunted: HAZ below -2	0.003 [0.090] N=1,021	-0.012 [0.094] N=906	-0.017 [0.103] N=778	0.040 [0.126] N=633	0.053 [0.154] N=490
Underweight: WAZ below -2	0.002 [0.036] N=1,157	-0.019 [0.038] N=1,036	-0.010 [0.044] N=890	-0.016 [0.047] N=721	-0.044 [0.047] N=552
Wasted: WLZ/WHZ below -2, 0-60 months	0.068 [0.052] N=1,011	0.066 [0.059] N=899	**0.123* [0.063] N=772**	-0.006 [0.066] N=626	0.031 [0.075] N=485
Overweight: WLZ/WHZ above +2, 0-60 months	**0.171*** [0.064] N=1,011**	**0.143** [0.065] N=899**	**0.146** [0.073] N=772**	**0.173* [0.089] N=626**	0.130 [0.106] N=485
Obese: WLZ/WHZ above +3, 0-60 months	**0.116** [0.046] N=1,011**	**0.091** [0.045] N=899**	0.082 [0.052] N=772	0.110 [0.070] N=626	-0.013 [0.081] N=485
CIAF: any stunting, underweight, or wasting, 0-60 months	0.041 [0.097] N=977	-0.001 [0.104] N=866	0.059 [0.117] N=743	-0.026 [0.132] N=604	0.044 [0.159] N=469
BMI-for-age z score below -2	**0.133** [0.051] N=983**	**0.130** [0.061] N=874**	**0.164** [0.068] N=753**	0.059 [0.066] N=612	0.084 [0.076] N=479

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.229	-0.280 [0.298] N=1,185	-0.354 [0.305] N=1,070	-0.320 [0.315] N=942	**-0.585* [0.347] N=797**	**-0.724* [0.384] N=654**
WHO weight-for-age z score	0.128	0.135 [0.153] N=1,334	0.182 [0.160] N=1,213	0.050 [0.160] N=1,067	0.089 [0.169] N=898	0.049 [0.189] N=729
WHO weight-for-length/height z score, 0-60 months	0.375	0.377 [0.231] N=1,172	0.327 [0.243] N=1,060	0.215 [0.254] N=933	0.410 [0.270] N=787	0.252 [0.292] N=646
WHO BMI-for-age z score	0.315	0.207 [0.258] N=1,136	0.163 [0.271] N=1,027	0.006 [0.287] N=906	0.160 [0.309] N=765	0.115 [0.341] N=632
Stunted: HAZ below -2	0.219 P=258	0.019 [0.054] N=1,185	0.017 [0.054] N=1,070	0.008 [0.057] N=942	0.037 [0.062] N=797	0.049 [0.070] N=654
Underweight: WAZ below -2	0.058 P=80	-0.001 [0.027] N=1,334	-0.013 [0.028] N=1,213	-0.007 [0.031] N=1,067	-0.004 [0.032] N=898	-0.008 [0.035] N=729
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=131	0.007 [0.039] N=1,172	0.005 [0.042] N=1,060	0.029 [0.043] N=933	-0.031 [0.043] N=787	-0.012 [0.046] N=646
Overweight: WLZ/WHZ above +2, 0-60 months	0.191 P=228	**0.125*** [0.047] N=1,172**	**0.110** [0.048] N=1,060**	**0.102** [0.051] N=933**	**0.111* [0.057] N=787**	0.096 [0.061] N=646
Obese: WLZ/WHZ above +3, 0-60 months	0.094 P=114	**0.064* [0.038] N=1,172**	0.050 [0.038] N=1,060	0.037 [0.042] N=933	0.051 [0.048] N=787	0.013 [0.048] N=646
CIAF: any stunting, underweight, or wasting, 0-60 months	0.305 P=344	0.018 [0.060] N=1,134	-0.003 [0.061] N=1,023	0.021 [0.064] N=900	-0.009 [0.065] N=761	0.042 [0.071] N=626
BMI-for-age z score below -2	0.121 P=137	0.031 [0.046] N=1,136	0.027 [0.050] N=1,027	0.036 [0.054] N=906	-0.010 [0.054] N=765	-0.000 [0.057] N=632

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.066 [0.422] N=1,021	0.001 [0.441] N=906	0.062 [0.485] N=778	-0.497 [0.585] N=633	-0.795 [0.710] N=490
WHO weight-for-age z score	0.191 [0.217] N=1,157	0.261 [0.230] N=1,036	0.023 [0.237] N=890	0.150 [0.260] N=721	0.118 [0.334] N=552
WHO weight-for-length/height z score, 0-60 months	0.260 [0.333] N=1,011	0.155 [0.355] N=899	-0.043 [0.396] N=772	0.416 [0.447] N=626	0.008 [0.550] N=485
WHO BMI-for-age z score	-0.127 [0.350] N=983	-0.235 [0.374] N=874	-0.550 [0.417] N=753	-0.232 [0.496] N=612	-0.404 [0.609] N=479
Stunted: HAZ below -2	0.008 [0.090] N=1,021	-0.009 [0.093] N=906	-0.012 [0.102] N=778	0.053 [0.127] N=633	0.077 [0.157] N=490
Underweight: WAZ below -2	0.003 [0.036] N=1,157	-0.017 [0.038] N=1,036	-0.007 [0.044] N=890	-0.009 [0.047] N=721	-0.031 [0.050] N=552
Wasted: WLZ/WHZ below -2, 0-60 months	0.065 [0.052] N=1,011	0.065 [0.058] N=899	**0.118* [0.062] N=772**	-0.011 [0.065] N=626	0.029 [0.074] N=485
Overweight: WLZ/WHZ above +2, 0-60 months	**0.173*** [0.063] N=1,011**	**0.145** [0.064] N=899**	**0.147** [0.072] N=772**	**0.178** [0.089] N=626**	0.135 [0.105] N=485
Obese: WLZ/WHZ above +3, 0-60 months	**0.116** [0.046] N=1,011**	**0.090** [0.045] N=899**	0.082 [0.051] N=772	0.113 [0.069] N=626	-0.007 [0.081] N=485
CIAF: any stunting, underweight, or wasting, 0-60 months	0.042 [0.097] N=977	-0.000 [0.103] N=866	0.057 [0.115] N=743	-0.022 [0.132] N=604	0.058 [0.159] N=469
BMI-for-age z score below -2	**0.130** [0.051] N=983**	**0.129** [0.060] N=874**	**0.161** [0.067] N=753**	0.056 [0.064] N=612	0.082 [0.074] N=479

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.