Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,523	-0.006 [0.022] N=1,618	0.010 [0.022] N=1,493	0.002 [0.023] N=1,340	-0.006 [0.025] N=1,139	-0.013 [0.030] N=935
Vegetables consumed at least weekly	0.901 P=1,457	0.014 [0.029] N=1,618	0.014 [0.030] N=1,493	-0.004 [0.030] N=1,340	-0.011 [0.030] N=1,139	-0.032 [0.034] N=935
Meat fish or poultry consumed at least weekly	0.803 P=1,306	0.053 [0.043] N=1,618	0.068 [0.045] N=1,493	0.057 [0.049] N=1,340	0.035 [0.054] N=1,139	0.013 [0.059] N=935
Legumes consumed at least weekly	0.802 P=1,302	**-0.068* [0.039] N=1,618**	**-0.082** [0.040] N=1,493**	**-0.089** [0.041] N=1,340**	**-0.078* [0.046] N=1,139**	**-0.120** [0.052] N=935**
Animal products consumed at least weekly	0.917 P=1,485	0.015 [0.029] N=1,618	0.012 [0.030] N=1,493	-0.002 [0.034] N=1,340	0.005 [0.038] N=1,139	-0.000 [0.042] N=935
Sweets consumed at least weekly	0.712 P=1,146	-0.024 [0.038] N=1,618	-0.009 [0.041] N=1,493	0.005 [0.043] N=1,340	-0.000 [0.045] N=1,139	0.017 [0.051] N=935
Snack foods consumed at least weekly	0.584 P=935	-0.031 [0.042] N=1,618	-0.021 [0.045] N=1,493	-0.014 [0.049] N=1,340	-0.028 [0.053] N=1,139	-0.041 [0.055] N=935
Packaged soft drinks consumed at least weekly	0.359 P=570	0.013 [0.051] N=1,618	0.027 [0.055] N=1,493	0.023 [0.059] N=1,340	0.041 [0.065] N=1,139	0.019 [0.077] N=935
Grains consumed at least weekly	0.927 P=1,497	0.007 [0.027] N=1,618	0.017 [0.029] N=1,493	0.002 [0.030] N=1,340	0.007 [0.033] N=1,139	-0.015 [0.039] N=935
Child was ever breastfed	0.963 P=1,556	0.001 [0.016] N=1,620	0.004 [0.018] N=1,495	0.005 [0.019] N=1,341	0.008 [0.021] N=1,140	0.008 [0.022] N=936

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.032] N=1,377	0.043 [0.032] N=1,252	0.033 [0.034] N=1,099	0.020 [0.042] N=898	0.031 [0.057] N=694
Vegetables consumed at least weekly	0.058 [0.043] N=1,377	0.068 [0.047] N=1,252	0.041 [0.051] N=1,099	0.027 [0.060] N=898	-0.005 [0.074] N=694
Meat fish or poultry consumed at least weekly	0.049 [0.057] N=1,377	0.076 [0.058] N=1,252	0.056 [0.067] N=1,099	-0.013 [0.077] N=898	-0.073 [0.088] N=694
Legumes consumed at least weekly	-0.013 [0.060] N=1,377	-0.030 [0.067] N=1,252	-0.022 [0.071] N=1,099	0.013 [0.088] N=898	-0.049 [0.118] N=694
Animal products consumed at least weekly	0.016 [0.038] N=1,377	0.011 [0.040] N=1,252	-0.018 [0.049] N=1,099	-0.006 [0.061] N=898	-0.023 [0.078] N=694
Sweets consumed at least weekly	-0.031 [0.053] N=1,377	-0.005 [0.057] N=1,252	0.022 [0.065] N=1,099	0.017 [0.071] N=898	0.083 [0.092] N=694
Snack foods consumed at least weekly	-0.020 [0.057] N=1,377	-0.002 [0.063] N=1,252	0.013 [0.076] N=1,099	-0.002 [0.091] N=898	-0.037 [0.105] N=694
Packaged soft drinks consumed at least weekly	-0.009 [0.058] N=1,377	0.015 [0.064] N=1,252	0.005 [0.069] N=1,099	0.024 [0.079] N=898	-0.044 [0.103] N=694
Grains consumed at least weekly	0.048 [0.043] N=1,377	0.078 [0.049] N=1,252	0.056 [0.053] N=1,099	0.092 [0.065] N=898	0.084 [0.091] N=694
Child was ever breastfed	0.014 [0.026] N=1,379	0.022 [0.030] N=1,254	0.025 [0.036] N=1,100	0.032 [0.045] N=899	0.071 [0.054] N=695

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,523	-0.006 [0.022] N=1,618	0.010 [0.022] N=1,493	0.003 [0.023] N=1,340	-0.006 [0.025] N=1,139	-0.013 [0.030] N=935
Vegetables consumed at least weekly	0.901 P=1,457	0.014 [0.029] N=1,618	0.014 [0.030] N=1,493	-0.004 [0.030] N=1,340	-0.011 [0.030] N=1,139	-0.032 [0.033] N=935
Meat fish or poultry consumed at least weekly	0.803 P=1,306	0.053 [0.043] N=1,618	0.068 [0.045] N=1,493	0.057 [0.049] N=1,340	0.035 [0.055] N=1,139	0.013 [0.060] N=935
Legumes consumed at least weekly	0.802 P=1,302	**-0.068* [0.039] N=1,618**	**-0.082** [0.040] N=1,493**	**-0.088** [0.042] N=1,340**	**-0.078* [0.047] N=1,139**	**-0.120** [0.052] N=935**
Animal products consumed at least weekly	0.917 P=1,485	0.014 [0.029] N=1,618	0.011 [0.030] N=1,493	-0.001 [0.034] N=1,340	0.005 [0.038] N=1,139	-0.000 [0.042] N=935
Sweets consumed at least weekly	0.712 P=1,146	-0.024 [0.039] N=1,618	-0.009 [0.041] N=1,493	0.006 [0.043] N=1,340	-0.000 [0.045] N=1,139	0.017 [0.051] N=935
Snack foods consumed at least weekly	0.584 P=935	-0.031 [0.043] N=1,618	-0.022 [0.045] N=1,493	-0.014 [0.049] N=1,340	-0.028 [0.053] N=1,139	-0.041 [0.056] N=935
Packaged soft drinks consumed at least weekly	0.359 P=570	0.013 [0.051] N=1,618	0.027 [0.055] N=1,493	0.023 [0.059] N=1,340	0.041 [0.065] N=1,139	0.019 [0.077] N=935
Grains consumed at least weekly	0.927 P=1,497	0.007 [0.027] N=1,618	0.017 [0.029] N=1,493	0.002 [0.030] N=1,340	0.007 [0.033] N=1,139	-0.015 [0.039] N=935
Child was ever breastfed	0.963 P=1,556	0.001 [0.017] N=1,620	0.004 [0.018] N=1,495	0.005 [0.019] N=1,341	0.008 [0.021] N=1,140	0.008 [0.022] N=936

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.011 [0.032] N=1,377	0.044 [0.032] N=1,252	0.034 [0.034] N=1,099	0.021 [0.042] N=898	0.033 [0.056] N=694
Vegetables consumed at least weekly	0.058 [0.043] N=1,377	0.068 [0.047] N=1,252	0.041 [0.051] N=1,099	0.029 [0.059] N=898	-0.004 [0.073] N=694
Meat fish or poultry consumed at least weekly	0.049 [0.057] N=1,377	0.076 [0.058] N=1,252	0.056 [0.067] N=1,099	-0.010 [0.077] N=898	-0.070 [0.088] N=694
Legumes consumed at least weekly	-0.013 [0.060] N=1,377	-0.029 [0.067] N=1,252	-0.020 [0.071] N=1,099	0.016 [0.088] N=898	-0.048 [0.118] N=694
Animal products consumed at least weekly	0.017 [0.037] N=1,377	0.012 [0.040] N=1,252	-0.015 [0.048] N=1,099	-0.003 [0.060] N=898	-0.020 [0.077] N=694
Sweets consumed at least weekly	-0.031 [0.053] N=1,377	-0.005 [0.057] N=1,252	0.022 [0.065] N=1,099	0.018 [0.071] N=898	0.084 [0.092] N=694
Snack foods consumed at least weekly	-0.019 [0.057] N=1,377	-0.000 [0.063] N=1,252	0.018 [0.075] N=1,099	0.003 [0.090] N=898	-0.033 [0.105] N=694
Packaged soft drinks consumed at least weekly	-0.009 [0.058] N=1,377	0.015 [0.064] N=1,252	0.006 [0.069] N=1,099	0.025 [0.079] N=898	-0.045 [0.102] N=694
Grains consumed at least weekly	0.049 [0.043] N=1,377	0.078 [0.049] N=1,252	0.057 [0.053] N=1,099	0.092 [0.065] N=898	0.083 [0.091] N=694
Child was ever breastfed	0.015 [0.026] N=1,379	0.022 [0.030] N=1,254	0.026 [0.036] N=1,100	0.031 [0.045] N=899	0.070 [0.054] N=695

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,523	-0.015 [0.022] N=1,618	0.000 [0.022] N=1,493	-0.006 [0.023] N=1,340	-0.014 [0.026] N=1,139	-0.022 [0.031] N=935
Vegetables consumed at least weekly	0.901 P=1,457	0.007 [0.029] N=1,618	0.005 [0.030] N=1,493	-0.011 [0.031] N=1,340	-0.019 [0.031] N=1,139	-0.042 [0.034] N=935
Meat fish or poultry consumed at least weekly	0.803 P=1,306	0.030 [0.042] N=1,618	0.041 [0.043] N=1,493	0.033 [0.048] N=1,340	0.010 [0.052] N=1,139	-0.016 [0.058] N=935
Legumes consumed at least weekly	0.802 P=1,302	**-0.076* [0.039] N=1,618**	**-0.091** [0.041] N=1,493**	**-0.096** [0.042] N=1,340**	**-0.084* [0.047] N=1,139**	**-0.124** [0.053] N=935**
Animal products consumed at least weekly	0.917 P=1,485	0.009 [0.030] N=1,618	0.005 [0.031] N=1,493	-0.007 [0.035] N=1,340	-0.001 [0.039] N=1,139	-0.009 [0.043] N=935
Sweets consumed at least weekly	0.712 P=1,146	-0.030 [0.039] N=1,618	-0.016 [0.041] N=1,493	-0.002 [0.043] N=1,340	-0.006 [0.045] N=1,139	0.009 [0.051] N=935
Snack foods consumed at least weekly	0.584 P=935	-0.029 [0.042] N=1,618	-0.019 [0.045] N=1,493	-0.013 [0.048] N=1,340	-0.025 [0.052] N=1,139	-0.035 [0.055] N=935
Packaged soft drinks consumed at least weekly	0.359 P=570	0.025 [0.050] N=1,618	0.044 [0.054] N=1,493	0.038 [0.057] N=1,340	0.057 [0.063] N=1,139	0.039 [0.074] N=935
Grains consumed at least weekly	0.927 P=1,497	-0.001 [0.027] N=1,618	0.010 [0.029] N=1,493	-0.004 [0.030] N=1,340	-0.002 [0.033] N=1,139	-0.022 [0.038] N=935
Child was ever breastfed	0.963 P=1,556	-0.004 [0.017] N=1,620	-0.002 [0.018] N=1,495	-0.001 [0.019] N=1,341	0.002 [0.021] N=1,140	0.002 [0.022] N=936

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.004 [0.031] N=1,377	0.035 [0.030] N=1,252	0.029 [0.032] N=1,099	0.025 [0.039] N=898	0.039 [0.055] N=694
Vegetables consumed at least weekly	0.054 [0.043] N=1,377	0.062 [0.046] N=1,252	0.039 [0.051] N=1,099	0.035 [0.059] N=898	0.002 [0.073] N=694
Meat fish or poultry consumed at least weekly	0.031 [0.055] N=1,377	0.049 [0.056] N=1,252	0.036 [0.067] N=1,099	-0.028 [0.076] N=898	-0.088 [0.090] N=694
Legumes consumed at least weekly	-0.020 [0.061] N=1,377	-0.040 [0.067] N=1,252	-0.029 [0.071] N=1,099	0.017 [0.090] N=898	-0.044 [0.120] N=694
Animal products consumed at least weekly	0.014 [0.037] N=1,377	0.007 [0.040] N=1,252	-0.019 [0.049] N=1,099	-0.002 [0.061] N=898	-0.019 [0.078] N=694
Sweets consumed at least weekly	-0.035 [0.053] N=1,377	-0.011 [0.058] N=1,252	0.018 [0.065] N=1,099	0.017 [0.070] N=898	0.085 [0.091] N=694
Snack foods consumed at least weekly	-0.019 [0.056] N=1,377	0.001 [0.063] N=1,252	0.018 [0.073] N=1,099	0.004 [0.087] N=898	-0.025 [0.103] N=694
Packaged soft drinks consumed at least weekly	0.004 [0.057] N=1,377	0.037 [0.062] N=1,252	0.025 [0.067] N=1,099	0.046 [0.078] N=898	-0.019 [0.099] N=694
Grains consumed at least weekly	0.041 [0.042] N=1,377	0.071 [0.048] N=1,252	0.050 [0.053] N=1,099	0.083 [0.064] N=898	0.069 [0.089] N=694
Child was ever breastfed	0.011 [0.026] N=1,379	0.017 [0.030] N=1,254	0.022 [0.036] N=1,100	0.029 [0.046] N=899	0.064 [0.054] N=695

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,523	-0.017 [0.022] N=1,618	-0.001 [0.022] N=1,493	-0.009 [0.024] N=1,340	-0.017 [0.027] N=1,139	-0.022 [0.031] N=935
Vegetables consumed at least weekly	0.901 P=1,457	0.007 [0.029] N=1,618	0.005 [0.030] N=1,493	-0.012 [0.031] N=1,340	-0.021 [0.031] N=1,139	-0.042 [0.034] N=935
Meat fish or poultry consumed at least weekly	0.803 P=1,306	0.026 [0.042] N=1,618	0.035 [0.043] N=1,493	0.024 [0.048] N=1,340	0.004 [0.052] N=1,139	-0.025 [0.058] N=935
Legumes consumed at least weekly	0.802 P=1,302	**-0.076* [0.040] N=1,618**	**-0.093** [0.042] N=1,493**	**-0.102** [0.043] N=1,340**	**-0.087* [0.048] N=1,139**	**-0.127** [0.054] N=935**
Animal products consumed at least weekly	0.917 P=1,485	0.007 [0.030] N=1,618	0.006 [0.031] N=1,493	-0.007 [0.035] N=1,340	-0.002 [0.039] N=1,139	-0.009 [0.042] N=935
Sweets consumed at least weekly	0.712 P=1,146	-0.026 [0.038] N=1,618	-0.014 [0.040] N=1,493	-0.002 [0.042] N=1,340	-0.004 [0.045] N=1,139	0.012 [0.051] N=935
Snack foods consumed at least weekly	0.584 P=935	-0.028 [0.043] N=1,618	-0.018 [0.045] N=1,493	-0.014 [0.049] N=1,340	-0.023 [0.053] N=1,139	-0.034 [0.055] N=935
Packaged soft drinks consumed at least weekly	0.359 P=570	0.032 [0.050] N=1,618	0.048 [0.053] N=1,493	0.042 [0.057] N=1,340	0.065 [0.062] N=1,139	0.050 [0.073] N=935
Grains consumed at least weekly	0.927 P=1,497	0.000 [0.027] N=1,618	0.011 [0.029] N=1,493	-0.005 [0.030] N=1,340	-0.000 [0.033] N=1,139	-0.020 [0.040] N=935
Child was ever breastfed	0.963 P=1,556	-0.004 [0.017] N=1,620	-0.002 [0.018] N=1,495	-0.002 [0.020] N=1,341	0.002 [0.022] N=1,140	0.002 [0.022] N=936

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.003 [0.032] N=1,377	0.035 [0.031] N=1,252	0.028 [0.034] N=1,099	0.020 [0.041] N=898	0.030 [0.056] N=694
Vegetables consumed at least weekly	0.055 [0.043] N=1,377	0.063 [0.047] N=1,252	0.038 [0.051] N=1,099	0.031 [0.060] N=898	-0.011 [0.073] N=694
Meat fish or poultry consumed at least weekly	0.027 [0.057] N=1,377	0.046 [0.058] N=1,252	0.033 [0.068] N=1,099	-0.022 [0.078] N=898	-0.093 [0.096] N=694
Legumes consumed at least weekly	-0.015 [0.060] N=1,377	-0.035 [0.067] N=1,252	-0.023 [0.070] N=1,099	0.030 [0.087] N=898	-0.040 [0.122] N=694
Animal products consumed at least weekly	0.011 [0.037] N=1,377	0.008 [0.040] N=1,252	-0.019 [0.049] N=1,099	-0.009 [0.061] N=898	-0.028 [0.080] N=694
Sweets consumed at least weekly	-0.030 [0.053] N=1,377	-0.007 [0.056] N=1,252	0.021 [0.063] N=1,099	0.032 [0.070] N=898	0.097 [0.092] N=694
Snack foods consumed at least weekly	-0.013 [0.056] N=1,377	0.005 [0.062] N=1,252	0.021 [0.072] N=1,099	0.021 [0.087] N=898	-0.009 [0.101] N=694
Packaged soft drinks consumed at least weekly	0.010 [0.056] N=1,377	0.035 [0.061] N=1,252	0.024 [0.067] N=1,099	0.053 [0.077] N=898	0.004 [0.097] N=694
Grains consumed at least weekly	0.041 [0.041] N=1,377	0.070 [0.046] N=1,252	0.050 [0.051] N=1,099	0.085 [0.060] N=898	0.067 [0.083] N=694
Child was ever breastfed	0.012 [0.027] N=1,379	0.018 [0.031] N=1,254	0.022 [0.036] N=1,100	0.032 [0.046] N=899	0.069 [0.056] N=695

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,421	-0.015 [0.023] N=1,511	-0.002 [0.023] N=1,401	-0.007 [0.024] N=1,255	-0.014 [0.027] N=1,070	-0.023 [0.031] N=880
Vegetables consumed at least weekly	0.901 P=1,361	0.001 [0.030] N=1,511	-0.002 [0.030] N=1,401	-0.021 [0.031] N=1,255	-0.023 [0.031] N=1,070	-0.039 [0.034] N=880
Meat fish or poultry consumed at least weekly	0.802 P=1,217	0.038 [0.042] N=1,511	0.043 [0.043] N=1,401	0.026 [0.047] N=1,255	0.003 [0.052] N=1,070	-0.031 [0.055] N=880
Legumes consumed at least weekly	0.798 P=1,209	**-0.083** [0.039] N=1,511**	**-0.101** [0.040] N=1,401**	**-0.108*** [0.040] N=1,255**	**-0.092** [0.046] N=1,070**	**-0.133** [0.053] N=880**
Animal products consumed at least weekly	0.918 P=1,388	0.013 [0.031] N=1,511	0.010 [0.032] N=1,401	-0.003 [0.035] N=1,255	-0.003 [0.039] N=1,070	-0.006 [0.044] N=880
Sweets consumed at least weekly	0.712 P=1,069	-0.006 [0.039] N=1,511	0.003 [0.041] N=1,401	0.012 [0.043] N=1,255	0.011 [0.046] N=1,070	0.031 [0.052] N=880
Snack foods consumed at least weekly	0.584 P=872	-0.032 [0.045] N=1,511	-0.021 [0.048] N=1,401	-0.017 [0.052] N=1,255	-0.029 [0.057] N=1,070	-0.023 [0.062] N=880
Packaged soft drinks consumed at least weekly	0.365 P=541	0.049 [0.051] N=1,511	0.065 [0.053] N=1,401	0.051 [0.058] N=1,255	0.065 [0.062] N=1,070	0.062 [0.073] N=880
Grains consumed at least weekly	0.926 P=1,396	0.001 [0.029] N=1,511	0.009 [0.031] N=1,401	-0.008 [0.031] N=1,255	-0.000 [0.035] N=1,070	-0.020 [0.042] N=880
Child was ever breastfed	0.964 P=1,454	-0.003 [0.018] N=1,513	0.003 [0.019] N=1,403	0.005 [0.020] N=1,256	0.011 [0.022] N=1,071	0.009 [0.024] N=881

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.008 [0.034] N=1,285	0.033 [0.035] N=1,175	0.025 [0.038] N=1,029	0.013 [0.046] N=844	0.029 [0.062] N=654
Vegetables consumed at least weekly	0.047 [0.045] N=1,285	0.051 [0.049] N=1,175	0.018 [0.054] N=1,029	0.016 [0.062] N=844	-0.000 [0.075] N=654
Meat fish or poultry consumed at least weekly	0.060 [0.058] N=1,285	0.068 [0.059] N=1,175	0.046 [0.068] N=1,029	-0.019 [0.078] N=844	-0.074 [0.088] N=654
Legumes consumed at least weekly	-0.019 [0.062] N=1,285	-0.047 [0.068] N=1,175	-0.045 [0.071] N=1,029	-0.007 [0.088] N=844	-0.067 [0.118] N=654
Animal products consumed at least weekly	0.018 [0.040] N=1,285	0.011 [0.043] N=1,175	-0.015 [0.051] N=1,029	-0.006 [0.063] N=844	-0.010 [0.082] N=654
Sweets consumed at least weekly	-0.010 [0.054] N=1,285	0.004 [0.060] N=1,175	0.015 [0.066] N=1,029	0.015 [0.074] N=844	0.055 [0.100] N=654
Snack foods consumed at least weekly	-0.027 [0.058] N=1,285	-0.008 [0.065] N=1,175	-0.003 [0.076] N=1,029	-0.026 [0.091] N=844	-0.059 [0.106] N=654
Packaged soft drinks consumed at least weekly	0.024 [0.059] N=1,285	0.046 [0.064] N=1,175	0.019 [0.070] N=1,029	0.024 [0.079] N=844	-0.018 [0.103] N=654
Grains consumed at least weekly	0.048 [0.046] N=1,285	0.073 [0.051] N=1,175	0.043 [0.054] N=1,029	0.089 [0.066] N=844	0.094 [0.095] N=654
Child was ever breastfed	0.008 [0.027] N=1,287	0.018 [0.030] N=1,177	0.021 [0.036] N=1,030	0.030 [0.045] N=845	0.059 [0.054] N=655

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.942 P=1,421	-0.015 [0.023] N=1,511	-0.002 [0.022] N=1,401	-0.008 [0.024] N=1,255	-0.016 [0.026] N=1,070	-0.026 [0.030] N=880
Vegetables consumed at least weekly	0.901 P=1,361	0.000 [0.030] N=1,511	-0.002 [0.031] N=1,401	-0.022 [0.031] N=1,255	-0.024 [0.031] N=1,070	-0.041 [0.035] N=880
Meat fish or poultry consumed at least weekly	0.802 P=1,217	0.042 [0.043] N=1,511	0.047 [0.044] N=1,401	0.033 [0.049] N=1,255	0.011 [0.054] N=1,070	-0.014 [0.059] N=880
Legumes consumed at least weekly	0.798 P=1,209	**-0.084** [0.039] N=1,511**	**-0.103** [0.040] N=1,401**	**-0.111*** [0.040] N=1,255**	**-0.098** [0.046] N=1,070**	**-0.146*** [0.053] N=880**
Animal products consumed at least weekly	0.918 P=1,388	0.012 [0.031] N=1,511	0.009 [0.032] N=1,401	-0.004 [0.035] N=1,255	-0.002 [0.039] N=1,070	-0.004 [0.043] N=880
Sweets consumed at least weekly	0.712 P=1,069	-0.005 [0.038] N=1,511	0.004 [0.041] N=1,401	0.015 [0.042] N=1,255	0.014 [0.044] N=1,070	0.034 [0.050] N=880
Snack foods consumed at least weekly	0.584 P=872	-0.032 [0.045] N=1,511	-0.022 [0.048] N=1,401	-0.017 [0.052] N=1,255	-0.030 [0.056] N=1,070	-0.030 [0.060] N=880
Packaged soft drinks consumed at least weekly	0.365 P=541	0.046 [0.051] N=1,511	0.062 [0.054] N=1,401	0.050 [0.058] N=1,255	0.062 [0.063] N=1,070	0.050 [0.074] N=880
Grains consumed at least weekly	0.926 P=1,396	0.000 [0.029] N=1,511	0.009 [0.031] N=1,401	-0.009 [0.032] N=1,255	-0.002 [0.035] N=1,070	-0.023 [0.041] N=880
Child was ever breastfed	0.964 P=1,454	-0.003 [0.018] N=1,513	0.002 [0.019] N=1,403	0.003 [0.020] N=1,256	0.009 [0.022] N=1,071	0.006 [0.023] N=881

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.007 [0.034] N=1,285	0.033 [0.034] N=1,175	0.026 [0.037] N=1,029	0.015 [0.045] N=844	0.026 [0.059] N=654
Vegetables consumed at least weekly	0.045 [0.045] N=1,285	0.050 [0.049] N=1,175	0.019 [0.054] N=1,029	0.015 [0.062] N=844	-0.004 [0.074] N=654
Meat fish or poultry consumed at least weekly	0.056 [0.057] N=1,285	0.064 [0.058] N=1,175	0.043 [0.067] N=1,029	-0.018 [0.078] N=844	-0.064 [0.087] N=654
Legumes consumed at least weekly	-0.022 [0.062] N=1,285	-0.050 [0.068] N=1,175	-0.041 [0.072] N=1,029	-0.002 [0.090] N=844	-0.071 [0.121] N=654
Animal products consumed at least weekly	0.016 [0.039] N=1,285	0.010 [0.042] N=1,175	-0.014 [0.051] N=1,029	-0.010 [0.063] N=844	-0.017 [0.082] N=654
Sweets consumed at least weekly	-0.020 [0.054] N=1,285	-0.010 [0.059] N=1,175	0.007 [0.066] N=1,029	0.011 [0.073] N=844	0.061 [0.098] N=654
Snack foods consumed at least weekly	-0.036 [0.057] N=1,285	-0.021 [0.064] N=1,175	-0.011 [0.076] N=1,029	-0.029 [0.091] N=844	-0.054 [0.107] N=654
Packaged soft drinks consumed at least weekly	0.020 [0.059] N=1,285	0.041 [0.064] N=1,175	0.017 [0.070] N=1,029	0.022 [0.079] N=844	-0.025 [0.103] N=654
Grains consumed at least weekly	0.047 [0.045] N=1,285	0.073 [0.051] N=1,175	0.048 [0.056] N=1,029	0.091 [0.069] N=844	0.092 [0.097] N=654
Child was ever breastfed	0.010 [0.028] N=1,287	0.020 [0.032] N=1,177	0.022 [0.038] N=1,030	0.033 [0.048] N=845	0.065 [0.058] N=655

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,366	-0.013 [0.027] N=1,446	-0.014 [0.029] N=1,316	-0.019 [0.032] N=1,150	-0.020 [0.032] N=964	-0.022 [0.038] N=771
Vegetables consumed at least weekly	0.903 P=1,304	0.010 [0.032] N=1,446	0.018 [0.034] N=1,316	0.020 [0.037] N=1,150	0.015 [0.041] N=964	0.017 [0.046] N=771
Meat fish or poultry consumed at least weekly	0.790 P=1,154	0.068 [0.042] N=1,446	**0.080* [0.044] N=1,316**	**0.082* [0.047] N=1,150**	**0.126*** [0.048] N=964**	**0.119** [0.056] N=771**
Legumes consumed at least weekly	0.802 P=1,167	-0.024 [0.045] N=1,446	-0.005 [0.049] N=1,316	-0.013 [0.052] N=1,150	-0.027 [0.056] N=964	-0.009 [0.067] N=771
Animal products consumed at least weekly	0.915 P=1,323	0.002 [0.034] N=1,446	0.002 [0.036] N=1,316	0.010 [0.038] N=1,150	-0.004 [0.041] N=964	0.019 [0.047] N=771
Sweets consumed at least weekly	0.719 P=1,041	0.024 [0.051] N=1,446	0.020 [0.054] N=1,316	0.009 [0.058] N=1,150	0.011 [0.062] N=964	-0.036 [0.071] N=771
Snack foods consumed at least weekly	0.583 P=838	0.002 [0.057] N=1,446	0.004 [0.060] N=1,316	0.019 [0.064] N=1,150	0.037 [0.067] N=964	-0.011 [0.077] N=771
Packaged soft drinks consumed at least weekly	0.346 P=496	0.001 [0.048] N=1,446	0.002 [0.050] N=1,316	0.040 [0.052] N=1,150	0.048 [0.056] N=964	0.045 [0.064] N=771
Grains consumed at least weekly	0.928 P=1,340	0.047 [0.037] N=1,446	0.051 [0.039] N=1,316	0.053 [0.043] N=1,150	0.055 [0.047] N=964	0.051 [0.054] N=771
Child was ever breastfed	0.961 P=1,390	-0.009 [0.021] N=1,447	-0.008 [0.022] N=1,317	-0.009 [0.022] N=1,151	-0.010 [0.023] N=965	-0.013 [0.027] N=772

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.035] N=1,259	-0.052 [0.038] N=1,129	-0.068 [0.045] N=963	-0.064 [0.046] N=777	-0.083 [0.068] N=584
Vegetables consumed at least weekly	-0.034 [0.038] N=1,259	-0.026 [0.041] N=1,129	-0.036 [0.048] N=963	-0.039 [0.054] N=777	-0.040 [0.073] N=584
Meat fish or poultry consumed at least weekly	0.004 [0.059] N=1,259	0.010 [0.062] N=1,129	-0.009 [0.068] N=963	0.073 [0.071] N=777	0.030 [0.091] N=584
Legumes consumed at least weekly	-0.045 [0.054] N=1,259	-0.016 [0.059] N=1,129	-0.029 [0.070] N=963	-0.037 [0.077] N=777	0.031 [0.092] N=584
Animal products consumed at least weekly	-0.048 [0.040] N=1,259	-0.047 [0.044] N=1,129	-0.046 [0.049] N=963	**-0.093* [0.051] N=777**	-0.079 [0.065] N=584
Sweets consumed at least weekly	0.029 [0.068] N=1,259	0.025 [0.074] N=1,129	0.009 [0.083] N=963	0.009 [0.093] N=777	-0.117 [0.117] N=584
Snack foods consumed at least weekly	0.029 [0.085] N=1,259	0.033 [0.092] N=1,129	0.064 [0.103] N=963	0.122 [0.117] N=777	0.044 [0.156] N=584
Packaged soft drinks consumed at least weekly	-0.027 [0.077] N=1,259	-0.023 [0.083] N=1,129	0.041 [0.094] N=963	0.057 [0.107] N=777	0.053 [0.137] N=584
Grains consumed at least weekly	0.026 [0.039] N=1,259	0.025 [0.041] N=1,129	0.020 [0.048] N=963	0.004 [0.059] N=777	-0.056 [0.083] N=584
Child was ever breastfed	0.020 [0.040] N=1,260	0.024 [0.042] N=1,130	0.028 [0.046] N=964	0.035 [0.052] N=778	0.047 [0.070] N=585

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,366	-0.012 [0.027] N=1,446	-0.014 [0.029] N=1,316	-0.018 [0.032] N=1,150	-0.020 [0.032] N=964	-0.022 [0.038] N=771
Vegetables consumed at least weekly	0.903 P=1,304	0.010 [0.032] N=1,446	0.019 [0.034] N=1,316	0.020 [0.037] N=1,150	0.015 [0.040] N=964	0.017 [0.046] N=771
Meat fish or poultry consumed at least weekly	0.790 P=1,154	0.069 [0.042] N=1,446	**0.081* [0.044] N=1,316**	**0.084* [0.047] N=1,150**	**0.126*** [0.048] N=964**	**0.119** [0.056] N=771**
Legumes consumed at least weekly	0.802 P=1,167	-0.024 [0.045] N=1,446	-0.006 [0.049] N=1,316	-0.014 [0.052] N=1,150	-0.027 [0.056] N=964	-0.009 [0.067] N=771
Animal products consumed at least weekly	0.915 P=1,323	0.005 [0.034] N=1,446	0.005 [0.036] N=1,316	0.013 [0.038] N=1,150	-0.003 [0.041] N=964	0.019 [0.047] N=771
Sweets consumed at least weekly	0.719 P=1,041	0.027 [0.052] N=1,446	0.023 [0.054] N=1,316	0.011 [0.058] N=1,150	0.011 [0.062] N=964	-0.036 [0.071] N=771
Snack foods consumed at least weekly	0.583 P=838	0.006 [0.057] N=1,446	0.007 [0.060] N=1,316	0.022 [0.064] N=1,150	0.038 [0.068] N=964	-0.011 [0.077] N=771
Packaged soft drinks consumed at least weekly	0.346 P=496	0.003 [0.047] N=1,446	0.003 [0.049] N=1,316	0.041 [0.052] N=1,150	0.048 [0.056] N=964	0.045 [0.064] N=771
Grains consumed at least weekly	0.928 P=1,340	0.048 [0.037] N=1,446	0.051 [0.039] N=1,316	0.053 [0.043] N=1,150	0.055 [0.047] N=964	0.051 [0.054] N=771
Child was ever breastfed	0.961 P=1,390	-0.009 [0.021] N=1,447	-0.008 [0.022] N=1,317	-0.010 [0.022] N=1,151	-0.010 [0.023] N=965	-0.013 [0.027] N=772

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.042 [0.035] N=1,259	-0.051 [0.038] N=1,129	-0.068 [0.046] N=963	-0.064 [0.047] N=777	-0.083 [0.068] N=584
Vegetables consumed at least weekly	-0.033 [0.037] N=1,259	-0.025 [0.041] N=1,129	-0.037 [0.048] N=963	-0.037 [0.054] N=777	-0.040 [0.073] N=584
Meat fish or poultry consumed at least weekly	0.007 [0.059] N=1,259	0.016 [0.062] N=1,129	-0.004 [0.068] N=963	0.078 [0.071] N=777	0.039 [0.094] N=584
Legumes consumed at least weekly	-0.046 [0.053] N=1,259	-0.018 [0.059] N=1,129	-0.032 [0.069] N=963	-0.039 [0.077] N=777	0.021 [0.091] N=584
Animal products consumed at least weekly	-0.042 [0.040] N=1,259	-0.040 [0.044] N=1,129	-0.040 [0.049] N=963	**-0.086* [0.051] N=777**	-0.069 [0.064] N=584
Sweets consumed at least weekly	0.035 [0.069] N=1,259	0.032 [0.075] N=1,129	0.015 [0.084] N=963	0.011 [0.093] N=777	-0.112 [0.117] N=584
Snack foods consumed at least weekly	0.039 [0.086] N=1,259	0.041 [0.093] N=1,129	0.072 [0.103] N=963	0.127 [0.118] N=777	0.055 [0.159] N=584
Packaged soft drinks consumed at least weekly	-0.023 [0.076] N=1,259	-0.021 [0.083] N=1,129	0.043 [0.094] N=963	0.058 [0.108] N=777	0.062 [0.137] N=584
Grains consumed at least weekly	0.027 [0.039] N=1,259	0.025 [0.042] N=1,129	0.020 [0.048] N=963	0.001 [0.059] N=777	-0.059 [0.083] N=584
Child was ever breastfed	0.020 [0.040] N=1,260	0.025 [0.042] N=1,130	0.028 [0.047] N=964	0.035 [0.053] N=778	0.047 [0.071] N=585

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,366	-0.014 [0.027] N=1,446	-0.015 [0.028] N=1,316	-0.018 [0.032] N=1,150	-0.021 [0.032] N=964	-0.023 [0.038] N=771
Vegetables consumed at least weekly	0.903 P=1,304	0.009 [0.032] N=1,446	0.018 [0.033] N=1,316	0.020 [0.036] N=1,150	0.015 [0.040] N=964	0.018 [0.046] N=771
Meat fish or poultry consumed at least weekly	0.790 P=1,154	0.063 [0.041] N=1,446	**0.077* [0.043] N=1,316**	**0.086* [0.045] N=1,150**	**0.126*** [0.046] N=964**	**0.121** [0.054] N=771**
Legumes consumed at least weekly	0.802 P=1,167	-0.026 [0.045] N=1,446	-0.007 [0.049] N=1,316	-0.014 [0.052] N=1,150	-0.028 [0.056] N=964	-0.009 [0.067] N=771
Animal products consumed at least weekly	0.915 P=1,323	0.003 [0.034] N=1,446	0.003 [0.036] N=1,316	0.013 [0.038] N=1,150	-0.004 [0.041] N=964	0.019 [0.047] N=771
Sweets consumed at least weekly	0.719 P=1,041	0.026 [0.052] N=1,446	0.022 [0.054] N=1,316	0.011 [0.058] N=1,150	0.010 [0.062] N=964	-0.037 [0.071] N=771
Snack foods consumed at least weekly	0.583 P=838	0.009 [0.058] N=1,446	0.009 [0.061] N=1,316	0.021 [0.064] N=1,150	0.036 [0.068] N=964	-0.013 [0.078] N=771
Packaged soft drinks consumed at least weekly	0.346 P=496	0.005 [0.049] N=1,446	0.003 [0.051] N=1,316	0.039 [0.053] N=1,150	0.046 [0.057] N=964	0.043 [0.066] N=771
Grains consumed at least weekly	0.928 P=1,340	0.046 [0.037] N=1,446	0.050 [0.039] N=1,316	0.054 [0.042] N=1,150	0.054 [0.047] N=964	0.051 [0.054] N=771
Child was ever breastfed	0.961 P=1,390	-0.009 [0.021] N=1,447	-0.008 [0.022] N=1,317	-0.009 [0.022] N=1,151	-0.010 [0.023] N=965	-0.012 [0.027] N=772

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.041 [0.035] N=1,259	-0.048 [0.037] N=1,129	-0.063 [0.044] N=963	-0.062 [0.047] N=777	-0.079 [0.068] N=584
Vegetables consumed at least weekly	-0.033 [0.037] N=1,259	-0.024 [0.040] N=1,129	-0.032 [0.046] N=963	-0.035 [0.053] N=777	-0.037 [0.073] N=584
Meat fish or poultry consumed at least weekly	0.008 [0.056] N=1,259	0.020 [0.059] N=1,129	0.011 [0.064] N=963	0.087 [0.070] N=777	0.047 [0.089] N=584
Legumes consumed at least weekly	-0.043 [0.052] N=1,259	-0.016 [0.058] N=1,129	-0.031 [0.070] N=963	-0.037 [0.076] N=777	0.019 [0.091] N=584
Animal products consumed at least weekly	-0.043 [0.040] N=1,259	-0.041 [0.043] N=1,129	-0.038 [0.048] N=963	**-0.089* [0.051] N=777**	-0.073 [0.064] N=584
Sweets consumed at least weekly	0.032 [0.069] N=1,259	0.028 [0.074] N=1,129	0.013 [0.083] N=963	0.004 [0.092] N=777	-0.123 [0.115] N=584
Snack foods consumed at least weekly	0.042 [0.086] N=1,259	0.042 [0.094] N=1,129	0.066 [0.104] N=963	0.118 [0.118] N=777	0.048 [0.162] N=584
Packaged soft drinks consumed at least weekly	-0.034 [0.078] N=1,259	-0.035 [0.085] N=1,129	0.023 [0.097] N=963	0.033 [0.111] N=777	0.042 [0.146] N=584
Grains consumed at least weekly	0.025 [0.038] N=1,259	0.025 [0.041] N=1,129	0.023 [0.046] N=963	0.000 [0.057] N=777	-0.063 [0.080] N=584
Child was ever breastfed	0.019 [0.040] N=1,260	0.024 [0.042] N=1,130	0.030 [0.047] N=964	0.038 [0.052] N=778	0.047 [0.070] N=585

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,366	-0.014 [0.027] N=1,446	-0.015 [0.028] N=1,316	-0.017 [0.032] N=1,150	-0.024 [0.032] N=964	-0.027 [0.037] N=771
Vegetables consumed at least weekly	0.903 P=1,304	0.010 [0.033] N=1,446	0.020 [0.034] N=1,316	0.022 [0.038] N=1,150	0.015 [0.041] N=964	0.019 [0.046] N=771
Meat fish or poultry consumed at least weekly	0.790 P=1,154	0.062 [0.043] N=1,446	**0.078* [0.044] N=1,316**	**0.083* [0.047] N=1,150**	**0.119** [0.048] N=964**	**0.111** [0.056] N=771**
Legumes consumed at least weekly	0.802 P=1,167	-0.022 [0.045] N=1,446	-0.003 [0.049] N=1,316	-0.009 [0.052] N=1,150	-0.024 [0.056] N=964	-0.005 [0.066] N=771
Animal products consumed at least weekly	0.915 P=1,323	0.010 [0.034] N=1,446	0.011 [0.036] N=1,316	0.021 [0.038] N=1,150	0.002 [0.041] N=964	0.024 [0.047] N=771
Sweets consumed at least weekly	0.719 P=1,041	0.033 [0.053] N=1,446	0.027 [0.055] N=1,316	0.014 [0.059] N=1,150	0.011 [0.063] N=964	-0.036 [0.073] N=771
Snack foods consumed at least weekly	0.583 P=838	0.016 [0.057] N=1,446	0.015 [0.060] N=1,316	0.026 [0.063] N=1,150	0.039 [0.067] N=964	-0.007 [0.077] N=771
Packaged soft drinks consumed at least weekly	0.346 P=496	0.016 [0.048] N=1,446	0.010 [0.050] N=1,316	0.044 [0.053] N=1,150	0.053 [0.057] N=964	0.051 [0.066] N=771
Grains consumed at least weekly	0.928 P=1,340	0.045 [0.037] N=1,446	0.048 [0.039] N=1,316	0.050 [0.042] N=1,150	0.051 [0.047] N=964	0.048 [0.054] N=771
Child was ever breastfed	0.961 P=1,390	-0.009 [0.021] N=1,447	-0.008 [0.022] N=1,317	-0.010 [0.022] N=1,151	-0.011 [0.024] N=965	-0.014 [0.028] N=772

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.044 [0.034] N=1,259	-0.051 [0.037] N=1,129	-0.064 [0.044] N=963	-0.067 [0.047] N=777	-0.093 [0.067] N=584
Vegetables consumed at least weekly	-0.033 [0.038] N=1,259	-0.023 [0.041] N=1,129	-0.030 [0.047] N=963	-0.034 [0.053] N=777	-0.035 [0.071] N=584
Meat fish or poultry consumed at least weekly	0.002 [0.058] N=1,259	0.016 [0.061] N=1,129	0.003 [0.066] N=963	0.069 [0.074] N=777	0.013 [0.092] N=584
Legumes consumed at least weekly	-0.042 [0.053] N=1,259	-0.014 [0.059] N=1,129	-0.022 [0.069] N=963	-0.029 [0.076] N=777	0.034 [0.088] N=584
Animal products consumed at least weekly	-0.037 [0.038] N=1,259	-0.033 [0.042] N=1,129	-0.031 [0.046] N=963	**-0.084* [0.050] N=777**	-0.067 [0.061] N=584
Sweets consumed at least weekly	0.039 [0.069] N=1,259	0.035 [0.075] N=1,129	0.014 [0.083] N=963	0.004 [0.093] N=777	-0.123 [0.114] N=584
Snack foods consumed at least weekly	0.046 [0.084] N=1,259	0.043 [0.092] N=1,129	0.067 [0.101] N=963	0.114 [0.115] N=777	0.046 [0.155] N=584
Packaged soft drinks consumed at least weekly	-0.012 [0.076] N=1,259	-0.018 [0.082] N=1,129	0.038 [0.094] N=963	0.054 [0.108] N=777	0.077 [0.141] N=584
Grains consumed at least weekly	0.023 [0.038] N=1,259	0.021 [0.041] N=1,129	0.015 [0.047] N=963	-0.007 [0.058] N=777	-0.070 [0.080] N=584
Child was ever breastfed	0.020 [0.039] N=1,260	0.025 [0.041] N=1,130	0.028 [0.046] N=964	0.034 [0.053] N=778	0.044 [0.072] N=585

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,275	-0.010 [0.029] N=1,350	-0.011 [0.030] N=1,227	-0.010 [0.034] N=1,076	-0.018 [0.034] N=901	-0.022 [0.040] N=724
Vegetables consumed at least weekly	0.904 P=1,220	0.013 [0.033] N=1,350	0.021 [0.035] N=1,227	0.033 [0.038] N=1,076	0.024 [0.042] N=901	0.011 [0.048] N=724
Meat fish or poultry consumed at least weekly	0.789 P=1,076	0.071 [0.044] N=1,350	**0.081* [0.046] N=1,227**	**0.087* [0.049] N=1,076**	**0.104** [0.051] N=901**	0.094 [0.059] N=724
Legumes consumed at least weekly	0.799 P=1,086	-0.021 [0.047] N=1,350	-0.001 [0.050] N=1,227	-0.007 [0.055] N=1,076	-0.026 [0.059] N=901	-0.015 [0.070] N=724
Animal products consumed at least weekly	0.915 P=1,237	0.007 [0.035] N=1,350	0.012 [0.038] N=1,227	0.022 [0.040] N=1,076	0.007 [0.042] N=901	0.029 [0.048] N=724
Sweets consumed at least weekly	0.726 P=980	0.025 [0.052] N=1,350	0.019 [0.055] N=1,227	0.003 [0.060] N=1,076	-0.001 [0.064] N=901	-0.043 [0.076] N=724
Snack foods consumed at least weekly	0.588 P=788	0.014 [0.059] N=1,350	0.010 [0.063] N=1,227	0.018 [0.066] N=1,076	0.034 [0.068] N=901	-0.006 [0.079] N=724
Packaged soft drinks consumed at least weekly	0.356 P=477	-0.006 [0.049] N=1,350	-0.011 [0.051] N=1,227	0.028 [0.054] N=1,076	0.039 [0.059] N=901	0.033 [0.069] N=724
Grains consumed at least weekly	0.927 P=1,250	0.051 [0.039] N=1,350	0.054 [0.041] N=1,227	0.058 [0.044] N=1,076	0.060 [0.049] N=901	0.055 [0.056] N=724
Child was ever breastfed	0.963 P=1,301	-0.010 [0.020] N=1,351	-0.011 [0.021] N=1,228	-0.015 [0.021] N=1,077	-0.021 [0.022] N=902	-0.026 [0.027] N=725

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.035 [0.037] N=1,174	-0.042 [0.040] N=1,051	-0.046 [0.050] N=900	-0.050 [0.050] N=725	-0.067 [0.067] N=548
Vegetables consumed at least weekly	-0.033 [0.038] N=1,174	-0.027 [0.042] N=1,051	-0.018 [0.049] N=900	-0.021 [0.058] N=725	-0.059 [0.072] N=548
Meat fish or poultry consumed at least weekly	0.028 [0.065] N=1,174	0.039 [0.068] N=1,051	0.032 [0.076] N=900	0.071 [0.081] N=725	0.027 [0.104] N=548
Legumes consumed at least weekly	-0.032 [0.055] N=1,174	-0.001 [0.059] N=1,051	-0.003 [0.072] N=900	-0.014 [0.081] N=725	0.036 [0.091] N=548
Animal products consumed at least weekly	-0.040 [0.042] N=1,174	-0.031 [0.046] N=1,051	-0.024 [0.051] N=900	-0.063 [0.054] N=725	-0.042 [0.065] N=548
Sweets consumed at least weekly	0.032 [0.071] N=1,174	0.025 [0.078] N=1,051	-0.008 [0.086] N=900	-0.023 [0.094] N=725	-0.151 [0.124] N=548
Snack foods consumed at least weekly	0.052 [0.090] N=1,174	0.047 [0.098] N=1,051	0.061 [0.111] N=900	0.112 [0.123] N=725	0.065 [0.167] N=548
Packaged soft drinks consumed at least weekly	-0.041 [0.078] N=1,174	-0.045 [0.084] N=1,051	0.012 [0.096] N=900	0.026 [0.109] N=725	0.001 [0.140] N=548
Grains consumed at least weekly	0.035 [0.043] N=1,174	0.034 [0.047] N=1,051	0.035 [0.055] N=900	0.019 [0.065] N=725	-0.042 [0.087] N=548
Child was ever breastfed	0.022 [0.038] N=1,175	0.025 [0.040] N=1,052	0.026 [0.044] N=901	0.023 [0.048] N=726	0.023 [0.066] N=549

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.947 P=1,275	-0.010 [0.029] N=1,350	-0.010 [0.030] N=1,227	-0.010 [0.034] N=1,076	-0.019 [0.034] N=901	-0.023 [0.039] N=724
Vegetables consumed at least weekly	0.904 P=1,220	0.011 [0.033] N=1,350	0.019 [0.035] N=1,227	0.031 [0.038] N=1,076	0.021 [0.042] N=901	0.009 [0.048] N=724
Meat fish or poultry consumed at least weekly	0.789 P=1,076	**0.072* [0.044] N=1,350**	**0.084* [0.045] N=1,227**	**0.088* [0.048] N=1,076**	**0.107** [0.049] N=901**	**0.095* [0.057] N=724**
Legumes consumed at least weekly	0.799 P=1,086	-0.021 [0.047] N=1,350	-0.000 [0.050] N=1,227	-0.007 [0.055] N=1,076	-0.029 [0.059] N=901	-0.020 [0.071] N=724
Animal products consumed at least weekly	0.915 P=1,237	0.007 [0.036] N=1,350	0.012 [0.038] N=1,227	0.021 [0.040] N=1,076	0.005 [0.042] N=901	0.028 [0.048] N=724
Sweets consumed at least weekly	0.726 P=980	0.025 [0.053] N=1,350	0.020 [0.055] N=1,227	0.004 [0.061] N=1,076	-0.001 [0.065] N=901	-0.045 [0.077] N=724
Snack foods consumed at least weekly	0.588 P=788	0.015 [0.058] N=1,350	0.011 [0.062] N=1,227	0.019 [0.065] N=1,076	0.034 [0.068] N=901	-0.006 [0.079] N=724
Packaged soft drinks consumed at least weekly	0.356 P=477	-0.006 [0.049] N=1,350	-0.011 [0.051] N=1,227	0.028 [0.054] N=1,076	0.037 [0.059] N=901	0.030 [0.069] N=724
Grains consumed at least weekly	0.927 P=1,250	0.051 [0.039] N=1,350	0.053 [0.041] N=1,227	0.058 [0.044] N=1,076	0.059 [0.049] N=901	0.053 [0.056] N=724
Child was ever breastfed	0.963 P=1,301	-0.010 [0.020] N=1,351	-0.012 [0.021] N=1,228	-0.013 [0.022] N=1,077	-0.016 [0.023] N=902	-0.022 [0.027] N=725

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.034 [0.038] N=1,174	-0.040 [0.041] N=1,051	-0.045 [0.050] N=900	-0.051 [0.050] N=725	-0.069 [0.070] N=548
Vegetables consumed at least weekly	-0.035 [0.038] N=1,174	-0.028 [0.043] N=1,051	-0.019 [0.050] N=900	-0.024 [0.058] N=725	-0.061 [0.073] N=548
Meat fish or poultry consumed at least weekly	0.030 [0.065] N=1,174	0.040 [0.069] N=1,051	0.031 [0.076] N=900	0.071 [0.082] N=725	0.027 [0.103] N=548
Legumes consumed at least weekly	-0.031 [0.055] N=1,174	0.001 [0.059] N=1,051	-0.002 [0.072] N=900	-0.019 [0.082] N=725	0.037 [0.090] N=548
Animal products consumed at least weekly	-0.040 [0.043] N=1,174	-0.031 [0.047] N=1,051	-0.024 [0.051] N=900	-0.066 [0.055] N=725	-0.044 [0.066] N=548
Sweets consumed at least weekly	0.034 [0.071] N=1,174	0.028 [0.078] N=1,051	-0.002 [0.087] N=900	-0.015 [0.095] N=725	-0.139 [0.124] N=548
Snack foods consumed at least weekly	0.053 [0.089] N=1,174	0.047 [0.098] N=1,051	0.064 [0.110] N=900	0.119 [0.121] N=725	0.069 [0.165] N=548
Packaged soft drinks consumed at least weekly	-0.041 [0.078] N=1,174	-0.046 [0.085] N=1,051	0.015 [0.095] N=900	0.030 [0.108] N=725	0.009 [0.140] N=548
Grains consumed at least weekly	0.035 [0.043] N=1,174	0.034 [0.046] N=1,051	0.035 [0.054] N=900	0.016 [0.064] N=725	-0.046 [0.086] N=548
Child was ever breastfed	0.022 [0.038] N=1,175	0.021 [0.040] N=1,052	0.025 [0.045] N=901	0.029 [0.050] N=726	0.026 [0.069] N=549

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.