Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Youngest Child

Age reference: 10 October 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.149	**0.441* [0.245] N=1,411**	**0.477* [0.263] N=1,306**	0.456 [0.285] N=1,172	0.479 [0.295] N=998	0.184 [0.324] N=825
WHO weight-for-age z score	0.232	-0.075 [0.136] N=1,610	-0.047 [0.147] N=1,486	-0.043 [0.151] N=1,333	-0.084 [0.168] N=1,134	-0.287 [0.181] N=931
WHO weight-for-length/height z score, 0-60 months	0.385	-0.047 [0.211] N=1,390	-0.107 [0.224] N=1,292	-0.211 [0.255] N=1,162	-0.285 [0.285] N=991	-0.244 [0.310] N=815
WHO BMI-for-age z score	0.348	-0.045 [0.242] N=1,356	-0.037 [0.262] N=1,260	-0.157 [0.299] N=1,130	-0.332 [0.330] N=962	-0.287 [0.360] N=795
Stunted: HAZ below -2	0.213 P=289	-0.062 [0.042] N=1,411	-0.071 [0.045] N=1,306	-0.047 [0.047] N=1,172	-0.035 [0.050] N=998	-0.011 [0.057] N=825
Underweight: WAZ below -2	0.053 P=90	0.025 [0.020] N=1,610	0.018 [0.021] N=1,486	0.019 [0.022] N=1,333	0.022 [0.024] N=1,134	0.048 [0.032] N=931
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=158	-0.024 [0.037] N=1,390	-0.018 [0.037] N=1,292	-0.016 [0.040] N=1,162	-0.007 [0.043] N=991	-0.028 [0.045] N=815
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=269	-0.002 [0.043] N=1,390	-0.013 [0.046] N=1,292	-0.020 [0.051] N=1,162	-0.014 [0.059] N=991	-0.004 [0.064] N=815
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.007 [0.031] N=1,390	-0.022 [0.033] N=1,292	-0.018 [0.035] N=1,162	-0.026 [0.039] N=991	-0.026 [0.043] N=815
CIAF: any stunting, underweight, or wasting, 0-60 months	0.293 P=389	**-0.079* [0.043] N=1,348**	**-0.083* [0.043] N=1,253**	-0.068 [0.046] N=1,127	-0.048 [0.046] N=959	-0.045 [0.054] N=791
BMI-for-age z score below -2	0.123 P=170	-0.006 [0.041] N=1,356	-0.003 [0.043] N=1,260	0.009 [0.047] N=1,130	0.047 [0.050] N=962	0.037 [0.055] N=795

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.387 [0.334] N=1,193	0.453 [0.379] N=1,088	0.491 [0.439] N=954	0.682 [0.482] N=780	-0.123 [0.636] N=607
WHO weight-for-age z score	-0.149 [0.183] N=1,370	-0.110 [0.202] N=1,246	-0.107 [0.206] N=1,093	-0.209 [0.264] N=894	**-0.880** [0.342] N=691**
WHO weight-for-length/height z score, 0-60 months	-0.012 [0.248] N=1,173	-0.099 [0.273] N=1,075	-0.389 [0.351] N=945	-0.646 [0.409] N=774	-0.793 [0.483] N=598
WHO BMI-for-age z score	0.106 [0.270] N=1,148	0.159 [0.304] N=1,052	-0.134 [0.395] N=922	-0.585 [0.453] N=754	-0.679 [0.536] N=587
Stunted: HAZ below -2	-0.026 [0.056] N=1,193	-0.038 [0.061] N=1,088	0.007 [0.066] N=954	0.039 [0.073] N=780	0.142 [0.109] N=607
Underweight: WAZ below -2	0.041 [0.026] N=1,370	0.031 [0.029] N=1,246	0.036 [0.030] N=1,093	0.055 [0.036] N=894	**0.159*** [0.057] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.011 [0.051] N=1,173	0.002 [0.053] N=1,075	0.019 [0.062] N=945	0.050 [0.073] N=774	0.020 [0.090] N=598
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.050] N=1,173	-0.023 [0.054] N=1,075	-0.053 [0.063] N=945	-0.060 [0.078] N=774	-0.079 [0.086] N=598
Obese: WLZ/WHZ above +3, 0-60 months	-0.001 [0.036] N=1,173	-0.024 [0.040] N=1,075	-0.013 [0.046] N=945	-0.029 [0.054] N=774	-0.039 [0.058] N=598
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.017 [0.064] N=1,138	-0.016 [0.066] N=1,043	0.031 [0.075] N=917	0.103 [0.082] N=749	0.185 [0.130] N=581
BMI-for-age z score below -2	-0.028 [0.051] N=1,148	-0.028 [0.052] N=1,052	0.001 [0.063] N=922	0.079 [0.076] N=754	0.067 [0.092] N=587

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.149	**0.441* [0.245] N=1,411**	**0.477* [0.263] N=1,306**	0.458 [0.285] N=1,172	0.479 [0.295] N=998	0.184 [0.325] N=825
WHO weight-for-age z score	0.232	-0.078 [0.136] N=1,610	-0.049 [0.147] N=1,486	-0.043 [0.152] N=1,333	-0.084 [0.168] N=1,134	-0.288 [0.181] N=931
WHO weight-for-length/height z score, 0-60 months	0.385	-0.048 [0.211] N=1,390	-0.109 [0.224] N=1,292	-0.211 [0.255] N=1,162	-0.287 [0.285] N=991	-0.247 [0.310] N=815
WHO BMI-for-age z score	0.348	-0.046 [0.242] N=1,356	-0.040 [0.262] N=1,260	-0.158 [0.299] N=1,130	-0.335 [0.330] N=962	-0.295 [0.359] N=795
Stunted: HAZ below -2	0.213 P=289	-0.062 [0.042] N=1,411	-0.071 [0.044] N=1,306	-0.048 [0.046] N=1,172	-0.035 [0.049] N=998	-0.010 [0.057] N=825
Underweight: WAZ below -2	0.053 P=90	0.026 [0.020] N=1,610	0.018 [0.021] N=1,486	0.019 [0.022] N=1,333	0.022 [0.024] N=1,134	0.048 [0.032] N=931
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=158	-0.024 [0.037] N=1,390	-0.017 [0.038] N=1,292	-0.016 [0.041] N=1,162	-0.006 [0.043] N=991	-0.027 [0.046] N=815
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=269	-0.002 [0.043] N=1,390	-0.012 [0.046] N=1,292	-0.020 [0.051] N=1,162	-0.014 [0.059] N=991	-0.004 [0.064] N=815
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.007 [0.031] N=1,390	-0.022 [0.033] N=1,292	-0.018 [0.035] N=1,162	-0.026 [0.039] N=991	-0.025 [0.043] N=815
CIAF: any stunting, underweight, or wasting, 0-60 months	0.293 P=389	**-0.078* [0.044] N=1,348**	**-0.082* [0.043] N=1,253**	-0.069 [0.046] N=1,127	-0.048 [0.047] N=959	-0.043 [0.054] N=791
BMI-for-age z score below -2	0.123 P=170	-0.006 [0.041] N=1,356	-0.002 [0.043] N=1,260	0.010 [0.047] N=1,130	0.049 [0.050] N=962	0.040 [0.054] N=795

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.385 [0.333] N=1,193	0.450 [0.378] N=1,088	0.484 [0.437] N=954	0.658 [0.480] N=780	-0.148 [0.638] N=607
WHO weight-for-age z score	-0.147 [0.182] N=1,370	-0.105 [0.201] N=1,246	-0.098 [0.206] N=1,093	-0.200 [0.264] N=894	**-0.865** [0.340] N=691**
WHO weight-for-length/height z score, 0-60 months	-0.013 [0.249] N=1,173	-0.101 [0.273] N=1,075	-0.383 [0.350] N=945	-0.641 [0.408] N=774	-0.793 [0.492] N=598
WHO BMI-for-age z score	0.106 [0.270] N=1,148	0.159 [0.305] N=1,052	-0.124 [0.393] N=922	-0.576 [0.453] N=754	-0.680 [0.547] N=587
Stunted: HAZ below -2	-0.026 [0.056] N=1,193	-0.039 [0.061] N=1,088	0.005 [0.066] N=954	0.040 [0.072] N=780	0.144 [0.109] N=607
Underweight: WAZ below -2	0.041 [0.026] N=1,370	0.031 [0.029] N=1,246	0.035 [0.030] N=1,093	0.054 [0.036] N=894	**0.159*** [0.057] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.010 [0.051] N=1,173	0.003 [0.053] N=1,075	0.016 [0.063] N=945	0.048 [0.074] N=774	0.020 [0.092] N=598
Overweight: WLZ/WHZ above +2, 0-60 months	-0.006 [0.050] N=1,173	-0.023 [0.054] N=1,075	-0.053 [0.063] N=945	-0.060 [0.078] N=774	-0.079 [0.086] N=598
Obese: WLZ/WHZ above +3, 0-60 months	-0.000 [0.037] N=1,173	-0.024 [0.040] N=1,075	-0.014 [0.046] N=945	-0.029 [0.055] N=774	-0.039 [0.058] N=598
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.017 [0.064] N=1,138	-0.017 [0.066] N=1,043	0.024 [0.074] N=917	0.099 [0.082] N=749	0.182 [0.131] N=581
BMI-for-age z score below -2	-0.028 [0.051] N=1,148	-0.028 [0.053] N=1,052	-0.002 [0.063] N=922	0.077 [0.076] N=754	0.067 [0.094] N=587

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.149	**0.408* [0.245] N=1,411**	0.423 [0.263] N=1,306	0.414 [0.284] N=1,172	0.437 [0.290] N=998	0.107 [0.315] N=825
WHO weight-for-age z score	0.232	-0.127 [0.132] N=1,610	-0.102 [0.143] N=1,486	-0.092 [0.148] N=1,333	-0.130 [0.164] N=1,134	**-0.346* [0.175] N=931**
WHO weight-for-length/height z score, 0-60 months	0.385	-0.072 [0.213] N=1,390	-0.126 [0.227] N=1,292	-0.220 [0.256] N=1,162	-0.289 [0.285] N=991	-0.265 [0.309] N=815
WHO BMI-for-age z score	0.348	-0.060 [0.244] N=1,356	-0.048 [0.264] N=1,260	-0.161 [0.299] N=1,130	-0.334 [0.328] N=962	-0.300 [0.358] N=795
Stunted: HAZ below -2	0.213 P=289	-0.055 [0.042] N=1,411	-0.059 [0.044] N=1,306	-0.038 [0.046] N=1,172	-0.027 [0.048] N=998	0.004 [0.055] N=825
Underweight: WAZ below -2	0.053 P=90	0.030 [0.019] N=1,610	0.025 [0.021] N=1,486	0.026 [0.021] N=1,333	0.030 [0.024] N=1,134	**0.057* [0.031] N=931**
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=158	-0.016 [0.037] N=1,390	-0.007 [0.038] N=1,292	-0.007 [0.041] N=1,162	-0.001 [0.043] N=991	-0.021 [0.046] N=815
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=269	0.001 [0.043] N=1,390	-0.007 [0.046] N=1,292	-0.015 [0.051] N=1,162	-0.008 [0.059] N=991	-0.001 [0.064] N=815
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.006 [0.031] N=1,390	-0.020 [0.033] N=1,292	-0.016 [0.035] N=1,162	-0.024 [0.039] N=991	-0.026 [0.042] N=815
CIAF: any stunting, underweight, or wasting, 0-60 months	0.293 P=389	-0.061 [0.043] N=1,348	-0.058 [0.043] N=1,253	-0.049 [0.045] N=1,127	-0.032 [0.046] N=959	-0.018 [0.054] N=791
BMI-for-age z score below -2	0.123 P=170	0.000 [0.041] N=1,356	0.006 [0.043] N=1,260	0.017 [0.047] N=1,130	0.053 [0.050] N=962	0.044 [0.055] N=795

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.352 [0.335] N=1,193	0.375 [0.382] N=1,088	0.426 [0.443] N=954	0.616 [0.481] N=780	-0.281 [0.625] N=607
WHO weight-for-age z score	-0.185 [0.178] N=1,370	-0.149 [0.197] N=1,246	-0.132 [0.205] N=1,093	-0.224 [0.266] N=894	**-0.911*** [0.337] N=691**
WHO weight-for-length/height z score, 0-60 months	-0.023 [0.254] N=1,173	-0.095 [0.282] N=1,075	-0.361 [0.356] N=945	-0.632 [0.414] N=774	-0.822 [0.501] N=598
WHO BMI-for-age z score	0.098 [0.274] N=1,148	0.163 [0.312] N=1,052	-0.111 [0.397] N=922	-0.587 [0.459] N=754	-0.709 [0.559] N=587
Stunted: HAZ below -2	-0.021 [0.056] N=1,193	-0.023 [0.062] N=1,088	0.015 [0.067] N=954	0.043 [0.072] N=780	0.163 [0.106] N=607
Underweight: WAZ below -2	**0.045* [0.026] N=1,370**	0.037 [0.029] N=1,246	0.041 [0.029] N=1,093	**0.059* [0.035] N=894**	**0.164*** [0.055] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.003 [0.051] N=1,173	0.013 [0.054] N=1,075	0.021 [0.063] N=945	0.053 [0.075] N=774	0.032 [0.095] N=598
Overweight: WLZ/WHZ above +2, 0-60 months	-0.001 [0.051] N=1,173	-0.013 [0.056] N=1,075	-0.047 [0.064] N=945	-0.055 [0.077] N=774	-0.070 [0.084] N=598
Obese: WLZ/WHZ above +3, 0-60 months	0.001 [0.037] N=1,173	-0.020 [0.041] N=1,075	-0.012 [0.047] N=945	-0.027 [0.055] N=774	-0.043 [0.059] N=598
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.002 [0.064] N=1,138	0.010 [0.066] N=1,043	0.040 [0.074] N=917	0.106 [0.082] N=749	**0.220* [0.125] N=581**
BMI-for-age z score below -2	-0.020 [0.051] N=1,148	-0.016 [0.054] N=1,052	0.006 [0.064] N=922	0.084 [0.078] N=754	0.081 [0.099] N=587

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.149	0.334 [0.239] N=1,411	0.351 [0.257] N=1,306	0.347 [0.277] N=1,172	0.359 [0.283] N=998	0.006 [0.307] N=825
WHO weight-for-age z score	0.232	-0.144 [0.132] N=1,610	-0.124 [0.143] N=1,486	-0.113 [0.148] N=1,333	-0.161 [0.166] N=1,134	**-0.387** [0.180] N=931**
WHO weight-for-length/height z score, 0-60 months	0.385	-0.034 [0.210] N=1,390	-0.092 [0.224] N=1,292	-0.196 [0.254] N=1,162	-0.271 [0.287] N=991	-0.245 [0.313] N=815
WHO BMI-for-age z score	0.348	-0.022 [0.239] N=1,356	-0.011 [0.259] N=1,260	-0.130 [0.295] N=1,130	-0.304 [0.331] N=962	-0.264 [0.364] N=795
Stunted: HAZ below -2	0.213 P=289	-0.040 [0.042] N=1,411	-0.044 [0.044] N=1,306	-0.022 [0.045] N=1,172	-0.007 [0.047] N=998	0.028 [0.053] N=825
Underweight: WAZ below -2	0.053 P=90	**0.033* [0.019] N=1,610**	0.027 [0.021] N=1,486	0.029 [0.022] N=1,333	0.033 [0.024] N=1,134	**0.061* [0.032] N=931**
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=158	-0.019 [0.037] N=1,390	-0.011 [0.038] N=1,292	-0.010 [0.041] N=1,162	-0.002 [0.044] N=991	-0.025 [0.046] N=815
Overweight: WLZ/WHZ above +2, 0-60 months	0.200 P=269	0.007 [0.043] N=1,390	-0.002 [0.046] N=1,292	-0.011 [0.051] N=1,162	-0.005 [0.059] N=991	-0.000 [0.064] N=815
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.004 [0.031] N=1,390	-0.019 [0.033] N=1,292	-0.015 [0.035] N=1,162	-0.023 [0.039] N=991	-0.025 [0.043] N=815
CIAF: any stunting, underweight, or wasting, 0-60 months	0.293 P=389	-0.047 [0.044] N=1,348	-0.044 [0.043] N=1,253	-0.031 [0.046] N=1,127	-0.007 [0.047] N=959	0.008 [0.055] N=791
BMI-for-age z score below -2	0.123 P=170	-0.002 [0.041] N=1,356	0.002 [0.043] N=1,260	0.014 [0.047] N=1,130	0.052 [0.050] N=962	0.039 [0.055] N=795

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.314 [0.335] N=1,193	0.363 [0.376] N=1,088	0.432 [0.438] N=954	0.616 [0.471] N=780	-0.359 [0.607] N=607
WHO weight-for-age z score	-0.207 [0.179] N=1,370	-0.174 [0.197] N=1,246	-0.161 [0.205] N=1,093	-0.257 [0.269] N=894	**-0.931*** [0.337] N=691**
WHO weight-for-length/height z score, 0-60 months	-0.028 [0.251] N=1,173	-0.105 [0.276] N=1,075	-0.382 [0.352] N=945	-0.622 [0.408] N=774	-0.732 [0.499] N=598
WHO BMI-for-age z score	0.101 [0.272] N=1,148	0.163 [0.306] N=1,052	-0.114 [0.395] N=922	-0.533 [0.448] N=754	-0.577 [0.545] N=587
Stunted: HAZ below -2	-0.017 [0.057] N=1,193	-0.025 [0.062] N=1,088	0.009 [0.067] N=954	0.033 [0.074] N=780	0.158 [0.108] N=607
Underweight: WAZ below -2	**0.048* [0.026] N=1,370**	0.039 [0.029] N=1,246	0.043 [0.029] N=1,093	0.060 [0.037] N=894	**0.167*** [0.058] N=691**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.001 [0.051] N=1,173	0.012 [0.054] N=1,075	0.022 [0.063] N=945	0.054 [0.076] N=774	0.029 [0.095] N=598
Overweight: WLZ/WHZ above +2, 0-60 months	0.001 [0.051] N=1,173	-0.013 [0.054] N=1,075	-0.045 [0.063] N=945	-0.048 [0.075] N=774	-0.053 [0.082] N=598
Obese: WLZ/WHZ above +3, 0-60 months	0.005 [0.038] N=1,173	-0.018 [0.041] N=1,075	-0.009 [0.047] N=945	-0.023 [0.055] N=774	-0.031 [0.058] N=598
CIAF: any stunting, underweight, or wasting, 0-60 months	0.006 [0.065] N=1,138	0.011 [0.067] N=1,043	0.041 [0.076] N=917	0.105 [0.086] N=749	**0.224* [0.129] N=581**
BMI-for-age z score below -2	-0.019 [0.051] N=1,148	-0.022 [0.054] N=1,052	0.002 [0.064] N=922	0.079 [0.078] N=754	0.075 [0.098] N=587

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.160	**0.439* [0.255] N=1,319**	0.421 [0.276] N=1,227	0.398 [0.295] N=1,098	0.424 [0.302] N=939	0.158 [0.322] N=776
WHO weight-for-age z score	0.229	-0.040 [0.141] N=1,503	-0.012 [0.150] N=1,394	0.002 [0.156] N=1,248	-0.016 [0.171] N=1,065	-0.191 [0.194] N=876
WHO weight-for-length/height z score, 0-60 months	0.392	0.064 [0.212] N=1,298	0.023 [0.224] N=1,214	-0.092 [0.254] N=1,090	-0.150 [0.281] N=934	-0.069 [0.303] N=768
WHO BMI-for-age z score	0.358	0.088 [0.243] N=1,265	0.089 [0.260] N=1,182	-0.025 [0.294] N=1,059	-0.165 [0.321] N=906	-0.072 [0.345] N=748
Stunted: HAZ below -2	0.214 P=270	-0.054 [0.045] N=1,319	-0.058 [0.048] N=1,227	-0.036 [0.050] N=1,098	-0.023 [0.052] N=939	0.005 [0.059] N=776
Underweight: WAZ below -2	0.054 P=86	0.025 [0.021] N=1,503	0.017 [0.022] N=1,394	0.018 [0.023] N=1,248	0.021 [0.025] N=1,065	0.043 [0.034] N=876
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=147	-0.030 [0.038] N=1,298	-0.026 [0.038] N=1,214	-0.020 [0.041] N=1,090	-0.013 [0.043] N=934	-0.033 [0.045] N=768
Overweight: WLZ/WHZ above +2, 0-60 months	0.201 P=252	0.009 [0.045] N=1,298	0.001 [0.048] N=1,214	-0.005 [0.053] N=1,090	0.002 [0.061] N=934	0.017 [0.068] N=768
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=122	-0.012 [0.033] N=1,298	-0.023 [0.035] N=1,214	-0.019 [0.037] N=1,090	-0.023 [0.041] N=934	-0.017 [0.045] N=768
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=364	-0.067 [0.047] N=1,260	-0.070 [0.046] N=1,178	-0.059 [0.049] N=1,057	-0.044 [0.051] N=904	-0.032 [0.059] N=745
BMI-for-age z score below -2	0.121 P=157	-0.020 [0.039] N=1,265	-0.019 [0.039] N=1,182	-0.007 [0.043] N=1,059	0.025 [0.046] N=906	0.010 [0.048] N=748

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.442 [0.340] N=1,114	0.451 [0.391] N=1,022	0.480 [0.453] N=893	0.682 [0.488] N=734	-0.003 [0.650] N=571
WHO weight-for-age z score	-0.198 [0.190] N=1,278	-0.167 [0.205] N=1,169	-0.164 [0.215] N=1,023	-0.237 [0.267] N=840	**-0.833** [0.376] N=651**
WHO weight-for-length/height z score, 0-60 months	-0.052 [0.254] N=1,094	-0.147 [0.269] N=1,010	-0.475 [0.341] N=886	-0.690 [0.418] N=730	-0.766 [0.500] N=564
WHO BMI-for-age z score	0.084 [0.272] N=1,070	0.081 [0.296] N=987	-0.210 [0.378] N=864	-0.602 [0.454] N=711	-0.621 [0.544] N=553
Stunted: HAZ below -2	-0.028 [0.060] N=1,114	-0.040 [0.065] N=1,022	0.002 [0.070] N=893	0.031 [0.075] N=734	0.113 [0.117] N=571
Underweight: WAZ below -2	**0.049* [0.027] N=1,278**	0.040 [0.030] N=1,169	0.045 [0.031] N=1,023	**0.065* [0.035] N=840**	**0.163*** [0.060] N=651**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.010 [0.052] N=1,094	0.001 [0.052] N=1,010	0.019 [0.061] N=886	0.034 [0.074] N=730	-0.015 [0.091] N=564
Overweight: WLZ/WHZ above +2, 0-60 months	-0.009 [0.053] N=1,094	-0.032 [0.055] N=1,010	-0.065 [0.066] N=886	-0.083 [0.083] N=730	-0.081 [0.093] N=564
Obese: WLZ/WHZ above +3, 0-60 months	-0.019 [0.039] N=1,094	-0.038 [0.043] N=1,010	-0.034 [0.048] N=886	-0.046 [0.057] N=730	-0.054 [0.059] N=564
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.002 [0.069] N=1,063	-0.005 [0.067] N=981	0.031 [0.077] N=860	0.075 [0.084] N=707	0.126 [0.138] N=548
BMI-for-age z score below -2	-0.020 [0.050] N=1,070	-0.019 [0.050] N=987	0.008 [0.060] N=864	0.071 [0.075] N=711	0.049 [0.094] N=553

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.160	**0.423* [0.254] N=1,319**	0.407 [0.275] N=1,227	0.374 [0.296] N=1,098	0.387 [0.309] N=939	0.095 [0.338] N=776
WHO weight-for-age z score	0.229	-0.053 [0.140] N=1,503	-0.023 [0.150] N=1,394	-0.013 [0.155] N=1,248	-0.039 [0.170] N=1,065	-0.229 [0.191] N=876
WHO weight-for-length/height z score, 0-60 months	0.392	0.050 [0.212] N=1,298	0.007 [0.224] N=1,214	-0.107 [0.255] N=1,090	-0.162 [0.286] N=934	-0.086 [0.313] N=768
WHO BMI-for-age z score	0.358	0.075 [0.243] N=1,265	0.073 [0.259] N=1,182	-0.041 [0.295] N=1,059	-0.183 [0.329] N=906	-0.102 [0.361] N=748
Stunted: HAZ below -2	0.214 P=270	-0.052 [0.045] N=1,319	-0.057 [0.047] N=1,227	-0.033 [0.049] N=1,098	-0.019 [0.052] N=939	0.006 [0.059] N=776
Underweight: WAZ below -2	0.054 P=86	0.026 [0.021] N=1,503	0.018 [0.022] N=1,394	0.019 [0.023] N=1,248	0.023 [0.025] N=1,065	0.047 [0.034] N=876
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=147	-0.029 [0.039] N=1,298	-0.025 [0.039] N=1,214	-0.021 [0.042] N=1,090	-0.016 [0.045] N=934	-0.040 [0.048] N=768
Overweight: WLZ/WHZ above +2, 0-60 months	0.201 P=252	0.009 [0.046] N=1,298	-0.001 [0.048] N=1,214	-0.008 [0.054] N=1,090	-0.004 [0.062] N=934	0.013 [0.068] N=768
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=122	-0.012 [0.033] N=1,298	-0.022 [0.035] N=1,214	-0.019 [0.037] N=1,090	-0.023 [0.041] N=934	-0.018 [0.046] N=768
CIAF: any stunting, underweight, or wasting, 0-60 months	0.294 P=364	-0.065 [0.046] N=1,260	-0.068 [0.045] N=1,178	-0.057 [0.048] N=1,057	-0.043 [0.050] N=904	-0.038 [0.058] N=745
BMI-for-age z score below -2	0.121 P=157	-0.018 [0.039] N=1,265	-0.017 [0.040] N=1,182	-0.007 [0.044] N=1,059	0.024 [0.047] N=906	0.007 [0.051] N=748

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.471 [0.343] N=1,114	0.470 [0.391] N=1,022	0.496 [0.449] N=893	0.679 [0.499] N=734	-0.046 [0.677] N=571
WHO weight-for-age z score	-0.171 [0.188] N=1,278	-0.133 [0.203] N=1,169	-0.141 [0.212] N=1,023	-0.227 [0.266] N=840	**-0.855** [0.378] N=651**
WHO weight-for-length/height z score, 0-60 months	-0.031 [0.252] N=1,094	-0.109 [0.268] N=1,010	-0.435 [0.340] N=886	-0.674 [0.418] N=730	-0.763 [0.500] N=564
WHO BMI-for-age z score	0.115 [0.270] N=1,070	0.131 [0.293] N=987	-0.163 [0.374] N=864	-0.562 [0.451] N=711	-0.604 [0.540] N=553
Stunted: HAZ below -2	-0.033 [0.059] N=1,114	-0.042 [0.064] N=1,022	0.001 [0.069] N=893	0.027 [0.075] N=734	0.110 [0.116] N=571
Underweight: WAZ below -2	**0.046* [0.027] N=1,278**	0.036 [0.030] N=1,169	0.042 [0.030] N=1,023	**0.063* [0.034] N=840**	**0.165*** [0.058] N=651**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.012 [0.052] N=1,094	-0.003 [0.051] N=1,010	0.016 [0.060] N=886	0.033 [0.072] N=730	-0.019 [0.087] N=564
Overweight: WLZ/WHZ above +2, 0-60 months	-0.009 [0.054] N=1,094	-0.028 [0.056] N=1,010	-0.059 [0.066] N=886	-0.075 [0.083] N=730	-0.082 [0.093] N=564
Obese: WLZ/WHZ above +3, 0-60 months	-0.021 [0.039] N=1,094	-0.039 [0.043] N=1,010	-0.033 [0.049] N=886	-0.046 [0.058] N=730	-0.051 [0.059] N=564
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.008 [0.067] N=1,063	-0.010 [0.066] N=981	0.028 [0.074] N=860	0.071 [0.082] N=707	0.119 [0.134] N=548
BMI-for-age z score below -2	-0.021 [0.050] N=1,070	-0.022 [0.050] N=987	0.005 [0.060] N=864	0.070 [0.075] N=711	0.045 [0.092] N=553

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.038	-0.072 [0.317] N=1,272	-0.177 [0.324] N=1,148	-0.199 [0.338] N=1,004	-0.442 [0.373] N=850	-0.497 [0.414] N=685
WHO weight-for-age z score	0.252	0.105 [0.141] N=1,439	0.138 [0.148] N=1,309	0.022 [0.148] N=1,143	0.097 [0.155] N=959	0.100 [0.183] N=768
WHO weight-for-length/height z score, 0-60 months	0.381	0.138 [0.206] N=1,260	0.165 [0.216] N=1,140	0.148 [0.235] N=998	0.381 [0.248] N=841	0.244 [0.279] N=679
WHO BMI-for-age z score	0.316	0.130 [0.240] N=1,219	0.172 [0.253] N=1,104	0.105 [0.276] N=968	0.308 [0.301] N=816	0.229 [0.337] N=662
Stunted: HAZ below -2	0.198 P=253	-0.003 [0.052] N=1,272	0.010 [0.053] N=1,148	0.006 [0.056] N=1,004	0.028 [0.061] N=850	0.016 [0.068] N=685
Underweight: WAZ below -2	0.049 P=75	0.018 [0.024] N=1,439	0.011 [0.024] N=1,309	0.017 [0.026] N=1,143	0.005 [0.028] N=959	-0.010 [0.032] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=141	0.026 [0.036] N=1,260	0.020 [0.038] N=1,140	0.037 [0.041] N=998	-0.040 [0.039] N=841	-0.025 [0.041] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=248	**0.095** [0.044] N=1,260**	**0.098** [0.047] N=1,140**	**0.106** [0.051] N=998**	**0.131** [0.058] N=841**	**0.115* [0.063] N=679**
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=123	0.037 [0.038] N=1,260	0.035 [0.038] N=1,140	0.027 [0.043] N=998	0.038 [0.050] N=841	0.004 [0.049] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.283 P=347	0.029 [0.056] N=1,221	0.027 [0.057] N=1,101	0.051 [0.061] N=965	0.001 [0.065] N=816	0.013 [0.068] N=659
BMI-for-age z score below -2	0.125 P=151	0.029 [0.043] N=1,219	0.022 [0.047] N=1,104	0.029 [0.052] N=968	-0.024 [0.051] N=816	-0.002 [0.054] N=662

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.242 [0.431] N=1,101	0.132 [0.464] N=977	0.155 [0.527] N=833	-0.381 [0.664] N=679	-0.398 [0.800] N=514
WHO weight-for-age z score	0.129 [0.194] N=1,253	0.189 [0.208] N=1,123	-0.021 [0.213] N=957	0.134 [0.228] N=773	0.192 [0.301] N=582
WHO weight-for-length/height z score, 0-60 months	-0.063 [0.284] N=1,093	-0.032 [0.307] N=973	-0.070 [0.358] N=831	0.466 [0.396] N=674	0.054 [0.458] N=512
WHO BMI-for-age z score	-0.235 [0.304] N=1,061	-0.202 [0.335] N=946	-0.372 [0.389] N=810	0.047 [0.454] N=658	-0.284 [0.522] N=504
Stunted: HAZ below -2	0.006 [0.086] N=1,101	0.020 [0.092] N=977	0.016 [0.103] N=833	0.080 [0.128] N=679	0.033 [0.155] N=514
Underweight: WAZ below -2	0.020 [0.036] N=1,253	0.007 [0.037] N=1,123	0.017 [0.042] N=957	-0.013 [0.044] N=773	-0.054 [0.049] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	**0.093* [0.048] N=1,093**	0.088 [0.054] N=973	**0.134** [0.065] N=831**	-0.019 [0.062] N=674	0.025 [0.067] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.108* [0.055] N=1,093**	**0.107* [0.059] N=973**	**0.134** [0.067] N=831**	**0.196** [0.084] N=674**	0.128 [0.094] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.082** [0.041] N=1,093**	**0.081* [0.042] N=973**	0.081 [0.049] N=831	**0.117* [0.068] N=674**	0.015 [0.079] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.090 [0.088] N=1,058	0.082 [0.094] N=938	0.148 [0.108] N=802	0.064 [0.128] N=653	0.071 [0.144] N=496
BMI-for-age z score below -2	**0.107** [0.048] N=1,061**	**0.102* [0.056] N=946**	**0.135** [0.067] N=810**	0.025 [0.065] N=658	0.098 [0.075] N=504

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.038	-0.079 [0.316] N=1,272	-0.183 [0.323] N=1,148	-0.205 [0.337] N=1,004	-0.452 [0.370] N=850	-0.505 [0.412] N=685
WHO weight-for-age z score	0.252	0.111 [0.140] N=1,439	0.144 [0.147] N=1,309	0.028 [0.146] N=1,143	0.100 [0.154] N=959	0.100 [0.183] N=768
WHO weight-for-length/height z score, 0-60 months	0.381	0.145 [0.207] N=1,260	0.172 [0.216] N=1,140	0.155 [0.234] N=998	0.386 [0.247] N=841	0.252 [0.277] N=679
WHO BMI-for-age z score	0.316	0.135 [0.240] N=1,219	0.176 [0.253] N=1,104	0.109 [0.276] N=968	0.310 [0.301] N=816	0.230 [0.338] N=662
Stunted: HAZ below -2	0.198 P=253	-0.003 [0.052] N=1,272	0.010 [0.053] N=1,148	0.005 [0.056] N=1,004	0.029 [0.061] N=850	0.017 [0.068] N=685
Underweight: WAZ below -2	0.049 P=75	0.017 [0.024] N=1,439	0.010 [0.024] N=1,309	0.016 [0.026] N=1,143	0.004 [0.028] N=959	-0.010 [0.032] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=141	0.023 [0.036] N=1,260	0.016 [0.039] N=1,140	0.032 [0.041] N=998	-0.042 [0.039] N=841	-0.027 [0.041] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=248	**0.096** [0.045] N=1,260**	**0.098** [0.047] N=1,140**	**0.106** [0.051] N=998**	**0.130** [0.058] N=841**	**0.116* [0.063] N=679**
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=123	0.035 [0.038] N=1,260	0.032 [0.038] N=1,140	0.023 [0.043] N=998	0.036 [0.050] N=841	0.003 [0.050] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.283 P=347	0.025 [0.057] N=1,221	0.021 [0.057] N=1,101	0.044 [0.061] N=965	-0.001 [0.065] N=816	0.012 [0.069] N=659
BMI-for-age z score below -2	0.125 P=151	0.026 [0.043] N=1,219	0.019 [0.046] N=1,104	0.026 [0.051] N=968	-0.025 [0.051] N=816	-0.002 [0.054] N=662

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.236 [0.429] N=1,101	0.128 [0.462] N=977	0.154 [0.526] N=833	-0.416 [0.655] N=679	-0.455 [0.798] N=514
WHO weight-for-age z score	0.143 [0.192] N=1,253	0.205 [0.204] N=1,123	-0.005 [0.208] N=957	0.157 [0.224] N=773	0.236 [0.299] N=582
WHO weight-for-length/height z score, 0-60 months	-0.046 [0.286] N=1,093	-0.013 [0.309] N=973	-0.049 [0.359] N=831	0.499 [0.398] N=674	0.156 [0.475] N=512
WHO BMI-for-age z score	-0.215 [0.305] N=1,061	-0.180 [0.334] N=946	-0.346 [0.386] N=810	0.082 [0.455] N=658	-0.184 [0.539] N=504
Stunted: HAZ below -2	0.004 [0.087] N=1,101	0.016 [0.092] N=977	0.009 [0.104] N=833	0.079 [0.129] N=679	0.034 [0.157] N=514
Underweight: WAZ below -2	0.018 [0.036] N=1,253	0.005 [0.038] N=1,123	0.015 [0.042] N=957	-0.015 [0.045] N=773	-0.058 [0.050] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	**0.086* [0.049] N=1,093**	0.077 [0.056] N=973	**0.120* [0.067] N=831**	-0.032 [0.063] N=674	0.003 [0.071] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.109* [0.056] N=1,093**	**0.109* [0.060] N=973**	**0.132* [0.068] N=831**	**0.193** [0.085] N=674**	0.133 [0.097] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.078* [0.041] N=1,093**	**0.076* [0.042] N=973**	0.072 [0.049] N=831	0.107 [0.068] N=674	0.005 [0.079] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.079 [0.090] N=1,058	0.066 [0.096] N=938	0.125 [0.110] N=802	0.048 [0.131] N=653	0.049 [0.148] N=496
BMI-for-age z score below -2	**0.100** [0.048] N=1,061**	0.092 [0.056] N=946	**0.123* [0.066] N=810**	0.014 [0.064] N=658	0.080 [0.076] N=504

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.038	-0.089 [0.312] N=1,272	-0.181 [0.320] N=1,148	-0.198 [0.334] N=1,004	-0.443 [0.362] N=850	-0.505 [0.400] N=685
WHO weight-for-age z score	0.252	0.104 [0.141] N=1,439	0.143 [0.147] N=1,309	0.033 [0.148] N=1,143	0.103 [0.155] N=959	0.105 [0.183] N=768
WHO weight-for-length/height z score, 0-60 months	0.381	0.142 [0.208] N=1,260	0.169 [0.217] N=1,140	0.160 [0.235] N=998	0.383 [0.246] N=841	0.252 [0.276] N=679
WHO BMI-for-age z score	0.316	0.128 [0.241] N=1,219	0.168 [0.254] N=1,104	0.108 [0.275] N=968	0.303 [0.298] N=816	0.231 [0.335] N=662
Stunted: HAZ below -2	0.198 P=253	-0.001 [0.051] N=1,272	0.009 [0.053] N=1,148	0.003 [0.054] N=1,004	0.027 [0.059] N=850	0.015 [0.066] N=685
Underweight: WAZ below -2	0.049 P=75	0.018 [0.024] N=1,439	0.010 [0.025] N=1,309	0.015 [0.026] N=1,143	0.004 [0.028] N=959	-0.010 [0.032] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=141	0.024 [0.037] N=1,260	0.016 [0.039] N=1,140	0.031 [0.042] N=998	-0.042 [0.039] N=841	-0.028 [0.042] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=248	**0.097** [0.044] N=1,260**	**0.098** [0.047] N=1,140**	**0.106** [0.051] N=998**	**0.129** [0.058] N=841**	**0.114* [0.064] N=679**
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=123	0.035 [0.038] N=1,260	0.032 [0.038] N=1,140	0.023 [0.043] N=998	0.035 [0.050] N=841	0.002 [0.050] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.283 P=347	0.030 [0.056] N=1,221	0.023 [0.057] N=1,101	0.043 [0.061] N=965	-0.003 [0.065] N=816	0.009 [0.068] N=659
BMI-for-age z score below -2	0.125 P=151	0.028 [0.044] N=1,219	0.021 [0.047] N=1,104	0.027 [0.051] N=968	-0.024 [0.052] N=816	-0.001 [0.055] N=662

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.249 [0.425] N=1,101	0.141 [0.461] N=977	0.200 [0.517] N=833	-0.373 [0.648] N=679	-0.388 [0.773] N=514
WHO weight-for-age z score	0.127 [0.194] N=1,253	0.197 [0.208] N=1,123	0.002 [0.213] N=957	0.158 [0.230] N=773	0.228 [0.304] N=582
WHO weight-for-length/height z score, 0-60 months	-0.070 [0.286] N=1,093	-0.025 [0.309] N=973	-0.063 [0.357] N=831	0.480 [0.396] N=674	0.115 [0.474] N=512
WHO BMI-for-age z score	-0.236 [0.305] N=1,061	-0.192 [0.336] N=946	-0.371 [0.382] N=810	0.056 [0.448] N=658	-0.207 [0.529] N=504
Stunted: HAZ below -2	-0.002 [0.085] N=1,101	0.010 [0.091] N=977	-0.002 [0.102] N=833	0.063 [0.127] N=679	0.016 [0.155] N=514
Underweight: WAZ below -2	0.017 [0.037] N=1,253	0.002 [0.038] N=1,123	0.010 [0.043] N=957	-0.020 [0.046] N=773	-0.064 [0.052] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	**0.088* [0.050] N=1,093**	0.077 [0.057] N=973	**0.120* [0.067] N=831**	-0.031 [0.063] N=674	0.004 [0.073] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.111* [0.056] N=1,093**	**0.111* [0.061] N=973**	**0.131* [0.069] N=831**	**0.192** [0.087] N=674**	0.131 [0.101] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.076* [0.042] N=1,093**	**0.074* [0.043] N=973**	0.070 [0.049] N=831	0.104 [0.069] N=674	-0.003 [0.081] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.079 [0.090] N=1,058	0.065 [0.097] N=938	0.119 [0.112] N=802	0.038 [0.132] N=653	0.045 [0.149] N=496
BMI-for-age z score below -2	**0.100** [0.049] N=1,061**	0.092 [0.057] N=946	**0.124* [0.065] N=810**	0.015 [0.064] N=658	0.079 [0.078] N=504

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.038	-0.031 [0.311] N=1,272	-0.116 [0.319] N=1,148	-0.132 [0.335] N=1,004	-0.413 [0.365] N=850	-0.469 [0.411] N=685
WHO weight-for-age z score	0.252	0.105 [0.143] N=1,439	0.148 [0.150] N=1,309	0.044 [0.151] N=1,143	0.110 [0.159] N=959	0.116 [0.187] N=768
WHO weight-for-length/height z score, 0-60 months	0.381	0.102 [0.207] N=1,260	0.135 [0.215] N=1,140	0.123 [0.233] N=998	0.376 [0.246] N=841	0.252 [0.279] N=679
WHO BMI-for-age z score	0.316	0.092 [0.239] N=1,219	0.137 [0.251] N=1,104	0.069 [0.271] N=968	0.292 [0.295] N=816	0.237 [0.337] N=662
Stunted: HAZ below -2	0.198 P=253	-0.011 [0.051] N=1,272	-0.001 [0.052] N=1,148	-0.008 [0.055] N=1,004	0.021 [0.060] N=850	0.009 [0.066] N=685
Underweight: WAZ below -2	0.049 P=75	0.018 [0.024] N=1,439	0.010 [0.025] N=1,309	0.013 [0.027] N=1,143	0.003 [0.029] N=959	-0.015 [0.033] N=768
Wasted: WLZ/WHZ below -2, 0-60 months	0.110 P=141	0.027 [0.036] N=1,260	0.020 [0.038] N=1,140	0.037 [0.041] N=998	-0.040 [0.038] N=841	-0.029 [0.041] N=679
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=248	**0.091** [0.044] N=1,260**	**0.094** [0.046] N=1,140**	**0.103** [0.050] N=998**	**0.131** [0.057] N=841**	**0.115* [0.062] N=679**
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=123	0.034 [0.037] N=1,260	0.034 [0.037] N=1,140	0.023 [0.042] N=998	0.038 [0.048] N=841	0.004 [0.048] N=679
CIAF: any stunting, underweight, or wasting, 0-60 months	0.283 P=347	0.026 [0.056] N=1,221	0.019 [0.057] N=1,101	0.039 [0.060] N=965	-0.004 [0.064] N=816	0.008 [0.067] N=659
BMI-for-age z score below -2	0.125 P=151	0.028 [0.043] N=1,219	0.021 [0.047] N=1,104	0.029 [0.051] N=968	-0.024 [0.051] N=816	-0.005 [0.055] N=662

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.308 [0.412] N=1,101	0.225 [0.444] N=977	0.291 [0.504] N=833	-0.346 [0.635] N=679	-0.456 [0.759] N=514
WHO weight-for-age z score	0.150 [0.190] N=1,253	0.225 [0.204] N=1,123	0.044 [0.206] N=957	0.190 [0.224] N=773	0.270 [0.291] N=582
WHO weight-for-length/height z score, 0-60 months	-0.098 [0.285] N=1,093	-0.059 [0.306] N=973	-0.105 [0.353] N=831	0.479 [0.400] N=674	0.192 [0.495] N=512
WHO BMI-for-age z score	-0.277 [0.304] N=1,061	-0.238 [0.332] N=946	-0.429 [0.376] N=810	0.035 [0.446] N=658	-0.129 [0.554] N=504
Stunted: HAZ below -2	-0.011 [0.084] N=1,101	-0.004 [0.090] N=977	-0.017 [0.102] N=833	0.055 [0.128] N=679	0.019 [0.155] N=514
Underweight: WAZ below -2	0.018 [0.036] N=1,253	0.002 [0.038] N=1,123	0.008 [0.043] N=957	-0.020 [0.046] N=773	-0.069 [0.053] N=582
Wasted: WLZ/WHZ below -2, 0-60 months	**0.091* [0.050] N=1,093**	0.083 [0.056] N=973	**0.129* [0.066] N=831**	-0.028 [0.063] N=674	-0.002 [0.072] N=512
Overweight: WLZ/WHZ above +2, 0-60 months	**0.102* [0.057] N=1,093**	**0.102* [0.060] N=973**	**0.124* [0.069] N=831**	**0.187** [0.087] N=674**	0.133 [0.102] N=512
Obese: WLZ/WHZ above +3, 0-60 months	**0.075* [0.041] N=1,093**	**0.076* [0.042] N=973**	0.069 [0.049] N=831	0.106 [0.069] N=674	0.003 [0.081] N=512
CIAF: any stunting, underweight, or wasting, 0-60 months	0.074 [0.089] N=1,058	0.058 [0.097] N=938	0.109 [0.112] N=802	0.029 [0.133] N=653	0.041 [0.148] N=496
BMI-for-age z score below -2	**0.103** [0.048] N=1,061**	**0.095* [0.057] N=946**	**0.130* [0.066] N=810**	0.018 [0.064] N=658	0.074 [0.078] N=504

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.048	-0.150 [0.314] N=1,189	-0.271 [0.325] N=1,072	-0.289 [0.340] N=942	-0.570 [0.368] N=795	-0.676 [0.410] N=642
WHO weight-for-age z score	0.243	0.154 [0.147] N=1,343	0.204 [0.154] N=1,220	0.061 [0.153] N=1,069	0.117 [0.162] N=896	0.071 [0.184] N=721
WHO weight-for-length/height z score, 0-60 months	0.384	0.257 [0.221] N=1,177	0.306 [0.236] N=1,064	0.251 [0.256] N=934	**0.468* [0.272] N=786**	0.317 [0.298] N=637
WHO BMI-for-age z score	0.315	0.184 [0.258] N=1,138	0.248 [0.277] N=1,030	0.137 [0.301] N=906	0.322 [0.326] N=763	0.190 [0.367] N=621
Stunted: HAZ below -2	0.200 P=238	0.004 [0.055] N=1,189	0.024 [0.056] N=1,072	0.015 [0.059] N=942	0.041 [0.064] N=795	0.041 [0.071] N=642
Underweight: WAZ below -2	0.051 P=72	0.020 [0.025] N=1,343	0.011 [0.026] N=1,220	0.015 [0.028] N=1,069	0.006 [0.030] N=896	-0.003 [0.035] N=721
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=132	0.022 [0.039] N=1,177	0.015 [0.040] N=1,064	0.035 [0.043] N=934	-0.033 [0.041] N=786	-0.016 [0.044] N=637
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=232	**0.115** [0.046] N=1,177**	**0.122** [0.049] N=1,064**	**0.123** [0.053] N=934**	**0.145** [0.061] N=786**	**0.130** [0.065] N=637**
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=118	0.047 [0.037] N=1,177	0.046 [0.038] N=1,064	0.036 [0.043] N=934	0.056 [0.050] N=786	0.026 [0.049] N=637
CIAF: any stunting, underweight, or wasting, 0-60 months	0.286 P=328	0.033 [0.058] N=1,142	0.033 [0.060] N=1,029	0.057 [0.063] N=905	0.014 [0.067] N=763	0.044 [0.070] N=618
BMI-for-age z score below -2	0.125 P=141	0.025 [0.044] N=1,138	0.014 [0.048] N=1,030	0.022 [0.052] N=906	-0.021 [0.053] N=763	0.008 [0.056] N=621

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.350 [0.429] N=1,028	0.231 [0.470] N=911	0.278 [0.533] N=781	-0.296 [0.651] N=634	-0.271 [0.776] N=481
WHO weight-for-age z score	0.246 [0.190] N=1,168	0.330 [0.200] N=1,045	0.071 [0.203] N=894	0.254 [0.212] N=721	0.287 [0.272] N=546
WHO weight-for-length/height z score, 0-60 months	0.046 [0.306] N=1,019	0.104 [0.335] N=906	-0.017 [0.385] N=776	0.491 [0.428] N=628	0.087 [0.502] N=479
WHO BMI-for-age z score	-0.160 [0.331] N=988	-0.096 [0.372] N=880	-0.355 [0.422] N=756	0.081 [0.490] N=613	-0.314 [0.582] N=471
Stunted: HAZ below -2	-0.015 [0.092] N=1,028	0.004 [0.099] N=911	-0.010 [0.110] N=781	0.045 [0.135] N=634	-0.003 [0.169] N=481
Underweight: WAZ below -2	0.019 [0.037] N=1,168	0.002 [0.039] N=1,045	0.010 [0.044] N=894	-0.022 [0.048] N=721	-0.060 [0.053] N=546
Wasted: WLZ/WHZ below -2, 0-60 months	0.083 [0.053] N=1,019	0.072 [0.059] N=906	**0.120* [0.068] N=776**	-0.027 [0.068] N=628	0.010 [0.078] N=479
Overweight: WLZ/WHZ above +2, 0-60 months	**0.128** [0.061] N=1,019**	**0.130* [0.066] N=906**	**0.142* [0.074] N=776**	**0.189** [0.094] N=628**	0.133 [0.106] N=479
Obese: WLZ/WHZ above +3, 0-60 months	**0.075* [0.044] N=1,019**	0.072 [0.045] N=906	0.064 [0.052] N=776	0.098 [0.073] N=628	-0.009 [0.084] N=479
CIAF: any stunting, underweight, or wasting, 0-60 months	0.063 [0.097] N=988	0.054 [0.105] N=875	0.114 [0.118] N=751	0.017 [0.142] N=609	0.031 [0.167] N=464
BMI-for-age z score below -2	**0.108** [0.051] N=988**	0.096 [0.061] N=880	**0.130* [0.068] N=756**	0.026 [0.069] N=613	0.109 [0.081] N=471

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.048	-0.176 [0.321] N=1,189	-0.296 [0.332] N=1,072	-0.309 [0.346] N=942	-0.612 [0.376] N=795	**-0.710* [0.420] N=642**
WHO weight-for-age z score	0.243	0.151 [0.148] N=1,343	0.195 [0.155] N=1,220	0.057 [0.154] N=1,069	0.109 [0.163] N=896	0.068 [0.186] N=721
WHO weight-for-length/height z score, 0-60 months	0.384	0.272 [0.223] N=1,177	0.312 [0.236] N=1,064	0.260 [0.256] N=934	**0.487* [0.270] N=786**	0.341 [0.301] N=637
WHO BMI-for-age z score	0.315	0.201 [0.262] N=1,138	0.255 [0.280] N=1,030	0.147 [0.305] N=906	0.344 [0.329] N=763	0.216 [0.376] N=621
Stunted: HAZ below -2	0.200 P=238	0.007 [0.054] N=1,189	0.024 [0.056] N=1,072	0.016 [0.058] N=942	0.044 [0.063] N=795	0.045 [0.071] N=642
Underweight: WAZ below -2	0.051 P=72	0.020 [0.025] N=1,343	0.011 [0.026] N=1,220	0.016 [0.028] N=1,069	0.006 [0.030] N=896	-0.002 [0.035] N=721
Wasted: WLZ/WHZ below -2, 0-60 months	0.111 P=132	0.020 [0.039] N=1,177	0.013 [0.041] N=1,064	0.032 [0.043] N=934	-0.038 [0.042] N=786	-0.020 [0.045] N=637
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=232	**0.117** [0.046] N=1,177**	**0.121** [0.048] N=1,064**	**0.123** [0.053] N=934**	**0.146** [0.060] N=786**	**0.134** [0.065] N=637**
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=118	0.047 [0.037] N=1,177	0.045 [0.038] N=1,064	0.036 [0.042] N=934	0.056 [0.050] N=786	0.027 [0.049] N=637
CIAF: any stunting, underweight, or wasting, 0-60 months	0.286 P=328	0.032 [0.058] N=1,142	0.031 [0.059] N=1,029	0.054 [0.062] N=905	0.011 [0.066] N=763	0.044 [0.069] N=618
BMI-for-age z score below -2	0.125 P=141	0.021 [0.046] N=1,138	0.012 [0.050] N=1,030	0.019 [0.054] N=906	-0.026 [0.055] N=763	0.004 [0.058] N=621

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.308 [0.439] N=1,028	0.201 [0.477] N=911	0.244 [0.542] N=781	-0.408 [0.669] N=634	-0.454 [0.820] N=481
WHO weight-for-age z score	0.231 [0.195] N=1,168	0.306 [0.208] N=1,045	0.054 [0.208] N=894	0.226 [0.223] N=721	0.248 [0.290] N=546
WHO weight-for-length/height z score, 0-60 months	0.040 [0.304] N=1,019	0.078 [0.331] N=906	-0.032 [0.382] N=776	0.524 [0.425] N=628	0.112 [0.497] N=479
WHO BMI-for-age z score	-0.159 [0.330] N=988	-0.121 [0.366] N=880	-0.368 [0.419] N=756	0.114 [0.485] N=613	-0.259 [0.583] N=471
Stunted: HAZ below -2	-0.011 [0.092] N=1,028	0.004 [0.099] N=911	-0.010 [0.110] N=781	0.056 [0.136] N=634	0.017 [0.170] N=481
Underweight: WAZ below -2	0.020 [0.038] N=1,168	0.004 [0.040] N=1,045	0.012 [0.044] N=894	-0.017 [0.048] N=721	-0.053 [0.054] N=546
Wasted: WLZ/WHZ below -2, 0-60 months	0.084 [0.053] N=1,019	0.077 [0.059] N=906	**0.122* [0.068] N=776**	-0.033 [0.067] N=628	0.003 [0.077] N=479
Overweight: WLZ/WHZ above +2, 0-60 months	**0.128** [0.060] N=1,019**	**0.128* [0.065] N=906**	**0.140* [0.073] N=776**	**0.193** [0.093] N=628**	0.134 [0.105] N=479
Obese: WLZ/WHZ above +3, 0-60 months	**0.074* [0.043] N=1,019**	0.070 [0.043] N=906	0.062 [0.051] N=776	0.100 [0.072] N=628	-0.005 [0.083] N=479
CIAF: any stunting, underweight, or wasting, 0-60 months	0.065 [0.097] N=988	0.057 [0.105] N=875	0.116 [0.118] N=751	0.019 [0.142] N=609	0.036 [0.165] N=464
BMI-for-age z score below -2	**0.109** [0.051] N=988**	**0.102* [0.060] N=880**	**0.134** [0.067] N=756**	0.022 [0.067] N=613	0.103 [0.079] N=471

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.