Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Youngest Child

Age reference: 12 November 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,520	-0.018 [0.021] N=1,618	-0.004 [0.021] N=1,494	-0.010 [0.023] N=1,344	-0.012 [0.025] N=1,146	-0.018 [0.030] N=939
Vegetables consumed at least weekly	0.902 P=1,458	0.012 [0.028] N=1,618	0.010 [0.028] N=1,494	-0.004 [0.029] N=1,344	-0.006 [0.030] N=1,146	-0.024 [0.033] N=939
Meat fish or poultry consumed at least weekly	0.796 P=1,293	**0.073* [0.042] N=1,618**	**0.084* [0.043] N=1,494**	0.077 [0.047] N=1,344	0.052 [0.052] N=1,146	0.037 [0.058] N=939
Legumes consumed at least weekly	0.795 P=1,290	-0.061 [0.041] N=1,618	**-0.078* [0.042] N=1,494**	-0.073 [0.044] N=1,344	-0.063 [0.050] N=1,146	-0.090 [0.057] N=939
Animal products consumed at least weekly	0.916 P=1,483	0.012 [0.031] N=1,618	0.012 [0.032] N=1,494	0.000 [0.036] N=1,344	0.003 [0.039] N=1,146	-0.002 [0.044] N=939
Sweets consumed at least weekly	0.689 P=1,109	-0.031 [0.039] N=1,618	-0.015 [0.041] N=1,494	0.003 [0.043] N=1,344	-0.010 [0.045] N=1,146	0.018 [0.051] N=939
Snack foods consumed at least weekly	0.562 P=901	-0.030 [0.043] N=1,618	-0.024 [0.046] N=1,494	-0.011 [0.050] N=1,344	-0.030 [0.053] N=1,146	-0.035 [0.056] N=939
Packaged soft drinks consumed at least weekly	0.339 P=538	0.026 [0.051] N=1,618	0.033 [0.055] N=1,494	0.031 [0.057] N=1,344	0.046 [0.063] N=1,146	0.037 [0.074] N=939
Grains consumed at least weekly	0.917 P=1,482	0.002 [0.028] N=1,618	0.008 [0.030] N=1,494	-0.008 [0.030] N=1,344	-0.008 [0.033] N=1,146	-0.016 [0.040] N=939
Child was ever breastfed	0.964 P=1,557	0.001 [0.016] N=1,620	0.005 [0.017] N=1,496	0.008 [0.018] N=1,345	0.011 [0.020] N=1,147	0.011 [0.020] N=940

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.004 [0.032] N=1,378	0.025 [0.031] N=1,254	0.018 [0.034] N=1,104	0.019 [0.040] N=906	0.028 [0.056] N=699
Vegetables consumed at least weekly	0.049 [0.042] N=1,378	0.056 [0.045] N=1,254	0.033 [0.050] N=1,104	0.035 [0.058] N=906	0.003 [0.075] N=699
Meat fish or poultry consumed at least weekly	0.072 [0.054] N=1,378	**0.092* [0.055] N=1,254**	0.081 [0.064] N=1,104	0.014 [0.072] N=906	-0.035 [0.083] N=699
Legumes consumed at least weekly	-0.013 [0.061] N=1,378	-0.038 [0.066] N=1,254	-0.009 [0.071] N=1,104	0.027 [0.088] N=906	-0.002 [0.119] N=699
Animal products consumed at least weekly	0.008 [0.039] N=1,378	0.008 [0.042] N=1,254	-0.015 [0.050] N=1,104	-0.009 [0.061] N=906	-0.027 [0.078] N=699
Sweets consumed at least weekly	-0.042 [0.055] N=1,378	-0.017 [0.060] N=1,254	0.015 [0.066] N=1,104	-0.014 [0.073] N=906	0.085 [0.097] N=699
Snack foods consumed at least weekly	-0.030 [0.058] N=1,378	-0.024 [0.065] N=1,254	0.002 [0.078] N=1,104	-0.039 [0.091] N=906	-0.058 [0.107] N=699
Packaged soft drinks consumed at least weekly	-0.002 [0.059] N=1,378	0.002 [0.066] N=1,254	-0.005 [0.070] N=1,104	0.008 [0.080] N=906	-0.040 [0.102] N=699
Grains consumed at least weekly	0.044 [0.044] N=1,378	0.066 [0.050] N=1,254	0.040 [0.053] N=1,104	0.061 [0.064] N=906	0.091 [0.097] N=699
Child was ever breastfed	0.009 [0.026] N=1,380	0.019 [0.030] N=1,256	0.024 [0.036] N=1,105	0.030 [0.044] N=907	0.059 [0.055] N=700

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,520	-0.018 [0.021] N=1,618	-0.004 [0.021] N=1,494	-0.009 [0.023] N=1,344	-0.012 [0.025] N=1,146	-0.017 [0.030] N=939
Vegetables consumed at least weekly	0.902 P=1,458	0.012 [0.028] N=1,618	0.010 [0.028] N=1,494	-0.003 [0.029] N=1,344	-0.005 [0.029] N=1,146	-0.024 [0.032] N=939
Meat fish or poultry consumed at least weekly	0.796 P=1,293	**0.073* [0.042] N=1,618**	**0.084* [0.043] N=1,494**	0.077 [0.047] N=1,344	0.053 [0.053] N=1,146	0.037 [0.059] N=939
Legumes consumed at least weekly	0.795 P=1,290	-0.061 [0.041] N=1,618	**-0.078* [0.042] N=1,494**	-0.073 [0.044] N=1,344	-0.063 [0.049] N=1,146	-0.090 [0.057] N=939
Animal products consumed at least weekly	0.916 P=1,483	0.012 [0.031] N=1,618	0.012 [0.032] N=1,494	0.002 [0.035] N=1,344	0.004 [0.039] N=1,146	-0.002 [0.044] N=939
Sweets consumed at least weekly	0.689 P=1,109	-0.031 [0.039] N=1,618	-0.015 [0.041] N=1,494	0.003 [0.043] N=1,344	-0.009 [0.045] N=1,146	0.018 [0.051] N=939
Snack foods consumed at least weekly	0.562 P=901	-0.030 [0.044] N=1,618	-0.023 [0.047] N=1,494	-0.009 [0.050] N=1,344	-0.030 [0.054] N=1,146	-0.035 [0.057] N=939
Packaged soft drinks consumed at least weekly	0.339 P=538	0.026 [0.051] N=1,618	0.033 [0.055] N=1,494	0.032 [0.057] N=1,344	0.047 [0.063] N=1,146	0.037 [0.074] N=939
Grains consumed at least weekly	0.917 P=1,482	0.002 [0.028] N=1,618	0.008 [0.030] N=1,494	-0.008 [0.030] N=1,344	-0.008 [0.032] N=1,146	-0.016 [0.040] N=939
Child was ever breastfed	0.964 P=1,557	0.001 [0.016] N=1,620	0.005 [0.017] N=1,496	0.008 [0.019] N=1,345	0.011 [0.020] N=1,147	0.011 [0.020] N=940

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.003 [0.032] N=1,378	0.027 [0.031] N=1,254	0.020 [0.034] N=1,104	0.020 [0.040] N=906	0.029 [0.057] N=699
Vegetables consumed at least weekly	0.049 [0.042] N=1,378	0.056 [0.045] N=1,254	0.034 [0.050] N=1,104	0.036 [0.058] N=906	0.004 [0.075] N=699
Meat fish or poultry consumed at least weekly	0.071 [0.054] N=1,378	**0.092* [0.055] N=1,254**	0.083 [0.064] N=1,104	0.016 [0.072] N=906	-0.031 [0.083] N=699
Legumes consumed at least weekly	-0.013 [0.061] N=1,378	-0.038 [0.066] N=1,254	-0.007 [0.071] N=1,104	0.030 [0.088] N=906	-0.000 [0.119] N=699
Animal products consumed at least weekly	0.010 [0.039] N=1,378	0.011 [0.042] N=1,254	-0.010 [0.049] N=1,104	-0.006 [0.060] N=906	-0.024 [0.077] N=699
Sweets consumed at least weekly	-0.043 [0.055] N=1,378	-0.016 [0.060] N=1,254	0.015 [0.066] N=1,104	-0.013 [0.073] N=906	0.085 [0.097] N=699
Snack foods consumed at least weekly	-0.028 [0.058] N=1,378	-0.020 [0.065] N=1,254	0.009 [0.078] N=1,104	-0.034 [0.092] N=906	-0.052 [0.107] N=699
Packaged soft drinks consumed at least weekly	-0.002 [0.059] N=1,378	0.003 [0.066] N=1,254	-0.005 [0.069] N=1,104	0.008 [0.079] N=906	-0.042 [0.102] N=699
Grains consumed at least weekly	0.044 [0.044] N=1,378	0.066 [0.050] N=1,254	0.040 [0.053] N=1,104	0.060 [0.063] N=906	0.089 [0.096] N=699
Child was ever breastfed	0.009 [0.027] N=1,380	0.019 [0.031] N=1,256	0.024 [0.036] N=1,105	0.029 [0.045] N=907	0.058 [0.055] N=700

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,520	-0.027 [0.021] N=1,618	-0.014 [0.021] N=1,494	-0.019 [0.023] N=1,344	-0.020 [0.026] N=1,146	-0.025 [0.031] N=939
Vegetables consumed at least weekly	0.902 P=1,458	0.006 [0.028] N=1,618	0.003 [0.028] N=1,494	-0.010 [0.029] N=1,344	-0.011 [0.029] N=1,146	-0.031 [0.032] N=939
Meat fish or poultry consumed at least weekly	0.796 P=1,293	0.049 [0.040] N=1,618	0.055 [0.041] N=1,494	0.053 [0.045] N=1,344	0.030 [0.050] N=1,146	0.012 [0.056] N=939
Legumes consumed at least weekly	0.795 P=1,290	-0.068 [0.041] N=1,618	**-0.087** [0.043] N=1,494**	**-0.080* [0.045] N=1,344**	-0.068 [0.050] N=1,146	-0.093 [0.057] N=939
Animal products consumed at least weekly	0.916 P=1,483	0.007 [0.031] N=1,618	0.006 [0.032] N=1,494	-0.004 [0.036] N=1,344	-0.001 [0.040] N=1,146	-0.010 [0.045] N=939
Sweets consumed at least weekly	0.689 P=1,109	-0.036 [0.039] N=1,618	-0.022 [0.042] N=1,494	-0.004 [0.044] N=1,344	-0.015 [0.045] N=1,146	0.011 [0.051] N=939
Snack foods consumed at least weekly	0.562 P=901	-0.028 [0.043] N=1,618	-0.021 [0.046] N=1,494	-0.009 [0.049] N=1,344	-0.027 [0.052] N=1,146	-0.031 [0.055] N=939
Packaged soft drinks consumed at least weekly	0.339 P=538	0.038 [0.049] N=1,618	0.049 [0.053] N=1,494	0.046 [0.055] N=1,344	0.060 [0.060] N=1,146	0.052 [0.071] N=939
Grains consumed at least weekly	0.917 P=1,482	-0.005 [0.028] N=1,618	0.001 [0.030] N=1,494	-0.014 [0.030] N=1,344	-0.016 [0.032] N=1,146	-0.023 [0.040] N=939
Child was ever breastfed	0.964 P=1,557	-0.004 [0.016] N=1,620	-0.000 [0.017] N=1,496	0.002 [0.019] N=1,345	0.006 [0.020] N=1,147	0.007 [0.020] N=940

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.011 [0.031] N=1,378	0.016 [0.029] N=1,254	0.014 [0.033] N=1,104	0.025 [0.038] N=906	0.039 [0.056] N=699
Vegetables consumed at least weekly	0.045 [0.042] N=1,378	0.050 [0.045] N=1,254	0.031 [0.050] N=1,104	0.042 [0.058] N=906	0.012 [0.075] N=699
Meat fish or poultry consumed at least weekly	0.051 [0.052] N=1,378	0.061 [0.053] N=1,254	0.061 [0.063] N=1,104	0.002 [0.070] N=906	-0.043 [0.084] N=699
Legumes consumed at least weekly	-0.021 [0.061] N=1,378	-0.050 [0.066] N=1,254	-0.015 [0.072] N=1,104	0.031 [0.089] N=906	0.004 [0.121] N=699
Animal products consumed at least weekly	0.006 [0.039] N=1,378	0.005 [0.042] N=1,254	-0.014 [0.050] N=1,104	-0.003 [0.061] N=906	-0.017 [0.077] N=699
Sweets consumed at least weekly	-0.046 [0.055] N=1,378	-0.022 [0.060] N=1,254	0.012 [0.067] N=1,104	-0.012 [0.072] N=906	0.090 [0.097] N=699
Snack foods consumed at least weekly	-0.027 [0.057] N=1,378	-0.017 [0.064] N=1,254	0.012 [0.076] N=1,104	-0.029 [0.088] N=906	-0.043 [0.104] N=699
Packaged soft drinks consumed at least weekly	0.010 [0.058] N=1,378	0.025 [0.064] N=1,254	0.014 [0.068] N=1,104	0.026 [0.079] N=906	-0.022 [0.099] N=699
Grains consumed at least weekly	0.037 [0.044] N=1,378	0.059 [0.049] N=1,254	0.033 [0.053] N=1,104	0.053 [0.063] N=906	0.079 [0.094] N=699
Child was ever breastfed	0.006 [0.027] N=1,380	0.014 [0.031] N=1,256	0.020 [0.036] N=1,105	0.029 [0.045] N=907	0.053 [0.056] N=700

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,520	-0.029 [0.022] N=1,618	-0.016 [0.022] N=1,494	-0.022 [0.023] N=1,344	-0.023 [0.026] N=1,146	-0.027 [0.031] N=939
Vegetables consumed at least weekly	0.902 P=1,458	0.005 [0.028] N=1,618	0.003 [0.028] N=1,494	-0.010 [0.030] N=1,344	-0.013 [0.030] N=1,146	-0.030 [0.033] N=939
Meat fish or poultry consumed at least weekly	0.796 P=1,293	0.045 [0.040] N=1,618	0.051 [0.041] N=1,494	0.045 [0.045] N=1,344	0.023 [0.050] N=1,146	0.003 [0.055] N=939
Legumes consumed at least weekly	0.795 P=1,290	-0.070 [0.042] N=1,618	**-0.089** [0.044] N=1,494**	**-0.087* [0.046] N=1,344**	-0.072 [0.052] N=1,146	-0.097 [0.058] N=939
Animal products consumed at least weekly	0.916 P=1,483	0.006 [0.031] N=1,618	0.007 [0.032] N=1,494	-0.003 [0.036] N=1,344	-0.002 [0.040] N=1,146	-0.010 [0.044] N=939
Sweets consumed at least weekly	0.689 P=1,109	-0.034 [0.039] N=1,618	-0.020 [0.041] N=1,494	-0.005 [0.043] N=1,344	-0.013 [0.045] N=1,146	0.012 [0.051] N=939
Snack foods consumed at least weekly	0.562 P=901	-0.028 [0.044] N=1,618	-0.021 [0.046] N=1,494	-0.011 [0.050] N=1,344	-0.027 [0.053] N=1,146	-0.032 [0.056] N=939
Packaged soft drinks consumed at least weekly	0.339 P=538	0.042 [0.049] N=1,618	0.051 [0.053] N=1,494	0.047 [0.055] N=1,344	0.065 [0.060] N=1,146	0.060 [0.071] N=939
Grains consumed at least weekly	0.917 P=1,482	-0.003 [0.028] N=1,618	0.003 [0.030] N=1,494	-0.013 [0.030] N=1,344	-0.014 [0.032] N=1,146	-0.020 [0.042] N=939
Child was ever breastfed	0.964 P=1,557	-0.004 [0.017] N=1,620	-0.000 [0.018] N=1,496	0.002 [0.019] N=1,345	0.005 [0.021] N=1,147	0.005 [0.021] N=940

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.012 [0.031] N=1,378	0.016 [0.030] N=1,254	0.012 [0.034] N=1,104	0.020 [0.039] N=906	0.027 [0.058] N=699
Vegetables consumed at least weekly	0.045 [0.042] N=1,378	0.050 [0.045] N=1,254	0.029 [0.050] N=1,104	0.037 [0.059] N=906	-0.003 [0.076] N=699
Meat fish or poultry consumed at least weekly	0.048 [0.053] N=1,378	0.058 [0.054] N=1,254	0.057 [0.063] N=1,104	0.003 [0.072] N=906	-0.054 [0.089] N=699
Legumes consumed at least weekly	-0.017 [0.061] N=1,378	-0.046 [0.066] N=1,254	-0.013 [0.071] N=1,104	0.042 [0.087] N=906	0.009 [0.123] N=699
Animal products consumed at least weekly	0.005 [0.039] N=1,378	0.007 [0.042] N=1,254	-0.014 [0.050] N=1,104	-0.011 [0.061] N=906	-0.033 [0.081] N=699
Sweets consumed at least weekly	-0.041 [0.054] N=1,378	-0.018 [0.059] N=1,254	0.014 [0.064] N=1,104	-0.001 [0.072] N=906	0.096 [0.096] N=699
Snack foods consumed at least weekly	-0.021 [0.056] N=1,378	-0.015 [0.063] N=1,254	0.012 [0.074] N=1,104	-0.018 [0.087] N=906	-0.031 [0.101] N=699
Packaged soft drinks consumed at least weekly	0.016 [0.057] N=1,378	0.024 [0.064] N=1,254	0.014 [0.068] N=1,104	0.033 [0.079] N=906	-0.003 [0.098] N=699
Grains consumed at least weekly	0.037 [0.043] N=1,378	0.058 [0.048] N=1,254	0.033 [0.051] N=1,104	0.052 [0.059] N=906	0.071 [0.089] N=699
Child was ever breastfed	0.007 [0.027] N=1,380	0.015 [0.031] N=1,256	0.020 [0.037] N=1,105	0.030 [0.046] N=907	0.059 [0.057] N=700

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,415	-0.026 [0.022] N=1,507	-0.016 [0.022] N=1,398	-0.021 [0.024] N=1,256	-0.023 [0.026] N=1,076	-0.032 [0.031] N=882
Vegetables consumed at least weekly	0.902 P=1,358	-0.002 [0.029] N=1,507	-0.005 [0.029] N=1,398	-0.022 [0.030] N=1,256	-0.018 [0.031] N=1,076	-0.034 [0.032] N=882
Meat fish or poultry consumed at least weekly	0.794 P=1,201	0.062 [0.040] N=1,507	0.064 [0.041] N=1,398	0.051 [0.044] N=1,256	0.022 [0.049] N=1,076	-0.010 [0.053] N=882
Legumes consumed at least weekly	0.791 P=1,195	**-0.074* [0.041] N=1,507**	**-0.096** [0.042] N=1,398**	**-0.094** [0.043] N=1,256**	-0.078 [0.050] N=1,076	**-0.107* [0.056] N=882**
Animal products consumed at least weekly	0.917 P=1,383	0.011 [0.031] N=1,507	0.010 [0.032] N=1,398	-0.001 [0.036] N=1,256	-0.004 [0.039] N=1,076	-0.011 [0.044] N=882
Sweets consumed at least weekly	0.692 P=1,035	-0.013 [0.040] N=1,507	-0.003 [0.043] N=1,398	0.011 [0.044] N=1,256	-0.000 [0.046] N=1,076	0.027 [0.053] N=882
Snack foods consumed at least weekly	0.563 P=840	-0.028 [0.047] N=1,507	-0.021 [0.050] N=1,398	-0.010 [0.054] N=1,256	-0.030 [0.058] N=1,076	-0.019 [0.064] N=882
Packaged soft drinks consumed at least weekly	0.344 P=510	0.060 [0.051] N=1,507	0.069 [0.054] N=1,398	0.058 [0.057] N=1,256	0.067 [0.061] N=1,076	0.073 [0.072] N=882
Grains consumed at least weekly	0.917 P=1,379	-0.003 [0.029] N=1,507	0.001 [0.031] N=1,398	-0.020 [0.031] N=1,256	-0.017 [0.033] N=1,076	-0.025 [0.041] N=882
Child was ever breastfed	0.965 P=1,451	-0.003 [0.017] N=1,509	0.004 [0.019] N=1,400	0.006 [0.020] N=1,257	0.013 [0.021] N=1,077	0.011 [0.022] N=883

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.006 [0.034] N=1,282	0.015 [0.033] N=1,173	0.007 [0.037] N=1,031	0.009 [0.043] N=851	0.012 [0.059] N=657
Vegetables consumed at least weekly	0.038 [0.044] N=1,282	0.038 [0.048] N=1,173	0.006 [0.053] N=1,031	0.018 [0.061] N=851	-0.007 [0.074] N=657
Meat fish or poultry consumed at least weekly	0.090 [0.055] N=1,282	0.092 [0.056] N=1,173	0.079 [0.064] N=1,031	0.010 [0.073] N=851	-0.051 [0.081] N=657
Legumes consumed at least weekly	-0.017 [0.062] N=1,282	-0.054 [0.066] N=1,173	-0.036 [0.070] N=1,031	0.005 [0.088] N=851	-0.033 [0.115] N=657
Animal products consumed at least weekly	0.013 [0.041] N=1,282	0.010 [0.044] N=1,173	-0.012 [0.052] N=1,031	-0.014 [0.063] N=851	-0.028 [0.080] N=657
Sweets consumed at least weekly	-0.019 [0.056] N=1,282	-0.004 [0.062] N=1,173	0.014 [0.067] N=1,031	-0.014 [0.075] N=851	0.058 [0.103] N=657
Snack foods consumed at least weekly	-0.030 [0.060] N=1,282	-0.023 [0.067] N=1,173	-0.003 [0.079] N=1,031	-0.059 [0.092] N=851	-0.081 [0.109] N=657
Packaged soft drinks consumed at least weekly	0.031 [0.062] N=1,282	0.037 [0.067] N=1,173	0.013 [0.071] N=1,031	0.010 [0.081] N=851	-0.015 [0.107] N=657
Grains consumed at least weekly	0.045 [0.046] N=1,282	0.061 [0.051] N=1,173	0.020 [0.053] N=1,031	0.049 [0.063] N=851	0.080 [0.095] N=657
Child was ever breastfed	0.002 [0.028] N=1,284	0.015 [0.031] N=1,175	0.019 [0.037] N=1,032	0.029 [0.044] N=852	0.048 [0.055] N=658

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.941 P=1,415	-0.027 [0.022] N=1,507	-0.017 [0.022] N=1,398	-0.021 [0.023] N=1,256	-0.023 [0.026] N=1,076	-0.033 [0.030] N=882
Vegetables consumed at least weekly	0.902 P=1,358	-0.003 [0.029] N=1,507	-0.006 [0.029] N=1,398	-0.023 [0.030] N=1,256	-0.019 [0.030] N=1,076	-0.036 [0.032] N=882
Meat fish or poultry consumed at least weekly	0.794 P=1,201	0.064 [0.041] N=1,507	0.066 [0.042] N=1,398	0.057 [0.045] N=1,256	0.030 [0.050] N=1,076	0.007 [0.056] N=882
Legumes consumed at least weekly	0.791 P=1,195	**-0.076* [0.041] N=1,507**	**-0.098** [0.042] N=1,398**	**-0.097** [0.044] N=1,256**	**-0.085* [0.050] N=1,076**	**-0.121** [0.056] N=882**
Animal products consumed at least weekly	0.917 P=1,383	0.010 [0.031] N=1,507	0.010 [0.032] N=1,398	-0.002 [0.036] N=1,256	-0.004 [0.039] N=1,076	-0.010 [0.044] N=882
Sweets consumed at least weekly	0.692 P=1,035	-0.013 [0.040] N=1,507	-0.003 [0.042] N=1,398	0.013 [0.043] N=1,256	0.003 [0.045] N=1,076	0.031 [0.051] N=882
Snack foods consumed at least weekly	0.563 P=840	-0.030 [0.046] N=1,507	-0.023 [0.050] N=1,398	-0.010 [0.053] N=1,256	-0.032 [0.057] N=1,076	-0.027 [0.062] N=882
Packaged soft drinks consumed at least weekly	0.344 P=510	0.058 [0.051] N=1,507	0.067 [0.054] N=1,398	0.057 [0.057] N=1,256	0.065 [0.061] N=1,076	0.064 [0.072] N=882
Grains consumed at least weekly	0.917 P=1,379	-0.004 [0.029] N=1,507	0.000 [0.031] N=1,398	-0.022 [0.031] N=1,256	-0.019 [0.034] N=1,076	-0.030 [0.040] N=882
Child was ever breastfed	0.965 P=1,451	-0.004 [0.017] N=1,509	0.003 [0.019] N=1,400	0.005 [0.020] N=1,257	0.011 [0.021] N=1,077	0.010 [0.021] N=883

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.008 [0.033] N=1,282	0.014 [0.032] N=1,173	0.009 [0.036] N=1,031	0.009 [0.042] N=851	0.011 [0.056] N=657
Vegetables consumed at least weekly	0.035 [0.044] N=1,282	0.036 [0.047] N=1,173	0.007 [0.053] N=1,031	0.017 [0.061] N=851	-0.009 [0.074] N=657
Meat fish or poultry consumed at least weekly	0.083 [0.054] N=1,282	0.087 [0.055] N=1,173	0.076 [0.063] N=1,031	0.012 [0.074] N=851	-0.038 [0.081] N=657
Legumes consumed at least weekly	-0.021 [0.062] N=1,282	-0.057 [0.067] N=1,173	-0.032 [0.072] N=1,031	0.008 [0.090] N=851	-0.040 [0.118] N=657
Animal products consumed at least weekly	0.010 [0.041] N=1,282	0.009 [0.044] N=1,173	-0.012 [0.051] N=1,031	-0.016 [0.062] N=851	-0.032 [0.079] N=657
Sweets consumed at least weekly	-0.031 [0.056] N=1,282	-0.019 [0.062] N=1,173	0.004 [0.068] N=1,031	-0.017 [0.075] N=851	0.066 [0.101] N=657
Snack foods consumed at least weekly	-0.040 [0.059] N=1,282	-0.035 [0.066] N=1,173	-0.010 [0.079] N=1,031	-0.060 [0.092] N=851	-0.077 [0.109] N=657
Packaged soft drinks consumed at least weekly	0.025 [0.061] N=1,282	0.031 [0.068] N=1,173	0.009 [0.072] N=1,031	0.008 [0.082] N=851	-0.025 [0.106] N=657
Grains consumed at least weekly	0.044 [0.046] N=1,282	0.062 [0.051] N=1,173	0.026 [0.055] N=1,031	0.052 [0.066] N=851	0.077 [0.096] N=657
Child was ever breastfed	0.003 [0.028] N=1,284	0.015 [0.032] N=1,175	0.018 [0.038] N=1,032	0.031 [0.047] N=852	0.053 [0.059] N=658

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,358	-0.015 [0.027] N=1,441	-0.016 [0.029] N=1,312	-0.021 [0.032] N=1,151	-0.020 [0.032] N=967	-0.019 [0.039] N=782
Vegetables consumed at least weekly	0.904 P=1,300	0.010 [0.032] N=1,441	0.021 [0.034] N=1,312	0.021 [0.037] N=1,151	0.014 [0.041] N=967	0.014 [0.046] N=782
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.075* [0.041] N=1,441**	**0.076* [0.043] N=1,312**	0.073 [0.046] N=1,151	**0.107** [0.047] N=967**	**0.096* [0.055] N=782**
Legumes consumed at least weekly	0.796 P=1,153	-0.031 [0.046] N=1,441	-0.014 [0.049] N=1,312	-0.023 [0.053] N=1,151	-0.045 [0.057] N=967	-0.026 [0.068] N=782
Animal products consumed at least weekly	0.913 P=1,317	-0.001 [0.034] N=1,441	0.003 [0.036] N=1,312	0.011 [0.038] N=1,151	-0.006 [0.041] N=967	0.018 [0.048] N=782
Sweets consumed at least weekly	0.694 P=1,001	0.007 [0.049] N=1,441	0.004 [0.052] N=1,312	-0.001 [0.056] N=1,151	0.003 [0.058] N=967	-0.044 [0.067] N=782
Snack foods consumed at least weekly	0.562 P=806	-0.005 [0.056] N=1,441	-0.001 [0.059] N=1,312	0.010 [0.062] N=1,151	0.027 [0.065] N=967	-0.021 [0.075] N=782
Packaged soft drinks consumed at least weekly	0.327 P=466	0.008 [0.046] N=1,441	0.003 [0.048] N=1,312	0.042 [0.051] N=1,151	0.050 [0.056] N=967	0.035 [0.063] N=782
Grains consumed at least weekly	0.917 P=1,320	0.032 [0.038] N=1,441	0.038 [0.040] N=1,312	0.036 [0.043] N=1,151	0.040 [0.048] N=967	0.037 [0.055] N=782
Child was ever breastfed	0.962 P=1,386	-0.010 [0.021] N=1,442	-0.008 [0.021] N=1,313	-0.009 [0.021] N=1,152	-0.009 [0.022] N=968	-0.017 [0.027] N=783

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.047 [0.034] N=1,253	-0.056 [0.037] N=1,124	**-0.075* [0.043] N=963**	-0.067 [0.045] N=779	-0.081 [0.063] N=594
Vegetables consumed at least weekly	-0.041 [0.036] N=1,253	-0.031 [0.040] N=1,124	-0.045 [0.046] N=963	-0.057 [0.052] N=779	-0.072 [0.062] N=594
Meat fish or poultry consumed at least weekly	0.017 [0.058] N=1,253	0.009 [0.061] N=1,124	-0.020 [0.067] N=963	0.037 [0.072] N=779	-0.012 [0.093] N=594
Legumes consumed at least weekly	-0.047 [0.053] N=1,253	-0.024 [0.059] N=1,124	-0.038 [0.070] N=963	-0.062 [0.076] N=779	0.001 [0.090] N=594
Animal products consumed at least weekly	-0.057 [0.039] N=1,253	-0.050 [0.043] N=1,124	-0.046 [0.046] N=963	**-0.093* [0.047] N=779**	-0.074 [0.061] N=594
Sweets consumed at least weekly	0.002 [0.065] N=1,253	-0.000 [0.070] N=1,124	-0.005 [0.080] N=963	-0.002 [0.090] N=779	-0.128 [0.114] N=594
Snack foods consumed at least weekly	0.009 [0.083] N=1,253	0.016 [0.090] N=1,124	0.038 [0.101] N=963	0.092 [0.114] N=779	0.001 [0.151] N=594
Packaged soft drinks consumed at least weekly	-0.024 [0.074] N=1,253	-0.029 [0.080] N=1,124	0.035 [0.092] N=963	0.052 [0.107] N=779	0.016 [0.134] N=594
Grains consumed at least weekly	0.004 [0.040] N=1,253	0.005 [0.042] N=1,124	-0.009 [0.049] N=963	-0.026 [0.061] N=779	-0.084 [0.080] N=594
Child was ever breastfed	0.016 [0.039] N=1,254	0.024 [0.040] N=1,125	0.027 [0.043] N=964	0.035 [0.046] N=780	0.031 [0.063] N=595

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,358	-0.014 [0.027] N=1,441	-0.016 [0.029] N=1,312	-0.020 [0.032] N=1,151	-0.020 [0.032] N=967	-0.019 [0.039] N=782
Vegetables consumed at least weekly	0.904 P=1,300	0.011 [0.031] N=1,441	0.022 [0.033] N=1,312	0.022 [0.036] N=1,151	0.014 [0.040] N=967	0.014 [0.045] N=782
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.076* [0.041] N=1,441**	**0.078* [0.043] N=1,312**	0.075 [0.046] N=1,151	**0.107** [0.047] N=967**	**0.096* [0.056] N=782**
Legumes consumed at least weekly	0.796 P=1,153	-0.031 [0.045] N=1,441	-0.014 [0.049] N=1,312	-0.023 [0.053] N=1,151	-0.045 [0.057] N=967	-0.026 [0.067] N=782
Animal products consumed at least weekly	0.913 P=1,317	0.002 [0.034] N=1,441	0.006 [0.036] N=1,312	0.013 [0.038] N=1,151	-0.006 [0.041] N=967	0.017 [0.047] N=782
Sweets consumed at least weekly	0.694 P=1,001	0.009 [0.049] N=1,441	0.006 [0.052] N=1,312	0.001 [0.056] N=1,151	0.003 [0.058] N=967	-0.045 [0.067] N=782
Snack foods consumed at least weekly	0.562 P=806	-0.001 [0.056] N=1,441	0.001 [0.059] N=1,312	0.012 [0.062] N=1,151	0.027 [0.065] N=967	-0.022 [0.075] N=782
Packaged soft drinks consumed at least weekly	0.327 P=466	0.009 [0.046] N=1,441	0.004 [0.048] N=1,312	0.044 [0.051] N=1,151	0.050 [0.056] N=967	0.034 [0.063] N=782
Grains consumed at least weekly	0.917 P=1,320	0.032 [0.038] N=1,441	0.038 [0.040] N=1,312	0.036 [0.043] N=1,151	0.040 [0.049] N=967	0.038 [0.056] N=782
Child was ever breastfed	0.962 P=1,386	-0.009 [0.021] N=1,442	-0.007 [0.021] N=1,313	-0.009 [0.021] N=1,152	-0.009 [0.022] N=968	-0.017 [0.027] N=783

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.046 [0.034] N=1,253	-0.054 [0.037] N=1,124	**-0.075* [0.044] N=963**	-0.067 [0.045] N=779	-0.082 [0.064] N=594
Vegetables consumed at least weekly	-0.040 [0.036] N=1,253	-0.030 [0.040] N=1,124	-0.044 [0.046] N=963	-0.055 [0.052] N=779	-0.071 [0.062] N=594
Meat fish or poultry consumed at least weekly	0.021 [0.058] N=1,253	0.014 [0.061] N=1,124	-0.015 [0.067] N=963	0.042 [0.073] N=779	-0.004 [0.095] N=594
Legumes consumed at least weekly	-0.046 [0.053] N=1,253	-0.024 [0.058] N=1,124	-0.039 [0.069] N=963	-0.062 [0.076] N=779	-0.005 [0.089] N=594
Animal products consumed at least weekly	-0.052 [0.038] N=1,253	-0.044 [0.042] N=1,124	-0.041 [0.046] N=963	**-0.089* [0.047] N=779**	-0.066 [0.060] N=594
Sweets consumed at least weekly	0.005 [0.066] N=1,253	0.005 [0.071] N=1,124	-0.001 [0.080] N=963	-0.002 [0.089] N=779	-0.125 [0.114] N=594
Snack foods consumed at least weekly	0.017 [0.084] N=1,253	0.023 [0.090] N=1,124	0.044 [0.102] N=963	0.094 [0.115] N=779	0.007 [0.153] N=594
Packaged soft drinks consumed at least weekly	-0.021 [0.074] N=1,253	-0.027 [0.080] N=1,124	0.038 [0.092] N=963	0.054 [0.108] N=779	0.023 [0.135] N=594
Grains consumed at least weekly	0.004 [0.040] N=1,253	0.004 [0.042] N=1,124	-0.011 [0.049] N=963	-0.029 [0.060] N=779	-0.089 [0.080] N=594
Child was ever breastfed	0.017 [0.039] N=1,254	0.025 [0.040] N=1,125	0.027 [0.043] N=964	0.035 [0.046] N=780	0.032 [0.064] N=595

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,358	-0.016 [0.027] N=1,441	-0.017 [0.029] N=1,312	-0.020 [0.032] N=1,151	-0.021 [0.033] N=967	-0.020 [0.039] N=782
Vegetables consumed at least weekly	0.904 P=1,300	0.010 [0.031] N=1,441	0.021 [0.033] N=1,312	0.022 [0.036] N=1,151	0.013 [0.040] N=967	0.013 [0.046] N=782
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.070* [0.039] N=1,441**	**0.072* [0.041] N=1,312**	**0.076* [0.044] N=1,151**	**0.108** [0.044] N=967**	**0.098* [0.051] N=782**
Legumes consumed at least weekly	0.796 P=1,153	-0.033 [0.045] N=1,441	-0.016 [0.049] N=1,312	-0.024 [0.053] N=1,151	-0.045 [0.057] N=967	-0.025 [0.068] N=782
Animal products consumed at least weekly	0.913 P=1,317	0.000 [0.034] N=1,441	0.003 [0.036] N=1,312	0.013 [0.038] N=1,151	-0.008 [0.041] N=967	0.017 [0.047] N=782
Sweets consumed at least weekly	0.694 P=1,001	0.009 [0.049] N=1,441	0.005 [0.052] N=1,312	0.001 [0.055] N=1,151	0.002 [0.058] N=967	-0.047 [0.067] N=782
Snack foods consumed at least weekly	0.562 P=806	0.003 [0.056] N=1,441	0.003 [0.059] N=1,312	0.012 [0.062] N=1,151	0.025 [0.066] N=967	-0.025 [0.076] N=782
Packaged soft drinks consumed at least weekly	0.327 P=466	0.011 [0.047] N=1,441	0.004 [0.049] N=1,312	0.040 [0.052] N=1,151	0.047 [0.057] N=967	0.031 [0.065] N=782
Grains consumed at least weekly	0.917 P=1,320	0.031 [0.038] N=1,441	0.037 [0.040] N=1,312	0.037 [0.043] N=1,151	0.040 [0.049] N=967	0.038 [0.056] N=782
Child was ever breastfed	0.962 P=1,386	-0.010 [0.021] N=1,442	-0.008 [0.021] N=1,313	-0.008 [0.021] N=1,152	-0.008 [0.022] N=968	-0.016 [0.027] N=783

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.034] N=1,253	-0.051 [0.036] N=1,124	-0.069 [0.042] N=963	-0.064 [0.045] N=779	-0.075 [0.063] N=594
Vegetables consumed at least weekly	-0.039 [0.036] N=1,253	-0.028 [0.039] N=1,124	-0.040 [0.044] N=963	-0.055 [0.050] N=779	-0.067 [0.062] N=594
Meat fish or poultry consumed at least weekly	0.022 [0.054] N=1,253	0.018 [0.057] N=1,124	-0.001 [0.062] N=963	0.057 [0.069] N=779	0.015 [0.085] N=594
Legumes consumed at least weekly	-0.043 [0.052] N=1,253	-0.022 [0.058] N=1,124	-0.038 [0.070] N=963	-0.062 [0.075] N=779	-0.005 [0.089] N=594
Animal products consumed at least weekly	-0.053 [0.038] N=1,253	-0.046 [0.042] N=1,124	-0.041 [0.045] N=963	**-0.094** [0.047] N=779**	-0.068 [0.060] N=594
Sweets consumed at least weekly	0.004 [0.066] N=1,253	0.002 [0.070] N=1,124	-0.004 [0.079] N=963	-0.010 [0.090] N=779	-0.134 [0.113] N=594
Snack foods consumed at least weekly	0.022 [0.085] N=1,253	0.024 [0.091] N=1,124	0.038 [0.103] N=963	0.084 [0.117] N=779	-0.003 [0.157] N=594
Packaged soft drinks consumed at least weekly	-0.031 [0.075] N=1,253	-0.042 [0.081] N=1,124	0.014 [0.094] N=963	0.021 [0.110] N=779	-0.004 [0.138] N=594
Grains consumed at least weekly	0.002 [0.039] N=1,253	0.003 [0.041] N=1,124	-0.008 [0.047] N=963	-0.031 [0.059] N=779	-0.090 [0.079] N=594
Child was ever breastfed	0.016 [0.039] N=1,254	0.023 [0.040] N=1,125	0.029 [0.044] N=964	0.041 [0.047] N=780	0.036 [0.063] N=595

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,358	-0.015 [0.027] N=1,441	-0.016 [0.028] N=1,312	-0.019 [0.032] N=1,151	-0.024 [0.033] N=967	-0.024 [0.038] N=782
Vegetables consumed at least weekly	0.904 P=1,300	0.011 [0.032] N=1,441	0.024 [0.034] N=1,312	0.024 [0.037] N=1,151	0.013 [0.041] N=967	0.014 [0.046] N=782
Meat fish or poultry consumed at least weekly	0.784 P=1,140	**0.071* [0.041] N=1,441**	**0.075* [0.042] N=1,312**	0.075 [0.045] N=1,151	**0.100** [0.047] N=967**	0.084 [0.054] N=782
Legumes consumed at least weekly	0.796 P=1,153	-0.027 [0.045] N=1,441	-0.011 [0.049] N=1,312	-0.018 [0.053] N=1,151	-0.043 [0.056] N=967	-0.023 [0.067] N=782
Animal products consumed at least weekly	0.913 P=1,317	0.007 [0.034] N=1,441	0.012 [0.036] N=1,312	0.022 [0.038] N=1,151	-0.001 [0.041] N=967	0.021 [0.048] N=782
Sweets consumed at least weekly	0.694 P=1,001	0.016 [0.050] N=1,441	0.012 [0.053] N=1,312	0.004 [0.057] N=1,151	0.004 [0.060] N=967	-0.044 [0.070] N=782
Snack foods consumed at least weekly	0.562 P=806	0.009 [0.056] N=1,441	0.009 [0.059] N=1,312	0.017 [0.062] N=1,151	0.029 [0.065] N=967	-0.016 [0.075] N=782
Packaged soft drinks consumed at least weekly	0.327 P=466	0.023 [0.046] N=1,441	0.012 [0.048] N=1,312	0.048 [0.052] N=1,151	0.055 [0.056] N=967	0.042 [0.065] N=782
Grains consumed at least weekly	0.917 P=1,320	0.030 [0.038] N=1,441	0.035 [0.040] N=1,312	0.034 [0.043] N=1,151	0.037 [0.048] N=967	0.035 [0.055] N=782
Child was ever breastfed	0.962 P=1,386	-0.009 [0.020] N=1,442	-0.007 [0.020] N=1,313	-0.009 [0.021] N=1,152	-0.009 [0.022] N=968	-0.017 [0.028] N=783

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.046 [0.033] N=1,253	-0.053 [0.035] N=1,124	-0.069 [0.042] N=963	-0.070 [0.046] N=779	-0.092 [0.064] N=594
Vegetables consumed at least weekly	-0.038 [0.037] N=1,253	-0.025 [0.040] N=1,124	-0.037 [0.046] N=963	-0.052 [0.052] N=779	-0.067 [0.063] N=594
Meat fish or poultry consumed at least weekly	0.021 [0.055] N=1,253	0.019 [0.058] N=1,124	-0.005 [0.063] N=963	0.039 [0.071] N=779	-0.029 [0.088] N=594
Legumes consumed at least weekly	-0.039 [0.052] N=1,253	-0.018 [0.058] N=1,124	-0.026 [0.070] N=963	-0.052 [0.075] N=779	0.004 [0.087] N=594
Animal products consumed at least weekly	-0.046 [0.037] N=1,253	-0.037 [0.041] N=1,124	-0.031 [0.043] N=963	**-0.086* [0.046] N=779**	-0.066 [0.057] N=594
Sweets consumed at least weekly	0.011 [0.066] N=1,253	0.008 [0.071] N=1,124	-0.001 [0.079] N=963	-0.010 [0.090] N=779	-0.139 [0.111] N=594
Snack foods consumed at least weekly	0.025 [0.082] N=1,253	0.025 [0.090] N=1,124	0.041 [0.099] N=963	0.083 [0.113] N=779	-0.002 [0.149] N=594
Packaged soft drinks consumed at least weekly	-0.009 [0.073] N=1,253	-0.026 [0.079] N=1,124	0.033 [0.091] N=963	0.045 [0.107] N=779	0.029 [0.136] N=594
Grains consumed at least weekly	0.001 [0.039] N=1,253	-0.000 [0.041] N=1,124	-0.015 [0.048] N=963	-0.036 [0.060] N=779	-0.099 [0.079] N=594
Child was ever breastfed	0.017 [0.038] N=1,254	0.025 [0.039] N=1,125	0.028 [0.042] N=964	0.037 [0.046] N=780	0.032 [0.065] N=595

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,264	-0.010 [0.029] N=1,341	-0.010 [0.031] N=1,219	-0.008 [0.034] N=1,073	-0.017 [0.035] N=902	-0.018 [0.041] N=732
Vegetables consumed at least weekly	0.905 P=1,212	0.011 [0.032] N=1,341	0.022 [0.035] N=1,219	0.034 [0.038] N=1,073	0.015 [0.042] N=902	-0.002 [0.047] N=732
Meat fish or poultry consumed at least weekly	0.783 P=1,059	**0.080* [0.044] N=1,341**	**0.080* [0.046] N=1,219**	**0.085* [0.049] N=1,073**	**0.091* [0.052] N=902**	0.078 [0.059] N=732
Legumes consumed at least weekly	0.794 P=1,071	-0.023 [0.047] N=1,341	-0.004 [0.051] N=1,219	-0.009 [0.055] N=1,073	-0.043 [0.060] N=902	-0.030 [0.072] N=732
Animal products consumed at least weekly	0.915 P=1,228	0.007 [0.035] N=1,341	0.017 [0.038] N=1,219	0.028 [0.039] N=1,073	0.005 [0.042] N=902	0.027 [0.048] N=732
Sweets consumed at least weekly	0.704 P=943	0.008 [0.050] N=1,341	0.006 [0.052] N=1,219	-0.007 [0.057] N=1,073	-0.011 [0.061] N=902	-0.053 [0.073] N=732
Snack foods consumed at least weekly	0.568 P=758	0.007 [0.057] N=1,341	0.005 [0.061] N=1,219	0.008 [0.064] N=1,073	0.024 [0.066] N=902	-0.013 [0.078] N=732
Packaged soft drinks consumed at least weekly	0.336 P=448	-0.000 [0.047] N=1,341	-0.010 [0.049] N=1,219	0.028 [0.052] N=1,073	0.037 [0.058] N=902	0.020 [0.067] N=732
Grains consumed at least weekly	0.917 P=1,228	0.041 [0.039] N=1,341	0.047 [0.041] N=1,219	0.048 [0.044] N=1,073	0.049 [0.050] N=902	0.041 [0.057] N=732
Child was ever breastfed	0.964 P=1,294	-0.016 [0.019] N=1,342	-0.017 [0.019] N=1,220	-0.020 [0.020] N=1,074	-0.023 [0.020] N=903	-0.028 [0.027] N=733

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.035 [0.037] N=1,163	-0.041 [0.040] N=1,041	-0.043 [0.049] N=895	-0.051 [0.049] N=724	-0.073 [0.064] N=554
Vegetables consumed at least weekly	-0.037 [0.038] N=1,163	-0.027 [0.042] N=1,041	-0.015 [0.048] N=895	-0.038 [0.056] N=724	-0.093 [0.061] N=554
Meat fish or poultry consumed at least weekly	0.044 [0.066] N=1,163	0.041 [0.070] N=1,041	0.031 [0.078] N=895	0.045 [0.085] N=724	-0.008 [0.106] N=554
Legumes consumed at least weekly	-0.026 [0.054] N=1,163	0.002 [0.058] N=1,041	0.004 [0.071] N=895	-0.033 [0.080] N=724	0.016 [0.093] N=554
Animal products consumed at least weekly	-0.045 [0.042] N=1,163	-0.027 [0.046] N=1,041	-0.013 [0.048] N=895	-0.064 [0.050] N=724	-0.038 [0.061] N=554
Sweets consumed at least weekly	0.002 [0.068] N=1,163	-0.001 [0.073] N=1,041	-0.030 [0.082] N=895	-0.047 [0.090] N=724	-0.182 [0.119] N=554
Snack foods consumed at least weekly	0.029 [0.088] N=1,163	0.028 [0.096] N=1,041	0.028 [0.109] N=895	0.079 [0.120] N=724	0.024 [0.160] N=554
Packaged soft drinks consumed at least weekly	-0.041 [0.075] N=1,163	-0.055 [0.081] N=1,041	0.001 [0.093] N=895	0.011 [0.111] N=724	-0.045 [0.142] N=554
Grains consumed at least weekly	0.021 [0.042] N=1,163	0.024 [0.045] N=1,041	0.019 [0.052] N=895	-0.004 [0.064] N=724	-0.071 [0.083] N=554
Child was ever breastfed	0.009 [0.035] N=1,164	0.012 [0.037] N=1,042	0.010 [0.039] N=896	0.012 [0.039] N=725	0.007 [0.059] N=555

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.945 P=1,264	-0.010 [0.029] N=1,341	-0.010 [0.031] N=1,219	-0.008 [0.034] N=1,073	-0.018 [0.035] N=902	-0.020 [0.040] N=732
Vegetables consumed at least weekly	0.905 P=1,212	0.011 [0.032] N=1,341	0.021 [0.035] N=1,219	0.032 [0.038] N=1,073	0.013 [0.042] N=902	-0.004 [0.047] N=732
Meat fish or poultry consumed at least weekly	0.783 P=1,059	**0.081* [0.043] N=1,341**	**0.083* [0.045] N=1,219**	**0.086* [0.048] N=1,073**	**0.094* [0.050] N=902**	0.078 [0.058] N=732
Legumes consumed at least weekly	0.794 P=1,071	-0.025 [0.047] N=1,341	-0.005 [0.051] N=1,219	-0.011 [0.055] N=1,073	-0.046 [0.060] N=902	-0.034 [0.072] N=732
Animal products consumed at least weekly	0.915 P=1,228	0.006 [0.035] N=1,341	0.016 [0.038] N=1,219	0.027 [0.039] N=1,073	0.003 [0.042] N=902	0.025 [0.048] N=732
Sweets consumed at least weekly	0.704 P=943	0.008 [0.051] N=1,341	0.006 [0.053] N=1,219	-0.007 [0.058] N=1,073	-0.010 [0.062] N=902	-0.053 [0.074] N=732
Snack foods consumed at least weekly	0.568 P=758	0.008 [0.056] N=1,341	0.007 [0.060] N=1,219	0.010 [0.063] N=1,073	0.025 [0.066] N=902	-0.014 [0.078] N=732
Packaged soft drinks consumed at least weekly	0.336 P=448	0.000 [0.047] N=1,341	-0.009 [0.049] N=1,219	0.028 [0.052] N=1,073	0.036 [0.058] N=902	0.017 [0.068] N=732
Grains consumed at least weekly	0.917 P=1,228	0.040 [0.039] N=1,341	0.046 [0.041] N=1,219	0.046 [0.044] N=1,073	0.047 [0.050] N=902	0.040 [0.057] N=732
Child was ever breastfed	0.964 P=1,294	-0.014 [0.019] N=1,342	-0.015 [0.020] N=1,220	-0.015 [0.021] N=1,074	-0.018 [0.021] N=903	-0.023 [0.028] N=733

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.035 [0.038] N=1,163	-0.040 [0.041] N=1,041	-0.043 [0.049] N=895	-0.053 [0.051] N=724	-0.072 [0.068] N=554
Vegetables consumed at least weekly	-0.038 [0.038] N=1,163	-0.028 [0.042] N=1,041	-0.017 [0.048] N=895	-0.042 [0.056] N=724	-0.093 [0.063] N=554
Meat fish or poultry consumed at least weekly	0.045 [0.065] N=1,163	0.044 [0.069] N=1,041	0.031 [0.077] N=895	0.046 [0.085] N=724	-0.006 [0.106] N=554
Legumes consumed at least weekly	-0.027 [0.054] N=1,163	0.003 [0.059] N=1,041	0.003 [0.071] N=895	-0.037 [0.080] N=724	0.020 [0.092] N=554
Animal products consumed at least weekly	-0.045 [0.042] N=1,163	-0.028 [0.047] N=1,041	-0.015 [0.048] N=895	-0.068 [0.051] N=724	-0.044 [0.062] N=554
Sweets consumed at least weekly	0.004 [0.068] N=1,163	0.003 [0.074] N=1,041	-0.022 [0.083] N=895	-0.034 [0.092] N=724	-0.164 [0.120] N=554
Snack foods consumed at least weekly	0.030 [0.087] N=1,163	0.028 [0.095] N=1,041	0.033 [0.108] N=895	0.087 [0.119] N=724	0.025 [0.159] N=554
Packaged soft drinks consumed at least weekly	-0.040 [0.075] N=1,163	-0.055 [0.081] N=1,041	0.006 [0.092] N=895	0.020 [0.108] N=724	-0.033 [0.140] N=554
Grains consumed at least weekly	0.020 [0.042] N=1,163	0.023 [0.045] N=1,041	0.017 [0.052] N=895	-0.008 [0.064] N=724	-0.076 [0.082] N=554
Child was ever breastfed	0.012 [0.036] N=1,164	0.012 [0.037] N=1,042	0.017 [0.041] N=896	0.019 [0.042] N=725	0.012 [0.062] N=555

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.