Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Youngest Child

Age reference: 12 November 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.383	0.284 [0.251] N=1,407	0.308 [0.271] N=1,303	0.289 [0.295] N=1,173	0.408 [0.298] N=1,005	0.153 [0.324] N=829
WHO weight-for-age z score	0.092	-0.113 [0.136] N=1,609	-0.087 [0.147] N=1,486	-0.088 [0.153] N=1,336	-0.118 [0.171] N=1,140	**-0.315* [0.187] N=934**
WHO weight-for-length/height z score, 0-60 months	0.391	-0.015 [0.211] N=1,387	-0.066 [0.224] N=1,290	-0.155 [0.255] N=1,164	-0.251 [0.283] N=996	-0.195 [0.310] N=818
WHO BMI-for-age z score	0.372	0.089 [0.244] N=1,353	0.114 [0.265] N=1,258	0.013 [0.300] N=1,133	-0.205 [0.330] N=969	-0.122 [0.363] N=799
Stunted: HAZ below -2	0.245 P=329	-0.053 [0.046] N=1,407	-0.065 [0.050] N=1,303	-0.044 [0.053] N=1,173	-0.053 [0.057] N=1,005	-0.018 [0.064] N=829
Underweight: WAZ below -2	0.068 P=112	0.018 [0.022] N=1,609	0.012 [0.023] N=1,486	0.009 [0.024] N=1,336	0.020 [0.027] N=1,140	0.052 [0.033] N=934
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=153	-0.029 [0.036] N=1,387	-0.024 [0.035] N=1,290	-0.022 [0.037] N=1,164	-0.008 [0.041] N=996	-0.026 [0.043] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=271	0.001 [0.046] N=1,387	-0.005 [0.049] N=1,290	-0.013 [0.055] N=1,164	-0.013 [0.062] N=996	-0.014 [0.069] N=818
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.012 [0.030] N=1,387	-0.023 [0.032] N=1,290	-0.018 [0.034] N=1,164	-0.021 [0.037] N=996	-0.013 [0.040] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	0.326 P=427	-0.077 [0.048] N=1,345	**-0.085* [0.049] N=1,250**	-0.066 [0.052] N=1,127	-0.061 [0.055] N=965	-0.034 [0.063] N=795
BMI-for-age z score below -2	0.118 P=163	-0.034 [0.038] N=1,353	-0.036 [0.039] N=1,258	-0.027 [0.041] N=1,133	0.012 [0.044] N=969	0.000 [0.048] N=799

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.190 [0.357] N=1,190	0.232 [0.404] N=1,086	0.250 [0.477] N=956	0.645 [0.515] N=788	0.040 [0.691] N=612
WHO weight-for-age z score	-0.190 [0.177] N=1,370	-0.161 [0.196] N=1,247	-0.171 [0.201] N=1,097	-0.249 [0.261] N=901	**-0.868** [0.336] N=695**
WHO weight-for-length/height z score, 0-60 months	0.015 [0.256] N=1,171	-0.073 [0.282] N=1,074	-0.312 [0.357] N=948	-0.609 [0.410] N=780	-0.683 [0.474] N=602
WHO BMI-for-age z score	0.239 [0.277] N=1,145	0.313 [0.315] N=1,050	0.085 [0.400] N=925	-0.439 [0.448] N=761	-0.356 [0.530] N=591
Stunted: HAZ below -2	-0.021 [0.058] N=1,190	-0.040 [0.066] N=1,086	-0.002 [0.071] N=956	-0.022 [0.079] N=788	0.078 [0.108] N=612
Underweight: WAZ below -2	0.016 [0.031] N=1,370	0.006 [0.034] N=1,247	-0.000 [0.036] N=1,097	0.028 [0.045] N=901	**0.138** [0.063] N=695**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.050] N=1,171	-0.009 [0.051] N=1,074	0.001 [0.057] N=948	0.041 [0.072] N=780	0.005 [0.083] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.014 [0.051] N=1,171	-0.028 [0.056] N=1,074	-0.057 [0.066] N=948	-0.076 [0.079] N=780	-0.132 [0.091] N=602
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.037] N=1,171	-0.042 [0.042] N=1,074	-0.034 [0.048] N=948	-0.046 [0.057] N=780	-0.050 [0.066] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.039 [0.067] N=1,136	-0.046 [0.070] N=1,041	-0.001 [0.076] N=918	0.022 [0.087] N=756	0.127 [0.120] N=586
BMI-for-age z score below -2	-0.054 [0.050] N=1,145	-0.060 [0.051] N=1,050	-0.045 [0.059] N=925	0.037 [0.073] N=761	0.004 [0.088] N=591

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.383	0.284 [0.252] N=1,407	0.310 [0.271] N=1,303	0.292 [0.294] N=1,173	0.406 [0.298] N=1,005	0.153 [0.325] N=829
WHO weight-for-age z score	0.092	-0.113 [0.136] N=1,609	-0.086 [0.148] N=1,486	-0.085 [0.153] N=1,336	-0.117 [0.171] N=1,140	**-0.314* [0.186] N=934**
WHO weight-for-length/height z score, 0-60 months	0.391	-0.015 [0.211] N=1,387	-0.066 [0.224] N=1,290	-0.154 [0.255] N=1,164	-0.250 [0.283] N=996	-0.195 [0.310] N=818
WHO BMI-for-age z score	0.372	0.089 [0.244] N=1,353	0.114 [0.265] N=1,258	0.015 [0.301] N=1,133	-0.204 [0.330] N=969	-0.123 [0.363] N=799
Stunted: HAZ below -2	0.245 P=329	-0.054 [0.046] N=1,407	-0.066 [0.049] N=1,303	-0.046 [0.052] N=1,173	-0.054 [0.056] N=1,005	-0.019 [0.064] N=829
Underweight: WAZ below -2	0.068 P=112	0.018 [0.022] N=1,609	0.012 [0.023] N=1,486	0.008 [0.024] N=1,336	0.020 [0.027] N=1,140	0.051 [0.033] N=934
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=153	-0.029 [0.036] N=1,387	-0.024 [0.036] N=1,290	-0.023 [0.038] N=1,164	-0.008 [0.041] N=996	-0.026 [0.043] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=271	0.001 [0.046] N=1,387	-0.005 [0.049] N=1,290	-0.013 [0.055] N=1,164	-0.013 [0.062] N=996	-0.014 [0.069] N=818
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.012 [0.030] N=1,387	-0.023 [0.032] N=1,290	-0.019 [0.034] N=1,164	-0.022 [0.037] N=996	-0.013 [0.040] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	0.326 P=427	-0.078 [0.048] N=1,345	**-0.086* [0.049] N=1,250**	-0.070 [0.051] N=1,127	-0.064 [0.055] N=965	-0.033 [0.062] N=795
BMI-for-age z score below -2	0.118 P=163	-0.034 [0.038] N=1,353	-0.036 [0.039] N=1,258	-0.028 [0.041] N=1,133	0.011 [0.045] N=969	0.001 [0.048] N=799

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.188 [0.356] N=1,190	0.230 [0.403] N=1,086	0.246 [0.473] N=956	0.622 [0.513] N=788	0.019 [0.691] N=612
WHO weight-for-age z score	-0.184 [0.176] N=1,370	-0.152 [0.196] N=1,247	-0.157 [0.201] N=1,097	-0.239 [0.262] N=901	**-0.850** [0.333] N=695**
WHO weight-for-length/height z score, 0-60 months	0.015 [0.256] N=1,171	-0.073 [0.282] N=1,074	-0.306 [0.356] N=948	-0.603 [0.411] N=780	-0.677 [0.481] N=602
WHO BMI-for-age z score	0.241 [0.276] N=1,145	0.316 [0.315] N=1,050	0.094 [0.398] N=925	-0.430 [0.448] N=761	-0.351 [0.538] N=591
Stunted: HAZ below -2	-0.022 [0.058] N=1,190	-0.042 [0.066] N=1,086	-0.006 [0.070] N=956	-0.024 [0.078] N=788	0.077 [0.107] N=612
Underweight: WAZ below -2	0.015 [0.031] N=1,370	0.004 [0.033] N=1,247	-0.003 [0.035] N=1,097	0.027 [0.045] N=901	**0.136** [0.063] N=695**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.022 [0.050] N=1,171	-0.009 [0.052] N=1,074	-0.003 [0.058] N=948	0.039 [0.073] N=780	0.003 [0.085] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.014 [0.051] N=1,171	-0.028 [0.056] N=1,074	-0.058 [0.066] N=948	-0.076 [0.079] N=780	-0.132 [0.091] N=602
Obese: WLZ/WHZ above +3, 0-60 months	-0.022 [0.037] N=1,171	-0.042 [0.042] N=1,074	-0.035 [0.048] N=948	-0.047 [0.057] N=780	-0.050 [0.066] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.041 [0.067] N=1,136	-0.049 [0.069] N=1,041	-0.012 [0.075] N=918	0.014 [0.088] N=756	0.124 [0.121] N=586
BMI-for-age z score below -2	-0.054 [0.050] N=1,145	-0.060 [0.051] N=1,050	-0.047 [0.059] N=925	0.034 [0.074] N=761	0.003 [0.090] N=591

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.383	0.251 [0.249] N=1,407	0.255 [0.270] N=1,303	0.250 [0.292] N=1,173	0.372 [0.291] N=1,005	0.085 [0.314] N=829
WHO weight-for-age z score	0.092	-0.165 [0.130] N=1,609	-0.142 [0.142] N=1,486	-0.136 [0.148] N=1,336	-0.160 [0.165] N=1,140	**-0.367** [0.177] N=934**
WHO weight-for-length/height z score, 0-60 months	0.391	-0.034 [0.213] N=1,387	-0.079 [0.227] N=1,290	-0.157 [0.256] N=1,164	-0.249 [0.282] N=996	-0.208 [0.309] N=818
WHO BMI-for-age z score	0.372	0.077 [0.246] N=1,353	0.109 [0.268] N=1,258	0.012 [0.302] N=1,133	-0.202 [0.329] N=969	-0.132 [0.364] N=799
Stunted: HAZ below -2	0.245 P=329	-0.044 [0.045] N=1,407	-0.050 [0.049] N=1,303	-0.035 [0.051] N=1,173	-0.046 [0.054] N=1,005	-0.003 [0.062] N=829
Underweight: WAZ below -2	0.068 P=112	0.025 [0.021] N=1,609	0.022 [0.022] N=1,486	0.018 [0.022] N=1,336	0.029 [0.026] N=1,140	**0.062** [0.031] N=934**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=153	-0.022 [0.035] N=1,387	-0.015 [0.035] N=1,290	-0.016 [0.038] N=1,164	-0.005 [0.041] N=996	-0.022 [0.043] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=271	0.004 [0.046] N=1,387	0.001 [0.050] N=1,290	-0.008 [0.055] N=1,164	-0.008 [0.062] N=996	-0.011 [0.069] N=818
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.012 [0.030] N=1,387	-0.021 [0.032] N=1,290	-0.016 [0.034] N=1,164	-0.021 [0.037] N=996	-0.014 [0.040] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	0.326 P=427	-0.057 [0.047] N=1,345	-0.057 [0.047] N=1,250	-0.048 [0.050] N=1,127	-0.048 [0.053] N=965	-0.007 [0.060] N=795
BMI-for-age z score below -2	0.118 P=163	-0.027 [0.038] N=1,353	-0.027 [0.039] N=1,258	-0.020 [0.042] N=1,133	0.015 [0.045] N=969	0.004 [0.048] N=799

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.152 [0.358] N=1,190	0.147 [0.407] N=1,086	0.178 [0.478] N=956	0.582 [0.509] N=788	-0.092 [0.677] N=612
WHO weight-for-age z score	-0.226 [0.171] N=1,370	-0.205 [0.192] N=1,247	-0.195 [0.199] N=1,097	-0.258 [0.263] N=901	**-0.877** [0.332] N=695**
WHO weight-for-length/height z score, 0-60 months	0.008 [0.260] N=1,171	-0.066 [0.290] N=1,074	-0.284 [0.361] N=948	-0.605 [0.412] N=780	-0.691 [0.484] N=602
WHO BMI-for-age z score	0.238 [0.280] N=1,145	0.326 [0.322] N=1,050	0.109 [0.401] N=925	-0.446 [0.450] N=761	-0.366 [0.543] N=591
Stunted: HAZ below -2	-0.015 [0.059] N=1,190	-0.023 [0.067] N=1,086	0.004 [0.072] N=956	-0.028 [0.078] N=788	0.088 [0.103] N=612
Underweight: WAZ below -2	0.022 [0.030] N=1,370	0.015 [0.032] N=1,247	0.006 [0.034] N=1,097	0.030 [0.045] N=901	**0.136** [0.062] N=695**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.016 [0.050] N=1,171	0.000 [0.052] N=1,074	0.002 [0.059] N=948	0.041 [0.074] N=780	0.010 [0.088] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.010 [0.052] N=1,171	-0.018 [0.058] N=1,074	-0.052 [0.067] N=948	-0.074 [0.077] N=780	-0.121 [0.086] N=602
Obese: WLZ/WHZ above +3, 0-60 months	-0.021 [0.038] N=1,171	-0.039 [0.042] N=1,074	-0.032 [0.049] N=948	-0.046 [0.057] N=780	-0.052 [0.065] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.024 [0.066] N=1,136	-0.016 [0.069] N=1,041	0.005 [0.074] N=918	0.014 [0.088] N=756	0.150 [0.112] N=586
BMI-for-age z score below -2	-0.047 [0.049] N=1,145	-0.049 [0.052] N=1,050	-0.040 [0.060] N=925	0.037 [0.076] N=761	0.011 [0.093] N=591

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.383	0.180 [0.244] N=1,407	0.191 [0.265] N=1,303	0.191 [0.289] N=1,173	0.303 [0.289] N=1,005	-0.005 [0.309] N=829
WHO weight-for-age z score	0.092	-0.181 [0.130] N=1,609	-0.166 [0.141] N=1,486	-0.159 [0.148] N=1,336	-0.191 [0.167] N=1,140	**-0.412** [0.182] N=934**
WHO weight-for-length/height z score, 0-60 months	0.391	-0.006 [0.210] N=1,387	-0.060 [0.224] N=1,290	-0.145 [0.254] N=1,164	-0.239 [0.285] N=996	-0.196 [0.313] N=818
WHO BMI-for-age z score	0.372	0.110 [0.244] N=1,353	0.135 [0.265] N=1,258	0.037 [0.300] N=1,133	-0.179 [0.333] N=969	-0.102 [0.367] N=799
Stunted: HAZ below -2	0.245 P=329	-0.030 [0.045] N=1,407	-0.036 [0.049] N=1,303	-0.020 [0.051] N=1,173	-0.027 [0.053] N=1,005	0.017 [0.059] N=829
Underweight: WAZ below -2	0.068 P=112	0.027 [0.021] N=1,609	0.024 [0.023] N=1,486	0.021 [0.023] N=1,336	0.032 [0.027] N=1,140	**0.066** [0.033] N=934**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=153	-0.025 [0.035] N=1,387	-0.018 [0.036] N=1,290	-0.017 [0.038] N=1,164	-0.005 [0.042] N=996	-0.025 [0.044] N=818
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=271	0.010 [0.046] N=1,387	0.005 [0.049] N=1,290	-0.006 [0.055] N=1,164	-0.006 [0.063] N=996	-0.011 [0.070] N=818
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=127	-0.010 [0.030] N=1,387	-0.022 [0.032] N=1,290	-0.017 [0.034] N=1,164	-0.021 [0.038] N=996	-0.015 [0.041] N=818
CIAF: any stunting, underweight, or wasting, 0-60 months	0.326 P=427	-0.045 [0.048] N=1,345	-0.044 [0.048] N=1,250	-0.032 [0.051] N=1,127	-0.025 [0.054] N=965	0.015 [0.061] N=795
BMI-for-age z score below -2	0.118 P=163	-0.029 [0.038] N=1,353	-0.029 [0.039] N=1,258	-0.021 [0.042] N=1,133	0.016 [0.045] N=969	0.002 [0.048] N=799

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.118 [0.354] N=1,190	0.140 [0.399] N=1,086	0.196 [0.473] N=956	0.595 [0.499] N=788	-0.152 [0.649] N=612
WHO weight-for-age z score	-0.247 [0.172] N=1,370	-0.229 [0.191] N=1,247	-0.226 [0.198] N=1,097	-0.284 [0.265] N=901	**-0.897*** [0.331] N=695**
WHO weight-for-length/height z score, 0-60 months	0.005 [0.257] N=1,171	-0.074 [0.283] N=1,074	-0.298 [0.357] N=948	-0.580 [0.412] N=780	-0.601 [0.492] N=602
WHO BMI-for-age z score	0.241 [0.278] N=1,145	0.327 [0.316] N=1,050	0.116 [0.398] N=925	-0.383 [0.445] N=761	-0.237 [0.542] N=591
Stunted: HAZ below -2	-0.009 [0.059] N=1,190	-0.023 [0.066] N=1,086	0.001 [0.071] N=956	-0.029 [0.077] N=788	0.090 [0.102] N=612
Underweight: WAZ below -2	0.025 [0.030] N=1,370	0.017 [0.033] N=1,247	0.009 [0.035] N=1,097	0.035 [0.046] N=901	**0.145** [0.065] N=695**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.014 [0.050] N=1,171	-0.002 [0.052] N=1,074	0.002 [0.059] N=948	0.041 [0.075] N=780	0.008 [0.088] N=602
Overweight: WLZ/WHZ above +2, 0-60 months	-0.007 [0.051] N=1,171	-0.018 [0.056] N=1,074	-0.050 [0.065] N=948	-0.067 [0.076] N=780	-0.105 [0.087] N=602
Obese: WLZ/WHZ above +3, 0-60 months	-0.017 [0.038] N=1,171	-0.035 [0.042] N=1,074	-0.029 [0.049] N=948	-0.040 [0.057] N=780	-0.039 [0.065] N=602
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.015 [0.067] N=1,136	-0.015 [0.069] N=1,041	0.008 [0.075] N=918	0.019 [0.089] N=756	0.161 [0.115] N=586
BMI-for-age z score below -2	-0.048 [0.050] N=1,145	-0.057 [0.052] N=1,050	-0.047 [0.060] N=925	0.029 [0.076] N=761	0.000 [0.093] N=591

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.387	0.276 [0.263] N=1,311	0.261 [0.286] N=1,221	0.237 [0.308] N=1,097	0.347 [0.306] N=946	0.113 [0.322] N=779
WHO weight-for-age z score	0.094	-0.062 [0.141] N=1,499	-0.036 [0.150] N=1,391	-0.030 [0.156] N=1,249	-0.044 [0.172] N=1,071	-0.205 [0.195] N=878
WHO weight-for-length/height z score, 0-60 months	0.397	0.110 [0.213] N=1,292	0.080 [0.224] N=1,209	-0.021 [0.256] N=1,090	-0.095 [0.282] N=939	-0.005 [0.307] N=770
WHO BMI-for-age z score	0.382	0.224 [0.248] N=1,260	0.245 [0.266] N=1,178	0.149 [0.300] N=1,061	-0.025 [0.323] N=914	0.115 [0.347] N=752
Stunted: HAZ below -2	0.246 P=307	-0.041 [0.049] N=1,311	-0.049 [0.052] N=1,221	-0.028 [0.055] N=1,097	-0.038 [0.058] N=946	-0.001 [0.065] N=779
Underweight: WAZ below -2	0.067 P=104	0.010 [0.023] N=1,499	0.005 [0.024] N=1,391	0.000 [0.025] N=1,249	0.011 [0.028] N=1,071	0.038 [0.036] N=878
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=142	-0.038 [0.036] N=1,292	-0.034 [0.036] N=1,209	-0.030 [0.038] N=1,090	-0.017 [0.041] N=939	-0.034 [0.043] N=770
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=252	0.015 [0.049] N=1,292	0.012 [0.052] N=1,209	0.004 [0.058] N=1,090	0.005 [0.066] N=939	0.008 [0.074] N=770
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=122	-0.013 [0.032] N=1,292	-0.020 [0.034] N=1,209	-0.014 [0.037] N=1,090	-0.016 [0.040] N=939	-0.005 [0.043] N=770
CIAF: any stunting, underweight, or wasting, 0-60 months	0.328 P=399	-0.068 [0.051] N=1,253	-0.073 [0.051] N=1,172	-0.059 [0.055] N=1,055	-0.064 [0.058] N=910	-0.031 [0.067] N=748
BMI-for-age z score below -2	0.118 P=151	-0.038 [0.038] N=1,260	-0.039 [0.039] N=1,178	-0.032 [0.042] N=1,061	0.003 [0.045] N=914	-0.007 [0.047] N=752

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.259 [0.365] N=1,107	0.259 [0.425] N=1,017	0.263 [0.500] N=893	0.662 [0.530] N=742	0.107 [0.721] N=575
WHO weight-for-age z score	-0.206 [0.185] N=1,275	-0.190 [0.200] N=1,167	-0.206 [0.208] N=1,025	-0.276 [0.261] N=847	**-0.824** [0.365] N=654**
WHO weight-for-length/height z score, 0-60 months	0.003 [0.253] N=1,089	-0.081 [0.272] N=1,006	-0.366 [0.343] N=887	-0.625 [0.407] N=736	-0.588 [0.480] N=567
WHO BMI-for-age z score	0.224 [0.282] N=1,065	0.253 [0.313] N=983	0.011 [0.394] N=866	-0.453 [0.448] N=719	-0.242 [0.538] N=557
Stunted: HAZ below -2	-0.023 [0.062] N=1,107	-0.043 [0.069] N=1,017	-0.007 [0.075] N=893	-0.038 [0.083] N=742	0.066 [0.116] N=575
Underweight: WAZ below -2	0.014 [0.032] N=1,275	0.007 [0.035] N=1,167	0.001 [0.038] N=1,025	0.032 [0.045] N=847	**0.139** [0.069] N=654**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.025 [0.050] N=1,089	-0.012 [0.050] N=1,006	0.000 [0.056] N=887	0.028 [0.072] N=736	-0.029 [0.083] N=567
Overweight: WLZ/WHZ above +2, 0-60 months	-0.014 [0.054] N=1,089	-0.030 [0.057] N=1,006	-0.068 [0.069] N=887	-0.094 [0.084] N=736	-0.126 [0.099] N=567
Obese: WLZ/WHZ above +3, 0-60 months	-0.034 [0.041] N=1,089	-0.047 [0.047] N=1,006	-0.042 [0.053] N=887	-0.056 [0.061] N=736	-0.053 [0.070] N=567
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.035 [0.071] N=1,057	-0.040 [0.072] N=976	-0.009 [0.079] N=859	-0.020 [0.090] N=714	0.074 [0.130] N=552
BMI-for-age z score below -2	-0.049 [0.049] N=1,065	-0.053 [0.048] N=983	-0.040 [0.056] N=866	0.029 [0.073] N=719	-0.015 [0.089] N=557

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.387	0.263 [0.262] N=1,311	0.250 [0.285] N=1,221	0.213 [0.307] N=1,097	0.309 [0.311] N=946	0.034 [0.335] N=779
WHO weight-for-age z score	0.094	-0.071 [0.140] N=1,499	-0.045 [0.149] N=1,391	-0.044 [0.155] N=1,249	-0.065 [0.171] N=1,071	-0.238 [0.193] N=878
WHO weight-for-length/height z score, 0-60 months	0.397	0.100 [0.212] N=1,292	0.067 [0.224] N=1,209	-0.033 [0.255] N=1,090	-0.104 [0.285] N=939	-0.005 [0.315] N=770
WHO BMI-for-age z score	0.382	0.214 [0.247] N=1,260	0.232 [0.265] N=1,178	0.137 [0.300] N=1,061	-0.037 [0.328] N=914	0.109 [0.361] N=752
Stunted: HAZ below -2	0.246 P=307	-0.039 [0.049] N=1,311	-0.047 [0.052] N=1,221	-0.025 [0.055] N=1,097	-0.034 [0.059] N=946	0.007 [0.066] N=779
Underweight: WAZ below -2	0.067 P=104	0.012 [0.022] N=1,499	0.006 [0.024] N=1,391	0.003 [0.025] N=1,249	0.015 [0.027] N=1,071	0.047 [0.035] N=878
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=142	-0.037 [0.036] N=1,292	-0.033 [0.036] N=1,209	-0.030 [0.038] N=1,090	-0.020 [0.043] N=939	-0.042 [0.045] N=770
Overweight: WLZ/WHZ above +2, 0-60 months	0.202 P=252	0.015 [0.049] N=1,292	0.011 [0.052] N=1,209	0.002 [0.058] N=1,090	-0.000 [0.066] N=939	0.007 [0.074] N=770
Obese: WLZ/WHZ above +3, 0-60 months	0.095 P=122	-0.013 [0.032] N=1,292	-0.020 [0.034] N=1,209	-0.013 [0.036] N=1,090	-0.016 [0.040] N=939	-0.003 [0.043] N=770
CIAF: any stunting, underweight, or wasting, 0-60 months	0.328 P=399	-0.064 [0.051] N=1,253	-0.069 [0.052] N=1,172	-0.055 [0.054] N=1,055	-0.060 [0.059] N=910	-0.025 [0.066] N=748
BMI-for-age z score below -2	0.118 P=151	-0.036 [0.038] N=1,260	-0.038 [0.039] N=1,178	-0.032 [0.042] N=1,061	0.002 [0.046] N=914	-0.012 [0.050] N=752

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.282 [0.365] N=1,107	0.277 [0.424] N=1,017	0.277 [0.495] N=893	0.659 [0.543] N=742	0.066 [0.749] N=575
WHO weight-for-age z score	-0.178 [0.182] N=1,275	-0.158 [0.198] N=1,167	-0.185 [0.205] N=1,025	-0.271 [0.261] N=847	**-0.855** [0.367] N=654**
WHO weight-for-length/height z score, 0-60 months	0.032 [0.251] N=1,089	-0.040 [0.271] N=1,006	-0.324 [0.342] N=887	-0.609 [0.409] N=736	-0.606 [0.478] N=567
WHO BMI-for-age z score	0.254 [0.277] N=1,065	0.295 [0.309] N=983	0.047 [0.388] N=866	-0.435 [0.444] N=719	-0.275 [0.529] N=557
Stunted: HAZ below -2	-0.024 [0.062] N=1,107	-0.041 [0.069] N=1,017	-0.005 [0.074] N=893	-0.037 [0.083] N=742	0.064 [0.115] N=575
Underweight: WAZ below -2	0.011 [0.032] N=1,275	0.001 [0.035] N=1,167	-0.005 [0.037] N=1,025	0.028 [0.044] N=847	**0.141** [0.066] N=654**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.027 [0.050] N=1,089	-0.015 [0.050] N=1,006	-0.003 [0.056] N=887	0.027 [0.071] N=736	-0.032 [0.080] N=567
Overweight: WLZ/WHZ above +2, 0-60 months	-0.010 [0.054] N=1,089	-0.024 [0.057] N=1,006	-0.059 [0.069] N=887	-0.087 [0.084] N=736	-0.129 [0.098] N=567
Obese: WLZ/WHZ above +3, 0-60 months	-0.034 [0.041] N=1,089	-0.047 [0.047] N=1,006	-0.040 [0.053] N=887	-0.054 [0.062] N=736	-0.052 [0.070] N=567
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.037 [0.070] N=1,057	-0.042 [0.071] N=976	-0.010 [0.078] N=859	-0.019 [0.089] N=714	0.070 [0.126] N=552
BMI-for-age z score below -2	-0.049 [0.049] N=1,065	-0.055 [0.049] N=983	-0.042 [0.057] N=866	0.028 [0.072] N=719	-0.017 [0.086] N=557

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.151 [0.307] N=1,269	-0.222 [0.315] N=1,147	-0.206 [0.325] N=1,005	-0.409 [0.359] N=852	-0.509 [0.393] N=698
WHO weight-for-age z score	0.116	0.103 [0.143] N=1,433	0.132 [0.149] N=1,304	0.031 [0.147] N=1,143	0.117 [0.156] N=961	0.095 [0.184] N=778
WHO weight-for-length/height z score, 0-60 months	0.383	0.198 [0.214] N=1,256	0.172 [0.223] N=1,137	0.115 [0.238] N=998	0.343 [0.249] N=843	0.208 [0.272] N=690
WHO BMI-for-age z score	0.330	0.150 [0.245] N=1,215	0.118 [0.257] N=1,100	-0.006 [0.277] N=967	0.179 [0.299] N=817	0.105 [0.330] N=672
Stunted: HAZ below -2	0.226 P=287	-0.006 [0.054] N=1,269	-0.006 [0.055] N=1,147	-0.007 [0.058] N=1,005	0.027 [0.064] N=852	0.028 [0.070] N=698
Underweight: WAZ below -2	0.059 P=88	0.006 [0.028] N=1,433	-0.002 [0.029] N=1,304	0.006 [0.031] N=1,143	0.002 [0.033] N=961	-0.004 [0.037] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=136	0.023 [0.037] N=1,256	0.023 [0.039] N=1,137	0.045 [0.043] N=998	-0.031 [0.040] N=843	-0.016 [0.043] N=690
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=247	**0.106** [0.046] N=1,256**	**0.101** [0.047] N=1,137**	**0.104** [0.051] N=998**	**0.119** [0.058] N=843**	0.100 [0.061] N=690
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=120	0.048 [0.039] N=1,256	0.040 [0.039] N=1,137	0.034 [0.043] N=998	0.041 [0.049] N=843	0.008 [0.047] N=690
CIAF: any stunting, underweight, or wasting, 0-60 months	0.309 P=375	0.012 [0.060] N=1,216	0.001 [0.060] N=1,098	0.029 [0.065] N=963	-0.010 [0.066] N=816	0.021 [0.072] N=670
BMI-for-age z score below -2	0.119 P=144	0.036 [0.045] N=1,215	0.035 [0.049] N=1,100	0.045 [0.055] N=967	-0.014 [0.054] N=817	-0.005 [0.057] N=672

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.071 [0.433] N=1,097	0.010 [0.460] N=975	0.054 [0.508] N=833	-0.388 [0.633] N=680	-0.627 [0.786] N=526
WHO weight-for-age z score	0.143 [0.199] N=1,246	0.185 [0.215] N=1,117	0.003 [0.215] N=956	0.192 [0.222] N=774	0.178 [0.304] N=591
WHO weight-for-length/height z score, 0-60 months	0.057 [0.307] N=1,087	-0.012 [0.330] N=968	-0.133 [0.376] N=829	0.356 [0.415] N=674	-0.027 [0.486] N=521
WHO BMI-for-age z score	-0.169 [0.329] N=1,056	-0.253 [0.355] N=941	-0.514 [0.403] N=808	-0.154 [0.466] N=658	-0.429 [0.540] N=513
Stunted: HAZ below -2	-0.011 [0.090] N=1,097	-0.022 [0.093] N=975	-0.015 [0.105] N=833	0.060 [0.131] N=680	0.069 [0.156] N=526
Underweight: WAZ below -2	0.009 [0.037] N=1,246	-0.003 [0.039] N=1,117	0.014 [0.043] N=956	0.002 [0.045] N=774	-0.012 [0.049] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	**0.086* [0.049] N=1,087**	0.092 [0.056] N=968	**0.147** [0.065] N=829**	-0.000 [0.060] N=674	0.037 [0.065] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.144** [0.057] N=1,087**	**0.133** [0.059] N=968**	**0.154** [0.066] N=829**	**0.201** [0.081] N=674**	0.142 [0.091] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.105** [0.041] N=1,087**	**0.091** [0.040] N=968**	**0.096** [0.048] N=829**	**0.120* [0.065] N=674**	0.021 [0.073] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.047 [0.094] N=1,051	0.025 [0.100] N=933	0.094 [0.115] N=798	0.020 [0.132] N=651	0.075 [0.153] N=505
BMI-for-age z score below -2	**0.135*** [0.050] N=1,056**	**0.141** [0.060] N=941**	**0.180** [0.073] N=808**	0.057 [0.068] N=658	0.085 [0.080] N=513

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.157 [0.305] N=1,269	-0.226 [0.314] N=1,147	-0.210 [0.323] N=1,005	-0.416 [0.356] N=852	-0.515 [0.391] N=698
WHO weight-for-age z score	0.116	0.108 [0.142] N=1,433	0.137 [0.148] N=1,304	0.035 [0.146] N=1,143	0.118 [0.157] N=961	0.093 [0.184] N=778
WHO weight-for-length/height z score, 0-60 months	0.383	0.203 [0.214] N=1,256	0.177 [0.223] N=1,137	0.118 [0.238] N=998	0.345 [0.249] N=843	0.211 [0.272] N=690
WHO BMI-for-age z score	0.330	0.151 [0.245] N=1,215	0.118 [0.257] N=1,100	-0.009 [0.276] N=967	0.179 [0.299] N=817	0.104 [0.330] N=672
Stunted: HAZ below -2	0.226 P=287	-0.007 [0.054] N=1,269	-0.008 [0.055] N=1,147	-0.009 [0.058] N=1,005	0.026 [0.064] N=852	0.028 [0.070] N=698
Underweight: WAZ below -2	0.059 P=88	0.004 [0.028] N=1,433	-0.003 [0.029] N=1,304	0.004 [0.031] N=1,143	0.002 [0.033] N=961	-0.003 [0.037] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=136	0.021 [0.037] N=1,256	0.019 [0.040] N=1,137	0.041 [0.043] N=998	-0.032 [0.040] N=843	-0.017 [0.043] N=690
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=247	**0.106** [0.046] N=1,256**	**0.101** [0.047] N=1,137**	**0.103** [0.051] N=998**	**0.119** [0.058] N=843**	0.100 [0.061] N=690
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=120	0.046 [0.039] N=1,256	0.037 [0.038] N=1,137	0.031 [0.043] N=998	0.040 [0.049] N=843	0.008 [0.048] N=690
CIAF: any stunting, underweight, or wasting, 0-60 months	0.309 P=375	0.006 [0.060] N=1,216	-0.007 [0.061] N=1,098	0.021 [0.066] N=963	-0.012 [0.067] N=816	0.020 [0.073] N=670
BMI-for-age z score below -2	0.119 P=144	0.035 [0.045] N=1,215	0.034 [0.049] N=1,100	0.043 [0.055] N=967	-0.014 [0.054] N=817	-0.005 [0.057] N=672

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.069 [0.432] N=1,097	0.012 [0.458] N=975	0.057 [0.505] N=833	-0.407 [0.628] N=680	-0.660 [0.783] N=526
WHO weight-for-age z score	0.154 [0.197] N=1,246	0.198 [0.211] N=1,117	0.015 [0.210] N=956	0.205 [0.219] N=774	0.208 [0.298] N=591
WHO weight-for-length/height z score, 0-60 months	0.068 [0.309] N=1,087	-0.000 [0.332] N=968	-0.126 [0.378] N=829	0.368 [0.419] N=674	0.027 [0.503] N=521
WHO BMI-for-age z score	-0.160 [0.330] N=1,056	-0.242 [0.355] N=941	-0.506 [0.402] N=808	-0.145 [0.468] N=658	-0.382 [0.558] N=513
Stunted: HAZ below -2	-0.014 [0.090] N=1,097	-0.028 [0.093] N=975	-0.022 [0.105] N=833	0.058 [0.131] N=680	0.070 [0.156] N=526
Underweight: WAZ below -2	0.006 [0.037] N=1,246	-0.006 [0.039] N=1,117	0.011 [0.043] N=956	-0.000 [0.045] N=774	-0.017 [0.049] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.080 [0.050] N=1,087	0.082 [0.058] N=968	**0.136** [0.067] N=829**	-0.008 [0.060] N=674	0.023 [0.068] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.145** [0.057] N=1,087**	**0.134** [0.059] N=968**	**0.152** [0.066] N=829**	**0.197** [0.081] N=674**	0.142 [0.092] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.102** [0.041] N=1,087**	**0.087** [0.040] N=968**	**0.090* [0.047] N=829**	**0.113* [0.065] N=674**	0.014 [0.072] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.035 [0.095] N=1,051	0.005 [0.101] N=933	0.071 [0.117] N=798	0.004 [0.133] N=651	0.055 [0.155] N=505
BMI-for-age z score below -2	**0.131** [0.051] N=1,056**	**0.135** [0.061] N=941**	**0.173** [0.073] N=808**	0.052 [0.068] N=658	0.078 [0.082] N=513

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.165 [0.300] N=1,269	-0.221 [0.309] N=1,147	-0.202 [0.319] N=1,005	-0.404 [0.347] N=852	-0.515 [0.380] N=698
WHO weight-for-age z score	0.116	0.098 [0.144] N=1,433	0.135 [0.150] N=1,304	0.038 [0.149] N=1,143	0.124 [0.159] N=961	0.101 [0.186] N=778
WHO weight-for-length/height z score, 0-60 months	0.383	0.201 [0.214] N=1,256	0.176 [0.224] N=1,137	0.123 [0.239] N=998	0.343 [0.249] N=843	0.209 [0.271] N=690
WHO BMI-for-age z score	0.330	0.147 [0.245] N=1,215	0.115 [0.258] N=1,100	-0.004 [0.276] N=967	0.177 [0.297] N=817	0.108 [0.328] N=672
Stunted: HAZ below -2	0.226 P=287	-0.004 [0.053] N=1,269	-0.008 [0.054] N=1,147	-0.011 [0.057] N=1,005	0.024 [0.061] N=852	0.026 [0.068] N=698
Underweight: WAZ below -2	0.059 P=88	0.006 [0.028] N=1,433	-0.003 [0.029] N=1,304	0.004 [0.031] N=1,143	0.001 [0.033] N=961	-0.004 [0.037] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=136	0.021 [0.037] N=1,256	0.019 [0.040] N=1,137	0.040 [0.043] N=998	-0.032 [0.041] N=843	-0.018 [0.044] N=690
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=247	**0.108** [0.046] N=1,256**	**0.101** [0.047] N=1,137**	**0.104** [0.051] N=998**	**0.118** [0.058] N=843**	0.099 [0.062] N=690
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=120	0.047 [0.038] N=1,256	0.037 [0.038] N=1,137	0.031 [0.043] N=998	0.039 [0.049] N=843	0.006 [0.048] N=690
CIAF: any stunting, underweight, or wasting, 0-60 months	0.309 P=375	0.012 [0.060] N=1,216	-0.005 [0.061] N=1,098	0.020 [0.065] N=963	-0.015 [0.067] N=816	0.015 [0.073] N=670
BMI-for-age z score below -2	0.119 P=144	0.036 [0.045] N=1,215	0.035 [0.050] N=1,100	0.043 [0.055] N=967	-0.014 [0.054] N=817	-0.004 [0.058] N=672

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.084 [0.425] N=1,097	0.031 [0.453] N=975	0.103 [0.493] N=833	-0.328 [0.612] N=680	-0.562 [0.751] N=526
WHO weight-for-age z score	0.131 [0.201] N=1,246	0.184 [0.216] N=1,117	0.014 [0.218] N=956	0.208 [0.228] N=774	0.215 [0.304] N=591
WHO weight-for-length/height z score, 0-60 months	0.050 [0.307] N=1,087	-0.005 [0.332] N=968	-0.131 [0.374] N=829	0.346 [0.415] N=674	-0.001 [0.496] N=521
WHO BMI-for-age z score	-0.175 [0.327] N=1,056	-0.249 [0.354] N=941	-0.525 [0.394] N=808	-0.187 [0.455] N=658	-0.416 [0.538] N=513
Stunted: HAZ below -2	-0.018 [0.088] N=1,097	-0.032 [0.092] N=975	-0.031 [0.102] N=833	0.039 [0.126] N=680	0.046 [0.150] N=526
Underweight: WAZ below -2	0.005 [0.038] N=1,246	-0.008 [0.040] N=1,117	0.007 [0.045] N=956	-0.006 [0.047] N=774	-0.024 [0.051] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	0.080 [0.050] N=1,087	0.080 [0.059] N=968	**0.135** [0.068] N=829**	-0.010 [0.062] N=674	0.019 [0.070] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.147** [0.058] N=1,087**	**0.136** [0.060] N=968**	**0.152** [0.066] N=829**	**0.193** [0.082] N=674**	0.138 [0.095] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.100** [0.042] N=1,087**	**0.084** [0.041] N=968**	**0.088* [0.048] N=829**	0.107 [0.066] N=674	0.008 [0.073] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.036 [0.095] N=1,051	0.005 [0.102] N=933	0.066 [0.118] N=798	-0.011 [0.132] N=651	0.036 [0.151] N=505
BMI-for-age z score below -2	**0.128** [0.052] N=1,056**	**0.131** [0.062] N=941**	**0.170** [0.073] N=808**	0.046 [0.069] N=658	0.069 [0.084] N=513

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.115 [0.300] N=1,269	-0.170 [0.309] N=1,147	-0.149 [0.321] N=1,005	-0.382 [0.351] N=852	-0.495 [0.390] N=698
WHO weight-for-age z score	0.116	0.106 [0.146] N=1,433	0.142 [0.152] N=1,304	0.049 [0.151] N=1,143	0.124 [0.162] N=961	0.100 [0.188] N=778
WHO weight-for-length/height z score, 0-60 months	0.383	0.176 [0.212] N=1,256	0.152 [0.221] N=1,137	0.097 [0.234] N=998	0.340 [0.244] N=843	0.223 [0.270] N=690
WHO BMI-for-age z score	0.330	0.124 [0.242] N=1,215	0.093 [0.253] N=1,100	-0.035 [0.269] N=967	0.166 [0.291] N=817	0.126 [0.327] N=672
Stunted: HAZ below -2	0.226 P=287	-0.014 [0.052] N=1,269	-0.017 [0.053] N=1,147	-0.019 [0.057] N=1,005	0.020 [0.061] N=852	0.025 [0.068] N=698
Underweight: WAZ below -2	0.059 P=88	0.003 [0.028] N=1,433	-0.005 [0.030] N=1,304	-0.000 [0.032] N=1,143	-0.001 [0.034] N=961	-0.010 [0.039] N=778
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=136	0.023 [0.037] N=1,256	0.021 [0.040] N=1,137	0.044 [0.043] N=998	-0.031 [0.040] N=843	-0.019 [0.043] N=690
Overweight: WLZ/WHZ above +2, 0-60 months	0.192 P=247	**0.104** [0.045] N=1,256**	**0.098** [0.046] N=1,137**	**0.102** [0.049] N=998**	**0.119** [0.055] N=843**	**0.102* [0.059] N=690**
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=120	0.048 [0.038] N=1,256	0.040 [0.037] N=1,137	0.032 [0.042] N=998	0.043 [0.047] N=843	0.011 [0.046] N=690
CIAF: any stunting, underweight, or wasting, 0-60 months	0.309 P=375	0.007 [0.059] N=1,216	-0.008 [0.060] N=1,098	0.018 [0.065] N=963	-0.012 [0.066] N=816	0.023 [0.071] N=670
BMI-for-age z score below -2	0.119 P=144	0.033 [0.045] N=1,215	0.032 [0.049] N=1,100	0.042 [0.055] N=967	-0.017 [0.053] N=817	-0.011 [0.057] N=672

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.144 [0.414] N=1,097	0.101 [0.440] N=975	0.175 [0.482] N=833	-0.347 [0.599] N=680	-0.668 [0.730] N=526
WHO weight-for-age z score	0.172 [0.195] N=1,246	0.222 [0.211] N=1,117	0.067 [0.209] N=956	0.236 [0.218] N=774	0.235 [0.283] N=591
WHO weight-for-length/height z score, 0-60 months	0.033 [0.307] N=1,087	-0.035 [0.328] N=968	-0.168 [0.368] N=829	0.349 [0.415] N=674	0.075 [0.516] N=521
WHO BMI-for-age z score	-0.200 [0.325] N=1,056	-0.282 [0.350] N=941	-0.567 [0.388] N=808	-0.183 [0.455] N=658	-0.322 [0.564] N=513
Stunted: HAZ below -2	-0.030 [0.087] N=1,097	-0.048 [0.090] N=975	-0.046 [0.102] N=833	0.037 [0.127] N=680	0.063 [0.150] N=526
Underweight: WAZ below -2	0.003 [0.037] N=1,246	-0.009 [0.039] N=1,117	0.002 [0.045] N=956	-0.007 [0.046] N=774	-0.030 [0.052] N=591
Wasted: WLZ/WHZ below -2, 0-60 months	**0.083* [0.050] N=1,087**	0.086 [0.058] N=968	**0.143** [0.066] N=829**	-0.006 [0.061] N=674	0.017 [0.068] N=521
Overweight: WLZ/WHZ above +2, 0-60 months	**0.140** [0.058] N=1,087**	**0.129** [0.059] N=968**	**0.146** [0.065] N=829**	**0.192** [0.081] N=674**	0.150 [0.094] N=521
Obese: WLZ/WHZ above +3, 0-60 months	**0.100** [0.042] N=1,087**	**0.086** [0.041] N=968**	**0.087* [0.047] N=829**	**0.111* [0.065] N=674**	0.016 [0.074] N=521
CIAF: any stunting, underweight, or wasting, 0-60 months	0.026 [0.094] N=1,051	-0.005 [0.101] N=933	0.053 [0.117] N=798	-0.012 [0.132] N=651	0.054 [0.151] N=505
BMI-for-age z score below -2	**0.130** [0.051] N=1,056**	**0.134** [0.061] N=941**	**0.175** [0.072] N=808**	0.052 [0.068] N=658	0.072 [0.082] N=513

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.188 [0.303] N=1,182	-0.274 [0.317] N=1,067	-0.224 [0.324] N=939	-0.471 [0.353] N=795	-0.640 [0.387] N=652
WHO weight-for-age z score	0.114	0.147 [0.149] N=1,334	0.187 [0.156] N=1,212	0.070 [0.154] N=1,066	0.138 [0.162] N=897	0.060 [0.183] N=729
WHO weight-for-length/height z score, 0-60 months	0.386	0.273 [0.230] N=1,170	0.250 [0.245] N=1,058	0.142 [0.261] N=931	0.366 [0.273] N=786	0.224 [0.295] N=645
WHO BMI-for-age z score	0.331	0.165 [0.262] N=1,132	0.136 [0.280] N=1,024	-0.039 [0.299] N=903	0.152 [0.321] N=763	0.044 [0.354] N=629
Stunted: HAZ below -2	0.227 P=268	-0.001 [0.057] N=1,182	0.003 [0.058] N=1,067	-0.006 [0.061] N=939	0.026 [0.066] N=795	0.040 [0.074] N=652
Underweight: WAZ below -2	0.058 P=81	0.001 [0.028] N=1,334	-0.009 [0.030] N=1,212	-0.005 [0.032] N=1,066	-0.005 [0.033] N=897	-0.003 [0.037] N=729
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=127	0.020 [0.039] N=1,170	0.019 [0.042] N=1,058	0.047 [0.044] N=931	-0.021 [0.043] N=786	-0.004 [0.046] N=645
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=230	**0.117** [0.047] N=1,170**	**0.112** [0.048] N=1,058**	**0.106** [0.052] N=931**	**0.117** [0.058] N=786**	0.096 [0.062] N=645
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=115	0.055 [0.038] N=1,170	0.045 [0.038] N=1,058	0.034 [0.042] N=931	0.048 [0.048] N=786	0.016 [0.048] N=645
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=352	0.017 [0.062] N=1,133	0.007 [0.064] N=1,022	0.034 [0.068] N=899	-0.004 [0.069] N=761	0.046 [0.075] N=626
BMI-for-age z score below -2	0.121 P=135	0.043 [0.047] N=1,132	0.041 [0.051] N=1,024	0.053 [0.056] N=903	-0.000 [0.056] N=763	0.016 [0.059] N=629

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.239 [0.430] N=1,019	0.172 [0.464] N=904	0.284 [0.507] N=776	-0.180 [0.608] N=632	-0.445 [0.753] N=489
WHO weight-for-age z score	0.260 [0.198] N=1,157	0.323 [0.210] N=1,035	0.112 [0.209] N=889	0.325 [0.212] N=720	0.266 [0.280] N=552
WHO weight-for-length/height z score, 0-60 months	0.125 [0.333] N=1,010	0.062 [0.364] N=898	-0.158 [0.405] N=771	0.324 [0.445] N=626	-0.048 [0.535] N=485
WHO BMI-for-age z score	-0.153 [0.358] N=980	-0.235 [0.396] N=872	-0.609 [0.439] N=751	-0.216 [0.502] N=611	-0.562 [0.603] N=477
Stunted: HAZ below -2	-0.036 [0.096] N=1,019	-0.046 [0.100] N=904	-0.053 [0.112] N=776	0.006 [0.135] N=632	0.014 [0.168] N=489
Underweight: WAZ below -2	-0.004 [0.038] N=1,157	-0.021 [0.040] N=1,035	-0.012 [0.044] N=889	-0.026 [0.046] N=720	-0.041 [0.047] N=552
Wasted: WLZ/WHZ below -2, 0-60 months	0.078 [0.054] N=1,010	0.080 [0.062] N=898	**0.141** [0.069] N=771**	-0.000 [0.066] N=626	0.033 [0.073] N=485
Overweight: WLZ/WHZ above +2, 0-60 months	**0.159** [0.062] N=1,010**	**0.146** [0.065] N=898**	**0.149** [0.072] N=771**	**0.182** [0.089] N=626**	0.128 [0.103] N=485
Obese: WLZ/WHZ above +3, 0-60 months	**0.103** [0.045] N=1,010**	**0.085* [0.045] N=898**	0.079 [0.051] N=771	0.106 [0.070] N=626	-0.001 [0.079] N=485
CIAF: any stunting, underweight, or wasting, 0-60 months	0.021 [0.102] N=977	-0.007 [0.110] N=866	0.058 [0.126] N=743	-0.040 [0.143] N=605	0.025 [0.169] N=470
BMI-for-age z score below -2	**0.144** [0.056] N=980**	**0.147** [0.067] N=872**	**0.189** [0.077] N=751**	0.070 [0.074] N=611	0.112 [0.087] N=477

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.263	-0.222 [0.311] N=1,182	-0.314 [0.323] N=1,067	-0.267 [0.330] N=939	-0.518 [0.360] N=795	**-0.689* [0.397] N=652**
WHO weight-for-age z score	0.114	0.145 [0.150] N=1,334	0.179 [0.157] N=1,212	0.063 [0.153] N=1,066	0.126 [0.163] N=897	0.053 [0.185] N=729
WHO weight-for-length/height z score, 0-60 months	0.386	0.296 [0.230] N=1,170	0.270 [0.244] N=1,058	0.168 [0.259] N=931	0.386 [0.270] N=786	0.246 [0.295] N=645
WHO BMI-for-age z score	0.331	0.192 [0.265] N=1,132	0.160 [0.283] N=1,024	-0.008 [0.302] N=903	0.175 [0.322] N=763	0.071 [0.360] N=629
Stunted: HAZ below -2	0.227 P=268	0.001 [0.057] N=1,182	0.004 [0.058] N=1,067	-0.004 [0.061] N=939	0.030 [0.066] N=795	0.044 [0.075] N=652
Underweight: WAZ below -2	0.058 P=81	0.001 [0.028] N=1,334	-0.009 [0.029] N=1,212	-0.004 [0.031] N=1,066	-0.004 [0.033] N=897	-0.002 [0.037] N=729
Wasted: WLZ/WHZ below -2, 0-60 months	0.108 P=127	0.016 [0.040] N=1,170	0.015 [0.042] N=1,058	0.040 [0.045] N=931	-0.027 [0.043] N=786	-0.009 [0.046] N=645
Overweight: WLZ/WHZ above +2, 0-60 months	0.193 P=230	**0.118** [0.046] N=1,170**	**0.113** [0.047] N=1,058**	**0.105** [0.051] N=931**	**0.118** [0.057] N=786**	0.099 [0.061] N=645
Obese: WLZ/WHZ above +3, 0-60 months	0.093 P=115	0.055 [0.038] N=1,170	0.044 [0.037] N=1,058	0.033 [0.042] N=931	0.048 [0.048] N=786	0.017 [0.047] N=645
CIAF: any stunting, underweight, or wasting, 0-60 months	0.312 P=352	0.014 [0.062] N=1,133	0.002 [0.063] N=1,022	0.029 [0.067] N=899	-0.008 [0.069] N=761	0.045 [0.075] N=626
BMI-for-age z score below -2	0.121 P=135	0.038 [0.048] N=1,132	0.036 [0.053] N=1,024	0.046 [0.058] N=903	-0.005 [0.057] N=763	0.012 [0.061] N=629

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.188 [0.442] N=1,019	0.122 [0.471] N=904	0.218 [0.514] N=776	-0.291 [0.626] N=632	-0.608 [0.782] N=489
WHO weight-for-age z score	0.249 [0.201] N=1,157	0.304 [0.217] N=1,035	0.092 [0.213] N=889	0.293 [0.220] N=720	0.235 [0.291] N=552
WHO weight-for-length/height z score, 0-60 months	0.135 [0.330] N=1,010	0.063 [0.359] N=898	-0.146 [0.402] N=771	0.355 [0.442] N=626	-0.027 [0.529] N=485
WHO BMI-for-age z score	-0.134 [0.356] N=980	-0.227 [0.390] N=872	-0.585 [0.437] N=751	-0.184 [0.498] N=611	-0.498 [0.603] N=477
Stunted: HAZ below -2	-0.031 [0.096] N=1,019	-0.044 [0.099] N=904	-0.048 [0.111] N=776	0.019 [0.136] N=632	0.038 [0.170] N=489
Underweight: WAZ below -2	-0.003 [0.038] N=1,157	-0.019 [0.040] N=1,035	-0.008 [0.044] N=889	-0.018 [0.046] N=720	-0.026 [0.049] N=552
Wasted: WLZ/WHZ below -2, 0-60 months	0.076 [0.054] N=1,010	0.080 [0.061] N=898	**0.137** [0.069] N=771**	-0.006 [0.065] N=626	0.028 [0.073] N=485
Overweight: WLZ/WHZ above +2, 0-60 months	**0.161*** [0.061] N=1,010**	**0.146** [0.064] N=898**	**0.148** [0.071] N=771**	**0.185** [0.088] N=626**	0.129 [0.102] N=485
Obese: WLZ/WHZ above +3, 0-60 months	**0.103** [0.045] N=1,010**	**0.085* [0.044] N=898**	0.078 [0.050] N=771	0.108 [0.070] N=626	0.002 [0.079] N=485
CIAF: any stunting, underweight, or wasting, 0-60 months	0.022 [0.102] N=977	-0.006 [0.109] N=866	0.057 [0.124] N=743	-0.035 [0.143] N=605	0.041 [0.169] N=470
BMI-for-age z score below -2	**0.141** [0.055] N=980**	**0.146** [0.066] N=872**	**0.186** [0.076] N=751**	0.066 [0.072] N=611	0.106 [0.086] N=477

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.